



**HAL**  
open science

## **Urinary Peptides as Potential Non-Invasive Biomarkers for Lupus Nephritis: Results of the Peptidu-LUP Study**

Maxence Tailliar, Joost Schanstra, Tim Dierckx, Benjamin Breuil, Guillaume Hanouna, Nicolas Charles, Jean-Loup Bascands, Bertrand Dussol, Alain Vazi, Laurent Chiche, et al.

### ► **To cite this version:**

Maxence Tailliar, Joost Schanstra, Tim Dierckx, Benjamin Breuil, Guillaume Hanouna, et al.. Urinary Peptides as Potential Non-Invasive Biomarkers for Lupus Nephritis: Results of the Peptidu-LUP Study. *Journal of Clinical Medicine*, 2021, 10 (8), pp.1-17. <10.3390/jcm10081690>. <hal-03268297>

**HAL Id: hal-03268297**

**<https://hal.science/hal-03268297v1>**

Submitted on 10 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Article

# Urinary Peptides as Potential Non-Invasive Biomarkers for Lupus Nephritis: Results of the Peptidu-LUP Study

Maxence Tailliar <sup>1,†</sup>, Joost P. Schanstra <sup>2,3,†</sup>, Tim Dierckx <sup>4</sup> , Benjamin Breuil <sup>2,3</sup>, Guillaume Hanouna <sup>5</sup>, Nicolas Charles <sup>6</sup> , Jean-Loup Bascands <sup>7</sup> , Bertrand Dussol <sup>1,8</sup>, Alain Vazi <sup>8</sup>, Laurent Chiche <sup>9</sup>, Justyna Siwy <sup>10</sup>, Stanislas Faguer <sup>2,3,11</sup>, Laurent Daniel <sup>12,13</sup>, Eric Daugas <sup>5,6</sup>, Noémie Jourde-Chiche <sup>1,13,\*</sup>   
and on behalf of the Groupe Coopératif sur le Lupus Rénal (GCLR) ‡

<sup>1</sup> AP-HM, Centre de Néphrologie et Transplantation Rénale, Hôpital de la Conception, 13005 Marseille, France; maxence.tailliar@ap-hm.fr (M.T.); bertrand.dussol@ap-hm.fr (B.D.)

<sup>2</sup> Institut National de la Santé et de la Recherche Médicale (INSERM), U1297, Institut of Metabolic and Cardiovascular Disease (I2MC), 31432 Toulouse, France; joost-peter.schanstra@inserm.fr (J.P.S.); benjamin.breuil@inserm.fr (B.B.); faguer.s@chu-toulouse.fr (S.F.)

<sup>3</sup> Université Toulouse III Paul-Sabatier, 31062 Toulouse, France

<sup>4</sup> Laboratory of Clinical and Epidemiological Virology, Department of Microbiology and Immunology, Rega Institute for Medical Research, KU Leuven, 3000 Leuven, Belgium; tim.dierckx@kuleuven.be

<sup>5</sup> AP-HP, Service de Néphrologie, Hôpital Bichat, DMU VICTOIRE, 75018 Paris, France; guillaume.hanouna@aphp.fr (G.H.); eric.daugas@aphp.fr (E.D.)

<sup>6</sup> Centre de Recherche sur l'Inflammation, Université de Paris, INSERM UMRS1149, CNRS ERL8252, Labex INFLAMEX, DHU FIRE, 75890 Paris, France; nicolas.charles@inserm.fr

<sup>7</sup> Institut National de la Santé et de la Recherche Médicale (INSERM), U1188-Université de La Réunion, 97490 Saint-Denis, France; jean-loup.bascands@inserm.fr

<sup>8</sup> Centre d'Investigation Clinique, CHU Conception, AP-HM, 13005 Marseille, France; alain.vazi@ap-hm.fr

<sup>9</sup> Médecine Interne, Hôpital Européen, 13003 Marseille, France; l.chiche@hopital-europeen.fr

<sup>10</sup> Mosaiques Diagnostics GmbH, 30659 Hannover, Germany; siwy@mosaiques.de

<sup>11</sup> CHU de Toulouse, Service de Néphrologie, 31300 Toulouse, France

<sup>12</sup> AP-HM, Laboratoire d'Anatomie Pathologique, Hôpital de la Timone, 13005 Marseille, France; laurent.daniel@ap-hm.fr

<sup>13</sup> Center for CardioVascular and Nutrition Research (C2VN), Aix-Marseille University, INSERM, INRAE, 13005 Marseille, France

\* Correspondence: noemie.jourde@ap-hm.fr

† These authors contributed equally to this work.

‡ Membership of the Groupe Coopératif sur le Lupus Rénal (GCLR) is provided in the Acknowledgements.



**Citation:** Tailliar, M.; Schanstra, J.P.; Dierckx, T.; Breuil, B.; Hanouna, G.; Charles, N.; Bascands, J.-L.; Dussol, B.; Vazi, A.; Chiche, L.; et al. Urinary Peptides as Potential Non-Invasive Biomarkers for Lupus Nephritis: Results of the Peptidu-LUP Study. *J. Clin. Med.* **2021**, *10*, 1690. <https://doi.org/10.3390/jcm10081690>

Academic Editor: Alberto Martínez-Castelao

Received: 15 February 2021

Accepted: 9 April 2021

Published: 14 April 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** *Background:* Lupus nephritis (LN) is a severe manifestation of Systemic Lupus Erythematosus (SLE). The therapeutic strategy relies on kidney biopsy (KB) results. We tested whether urinary peptidome analysis could non-invasively differentiate active from non-active LN. *Design:* Urinary samples were collected from 93 patients (55 with active LN and 38 with non-active LN), forming a discovery ( $n = 42$ ) and an independent validation ( $n = 51$ ) cohort. Clinical characteristics were collected at inclusion and prospectively for 24 months. The urinary peptidome was analyzed by capillary-electrophoresis coupled to mass-spectrometry, comparing active LN to non-active LN, and assessing chronic lesions and response to therapy. The value of previously validated prognostic (CKD273) and differential diagnostic (LN172) signatures was evaluated. *Results:* Urinary peptides could not discriminate between active and non-active LN or predict early response to therapy. Tubulo-interstitial fibrosis was correlated to the CKD273. The LN172 score identified 92.5% of samples as LN. Few patients developed new-onset CKD. *Conclusions:* We validated the CKD273 and LN172 classifiers but did not identify a robust signature that could predict active LN and replace KB. The value of urinary peptidome to predict long-term CKD, or renal flares in SLE, remains to be evaluated.

**Keywords:** lupus nephritis; systemic lupus erythematosus; non-invasive; biomarker; proteomics; peptidomics; urine; kidney biopsy; classification; prognosis

## 1. Introduction

Systemic Lupus Erythematosus (SLE) is a systemic auto-immune disease which can affect many organs. Lupus Nephritis (LN), which occurs in 20–70% of patients according to ethnicity [1], worsens the prognosis of SLE [2] and leads to End Stage Kidney Disease (ESKD) in 10% of patients after 10 years [3]. The treatment of LN is based on histopathological classification [4,5] which mainly distinguishes patients with active proliferative lesions (class III-A or class IV-A  $\pm$  C  $\pm$  V LN according to the ISN/RPS classification [5]) from patients with non-active LN (other LN classes). Patients with non-active LN will receive antimalarials and renin angiotensin aldosterone system (RAAS) blockers. Active proliferative LN requires additional immunosuppressive therapy, until remission is achieved, followed by maintenance immunosuppressive therapy to avoid relapses [6,7]. Lesions of tubulointerstitial nephritis (TIN) can be associated to the specific glomerular rearrangements and their impact on LN outcome has been highlighted recently [8]. Interstitial inflammation on initial KB in patients with LN can predict long-term renal survival [9], even among patients with class IV LN. The severity of interstitial fibrosis and tubular atrophy (IF/TA) also independently predicts long-term renal survival. In a murine model of LN (NZB/W mice), the infiltration of renal tubulointerstitium by anti-dsDNA-secreting plasma cells could suggest a specific role of TIN in the onset of LN [10]. Because proteinuria, urinary sediment, serum creatinine or immunological markers (anti-dsDNA, complement fractions) are poorly correlated with pathological lesions and because there is to date no specific biomarker of TIN lesions, a kidney biopsy (KB) is required for every suspected flare of LN [11]. Nevertheless, KB is associated with significant complications [12] such as macroscopic hematuria, pain, hematoma, or in exceptional cases, nephrectomy or death. As such, there is a need for non-invasive biomarkers, such as urinary biomarkers, to predict the pathological severity of LN and to predict the response to therapy and the risk of chronic kidney disease (CKD) in patients with LN.

For example, promising predictive values for LN activity/severity and/or the prediction of renal flares have been reported for urinary cytokines or markers of tubular injury, either alone [13] or combined [14], or the quantification of inflammatory cell markers [15]. However, the validation of these urinary biomarkers in independent cohorts is still awaited. Because LN pathogenicity is a complex problem, it is likely that a combination of non-invasive biomarkers is to be preferred [16]. As such, urinary proteomics or peptidomics (i.e., small molecular weight proteins) is an appealing approach for the evaluation of local tissue inflammation and damage.

The analysis of urinary peptides shows a strong reproducibility under different storage conditions [17]. Using capillary-electrophoresis coupled to mass-spectrometry (CE-MS), urinary peptidome analysis has been used as a prognostic or diagnostic tool in fetal medicine [18–20], in CKD with the use of the urinary peptide CKD273 classifier [20–23] and in different glomerular diseases [24–26]. In the field of LN, previous studies investigating the urinary proteome used older technologies, small study populations or lacked validation cohorts [27–32]. Studies focusing on the urinary peptidome did not consider the difference between active LN and non-active LN, and instead aimed to discriminate LN from healthy controls or from other renal diseases [33,34]. The LN172 urinary peptide classifier, in particular, was able to differentiate LN from CKD with other etiologies (focal segmental glomerulosclerosis, minimal-change disease, membranous nephropathy, diabetic nephropathy, hypertensive nephrosclerosis and vasculitis-induced kidney disease) with a good accuracy (AUC 0.82) [35].

In the present study, we aimed to assess, using urinary peptidomics, whether we could identify urinary peptides that discriminated between active and non-active LN, in patients undergoing a KB for a suspicion of LN flare. The secondary objectives were to investigate if the urinary peptidome could predict: (1) glomerular chronic lesions, (2) IF/TA, (3) renal response to treatment, (4) renal flares and (5) renal function after 24 months. We also evaluated the performances of the previously established CKD273 and LN172 classifiers in this cohort.

## 2. Materials and Methods

### 2.1. Patients' Characteristics

Patients were included from July 2011 to December 2016 in 3 French centers participating in the "Groupe Coopératif sur le Lupus Rénal" (GCLR). The patients of the discovery cohort ( $n = 42$ ) were included from July 2011 to November 2012, and the patients of the validation cohort ( $n = 51$ ) were included from September 2014 to December 2016. All patients signed an informed consent before any study procedure, and the samples were included in the biological collections of these 3 centers (Marseille: DC 2012-1704; Paris Bichat: ID-RCB 2014-A00809-38; and Toulouse: DC 2011-1388). Inclusion criteria were: age  $\geq 18$  years, SLE according to the ACR 1997 [36] or SLICC 2012 [37] classification criteria, urinary sample collected at the time of KB, biopsy including  $\geq 10$  glomeruli. Clinical data and laboratory results were collected on the day of KB, and during the follow-up.

Initial clinical characteristics included age, sex and ethnicity. Serum creatinine and estimated-Glomerular Filtration-Rate (eGFR) using the MDRD formula [38], urinary protein/creatinine ratio (UPCR), urinary sediment, complement and anti-dsDNA antibodies were collected at the time of KB and during the longitudinal follow-up. Therapeutic regimens for SLE before the renal flare, at the time of KB, and for the flare of LN, were recorded. The renal response to treatment was evaluated using eGFR and UPCR at M6, M12 and M18, and the long-term renal function was estimated from the same parameters measured at M24 and at the end of follow-up. Remission was defined with 3 possible thresholds of UPCR (0.2 g/g, 0.5 g/g or 0.7 g/g), together with a return to baseline eGFR or a decrease in eGFR not exceeding 20%. CKD was defined by a persistent decline of eGFR  $< 60$  mL/min/1.73 m<sup>2</sup>. LN relapse was defined as a new KB showing active lesions and leading to therapeutic implementation.

### 2.2. Kidney Biopsy Pathological Analysis

Pathological data were interpreted according to the ISN/RPS 2003 classification [5], with the addition of IF/TA evaluation (F0:  $\leq 5\%$ ; F1: 6–25%; F2: 26–50%; F3:  $>50\%$ ) according to the Banff classification [39]. Active LN was defined as class III or IV LN with active lesions,  $\pm$  associated class V. Non-active LN was defined as class II LN, isolated class V LN or the presence of chronic lesions only.

### 2.3. Sample Preparation

Fresh urine was collected on the morning of the kidney biopsy, centrifuged to remove the cell pellet, and stored at  $-80$  °C. Urine samples were thawed immediately before use. A volume of 0.7 mL was diluted with 0.7 mL 2 M urea, 10 mM NH<sub>4</sub>OH and 0.02% sodium dodecyl sulphate (SDS). In order to remove high molecular weight polypeptides, samples were filtered using Centriscart ultracentrifugation filter devices (20 kDa molecular weight cut-off; Sartorius, Goettingen, Germany) at  $3000 \times g$  until 1.1 mL of filtrate was obtained. The filtrate was desalted with PD-10 column (GE Healthcare, Sweden) equilibrated in 0.01% NH<sub>4</sub>OH in HPLC-grade water. The prepared samples were lyophilized and stored at 4 °C. Shortly before CE-MS analysis, lyophilized samples were resuspended in HPLC-grade water (Merck KGaA, Darmstadt, Germany). The preparation method has previously been described in more detail [40].

### 2.4. CE-MS Analysis and Data Processing

CE-MS analysis was performed as previously described [41]. Briefly, CE-MS analyses were performed using a Beckman Coulter Proteome Lab PA800 capillary electrophoresis system (Beckman Coulter, Fullerton, CA, USA) on-line coupled to a micrOTOF II MS (Bruker Daltonic, Bremen, Germany). The electro-ionization sprayer (Agilent Technologies, Palo Alto, CA, USA) was grounded, and the ion spray interface potential was set to  $-4.5$  kV. Data acquisition and MS acquisition methods were automatically controlled by the CE via contact-close-relays. Spectra were accumulated every 3 s, over a range of  $m/z$  350 to 3000. In the next step the MosaiquesVisu software package was applied to deconvolute

mass spectral ion peaks, because ionization produced ions at different charged states from the original urinary peptides. MosaiquesVisu was used to deconvolute mass spectral peaks representing identical molecules into singles masses. The obtained peak list of each polypeptide is characterized by molecular mass (in Daltons), CE-migration time (in minutes), and normalized ion signal intensity. MS signal intensities were used as measure of relative abundance and normalized using 29 internal standard peptides as described by Jantos-Siwy et al. [42]. All detected peptides were deposited, matched and annotated in a Microsoft SQL database, permitting further correlation and statistical analysis. Raw peptide data can be found in Supplementary Table S1.

### 2.5. Biomarker Selection and Modelling

For the identification of new candidate urinary biomarkers for LN activity, the reported *p*-values were calculated using the Wilcoxon rank sum test (R software, version 3.1.3) comparing active LN to non-active LN samples followed by adjustment for multiple testing (Benjamini and Hochberg, 1995). Peptides that were detectable in at least 60% of the samples in one of the two groups (active LN or non-active LN) and reached an adjusted *p*-value of <0.05 were considered as potential biomarker peptides. The R package randomForest (version 4.6-14) was used to generate biomarker models that consisted of an ensemble of 500 trees sampling 5 predictors per split. The overall yield of the polypeptide pattern was evaluated by receiver operating characteristic (ROC) and area under curve (AUC) plots using the Prism 7.00 GraphPad software.

We also tested previously validated peptidome classifiers, using the same algorithms as in the original publications: CKD273 [21,22,43] as a continuous variable, and LN172 both as a continuous and categorical variable [35].

## 3. Results

### 3.1. Patients' Characteristics

The whole cohort comprised 86 women (92.5%) and 7 men (7.5%). Median age was 35 years, most patients were of European (43%) or North African (24.7%) origin. The demographic and renal characteristics at baseline are described in Table 1. There was a majority of active LN (59.1% of class III or IV ± V with active lesions). A large majority of patients had previously received hydroxychloroquine (HCQ) (80.6%) or corticosteroids (CS) (77.4%), and 33.3% had been treated with Cyclophosphamide. At the time of KB, 68.8% of patients were prescribed HCQ, 70.1% CS, 17.2% mycophenolate mofetil (MMF) and 9.7% Azathioprine (AZA).

**Table 1.** Patients' baseline characteristics.

Characteristics	
Patients, <i>n</i>	93
Women— <i>n</i> (%)	86 (92.4)
Age—years	35 ± 11
Range—years	19–58
Ethnicity— <i>n</i> (%)	
European	40 (43)
North African	23 (24.7)
African	23 (23.7)
Asian	8 (8.6)
Characteristics at kidney biopsy	
Serum creatinine— $\mu\text{mol/L}$	92.1 ± 68.5
eGFR— $\text{mL}/\text{min}/1.73 \text{ m}^2$	91.4 ± 39.2
UPCR— $\text{g}/\text{g}$	2.34 ± 2.23

Table 1. Cont.

Characteristics	
SLE vintage	
First flare of SLE— <i>n</i> (%)	21 (22.6)
Pre-existing SLE— <i>n</i> (%)	72 (77.4)
Disease duration—years	10.6 ± 8.5
First renal flare— <i>n</i> (%)	55 (59.1)
LN duration—years	8.2 ± 6.9
Kidney biopsies— <i>n</i> (%)	
Class ISN/RPS 2003	
I	1 (1.1)
II	6 (6.5)
III-A or -A/C ± V	22 (23.7)
III-C ± V	5 (5.4)
IV-A or -A/C ± V	33 (35.5)
IV-C ± V	2 (2.2)
Pure V	21 (22.6)
VI	3 (3.2)
Group	
Active LN	55 (59.1)
Non-active LN	38 (40.9)
IF/TA	
F0	41 (44.1)
F1	33 (35.5)
F2	12 (12.9)
F3	7 (7.5)
Previous treatment for SLE (%)	
Hydroxychloroquine	80.6
Corticosteroids	77.4
MMF/Mycophenolic acid	25.8
Azathioprine	19.4
Cyclophosphamide	33.3
Rituximab	8.6
Other	14
Treatment for SLE at inclusion (%)	
Hydroxychloroquine	68.8
Corticosteroids	70.1
MMF/Mycophenolic acid	17.2
Azathioprine	9.7
Cyclophosphamide	1.1
Rituximab	1.1
Other	2.2
Patients/center— <i>n</i> (%)	
Marseille	37 (39.8)
Paris	53 (57)
Toulouse	3 (3.2)

Continuous values expressed in mean value ± SD. MMF, mycophenolate mofetil; eGFR, estimated glomerular filtration rate; UPCR, urine protein to creatinine ratio; SLE, Systemic Lupus Erythematosus; LN, Lupus nephritis.

The mean follow-up was 43.2 months; 9 patients were lost to follow-up (Table 2). At least one relapse of LN occurred in 19 (20.4%) patients after a median of 12 months; 68.4% of relapses occurred within two years of the previous flare. At the end of the follow-up, 12 (12.9%) patients displayed CKD, among whom 9 had preexisting CKD at inclusion (only 3 patients developed new-onset CKD); 5 patients progressed to ESKD requiring renal replacement therapy (4 renal transplantations, 1 hemodialysis); 4 (4.3%) patients died during the follow-up.

**Table 2.** Outcomes of patients from the discovery and validation cohorts.

	Discovery Cohort	Validation Cohort	<i>p</i>
Follow-up—months, median [IQR]	63.5 (51–69)	32 (24–39)	<0.001
Renal function at M24			
Serum creatinine— $\mu\text{mol/L}$ , mean $\pm$ SD	64.8 $\pm$ 12.5	75.3 $\pm$ 19.9	0.12
eGFR— $\text{mL}/\text{min}/1.73 \text{ m}^2$ , mean $\pm$ SD	109 $\pm$ 22.1	96.6 $\pm$ 23.8	0.12
UPCR— $\text{g}/\text{g}$ , mean $\pm$ SD	0.55 $\pm$ 0.57	0.42 $\pm$ 0.36	0.73
Renal function at last follow-up			
Serum creatinine— $\mu\text{mol/L}$ , mean $\pm$ SD	73.5 $\pm$ 23.1	76.2 $\pm$ 20.8	0.31
eGFR— $\text{mL}/\text{min}/1.73 \text{ m}^2$ , mean $\pm$ SD	106.8 $\pm$ 26	96.1 $\pm$ 19.6	0.22
UPCR— $\text{g}/\text{g}$ , mean $\pm$ SD	0.56 $\pm$ 0.55	0.37 $\pm$ 0.35	0.44
Relapse of LN			
Total— <i>n</i> (%)	11 (26.2)	8 (15.7)	0.21
Early relapse <M24— <i>n</i> (%)	5 (11.9)	8 (15.7)	1
Time until relapse—months, mean $\pm$ SD	29.3 $\pm$ 19.6	12.3 $\pm$ 3.6	0.016
CKD (eGFR < 60 $\text{mL}/\text{min}/1.73 \text{ m}^2$ )			
Total, <i>n</i> (%)	6 (14.3)	6 (11.8)	0.56
New onset CKD at last follow-up	2 (4.8)	1 (2)	0.58
New onset CKD at M24	0	0	NS
Time until CKD—months, mean $\pm$ SD	27.5 $\pm$ 1.5	35	NS
ESKD			
At M24— <i>n</i> (%)	2 (4.8)	0	NS
At the end of follow up— <i>n</i> (%)	3 (7.1)	2 (3.9)	NS
Time until ESKD—months, mean $\pm$ SD	17.3 $\pm$ 8.4	23 $\pm$ 11	0.55
Death— <i>n</i> (%)			
Time until death—months mean $\pm$ SD	3 (7)	1 (2)	NS
	26.3 $\pm$ 13.1	39	NS

CKD: Chronic Kidney Disease; eGFR: estimated glomerular filtration rate; ESKD: End Stage Kidney Disease; LN: Lupus nephritis; M24: 24 months of follow-up; UPCR: urine protein to creatinine ratio.

There was no significant difference in gender, age, ethnicity or renal function between the discovery cohort and the validation cohort (Appendix A, Table A1), except for a difference in the center of origin (more patients from Paris Bichat in the discovery cohort, more patients from Marseille in the validation cohort).

### 3.2. Absence of Urinary Peptides Predicting Proliferative LN

In the discovery cohort of 42 samples, comprising 22 samples from patients with active LN and 20 samples from patients with non-active LN, we identified 88 significant peptides ( $p < 0.05$ , Wilcoxon) that differentiated active from non-active LN. However, these peptides lost significance after multiple testing correction suggesting limited discriminative power of these peptides. This was confirmed by attempts to validate these peptides in the validation cohort (comprising 33 patients with active LN and 18 with non-active LN). While a random forest model of these 88 peptides performed well in the discovery cohort (area under the curve (AUC) of 0.834), it performed poorly in the validation cohort (AUC 0.542). The list of these 88 peptides is provided in Supplementary Table S2.

Assuming that glomerular activity was a continuous dimension, we hypothesized that the limit of our analysis may be the dichotomous distinction between active and non-active LN and that urinary peptidome may be correlated to the amount of active glomerular lesions, but Spearman correlation analysis showed no peptide was significantly correlated to glomerular activity after multiple testing correction.

Routine laboratory parameters were also tested as potential discriminative markers between active LN and non-active LN (Table 3). Complement consumption, anti-dsDNA antibodies and pyuria were significantly associated to active LN ( $p = 0.001$ ,  $p = 0.007$  and  $p = 0.03$ , respectively). In non-active LN, serum creatinine was significantly lower ( $87.1 \pm 71$  versus  $96.7 \pm 71$ ;  $p = 0.01$ ) and eGFR significantly higher ( $106.7 \pm 45.8$  versus

85.9 ± 31;  $p = 0.009$ ). Anti-dsDNA antibodies offered the best sensitivity (96%) and pyuria the best specificity (67%) for active LN. UPCR and hematuria did not differ significantly between active and non-active LN.

**Table 3.** Comparison of biological parameters between patients with active and non-active lupus nephritis and discriminative power of routine SLE markers.

	Active LN	Non-Active LN	<i>p</i>	Sensitivity	Specificity	PPV	NPV
Peptidomic		No profile		NA	NA	NA	NA
Complement consumption (%)	84	46.1	0.001	0.84	0.54	0.78	0.64
Anti-DNA antibodies (%)	95.8	71.9	0.007	0.96	0.28	0.67	0.82
Hematuria (%)	53.1	40.0	0.37	0.53	0.60	0.68	0.44
Pyuria (%)	64.3	33.3	0.03	0.64	0.67	0.77	0.52
Mean eGFR (mL/min/1.73 m <sup>2</sup> )	82.7 ± 31	104.4 ± 47	0.01	NA	NA	NA	NA
Mean UPCR (g/g)	2.63 ± 2.3	1.97 ± 1.7	0.09	NA	NA	NA	NA
Mean serum creatinine (μmol/L)	96.7 ± 71	87.1 ± 71	0.01	NA	NA	NA	NA


eGFR: estimated glomerular filtration rate, calculated with the MDRD formula; NA: not applicable; NPV: negative predictive value; PPV: positive predictive value; SLE: Systemic lupus erythematosus; UPCR: Urine Protein to Creatinine Ratio.

### 3.3. The Urinary Peptidome Does Not Predict Glomerulosclerosis

Because the urine peptidome can reflect glomerular rearrangements and fibrosis [44], we hypothesized that severe glomerular sclerosis may significantly change the composition of peptides found in patients developing LN. We tried to identify a peptide signature able to predict glomerular chronicity defined by glomerulosclerosis  $\geq 25\%$  (or  $\geq 50\%$  in a second analysis). After the multiple testing correction, 177 peptides were predictive of glomerular chronicity in the discovery cohort, but none of them could be confirmed in the validation cohort. The list of these 177 peptides is provided in Supplementary Table S3.

### 3.4. CKD273 Is Correlated to Tubulo-Interstitial Chronicity

No peptide profile appeared reliable to predict tubulo-interstitial chronicity or its severity, but a significant correlation was found between the level of IF/TA and the CKD273 score ( $r = 0.3139$ ;  $p = 0.0015$ ) (Figure 1). Several clinical characteristics were strongly correlated to IF/TA (Table 4), such as SLE duration ( $r = 0.408$ ;  $p < 0.0001$ ) and LN duration ( $r = 0.450$ ;  $p < 0.0001$ ). Age was also associated to IF/TA ( $r = 0.231$ ;  $p = 0.026$ ). No correlation was found with hematuria or UPCR, but the level of IF/TA at inclusion was correlated with serum creatinine and eGFR ( $r = 0.274$ ;  $p = 0.008$  and  $r = -0.290$ ;  $p = 0.0049$ , respectively). The application of CKD273 with the established cut-off for early CKD detection [45,46] scored 82 LN patients of the whole cohort (88.2%) as a high risk group for CKD development. The eleven patients that scored negative (classification score below 0.154) had a higher mean eGFR of 112.5 (SD = 40.2) than the high risk group patients (mean eGFR = 88.6 (SD = 38.4),  $p = 0.057$ ).



**Figure 1.** Correlation between Interstitial Fibrosis/Tubular Atrophy and (A) CKD273, (B) serum creatinine at inclusion, (C) SLE duration, (D) eGFR at inclusion, (E) LN duration and (F) age. eGFR: estimate glomerular filtration rate calculated with the MDRD formula; LN: Lupus Nephritis; SLE: Systemic Lupus Erythematosus.

**Table 4.** Correlation between peptidomic and clinico-biological characteristics with the severity of IF/TA.

Parameter	<i>r</i>	<i>p</i>
Peptidomic CKD273	0.314	No profile 0.0015
SLE duration	0.408	<0.0001
LN duration	0.450	<0.0001
Age	0.231	0.026
Serum creatinine	0.274	0.008
eGFR (MDRD)	−0.290	0.005
UPCR	0.164	0.121
Hematuria	−0.031	0.783

LN: Lupus nephritis; SLE: Systemic lupus erythematosus; UPCR: Urine Protein to Creatinine Ratio.


### 3.5. LN 172 Is a Sensible Predictor for LN

The previously established LN172 classifier [35] identified 86 out of the 93 patients as patients with LN (LN172+, sensitivity 92.5%). The diagnostics threshold of 0.013 was

established using the independent validation set ( $n = 474$ ) of Siwy et al. data [35] and resulted there in a sensitivity of 79.3% and specificity of 75.1% (AUC = 0.82). Among the 7 LN patients who were not identified as LN patients by the LN172 classifier (LN172-), 6 had active LN and 1 non-active LN. The LN172- patients had lower eGFR at inclusion ( $46.2 \pm 29$  vs.  $94.9 \pm 38$  mL/min/1.73 m<sup>2</sup>;  $p = 0.001$ ) and at the end of the follow-up ( $67.5 \pm 32$  vs.  $104.8 \pm 30$  mL/min/1.73 m<sup>2</sup>;  $p = 0.004$ ), and were more likely to display baseline CKD (42.9% vs. 7%) than LN172+ patients. Their CKD273 score was significantly higher ( $p = 0.02$ ) (Appendix A, Table A2). Accordingly, the LN172 score was inversely correlated with the CKD273 score in the global cohort (Pearson correlation,  $r = -0.3$ ;  $p = 0.004$ ).

### 3.6. The Urinary Peptidome Does Not Predict Early Remission

We hypothesized that urinary peptides may predict early renal remission and thus renal response to treatment. No peptide profile was significantly predictive of renal remission at M6, whatever the UPCR threshold used to define remission (0.2, 0.5 or 0.7 g/g) (Figure 2).



**Figure 2.** Renal remission over time according to three different thresholds for proteinuria.

The relation between urinary peptidome and long term renal evolution could not be tested as only three patients developed new-onset CKD during follow-up. As such, statistical analysis could not be performed to predict renal long-term evolution.

## 4. Discussion

In this study, comprising an independent discovery and a validation cohort, no urinary peptide signature could discriminate active LN from non-active LN. Previous works aimed to discriminate LN from other causes of renal damage [35] or renal SLE from non-renal SLE [34]. To our knowledge, this is the first study using urinary peptidomics to address the specific distinction between active and non-active LN, which is crucial for the therapeutic strategy in patients with LN.

Other studies have identified significant changes in urinary proteome in the course of LN. For instance, Zhan et al. [28] highlighted the modification of the urinary low-molecular weight proteins during a flare cycle (baseline, preflare, flare, post-flare) in sequential analyses by SELDI-TOF of urinary samples from 19 patients. Though these results have not been validated in an independent cohort, this work supported the intuitive hypothesis that molecular changes occur during the different phases of LN and that

urine proteome/peptidome analysis could allow the diagnosis of severe flares. Wei et al. identified 300 peptides discriminating LN from non-renal SLE and controls [34]. The LN172 urinary peptide classifier developed by Siwy et al. [35] aimed to discriminate LN from other kidney diseases. In a cohort comprising 92 patients with LN and 1088 patients with other kidney diseases, the LN172 score correctly classified LN samples with an AUC of 0.82 and sensitivity of 79.3% and specificity of 75.1%. We confirmed in an independent cohort that, indeed, the LN172 classifier displayed a high sensitivity to detect LN (92.5%). Interestingly, the 7 patients that were not classified as LN by the LN172 classifier had lower eGFR and 3 of the 7 patients displayed CKD at baseline (eGFR < 60 mL/min/1.73 m<sup>2</sup>). In the setting of more advanced kidney disease, one can hypothesize that the urinary peptidome from these 7 patients was enriched with “regular CKD” peptides and thereby modifying its LN specific profile. This is confirmed by the fact that the LN172- patients had a higher CDK273 score, which targets chronic kidney rearrangements, and the inverse correlation between these two classifiers reinforces this hypothesis.

A possible pitfall for urinary peptidome analysis in this work, as in real life, is the baseline immunosuppressive therapy of some patients at the time of KB that could mitigate some markers of active LN. Because LN is not always diagnosed early, and 45% of patients from this study had a previous history of LN flare, our second hypothesis was that urinary peptidome analysis would reflect the severity of chronic kidney damage, either glomerular or tubulo-interstitial. Indeed, the urinary peptidome has been tested and validated through the development of the CDK273 classifier in CKD diagnosis and prediction [22], and in diabetic kidney disease (early prediction, diagnosis and CKD prediction) [47]. However, in this work, no robust “SLE-specific” peptide signature predicted glomerular chronicity or tubulo-interstitial fibrosis. The application of CKD273 for the early detection of high-risk patients like performed by Tofte et al. resulted in 88.2% high risk patients for CKD development in our cohort. Moreover, CKD273 signature was positively correlated to the level of IF/TA in patients with LN, supporting the hypothesis that peptide markers can reflect chronic TIN rearrangement. While the therapeutic guidelines focus on glomerular lesions through the ISN/RPS classification, TIN lesions, whether acute (such as tubulitis) or chronic (such as IF/TA), are strong independent prognostic factors of kidney survival in LN [9,48,49]. In particular, in a study from Hsieh et al., interstitial nephritis was found to be a better predictor of renal failure than the ISN/RPS class [50]. The association between the severity of TIN lesions and the ISN/RPS class is inconstant [50,51] which suggests that they may result from independent inflammatory pathways and thus may require specific therapeutic adjustments. Taken together, these results explain the better prognostic performance of global histologic assessments compared to the glomerulo-centric ISN/RPS classification [51,52]. The association between the CKD273 score and IF/TA encourages the development of the urinary peptidome analysis as a prognostic factor through the non-invasive assessment of TIN lesions. In our study, despite a minimum follow-up of 2 years (with a median of 4 years), we could not test the predictive value of urinary peptidome for the development of CKD, since only 3 patients developed new-onset CKD after a mean follow-up of 4 years. Moreover, it can be noted that SLE vintage and LN duration, as well as serum creatinine were better predictors of renal fibrosis than CKD273 in this cohort.

Another possible pitfall that could blunt the predictive value of urinary peptidome analysis in LN is the abundance of proteinuria. Mean UPCR was over 2 g/g in both the discovery and validation cohorts in the present study. It is possible that high proteinuria is a confounding factor on the relation between the peptidome and the clinical classification of the patients. However, among the 177 peptides which appeared correlated to glomerular chronicity in the discovery cohort, only 6 were correlated to UPCR. Similarly, among the 240 peptides identified by Wei et al. [34] as significantly increased in LN compared with non-renal SLE patients and controls, only 13 were correlated to proteinuria. In addition to proteinuria, the low number of patients used in the discovery set ( $n = 42$ ) may have limited the chance to find robust discriminatory peptides between the active and non-active form of LN.

Anti-dsDNA antibodies, complement consumption and pyuria were significantly associated to active LN, but with poor diagnostic performances. A combination of 12 clinical and biological markers has been previously used to predict active LN [53], with a ROC curve AUC of 0.83, offering a fine orientation but still insufficient to replace KB.

The poor correlation of molecular markers and renal pathological results described here has also been reported within the renal tissue, in transcriptomic analysis of dissected glomeruli and tubules from KB of patients with LN [54,55]. Single-cell RNA sequencing from KB of patients with LN and controls [56] has shown specific gene expression profiles associated with LN, strongly correlated to the profile found in urinary cells. Similarly, Abedini et al. recently demonstrated that the single-cell transcriptome analysis could detect and quantify almost all kidney cells in urinary samples from patients with diabetic kidney disease (DKD) and controls and that the distribution of the detected cells was different between DKD and controls [57]. These results suggest a possible use of urinary cells transcriptomics to identify immunological pathways activated within the kidney and assess non-invasively infra-clinical rearrangements occurring in the course of CKD.

Although urinary peptidomics did not differentiate between active LN and non-active LN patients in this cohort, there are other potential applications of urinary peptidomics in LN that were not explored in this study. First, early detection of LN flare through the detection of early renal rearrangements could be investigated using urinary peptidomics. Some immunological parameters are associated with the risk of LN in SLE [58–60] but remain poorly predictive of flares [61,62]. Moreover, some patients show active renal lesions while no proteinuria, hematuria or pyuria is detected [63,64]. Here, all patients were sampled while clinically flaring and had a renal biopsy because of proteinuria. Second, as insufficient data on CKD and ESKD were available, their relation to the value of urinary peptidomics also remains untested.

## 5. Conclusions

We were not able to identify a urinary peptide profile predictive of active LN, of glomerular activity or chronicity in this cohort of patients with LN. However, the LN172 signature displayed high sensitivity for the diagnosis of LN while the CKD273 classifier was correlated to IF/TA which paves the way to the development of a new prognostic tool in the care of patient with LN. Whether urinary peptidomics will allow the early detection of renal flares in SLE remains to be determined. To date, kidney biopsy remains the cornerstone of LN management and could provide useful information on the molecular pathways activated at a given time, in addition to the pathological lesions observed [7]. Nonetheless, non-invasive biomarkers such as the urinary peptidome are still forecasted to prove their added-value for a more personalized care of patients with LN, yet their development and validation will require larger cohorts.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/article/10.3390/jcm10081690/s1>. Table S1: raw peptides data from the discovery and the validation cohorts; Table S2: list of the 88 peptides significantly different between samples with active LN and samples with inactive LN in the discovery cohort; Table S3: list of the 177 peptides significantly different between samples with  $\geq 25\%$  sclerotic glomeruli and samples with  $<25\%$  sclerotic glomeruli in the discovery cohort.

**Author Contributions:** Conceptualization, S.F., E.D. and N.J.-C.; Data curation, T.D., A.V. and L.D.; Formal analysis, M.T., J.P.S. and T.D.; Funding acquisition, N.J.-C.; Methodology, J.P.S.; Project administration, J.P.S., E.D. and N.J.-C.; Software, T.D.; Supervision, N.J.-C.; Validation, B.B., G.H., N.C., J.-L.B. and B.D.; Visualization, M.T.; Writing—original draft, M.T.; Writing—review & editing, J.P.S., L.C., J.S., S.F., L.D., E.D. and N.J.-C. All authors have read and agreed to the published version of the manuscript.

**Funding:** The Peptidu-LUP study was funded by a grant from the “Fondation pour la Recherche Médicale” (FRM) and the “Fondation du Rein” (FDR) attributed by the Société Française de Néphrologie

Dialyse et Transplantation (SFNDT). J.P.S. was supported by a grant from the “Fondation pour la Recherche Médicale” (Grant No. DEQ20170336759).

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki. The samples were included in the biological collections of the 3 centers participating in the study (Marseille: DC 2012-1704, approval date: 4 February 2013; Paris Bichat: ID-RCB 2014-A00809-38 approval date 21 May 2014; and Toulouse: DC2011-1388, approval date 1 January 2011).

**Informed Consent Statement:** All patients signed an informed consent before any study procedure.

**Data Availability Statement:** The authors confirm that the data supporting the findings of this study are available within the article and its Supplementary Materials. Complementary data are available on request from the corresponding author, [N.J.C].

**Acknowledgments:** We thank Laurent Samson and Julien Faraut for their help with the samples. Members of the Groupe Coopératif sur le Lupus Rénal (GCLR) who also contributed to this work: Lafita Azeroual; Julie Belliere; Laure Champion; Dominique Chauveau; Jonathan Chemouny; Mathieu Jablonski; Bénédicte Levy; Jacques Pourrat; Quentin Raimbourg; Florence Vende; Marion Venot.

**Conflicts of Interest:** J.S. is employed by Mosaiques Diagnostics GmbH.

## Appendix A

**Table A1.** Baseline characteristics of patients from the discovery and validation cohorts.

	Discovery Cohort	Validation Cohort	<i>p</i>
Patients, <i>n</i>	42	51	
Women— <i>n</i> (%)	40 (95.2)	46 (90.2)	0.45
Mean age ± SD	34 ± 7.7	36 ± 10	0.37
Ethnicity— <i>n</i> (%)			0.66
European	16 (38.1)	24 (47.1)	0.38
North African	10 (23.8)	13 (25.5)	1
African	11 (26.2)	11 (21.6)	0.78
Asian	5 (11.9)	3 (5.9)	0.46
Characteristics at kidney biopsy			
Serum creatinine—μmol/L, mean ± SD	99.4 ± 55	86.8 ± 32	0.82
eGFR—mL/min/1.73 m <sup>2</sup> , mean ± SD	90.4 ± 32	92.3 ± 28.6	0.81
UPCR—g/g, mean ± SD	2.12 ± 1.25	2.53 ± 1.7	0.74
SLE vintage			
First flare of SLE— <i>n</i> (%)	10 (23.8)	11 (21.6)	0.99
Pre-existing SLE— <i>n</i> (%)	32 (76.2)	40 (78.4)	0.99
Disease duration—years, mean ± SD	9.8 ± 7	11.33 ± 9.6	0.78
First renal flare— <i>n</i> (%)	22 (52.4)	33 (64.7)	0.32
LN duration—years, mean ± SD	6.6 ± 5	10.5 ± 8.2	0.22
Kidney biopsies, <i>n</i>			
Class, <i>n</i> (%)			0.15
I	0 (0)	1 (2.0)	1
II	4 (9.5)	2 (3.9)	0.40
III-A or -A/C ± V	12 (28.6)	10 (19.6)	0.59
III-C ± V	2 (4.8)	3 (5.9)	1
IV-A or -A/C ± V	10 (23.8)	23 (45.1)	0.06
IV-C ± V	1 (2.4)	1 (2.0)	0.45
Pure V	12 (28.6)	9 (17.6)	0.32
VI	1 (2.4)	2 (3.9)	1
IF/TA, <i>n</i> —(%)			0.70
F0	20 (44.6)	21 (41.2)	0.68
F1	14 (33.3)	19 (37.3)	0.86

Table A1. Cont.

	Discovery Cohort	Validation Cohort	<i>p</i>
F2	4 (9.5)	8 (15.7)	0.54
F3	4 (9.5)	3 (5.9)	0.7
Group			
Active LN	22(52.4)	33 (64.7)	0.36
Non-active LN	20 (47.6)	18 (35.3)	0.36
Patients/center— <i>n</i> (%)			
Marseille	7 (16.7)	30 (58.8)	<0.001
Paris	35 (83.3)	18 (35.3)	<0.001
Toulouse	0 (0)	3 (5.9)	0.25
Previous treatment for SLE—%			
Hydroxychloroquine	78.6	82.4	0.84
Corticosteroids	80.1	74.5	0.62
MMF/Mycophenolic acid	30.9	21.6	0.31
Azathioprine	16.7	21.6	0.74
Cyclophosphamide	35.7	31.4	0.82
Rituximab	4.8	11.8	0.29
Other	14.3	13.7	1
Treatment for SLE at inclusion—%			
Hydroxychloroquine	66.7	70.6	0.86
Corticosteroids	76.2	66.7	0.44
MMF/Mycophenolic acid	21.4	13.7	0.48
Azathioprine	14.3	5.9	0.29
Cyclophosphamide	2.4	0	0.45
Rituximab	0	2	1
Methotrexate	2.4	2	1

Table A2. Characteristics of patients according to their classification with the LN172 score.

	LN172+	LN172–	<i>p</i>
Patients, <i>n</i> (%)	86 (92.5)	7 (7.5)	
Women— <i>n</i> (%)	80 (93)	6 (85.7)	0.43
Mean age ± SD	35 ± 11	36 ± 10	0.80
Ethnicity— <i>n</i> (%)			0.07
European	35 (40.7)	5 (71.4)	0.13
North African	22 (25.6)	1 (14.3)	0.67
African	21 (24.4)	1 (14.3)	1
Asian	8 (9.3)	0 (0)	1
Characteristics of kidney biopsy			
Serum creatinine— $\mu\text{mol/L}$ , mean ± SD	80 ± 61	184.6 ± 114	0.003
eGFR— $\text{mL/min/1.73 m}^2$ , mean ± SD	94.9 ± 38	46.2 ± 29	0.001
UPCR— $\text{g/g}$ , mean ± SD	2.29 ± 2.03	2.95 ± 2.48	0.79
CKD— <i>n</i> (%)	6 (7.0)	3 (42.9)	0.01
Complement consumption (%)	71.4	66.7	1
Anti-DNA antibodies (%)	86.3	85.7	1
CKD273 score	0.56 ± 0.35	0.86 ± 0.13	0.02
SLE vintage			
First flare of SLE— <i>n</i> (%)	20 (23.3)	1 (14.3)	1
Pre-existing SLE— <i>n</i> (%)	66 (76.7)	6 (85.7)	1
Disease duration—years, mean ± SD	10.5 ± 8.6	11.7 ± 7.5	0.56
First renal flare— <i>n</i> (%)	53 (61.6)	2 (28.6)	0.11
LN duration—years, mean ± SD	8.0 ± 6.9	9.8 ± 7.8	0.58

Table A2. Cont.

	LN172+	LN172−	<i>p</i>
Kidney biopsies, <i>n</i>			
Class, <i>n</i> (%)			0.37
I	1 (1.2)	0 (0)	1
II	6 (7.0)	0 (0)	1
III-A or -A/C ± V	19 (22.1)	3 (42.9)	0.36
III-C ± V	5 (5.8)	0 (0)	1
IV-A or -A/C ± V	30 (34.9)	3 (42.9)	0.70
IV-C ± V	2 (1.2)	0 (0)	1
V	21 (24.4)	0 (0)	0.34
VI	2 (2.3)	1 (1.2)	0.21
IF/TA, <i>n</i> —(%)			0.16
F0	40 (46.5)	1 (14.3)	0.13
F1	30 (34.9)	3 (42.9)	0.70
F2	10 (11.6)	2 (28.6)	0.22
F3	6 (7.0)	1 (14.3)	0.43
Group			
Active LN	49 (57.0)	6 (85.7)	0.23
Non-active LN	37 (43.0)	1 (14.3)	0.23
Patients/center— <i>n</i> (%)			0.18
Marseille	50 (58.1)	3 (42.9)	0.46
Paris	34 (39.5)	3 (42.9)	1
Toulouse	2 (2.3)	1 (14.3)	0.21
Previous treatment for SLE—%			
Hydroxychloroquine	81.2	100	0.59
Corticosteroids	78.8	83.3	1
MMF/Mycophenolic acid	23.8	66.7	0.04
Azathioprine	17.9	50	0.09
Cyclophosphamide	32.1	66.7	0.18
Rituximab	7.1	33.3	0.09
Other	11.8	50	0.04
Treatment for SLE at inclusion—%			
Hydroxychloroquine	69.4	83.3	0.67
Corticosteroids	71.8	83.3	1
MMF/Mycophenolic acid	16.7	33.3	0.29
Azathioprine	9.5	16.7	0.48
Cyclophosphamide	0	16.7	0.07
Rituximab	1.2	0	1
Methotrexate	2.4	0	1
Relapse of LN			
Total— <i>n</i> (%)	19 (22.1)	0 (0)	0.33
Renal function at last follow-up			
Serum creatinine— $\mu\text{mol/L}$ , mean $\pm$ SD	70.7 $\pm$ 32.7	128.7 $\pm$ 109.8	0.064
eGFR— $\text{mL}/\text{min}/1.73 \text{ m}^2$ , mean $\pm$ SD	104.8 $\pm$ 29.9	67.5 $\pm$ 32	0.004
UPCR— $\text{g}/\text{g}$ , mean $\pm$ SD	0.42 $\pm$ 0.55	0.68 $\pm$ 1.52	0.17

## References

1. Cervera, R. Systemic lupus erythematosus in Europe at the change of the millennium: Lessons from the “Euro-Lupus Project”. *Autoimmun. Rev.* **2006**, *5*, 180–186. [[CrossRef](#)]
2. Cervera, R.; Khamashta, M.A.; Font, J.; Sebastiani, G.D.; Gil, A.; Lavilla, P.; Mejía, J.C.; Aydintug, A.O.; Chwalinska-Sadowska, H.; de Ramón, E.; et al. Morbidity and Mortality in Systemic Lupus Erythematosus During a 10-Year Period. *Medicine* **2003**, *82*, 299–308. [[CrossRef](#)]
3. Houssiau, F.A.; Vasconcelos, C.; D’Cruz, D.; Sebastiani, G.D.; Garrido, E.D.R.; Danieli, M.G.; Abramovicz, D.; Blockmans, D.; Cauli, A.; Direskeneli, H.; et al. The 10-year follow-up data of the Euro-Lupus Nephritis Trial comparing low-dose and high-dose intravenous cyclophosphamide. *Ann. Rheum. Dis.* **2009**, *69*, 61–64. [[CrossRef](#)] [[PubMed](#)]

4. Bajema, I.M.; Wilhelmus, S.; Alpers, C.E.; Bruijn, J.A.; Colvin, R.B.; Cook, H.T.; D'Agati, V.D.; Ferrario, F.; Haas, M.; Jennette, J.C.; et al. Revision of the International Society of Nephrology/Renal Pathology Society classification for lupus nephritis: Clarification of definitions, and modified National Institutes of Health activity and chronicity indices. *Kidney Int.* **2018**, *93*, 789–796. [[CrossRef](#)] [[PubMed](#)]
5. Weening, J.J.; D'Agati, V.D.; Schwartz, M.M.; Seshan, S.V.; Alpers, C.E.; Appel, G.B.; Balow, J.E.; Bruijn, J.A.; Cook, T.; Ferrario, F.; et al. The classification of glomerulonephritis in systemic lupus erythematosus revisited. *Kidney Int.* **2004**, *65*, 521–530. [[CrossRef](#)] [[PubMed](#)]
6. Bertsias, G.K.; Tektonidou, M.; Amoura, Z.; Aringer, M.; Bajema, I.; Berden, J.H.M.; Boletis, J.; Cervera, R.; Doerner, T.; Doria, A.; et al. Joint European League Against Rheumatism and European Renal Association–European Dialysis and Transplant Association (EULAR/ERA-EDTA) recommendations for the management of adult and paediatric lupus nephritis. *Ann. Rheum. Dis.* **2012**, *71*, 1771–1782. [[CrossRef](#)]
7. Dörner, T.; Furie, R. Novel paradigms in systemic lupus erythematosus. *Lancet* **2019**, *393*, 2344–2358. [[CrossRef](#)]
8. Hong, S.; Healy, H.; Kassianos, A.J. The Emerging Role of Renal Tubular Epithelial Cells in the Immunological Pathophysiology of Lupus Nephritis. *Front. Immunol.* **2020**, *11*, 578952. [[CrossRef](#)]
9. Wilson, P.C.; Kashgarian, M.; Moeckel, G. Interstitial inflammation and interstitial fibrosis and tubular atrophy predict renal survival in lupus nephritis. *Clin. Kidney J.* **2018**, *11*, 207–218. [[CrossRef](#)]
10. Espeli, M.; Bökers, S.; Giannico, G.; Dickinson, H.A.; Bardsley, V.; Fogo, A.B.; Smith, K.G. Local Renal Autoantibody Production in Lupus Nephritis. *J. Am. Soc. Nephrol.* **2010**, *22*, 296–305. [[CrossRef](#)]
11. Narváez, J.; Ricse, M.; Gomà, M.; Mitjavila, F.; Fulladosa, X.; Capdevila, O.; Torras, J.; Juanola, X.; Pujol-Farriols, R.; Nolla, J.M. The value of repeat biopsy in lupus nephritis flares. *Medicine* **2017**, *96*, e7099. [[CrossRef](#)]
12. Bataille, S.; Jourde, N.; Daniel, L.; Mondain, J.-R.; Faure, M.; Gobert, P.; Alcheikh-Hassan, Z.; Lankester, M.; Giaime, P.; Gaudart, J.; et al. Comparative Safety and Efficiency of Five Percutaneous Kidney Biopsy Approaches of Native Kidneys: A Multicenter Study. *Am. J. Nephrol.* **2012**, *35*, 387–393. [[CrossRef](#)]
13. Soliman, S.; Mohan, C. Lupus nephritis biomarkers. *Clin. Immunol.* **2017**, *185*, 10–20. [[CrossRef](#)]
14. Gulati, G.; Bennett, M.R.; Abulaban, K.; Song, H.; Zhang, X.; Ma, Q.; Brodsky, S.V.; Nadasdy, T.; Haffner, C.; Wiley, K.; et al. Prospective validation of a novel renal activity index of lupus nephritis. *Lupus* **2016**, *26*, 927–936. [[CrossRef](#)]
15. Kitagawa, A.; Tsuboi, N.; Yokoe, Y.; Katsuno, T.; Ikeuchi, H.; Kajiyama, H.; Endo, N.; Sawa, Y.; Suwa, J.; Sugiyama, Y.; et al. Urinary levels of the leukocyte surface molecule CD11b associate with glomerular inflammation in lupus nephritis. *Kidney Int.* **2019**, *95*, 680–692. [[CrossRef](#)]
16. Ding, Y.; Nie, L.-M.; Pang, Y.; Wu, W.-J.; Tan, Y.; Yu, F.; Zhao, M.-H. Composite urinary biomarkers to predict pathological tubulointerstitial lesions in lupus nephritis. *Lupus* **2018**, *27*, 1778–1789. [[CrossRef](#)] [[PubMed](#)]
17. Klein, J.; Bascands, J.-L.; Mischak, H.; Schanstra, J.P. The role of urinary peptidomics in kidney disease research. *Kidney Int.* **2016**, *89*, 539–545. [[CrossRef](#)] [[PubMed](#)]
18. Klein, J.; Lacroix, C.; Caubet, C.; Siwy, J.; Zürbig, P.; Dakna, M.; Muller, F.; Breuil, B.; Stalmach, A.; Mullen, W.; et al. Fetal Urinary Peptides to Predict Postnatal Outcome of Renal Disease in Fetuses with Posterior Urethral Valves (PUV). *Sci. Transl. Med.* **2013**, *5*, 198ra106. [[CrossRef](#)] [[PubMed](#)]
19. Klein, J.; Buffin-Meyer, B.; Boizard, F.; Moussaoui, N.; Lescat, O.; Breuil, B.; Fedou, C.; Feuillet, G.; Casemayou, A.; Neau, E.; et al. Amniotic fluid peptides predict postnatal kidney survival in developmental kidney disease. *Kidney Int.* **2021**, *99*, 737–749. [[CrossRef](#)]
20. Desveaux, C.; Klein, J.; Leruez-Ville, M.; Ramirez-Torres, A.; Lacroix, C.; Breuil, B.; Froment, C.; Bascands, J.-L.; Schanstra, J.P.; Ville, Y. Identification of Symptomatic Fetuses Infected with Cytomegalovirus Using Amniotic Fluid Peptide Biomarkers. *PLoS Pathog.* **2016**, *12*, e1005395. [[CrossRef](#)] [[PubMed](#)]
21. Good, D.M.; Zürbig, P.; Argilés, À.; Bauer, H.W.; Behrens, G.; Coon, J.J.; Dakna, M.; Decramer, S.; Delles, C.; Dominiczak, A.F.; et al. Naturally Occurring Human Urinary Peptides for Use in Diagnosis of Chronic Kidney Disease. *Mol. Cell. Proteom.* **2010**, *9*, 2424–2437. [[CrossRef](#)] [[PubMed](#)]
22. Pontillo, C.; Zhang, Z.-Y.; Schanstra, J.P.; Jacobs, L.; Zürbig, P.; Thijs, L.; Ramírez-Torres, A.; Heerspink, H.J.; Lindhardt, M.; Klein, R.; et al. Prediction of Chronic Kidney Disease Stage 3 by CKD273, a Urinary Proteomic Biomarker. *Kidney Int. Rep.* **2017**, *2*, 1066–1075. [[CrossRef](#)] [[PubMed](#)]
23. Schanstra, J.P.; Zürbig, P.; Alkhalaf, A.; Argiles, A.; Bakker, S.J.; Beige, J.; Bilo, H.J.; Chatzikyrkou, C.; Dakna, M.; Dawson, J.; et al. Diagnosis and Prediction of CKD Progression by Assessment of Urinary Peptides. *J. Am. Soc. Nephrol.* **2015**, *26*, 1999–2010. [[CrossRef](#)] [[PubMed](#)]
24. Haubitz, M.; Wittke, S.; Weissinger, E.M.; Walden, M.; Rupperecht, H.D.; Floege, J.; Haller, H.; Mischak, H. Urine protein patterns can serve as diagnostic tools in patients with IgA nephropathy. *Kidney Int.* **2005**, *67*, 2313–2320. [[CrossRef](#)] [[PubMed](#)]
25. Wittke, S.; Mischak, H.; Walden, M.; Kolch, W.; Rädler, T.; Wiedemann, K. Discovery of biomarkers in human urine and cerebrospinal fluid by capillary electrophoresis coupled to mass spectrometry: Towards new diagnostic and therapeutic approaches. *Electrophoresis* **2005**, *26*, 1476–1487. [[CrossRef](#)]
26. Weissinger, E.M.; Wittke, S.; Kaiser, T.; Haller, H.; Bartel, S.; Krebs, R.; Golovko, I.; Rupperecht, H.D.; Haubitz, M.; Hecker, H.; et al. Proteomic patterns established with capillary electrophoresis and mass spectrometry for diagnostic purposes. *Kidney Int.* **2004**, *65*, 2426–2434. [[CrossRef](#)]

27. Varghese, S.A.; Powell, T.B.; Budisavljevic, M.N.; Oates, J.C.; Raymond, J.R.; Almeida, J.S.; Arthur, J.M. Urine Biomarkers Predict the Cause of Glomerular Disease. *J. Am. Soc. Nephrol.* **2007**, *18*, 913–922. [[CrossRef](#)]
28. Zhang, X.; Jin, M.; Wu, H.; Nadasdy, T.; Nadasdy, G.; Harris, N.; Green-Church, K.; Nagaraja, H.; Birmingham, D.J.; Yu, C.-Y.; et al. Biomarkers of lupus nephritis determined by serial urine proteomics. *Kidney Int.* **2008**, *74*, 799–807. [[CrossRef](#)]
29. Suzuki, M.; Ross, G.F.; Wiers, K.; Nelson, S.; Bennett, M.; Passo, M.H.; Devarajan, P.; Brunner, H.I. Identification of a urinary proteomic signature for lupus nephritis in children. *Pediatr. Nephrol.* **2007**, *22*, 2047–2057. [[CrossRef](#)]
30. Suzuki, M.; Wiers, K.; Brooks, E.B.; Greis, K.D.; Haines, K.; Klein-Gitelman, M.S.; Olson, J.; Onel, K.; O’Neil, K.M.; Silverman, E.D.; et al. Initial Validation of a Novel Protein Biomarker Panel for Active Pediatric Lupus Nephritis. *Pediatr. Res.* **2009**, *65*, 530–536. [[CrossRef](#)]
31. Oates, J.; Varghese, S.; Bland, A.M.; Taylor, T.P.; Self, S.E.; Stanislaus, R.; Almeida, J.S.; Arthur, J.M. Prediction of urinary protein markers in lupus nephritis. *Kidney Int.* **2005**, *68*, 2588–2592. [[CrossRef](#)]
32. Mosley, K.; Tam, F.W.K.; Edwards, R.J.; Crozier, J.; Pusey, C.D.; Lightstone, L. Urinary proteomic profiles distinguish between active and inactive lupus nephritis. *Rheumatology* **2006**, *45*, 1497–1504. [[CrossRef](#)] [[PubMed](#)]
33. Pejchinovski, M.; Siwy, J.; Mullen, W.; Mischak, H.; Petri, M.A.; Burkly, L.C.; Wei, R. Urine peptidomic biomarkers for diagnosis of patients with systematic lupus erythematosus. *Lupus* **2018**, *27*, 6–16. [[CrossRef](#)]
34. Wei, R.; Gao, B.; Shih, F.; Ranger, A.; Dearth, A.; Mischak, H.; Siwy, J.; Wisniacki, N.; Petri, M.; Burkly, L.C. Alterations in urinary collagen peptides in lupus nephritis subjects correlate with renal dysfunction and renal histopathology. *Nephrol. Dial. Transplant.* **2017**, *32*, 1468–1477. [[CrossRef](#)] [[PubMed](#)]
35. Siwy, J.; Zürgbig, P.; Argiles, A.; Beige, J.; Haubitz, M.; Jankowski, J.; Julian, B.A.; Linde, P.G.; Marx, D.; Mischak, H.; et al. Noninvasive diagnosis of chronic kidney diseases using urinary proteome analysis. *Nephrol. Dial. Transplant.* **2016**, *32*, 2079–2089. [[CrossRef](#)]
36. Hochberg, M.C. Updating the American college of rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum.* **1997**, *40*, 1725. [[CrossRef](#)] [[PubMed](#)]
37. Petri, M.; Orbai, A.-M.; Alarcón, G.S.; Gordon, C.; Merrill, J.T.; Fortin, P.R.; Bruce, I.N.; Isenberg, D.; Wallace, D.J.; Nived, O.; et al. Derivation and validation of the Systemic Lupus International Collaborating Clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum.* **2012**, *64*, 2677–2686. [[CrossRef](#)] [[PubMed](#)]
38. Levey, A.S.; Coresh, J.; Greene, T.; Stevens, L.A.; Zhang, Y.L.; Hendriksen, S.; Kusek, J.W.; Van Lente, F. Using Standardized Serum Creatinine Values in the Modification of Diet in Renal Disease Study Equation for Estimating Glomerular Filtration Rate. *Ann. Intern. Med.* **2006**, *145*, 247–254. [[CrossRef](#)]
39. Haas, M. The Revised (2013) Banff Classification for Antibody-Mediated Rejection of Renal Allografts: Update, Difficulties, and Future Considerations. *Arab. Archaeol. Epigr.* **2016**, *16*, 1352–1357. [[CrossRef](#)]
40. Theodorescu, D.; Wittke, S.; Ross, M.M.; Walden, M.; Conaway, M.; Just, I.; Mischak, H.; Frierson, H.F. Discovery and validation of new protein biomarkers for urothelial cancer: A prospective analysis. *Lancet Oncol.* **2006**, *7*, 230–240. [[CrossRef](#)]
41. Mischak, H.; Vlahou, A.; Ioannidis, J.P. Technical aspects and inter-laboratory variability in native peptide profiling: The CE-MS experience. *Clin. Biochem.* **2013**, *46*, 432–443. [[CrossRef](#)]
42. Jantos-Siwy, J.; Schiffer, E.; Brand, K.; Schumann, G.; Rossing, K.; Delles, C.; Mischak, H.; Metzger, J. Quantitative Urinary Proteome Analysis for Biomarker Evaluation in Chronic Kidney Disease. *J. Proteome Res.* **2009**, *8*, 268–281. [[CrossRef](#)]
43. Pontillo, C.; Mischak, H. Urinary peptide-based classifier CKD273: Towards clinical application in chronic kidney disease. *Clin. Kidney J.* **2017**, *10*, 192–201. [[CrossRef](#)]
44. Magalhães, P.; Pejchinovski, M.; Markoska, K.; Banasik, M.; Klinger, M.; Švec-Billá, D.; Rychlík, I.; Rroji, M.; Restivo, A.; Capasso, G.; et al. Association of kidney fibrosis with urinary peptides: A path towards non-invasive liquid biopsies? *Sci. Rep.* **2017**, *7*, 1–8. [[CrossRef](#)] [[PubMed](#)]
45. Tofte, N.; Lindhardt, M.; Adamova, K.; Bakker, S.J.L.; Beige, J.; Beulens, J.W.J.; Birkenfeld, A.L.; Currie, G.; Delles, C.; Dimos, I.; et al. Early detection of diabetic kidney disease by urinary proteomics and subsequent intervention with spironolactone to delay progression (PRIORITY): A prospective observational study and embedded randomised placebo-controlled trial. *Lancet Diabetes Endocrinol.* **2020**, *8*, 301–312. [[CrossRef](#)]
46. Lindhardt, M.; Persson, F.; Currie, G.; Pontillo, C.; Beige, J.; Delles, C.; Von Der Leyen, H.; Mischak, H.; Navis, G.; Noutsou, M.; et al. Proteomic prediction and Renin angiotensin aldosterone system Inhibition prevention of early diabetic nephropathy in Type 2 diabetic patients with normoalbuminuria (PRIORITY): Essential study design and rationale of a randomised clinical multicentre trial. *BMJ Open* **2016**, *6*, e010310. [[CrossRef](#)] [[PubMed](#)]
47. Lindhardt, M.; Persson, F.; Zürgbig, P.; Stalmach, A.; Mischak, H.; De Zeeuw, D.; Heerspink, H.L.; Klein, R.; Orchard, T.; Porta, M.; et al. Urinary proteomics predict onset of microalbuminuria in normoalbuminuric type 2 diabetic patients, a sub-study of the DIRECT-Protect 2 study. *Nephrol. Dial. Transplant.* **2016**, *32*, 1866–1873. [[CrossRef](#)]
48. Rijnink, E.C.; Teng, Y.O.; Wilhelmus, S.; Almekinders, M.; Wolterbeek, R.; Cransberg, K.; Bruijn, J.A.; Bajema, I.M. Clinical and Histopathologic Characteristics Associated with Renal Outcomes in Lupus Nephritis. *Clin. J. Am. Soc. Nephrol.* **2017**, *12*, 734–743. [[CrossRef](#)] [[PubMed](#)]
49. Obrisca, B.; Jurubița, R.; Andronesi, A.; Sorohan, B.; Achim, C.; Bobeica, R.; Gherghiceanu, M.; Mandache, E.; Ismail, G. Histological predictors of renal outcome in lupus nephritis: The importance of tubulointerstitial lesions and scoring of glomerular lesions. *Lupus* **2018**, *27*, 1455–1463. [[CrossRef](#)]

50. Hsieh, C.; Chang, A.; Brandt, D.; Guttikonda, R.; Utset, T.O.; Clark, M.R. Predicting outcomes of lupus nephritis with tubulointerstitial inflammation and scarring. *Arthritis Rheum.* **2011**, *63*, 865–874. [[CrossRef](#)]
51. Leatherwood, C.; Speyer, C.B.; Feldman, C.H.; D’Silva, K.; Gómez-Puerta, J.A.; Hoover, P.J.; Waikar, S.S.; McMahon, G.M.; Rennke, H.G.; Costenbader, K.H. Clinical characteristics and renal prognosis associated with interstitial fibrosis and tubular atrophy (IFTA) and vascular injury in lupus nephritis biopsies. *Semin. Arthritis Rheum.* **2019**, *49*, 396–404. [[CrossRef](#)]
52. Pagni, F.; Galimberti, S.; Galbiati, E.; Rebori, P.; Pietropaolo, V.; Pieruzzi, F.; Smith, A.J.; Ferrario, F. Tubulointerstitial lesions in lupus nephritis: International multicentre study in a large cohort of patients with repeat biopsy. *Nephrology* **2016**, *21*, 35–45. [[CrossRef](#)] [[PubMed](#)]
53. Mavragani, C.P.; Fragoulis, G.E.; Somarakis, G.; Drosos, A.; Tzioufas, A.G.; Moutsopoulos, H.M. Clinical and Laboratory Predictors of Distinct Histopathological Features of Lupus Nephritis. *Medicine* **2015**, *94*, e829. [[CrossRef](#)] [[PubMed](#)]
54. Mejia-Vilet, J.M.; Parikh, S.V.; Song, H.; Fadda, P.; Shapiro, J.P.; Ayoub, I.; Yu, L.; Zhang, J.; Uribe-Uribe, N.; Rovin, B.H. Immune gene expression in kidney biopsies of lupus nephritis patients at diagnosis and at renal flare. *Nephrol. Dial. Transplant.* **2019**, *34*, 1197–1206. [[CrossRef](#)] [[PubMed](#)]
55. Almaani, S.; Prokopec, S.D.; Zhang, J.; Yu, L.; Avila-Casado, C.; Wither, J.; Scholey, J.W.; Alberton, V.; Malvar, A.; Parikh, S.V.; et al. Rethinking Lupus Nephritis Classification on a Molecular Level. *J. Clin. Med.* **2019**, *8*, 1524. [[CrossRef](#)] [[PubMed](#)]
56. Arazi, A.; Rao, D.A.; Berthier, C.C.; Davidson, A.; Liu, Y.; Hoover, P.J.; Chicoine, A.; Eisenhaure, T.M.; Jonsson, A.H.; Li, S.; et al. The immune cell landscape in kidneys of patients with lupus nephritis. *Nat. Immunol.* **2019**, *20*, 902–914. [[CrossRef](#)]
57. Abedini, A.; Zhu, Y.O.; Chatterjee, S.; Halasz, G.; Devalaraja-Narashimha, K.; Shrestha, R.; Balzer, M.S.; Park, J.; Zhou, T.; Ma, Z.; et al. Urinary Single-Cell Profiling Captures the Cellular Diversity of the Kidney. *J. Am. Soc. Nephrol.* **2021**, *32*, 614–627. [[CrossRef](#)] [[PubMed](#)]
58. Birmingham, D.J.; Irshaid, F.; Nagaraja, H.N.; Zou, X.; Tsao, B.P.; Wu, H.; Yu, C.Y.; Hebert, L.A.; Rovin, B.H. The complex nature of serum C3 and C4 as biomarkers of lupus renal flare. *Lupus* **2010**, *19*, 1272–1280. [[CrossRef](#)]
59. Mok, C.C.; Ying, K.Y.; Tang, S.; Leung, C.Y.; Lee, K.W.; Ng, W.L.; Wong, R.W.S.; Lau, C.S. Predictors and outcome of renal flares after successful cyclophosphamide treatment for diffuse proliferative lupus glomerulonephritis. *Arthritis Rheum.* **2004**, *50*, 2559–2568. [[CrossRef](#)]
60. Linnik, M.D.; Hu, J.Z.; Heilbrunn, K.R.; Strand, V.; Hurley, F.L.; Joh, T. LJP 394 Investigator Consortium Relationship between anti-double-stranded DNA antibodies and exacerbation of renal disease in patients with systemic lupus erythematosus. *Arthritis Rheum.* **2005**, *52*, 1129–1137. [[CrossRef](#)]
61. Steiman, A.J.; Gladman, D.D.; Ibañez, M.; Urowitz, M.B. Prolonged Serologically Active Clinically Quiescent Systemic Lupus Erythematosus: Frequency and Outcome. *J. Rheumatol.* **2010**, *37*, 1822–1827. [[CrossRef](#)] [[PubMed](#)]
62. Gensous, N.; Marti, A.; Barnette, T.; Blanco, P.; Lazaro, E.; Seneschal, J.; Truchetet, M.-E.; Duffau, P.; Richez, C. Predictive biological markers of systemic lupus erythematosus flares: A systematic literature review. *Arthritis Res.* **2017**, *19*, 238. [[CrossRef](#)] [[PubMed](#)]
63. De Rosa, M.; Azzato, F.; Toblli, J.E.; De Rosa, G.; Fuentes, F.; Nagaraja, H.N.; Nash, R.; Rovin, B.H. A prospective observational cohort study highlights kidney biopsy findings of lupus nephritis patients in remission who flare following withdrawal of maintenance therapy. *Kidney Int.* **2018**, *94*, 788–794. [[CrossRef](#)] [[PubMed](#)]
64. Piñeiro, G.J.; Arrizabalaga, P.; Solé, M.; Abellana, R.M.; Espinosa, G.; Cervera, R. Repeated Renal Biopsy—A Predictive Tool to Assess the Probability of Renal Flare in Lupus Nephritis. *Am. J. Nephrol.* **2016**, *44*, 439–446. [[CrossRef](#)]