



**HAL**  
open science

## On homology torsion growth

Miklos Abert, Nicolas Bergeron, Mikolaj Fraczyk, Damien Gaboriau

► **To cite this version:**

Miklos Abert, Nicolas Bergeron, Mikolaj Fraczyk, Damien Gaboriau. On homology torsion growth. Journal of the European Mathematical Society, In press. hal-03270482v1

**HAL Id: hal-03270482**

**<https://hal.science/hal-03270482v1>**

Submitted on 24 Jun 2021 (v1), last revised 27 Jun 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON HOMOLOGY TORSION GROWTH

MIKLOS ABERT, NICOLAS BERGERON, MIKOLAJ FRACZYK, AND DAMIEN GABORIAU

ABSTRACT. We prove new vanishing results on the growth of higher torsion homologies for suitable arithmetic lattices, Artin groups and mapping class groups. The growth is understood along Farber sequences, in particular, along residual chains. For principal congruence subgroups, we also obtain strong asymptotic bounds for the torsion growth.

As a central tool, we introduce a quantitative homotopical method called effective rebuilding. This constructs small classifying spaces of finite index subgroups, at the same time controlling the complexity of the homotopy. The method easily applies to free abelian groups and then extends recursively to a wide class of residually finite groups.

## CONTENTS

1. Introduction	1
2. The Borel construction and Geoghegan rebuilding	8
3. Cellular homology	10
4. Effective rebuilding	12
5. Proof of Proposition 4.1	17
6. Quality of rebuilding for nilpotent groups (Proof of Theorem E)	22
7. Quality of rebuilding of extensions by unipotent lattices	30
8. A general quantitative rebuilding theorem	31
9. Bounding torsion - A proof of Gabber's proposition 9.1	33
10. Farber sequences and cheap rebuilding property	34
11. Lattices in semi-simple Lie groups (Proof of Theorem B)	45
12. Application to mapping class groups (Proof of Theorem D)	47
References	48

## 1. INTRODUCTION

Let  $M$  be a finite volume Riemannian manifold or a finite CW-complex. One of the most powerful and well studied invariants of  $M$  are its homology groups. These abelian groups decompose to a free and a torsion part. The ranks of the free parts give the rational Betti numbers of  $M$ . One can also compute the mod  $p$  Betti numbers, from the full homology group, and the Euler characteristic, as the alternating sum of Betti numbers. For a finite sheeted cover of  $M$ , the Euler characteristic is multiplicative in the index of the cover, but already the rational Betti numbers can behave quite erratically in this respect. To smooth this behavior out, one can consider the  $j$ -th  $\ell^2$  homology, and measure the dimension of the corresponding Hilbert space using von Neumann dimension (Atiyah [Ati76]). Alternatively, one can consider the **growth** of the  $j$ -th rational Betti numbers over a suitable sequence of finite sheeted covers of  $M$ . As shown by the Lück approximation theorem [Lüc94] these two attempts give the same result, called the  $j$ -th  $\ell^2$  Betti number of

---

*Date:* June 24, 2021.

*2020 Mathematics Subject Classification.* 22E40; 57M07; 11F75; 20J05; 20J06; 20E26; 20F69; 20F36; 20G25.

$M$ . This result naturally led to studying the growth of other homological invariants as well, like the growth of the mod  $p$  Betti numbers and the growth of the torsion.

Over the years, the interest of the community has shifted from the study of the homology of spaces  $M$  to the homology of their fundamental groups. Our results are also expressed in terms of group homology.

Since the torsion grows at most exponentially in the index of the cover, the right definition of the  **$j$ -th torsion growth of  $M$**  is to take the logarithm of the cardinality of the  $j$ -th torsion group of the covering space, normalized by the index of the cover and consider its limit, for a suitable sequence of covers.

As of now, control of the torsion (and also the mod  $p$  Betti numbers) is much weaker than for the rational Betti numbers. In particular, we do not have the analogue of the Lück approximation theorem, even in the most natural settings. More than that, strikingly, we do not know a single example of a **finitely presented** group with positive **first** homology torsion growth over a decreasing sequence of finite index normal subgroups with trivial intersection, while, at the same time, it is a well-accepted conjecture that all sequences of congruence subgroups of arithmetic hyperbolic 3-manifold group have this property, see [BSV16, BD15, Ber18].

In this paper we prove vanishing results on the torsion growth of higher homology groups of a natural class of residually finite groups, using a new homotopical method called effective rebuilding. We apply this method to non-uniform higher rank lattices, mapping class groups and various Artin groups, among others. For the lattice case, we also obtain explicit estimates of the convergence for covering maps with respect to principal congruence subgroups.

**1.1. Main results.** We start by stating two theorems in their simplest forms that are good showcases for the more general and technical results of the paper. For a group  $\Gamma$ , its homology groups decompose as  $H_j(\Gamma, \mathbb{Z}) = \mathbb{Z}^{b_j(\Gamma)} \oplus H_j(\Gamma, \mathbb{Z})_{\text{tors}}$  where  $b_j(\Gamma)$  is the  $j$ -th Betti number and where  $H_j(\Gamma, \mathbb{Z})_{\text{tors}}$  is the torsion subgroup. Let  $|H_j(\Gamma, \mathbb{Z})_{\text{tors}}|$  denote its cardinality.

Our first result deals with an arbitrary sequence of finite index subgroups.

**Theorem A.** *Let  $\Gamma = \text{SL}_d(\mathbb{Z})$  ( $d \geq 3$ ) and let  $(\Gamma_n)_{n \in \mathbb{N}}$  be an injective sequence of finite index subgroups of  $\Gamma$ . Then for every degree  $j \leq d - 2$ , we have:*

$$\frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma_n]} \longrightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (1.1)$$

There are deep number-theoretical motivations to study torsion in the homology of arithmetic groups. It is directly related to algebraic K-theory and the Vandiver conjecture, see e.g. [Sou99, Eme14]. More recently it has attracted further attention since, thanks to the deep work of Scholze [Sch15], one can roughly say that mod  $p$  torsion classes in the homology of congruence subgroups of  $\text{SL}_d(\mathbb{Z})$  parametrize field extensions  $K/\mathbb{Q}$  whose Galois groups are subgroups of  $\text{PGL}_d(\overline{\mathbb{F}}_p)$ . Theorem A confirms a part of a general related conjecture of Venkatesh and the second author that postulates that the limit in (1.1) is zero for all  $d \geq 3$  and all  $j$  **except** when  $(d, j) = (3, 2)$  or  $(d, j) = (4, 4)$ , see [BV13, Ber18, AGMY20].

When restricting our attention to **principal congruence subgroups**

$$\Gamma(N) = \ker(\text{SL}_d(\mathbb{Z}) \rightarrow \text{SL}_d(\mathbb{Z}/N\mathbb{Z}))$$

our second result gives quantitative upper bounds on the torsion growth.

**Theorem B.** *Let  $\Gamma = \text{SL}_d(\mathbb{Z})$  ( $d \geq 3$ ). Then for all  $N \geq 1$  and  $j \leq d - 2$  we have*

$$\frac{\log |H_j(\Gamma(N), \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma(N)]} = O\left(\frac{\log N}{N^{d-1}}\right), \quad \text{as } N \rightarrow \infty. \quad (1.2)$$

We refer to Section 11 for the more general Theorem 11.1 about non-uniform lattices in semisimple Lie groups.

The first homology case of Theorem A, that is the case  $j = 1$ , is known, in particular it follows from [AGN17, Theorem 4]. Using the less elementary, but more classical machinery of **congruence subgroup property** (CSP) on these groups, they also show the stronger estimate that there exists a constant  $c$  depending only on  $d$  such that for all finite index subgroups  $\Gamma_0$  of  $\mathrm{SL}_d(\mathbb{Z})$  we have

$$|H_1(\Gamma_0, \mathbb{Z})_{\mathrm{tors}}| = |H_1(\Gamma_0, \mathbb{Z})| = |\Gamma_0^{\mathrm{ab}}| \leq [\mathrm{SL}_d(\mathbb{Z}) : \Gamma_0]^c, \quad (1.3)$$

see [AGN17, §5.1].

For  $d = 3$ , we expect that the degree bound  $j \leq d - 2$  is sharp for both theorems above. The general conjecture of [BV13] indeed postulates an exponential growth of torsion for  $\mathrm{SL}_3(\mathbb{Z})$  in degree 2. As an indication, big torsion groups do show up in the recent computational work of Ash, Gunnells, McConnell and Yasaki [AGMY20] for both cases. The degree bound  $j \leq d - 2$  is probably not optimal for higher degree. We shall see that it is a natural stopping point for our approach, though.

The non-effective version of our methods already gives applications for the growth of mod  $p$  Betti numbers for any prime  $p$ . Since we obtain a good control over the number of cells of finite covers, it follows that the growth of the  $j$ -th mod  $p$  Betti number is zero, under the same conditions as in Theorem A. The supplementary control offered by principal congruence subgroups leads as in Theorem B to explicit estimates on the growth of the mod  $p$  Betti numbers (see Theorem 11.1):

**Theorem C.** *Let  $\Gamma = \mathrm{SL}_d(\mathbb{Z})$  ( $d \geq 3$ ) and let  $(\Gamma_n)_{n \in \mathbb{N}}$  be an injective sequence of finite index subgroups of  $\Gamma$ . Then, for every field  $K$  and for every degree  $j \leq d - 2$ , we have*

$$\frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (1.4)$$

For the principal congruence subgroups  $\Gamma(N)$ , we have

$$\frac{\dim_K H_j(\Gamma(N), K)}{[\Gamma : \Gamma(N)]} = O\left(N^{(1-d)}\right), \quad \text{as } N \rightarrow \infty. \quad (1.5)$$

Note that the  $j = 1$  case of (1.4) again follows from known results. The first mod  $p$  Betti number is a lower bound for the minimal number of generators, hence the rank gradient dominates the growth of the first mod  $p$  Betti number. It is shown for instance in [CGdS19, Theorem 4.14(3)] that for  $\mathrm{SL}_d(\mathbb{Z})$  ( $d \geq 3$ ), the rank gradient vanishes for arbitrary injective sequences of finite index subgroups, using the work of Gaboriau on cost [Gab00]. In a somewhat different direction, Fraczyk proved that, for the finite field  $\mathbb{F}_2$ , the estimate  $\frac{\dim_{\mathbb{F}_2} H_1(\Gamma, K)}{\mathrm{vol}(\Gamma \backslash G)} = o(1)$  holds uniformly for all lattices in any simple higher rank real Lie group  $G$  [Fra18]. We wonder whether the methods implemented in our paper can be adapted to deal similarly with arbitrary sequences of lattices in a simple higher rank real Lie group  $G$ . For the rank 1 case, see the nice papers [Sau16, BGS20].

The results of Theorem C are new for higher homology groups. The closest result we are aware of is due to Calegari and Emerton and concerns  $p$ -adic chains like  $p^n$ -congruence subgroups  $\Gamma(p^n)$  in  $\mathrm{SL}_d(\mathbb{Z})$ , see [CE09, CE11, BLLS14] where the existence of a limit is established (not its value) and the error term is estimated. In the small degrees that our methods address, our error term is better, but note that their paper also deals with the difficult case around the middle dimension. Recent works [CE16, Cal15] and conjectures (personal communication) of Calegari and Emerton suggest that in reality the bound on  $|H_j(\Gamma(N), \mathbb{Z})_{\mathrm{tors}}|$  (rather than its logarithm) should be polynomial in  $N$  in degrees  $\leq d - 2$ . Let us also mention the paper of Kar, Kropholler and Nikolov [KKN17], in which they prove the vanishing of the  $j$ -th torsion growth for a wide class of amenable groups.

Although the results above are homological, we build a homotopical machinery to obtain them. This approach traces back to [Gab00, KN16, BK17]. Since we use a general topological approach, our work also applies to a wide class of groups such as torsion-free nilpotent groups, infinite polycyclic groups, Baumslag-Solitar groups  $\mathrm{BS}(1, n)$  and  $\mathrm{BS}(n, n)$  for any non-zero integer  $n$ , residually finite Artin groups,.. and does not need or assume underlying deep results like the CSP.

In particular, our method applies to mapping class groups of higher genus surfaces. For  $g, b \in \mathbb{N}$  let  $\alpha(g, b) = 2g - 2$  if  $g > 0$  and  $b = 0$ ,  $\alpha(g, b) = 2g - 3 + b$  if  $g > 0$  and  $b > 0$ , and  $\alpha(g, b) = b - 4$  if  $g = 0$ .

**Theorem D.** *Let  $S$  be an orientable surface of genus  $g > 0$  with  $b$  boundary components and let  $\Gamma = \mathcal{MCG}(S)$  be its mapping class group. Let  $(\Gamma_n)_{n \in \mathbb{N}}$  be a Farber sequence of finite index subgroups of  $\Gamma$ . Then for every field coefficient  $K$  and every degree  $j \leq \alpha(g, b)$ , we have*

$$\frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]} \longrightarrow 0 \quad \text{and} \quad \frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma_n]} \longrightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (1.6)$$

See Definition 10.1 for the definition of a Farber sequence. Examples are decreasing sequences of finite index normal subgroups with trivial intersection. Observe that for lattices in higher rank simple Lie groups, it follows from the Stuck–Zimmer Theorem [SZ94] via [ABB<sup>+</sup>17] that any injective sequence is indeed a Farber sequence. Both  $\text{SL}_d(\mathbb{Z})$  and  $\mathcal{MCG}(S)$  are then handled using the same method.

**1.2. Structure, arguments and further results.** Throughout this article, any group action on a CW-complex is required to respect the CW-complex structure. Our starting point is the obvious observation that, while the number of cells of a finite index cover  $\bar{\Sigma}$  of a CW-complex  $\Sigma$  is proportional to the index, it turns out that, for tori, the space  $\bar{\Sigma}$  can be **rebuilt** with a much simpler cell structure; indeed, with the same number of cells as  $\Sigma$  itself.

At the origin of the theory of “(co)homology of groups”, the standard homological invariants of a group  $\Gamma$  are obtained as the homological invariants of a (and hence of any) compact space  $\Sigma$  with fundamental group  $\Gamma$  and contractible universal cover  $\tilde{\Sigma}$ , whenever such a  $\Sigma$  exists. In fact, instead of contractibility, the  $\alpha$ -connectedness of  $\tilde{\Sigma}$  is enough to compute the homological invariants up to degree  $\alpha$  from the cellular chain complex of  $\Sigma$ .

Our strategy is to exploit the existence for certain groups  $\Gamma$  (like those appearing in Theorems A and D) of such a complex  $\Sigma$  with nicely embedded tori-like sub-complexes so as to capitalize on the “obvious observation” above.

Let  $\Gamma$  be a countable group acting on a CW-complex  $\Omega$  and let  $\alpha$  be a positive integer. Suppose that the CW-complex  $\Omega$  is  $\alpha$ -connected and that for every cell  $\omega \subset \Omega$  the stabilizer  $\Gamma_\omega$  acts trivially on the cell  $\omega$ . We may be led to consider a finite index subgroup first or to consider a barycentric subdivision so as to ensure this last condition. A general construction, called the **Borel construction** then associates to the action  $\Gamma \curvearrowright \Omega$  a  $\alpha$ -aspherical CW-complex  $\Sigma$  whose fundamental group is  $\Gamma$ . The construction in fact naturally leads to a **stack of CW-complexes**  $\Pi : \Sigma \rightarrow \Gamma \backslash \Omega$ , and, roughly speaking, the fiber over a cell  $\Gamma\omega$  of  $\Gamma \backslash \Omega$  is a classifying space for the stabilizer  $\Gamma_\omega$  of  $\omega$  under the action  $\Gamma \curvearrowright \Omega$ .

$$\begin{array}{ccccc} \Gamma_\omega \curvearrowright \widetilde{\Pi^{-1}(\Gamma\omega)} & & \Gamma \curvearrowright \tilde{\Sigma} & \longrightarrow & \Gamma \curvearrowright \Omega \\ \downarrow & & \downarrow & & \downarrow \\ B(\Gamma_\omega, 1) = \Pi^{-1}(\Gamma\omega) & \longrightarrow & \Sigma & \xrightarrow{\Pi} & \Gamma \backslash \Omega \end{array}$$

We briefly recall all these notions in Section 2 (in particular Proposition 2.1) as it will be useful for us. We refer to the excellent book of Geoghegan [Geo08, §6.1] for more details.

At the risk of a spoiler, we indicate right away that we shall use the complex  $\Omega$  to be the *rational Tits building*  $\Omega$  for  $\Gamma = \text{SL}_d(\mathbb{Z})$ . Note that this complex  $\Omega$  is  $(d - 3)$ -connected (see Section 11), which is the sole reason why we can only treat the homology groups in the range  $j \leq d - 2$  in Theorems B and A. Similarly in Theorem D for the mapping class group  $\mathcal{MCG}(S)$ , for which we shall use the *curve complex* as  $\Omega$  (see Section 12). The key aspect of our proof is that the stabilizers  $\Gamma_\omega$  in these actions  $\Gamma \curvearrowright \Omega$  contain non-trivial free abelian normal subgroups, so that each fiber is

itself a torus bundle. However, in its original form the Borel construction is not yet suitable for us. More precisely, in order to build “nicer classifying spaces” for the finite index subgroups that allow exploiting the structure of the stabilizers out of the total space  $\Sigma$  of the stack  $\Pi : \Sigma \rightarrow \Gamma \backslash \Omega$ , we make use of Geoghegan’s **Rebuilding Lemma** [Geo08, Prop. 6.1.4] (see Proposition 2.2 below). This is not yet enough. In order to get a grip on the torsion part of the homology, we make use of a proposition attributed to Gabber (see [Sou99, Prop. 3, p. 214]):

$$\log |H_j(\Sigma, \mathbb{Z})_{\text{tors}}| \leq (\# \text{ of } j\text{-cells}) \times \sup(\log \|\partial_{j+1}\|, 0). \quad (1.7)$$

See Section 9 where, for the convenience of the reader, we give a proof of this inequality following the point of view of [BV13]. This reduces the problem of estimating the torsion to bounding the number of cells and the norms of the boundary maps. Thus, to be able to really make use of Gabber’s proposition, we have to turn the Rebuilding Lemma into an effective statement. This is the content of our Proposition 4.2. In the end, this **Effective Rebuilding Lemma** is a machine that provides an explicit rebuilding of the total space  $\Sigma$  of a stack of complexes  $\Sigma \rightarrow \Gamma \backslash \Omega$  given a rebuilding of its fibers, and that moreover provides bounds on the number of cells and norms of the chain boundary maps.

In various interesting situations, each fiber of  $\Pi : \Sigma \rightarrow \Gamma \backslash \Omega$  is itself a torus bundle. The standard CW structure on the torus and an inductive procedure using the effective rebuilding lemma yield “classifying spaces” for the finite index subgroups which will be sufficiently “small” to prove our asymptotic theorems.

One needs to be able to rebuild efficiently not only free abelian groups but also finitely generated torsion-free nilpotent groups. This is the content of the general Theorem E stated below which we believe to be of independent interest.

First let us give a precise definition of a rebuilding (of good quality). This is a central notion of this paper.

For a CW-complex  $X$  we denote by  $X^{(\alpha)}$  its  $\alpha$ -skeleton and  $|X^{(\alpha)}|$  its number of  $\alpha$ -cells. By convention  $X^{(-1)}$  is the empty set.

**Definition 1** (Rebuilding). Let  $\alpha \in \mathbb{N}$  and let  $X$  be a CW-complex with finite  $\alpha$ -skeleton. A  $\alpha$ -**rebuilding** of  $X$  consists of the following data  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$ :

- (1) a CW-complex  $X'$  with finite  $\alpha$ -skeleton;
- (2) two cellular maps  $\mathbf{g} : X^{(\alpha)} \rightarrow X'^{(\alpha)}$  and  $\mathbf{h} : X'^{(\alpha)} \rightarrow X^{(\alpha)}$  that are homotopy inverse to each other up to dimension  $\alpha - 1$ , i.e.,  $\mathbf{h} \circ \mathbf{g}|_{X^{(\alpha-1)}} \sim \text{id}|_{X^{(\alpha-1)}}$  within  $X^{(\alpha)}$  and  $\mathbf{h}|_{X'^{(\alpha-1)}} \sim \text{id}|_{X'^{(\alpha-1)}}$  within  $X'^{(\alpha)}$ ;
- (3) a cellular homotopy  $\mathbf{P} : [0, 1] \times X^{(\alpha-1)} \rightarrow X^{(\alpha)}$  between the identity and  $\mathbf{h} \circ \mathbf{g}$ , i.e.,  $\mathbf{P}(0, \cdot) = \text{id}|_{X^{(\alpha-1)}}$  and  $\mathbf{P}(1, \cdot) = \mathbf{h} \circ \mathbf{g}|_{X^{(\alpha-1)}}$

**Definition 2** (Quality of a rebuilding). Given real numbers  $T, \kappa \geq 1$ , we say that  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$  is a  $\alpha$ -rebuilding of **quality**  $(T, \kappa)$  if we have

$$\forall j \leq \alpha, \quad |X'^{(j)}| \leq \kappa T^{-1} |X^{(j)}| \quad (\text{cells bound})$$

$$\forall j \leq \alpha, \quad \log \|g_j\|, \log \|h_j\|, \log \|\rho_{j-1}\|, \log \|\partial'_j\| \leq \kappa(1 + \log T) \quad (\text{norms bound})$$

where  $\partial'$  is the boundary map on  $X'$ , we denote by  $g$  and  $h$  the chain maps respectively associated to  $\mathbf{g}$  and  $\mathbf{h}$ , and  $\rho : C_\bullet(X) \rightarrow C_{\bullet+1}(X)$  is the chain homotopy induced by  $\mathbf{P}$  in the cellular chain complexes:

$$\begin{array}{ccccccc} C_\alpha(X) & \xrightarrow{\partial_\alpha} & \dots & \xrightarrow{\partial_2} & C_1(X) & \xrightarrow{\partial_1} & C_0(X) \\ g_\alpha \uparrow & \leftarrow \rho_{\alpha-1} & & \leftarrow \rho_1 & g_1 \uparrow & \leftarrow \rho_0 & g_0 \uparrow \\ & h_\alpha & & h_1 & & h_0 & \\ C_\alpha(X') & \xrightarrow{\partial'_\alpha} & \dots & \xrightarrow{\partial'_2} & C_1(X') & \xrightarrow{\partial'_1} & C_0(X') \end{array} \quad (1.8)$$

and the norms  $\|\cdot\|$  are derived from the canonical  $\ell^2$ -norms on the cellular chain complexes.

We will simplify the notation and write  $(X, X')$  instead of  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$  when the explicit cellular maps are not relevant.

The definition above captures an intrinsic tension between “having few cells” and “maintaining tame norms”.

Given a finite cover  $X_1 \rightarrow X$  (of large degree) our main task is to construct a rebuilding  $(X_1, X'_1)$  of sufficiently good quality  $(T_1, \kappa_1)$ . In some cases it is possible to take  $\kappa_1 = \kappa(X)$  independently of the cover and  $T_1$  linear in the degree, see Sections 6 and 7. In particular for finitely generated torsion-free nilpotent groups (called **unipotent lattices** in the text) the precise efficient rebuilding we obtain can be stated as follows.

**Theorem E.** *Let  $\Lambda$  be a finitely generated torsion-free nilpotent group. If  $Y_0$  is a compact  $K(\Lambda, 1)$  space, there exists a constant  $\kappa_{Y_0} \geq 1$  such that for every finite index subgroup  $\Lambda_1 \leq \Lambda$ , the cover  $Y_1 = \Lambda_1 \backslash \tilde{Y}_0$  admits a  $\alpha$ -rebuilding  $(Y_1, Y'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  of quality  $([\Lambda : \Lambda_1], \kappa_{Y_0})$  for every  $\alpha$ .*

Recall that a  $K(\Lambda, 1)$  space is a CW complex with fundamental group isomorphic with  $\Lambda$  and whose universal cover is contractible.

*Remark.* Since the number of  $j$ -cells of  $Y_1$  is  $|Y_1^{(j)}| = [\Lambda : \Lambda_1] |Y_0^{(j)}|$ , the number of  $j$ -cells of the  $K(\Lambda_1, 1)$  space  $Y'_1$  satisfies the bound  $|Y'_1{}^{(j)}| \leq \kappa_{Y_0} [\Lambda : \Lambda_1]^{-1} |Y_1^{(j)}| = \kappa_{Y_0} |Y_0^{(j)}|$ , independent of  $\Lambda_1$ .

*Remark.* The proof of Theorem E (in Section 6) yields for various  $\Lambda_1$  a rebuilding  $Y'_1$  with only  $O(2^{\mathfrak{h}})$  cells, where  $\mathfrak{h}$  is the Hirsch length of  $\Lambda$ . This is in fact “the” minimal number of cells that a  $K(\Lambda, 1)$  CW-complex can have, since  $\sum_j \dim_{\mathbb{Q}} H_j(\Lambda, \mathbb{Q}) = 2^{\mathfrak{h}}$ .

Theorem E is used for the proof of Theorem B; the latter is in fact deduced from a general result: Theorem 8.1 proved in Section 8. One key feature of the **principal** congruence subgroup  $\Gamma(N)$  of Theorem B is that it intersects every infinite unipotent subgroup of  $\mathrm{SL}_d(\mathbb{Z})$  along a subgroup of index at least  $N$ . Note that general sequences of finite index subgroups of  $\mathrm{SL}_d(\mathbb{Z})$  do not have this property, e.g., for each positive integer  $N$  the **non-principal** congruence subgroup

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & D \end{pmatrix} \in \mathrm{SL}_d(\mathbb{Z}) : a \in \mathbb{Z}, b, c \in \mathbb{Z}^{d-1}, D \in \mathrm{GL}_{d-1}(\mathbb{Z}), \text{ and } N|c \right\} \subset \mathrm{SL}_d(\mathbb{Z})$$

contains the whole group of upper-triangular unipotent matrices with integer coefficients. To prove Theorem A we need to show that if  $(\Gamma_n)$  is an injective sequence of finite index subgroups of  $\mathrm{SL}_d(\mathbb{Z})$ , then  $\Gamma_n$  intersects most of the infinite unipotent subgroups of  $\mathrm{SL}_d(\mathbb{Z})$  along subgroups of large index.

In general to deal with arbitrary Farber sequences of a given residually finite group  $\Gamma$  we introduce a property that we believe to be of independent interest. We say that  $\Gamma$  has the **cheap  $\alpha$ -rebuilding property** (Definition 10.6) if it admits a  $K(\Gamma, 1)$  space  $X$  with finite  $\alpha$ -skeleton and a constant  $\kappa_X \geq 1$  that satisfy the following property. For every Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$  for  $\Gamma$  and  $T \geq 1$ , there exists  $n_0$  such that for  $n \geq n_0$  the cover  $X_n := \Gamma_n \backslash \tilde{X}$  admits a  $\alpha$ -rebuilding  $(X_n, Y_n)$  of quality  $(T, \kappa_X)$ . We note that cheap 0-rebuilding simply means that  $\Gamma$  is infinite.

We establish a robust bootstrapping criterion for a group to have the cheap  $\alpha$ -rebuilding property:

**Theorem F** (Theorem 10.10). *Let  $\Gamma$  be a residually finite group acting on a CW-complex  $\Omega$  in such a way that any element stabilizing a cell fixes it pointwise. Let  $\alpha \in \mathbb{N}$  and assume that the following conditions hold:*

- (1)  $\Gamma \backslash \Omega$  has finite  $\alpha$ -skeleton;
- (2)  $\Omega$  is  $(\alpha - 1)$ -connected;
- (3) The stabilizer of each cell of dimension  $j \leq \alpha$  has the cheap  $(\alpha - j)$ -rebuilding property.

*Then,  $\Gamma$  itself has the cheap  $\alpha$ -rebuilding property.*

As a corollary we get that the cheap  $\alpha$ -rebuilding property is an invariant of commensurability. Also, we get that it holds for all infinite polycyclic groups for all  $\alpha$ . It also passes from a normal subgroup  $N \triangleleft \Gamma$  to  $\Gamma$  if  $N \setminus \Gamma$  is of type  $F_\alpha$ . See Corollary 10.14. More significantly, we show

**Theorem G.** *Let  $\alpha \geq 0$ . The following groups have the cheap  $\alpha$ -rebuilding property:*

- (1) *Arithmetic lattices of  $\mathbb{Q}$ -rank at least  $\alpha + 1$  (Theorem 11.3).*
- (2) *Finitely generated residually finite Artin groups satisfying the  $K(\pi, 1)$  conjecture and whose nerve is  $(\alpha - 1)$ -connected (Theorem 10.18).*
- (3) *Mapping class groups  $\mathcal{MGC}(S)$  where  $S$  is a surface of genus  $g > 0$  with  $b > 0$  boundary components and  $2g - 3 + b \geq \alpha$  or  $g > 0$ ,  $b = 0$  and  $2g - 2 \geq \alpha$  (Theorem 12.1).*

The “cells bound” condition of Definition 2 in the cheap  $\alpha$ -rebuilding property is very much related to the notion of slowness introduced in [BK17]. This condition already implies the vanishing of the growth of homology over arbitrary fields for a wide class of groups. In particular a group with the cheap 2-rebuilding property is finitely presented and economical, in the sense of [KN16], with respect to any Farber chain of finite index subgroups. For controlling the torsion, however, we need the full force of the cheap  $\alpha$ -rebuilding property:

**Theorem H** (Theorem 10.21). *Let  $\Gamma$  be a countable group of type  $F_{\alpha+1}$  that has the cheap  $\alpha$ -rebuilding property for some non-negative integer  $\alpha$ . Then, for every Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$ , coefficient field  $K$  and  $0 \leq j \leq \alpha$  we have*

$$\lim_{n \rightarrow \infty} \frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma_n]} = 0.$$

In analogy with the Lück approximation theorem [Lüc94] which identifies the first limit with the usual  $\ell^2$ -Betti numbers  $b_j^{(2)}(\Gamma; K)$  when  $K = \mathbb{Q}$ , we will loosely write

$$b_j^{(2)}(\Gamma; K) = \limsup_{n \rightarrow \infty} \frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]}$$

once the sequence  $(\Gamma_n)$  is fixed. Note that non-abelian finitely generated free groups do not have the cheap 1-rebuilding property (since  $b_1^{(2)}(\mathbf{F}_r; \mathbb{Q}) = r - 1 \neq 0$ ).

According to Theorem G (1)  $\text{SL}_d(\mathbb{Z})$  has the cheap  $(d - 2)$ -rebuilding property. Theorem H therefore implies Theorem A. Similarly Theorem G (3) and Theorem H imply Theorem D.

Theorem H is related to [AGN17, Theorems 7 and 9] where Abert, Gelander and Nikolov investigate the first homology torsion growth for **right-angled groups**, i.e. groups that admit a finite generating list  $\{\gamma_1, \dots, \gamma_m\}$  of elements of infinite order such that  $[\gamma_i, \gamma_{i+1}] = 1$  for  $i = 1, \dots, m - 1$ . This is a class of groups featured in [Gab00] where  $\text{SL}_d(\mathbb{Z})$  ( $d \geq 3$ ) being a right-angled group is exploited to compute its cost. These groups have the cheap 1-rebuilding property, see Proposition 10.16. We therefore recover that finitely presented right-angled groups have vanishing first homology torsion growth along any Farber sequence. Note that already in degree  $j = 1$ , Theorem H also establishes the vanishing of the homology torsion growth for a natural class of groups where [AGN17] doesn't apply (see Example 10.17). Moreover, thanks to Theorem G (2) it applies to quite general Artin groups in higher degrees and yields:

**Theorem I.** *Let  $\Gamma$  be a finitely generated residually finite Artin group satisfying the  $K(\pi, 1)$  conjecture and whose nerve is a  $(\alpha - 1)$ -connected. Then, for every Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$ , coefficient field  $K$  and  $0 \leq j \leq \alpha$  we have*

$$b_j^{(2)}(\Gamma; K) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma_n]} = 0.$$

When  $\Gamma$  is a right-angled Artin group (**RAAG**) the first part of the theorem is not new: In fact Avramidi, Okun and Schreve [AOS20] have even computed all the  $\ell^2$ -Betti numbers  $b_j^{(2)}(\Gamma; K)$ .



Their result is that  $b_j^{(2)}(\Gamma; K) = \bar{b}_{j-1}(L; K)$ , the reduced Betti number of the nerve  $L$  of  $\Gamma$ . Thus  $b_j^{(2)}(\Gamma; K)$  is indeed equal to 0 when  $L$  is  $(j-1)$ -connected. This shows that Theorem [G \(2\)](#) is optimal for RAAGs. Building on their computation of the  $\ell^2$ -Betti numbers  $b_j^{(2)}(\Gamma; \mathbb{F}_2)$ , Avramidi, Okun and Schreve also prove that a RAAG  $\Gamma$  whose nerve is a flag triangulation of  $\mathbb{R}P^2$  has exponential homology torsion growth in degree  $j = 2$ , see [[AOS20](#), Corollary 3]. Since  $\Gamma$  has the cheap 1-rebuilding property (it is in fact right-angled) this shows that Theorem [H](#) is in a certain sense optimal: for  $\alpha = 1$ , there are groups that have the cheap  $\alpha$ -rebuilding property and have exponential torsion growth in degree  $j = \alpha + 1$ .

To conclude this introduction, we finally raise the following:

**Question 1.1.** *Let  $\Gamma$  be the fundamental group of a closed, or finite volume, hyperbolic 3-manifold. Does  $\Gamma$  have the cheap 1-rebuilding property?*

It is tempting to believe that the answer is no. These groups have zero first  $\ell^2$ -Betti number, so the Betti number criterion does not apply. But it is conjectured that for arithmetic hyperbolic 3-manifolds groups the torsion in degree 1 grows exponentially along Farber sequences of congruence subgroups, see [[BSV16](#), [BD15](#), [Ber18](#)].

**Acknowledgement.** We would like to thank Ian Biringer, Ross Geoghegan, Grégory Ginot, Athanase Papadopoulos, and Shmuel Weinberger for helpful conversations. We especially thank Boris Okun and Kevin Schreve for reading a preliminary version of this paper and making valuable comments. We also thank Emma Bergeron for drawing our pictures. M. A. and M. F. are partially supported by ERC Consolidator Grant 648017. D. G. was partially supported by the ANR project GAMME (ANR-14-CE25-0004) and by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). M. F. would like to thank IAS Princeton for an excellent working environment during a crucial stage of this project.

## 2. THE BOREL CONSTRUCTION AND GEOGHEGAN REBUILDING

Let  $\Gamma$  be a countable group acting on a CW-complex  $\Omega$ .

Technically speaking, the **Borel construction** is a trick that converts an action of a group  $\Gamma$  on a space  $\Omega$  into a **free action** of  $\Gamma$  on a homotopy equivalent space  $\Omega'$ . See for instance [[Hat02](#), 3.G.2]. Namely, take  $\Omega' = \Omega \times E\Gamma$  with the diagonal action of  $\Gamma$ ,  $\gamma(y, z) = (\gamma y, \gamma z)$  where  $\Gamma$  acts on  $E\Gamma$  (the universal cover of some classifying space  $B\Gamma$ ) as deck transformations. The diagonal action is free, in fact a covering space action, since this is true for the action in the second coordinate. The orbit space of this diagonal action is usually denoted  $\Omega \times_{\Gamma} E\Gamma$ . We now briefly explain the “stack” interpretation of Geoghegan [[Geo08](#), §6.1] alluded to in the introduction.

A cellular map  $\Pi : \Sigma \rightarrow \Phi$  between two CW-complexes is a **stack of CW-complexes** with **base space**  $\Phi$ , **total space**  $\Sigma$  and CW-complexes called **fibers**  $F_e$  over  $e$ , if for each  $n \geq 1$  (denoting by  $E_n$  the set of  $n$ -cells of  $\Phi$ ) there is a cellular map

$$f_n : \bigsqcup_{e \in E_n} F_e \times \mathbb{S}^{n-1} \rightarrow \Pi^{-1}(\Phi^{(n-1)}) \quad (2.1)$$

and a homeomorphism

$$k_n : \Pi^{-1}(\Phi^{(n-1)}) \cup_{f_n} \left( \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n \right) \rightarrow \Pi^{-1}(\Phi^{(n)}) \quad (2.2)$$

satisfying:

- (1)  $k_n$  agrees with the inclusion on  $\Pi^{-1}(\Phi^{(n-1)})$ ,
- (2)  $k_n$  maps each cell onto a cell, and

(3) the following diagram commutes up to homotopy relative to  $\Pi^{-1}(\Phi^{(n-1)})$ :

$$\begin{array}{ccc}
 \Pi^{-1}(\Phi^{(n-1)}) \sqcup \left( \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n \right) & & (2.3) \\
 \downarrow \text{quotient} & \searrow^{(\Pi|, \chi_e \circ \text{pr}_2)} & \\
 \Pi^{-1}(\Phi^{(n-1)}) \cup_{f_n} \left( \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n \right) & & \Phi^{(n)} \\
 \downarrow k_n & \searrow^{\Pi|} & \\
 \Pi^{-1}(\Phi^{(n)}) & & 
 \end{array}$$

where  $(\Pi|, \chi_e \circ \text{pr}_2) : \Pi^{-1}(\Phi^{(n-1)}) \sqcup \left( \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n \right) \rightarrow \Phi^{(n)}$  is the restriction of  $\Pi$  on  $\Pi^{-1}(\Phi^{(n-1)})$ , while on each  $F_e \times \mathbb{B}^n$ , it is the composition of the projection onto  $\mathbb{B}^n$  followed by the characteristic map  $\chi_e$  of the cell  $e \in E_n$  in the CW-complex  $\Phi$ . Thus there is a homotopy

$$\mathfrak{H} : \left( \Pi^{-1}(\Phi^{(n-1)}) \sqcup \left( \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n \right) \right) \times [0, 1] \rightarrow \Phi^{(n)}$$

such that

$$\mathfrak{H}|_{\Pi^{-1}(\Phi^{(n-1)}) \times [0, 1]} = \Pi|, \quad \mathfrak{H}(\cdot, 0) = (\Pi|, \chi_e \circ \text{pr}_2), \quad \text{and} \quad \mathfrak{H}(\cdot, 1) = \Pi| \circ k_n \circ \text{quotient}.$$

Note that the strong commutation of diagram (2.3) required in the definition of stacks in [Geo08, Section 6.1, p. 147] (2008 version of the book) is in fact too strong and that [Geo08, Proposition 6.1.4] needs the commutation to occur up to homotopy. Both the flaw and the way to fix it have been pointed out to us by Boris Okun and Kevin Schreve. We are extremely grateful to them for that.

Following [Geo08, §6.1], the Borel construction for  $\Gamma \curvearrowright \Omega$  eventually takes the form of a stack  $\Pi : \Sigma \rightarrow \Gamma \backslash \Omega$  below:

**Proposition 2.1.** *Let  $\Gamma$  be a countable group acting on a simply connected CW-complex  $\Omega$  so that for every cell  $\omega \subset \Omega$  the stabilizer  $\Gamma_\omega$  acts trivially on  $\omega$ . Write  $\Phi = \Gamma \backslash \Omega$ . Then there exists a stack of CW-complexes  $\Pi : \Sigma \rightarrow \Phi$  with fiber  $F_e$  over  $e$  such that*

- (1) *The fundamental group  $\pi_1(\Sigma)$  is isomorphic to  $\Gamma$ .*
- (2) *For each cell  $e$  of  $\Phi$  the fiber  $F_e$  is contractible and  $\pi_1(F_e) \cong \Gamma_\omega$ , where  $\omega$  is any cell in  $\Omega$  above  $e$ .*

Moreover, if  $\Omega$  is  $n$ -connected then the universal cover  $\tilde{\Sigma}$  of the total space  $\Sigma$  is  $n$ -connected.

Recall that a CW-complex space  $Y$  is said to be  **$n$ -connected** if it is connected and all its homotopy groups  $\pi_j Y$  are trivial for  $0 \leq j \leq n$ . By convention, a  $(-1)$ -connected space will just be an arbitrary topological space.

Note that if  $\omega$  and  $\omega'$  are two cells of  $\Omega$  above a cell  $e$  of  $\Phi$ , they are in the same  $\Gamma$ -orbit and thus the stabilizers  $\Gamma_\omega$  and  $\Gamma_{\omega'}$  are conjugate in  $\Gamma$ . In particular these stabilizers are isomorphic, so there is no harm in writing  $\Gamma_e := \Gamma_\omega$ .

In practice it is often desirable to replace the fibers  $F_e$  by different CW-complexes of the same homotopy types. Geoghegan's stack decomposition and his **rebuilding Lemma** [Geo08, Proposition 6.1.4] makes this possible:

**Proposition 2.2** (Rebuilding Lemma, Geoghegan [Geo08, Proposition 6.1.4]). *Let  $\Pi : \Sigma \rightarrow \Phi$  be a stack of CW-complexes. If for each cell  $e$  of  $\Phi$  we are given a CW-complex  $F'_e$  of the same homotopy type as  $F_e$ , then there is a stack of CW-complexes  $\Pi' : \Sigma' \rightarrow \Phi$  with fiber  $F'_e$  over  $e$ , and a homotopy equivalence  $\mathfrak{g}$  making the following diagram commute up to homotopy over each closed*

cell:

$$\begin{array}{ccc} \Sigma & \xrightarrow{\mathbf{g}} & \Sigma' \\ & \searrow \Pi & \swarrow \Pi' \\ & & \Phi \end{array}$$

More precisely,  $\Pi' \circ \mathbf{g}$  is homotopy equivalent to  $\Pi$  via a homotopy equivalence such that for every closed cell  $e$  of  $\Phi$ , the restriction of the homotopy to  $\Pi^{-1}(e) \times [0, 1]$  has its image in  $e$ .

Proposition 2.2 allows one, up to homotopy equivalence, to replace each fiber  $F_e$  in Proposition 2.1 by a prescribed classifying space for  $\Gamma_\omega$ .

Recall that a group  $\Gamma$  is of type  $F_n$  if it admits a classifying space whose  $n$ -skeleton is finite; equivalently if it admits a  $(n-1)$ -aspherical CW-complex  $X$  with finite  $n$ -skeleton and  $\pi_1(X) \simeq \Gamma$  (since turning  $X$  into an aspherical complex can be made by adding only cells of dimension  $\geq n+1$ ).

Let  $\alpha \in \mathbb{N}$ . Suppose that each group  $\Gamma_e$  is of type  $F_\alpha$ . Thanks to the rebuilding lemma we may then assume that:

Each fiber  $F_e$  of the stack  $\Pi : \Sigma \rightarrow \Phi$  has a finite  $\alpha$ -skeleton.

If we suppose furthermore that  $\Omega$  is  $(\alpha-1)$ -connected then the universal cover of the total space  $\tilde{\Sigma}$  is also  $(\alpha-1)$ -connected. The cellular chain complex  $C_q(\tilde{\Sigma})$  therefore gives a partial resolution

$$C_\alpha(\tilde{\Sigma}) \rightarrow \cdots \rightarrow C_0(\tilde{\Sigma}) \rightarrow \mathbb{Z} \rightarrow 0$$

of  $\mathbb{Z}$  by free  $\mathbb{Z}[\Gamma]$ -modules of finite rank.

*Remark 2.3.* If  $\bar{\Gamma} \leq \Gamma$  is a subgroup of finite index, then in the commuting diagram

$$\begin{array}{ccc} \bar{\Sigma} = \bar{\Gamma} \backslash \tilde{\Sigma} & \xrightarrow{\bar{\Pi}} & \bar{\Phi} = \bar{\Gamma} \backslash \Omega \\ \downarrow \zeta & \circlearrowleft & \downarrow \\ \Sigma = \Gamma \backslash \tilde{\Sigma} & \xrightarrow{\Pi} & \Phi = \Gamma \backslash \Omega \end{array} \quad (2.4)$$

the map  $\bar{\Pi}$  is naturally a stack satisfying (1) and (2) of Proposition 2.1, the map  $\zeta$  is a finite cover. For any  $n \in \mathbb{N}$  we thus have

$$|\bar{\Sigma}^{(n)}| = [\Gamma : \bar{\Gamma}] |\Sigma^{(n)}|.$$

Proposition 2.2 gives a way to improve the covering space  $\bar{\Sigma}$  by simplifying the fibers of the stack  $\bar{\Pi}$ . Our next goal is to make this procedure explicit enough to control the boundary maps in (cellular) homology. We first recall some basic facts about the latter.

### 3. CELLULAR HOMOLOGY

The material in this section covers the basic properties of cellular homology in the context of stacks of CW-complexes. We review it to set up the notation for the following sections. Let  $\Pi : \Sigma \rightarrow \Phi$  be a stack of CW-complexes with fiber over each cell  $e$  of  $\Phi$  denoted by  $F_e$ .

Recall that by definition, the  $\mathbb{Z}$ -module of the degree  $a$  cellular chains is

$$C_a(\Sigma) = H_a(\Sigma^{(a)}, \Sigma^{(a-1)}).$$

Now each homeomorphism (2.2) induces a map of pairs

$$(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n) \rightarrow (\Sigma^{(n+k)}, \Sigma^{(n+k-1)})$$

which induces an injective map

$$H_{n+k}(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n) \rightarrow C_{n+k}(\Sigma). \quad (3.1)$$

Considering the long exact sequence for the pair

$$(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n),$$

Künneth theorem yields a natural isomorphism of  $\mathbb{Z}$ -modules:

$$H_k(F_e^{(k)}, F_e^{(k-1)}) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \xrightarrow{\cong} H_{n+k}(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n). \quad (3.2)$$

By definition, we have

$$H_k(F_e^{(k)}, F_e^{(k-1)}) = C_k(F_e).$$

The  $\mathbb{Z}$ -module  $H_n(\mathbb{B}^n, \mathbb{S}^{n-1})$  is free of rank one generated by the relative fundamental class  $[\mathbb{B}^n, \mathbb{S}^{n-1}]$ . From now on we will identify

$$C_k(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \quad (3.3)$$

with the subspace of  $C_{n+k}(\Sigma)$  spanned by the image of the composition of the maps (3.2) and (3.1).

The ascending filtration of  $\Phi$  by its  $n$ -skeleta  $\Phi^{(n)}$  induced via  $\Pi : \Sigma \rightarrow \Phi$  then yields an ascending filtration

$$F^n C_\bullet(\Sigma) = \bigoplus_{e \in \Phi^{(n)}} C_{\bullet - \dim(e)}(F_e) \otimes H_{\dim e}(\mathbb{B}^{\dim e}, \mathbb{S}^{\dim e - 1}) \quad (3.4)$$

on the cellular chain complex  $C_\bullet(\Sigma)$ . Recall that the boundary operator  $\partial : C_a(\Sigma) \rightarrow C_{a-1}(\Sigma)$  is defined using the homology long exact sequences of the pairs  $(\Sigma^{(a)}, \Sigma^{(a-1)})$  and  $(\Sigma^{(a-1)}, \Sigma^{(a-2)})$ :

$$\begin{array}{ccc} C_a(\Sigma) & \xrightarrow{\partial} & C_{a-1}(\Sigma) \\ \downarrow \cong & & \downarrow \cong \\ H_a(\Sigma^{(a)}, \Sigma^{(a-1)}) & \xrightarrow{\delta} & H_{a-1}(\Sigma^{(a-1)}, \Sigma^{(a-2)}). \end{array}$$

In restriction to the image of (3.1) the composition of maps in the diagram above is the composition of the map

$$H_{n+k}(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n) \xrightarrow{\delta = \delta_1 \oplus \delta_2} H_{n+k-1}(F_e^{(k)} \times \mathbb{S}^{n-1}) \oplus H_{n+k-1}(F_e^{(k-1)} \times \mathbb{B}^n)$$

by the direct sum of the two natural maps

$$\begin{aligned} H_{n+k-1}(F_e^{(k)} \times \mathbb{S}^{n-1}) &\longrightarrow H_{n+k-1}(F_e^{(k)} \times \mathbb{S}^{n-1}, F_e^{(k-1)} \times \mathbb{S}^{n-1}) \\ H_{n+k-1}(F_e^{(k-1)} \times \mathbb{B}^n) &\longrightarrow H_{n+k-1}(F_e^{(k-1)} \times \mathbb{B}^n, F_e^{(k-1)} \times \mathbb{S}^{n-1}). \end{aligned}$$

**Definition 3.1** (In the fiber,  $\partial^{\text{vert}}$ ). Let

$$\partial^{\text{vert}} : C_k(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \rightarrow C_{k-1}(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1})$$

be the map that makes the following diagram commute:

$$\begin{array}{ccc} C_k(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) & \xrightarrow{\partial^{\text{vert}}} & C_{k-1}(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \\ \downarrow \cong & & \uparrow \cong \\ H_{n+k}(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n) & \xrightarrow{\delta_2} & H_{n+k-1}(F_e^{(k-1)} \times \mathbb{B}^n) \longrightarrow H_{n+k-1}(F_e^{(k-1)} \times \mathbb{B}^n, F_e^{(k-1)} \times \mathbb{S}^{n-1}). \end{array}$$

Given  $e \in E_n$ , we denote by

$$f_e : F_e \times \mathbb{S}^{n-1} \rightarrow \Sigma$$

the map obtained by restricting  $f_n$ . By construction it induces a map of pairs

$$(F_e^{(k)} \times \mathbb{S}^{n-1}, F_e^{(k-1)} \times \mathbb{S}^{n-1}) \longrightarrow (\Sigma^{(n+k-1)}, \Sigma^{(n+k-2)}).$$

**Definition 3.2** (In the base,  $\partial^{\text{hor}}$ ). Let

$$\partial^{\text{hor}} : C_k(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \rightarrow C_{n+k-1}(\Sigma)$$

be the map that makes the following diagram commute:

$$\begin{array}{ccc} C_k(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) & \xrightarrow{\partial^{\text{hor}}} & C_{n+k-1}(\Sigma) \\ \downarrow \cong & & \uparrow (f_e)_* \\ H_{n+k}(F_e^{(k)} \times \mathbb{B}^n, F_e^{(k)} \times \mathbb{S}^{n-1} \cup F_e^{(k-1)} \times \mathbb{B}^n) & \xrightarrow{\delta_1} & H_{n+k-1}(F_e^{(k)} \times \mathbb{S}^{n-1}) \longrightarrow H_{n+k-1}(F_e^{(k)} \times \mathbb{S}^{n-1}, F_e^{(k-1)} \times \mathbb{S}^{n-1}). \end{array}$$

By naturality we get the following:

**Lemma 3.3.** *The boundary map  $\partial : C_\bullet(\Sigma) \rightarrow C_{\bullet-1}(\Sigma)$  decomposes as*

$$\partial = \partial^{\text{vert}} + \partial^{\text{hor}} \quad (3.5)$$

where  $\partial^{\text{vert}}$  preserves each summand  $C_\bullet(F_e) \otimes H_{\dim e}(\mathbb{B}^{\dim e}, \mathbb{S}^{\dim e-1})$  and acts on it by the boundary operator of the cellular chain complex  $C_\bullet(F_e)$ , and where

$$\partial^{\text{hor}} : F^n C_\bullet(\Sigma) \rightarrow F^{n-1} C_{\bullet-1}(\Sigma)$$

maps  $c \otimes [\mathbb{B}^n, \mathbb{S}^{n-1}] \in C_\bullet(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1})$  (with  $e \in E_n$ ) to

$$(-1)^{\dim c} (f_e)_*(c \otimes [\mathbb{S}^{n-1}]).$$

*Proof.* The sign here comes from the isomorphism in the first vertical arrow of the commutative diagram of Definition 3.2. <sup>1</sup>  $\square$

#### 4. EFFECTIVE REBUILDING

Let  $\Phi$  be a finite CW-complex and let  $\Sigma \rightarrow \Phi$  be a stack of CW-complexes with fibers  $F_e$ . Now, if for each cell  $e$  of  $\Phi$  we are given a CW-complex  $F'_e$  of the same homotopy type as  $F_e$ , Proposition 2.2 implies that there exists a stack of CW-complexes  $\Sigma' \rightarrow \Phi$  with fiber  $F'_e$  over  $e$ , and a homotopy equivalence  $\mathbf{g}$  making the following diagram commute up to homotopy over each cell:

$$\begin{array}{ccc} \Sigma & \xrightarrow{\mathbf{g}} & \Sigma' \\ & \searrow \Pi & \swarrow \Pi' \\ & & \Phi \end{array} \quad (4.1)$$

For each cell  $e$  of  $\Phi$  let  $\mathbf{k}_e : F_e \rightarrow F'_e$  be a cellular map that induces a homotopy equivalence. In particular there exists a cellular map  $\mathbf{l}_e : F'_e \rightarrow F_e$  and a homotopy

$$\Sigma_e : [0, 1] \times F_e \rightarrow F_e \quad (4.2)$$

between  $\Sigma_e(0, \cdot) = \mathbf{l}_e \circ \mathbf{k}_e$  and  $\Sigma_e(1, \cdot) = \text{id}_{F_e}$ .

The maps  $\mathbf{k}_e$  and  $\mathbf{l}_e$  induce chain maps

$$k_e : C_\bullet(F_e) \rightarrow C_\bullet(F'_e) \quad \text{and} \quad l_e : C_\bullet(F'_e) \rightarrow C_\bullet(F_e).$$

The homotopy  $\Sigma_e$  induces a map  $\sigma_e : C_\bullet(F_e) \rightarrow C_{\bullet+1}(F_e)$ , given by

$$\sigma_e(c) = (\Sigma_e)_*([I, \partial I] \otimes c), \quad \text{for any cell } c \subset F_e.$$

The map  $\sigma_e$  is then a chain homotopy between 1 and  $l_e \circ k_e$ , i.e.

$$l_e \circ k_e - 1 = \partial \sigma_e + \sigma_e \partial.$$

We denote by

$$k : C_\bullet(\Sigma) \rightarrow C_\bullet(\Sigma'), \quad l : C_\bullet(\Sigma') \rightarrow C_\bullet(\Sigma)$$

<sup>1</sup>Analogously, the cellular chain complex of a product  $X \times Y$  of two CW-complexes is the tensor product  $C_\bullet(X) \otimes C_\bullet(Y)$  equipped with the boundary operator  $\partial \otimes \text{id} + (-1)^j \text{id} \otimes \partial$  on  $C_i(X) \otimes C_j(Y)$ .

and

$$\sigma : C_\bullet(\Sigma) \rightarrow C_{\bullet+1}(\Sigma)$$

the (“vertical”) maps induced by  $k_e$ ,  $l_e$  and  $\sigma_e$  on each subspace (3.3).

The three goals of this section are the following.

- (1) To give explicit formulas for the chain maps  $g : C_\bullet(\Sigma) \rightarrow C_\bullet(\Sigma')$  and  $h : C_\bullet(\Sigma') \rightarrow C_\bullet(\Sigma)$  respectively associated to  $\mathbf{g}$  and a homotopy inverse  $\mathbf{h}$ , in terms of  $k$ ,  $l$  and  $\sigma$ .
- (2) To describe an explicit chain homotopy  $\rho : C_\bullet(\Sigma) \rightarrow C_{\bullet+1}(\Sigma)$  between 1 and  $h \circ g$ , in terms of  $k$ ,  $l$  and  $\sigma$ .
- (3) To give an explicit formula for the boundary operator  $\partial'$  on the cellular chain complex  $C_\bullet(\Sigma')$  in terms of  $k$ ,  $l$ ,  $\sigma$ , the boundary operator  $\partial$  on the cellular chain complex  $C_\bullet(\Sigma)$ , and the vertical boundary operator  $(\partial')^{\text{vert}}$ .

The precise result is Proposition 4.2 below; it is a (homological) effective version of Proposition 2.2. Note that we do not give effective formulas for the homotopy between  $\mathbf{g} \circ \mathbf{h}$  and the identity as we will not use it. For the same reason we do not even name the homotopy between  $g \circ h$  and the identity.

The proof consists in explicating the construction of the homotopy equivalence between  $\Sigma$  and  $\Sigma'$ : both the stack of CW-complexes  $\Pi' : \Sigma' \rightarrow \Phi$  and the map  $\mathbf{g} : \Sigma \rightarrow \Sigma'$  are constructed by induction on the dimension of the cells. Suppose that

$$\mathbf{g}_{n-1} : \Pi^{-1}(\Phi^{(n-1)}) \rightarrow (\Pi')^{-1}(\Phi^{(n-1)})$$

has been constructed. Then over  $n$ -cells the stack  $\Sigma'$  is built from the composition of the maps

$$f'_n : \bigsqcup_{e \in E_n} F'_e \times \mathbb{S}^{n-1} \xrightarrow{\mathbf{l}_n} \bigsqcup_{e \in E_n} F_e \times \mathbb{S}^{n-1} \xrightarrow{f_n} \Pi^{-1}(\Phi^{(n-1)}) \xrightarrow{\mathbf{g}^{(n-1)}} (\Pi')^{-1}(\Phi^{(n-1)}) \quad (4.3)$$

where

$$\mathbf{l}_n = \bigsqcup_{e \in E_n} \mathbf{l}_e \times \text{id}_{\mathbb{S}^{n-1}}.$$

For each  $n$  we denote

$$X_n = \bigsqcup_{e \in E_n} F_e \times \mathbb{B}^n, \quad A_n = \bigsqcup_{e \in E_n} F_e \times \mathbb{S}^{n-1} \quad \text{and} \quad Y_n = \Pi^{-1}(\Phi^{(n)})$$

and

$$X'_n = \bigsqcup_{e \in E_n} F'_e \times \mathbb{B}^n, \quad A'_n = \bigsqcup_{e \in E_n} F'_e \times \mathbb{S}^{n-1} \quad \text{and} \quad Y'_n = (\Pi')^{-1}(\Phi^{(n)}).$$

By construction we have:

$$Y_n = Y_{n-1} \cup_{f_n} X_n \quad \text{and} \quad Y'_n = Y'_{n-1} \cup_{f'_n} X'_n.$$

Now the map  $\mathbf{g}$ , its homotopy inverse  $\mathbf{h}$  and an explicit homotopy

$$\mathbf{P} : [0, 1] \times \Sigma \rightarrow \Sigma, \quad \mathbf{P}(0, \cdot) = \text{Id}_\Sigma, \quad \mathbf{P}(1, \cdot) = \mathbf{h} \circ \mathbf{g},$$

are obtained as the direct limits of maps

$$\mathbf{g}_n : Y_n \rightarrow Y'_n, \quad \mathbf{h}_n : Y'_n \rightarrow Y_n \quad \text{and} \quad \mathbf{P}_n : [0, 1] \times Y_n \rightarrow Y_n$$

inductively constructed by considering the diagram

$$\begin{array}{ccccc} X_n & \xleftarrow{\iota} & A_n & \xrightarrow{f_n} & Y_{n-1} \\ \mathbf{k}_n \updownarrow \mathbf{l}_n & & \mathbf{k}_n \updownarrow \mathbf{l}_n & & \mathbf{g}_{n-1} \updownarrow \mathbf{h}_{n-1} \\ X'_n & \xleftarrow{\iota} & A'_n & \xrightarrow{f'_n} & Y_{n-1}. \end{array} \quad (4.4)$$

Here  $\iota$  denotes the inclusion map, we have  $f'_n = \mathbf{g}_{n-1} \circ f_n \circ \mathbf{l}_n$  and the maps  $\mathbf{l}_n \circ \mathbf{k}_n$  and  $\mathbf{h}_{n-1} \circ \mathbf{g}_{n-1}$  are homotopic to the identity. From these data the proposition proved in the next paragraph gives explicit formulas for the chain maps induced by  $\mathbf{g}_n$  and  $\mathbf{h}_n$ . This provides a (homological) effective version of [Geo08, Theorem 4.1.8].

**4.1. Explicit gluing.** Let  $X, X', A, A', Y, Y'$  be CW-complexes fitting into the following diagram of cellular maps

$$\begin{array}{ccccc} X & \xleftarrow{\iota} & A & \xrightarrow{f} & Y \\ \mathbf{k} \downarrow \uparrow \mathbf{l} & & \mathbf{k} \downarrow \uparrow \mathbf{l} & & \mathbf{g} \downarrow \uparrow \mathbf{h} \\ X' & \xleftarrow{\iota} & A' & \xrightarrow{f'} & Y' \end{array}$$

where the  $\iota$ 's denote inclusions,  $f' = \mathbf{g} \circ f \circ \mathbf{l}$  and the maps  $\mathbf{l} \circ \mathbf{k}$  and  $\mathbf{h} \circ \mathbf{g}$  are homotopic to the identity. Assume furthermore that any cell  $e$  of  $X$  that intersects  $A$  is contained in  $A$ , similarly for  $X'$  and  $A'$ , and that  $\mathbf{k}^{-1}(A') = A$  and  $\mathbf{l}^{-1}(A) = A'$ .

Let  $I = [0, 1]$ . Fix explicit homotopies

$$\Sigma: I \times X \rightarrow X \quad \text{and} \quad \mathbf{P}: I \times Y \rightarrow Y,$$

$$\Sigma(0, x) = x, \quad \Sigma(1, x) = \mathbf{l} \circ \mathbf{k}(x), \quad \mathbf{P}(0, y) = y, \quad \mathbf{P}(1, y) = \mathbf{h} \circ \mathbf{g}(y),$$

such that  $\Sigma(\cdot, A) \subset A$ .

The cellular chain complex  $C_\bullet(X \sqcup_f Y)$  can be naturally identified with

$$C_\bullet(X) \oplus C_\bullet(Y) / (\iota, -f)C_\bullet(A).$$

In the following we use the same letter for the maps and the induced maps on a cellular chain complex except that the chain maps are not bold faced.

For any cell  $e \subset X^{(n)}$  let  $[e] := e_*([\mathbb{B}^n, \mathbb{S}^{n-1}])$ . The homotopy  $\Sigma$  induces a map  $\sigma: C_\bullet(X) \rightarrow C_{\bullet+1}(X)$ , given by

$$\sigma([e]) = (\Sigma)_*([I, \partial I] \otimes [e]), \quad \text{for any cell } e \subset X.$$

The map  $\sigma$  is then a chain homotopy between  $1$  and  $l \circ k$ , i.e.

$$l \circ k - 1 = \partial\sigma + \sigma\partial.$$

The same discussion applies to  $Y$ ; we denote by  $\rho$  the corresponding chain homotopy.

Decompose the cellular chain complex  $C_\bullet(X)$  as

$$C_\bullet(X) = \sum_{e \notin A} \mathbb{Z}[e] \oplus \iota_* C_\bullet(A)$$

and let  $1_A: C_\bullet(X) \rightarrow \iota_* C_\bullet(A)$  be the projection onto the second component. Define analogously  $1_{A'}: C_\bullet(X') \rightarrow \iota'_* C_\bullet(A')$ . It follows from our assumptions  $k^{-1}(A') \subset A$  and  $l^{-1}(A) \subset A'$  that

$$k \circ 1_A = 1_{A'} \circ k \quad \text{and} \quad l \circ 1_{A'} = 1_A \circ l.$$

In order to lighten the notations we will suppress  $\iota_*$  and identify  $C_\bullet(A)$  with a sub-complex of  $C_\bullet(X)$ .

**Proposition 4.1** (Explicit gluing Lemma). *1. There exist cellular maps*

$$\tilde{\mathbf{g}}: X \cup_f Y \rightarrow X' \cup_{f'} Y' \quad \text{and} \quad \tilde{\mathbf{h}}: X' \cup_{f'} Y' \rightarrow X \cup_f Y$$

such that  $\tilde{\mathbf{g}}|_Y = \mathbf{g}$ ,  $\tilde{\mathbf{h}}|_Y = \mathbf{h}$  and  $\tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}$  is homotopy equivalent to the identity.

*2. The chain maps  $\tilde{g}: C_\bullet(X \cup_f Y) \rightarrow C_\bullet(X' \cup_{f'} Y')$  and  $\tilde{h}: C_\bullet(X' \cup_{f'} Y') \rightarrow C_\bullet(X \cup_f Y)$  respectively associated to  $\tilde{\mathbf{g}}$  and  $\tilde{\mathbf{h}}$  are induced by maps defined on  $C_\bullet(X) \oplus C_\bullet(Y)$  and  $C_\bullet(X') \oplus C_\bullet(Y')$  by the following formulas*

$$\tilde{g} = (k \circ (1 - 1_A) + g \circ f \circ 1_A - g \circ f \circ \sigma \circ (1_A \partial - \partial 1_A)) \oplus g$$

and

$$\tilde{h} = (l \circ (1 - 1_{A'}) + h \circ f' \circ 1_{A'} + \rho \circ f \circ l \circ (1_{A'} \partial - \partial 1_{A'})) \oplus h,$$

where  $g$  and  $h$  are the chain maps respectively induced by  $\mathbf{g}$  and  $\mathbf{h}$ .

3. There exists a homotopy map

$$\tilde{\mathbf{P}}: I \times (X \cup_f Y) \rightarrow X \cup_f Y$$

such that  $\tilde{\mathbf{P}}(0, z) = z$  and  $\tilde{\mathbf{P}}(1, z) = \tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}(z)$ .

4. The chain homotopy map  $\tilde{\rho}(c) = \tilde{\mathbf{P}}_*([I, \partial I] \otimes c)$  associated to  $\tilde{\mathbf{P}}$  is induced by a map defined on  $C_\bullet(X) \oplus C_\bullet(Y)$  by the following formula

$$\tilde{\rho} = (\sigma \circ (1 - 1_A) + \rho \circ f \circ 1_A - \rho \circ f \circ \sigma \circ (1_A \partial - \partial 1_A)) \oplus \rho.$$

The proof of Proposition 4.1 consists of explicating the construction of the maps  $\tilde{\mathbf{h}}$  and  $\tilde{\mathbf{g}}$ . We postpone it until Section 5. The construction is quite technical so the reader may want to skip Section 5 on a first reading. It is instructive to check that our formulas work at the level of homology groups.

4.2. **Rebuilding of  $\Sigma$ .** Starting from

$$Y_0 = \sqcup_{e \in E_0} F_e, \quad Y'_0 = \sqcup_{e \in E_0} F'_e, \quad \mathbf{g}_0 = \mathbf{k}_0, \quad \mathbf{h}_0 = \mathbf{l}_0, \quad \text{and} \quad \mathbf{P}_0 = \Sigma_0,$$

we inductively apply Proposition 4.1 to (4.4) and construct the desired extensions  $\mathbf{g}_n$ ,  $\mathbf{h}_n$  and  $\mathbf{P}_n$  of  $\mathbf{g}_{n-1}$ ,  $\mathbf{h}_{n-1}$  and  $\mathbf{P}_{n-1}$ .

We identify

$$C_\bullet(X_n) = \bigoplus_{e \in E_n} C_\bullet(F_e) \otimes C_\bullet(\mathbb{B}^n)$$

and let

$$[\mathbb{B}^n, \mathbb{S}^n] \in H_n(\mathbb{B}^n, \mathbb{S}^n) = C_n(\mathbb{B}^n)$$

be the relative fundamental class that generates the  $\mathbb{Z}$ -module of rank one  $C_n(\mathbb{B}^n)$ . Note that given  $c \in C_\bullet(F_e)$  we have:

$$1_{A_n}(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) = 0 \quad \text{and} \quad (1_{A_n} \circ \partial - \partial \circ 1_{A_n})(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) = 1_{A_n} \partial(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) = (-1)^{\dim c} c \otimes [\mathbb{S}^{n-1}]$$

where the last equality follows from Lemma 3.3. There are analogous formulas for  $c' \in C_\bullet(F'_e)$  replacing  $A_n$  by  $A'_n$ .

Denoting by  $g_n$  and  $h_n$  the chain maps respectively associated to  $\mathbf{g}_n$  and  $\mathbf{h}_n$  and by  $\sigma_n$  and  $\rho_n$  the chain homotopies respectively associated to  $\Sigma_n$  and  $\mathbf{P}_n$ , Proposition 4.1 gives

$$\begin{aligned} g_n(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) &= k_n(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) - (-1)^{\dim c} g_{n-1} \circ f \circ \sigma_n(c \otimes [\mathbb{S}^{n-1}]), \\ h_n(c' \otimes [\mathbb{B}^n, \mathbb{S}^n]) &= l_n(c' \otimes [\mathbb{B}^n, \mathbb{S}^n]) + (-1)^{\dim c} \rho_{n-1} \circ f \circ l_{n-1}(c' \otimes [\mathbb{S}^{n-1}]) \end{aligned}$$

and

$$\rho_n(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) = \sigma_n(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) - (-1)^{\dim c} \rho_{n-1} \circ f \circ \sigma_n(c \otimes [\mathbb{S}^{n-1}]).$$

We can simplify these formulas by identifying

$$C_\bullet(Y_n) = (C_\bullet(X_n) \oplus C_\bullet(Y_{n-1})) / (1, -f_*) C_\bullet(A_n),$$

with

$$\bigoplus_{e \in E_n} C_\bullet(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^{n-1}) \subset F^n C_\bullet(\Sigma).$$

The map  $\mathbf{g}: \Sigma \rightarrow \Sigma'$  induces chain maps  $g_n: F^n C_\bullet(\Sigma) \rightarrow F^n C_\bullet(\Sigma')$ , and by Lemma 3.3 the chain maps  $g_n$  are inductively defined as follows:

- The restriction of  $g_n$  to  $F^{n-1} C_\bullet(\Sigma)$  is equal to  $g_{n-1}$ .
- If  $e$  is a  $n$ -cell of  $\Phi$ , the restriction of  $g_n$  to  $C_\bullet(F_e) \otimes H_n(\mathbb{B}^n, \mathbb{S}^n)$  is given by

$$g_n(c \otimes [\mathbb{B}^n, \mathbb{S}^n]) = k_e(c) \otimes [\mathbb{B}^n, \mathbb{S}^{n-1}] + g_{n-1} \circ \partial^{\text{hor}}(\sigma_e(c) \otimes [\mathbb{B}^n, \mathbb{S}^{n-1}]) \quad (c \in C_\bullet(F_e)). \quad (4.5)$$



The sign changed because  $\dim \sigma_e(c) = \dim c + 1$ , so  $(-1)^{\dim c} f \circ \sigma_n(c \otimes [\mathbb{S}^{n-1}]) = -\partial^{\text{hor}}(\sigma_e(c) \otimes [\mathbb{B}^n, \mathbb{S}^{n-1}])$ , by Lemma 3.3. By induction, we get:

$$g = k \circ \left( \sum_{i=0}^n (\partial^{\text{hor}} \circ \sigma)^i \right). \quad (4.6)$$

Note that the map  $\sigma$  preserves the filtration  $F^n C_\bullet(\Sigma)$  and that the map  $\partial^{\text{hor}}$  maps  $F^n C_\bullet(\Sigma)$  to  $F^{n-1} C_{\bullet-1}(\Sigma)$ . It follows that  $(\partial^{\text{hor}} \circ \sigma)^i$  vanishes on  $C_n(\Sigma)$  for  $i \geq n + 1$ .

We similarly get formulas for  $h$  and  $\rho$ . To sum up we get:

**Proposition 4.2** (Effective rebuilding Lemma). *We have*

- (1) *The rebuilding lemma yields a CW-stack  $\Pi' : \Sigma' \rightarrow \Phi$ , cellular maps  $\mathbf{g} : \Sigma \rightarrow \Sigma'$ ,  $\mathbf{h} : \Sigma' \rightarrow \Sigma$  and a homotopy  $\mathbf{P} : I \times \Sigma \rightarrow \Sigma$  such that*

$$\mathbf{P}(0, \cdot) = \text{Id}_\Sigma \quad \text{and} \quad \mathbf{P}(1, \cdot) = \mathbf{h} \circ \mathbf{g}.$$

- (2) *The chain maps  $g : C_\bullet(\Sigma) \rightarrow C_\bullet(\Sigma')$ ,  $h : C_\bullet(\Sigma') \rightarrow C_\bullet(\Sigma)$  and  $\rho : C_\bullet(\Sigma) \rightarrow C_\bullet(\Sigma)$  respectively associated to  $\mathbf{g}$ ,  $\mathbf{h}$  and  $\mathbf{P}$  are respectively given by*

$$g = k \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right) \quad \text{on } C_\bullet(\Sigma), \quad (4.7)$$

$$h = \left( \sum_{i=0}^{\infty} (\sigma \circ \partial^{\text{hor}})^i \right) \circ l \quad \text{on } C_\bullet(\Sigma'), \quad (4.8)$$

and

$$\rho = \sigma \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right) \quad \text{on } C_\bullet(\Sigma). \quad (4.9)$$

- (3) *The formula for the boundary operator  $\partial'$  on the cellular chain complex  $C_\bullet(\Sigma')$  of the rebuilt CW-complex  $\Sigma'$  is given on  $C_\bullet(\Sigma')$  by*

$$\partial' = (\partial')^{\text{vert}} + k \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right) \circ \partial^{\text{hor}} \circ l. \quad (4.10)$$

*Proof.* It only remains to prove (3). Lemma 3.3 applies to the stack  $\Pi' : \Sigma' \rightarrow \Phi$  so that if  $e$  is an  $n$ -cell of  $\Phi$  and  $c \in C_\bullet(F_e)$  we have:

$$((\partial')^{\text{hor}})(c \otimes e) = (-1)^{\dim c} f'_e(c \otimes [\mathbb{S}^{n-1}]). \quad (4.11)$$

On the other hand it follows from the construction of  $\Pi' : \Sigma' \rightarrow \Phi$  that

$$f'_e(c \otimes [\mathbb{S}^{n-1}]) = g_{n-1}(f_e(l_e(c) \otimes [\mathbb{S}^{n-1}])). \quad (4.12)$$

We conclude that

$$\begin{aligned} ((\partial')^{\text{hor}})(c \otimes e) &= (-1)^{\dim c} f'_e(c \otimes [\mathbb{S}^{n-1}]) \\ &= (-1)^{\dim c} g_{n-1}(f_e(l_e(c) \otimes [\mathbb{S}^{n-1}])) \\ &= g_{n-1} \circ \partial^{\text{hor}} \circ l(c \otimes e), \end{aligned}$$

and (4.10) follows from (4.7).  $\square$

## 5. PROOF OF PROPOSITION 4.1

Let  $X$  be a CW-complex and let  $A \subset X$  be a sub-complex. Then  $(\{0\} \times X) \cup (I \times A)$  is a strong deformation retract of  $I \times X$ ; see e.g. [Geo08, 1.3.15]. We refine this property in the following lemma.

**Lemma 5.1.** *There exists a cellular map*

$$p: I \times X \rightarrow \{0\} \times X \cup I \times A$$

such that for all  $x \in X$ ,  $a \in A$  and  $s \in ]0, 1]$ ,

- (1)  $p(0, x) = (0, x) \in \{0\} \times X$ ;
- (2)  $p(s, x) = (p^1(s, x), p^2(s, x)) \in [0, s] \times X$ ;
- (3)  $p(s, a) = (s, a) \in I \times A$

and the following formulas hold for any chain  $c \in C_\bullet(X)$ :

$$\begin{aligned} p_*([I, \partial I] \otimes c) &= [I, \partial I] \otimes 1_A c, \\ \delta_*([I, \partial I] \otimes c) &= [I, \partial I] \otimes (1 - 1_A)c, \end{aligned}$$

where

$$\delta: I \times X \rightarrow I \times X, \quad \delta(s, x) = (s - p^1(s, x), p^2(s, x)).$$

*Proof.* We define the map  $p$  cell by cell, starting from 0-dimensional cells and attaching any available cell of the lowest dimension as pictured on Figure 1.




FIGURE 1. The map  $p$

The proof reduces to the following statement. Let  $Z$  be a CW-sub-complex of  $X$  of dimension at most  $n$ , and let  $e: \mathbb{B}^n \rightarrow X$  be an  $n$  cell with  $e(\mathbb{S}^{n-1}) \subset Z^{(n-1)}$ . Suppose we are given a cellular map

$$p: I \times Z \rightarrow \{0\} \times Z \cup I \times (A \cap Z)$$

satisfying all the desired properties. Put  $Z' = \mathbb{B}^n \sqcup_e Z$ . We shall construct an extension

$$p: I \times Z' \rightarrow \{0\} \times Z' \cup I \times (A \cap Z')$$


of  $p$ , satisfying all the required properties. To do so it is enough to define  $p$  on  $I \times \mathbb{B}^n$  so that

$$\forall s \in I, \forall x \in \mathbb{S}^{n-1}, \quad p(s, x) = p(s, e(x)).$$

Now if  $e \subset A$  we put

$$p(s, e(x)) = (s, e(x)) \in I \times A. \tag{5.1}$$

Otherwise  $p(s, e(x))$  is already defined for all  $x \in \mathbb{S}^{n-1}$  so we put  $p(s, x) = p(s, e(x))$  for  $x \in \mathbb{S}^{n-1}$ ,  $p(0, x) = (0, x)$  for  $x \in \mathbb{B}^n$  and extend the map to  $I \times \mathbb{B}^n$  using the homotopy extension property

FIGURE 2. The map  $p(1, -)$ 

for the pair  $\{0\} \times \mathbb{B}^n \cup I \times \mathbb{S}^{n-1}$  (cf. [Geo08, 1.3.15]). By replacing  $p^1(s, x)$  with  $\max\{s, p^1(s, x)\}$  we can ensure that the second condition is satisfied.

By construction, the maps  $p$  and  $\delta$  are cellular and both  $p(I \times e)$  and  $\delta(I \times e)$  are contained in  $I \times e$ . Write  $[e] = e_*([\mathbb{B}^n, \mathbb{S}^n])$  and  $[\partial e] = e_*([\mathbb{S}^{n-1}])$ . We check the formulas for  $p_*$  and  $\delta_*$  on  $[I, \partial I] \otimes [e]$ . When  $e \subset A$ , the formula for  $p_*$  follows from the definition (see (5.1)). If  $e \not\subset A$  then the  $(n+1)$ -cell  $I \times e$  is mapped by  $p$  into

$$\{0\} \times (Z')^{(n)} \cup I \times Z^{(n-1)}$$


which is contained in the  $n$ -skeleton of  $I \times Z'$ . It follows that  $p_*([I, \partial I] \otimes [e]) = 0$ . This proves the formula for  $p_*$ .

Using that  $p_* \circ \partial = \partial \circ p_*$ , the formula for  $p_*$  now implies that

$$p_*([1] \otimes c) = [0] \otimes (1 - 1_A)c + [I, \partial I] \otimes (1_A \partial - \partial 1_A)c + [1] \otimes 1_A c \quad (5.2)$$

and consequently that

$$\delta_*([1] \otimes c) = [1] \otimes (1 - 1_A)c - [I, \partial I] \otimes (1_A \partial - \partial 1_A)c + [0] \otimes 1_A c. \quad (5.3)$$

FIGURE 3. The map  $\delta$ 

We finally derive the general formula for  $\delta_*([I, \partial I] \otimes [e])$  from (5.3): we know that

$$\delta_*([I, \partial I] \otimes [e]) \in C_n(I \times Z') = C_0(I) \otimes C_{n+1}(Z') \oplus C_1(I) \otimes C_n(Z') = C_1(I) \otimes C_n(Z'),$$

since  $Z'$  is  $n$ -dimensional. As  $\delta(I \times e) \subset I \times e$  we must have

$$\delta_*([I, \partial I] \otimes [e]) = \alpha [I, \partial I] \otimes [e]$$

for some  $\alpha \in \mathbb{Z}$ . To find  $\alpha$  we compute  $\partial\delta_*([I, \partial I] \otimes [e])$  using (5.3) and, by induction, the formula for  $\delta_*$  on  $[I, \partial I] \otimes [\partial e]$ :

$$\begin{aligned} \partial\delta_*([I, \partial I] \otimes [e]) &= \delta_*(\partial([I, \partial I] \otimes [e])) \\ &= \delta_*([1] \otimes [e]) - \delta_*([0] \otimes [e]) - \delta_*([I, \partial I] \otimes [\partial e]) \\ &= \partial([I, \partial I] \otimes (1 - 1_A)[e]). \end{aligned}$$

Hence  $\alpha = 0$  if  $e \subset A$  and  $\alpha = 1$  otherwise. This proves the formula for  $\delta_*([I, \partial I] \otimes [e])$ .  $\square$

We now come back to the notations of Proposition 4.1.

Let

$$p: I \times (X \cup_\iota A) \rightarrow (\{0\} \times X) \cup (I \times A)$$

be the map afforded by Lemma 5.1 applied to the pair  $(X, A)$ . We write

$$p_s(x) = p(s, x), \quad (s \in I, x \in X).$$

We have

$$C_\bullet(\{0\} \times X \cup I \times A) = [0] \otimes C_\bullet(X) + [I, \partial I] \otimes C_\bullet(A) + [1] \otimes C_\bullet(A)$$

and, according to Lemma 5.1 and Equation (5.2), the map  $p$  can be chosen so that

$$\begin{aligned} p_*([I, \partial I] \otimes c) &= [I, \partial I] \otimes 1_A c, \\ (p_1)_*(c) &= [0] \otimes (1 - 1_A)c + [1] \otimes 1_A c + [I, \partial I] \otimes (1_A \partial - \partial 1_A)c \\ &= [0] \otimes c + \partial[I, \partial I] \otimes 1_A c + [I, \partial I] \otimes (1_A \partial - \partial 1_A)c. \end{aligned}$$

As in Lemma 5.1 we write  $p_s^1: I \times X \rightarrow I$  and  $p_s^2: I \times X \rightarrow X$  for the coordinates of  $p_s$ . Replacing  $p_s^1(x)$  with  $\min\{s, p_s^1(x)\}$  we may assume that  $p_s^1(x) \leq s$  for  $s \in I$  (actually the construction in Lemma 5.1 already gives such  $p$ ).

Similarly, using Lemma 5.1 we choose a map

$$q: I \times (X' \cup_\iota A') \rightarrow (\{0\} \times X') \cup (I \times A').$$

For the readers' convenience we spell out the relevant properties of  $q$ . For all  $x' \in X'$ ,  $a' \in A'$  and  $s \in I$

$$q(0, x') = (0, x'), \quad q(s, a') = (s, a')$$

and

$$\begin{aligned} q_*([I, \partial I] \otimes c) &= [I, \partial I] \otimes 1_{A'} c, \\ (q_1)_*(c) &= [0] \otimes (1 - 1_{A'})c + [1] \otimes 1_{A'} c + [I, \partial I] \otimes (1_{A'} \partial - \partial 1_{A'})c \\ &= [0] \otimes c + \partial[I, \partial I] \otimes 1_{A'} c + [I, \partial I] \otimes (1_{A'} \partial - \partial 1_{A'})c. \end{aligned}$$

We define the map

$$\tilde{\mathbf{g}}: X \cup_f Y \rightarrow X' \cup_{f'} Y'$$

to be equal to  $\mathbf{g}$  on  $Y$  and for  $x \in X$  by the formula

$$\tilde{\mathbf{g}}(x) = \begin{cases} \mathbf{k} \circ p_1^2(x) & \text{if } p_1(x) \in \{0\} \times X, \\ \mathbf{g} \circ f \circ \Sigma(1 - p_1^1(x), p_1^2(x)) & \text{if } p_1(x) \in I \times A. \end{cases}$$

To check that  $\tilde{\mathbf{g}}$  is a well defined continuous map it is enough to verify that the partial formulas coincide on

$$\{x \in X : p_1(x) \in \{0\} \times X\}$$

and that for all  $a \in A$  we have  $\tilde{\mathbf{g}}(a) = \mathbf{g} \circ f(a)$ . If  $p_1(x) \in \{0\} \times A$  we have

$$\mathbf{k} \circ p_1^2(x) = f' \circ \mathbf{k} \circ p_1^2(x) \quad \text{in } X' \cup_{f'} Y'$$

and

$$f' \circ \mathbf{k} \circ p_1^2(x) = \mathbf{g} \circ f \circ \mathbf{l} \circ \mathbf{k} \circ p_1^2(x) = \mathbf{g} \circ f \circ \Sigma(1, p_1^2(x)) = \mathbf{g} \circ f \circ \Sigma(1 - p_1^1(x), p_1^2(x)).$$

For  $a \in A$  we have  $p_1(a) = (1, a)$  so

$$\tilde{\mathbf{g}}(a) = \mathbf{g} \circ f \circ \Sigma(0, a) = \mathbf{g} \circ f(a),$$

as desired.

Analogously we define the map

$$\tilde{\mathbf{h}} : X' \cup_{f'} Y' \rightarrow X \cup_f Y$$

to be equal to  $\mathbf{h}$  on  $Y'$  and for  $x \in X'$  by the formula

$$\tilde{\mathbf{h}}(x) = \begin{cases} \mathbf{l} \circ q_1^2(x) & \text{if } q_1(x) \in \{0\} \times X', \\ \Sigma(q_1^1(x), f \circ \mathbf{l} \circ q_1^2(x)) & \text{if } q_1(x) \in I \times A'. \end{cases}$$

The verification that  $\tilde{\mathbf{h}}$  is well defined and continuous is completely analogous to what we did for  $\tilde{\mathbf{g}}$ .

The formulas for the chain maps  $\tilde{g}$  and  $\tilde{h}$  respectively associated to  $\tilde{\mathbf{g}}$  and  $\tilde{\mathbf{h}}$  now follow from those for  $(p_1)_*$  and  $(q_1)_*$ ; more specifically the identity (5.2).

It remains to construct an explicit homotopy between the identity map and  $\tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}$ , which is the main content of the proposition. On  $X$  we have

$$\tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}(x) = \begin{cases} \mathbf{l} \circ q_1^2 \circ \mathbf{k} \circ p_1^2(x) & \text{if } p_1(x) \in \{0\} \times X \text{ and } q_1 \circ \mathbf{k} \circ p_1^2(x) \in \{0\} \times X', \\ \mathbf{P}(q_1^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \mathbf{l} \circ q_1^2 \circ \mathbf{k} \circ p_1^2(x)) & \text{if } p_1(x) \in \{0\} \times X \text{ and } q_1 \circ \mathbf{k} \circ p_1^2(x) \in I \times A', \\ \mathbf{h} \circ \mathbf{g} \circ f \circ \Sigma(1 - p_1^1(x), p_1^2(x)) & \text{if } p_1(x) \in I \times A, \end{cases}$$

while on  $Y$  we have

$$\tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}(y) = \mathbf{h} \circ \mathbf{g}(y), \quad \text{for all } y \in Y.$$

Note that for  $p_1(x) \in I \times A$ , we have  $\mathbf{k} \circ p_1^2(x) \in A'$  so that  $q_1(\mathbf{k} \circ p_1^2(x)) = (1, \mathbf{k} \circ p_1^2(x))$  and therefore

$$\begin{aligned} \mathbf{h} \circ \mathbf{g} \circ f \circ \Sigma(1 - p_1^1(x), p_1^2(x)) &= \mathbf{P}(1, f \circ \Sigma(1 - p_1^1(x), p_1^2(x))) \\ &= \mathbf{P}(q_1^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \Sigma(1 - p_1^1(x), p_1^2(x))). \end{aligned}$$

This already suggests the rough form of the homotopy between 1 and  $\tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}$ . We want to construct a map

$$\tilde{\mathbf{P}} : I \times (X \cup_f Y) \rightarrow X \cup_f Y \quad \text{with} \quad \tilde{\mathbf{P}}(0, z) = z \quad \text{and} \quad \tilde{\mathbf{P}}(1, z) = \tilde{\mathbf{h}} \circ \tilde{\mathbf{g}}(z).$$

We define it piece by piece starting with  $[1/2, 1] \times (X \cup_f Y)$ . Define subsets

$$\begin{aligned} \mathcal{C}_1 &:= \{(s, x) \in [0, 1/2] \times X \mid p_1(x) \in \{0\} \times X \text{ and } q_{2s} \circ \mathbf{k} \circ p_1^2(x) \in \{0\} \times X'\}, \\ \mathcal{C}_2 &:= \{(s, x) \in [0, 1/2] \times X \mid p_1(x) \in \{0\} \times X \text{ and } q_{2s} \circ \mathbf{k} \circ p_1^2(x) \in I \times A'\}, \\ \mathcal{C}_3 &:= \{(s, x) \in [0, 1/2] \times X \mid p_1(x) \in I \times A\}. \end{aligned}$$

For all  $s \in [0, 1/2]$  we set

$$\tilde{\mathbf{P}}(1/2 + s, y) = \mathbf{P}(2s, y) \quad \text{if } y \in Y$$

and define

$$\tilde{\mathbf{P}}(1/2 + s, x) = \begin{cases} \mathbf{l} \circ q_{2s}^2 \circ \mathbf{k} \circ p_1^2(x) & \text{if } (s, x) \in \mathcal{C}_1, \\ \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \mathbf{l} \circ q_{2s}^2 \circ \mathbf{k} \circ p_1^2(x)) & \text{if } (s, x) \in \mathcal{C}_2, \\ \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \Sigma(1 - p_1^1(x), p_1^2(x))) & \text{if } (s, x) \in \mathcal{C}_3, \end{cases}$$

for all  $x \in X$ . Let's check that the partial maps agree on the common boundaries.

- The common boundary to  $\mathcal{C}_1$  and  $\mathcal{C}_2$  is the set

$$\{(1/2 + s, x) : p_1(x) \in \{0\} \times X \text{ and } q_{2s} \circ \mathbf{k} \circ p_1^2(x) \in \{0\} \times A'\}.$$

For  $(1/2 + s, x)$  therein, we have  $\mathbf{l} \circ q_{2s}^2 \circ \mathbf{k} \circ p_1^2(x) \in A$  so that

$$\mathbf{l} \circ q_{2s} \circ \mathbf{k} \circ p_1^2(x) = f \circ \mathbf{l} \circ q_{2s} \circ \mathbf{k} \circ p_1^2(x) = \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \mathbf{l} \circ q_{2s} \circ \mathbf{k} \circ p_1^2(x)).$$

- The common boundary of  $\mathcal{C}_2$  and  $\mathcal{C}_3$  is

$$\{(1/2 + s, x) : p_1(x) \in \{0\} \times A \text{ and } q_{2s} \circ \mathbf{k} \circ p_1^2(x) \in I \times A'\}.$$

Note that  $p_1(x) \in \{0\} \times A$  forces  $\mathbf{k} \circ p_1^2(x) \in A'$  so that  $q_{2s} \circ \mathbf{k} \circ p_1^2(x) = (2s, \mathbf{k} \circ p_1^2(x))$ . We then have

$$\begin{aligned} \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \mathbf{l} \circ q_{2s}^2 \circ \mathbf{k} \circ p_1^2(x)) &= \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \mathbf{l} \circ \mathbf{k} \circ p_1^2(x)) \\ &= \mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(x), f \circ \Sigma(1 - p_1^1(x), p_1^2(x))). \end{aligned}$$

- The common boundary of  $\mathcal{C}_1$  and  $\mathcal{C}_3$  is

$$\{(1/2, x) : p_1(x) \in \{0\} \times A\}.$$

It is contained in  $\mathcal{C}_1 \cap \mathcal{C}_2 \cap \mathcal{C}_3$  so that this case follows from the two previous ones.

- Finally  $Y$  intersects non-trivially only  $\mathcal{C}_3$  and their common subset is

$$\{(1/2 + s, a) : a \in A\}.$$

There we have

$$q_{2s}^1 \circ \mathbf{k} \circ p_1^2(a) = 2s, \quad p_1^1(a) = 1 \quad \text{and} \quad p_1^2(a) = a$$

so that

$$\mathbf{P}(q_{2s}^1 \circ \mathbf{k} \circ p_1^2(a), f \circ \Sigma(1 - p_1^1(a), p_1^2(a))) = \mathbf{P}(2s, f(a)).$$

Now that we have a well defined continuous map  $\tilde{\mathbf{P}}(s, \cdot)$  for  $s \in [1/2, 1]$ , we proceed to define  $\tilde{\mathbf{P}}$  for  $s \in [0, 1/2]$ . First note that  $\tilde{\mathbf{P}}(1/2, \cdot)$  can be more simply defined by

$$\tilde{\mathbf{P}}(1/2, y) = y, \quad \text{if } y \in Y, \quad \text{and} \quad \tilde{\mathbf{P}}(1/2, x) = \Sigma(1 - p_1^1(x), p_1^2(x)), \quad \text{if } x \in X.$$

The last expression is indeed equal to

$$\mathbf{l} \circ \mathbf{k} \circ p_1^2(x), \quad \text{if } p_1(x) \in \{0\} \times X, \quad \text{and} \quad f \circ \Sigma(1 - p_1^1(x), p_1^2(x)), \quad \text{if } p_1(x) \in I \times A.$$

For  $s \in [0, 1/2]$  we then set

$$\tilde{\mathbf{P}}(s, y) = y, \quad \text{if } y \in Y, \quad \text{and} \quad \tilde{\mathbf{P}}(s, x) = \Sigma(2s - p_{2s}^1(x), p_{2s}^2(x)), \quad \text{if } x \in X.$$

The total map  $\tilde{\mathbf{P}}: I \times (X \cup_f Y) \rightarrow X \cup_f Y$  is continuous by construction. It remains to compute

$$\tilde{\rho}(c) = \tilde{\mathbf{P}}_*([I, \partial I] \otimes c) \quad \text{for all } c \in C_\bullet(X \cup_f Y).$$

Let us refine the CW-complex structure of  $I$  by taking the first barycentric subdivision of the original one. There are now two 1-dimensional cells  $I_1 = (0, 1/2)$  and  $I_2 = (1/2, 1)$  such that  $[I, \partial I] = [I_1, \partial I_1] + [I_2, \partial I_2]$ . It follows that

$$\tilde{\mathbf{P}}_*([I, \partial I] \otimes c) = \tilde{\mathbf{P}}_*([I_1, \partial I_1] \otimes c) + \tilde{\mathbf{P}}_*([I_2, \partial I_2] \otimes c).$$

Given  $c \in C_\bullet(Y)$  we have

$$\tilde{\mathbf{P}}_*([I_1, \partial I_1] \otimes c) = 0 \quad \text{and} \quad \tilde{\mathbf{P}}_*([I_2, \partial I_2] \otimes c) = \rho(c),$$

whence  $\tilde{\mathbf{P}}_*([I, \partial I] \otimes c) = \rho(c)$ .

Now let  $c \in C_\bullet(X)$ . We first compute  $\tilde{\mathbf{P}}_*([I_2, \partial I_2] \otimes c)$ . Recall that it follows from (5.2) that

$$(p_1)_*(c) = [0] \otimes (1 - 1_A)c + [I, \partial I] \otimes (1_A \partial - \partial 1_A)c + [1] \otimes 1_{AC}.$$

From this and the definition of  $\tilde{\mathbf{P}}$  on  $[1/2, 1] \times (X \cup_f Y)$  we get

$$\begin{aligned} \tilde{\mathbf{P}}_*([I_2, \partial I_2] \otimes c) &= \alpha_* \beta_*([I, \partial I] \otimes (1 - 1_A)c) \\ &\quad + \mathbf{P}_*(\gamma_*([I, \partial I] \otimes [I, \partial I] \otimes (1_A \partial - \partial 1_A)c)) \\ &\quad + \mathbf{P}_*([I, \partial I] \otimes f \circ 1_A(c)), \end{aligned} \quad (5.4)$$

where

$\alpha: (\{0\} \times X') \cup (I \times A') \rightarrow X \cup_f Y$ ,  $\beta: I \times X \rightarrow (\{0\} \times X') \cup (I \times A')$  and  $\gamma: I \times I \times A \rightarrow I \times Y$  are respectively given by

$$\alpha(s, x) = \begin{cases} \mathbf{1}(x), & \text{if } (s, x) \in \{0\} \times X', \\ \mathbf{P}(s, f \circ \mathbf{1}(x)), & \text{if } (s, x) \in I \times A', \end{cases} \quad \beta(s, x) = q_s \circ \mathbf{k}(x)$$

and

$$\gamma(s, t, a) = (q_s^1 \circ \mathbf{k}(a), f \circ \Sigma(1 - t, a)) = (s, f \circ \Sigma(1 - t, a)).$$

Recall the notations

$$\sigma(u) = \Sigma_*([I, \partial I] \otimes u) \quad \text{and} \quad \rho(u) = \mathbf{P}_*([I, \partial I] \otimes u)$$

for respectively  $u \in C_\bullet(X)$  and  $C_\bullet(Y)$ .

For all  $u \in C_\bullet(X)$  we have

$$\beta_*([I, \partial I] \otimes u) = q_*([I, \partial I] \otimes k(u)) = [I, \partial I] \otimes 1_{A'} \circ k(u)$$

and

$$\gamma_*([I, \partial I] \otimes [I, \partial I] \otimes u) = -[I, \partial I] \otimes f \circ \sigma(u).$$

Note that it follows from the formula for  $\beta_*$  that

$$\beta_*([I, \partial I] \otimes (1 - 1_A)(c)) = [I, \partial I] \otimes 1_{A'} \circ k \circ (1 - 1_A)(c) = [I, \partial I] \otimes k \circ 1_A \circ (1 - 1_A)(c) = 0.$$

So we do not need to compute  $\alpha_*$  in (5.4) to conclude that

$$\tilde{\mathbf{P}}_*([I_2, \partial I_2] \otimes c) = -\rho \circ f \circ \sigma \circ (1_A \partial - \partial 1_A)(c) + \rho \circ f \circ 1_A(c).$$

We finally compute  $\tilde{\mathbf{P}}_*([I_1, \partial I_1] \otimes c)$ . Let

$$\delta: I \times X \rightarrow I \times X, \quad \delta(s, x) = (s - p_s^1(x), p_s^2(x)).$$

By Lemma 5.1, we can choose  $p$  so that  $\delta$  is cellular and

$$\delta_*([I, \partial I] \otimes c) = [I, \partial I] \otimes (1 - 1_A)c.$$

It then follows from the definition of  $\tilde{\mathbf{P}}$  on  $[0, 1/2] \times (X \cup_f Y)$  that

$$\tilde{\mathbf{P}}_*([I_1, \partial I_1] \otimes c) = \Sigma_* \circ \delta_*([I, \partial I] \otimes c) = \sigma \circ (1 - 1_A)(c).$$

We conclude that

$$\tilde{\rho} = (\sigma \circ (1 - 1_A) - \rho \circ f \circ \sigma \circ (1_A \partial - \partial 1_A) + \rho \circ f \circ 1_A) \oplus \rho.$$

This finishes the proof of Proposition 4.1.  $\square$

## 6. QUALITY OF REBUILDING FOR NILPOTENT GROUPS (PROOF OF THEOREM E)

The goal of the section is to prove the following:

**Theorem 6.1** (Theorem E). *Let  $\Lambda$  be a finitely generated torsion-free nilpotent group. If  $Y_0$  is a compact  $K(\Lambda, 1)$  space, there exists a constant  $\kappa \geq 1$  such that for every finite index subgroup  $\Lambda_1 \leq \Lambda$ , the cover  $Y_1 = \Lambda_1 \backslash \tilde{Y}_0$  admits a  $\alpha$ -rebuilding  $(Y_1, Y'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  of quality  $([\Lambda : \Lambda_1], \kappa)$  for every  $\alpha$ .*

**6.1. Generalities on rebuildings and quality.** Let  $\alpha \in \mathbb{N}$  and let  $X$  be a CW-complex with finite  $\alpha$ -skeleton. Recall (Definition 1 of the introduction) that a  $\alpha$ -**rebuilding** of  $X$  is a collection  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$  that consists of a CW-complex with finite  $\alpha$ -skeleton  $X'$ , two cellular maps between the  $\alpha$ -skeleta

$$\mathbf{g}: X^{(\alpha)} \rightarrow X'^{(\alpha)} \quad \text{and} \quad \mathbf{h}: X'^{(\alpha)} \rightarrow X^{(\alpha)}$$

that are homotopy inverse to each other up to dimension  $\alpha - 1$ , and a cellular homotopy  $\mathbf{P}: [0, 1] \times X^{(\alpha-1)} \rightarrow X^{(\alpha)}$  between the identity and  $\mathbf{h} \circ \mathbf{g}$  on  $X^{\alpha-1}$ .

Recall furthermore (Definition 2 of the introduction) that given two real numbers  $T, \kappa \geq 1$ , a  $\alpha$ -rebuilding  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$  is of **quality**  $(T, \kappa)$  if

$$\forall j \leq \alpha, \quad |X'^{(j)}| \leq \kappa T^{-1} |X^{(j)}| \quad (\text{cells bound})$$

$$\forall j \leq \alpha, \quad \log \|g_j\|, \log \|h_j\|, \log \|\rho_{j-1}\|, \log \|\partial'_j\| \leq \kappa(1 + \log T) \quad (\text{norms bound})$$

where  $g, h$  are the chain maps respectively associated to  $\mathbf{g}, \mathbf{h}$  and where  $\rho: C_\bullet(X) \rightarrow C_{\bullet+1}(X)$  is the chain homotopy induced by  $\mathbf{P}$  in the cellular chain complexes (1.8).

In this paragraph we make two general observations regarding rebuildings and their quality.

First note that if  $X_1 \rightarrow X$  is a finite cover, every  $\alpha$ -rebuilding  $(X, X', \mathbf{g}, \mathbf{h}, \mathbf{P})$  of  $X$  induces (via the lifting property [Hat02, Prop. 1.33 and 1.34]) a  $\alpha$ -rebuilding  $(X_1, X'_1, \mathbf{g}_1, \mathbf{h}_1, \mathbf{P}_1)$  with  $X'_1 = \widetilde{X}'/\pi_1(X_1)$ .

**Lemma 6.2** (Rebuilding induced to finite cover). *Let  $X$  be a finite CW-complex. There is a constant  $\delta_X$  such that for every  $\alpha$ -rebuilding  $(X, X')$  of quality  $(T, \kappa)$  and for every finite cover  $X_1 \rightarrow X$ , the induced  $\alpha$ -rebuilding  $(X_1, X'_1)$  is of quality  $(T, \kappa\delta_X)$ .*

*Proof.* A covering of CW-complexes induces a trivial covering over each open cell. In particular:

- (1) both sides of the cells bounds are multiplied by the degree of the cover leaving the quality of the cells bounds unchanged, and
- (2) the degrees of the attaching map of each cell remains bounded along coverings by a constant depending only on  $X$ .

It follows from this last observation that the norms (induced by the  $\ell^2$ -norms) of the boundary map and of the maps  $g, h$  and  $\rho$  remain bounded by a constant  $\delta_X$  along coverings.  $\square$

**Lemma 6.3** (Composition of rebuildings). *Let  $(X, X', \mathbf{g}_1, \mathbf{h}_1, \mathbf{P}_1)$  and  $(X', X'', \mathbf{g}_2, \mathbf{h}_2, \mathbf{P}_2)$  be two  $\alpha$ -rebuildings of respective quality  $(T_1, \kappa_1)$  and  $(T_2, \kappa_2)$ , with  $T_1, T_2, \kappa_1, \kappa_2 \geq 1$ . Let*

$$\mathbf{g}_3 = \mathbf{g}_2 \circ \mathbf{g}_1, \quad \mathbf{h}_3 = \mathbf{h}_1 \circ \mathbf{h}_2, \quad \text{and} \quad \mathbf{P}_3(t, x) = \begin{cases} \mathbf{P}_1(2t, x) & \text{if } 0 \leq t \leq 1/2, \\ \mathbf{h}_1 \circ \mathbf{P}_2(2t - 1, \mathbf{g}_1(x)) & \text{if } 1/2 < t \leq 1. \end{cases}$$

*Then  $(X, X'', \mathbf{g}_3, \mathbf{h}_3, \mathbf{P}_3)$  is a  $\alpha$ -rebuilding of quality  $(T_1 T_2, 4\kappa_1 \kappa_2)$ .*

*Proof.* The fact that  $(X, X'', \mathbf{g}_3, \mathbf{h}_3, \mathbf{P}_3)$  is a  $\alpha$ -rebuilding follows from the definition. The quality of the cells bounds being multiplicative, it remains to check the norms bounds. Since we have

$$\begin{aligned} \log \|g_3\| &\leq \log \|g_1\| + \log \|g_2\| \leq \kappa_1(1 + \log T_1) + \kappa_2(1 + \log T_2) \\ &\leq \kappa_1 + \kappa_2 + \kappa_1 \log T_1 + \kappa_2 \log T_2 \\ &\leq 2\kappa_1 \kappa_2(1 + \log T_1 T_2), \end{aligned}$$

similarly for  $h_3$ , and

$$\begin{aligned} \log \|\rho_3\| &= \log \|\rho_1 + h_1 \circ \rho_2 \circ g_1\| \leq \log 2 + \max\{\log \|\rho_1\|, \log \|h_1 \circ \rho_2 \circ g_1\|\} \\ &\leq \log 2 + \kappa_1(1 + \log T_1) + \kappa_2(1 + \log T_2) + \kappa_1(1 + \log T_1) \\ &\leq 4\kappa_1 \kappa_2(1 + \log T_1 T_2), \end{aligned}$$

Here the degrees match well, e.g.  $(\rho_3)_{\alpha-1} = (\rho_1)_{\alpha-1} + (h_1)_\alpha \circ (\rho_2)_{\alpha-1} \circ (g_1)_{\alpha-1}$  in top degree. The lemma follows.  $\square$



**6.2. Unipotent lattices.** Now let  $\Gamma$  be a finitely generated, torsion free nilpotent group, equivalently — by a theorem of Malcev [Mal49] — the group  $\Gamma$  is a **unipotent lattice**, i.e., it is isomorphic to a lattice in a connected, finite dimensional unipotent Lie group.

We define a central series  $L_i(\Gamma)$  by

$$L_0(\Gamma) = \Gamma \quad \text{and} \quad L_{i+1}(\Gamma) = \ker(L_i(\Gamma) \rightarrow (L_i(\Gamma)/[\Gamma, L_i(\Gamma)]) \otimes_{\mathbb{Z}} \mathbb{Q}).$$

By definition, the quotients  $L_i(\Gamma)/L_{i+1}(\Gamma)$  are torsion free abelian. We define the graded group

$$\text{gr}\Gamma = \bigoplus_{i=0}^{\infty} L_i(\Gamma)/L_{i+1}(\Gamma).$$

We have  $\text{gr}\Gamma \simeq \mathbb{Z}^{\mathfrak{h}}$  where  $\mathfrak{h}$  is the **Hirsch length** of  $\Gamma$ . For the purpose of this paper this can be taken as the definition of Hirsch length.

The goal of this section is to prove Theorem E according to which finite covers of classifying spaces of unipotent lattices admit a rewiring of quality proportional to the degree of the cover. This is, up to a constant, the best quality one could hope for.

To prepare for the proof we will need a few simple lemmas.

**Definition 6.4.** An automorphism  $\sigma$  of an unipotent lattice  $\Gamma$  is a **unipotent automorphism** if the induced automorphism of the graded group

$$\text{gr}\sigma \in \text{Aut}(\text{gr}\Gamma) \simeq \text{GL}_{\mathfrak{h}}(\mathbb{Z})$$

is unipotent.

Equivalently  $\sigma \in \text{Aut}(\Gamma)$  is unipotent if and only if the associated semi-direct product

$$\Gamma \rtimes_{\sigma} \mathbb{Z} = \langle (\gamma, t) \mid t\gamma t^{-1} = \sigma(\gamma) \rangle$$

is nilpotent.

**Lemma 6.5.** *Let  $\Gamma$  be a non-trivial unipotent lattice and  $\sigma$  be a unipotent automorphism of  $\Gamma$ . There exists a normal subgroup  $\Gamma' \leq \Gamma$  with quotient  $\Gamma/\Gamma' \simeq \mathbb{Z}$  such that  $\Gamma'$  is preserved by  $\sigma$  and  $\sigma$  acts trivially on  $\Gamma/\Gamma'$ .*

*Proof.* The group  $L_1(\Gamma)$  is preserved by  $\sigma$  and by construction we have  $\Gamma/L_1(\Gamma) \simeq \mathbb{Z}^d$  for some  $d > 0$ . Write  $\sigma_0 \in \text{GL}(d, \mathbb{Z})$  for the automorphism induced on  $\Gamma/L_1(\Gamma) \simeq \mathbb{Z}^d$ . Since  $\sigma_0$  is unipotent we can find a codimension 1 subspace  $W \subset V = \mathbb{Q}^d$  such that  $\sigma_0 W = W$  and the induced action on  $V/W$  is trivial. Let  $U = \mathbb{Z}^d \cap W$  and put  $\Gamma' = UL_1(\Gamma)$ . Then  $\sigma(\Gamma') = \sigma_0(U)L_1(\Gamma) = \Gamma'$ , the quotient

$$\Gamma/\Gamma' \simeq \mathbb{Z}^d/\mathbb{Z}^d \cap W \simeq \mathbb{Z}$$

and the induced action of  $\sigma$  on this quotient is trivial. □

**Lemma 6.6.** *Let  $(X_0, x_0)$  be a pointed CW-complex. Let  $G = \pi_1(X_0, x_0)$  and assume that  $X_0$  is a classifying space for  $G$ . Let  $\alpha$  and  $\beta: (X_0, x_0) \rightarrow (X_0, x_0)$  be two cellular maps such that the induced morphisms*

$$\alpha_* \quad \text{and} \quad \beta_* : G \rightarrow G$$

*are conjugated by some element  $g_0 \in G : \alpha_* = g_0 \beta_* g_0^{-1}$ . Then, there exists a cellular homotopy*

$$H : I \times X \rightarrow X \quad \text{such that} \quad H(0, \cdot) = \alpha, \quad H(1, \cdot) = \beta$$

*and the loop*

$$[0, 1] \rightarrow X_0, \quad s \mapsto H(s, x_0)$$

*represents  $g_0 \in G$ .*

*Proof.* We first construct an explicit cellular map  $q: (X_0, x_0) \rightarrow (X_0, x_0)$  that induces the morphism

$$q_* : G \rightarrow G, \quad g \mapsto g_0^{-1} g g_0$$

and a (non-pointed!) homotopy between the identity map and  $q$ . Start with a map

$$q_0 : (\{0\} \times X_0) \cup (I \times \{x_0\}) \rightarrow X_0, \quad q_0(0, \cdot) = \text{id}_{X_0} \quad \text{and} \quad q_0(\cdot, x_0) = s$$

where  $s: I \rightarrow X_0$  is some loop based at  $x_0$  and representing  $g_0^{-1} \in G$ . By Lemma 5.1 there is a map

$$p: I \times X_0 \rightarrow (\{0\} \times X_0) \cup (I \times \{x_0\}) \quad \text{such that} \quad p(0, \cdot) = \{0\} \times \text{id}_{X_0} \quad \text{and} \quad p(1, x_0) = (1, x_0).$$

Let

$$q: (X_0, x_0) \rightarrow (X_0, x_0); \quad x \mapsto q_0(p(1, x)).$$

It follows from the construction that  $q$  induces the morphism

$$q_* : G \rightarrow G, \quad g \mapsto g_0^{-1} g g_0$$

and the map

$$H_0: I \times X_0 \rightarrow X_0; (t, x) \mapsto q_0(p(t, x))$$

gives a homotopy between the identity map and  $q$ . The maps

$$q \circ \alpha \quad \text{and} \quad \beta: (X_0, x_0) \rightarrow (X_0, x_0)$$

induce the same endomorphisms of  $G$ . It therefore follows from e.g. [Geo08, Proposition 7.1.6] that there exists a homotopy

$$H_1: I \times X_0 \rightarrow X_0 \quad \text{such that} \quad H(0, \cdot) = q \circ \alpha, \quad H(1, \cdot) = \beta \quad \text{and} \quad H(\cdot, x_0) = x_0.$$

Therefore, the map

$$H(t, x) = \begin{cases} H_0(2t, \alpha(x)) & \text{if } 0 \leq t \leq 1/2, \\ H_1(2t - 1, x) & \text{if } 1/2 \leq t \leq 1, \end{cases}$$

is a homotopy between  $\alpha$  and  $\beta$  and the loop  $s \mapsto H(s, x_0)$  represents  $g_0$ .  $\square$

**Lemma 6.7.** *Let  $\Gamma$  be a unipotent lattice of Hirsch length  $\mathfrak{h}$  and let  $\sigma: \Gamma \rightarrow \Gamma$  be a unipotent automorphism. There exists a  $K(\Gamma, 1)$  CW-complex  $X_0$  with a distinguished point  $x_0$  and a cellular map  $\theta: X_0 \rightarrow X_0$  fixing  $x_0$  such that for every  $\gamma \in \Gamma$ ,*

$$\theta_*(\gamma) = \sigma(\gamma).$$

and for every  $m \in \mathbb{N}$  we have

$$\log \|\theta^m\| \leq \mathfrak{h} \log |m| + O_\Gamma(1),$$

where  $\|\theta^m\|$  denotes the norm induced by the  $\ell^2$ -norm of the action of  $\theta^m$  on the cellular chain complex associated to  $X_0$ .

*Proof.* We prove the lemma by induction on the Hirsch length of  $\Gamma$ . The base case  $\mathfrak{h} = 0$  corresponds to the trivial group and one can take  $X_0$  to be the CW-complex that consists of exactly one 0-cell.

Suppose now that the lemma is proved up to Hirsch length  $\mathfrak{h} - 1$  and consider a unipotent lattice  $\Gamma$  of Hirsch length  $\mathfrak{h}$ . By Lemma 6.5 there exists a  $\sigma$ -invariant normal subgroup  $\Gamma' \leq \Gamma$  such that  $\Gamma/\Gamma' \simeq \mathbb{Z}$  and the induced action of  $\sigma$  on  $\Gamma/\Gamma'$  is trivial. Choose  $t_0 \in \Gamma$  such that  $t_0\Gamma'$  generates  $\Gamma/\Gamma'$ .

By the inductive hypothesis there exists a  $K(\Gamma', 1)$  space  $X'_0$  with a distinguished point  $x'_0$  and a cellular automorphism  $\theta': X'_0 \rightarrow X'_0$  that fixes  $x'_0$ , induces  $\sigma'$  on  $\pi_1(X'_0, x'_0) = \Gamma'$  and satisfies

$$\log \|(\theta')^m\| \leq (\mathfrak{h} - 1) \log m + O_{\Gamma'}(1)$$

for  $m \in \mathbb{N}$ . Choose a cellular map  $\tau_0: X'_0 \rightarrow X'_0$  which fixes  $x'_0$  and induces the automorphism  $\gamma \mapsto t_0^{-1}\gamma t_0$  on the fundamental group. Take  $X_0$  to be the quotient

$$X_0 = (X'_0 \times [0, 1]) / \sim \quad \text{where } \sim \text{ is generated by } (x, 1) \sim (\tau_0(x), 0) \ (x \in X'_0)$$

and let  $x_0$  be the image of  $(x'_0, 0)$  in the quotient. The CW-complex  $X_0$  is a  $K(\Gamma, 1)$  space and the image of  $\{x'_0\} \times [0, 1]$  in  $X_0$  is a loop based at  $x_0$  that represents  $t_0$  in  $\pi_1(X_0) \simeq \Gamma$ . We proceed to construct the desired cellular map  $\theta: X_0 \rightarrow X_0$ . Since  $\sigma$  acts trivially on  $\Gamma/\Gamma'$  the two maps

$$\tau_0 \circ \theta' \quad \text{and} \quad \theta' \circ \tau_0 : (X_0, x_0) \rightarrow (X_0, x_0)$$

induce two endomorphisms of  $\Gamma$  that are conjugated by  $t_0^{-1}\sigma(t_0) \in \Gamma'$ . Lemma 6.6 therefore implies that there exists a homotopy

$$H: I \times X'_0 \rightarrow X'_0 \quad \text{such that} \quad H(0, \cdot) = \tau_0 \circ \theta', \quad H(1, \cdot) = \theta' \circ \tau_0$$

and the loop

$$[0, 1] \rightarrow X_0, \quad s \mapsto H(s, x_0)$$

represents  $t_0^{-1}\sigma(t_0) \in \Gamma'$ . We define  $\theta: X_0 \rightarrow X_0$  by

$$\theta(x, t) = \begin{cases} (\theta'(x), 2t) & \text{if } 0 \leq t \leq 1/2 \\ (H(2t - 1, x), 0) & \text{if } 1/2 < t \leq 1. \end{cases}$$

We claim that the endomorphism of  $\Gamma$  induced by  $\theta$  is  $\sigma$ . Since we already know that  $\theta$  induces  $\sigma$  on  $\Gamma'$  it is enough to compute its value on  $t_0$ . By construction,  $t_0$  is represented by the loop in  $X_0$  image of  $\{x'_0\} \times [0, 1]$ . This loop is sent by  $\theta$  to the concatenation of  $s \mapsto (x_0, s)$  with  $s \mapsto H(s, x_0)$ . The latter represents  $t_0^{-1}\sigma(t_0) \in \Gamma'$  so the concatenation represents  $\sigma(t_0)$  as desired.

It remains to compute the chain maps and to check the bound on the norm. Recall that as  $\mathbb{Z}$ -modules,

$$C_\bullet(X_0) = (C_\bullet(X'_0) \otimes [0]) \oplus (C_\bullet(X'_0)) \otimes [I, \partial I].$$

In these coordinates, the chain map induced by  $\theta$  is given by the following formulas

$$\theta(c \otimes [0]) = \theta'(c) \otimes [0] \quad \text{and} \quad \theta(c \otimes [I, \partial I]) = \theta'(c) \otimes [I, \partial I] + r(c) \otimes [0],$$

where  $r(c) = H([I, \partial I] \otimes c)$ . By induction we have

$$\theta^m(c \otimes [0]) = \theta'^m(c) \otimes [0] \quad \text{and} \quad \theta^m(c \otimes [I, \partial I]) = \theta'^m(c) \otimes [I, \partial I] + \sum_{i=0}^{m-1} \theta'^i(r(c)) \otimes [0].$$

It follows that

$$\|\theta^m\| \leq m \sup\{\|\theta'^m\|, \|\theta'^i \circ r\| : i = 1, \dots, m-1\} + \|r\|$$

which, by induction, gives

$$\log \|\theta^m\| \leq \mathfrak{h} \log m + 2 \log \|r\| + O_{\Gamma', \sigma'}(1) = \mathfrak{h} \log m + O_{\Gamma, \sigma}(1).$$

□

### 6.3. The Proof of Theorem E.

*Proof of Theorem E.* We prove the theorem by induction on the Hirsch length  $\mathfrak{h}$ . The base case  $\mathfrak{h} = 0$  holds trivially since then  $\Gamma$  is the trivial group.

Suppose now that the lemma is proved up to Hirsch length  $\mathfrak{h} - 1$  and consider a unipotent lattice  $\Gamma$  of Hirsch length  $\mathfrak{h} > 0$ . Thanks to Lemma 6.2 (Rebuilding induced to finite cover) and Lemma 6.3 (Composition of rebuildings) we only need to prove the theorem for a single classifying space of  $\Gamma$ . Therefore our first step is the construction of a convenient  $Y_0$ .

Let  $\Lambda$  be a normal subgroup of  $\Gamma$  such that  $\Gamma/\Lambda \simeq \mathbb{Z}$ . Choose  $t_0 \in \Gamma$  such that  $t_0\Lambda$  generates  $\Gamma/\Lambda$ . Then  $\Gamma$  decomposes as a semidirect product

$$\Gamma = \langle t_0 \rangle \rtimes \Lambda.$$

The automorphism

$$\Lambda \rightarrow \Lambda; \quad \lambda \mapsto t_0 \lambda t_0^{-1}$$

is unipotent. By Lemma 6.7 there exists a pointed CW-complex  $(X_0, x_0)$  which is a  $K(\Lambda, 1)$  space and a cellular map  $\theta: (X_0, x_0) \rightarrow (X_0, x_0)$  such that the induced endomorphism  $\theta_*: \Lambda \rightarrow \Lambda$  is precisely equal to the unipotent automorphism  $\lambda \mapsto t_0 \lambda t_0^{-1}$  and

$$\|\theta^m\| \leq (\mathfrak{h} - 1) \log m + O_{\Gamma', t_0}(1).$$

We let  $Y_0$  be the quotient

$$Y_0 = (X_0 \times [0, 1]) / (x, 1) \sim (\theta(x), 0)$$

and take the image  $y_0$  of  $(x_0, 0)$  as basepoint. Then  $\pi_1(Y_0, y_0)$  is isomorphic to  $\Gamma$  and  $Y_0$  is a  $K(\Gamma, 1)$  space. Note for future reference that the projection of  $\{x_0\} \times [0, 1]$  into  $Y_0$  is a loop that represents the element  $t_0$  in  $\Gamma$ .

Now let  $\Gamma_1 \leq \Gamma$  be a finite index subgroup. Set  $\ell = [\Gamma : \Lambda\Gamma_1]$  and let  $\Lambda_1 = \Lambda \cap \Gamma_1$ . Since  $t_0^\ell \in \Lambda\Gamma_1$  we can choose  $a \in \Lambda$  such that  $t_1 = at_0^\ell \in \Gamma_1$ . The group  $\Gamma_1$  then decomposes as a semi-direct product  $\Gamma_1 = \langle t_1 \rangle \rtimes \Lambda_1$ . It follows that  $[\Gamma : \Gamma_1] = \ell[\Lambda : \Lambda_1]$ . Put

$$Y_1 = \Gamma_1 \backslash \tilde{Y}_0.$$

This is the complex we want to rebuild. It is naturally a stack over the CW-complex obtained by lifting the standard CW-complex structure on the circle by a  $\ell$ -fold self-covering map. We rebuild  $Y_1$  in two steps. The first rebuilding  $(Y_1, Y_1')$  consists of changing the base of the stack to get a stack over the standard CW-complex on the circle. It decreases the number of cells to  $|Y_1'| = |Y_1| \ell^{-1}$ . The second rebuilding  $(Y_1', Y_1'')$  is obtained by applying our effective version of the rebuilding lemma after applying the inductive hypothesis to the fibers of the stack. This second rebuilding will bring the number of cells down to  $|Y_1''| = |Y_1| [\Gamma : \Gamma_1] = |Y_0|$ .

Before performing the two rebuildings we give an explicit description of the stack structure of  $Y_1$ : the map  $\theta: X_0 \rightarrow X_0$  lifts to a map  $\theta: \tilde{X}_0 \rightarrow \tilde{X}_0$  such that for all  $\lambda \in \Lambda$  and for all  $\tilde{x} \in \tilde{X}_0$ , we have

$$\theta(\lambda \cdot \tilde{x}) = \theta_*(\lambda) \cdot \tilde{x}.$$

In particular  $\theta$  maps each  $(t_0^i \Lambda_1 t_0^{-i})$ -orbit of  $\tilde{x}$  (with  $i \in \mathbb{Z}$ ) to the  $(t_0^{i+1} \Lambda_1 t_0^{-i-1})$ -orbit of  $\tilde{x}$ . Writing

$$X_{1,(i)} = (t_0^i \Lambda_1 t_0^{-i}) \backslash \tilde{X}_0 \quad \text{and simply} \quad X_1 = X_{1,(0)},$$

we conclude that  $\theta$  induces a well-defined map

$$\theta: X_{1,(i)} \rightarrow X_{1,(i+1)}.$$

Finally since

$$a(t_0^\ell \Lambda_1 t_0^{-\ell}) a^{-1} = t_1 \Lambda_1 t_1^{-1} = \Lambda_1,$$

the map

$$\tilde{X}_0 \rightarrow \tilde{X}_0; \quad \tilde{x} \mapsto a \cdot \tilde{x}$$

induces a well-defined map

$$a: X_{1,(\ell)} \rightarrow X_0; \quad (t_0^\ell \Lambda_1 t_0^{-\ell}) \tilde{x} \mapsto \Lambda_1 a \tilde{x}.$$

It follows that

$$Y_1 \simeq \bigsqcup_{i=0}^{\ell-1} (X_{1,(i)} \times [0, 1]) / \sim, \tag{6.1}$$

where  $\sim$  is generated by

$$(x_{(i)}, 1) \sim (\theta(x_{(i)}), 0), \quad \text{for } i \in \{0, \dots, \ell-2\}, \quad \text{and} \quad (x_{(\ell-1)}, 1) \sim (a \circ \theta(x_{(\ell-1)}), 0).$$

Indeed: we have

$$\Lambda_1 \backslash \tilde{Y}_0 = \bigsqcup_{i \in \mathbb{Z}} (X_{1,(i)} \times [0, 1]) / (x_{(i)}, 1) \sim (\theta(x_{(i)}), 0)$$

and quotienting it further by the action of  $\langle t_1 \rangle$ , which has the effect of identifying  $(x_{(i)}, t)$  with  $(a(x_{(i+\ell)}), t)$ , we get (6.1).

**First rebuilding.** Let

$$Y'_1 = X_1 \times [0, 1] / (x, 1) \sim (a \circ \theta^\ell(x), 0).$$

It is aspherical and  $\pi_1(Y'_1) \simeq \langle t_1 \rangle \rtimes \Lambda_1 = \Gamma_1$  so  $Y'_1$  is another  $K(\Gamma_1, 1)$  space. Note that  $|Y'_1| = \ell^{-1}|Y_1|$ . We define three maps

$$\mathbf{g}: Y_1 \rightarrow Y'_1, \quad \mathbf{h}: Y'_1 \rightarrow Y_1 \quad \text{and} \quad \mathbf{P}: I \times Y_1 \rightarrow Y_1$$

by the following formulas, with  $x_{(i)} \in X_{1,(i)}$ ,  $x \in X_1$  and  $s, t \in [0, 1]$ ,

$$\mathbf{g}(x_{(i)}, t) = \begin{cases} (x_{(i)}, t) & \text{if } i = 0, \\ (a \circ \theta^{\ell-i}(x_{(i)}), 0) & \text{if } i = 1, \dots, \ell - 1, \end{cases}$$

$$\mathbf{h}(x, t) = (\theta^{\lfloor \ell t \rfloor}(x), \ell t - \lfloor \ell t \rfloor),$$

and

$$\mathbf{P}(s, (x_{(i)}, t)) = \begin{cases} (\theta^{\lfloor \alpha \rfloor}(x_{(i)}), \alpha - \lfloor \alpha \rfloor) & \text{if } i + t \leq (\ell - 1)s \text{ and } \alpha = \frac{\ell(i+t-s(\ell-1))}{\ell-s(\ell-1)}, \\ (a \circ \theta^{\ell-i}(x_{(i)}), 0) & \text{if } i + t \geq (\ell - 1)s. \end{cases}$$

The maps  $\mathbf{g}$  and  $\mathbf{h}$  are cellular and  $\mathbf{P}$  is a cellular homotopy between the identity and  $\mathbf{h} \circ \mathbf{g}$ . The composition  $\mathbf{g} \circ \mathbf{h}$  is also homotopic to the identity since it induces the identity on  $\pi_1(Y'_1)$ . Therefore  $(Y_1, Y'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  is a rebuilding.

We now compute the chain maps induced by  $\mathbf{g}$ ,  $\mathbf{h}$  and  $\mathbf{P}$  as well as the boundary map on  $C_\bullet(Y'_1)$ , which we shall denote by  $\partial_{Y'_1}$ . First remark that

$$C_\bullet(Y_1) = \bigoplus_{i=0}^{\ell-1} (C_\bullet(X_{1,(i)}) \otimes [0] \oplus C_\bullet(X_{1,(i)}) \otimes [I]) \quad \text{and} \quad C_\bullet(Y'_1) \simeq C_\bullet(X_1) \otimes [0] \oplus C_\bullet(X_1) \otimes [I],$$

as  $\mathbb{Z}$ -modules. In these coordinates, for  $c \in C_\bullet(X_1)$  we have

$$\partial_{Y'_1}(c \otimes [0]) = \partial_{X_1} c \otimes [0] \quad \text{and} \quad \partial_{Y'_1}(c \otimes [I]) = \partial_{X_1} c \otimes [I] + (-1)^{\dim c} (-c \otimes [0] + a \circ \theta^\ell(c) \otimes [0]),$$

for  $c_{(i)} \in C_\bullet(X_{1,(i)})$  we have

$$g(c_{(i)} \otimes [0]) = \begin{cases} c_{(i)} \otimes [0] & \text{if } i = 0, \\ a \circ \theta^{\ell-i}(c_{(i)}) \otimes [0] & \text{if } i > 0, \end{cases} \quad \text{and} \quad g(c_{(i)} \otimes [I]) = \begin{cases} c_{(i)} \otimes [I] & \text{if } i = 0, \\ 0 & \text{if } i > 0, \end{cases}$$

for  $c \in C_\bullet(X_1)$  we have

$$h(c \otimes [0]) = c \otimes [0] \quad \text{and} \quad h(c \otimes [I]) = \sum_{i=0}^{\ell-1} \theta^i(c) \otimes [I],$$

and for  $c_{(i)} \in C_\bullet(X_{1,(i)})$  we have

$$\rho(c_{(i)} \otimes [0]) = \begin{cases} 0 & \text{if } i = 0, \\ \sum_{j=i}^{\ell-1} \theta^j(c) \otimes [I], & \text{if } i > 0, \end{cases} \quad \text{and} \quad \rho(c_{(i)} \otimes [I]) = 0.$$

The map  $a: C_\bullet(X_{1,(\ell)}) \rightarrow C_\bullet(X_1)$  is unitary and by Lemma 6.7  $\|\theta^\ell\| \leq O_{X_0, \sigma}(1 + \log \ell)$ . We deduce that

$$\|\partial_{Y'_1}\|, \quad \|g\|, \quad \|h\|, \quad \|\rho\| \leq O_{Y_0}(1 + \log \ell). \quad (6.2)$$

This concludes the first step.

**Second rebuilding.** By induction there exists a rebuilding  $(X_1, X'_1, \mathbf{k}, \mathbf{l}, \Sigma)$  of quality  $([\Lambda : \Lambda_1], O_{X_0}(1))$ . To shorten notation we shall write  $\theta_1 = a \circ \theta^\ell$ . Let

$$A = X_1 \times \{0, 1\} \quad \text{and} \quad A' = X'_1 \times \{0, 1\}.$$

Define a map  $f: A \rightarrow X_1$  by

$$f(x, 0) = x \quad \text{and} \quad f(x, 1) = \theta_1(x).$$

We then have

$$Y'_1 = X_0 \times [0, 1]/(x, 1) \sim (\theta_1(x), 0) = (X_1 \times [0, 1]) \cup_f X_1.$$

Now consider the following diagram

$$\begin{array}{ccccc} X_1 \times [0, 1] & \longleftarrow & A & \xrightarrow{f} & X_1 \\ \mathbf{k} \times \text{id} \downarrow \uparrow \mathbf{l} \times \text{id} & & \mathbf{k} \times \text{id} \downarrow \uparrow \mathbf{l} \times \text{id} & & \mathbf{k} \downarrow \uparrow \mathbf{l} \\ X'_1 \times [0, 1] & \longleftarrow & A' & \xrightarrow{f'} & X'_1, \end{array}$$

where  $f' = \mathbf{k} \circ f \circ (\mathbf{l} \times \text{id})$ . Define

$$Y''_1 = (X'_1 \times [0, 1]) \cup_{f'} X'_1.$$

The number of cells in  $Y''_1$  is  $2|X'_1| = O_{X_0}(1)$  and

$$C_\bullet(Y''_1) = C_\bullet(X_1) \otimes [I] \oplus C_\bullet(X_1),$$

as  $\mathbb{Z}$ -modules. The boundary map is given by

$$\partial_{Y'_1}(c \otimes [I]) = \partial_{X'_1}(c) \otimes [I] + (-1)^{\dim c} (-k(\theta_1(l(c))) + k(l(c))) \quad \text{and} \quad \partial_{Y'_1}(c) = \partial_{X'_1}(c),$$

for  $c \in C_\bullet(X'_1)$ .

Let

$$\mathbf{g}' : Y'_1 \rightarrow Y''_1 \quad \text{and} \quad \mathbf{h}' : Y''_1 \rightarrow Y'_1$$

be the cellular maps afforded by Proposition 4.1 and let

$$\mathbf{P}' : I \times Y'_1 \rightarrow Y'_1$$

be the homotopy between the identity and  $\mathbf{h}' \circ \mathbf{g}'$ , also provided by Proposition 4.1. Let

$$\sigma : C_\bullet(X_1) \rightarrow C_{\bullet+1}(X_1) \quad \text{and} \quad \rho' : C_\bullet(Y'_1) \rightarrow C_{\bullet+1}(Y'_1)$$

be the chain homotopies respectively induced by  $\Sigma$  and  $\mathbf{P}'$ .

By induction we have

$$\log \|k\|, \quad \log \|l\|, \quad \log \|\sigma\| \leq O_{X_0}(1 + \log[\Lambda : \Lambda_1]) \quad \text{and} \quad \log \|f\| \leq O_{Y_0}(1 + \log \ell).$$

Using the explicit formulas for the chain maps from Lemma 4.1 we deduce that

$$\log \|\partial_{Y'_1}\|, \log \|g'\|, \log \|h'\|, \log \|\rho'\| \ll_{Y_0} \log \ell + \log[\Lambda : \Lambda_1] + 1 = \log[\Gamma : \Gamma_1] + 1. \quad (6.3)$$

Let

$$\mathbf{g}'' = \mathbf{g}' \circ \mathbf{g}, \quad \mathbf{h}'' = \mathbf{h} \circ \mathbf{h}'$$

and

$$\mathbf{P}''(s, y) = \begin{cases} \mathbf{P}(2s, y) & \text{if } 0 \leq s \leq 1/2, \\ \mathbf{h}(\mathbf{P}'(2t-1, g(y))) & \text{if } 1/2 \leq s \leq 1, \end{cases} \quad \text{for } y \in Y_1.$$

Using (6.2) and (6.3) we see that  $(Y_1, Y''_1, \mathbf{g}'', \mathbf{h}'', \mathbf{P}'')$  is a rebuilding of quality  $([\Gamma : \Gamma_1], O_{Y_0}(1))$ . This proves Theorem E.  $\square$

## 7. QUALITY OF REBUILDING OF EXTENSIONS BY UNIPOTENT LATTICES

Consider a countable group  $G$  that contains a finitely generated, torsion-free, nilpotent, **normal** subgroup  $A$ . Suppose furthermore that  $G/A$  is of type  $F_\alpha$  for some integer  $\alpha \geq 0$ .

Let  $B$  be a classifying space (CW-complex) for  $G/A$  with finite  $\alpha$ -skeleton. The group  $G$  acts on its universal cover  $\tilde{B}$  with cell-stabilizers all equal to  $A$ .

Let  $Y$  be a finite classifying space for  $A$ . The Borel construction followed by the rebuilding Lemma 2.2 yields a stack of CW-complexes  $D \rightarrow B$  with base  $B$  where each fiber is  $Y$ .

The goal of this section is to prove the following effective rebuilding statement.

**Proposition 7.1.** *Let  $G_1 \leq G$  be a finite index subgroup, let  $D_1 \rightarrow B_1$  be the associated stack and let  $A_1 = A \cap G_1$ . Then the total space  $D_1$  is a classifying space for  $G_1$  and there exists a  $\alpha$ -rebuilding  $(D_1, D'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  of quality  $([A : A_1], O_{Y,D}(1))$ .*

*Proof.* Each fiber of  $D \rightarrow B$  has the same number of cells  $n_i \in \mathbb{N}$  in each dimension  $i$  (with  $n_i = 0$  above the dimension of  $Y$ ).

The CW-complex  $B$  has finitely many cells in each dimension  $j \leq \alpha$ ; we denote by  $m_j$  this number. The total number of cells of dimension  $\ell \leq \alpha$  of the total complex  $D$  is then finite and equal to

$$N_\ell = \sum_{i+j=\ell} n_i m_j.$$

Now consider the stack  $D_1 \rightarrow B_1$ . The total space  $D_1$  has  $[G : G_1]N_\ell$  cells in each dimension  $\ell \leq \alpha$ . More precisely, each fiber  $Y_1$  of  $D_1 \rightarrow B_1$  is a classifying space for  $A_1 = A \cap G_1$  which contains  $[A : A_1]n_i$  cells in each dimension  $i$  and  $B_1$  contains  $[G/A : G_1/A_1]m_j$  cells in each dimension  $j \leq \alpha$ . Thus:

$$N_\ell[G : G_1] = \sum_{i+j=\ell} [A : A_1]n_i [G/A : G_1/A_1]m_j.$$

The group  $A$  being finitely generated, torsion-free and nilpotent, Theorem E implies that there exists a  $\alpha$ -rebuilding  $(Y_1, Y'_1, \mathbf{k}, \mathbf{l}, \mathbf{\Sigma})$  of quality  $([A : A_1], O_Y(1))$ . In particular  $Y'_1$  is a classifying space for  $A_1$  and letting  $D'_1 \rightarrow B_1$  be the rebuilding of the stack  $D_1 \rightarrow B_1$  associated to the rebuilding  $Y'_1$  of the fibers we conclude that  $D'_1$  has

$$O_Y \left( \sum_{i+j=\ell} n_i [G/A : G_1/A_1]m_j \right) = O_Y \left( \frac{N_\ell[G : G_1]}{[A : A_1]} \right)$$

cells in each dimension  $\ell \leq \alpha$ .

In fact Proposition 4.2 provides  $(D_1, D'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  and expresses the boundary map  $\partial'$  on  $C_n(D'_1)$ , the chain maps induced by  $\mathbf{g}$  and  $\mathbf{h}$ , and the chain homotopy induced by  $\mathbf{P}$  in terms of  $k, l, \sigma$ , the (vertical) boundary map on  $C_\bullet(Y)$  and the horizontal boundary map on  $C_\bullet(B)$ . The norms of the two boundary maps are bounded by  $O_{Y,D}(1)$  and, since the rebuilding  $(Y_1, Y'_1, \mathbf{k}, \mathbf{l}, \mathbf{\Sigma})$  is of quality  $([A : A_1], O_Y(1))$ , the log of the norms of  $k, l$  and  $\sigma$  are bounded by  $O_{Y,D}(1 + \log[A : A_1])$ . The formulas (4.10) therefore imply that  $(D_1, D'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  is of quality  $([A : A_1], O_{Y,D}(1))$ . Indeed, consider for example the boundary map  $\partial'$  on  $C_j(D'_1)$  for some  $j \leq \alpha$ . We have:

$$\begin{aligned} \|\partial'\| &\leq \|(\partial')^{\text{vert}}\| + \|k\| \left\| \left( \sum_{i=0}^j (\partial^{\text{hor}} \circ \sigma)^i \right) \right\| \|\partial^{\text{hor}}\| \|l\| \\ &\leq \|(\partial')^{\text{vert}}\| + P_j([A : A_1]) \end{aligned}$$

where  $P_j$  is a polynomial of degree  $O_{Y,D}(1)$  whose coefficients do not depend on the subgroup  $A_1$ . It follows that

$$\log \|\partial'\| = O_{Y,D}(1 + \log[A : A_1]).$$

The bounds for  $g$ ,  $h$  and  $\rho$  are obtained similarly.  $\square$

### 8. A GENERAL QUANTITATIVE REBUILDING THEOREM

Let  $\alpha$  be a positive integer and let  $\Gamma \curvearrowright \Omega$  be a CW-complex action of a countable group  $\Gamma$  that satisfies the following assumptions.

- (**Cond 1**) : The CW-complex  $\Omega$  is  $(\alpha - 1)$ -connected.
- (**Cond 2**) : For every cell  $\omega \subset \Omega$  the stabilizer  $\Gamma_\omega$  acts trivially on  $\omega$ .
- (**Cond 3**) : The quotient CW-complex  $\Phi := \Gamma \backslash \Omega$  has finite  $\alpha$ -skeleton.
- (**Cond 4**) : The group  $\Gamma$  is of type  $F_{\alpha+1}$ .
- (**Cond 5**) : Each stabilizer group  $\Gamma_\omega$  is of type  $F_\alpha$ .

Recall from Proposition 2.1 that, as long as  $\Omega$  is assumed to be simply connected, the Borel construction associates to this action a stack of CW-complexes  $\Sigma \rightarrow \Phi := \Gamma \backslash \Omega$  such that  $\Sigma$  has a  $(\alpha - 1)$ -connected universal cover and the fundamental group  $\pi_1(\Sigma)$  isomorphic to  $\Gamma$ .

**Theorem 8.1** (Quantitative rebuilding). *Suppose that  $\Gamma$  is finitely presented and that for each cell  $e$  of  $\Phi^{(\alpha)}$  the group  $\Gamma_e$  is finitely generated and contains a finitely generated, torsion-free, nilpotent, normal subgroup  $Z_e$ . There exists a constant  $\kappa$  such that for every finite index normal subgroup  $\Gamma_1 \leq \Gamma$ , there exists a CW-complex  $\Sigma_1^+$  with finite  $(\alpha + 1)$ -skeleton such that the following hold.*

- (1) *The CW-complex  $\Sigma_1^+$  has a  $\alpha$ -connected universal cover.*
- (2) *The fundamental group  $\pi_1(\Sigma_1^+)$  is isomorphic to  $\Gamma_1$ .*
- (3) *In each dimension  $\leq \alpha$ , the total number of cells of  $\Sigma_1^+$  is*

$$\leq \kappa \sum_{e \in \Phi^{(\alpha)}} \frac{[\Gamma : \Gamma_1]}{[Z_e : \Gamma_1 \cap Z_e]}. \quad (8.1)$$

- (4) *In each degree  $\leq \alpha + 1$ , the norm of the boundary operator on the chain complex  $C_\bullet(\Sigma_1^+)$  is*

$$\leq \kappa [\Gamma : \Gamma_1]^\kappa. \quad (8.2)$$

To prove the theorem we will consider the stack of CW-complexes  $\Sigma \rightarrow \Phi := \Gamma \backslash \Omega$ . However for the fundamental group of the latter to be isomorphic to  $\Gamma$ , we need  $\Omega$  to be simply connected. This follows from the hypothesis for  $\alpha \geq 2$  but in certain interesting cases, e.g., for  $\Gamma = \mathrm{SL}_3(\mathbb{Z})$ , the natural candidate for  $\Omega$  is the Tits complex which fails to be simply connected. We need a way to “fix” these spaces before we start using them. This is the content of the following:

**Lemma 8.2.** *Let  $\Gamma$  be a finitely presented group. Consider a 1-dimensional co-compact  $\Gamma$ -CW-complex  $\Omega$  whose cells are fixed pointwise by their stabilizers. Assume that all vertex stabilizers are finitely generated. Then there exists a simply connected co-compact  $\Gamma$ -CW-complex  $\Omega^+$  whose 1-skeleton is  $\Omega$ .*

*Proof.* Up to refining the structure of the CW-complex on  $\Omega$ , we may assume that it is a graph. The point is to kill the fundamental group of  $\Omega$  by gluing some 2-cells along finitely many  $\Gamma$ -orbits of loops in  $\Omega$ . By [DD89, Theorem 9.2, p. 39] the group  $\Gamma$  sits in a short exact sequence

$$1 \rightarrow \pi_1(\Omega) \rightarrow G \xrightarrow{\zeta} \Gamma \rightarrow 1 \quad (8.3)$$

where  $G$  is the fundamental group of the natural quotient graph of groups associated with the action  $\Gamma \curvearrowright \Omega$ . The assumption that the quotient space is compact and the vertex stabilizers are finitely generated imply that  $G$  is finitely generated. Thus  $G$  is the quotient of a free group  $\mathbf{F}(\tilde{S})$  (where  $\tilde{S}$  is finite). By (8.3),  $\Gamma$  is also such a quotient. Thus we have surjective morphisms:

$$\mathbf{F}(\tilde{S}) \xrightarrow{\eta} G \xrightarrow{\zeta} \Gamma.$$



By standard arguments, since it is finitely presented,  $\Gamma$  admits a finite presentation with these generators  $\tilde{S}$ .

$$\Gamma = \langle \tilde{S} | r_1, r_2, \dots, r_p \rangle.$$

It follows that the normal subgroup of  $G$  generated by  $\eta(r_1), \eta(r_2), \dots, \eta(r_p)$  is exactly the kernel of  $\zeta$ , i.e., it is  $\pi_1(\Omega)$ .

Consider now the action of  $G$  on its Bass-Serre tree  $T$ , a base vertex  $v$  in  $T$  and the paths  $c_j$  in  $T$  from  $v$  to  $\eta(r_j)(v)$ . They project to loops  $\sigma_1, \sigma_2, \dots, \sigma_p$  in  $\Omega = \pi_1(\Omega) \backslash T$ . Since these loops normally generate  $\pi_1(\Omega)$ , gluing a 2-cell along the  $\sigma_i$  and extending  $\Gamma$ -equivariantly produces the required simply connected  $\Gamma$ -cocompact  $\Omega^+$ .  $\square$

*Proof of the Quantitative Rebuilding Theorem 8.1.* We need to work with an  $\Omega$  that is simply connected. If  $\alpha \geq 2$  this follows from the assumptions. For  $\alpha = 1$  we can use Lemma 8.2 to replace  $\Omega$  by (its 1-skeleton and then by) a simply connected 2-dimensional complex  $\Omega^+$  satisfying all the assumptions of the theorem. In any case, the Borel construction followed by the rebuilding Lemma 2.2 then associates to the action of  $\Gamma$  on  $\Omega$  a stack of CW-complexes  $\Sigma \rightarrow \Phi := \Gamma \backslash \Omega$  such that the fundamental group of  $\Sigma$  is isomorphic to  $\Gamma$  and such that the fiber over each cell  $e = \Gamma \omega$  is a classifying space  $F_e$  of  $\Gamma_\omega$  with finite  $\alpha$ -skeleton.

Now consider the stack of CW-complexes

$$\Sigma_1 \rightarrow \Phi_1 \tag{8.4}$$

associated to the finite index subgroup  $\Gamma_1 \leq \Gamma$ , so that

$$\Sigma_1 = \Gamma_1 \backslash \tilde{\Sigma} \quad \text{and} \quad \Phi_1 = \Gamma_1 \backslash \Omega.$$

Each cell  $e \in \Phi$  is covered by  $\#(\Gamma_e \backslash \Gamma / \Gamma_1)$  cells in  $\Gamma_1 \backslash \Omega$  and, since  $\Gamma_1$  is a normal subgroup of  $\Gamma$ , the fibers of (8.4) over each of these cells are all isomorphic to the finite cover  $D_e$  of  $F_e$  associated to  $\Gamma_1 \cap \Gamma_e \leq \Gamma_e$ .

By hypothesis, as long as  $\dim e \leq \alpha$ , the group  $\Gamma_e$  contains a finitely generated, torsion-free, nilpotent, normal subgroup  $Z_e$ . We may therefore apply Proposition 7.1 with  $G = \Gamma_e$  and  $G_1 = \Gamma_e \cap \Gamma_1$ . It follows that there exists a rebuilding  $(D_e, D'_e, \mathbf{k}_e, \mathbf{l}_e, \Sigma_e)$  of quality  $([Z_e : \Gamma_1 \cap Z_e], O_{\Gamma, \Omega}(1))$  of each fiber  $D_e$  of (8.4).

Applying the rebuilding lemma (Proposition 4.2) to (8.4) we get a stack of CW-complexes

$$\Pi' : \Sigma'_1 \rightarrow \Phi_1,$$

cellular homotopy equivalences  $\mathbf{g}_1 : \Sigma_1 \rightarrow \Sigma'_1$ ,  $\mathbf{h}_1 : \Sigma'_1 \rightarrow \Sigma_1$  and a homotopy  $\mathbf{P}_1$  between the identity and  $\mathbf{h}_1 \circ \mathbf{g}_1$ . Note that over each cell of  $\Phi_1$  covering a cell  $e$  of  $\Phi$ , the fiber of  $\Pi'$  is  $D'_e$ .

The total space  $\Sigma'_1$  is homotopy equivalent to  $\Sigma_1$  and therefore has an  $(\alpha-1)$ -connected universal cover and a fundamental group isomorphic to  $\Gamma_1$ . Moreover, in each dimension  $n \leq \alpha$ , the total number of cells of  $\Sigma'_1$  is

$$O \left( \sum_{e \in \Phi^{(n)}} \#(\Gamma_e \backslash \Gamma / \Gamma_1) \frac{[\Gamma_e : \Gamma_1 \cap \Gamma_e]}{[Z_e : \Gamma_1 \cap Z_e]} \right) = O \left( \sum_{e \in \Phi^{(n)}} \frac{[\Gamma : \Gamma_1]}{[Z_e : \Gamma_1 \cap Z_e]} \right).$$

To control the norm of the boundary operator on the chain complex  $C_\bullet(\Sigma'_1)$  we apply Proposition 4.2 as in the proof of Proposition 7.1. Equation (4.10) implies that the boundary operator  $\partial'$  of the (rebuilt) chain complex  $C_\bullet(\Sigma'_1)$  can be written in terms of the (vertical) boundary operator in the fibers, the (vertical) maps  $(k, l$  and  $\sigma)$  and the boundary operators  $\partial$  and  $\partial^{\text{vert}}$  acting on the chain complex  $C_\bullet(\Sigma_1) = C_\bullet(\Gamma_1 \backslash \tilde{\Sigma})$  (before rebuilding).

The norm of the boundary operators  $\partial$  and  $\partial^{\text{vert}}$  are bounded by constants that depend only on the local combinatorial structure of  $\Sigma$ , see Lemma 6.2.

Now over each cell  $e$  of  $\Phi$  of dimension  $\leq \alpha$ , the rebuilding  $(D_e, D'_e, \mathbf{k}_e, \mathbf{l}_e, \Sigma_e)$  is of quality  $([Z_e : \Gamma_1 \cap Z_e], O_{\Gamma, \Omega}(1))$  and it follows that the norm of the vertical boundary operator  $(\partial')^{\text{vert}}$  is

$O([\Gamma_e : \Gamma_e \cap \Gamma_1]^{O(1)})$  and therefore  $O([\Gamma : \Gamma_1]^{O(1)})$ . For the same reason, the norms of the vertical maps  $k_e, l_e$  and  $\sigma_e$  are bounded by  $O([\Gamma_e : \Gamma_e \cap \Gamma_1]^{O(1)})$ , and formula (4.10) finally implies that the norm of the boundary operator on the chain complex  $C_{\leq \alpha}(\Sigma'_1)$  is a  $O([\Gamma : \Gamma_1]^{O(1)})$ .

The universal cover of  $\Sigma'_1$  is ‘only’  $(\alpha - 1)$ -connected. However, since  $\Gamma$  is of type  $F_{\alpha+1}$ , it follows from [Geo08, Theorem 8.2.1] that it is possible to attach *finitely many*  $\Gamma$ -orbits of  $(\alpha + 1)$ -cells to  $\widetilde{\Sigma}$  to make a  $\alpha$ -connected  $\Gamma$ -CW complex. Write the quotient as  $\Sigma^+ = \Sigma \sqcup_f \Phi$ , with  $\Phi = \sqcup_J \mathbb{B}^{\alpha+1}$  a finite collection of  $(\alpha + 1)$ -cells and  $f: \partial\Phi = \sqcup_J \mathbb{S}^\alpha \rightarrow \Sigma^{(\alpha)}$  the map that attaches these  $(\alpha + 1)$ -cells to  $\Sigma$ . Then write

$$\Gamma_1 \backslash \widetilde{\Sigma}^+ = \Sigma_1 \sqcup_{f_1} \Phi_1$$

where  $\Phi_1 = \sqcup_{J_1} \mathbb{B}^{\alpha+1}$  is the preimage of  $\Phi$  in  $\Sigma_1^+$  and  $f_1: \partial\Phi_1 = \sqcup_{J_1} \mathbb{S}^\alpha \rightarrow \Sigma_1^{(\alpha)}$  is the lift of  $f$ . We have a diagram

$$\begin{array}{ccccc} \Phi_1 & \longleftarrow & \partial\Phi_1 & \xrightarrow{f_1} & \Sigma_1 \\ \text{id} \updownarrow \text{id} & & \text{id} \updownarrow \text{id} & & g \updownarrow h \\ \Phi_1 & \longleftarrow & \partial\Phi_1 & \xrightarrow{\varphi} & \Sigma'_1, \end{array}$$

with  $\varphi = g \circ f_1$ . It then follows from Proposition 4.1 that the space

$$\Sigma_1^+ = \Sigma'_1 \sqcup_\varphi \Phi_1$$

is homotopy equivalent to  $\Sigma_1 \sqcup_{f_1} \Phi_1$ . The map  $f_1$ , being a lift of  $f$ , is of norm  $\|f_1\| \leq \|f\|$ . Finally the norm of the boundary map on  $\Sigma_1^+$  in degree  $\alpha + 1$  is bounded by  $\|\varphi\| \leq \|f_1\| \|g\|$ . The latter being of norm  $O([\Gamma : \Gamma_1]^{O(1)})$  we conclude that the norm of the degree  $\alpha + 1$  boundary operator on the chain complex  $C_\bullet(\Sigma_1^+)$  is  $O([\Gamma : \Gamma_1]^{O(1)})$ .  $\square$

## 9. BOUNDING TORSION - A PROOF OF GABBER’S PROPOSITION 9.1

In this section we prove the following useful proposition attributed to Gabber (see [Sou99, Prop. 3, p. 214]). Our proof here follows the viewpoint of [BV13, Section 2].

Let  $(C_\bullet, \partial)$  be the cellular chain complex associated to a finite CW-complex  $\Sigma$ . Each  $C_j$  is a free  $\mathbb{Z}$ -module of finite rank with a canonical basis associated to the  $j$ -cells of  $\Sigma$ . We equip each finite dimensional vector space  $C_j \otimes \mathbb{R}$  with the associated Euclidean norm. For any coefficient field  $K$  and for any integer  $j \geq 0$ , it follows from the definition of the homology groups that

$$\dim_K H_j(C_\bullet \otimes K) \leq \text{rank } C_j. \quad (9.1)$$

To bound the torsion homology we will use the following analogous observation.

**Proposition 9.1** (Gabber). *For any  $j \geq 0$ ,*

$$\log |H_j(C_\bullet)_{\text{tors}}| \leq (\text{rank } C_j) \times \sup(\log \|\partial_{j+1}\|, 0). \quad (9.2)$$

Here  $\|\partial_{j+1}\|$  denotes the operator norm.

*Proof.* Given a finite rank free  $\mathbb{Z}$ -module  $A$ , such that  $A_{\mathbb{R}} = A \otimes \mathbb{R}$  is endowed with a positive definite inner product  $(\cdot, \cdot)$  (a **metric** for short), we define  $\text{vol}(A)$  to be the volume of  $A_{\mathbb{R}}/A$ . When considered without further precision, the free  $\mathbb{Z}$ -module  $\mathbb{Z}^a$  ( $a \in \mathbb{N}^*$ ) will denote the standard one where  $\mathbb{R}^a = \mathbb{Z}^a \otimes \mathbb{R}$  is endowed with the canonical metric.

Let  $a > 0, b > 0$  be integers and

$$f: \mathbb{Z}^a \rightarrow \mathbb{Z}^b \quad (9.3)$$

be a  $\mathbb{Z}$ -linear map. We set  $\det'(f)$  to be the product of all nonzero singular values of  $f$ . Recall that the nonzero singular values of  $f$  are — with multiplicity — the positive square roots of the nonzero eigenvalues of  $ff^*$ . Note that we have

$$\det'(f) \leq \sup(\|f\|, 1)^{\min(a,b)}$$

where  $\|f\|$  denotes the operator norm of  $f_{\mathbb{R}} : \mathbb{R}^a \rightarrow \mathbb{R}^b$ .

Now recall from [BV13, (2.1.1)] the ‘‘metric rank formula’’

$$\det'(f) = \text{vol}(\text{image } f) \text{vol}(\ker f). \quad (9.4)$$

Here we understand the metrics on  $(\ker f) \otimes \mathbb{R}$  and  $(\text{image } f) \otimes \mathbb{R}$  as those induced from  $\mathbb{R}^a$  and  $\mathbb{R}^b$ . Let  $Q = \text{coker } f$  be the cokernel of  $f$ . It is a finitely generated  $\mathbb{Z}$ -module. Write  $Q = Q_{\text{tors}} \oplus Q_{\text{free}}$  its decomposition into a torsion and a free part.

Writing

$$\mathbb{R}^b = (\text{image } f) \otimes \mathbb{R} \oplus (\text{image } f)^{\perp} \quad (9.5)$$

and applying [BV13, (2.1.1)] to the orthogonal projection  $\mathbb{Z}^b \rightarrow (\text{image } f)^{\perp}$  we conclude that the quotient  $1/\text{vol}(\text{image } f)$  is the product of  $|Q_{\text{tors}}|^{-1}$  with the ‘regulator’  $\text{vol}(Q_{\text{free}})$ , where the metric on  $Q_{\text{free}} \otimes \mathbb{R}$  is obtained by identifying it as a subspace of  $(\text{image } f)^{\perp}$ . In summary,

$$\frac{1}{\text{vol}(\text{image } f)} = \frac{\text{vol}(Q_{\text{free}})}{|Q_{\text{tors}}|^{-1}}. \quad (9.6)$$

It follows from (9.4) and (9.6) that

$$|Q_{\text{tors}}| = \det'(f) \frac{\text{vol}(Q_{\text{free}})}{\text{vol}(\ker f)} \leq \det'(f). \quad (9.7)$$

The last inequality follows from two facts: (1)  $\ker f$  being a sub-lattice of  $\mathbb{Z}^a$  we have  $\text{vol}(\ker f) \geq 1$ , and (2)  $\text{vol}(Q_{\text{free}}) \leq 1$  since  $Q_{\text{free}}$  is spanned by vectors of length at most one.

Proposition 9.1 follows from the equation (9.7) applied to

$$Q = \text{coker}(\partial_{j+1} : C_{j+1} \rightarrow C_j).$$

Indeed, the homology group  $H_j(C_{\bullet})$  is contained in  $Q$  and it follows from (9.7) that the size of the torsion part of  $Q$  is smaller than

$$\det'(\partial_{j+1}) \leq \sup(\|\partial_{j+1}\|, 1)^{\text{rank } C_j}.$$

□

## 10. FARBER SEQUENCES AND CHEAP REBUILDING PROPERTY

**10.1. Farber neighborhoods.** Let  $\Gamma$  be a countable group. Let  $\text{Sub}_{\Gamma}$  denote the space of subgroups of  $\Gamma$  equipped with the topology induced from the topology of pointwise convergence on  $\{0, 1\}^{\Gamma}$ . The subset  $\text{Sub}_{\Gamma}^{f.i.} \subseteq \text{Sub}_{\Gamma}$  of finite index subgroups is equipped with the induced topology. It is countable when  $\Gamma$  is finitely generated. The group  $\Gamma$  continuously acts by conjugation on both  $\text{Sub}_{\Gamma}$  and  $\text{Sub}_{\Gamma}^{f.i.}$ .

We consider the fixed point ratio function defined for finite index subgroups  $\Gamma' \leq \Gamma$ :

$$\text{fx}_{\Gamma, \gamma} : \text{Sub}_{\Gamma}^{f.i.} \rightarrow [0, 1], \quad \Gamma' \mapsto \frac{|\{g\Gamma' | \gamma g\Gamma' = g\Gamma'\}|}{[\Gamma : \Gamma']}.$$

Thus  $\text{fx}_{\Gamma, \gamma}(\Gamma')$  is just the proportion of fixed points of  $\gamma \in \Gamma$  in the action  $\Gamma \curvearrowright \Gamma/\Gamma'$ .

**Definition 10.1.** A sequence  $(\Gamma_n)_{n \in \mathbb{N}}$  of subgroups of  $\Gamma$  is called a **Farber sequence** if it consists of finite index subgroups and for every  $\gamma \in \Gamma \setminus \{1\}$  we have  $\lim_{n \rightarrow \infty} \text{fx}_{\Gamma, \gamma}(\Gamma_n) = 0$ .

Though we won’t use it, note that if  $\Gamma$  is finitely generated and  $S \Subset \Gamma$  is a finite symmetric generating set, the sequence  $(\Gamma_n)_{n \in \mathbb{N}}$  is Farber if and only if the sequence of Schreier graphs  $\text{Sch}(\Gamma_n \backslash \Gamma, S)$  converges to the Cayley graph  $\text{Cay}(\Gamma, S)$  in the Benjamini–Schramm topology [BS01]; if and only if the sequence of actions  $\Gamma \curvearrowright \Gamma/\Gamma_n$  defines a sofic approximation of  $\Gamma$ .

Observe that  $\Gamma$  admits Farber sequences if and only if it is residually finite. The notion is designed to encompass non-normal finite index subgroups.

**Definition 10.2** (Farber neighborhood). Let  $\Gamma$  be a residually finite group. An open subset  $U$  of  $\text{Sub}_\Gamma^{fi}$  is a  $\Gamma$ -**Farber neighborhood** if it is  $\Gamma$ -invariant and every Farber sequence in  $\text{Sub}_\Gamma^{fi}$  eventually belongs to  $U$ .

We can think of these  $\Gamma$ -Farber neighborhoods as neighborhoods of  $\{\text{id}\}$  in  $\text{Sub}_\Gamma^{fi}$ , except that  $\{\text{id}\} \notin \text{Sub}_\Gamma^{fi}$ .

**Example 10.3.** Assume  $\Gamma$  is residually finite. The sets

$$U_{\Gamma, S, \delta} = \left\{ \Gamma' \in \text{Sub}_\Gamma^{fi} \mid \text{fx}_{\Gamma, \gamma}(\Gamma') < \delta \text{ for } \gamma \in S \right\},$$

where  $S \subseteq \Gamma \setminus \{1\}$  is a finite subset and  $\delta > 0$ , are non-empty and form a basis of  $\Gamma$ -Farber neighborhoods. Let  $(\gamma_j)_{j \in \mathbb{N}}$  be an enumeration of  $\Gamma$  and  $S_n = \{\gamma_0, \gamma_1, \dots, \gamma_n\}$ , then the same holds for  $U_{\Gamma, S_n, \frac{1}{n}}$ . If  $(\Gamma_n)_{n \in \mathbb{N}}$  is a Farber sequence. It follows that if  $V \subseteq \text{Sub}_\Gamma^{fi}$  does not contain any  $U_{\Gamma, S, \delta}$ , then we can construct a Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$  as above that does not meet  $V$ .

**Lemma 10.4.** Let  $\Gamma$  be a residually finite group and let  $\Lambda \leq \Gamma$  be an infinite subgroup. For every  $\Lambda$ -Farber neighborhood  $U \subseteq \text{Sub}_\Lambda^{fi}$  and  $\delta > 0$ , there exists a  $\Gamma$ -Farber neighborhood  $V \subseteq \text{Sub}_\Gamma^{fi}$  such that for any  $\Gamma' \in V$  we have

$$\frac{|\{\gamma \in \Gamma/\Gamma' : \gamma\Gamma'\gamma^{-1} \cap \Lambda \in U\}|}{[\Gamma : \Gamma']} \geq 1 - \delta.$$

In words, the lemma says that the finite index subgroups  $(\Gamma')^\gamma \cap \Lambda$  of  $\Lambda$  belong to a prescribed  $\Lambda$ -Farber neighborhood for a large proportion of the conjugates  $(\Gamma')^\gamma$ , as soon as  $\Gamma'$  belongs to a small enough  $\Gamma$ -Farber neighborhood.

*Proof.* It is enough to prove the statement for  $U = U_{\Lambda, S, \delta}$  (from Example 10.3) for any finite  $S \subseteq \Lambda$  and  $1 > \delta > 0$ .

Let  $V = U_{\Gamma, S, \frac{\delta^2}{|S|}}$ . Then, for  $\Gamma' \in V$  and each  $\gamma \in S$  simple combinatorics arguments give

$$\begin{aligned} \frac{\delta^2}{|S|} &\geq \text{fx}_{\Gamma, \gamma}(\Gamma') = \frac{1}{[\Gamma : \Gamma']} \sum_{g \in \Gamma/\Gamma'} 1_{(\Gamma')^g}(\gamma) = \frac{1}{[\Gamma : \Gamma']} \sum_{g \in \Lambda \setminus \Gamma/\Gamma'} \sum_{\lambda \in \Lambda / ((\Gamma')^g \cap \Lambda)} 1_{((\Gamma')^g \cap \Lambda)^\lambda}(\gamma) \\ &= \frac{1}{[\Gamma : \Gamma']} \sum_{g \in \Gamma/\Gamma'} \left( \frac{1}{[\Lambda : (\Gamma')^g \cap \Lambda]} \sum_{\lambda \in \Lambda / ((\Gamma')^g \cap \Lambda)} 1_{((\Gamma')^g \cap \Lambda)^\lambda}(\gamma) \right) \\ &= \frac{1}{[\Gamma : \Gamma']} \sum_{g \in \Gamma/\Gamma'} \text{fx}_{\Lambda, \gamma}((\Gamma')^g \cap \Lambda). \end{aligned}$$

Hence,

$$|\{g \in \Gamma/\Gamma' : \text{fx}_{\Lambda, \gamma}((\Gamma')^g \cap \Lambda) \geq \delta\}| \leq [\Gamma : \Gamma'] \frac{\delta}{|S|}$$

and

$$|\{g \in \Gamma/\Gamma' : \exists \gamma \in S \text{ s. t. } \text{fx}_{\Lambda, \gamma}((\Gamma')^g \cap \Lambda) \geq \delta\}| \leq [\Gamma : \Gamma'] \delta.$$

Thus

$$|\{g \in \Gamma/\Gamma' : \text{fx}_{\Lambda, \gamma}((\Gamma')^g \cap \Lambda) \leq \delta \text{ for all } \gamma \in S\}| \geq [\Gamma : \Gamma'](1 - \delta).$$

We conclude that

$$\frac{1}{[\Gamma : \Gamma']} \sum_{g \in \Gamma/\Gamma'} 1_{U_{\Lambda, S, \delta}}((\Gamma')^g \cap \Lambda) \geq 1 - \delta,$$

as desired.  $\square$

## 10.2. The cheap rebuilding property.

**Definition 10.5** (Cheap  $\alpha$ -Rebuilding Property, Farber sequences). Let  $\Gamma$  be a residually finite group and let  $\alpha$  be a non-negative integer. A Farber sequence  $(\Gamma_n)_n$  of  $\Gamma$  has the **Cheap  $\alpha$ -Rebuilding Property** if there exists a  $K(\Gamma, 1)$  space  $X$  with finite  $\alpha$ -skeleton and a constant  $\kappa_X \geq 1$  such that for every real number  $T \geq 1$ , there is  $n_0$  such that when  $n \geq n_0$  the finite covers  $Y_n \rightarrow X$  with  $\pi_1(Y_n) = \Gamma_n$  admit a  $\alpha$ -rebuilding  $(Y_n, Y'_n)$  of quality  $(T, \kappa_X)$ .

The group  $\Gamma$  itself has the **Cheap  $\alpha$ -Rebuilding Property** if the existence of the complex  $X$  and of the constant  $\kappa_X$  holds in a “uniform way” for all Farber sequences. More precisely:

**Definition 10.6** (Cheap  $\alpha$ -Rebuilding Property, groups). Let  $\Gamma$  be a countable group and let  $\alpha$  be a non-negative integer. The group  $\Gamma$  has the **Cheap  $\alpha$ -Rebuilding Property** if it is residually finite and there exists a  $K(\Gamma, 1)$  space  $X$  with finite  $\alpha$ -skeleton and a constant  $\kappa_X \geq 1$  such that for every real number  $T \geq 1$ , there exists a  $\Gamma$ -Farber neighborhood  $U = U(X, T) \subseteq \text{Sub}_\Gamma^{fi}$  such that every finite cover  $Y \rightarrow X$  with  $\pi_1(Y) \in U$  admits a  $\alpha$ -rebuilding  $(Y, Y')$  of quality  $(T, \kappa_X)$ .

*Remark 10.7.* Note that a residually finite group has the cheap 0-rebuilding property if and only if it is infinite.

**Lemma 10.8.** *One can replace in both Definitions “there exists a  $K(\Gamma, 1)$  space with finite  $\alpha$ -skeleton and there is a constant  $\kappa_X$ ” by “for every  $K(\Gamma, 1)$  space with finite  $\alpha$ -skeleton, there is a constant  $\kappa_X$ ”.*

*Proof.* Let  $X$  and  $X'$  be  $k$ -aspherical CW-complexes with finite  $\alpha$ -skeleta and  $\pi_1(X) \simeq \pi_1(X') \simeq \Gamma$ . Assume that  $X'$  satisfies the Definition 10.5, then by basic obstruction theory, there exist cellular maps

$$\mathbf{g}_1: X^{(\alpha)} \rightarrow X'^{(\alpha)} \quad \text{and} \quad \mathbf{h}_1: X'^{(\alpha)} \rightarrow X^{(\alpha)}$$

that are homotopy inverse to each other up to dimension  $\alpha - 1$ , and a cellular homotopy  $\mathbf{P}_1: [0, 1] \times X^{(\alpha-1)} \rightarrow X^{(\alpha)}$  between the identity of  $X^{(\alpha-1)}$  and the restriction of  $\mathbf{h}_1 \circ \mathbf{g}_1$  to  $X^{(\alpha-1)}$ . Observe for the case  $\alpha = 1$  that the isomorphism between the fundamental groups allows us to define  $\mathbf{g}_1$  and  $\mathbf{h}_1$  up to the 2-skeleta.

Let  $\kappa_1$  be an upper bound for all the norms  $\|(h_1)_j\|$ ,  $\|(g_1)_j\|$  for  $j \in \{0, \dots, \alpha\}$  and  $\|(\rho_1)_j\|$  for  $j = 0, 1, \dots, \alpha - 1$  that moreover satisfies  $|X'^{(j)}| \leq \kappa_1 |X^{(j)}|$  for all  $j \in \{0, \dots, \alpha\}$ . Thus  $(X, X')$  is a  $\alpha$ -rebuilding of quality  $(1, \kappa_1)$ .

For every real number  $T \geq 1$ , there exists a  $\Gamma$ -Farber neighborhood  $U = U(X', T) \subseteq \text{Sub}_\Gamma^{fi}$  such that every finite cover  $Y' \rightarrow X'$  with  $\pi_1(Y') \in U$  admits a  $\alpha$ -rebuilding  $(Y', Y'')$  of quality  $(T, \kappa_{X'})$ . Let  $Y \rightarrow X$  be the finite cover associated with  $(\mathbf{h}_1)_*(\pi_1(Y')) \leq \pi_1(X) \simeq \Gamma$ . By Lemma 6.2 (Rebuilding induced to finite cover) the  $\alpha$ -rebuilding  $(X, X')$  induces a rebuilding  $(Y, Y')$  of quality  $(1, \kappa_1 \delta_X)$ . Applying Lemma 6.3 (Composition of rebuildings) to the rebuildings  $(Y, Y')$  and  $(Y', Y'')$  we get a  $\alpha$ -rebuilding  $(Y, Y'')$  of quality  $(T, 4\kappa_1 \delta_X \kappa_{X'})$ .  $\square$

**Example 10.9.** *It is an interesting exercise to check that  $\mathbb{Z}$  has the cheap  $\alpha$ -rebuilding property for every  $\alpha$ . See Lemma 10.13 for a detailed solution. Many other groups have this cheap  $\alpha$ -rebuilding property. See Corollary 10.14 for some examples. Finite groups do not have the cheap  $\alpha$ -rebuilding property, for any  $\alpha$ .*

**Theorem 10.10.** *Let  $\Gamma$  be a finitely presented residually finite group acting on a CW-complex  $\Omega$  in such a way that any element stabilizing a cell fixes it pointwise. Let  $\alpha$  be a non-negative integer and assume that the following conditions hold:*

- (1)  $\Gamma \backslash \Omega$  has finite  $\alpha$ -skeleton;
- (2)  $\Omega$  is  $(\alpha - 1)$ -connected;
- (3) For each cell  $\omega \in \Omega$  of dimension  $j \leq \alpha$  the stabilizer  $\Gamma_\omega$  has the cheap  $(\alpha - j)$ -rebuilding property.

Then,  $\Gamma$  itself has the cheap  $\alpha$ -rebuilding property.

*Proof.* If  $\alpha = 0$ , this follows from Remark 10.7 and (3): the stabilizer of any 0-cell is infinite, thus so is  $\Gamma$ .

From now on we assume  $\alpha \geq 1$ . As in the proof of Theorem 8.1 we need to work with a simply connected  $\Omega$ . For  $\alpha \geq 2$  this follows from the assumptions. For  $\alpha = 1$ , we note that the vertex stabilizers ( $j = 0$ ) have the 1-rebuilding property and thus are finitely generated, so we can use Lemma 8.2 to replace  $\Omega$  by (its 1-skeleton and then by) a 1-connected 2-dimensional complex  $\Omega^+$  satisfying all the assumptions of the theorem.

Since all the information we need is located in the  $\alpha$ -skeleton (or 2-skeleton for  $\alpha = 1$ ), we may assume that  $\Omega$  has dimension at most  $\alpha$  (or 2 for  $\alpha = 1$ ).

Given a contractible CW-complex  $E\Gamma$  with a free action of  $\Gamma$  (i.e., the universal cover of some classifying space for  $\Gamma$ ), the Borel construction (Section 2) considers the product  $\Omega \times E\Gamma$  with the diagonal action. It is  $(\alpha - 1)$ -connected, the fundamental group of the quotient  $\Gamma \backslash (\Omega \times E\Gamma)$  is isomorphic to  $\Gamma$  and the projection map  $\Gamma \backslash (\Omega \times E\Gamma) \rightarrow \Gamma \backslash \Omega$  is interpreted (by Proposition 2.1) as a stack of CW-complexes with fiber  $\simeq \omega \times (\Gamma_\omega \backslash E\Gamma) \simeq \Gamma_\omega \backslash E\Gamma$  over each cell  $\Gamma\omega$  of  $\Gamma \backslash \Omega$ .

Let  $[\Omega] := \{\omega_1, \dots, \omega_N\} \subseteq \Omega$  be a list of distinct representatives of the  $\Gamma$ -orbits of cells of dimension  $\leq \alpha$ ; there are finitely many of them, by hypothesis (1). Let  $\omega \in [\Omega]$  be a cell of dimension  $j \leq \alpha$ . By hypothesis the stabilizer  $\Gamma_\omega$  has the cheap  $(\alpha - j)$ -rebuilding property and is therefore of type  $F_{\alpha-j}$ . Let  $X_\omega$  be a classifying space for  $\Gamma_\omega$  with finite  $(\alpha - j)$ -skeleton. Since  $X_\omega$  and  $\Gamma_\omega \backslash E\Gamma$  are homotopy equivalent, the Geoghegan's rebuilding Lemma (Proposition 2.2) yields a stack  $\Sigma \rightarrow \Gamma \backslash \Omega$  with fiber  $X_\omega$  over each cell  $\Gamma\omega$  of  $\Gamma \backslash \Omega$ . Per condition (1), the quotient  $\Gamma \backslash \Omega$  has finite  $\alpha$ -skeleton so the whole stack  $\Sigma$  has finite  $\alpha$ -skeleton. At this point we might as well forget how we constructed  $\Sigma$ , the only important properties to keep in mind are that it fits into a stack  $\Pi : \Sigma \rightarrow \Gamma \backslash \Omega$ , it has a  $(\alpha - 1)$ -connected universal cover, it has finite  $\alpha$ -skeleton, that  $\pi_1(\Sigma)$  is isomorphic to  $\Gamma$  and each fiber over  $\Gamma\omega$  is a CW-complex  $X_\omega$  with  $(\alpha - 1)$ -connected universal cover, finite  $\alpha$ -skeleton, and  $\pi_1(X_\omega) \simeq \Gamma_\omega$ .

Now let  $\Gamma_1 \leq \Gamma$  be a finite index subgroup of  $\Gamma$  and let  $\Sigma_1$  be the finite cover of  $\Sigma$  corresponding to  $\Gamma_1$ . Any  $\Gamma$ -orbit of cells  $\Gamma\omega \subseteq \Omega$  splits into a family of  $\Gamma_1$ -orbits indexed by the double cosets:

$$\Gamma\omega = \bigsqcup_{\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_\omega} \Gamma_1 \gamma \omega$$

The complex  $\Sigma_1$  is naturally the total space of a stack over  $\Gamma_1 \backslash \Omega$ . If  $\gamma\omega$  is any cell of  $\Omega$  ( $\gamma \in \Gamma$ ) and  $\Gamma_{\gamma\omega} \cap \Gamma_1 = \gamma \Gamma_\omega \gamma^{-1} \cap \Gamma_1$  is its stabilizer for the  $\Gamma_1$ -action, then the fiber of this stack over the cell  $\Gamma_1 \gamma \omega$  of  $\Gamma_1 \backslash \Omega$  takes the form  $X_{1,\gamma\omega} = (\gamma \Gamma_\omega \gamma^{-1} \cap \Gamma_1) \backslash \tilde{X}_{\gamma\omega} \simeq (\Gamma_\omega \cap \gamma^{-1} \Gamma_1 \gamma) \backslash \tilde{X}_\omega$ .

$$\begin{array}{ccccc} \tilde{\Sigma} & \longrightarrow & \Sigma_1 = \Gamma_1 \backslash \tilde{\Sigma} & \longrightarrow & \Sigma = \Gamma \backslash \tilde{\Sigma} \\ \downarrow & & \downarrow & & \downarrow \\ \Omega & \longrightarrow & \Gamma_1 \backslash \Omega & \longrightarrow & \Gamma \backslash \Omega \end{array}$$

We remark that since  $\Sigma_1$  is a cover of  $\Sigma$ , we know that the norm (induced from the  $\ell^2$ -norm) of the boundary map  $\partial : C_\bullet(\Sigma_1) \rightarrow C_{\bullet-1}(\Sigma_1)$  is bounded in degrees  $\bullet \leq \alpha$  by a constant depending only on  $\Sigma$ , not on  $\Gamma_1$ .

We will perform some rebuilding of  $\Sigma_1$  and determine conditions on  $\Gamma_1$  under which its quality is good enough.

**Step 1. Rebuilding the stack.** The cheap -rebuilding property (assumption (3)) of the stabilizers  $\Gamma_{\omega_i}$  applied to  $X_{\omega_i}$  (by Lemma 10.8, we have the freedom of the space in Definition 10.6) gives constants  $c_{X_{\omega_i}} \geq 1$  for which the following choice is possible:

Let  $T \geq 1$  be a real number. We choose for each  $i = 1, \dots, N$  a  $\Gamma_{\omega_i}$ -Farber neighborhood

$U_i \subseteq \text{Sub}_{\Gamma_{\omega_i}}^{f_i}$  and for every finite index subgroup  $\Lambda \leq \Gamma_{\omega_i}$  with  $\Lambda \in U_i$  we choose an  $(\alpha - \dim(\omega_i))$ -rebuilding  $R(\omega_i, T, \Lambda)$  of  $\Lambda \backslash \tilde{X}_{\omega_i}$  of quality  $(T, \kappa_{X_{\omega_i}})$ .

We want to use the effective rebuilding Lemma (Proposition 4.2) to rebuild the stack  $\Sigma_1 = \Gamma_1 \backslash \tilde{\Sigma}$ . To do that, we need to specify a rebuilding of each fiber  $X_{1,\omega}$ , with  $\Gamma\omega \subset \Gamma \backslash \Omega$ . The full set of fibers of  $\Sigma_1 \rightarrow \Gamma_1 \backslash \Omega$  consists of the CW-complexes  $X_{1,\gamma\omega_i}$ , with  $i = 1, \dots, N$  and  $\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_{\omega_i}$ . Note that

$$X_{1,\gamma\omega_i} = (\Gamma_{\gamma\omega_i} \cap \Gamma_1) \backslash \tilde{X}_{\gamma\omega_i} \simeq (\Gamma_{\omega_i} \cap \gamma^{-1}\Gamma_1\gamma) \backslash \tilde{X}_{\omega_i}.$$

We rebuilt  $X_{1,\gamma\omega_i}$  according to whether  $\Gamma_{1,\omega_i,\gamma} := (\Gamma_{\omega_i} \cap \gamma^{-1}\Gamma_1\gamma) \in U_i \subseteq \text{Sub}_{\Gamma_{\omega_i}}^{f_i}$  or not by using the rebuilding:

$$(X_{1,\gamma\omega_i}, X'_{1,\gamma\omega_i}, \mathbf{k}_{\gamma\omega_i}, \mathbf{l}_{\gamma\omega_i}, \Sigma_{\gamma\omega_i}) := \begin{cases} R(\omega_i, T, \Gamma_{1,\omega_i,\gamma}) & \text{if } \Gamma_{1,\omega_i,\gamma} \in U_i \subseteq \text{Sub}_{\Gamma_{\omega_i}}^{f_i} \\ X'_{1,\gamma\omega_i} = X_{1,\gamma\omega_i} & \text{if } \Gamma_{1,\omega_i,\gamma} \notin U_i \end{cases} \quad (10.1)$$

i.e., we simply use the trivial rebuilding  $(X_{1,\gamma\omega_i}, X_{1,\gamma\omega_i}, id, id, 0)$  when  $\Gamma_{1,\omega_i,\gamma} \notin U_i$ .

By virtue of Proposition 4.2 we obtain a global rebuilding  $(\Sigma_1, \Sigma'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$ . Our goal is now to estimate its quality. Recall that the tension in Definition 2 of quality (in the introduction) is between “having few cells” and “maintaining tame norms”. Observe that for the fibers associated with  $\Gamma_{1,\omega_i,\gamma} \in U_i$ , both are controlled by definition of the  $\Gamma_{\omega_i}$ -Farber neighborhood. In particular, the norms of the vertical maps in these fibers are bounded by a polynomial in  $T$ . As for the fibers associated with  $\Gamma_{1,\omega_i,\gamma} \notin U_i$ , the quality is “very good” as far as the “norms bound” is concerned. The number of cells will be controlled later on in Step 2.

Let us simply denote by  $k: C_\bullet(X_{1,\omega}) \rightarrow C_\bullet(X'_{1,\omega})$  and  $l: C_\bullet(X'_{1,\omega}) \rightarrow C_\bullet(X_{1,\omega})$  the maps respectively induced by  $\mathbf{k}_\omega$  and  $\mathbf{l}_\omega$ . Similarly, let  $\sigma: C_\bullet(X_{1,\omega}) \rightarrow C_{\bullet+1}(X_{1,\omega})$  be the chain homotopy map induced by  $\Sigma_\omega$ . Finally let  $g: C_\bullet(\Sigma_1) \rightarrow C_\bullet(\Sigma'_1)$  and  $h: C_\bullet(\Sigma'_1) \rightarrow C_\bullet(\Sigma_1)$  be the chain maps induced by  $\mathbf{g}$  and  $\mathbf{h}$  and let  $\rho: C_\bullet(\Sigma_1) \rightarrow C_{\bullet+1}(\Sigma_1)$  be the chain homotopy map induced by  $\mathbf{P}$ .

As in the paragraphs preceding and preparing to Proposition 4.2, we consider the decomposition

$$C_\bullet(\Sigma_1) = \bigoplus_{\Gamma_1\omega \in \Gamma_1 \backslash \Omega} [\omega] \otimes C_\bullet(X_{1,\omega}) \quad \text{and} \quad C_\bullet(\Sigma'_1) = \bigoplus_{\Gamma_1\omega \in \Gamma_1 \backslash \Omega} [\omega] \otimes C_\bullet(X'_{1,\omega}).$$

By a slight abuse of notation we will simply write  $\heartsuit$  instead of the  $1 \otimes \heartsuit$  for  $\heartsuit = k, k_\omega, l, l_\omega, \sigma$  or  $\sigma_\omega$ . We write  $\partial, \partial^{\text{hor}}, \partial^{\text{vert}}$  for the boundary, horizontal boundary and vertical boundary maps in the stack  $C_\bullet(\Sigma_1)$  and  $\partial', (\partial')^{\text{hor}}, (\partial')^{\text{vert}}$  for the boundary, horizontal boundary and vertical boundary maps on  $C_\bullet(\Sigma'_1)$ . By Proposition 4.2 formulae (4.7)–(4.10) we have

$$\begin{aligned} g &= k \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right), \\ h &= \left( \sum_{i=0}^{\infty} (\sigma \circ \partial^{\text{hor}})^i \right) \circ l, \\ \rho &= \sigma \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right), \\ \partial' &= (\partial')^{\text{vert}} + k \circ \left( \sum_{i=0}^{\infty} (\partial^{\text{hor}} \circ \sigma)^i \right) \circ \partial^{\text{hor}} \circ l. \end{aligned}$$

Recall that in each of these expressions the internal sum is in fact finite since the summands vanish for  $i$  large enough. The norms of  $(\partial')^{\text{vert}}, k, l, \sigma$  and  $\partial^{\text{hor}}$  are bounded by polynomials in  $T$  with coefficients depending on  $\Sigma$  and on the constants  $\kappa_{X_{\omega_i}}$ , but independent of  $\Gamma_1$ . This is thus also the case for the norms of  $g, h, \rho$  and  $\partial'$  the chain maps appearing in the definition of the quality

of a rebuilding. Their log is thus bounded by a constant (independent of  $\Gamma_1$ ) times  $(1 + \log T)$ . This ensures the norms bound condition for every  $\Gamma_1$ .

**Step 2. Counting cells.** It remains to ensure a control on the number of cells of  $\Sigma'_1$ . Let  $\#^{(d)}(X)$  denote the number of  $d$ -cells of  $X$ . We need to exhibit a  $\Gamma$ -Farber neighborhood  $V \subseteq \text{Sub}_\Gamma^{fi}$  such that if  $\Gamma_1$  belongs to  $V$  then

$$\#^{(j)}(\Sigma'_1) \leq O(\#^{(j)}(\Sigma_1)/T) \quad \text{for each } j \in \{0, \dots, \alpha\}.$$

Given an arbitrary finite index  $\Gamma_1 \leq \Gamma$ , we intend to count the number of cells in  $\Sigma'_1$ . For that purpose we introduce  $N$  functions  $F_i : \Gamma \rightarrow \mathbb{R}$ , with  $i \in \{1, \dots, N\}$ , corresponding to the chosen distinct representatives  $[\Omega] := \{\omega_1, \dots, \omega_N\} \subseteq \Omega$  of the  $\Gamma$ -orbits of cells in  $\Omega$ , by the following formulas

$$F_i(\gamma) = \begin{cases} T^{-1} & \text{if } (\Gamma_{\omega_i} \cap \gamma^{-1}\Gamma_1\gamma) \in U_i \\ 1 & \text{otherwise,} \end{cases}$$

where  $U_i$  is the  $\Gamma_{\omega_i}$ -Farber neighborhood chosen at the beginning of Step 1. Denoting  $q := \dim(\omega_i)$ , recall to that each  $\Lambda \in U_i \subseteq \text{Sub}_{\Gamma_{\omega_i}}^{fi}$  we have associated a cheap  $(\alpha - q)$ -rebuilding  $(X_{1,\gamma\omega_i}, X'_{1,\gamma\omega_i})$  in equation (10.1) of quality  $(T, \kappa_{X_{\omega_i}})$ . In particular, the cells bound gives  $\#^{(\ell)}(X'_{1,\gamma\omega_i}) \leq \kappa_{X_{\omega_i}} \times F_i(\gamma) \times \#^{(\ell)}(X_{1,\gamma\omega_i})$ , for every  $\ell \in \{0, 1, \dots, \alpha - q\}$ .

By our choices of fibers rebuildings we thus get:

$$\begin{aligned} \#^{(j)}(\Sigma'_1) &= \sum_{q=0}^j \sum_{\substack{\omega \in \Gamma_1 \setminus \Omega \\ \dim(\omega)=q}} \#^{(j-q)}(X'_{1,\omega}) = \sum_{\substack{\omega \in \Gamma_1 \setminus \Omega \\ \dim(\omega) \leq j}} \#^{(j-\dim(\omega))}(X'_{1,\omega}) \\ &= \sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} \sum_{\gamma \in \Gamma_1 \setminus \Gamma/\Gamma_{\omega_i}} \#^{(j-q)}(X'_{1,\gamma\omega_i}) \\ &\leq \sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} \sum_{\gamma \in \Gamma_1 \setminus \Gamma/\Gamma_{\omega_i}} \kappa_{X_{\omega_i}} \times F_i(\gamma) \times \#^{(j-q)}(X_{1,\gamma\omega_i}) \\ &\leq \kappa^\Omega \sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} \left( \underbrace{\sum_{\gamma \in \Gamma_1 \setminus \Gamma/\Gamma_{\omega_i}} \frac{\#^{(j-q)}(X_{1,\gamma\omega_i})}{T}}_{(A)} + \left(1 - \frac{1}{T}\right) \underbrace{\sum_{\substack{\gamma \in \Gamma_1 \setminus \Gamma/\Gamma_{\omega_i} \\ F_i(\gamma)=1}} \#^{(j-q)}(X_{1,\gamma\omega_i})}_{(B)} \right) \end{aligned}$$

where  $\kappa^\Omega := \max\{\kappa_{X_{\omega_i}} : i \in \{1, \dots, N\}\}$ .

In order to bound the part associated with (A), observe that

$$\sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} \sum_{\gamma \in \Gamma_1 \setminus \Gamma/\Gamma_{\omega_i}} \#^{(j-q)}(X_{1,\gamma\omega_i}) = \#^{(j)}(\Sigma_1).$$

In order to bound the part associated with (B), observe that

- the covering  $\Sigma_1 \rightarrow \Sigma$  induces the covering  $X_{1,\gamma\omega_i} \rightarrow X_{\omega_i}$ ; and thus

$$\#^{(j-q)}(X_{1,\gamma\omega_i}) = [\Gamma_{\gamma\omega_i} : \Gamma_{\gamma\omega_i} \cap \Gamma_1] \#^{(j-q)}(X_{\omega_i});$$



- $[\Gamma_{\gamma\omega_i} : \Gamma_{\gamma\omega_i} \cap \Gamma_1] = [\Gamma_{\omega_i} : \gamma^{-1}\Gamma_1\gamma \cap \Gamma_{\omega_i}]$  is exactly the number of  $\Gamma_1 \backslash \Gamma$ -classes that are gathered together to form the  $\Gamma_1 \backslash \Gamma / \Gamma_{\omega_i}$ -class of  $\gamma$ ;
- the invariance under conjugation of the  $\Gamma_{\omega_i}$ -Farber neighborhoods ensures that  $F_i(\gamma) = F_i(\gamma\lambda)$  for every  $\lambda \in \Gamma_{\omega_i}$ . It follows that

$$\sum_{\substack{\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_{\omega_i} \\ F_i(\gamma)=1}} \#^{(j-q)}(X_{1,\gamma\omega_i}) = \sum_{\substack{\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_{\omega_i} \\ F_i(\gamma)=1}} [\Gamma_{\gamma\omega_i} : \Gamma_{\gamma\omega_i} \cap \Gamma_1] \#^{(j-q)}(X_{\omega_i}) = \sum_{\substack{\gamma \in \Gamma_1 \backslash \Gamma \\ F_i(\gamma)=1}} \#^{(j-q)}(X_{\omega_i}).$$

By Lemma 10.4, there exists a  $\Gamma$ -Farber neighborhood  $V \subseteq \text{Sub}_\Gamma^{fi}$  such that for  $\Gamma_1 \in V$  and  $i \in \{1, \dots, N\}$  we have

$$|\{\gamma \in \Gamma_1 \backslash \Gamma : F_i(\gamma) = 1\}| = |\{\gamma \in \Gamma / \Gamma_1 : \gamma\Gamma_1\gamma^{-1} \cap \Gamma_{\omega_i} \notin U_i\}| \leq T^{-1}[\Gamma : \Gamma_1].$$

Thus

$$\begin{aligned} \sum_{\substack{\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_{\omega_i} \\ F_i(\gamma)=1}} \#^{(j-q)}(X_{1,\gamma\omega_i}) &\leq T^{-1}[\Gamma : \Gamma_1] \#^{(j-q)}(X_{\omega_i}) \\ \sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} \sum_{\substack{\gamma \in \Gamma_1 \backslash \Gamma / \Gamma_{\omega_i} \\ F_i(\gamma)=1}} \#^{(j-q)}(X_{1,\gamma\omega_i}) &\leq \underbrace{\sum_{q=0}^j \sum_{\substack{\omega_i \in [\Omega] \\ \dim(\omega_i)=q}} T^{-1}[\Gamma : \Gamma_1] \#^{(j-q)}(X_{\omega_i})}_{=T^{-1}\#^{(j)}(\Sigma_1)}. \end{aligned}$$

This finishes the proof of Theorem 10.10.  $\square$

As particular cases of Theorem 10.10 we obtain:

**Example 10.11** (Graphs of groups). *Let  $\Gamma$  be a residually finite group that splits as a finite graph of groups with edge and vertex stabilizers that satisfy the cheap  $(\alpha - 1)$ -rebuilding property (and cheap  $\alpha$ -rebuilding property respectively) for every  $\alpha$ . Considering its Bass-Serre tree  $\Omega$ , we obtain that  $\Gamma$  itself has the cheap  $\alpha$ -rebuilding property. This applies for instance to the residually finite Baumslag-Solitar groups  $\text{BS}(1, n)$  and  $\text{BS}(n, n)$ , for any non-zero integer  $n$ .*

**Example 10.12** (Groups acting on graphs). *If a residually finite group  $\Gamma$  acts co-compactly on a connected graph  $\mathcal{G}$  such that*

- *vertex stabilizers have the cheap 1-rebuilding property;*
- *edge stabilizers are infinite,*

*then  $\Gamma$  has the cheap 1-rebuilding property.*

### 10.3. First example.

**Lemma 10.13.**  *$\mathbb{Z}$  has the cheap  $\alpha$ -rebuilding property for any  $\alpha$ .*

*Proof.* We will verify the property for  $X = [0, 1] / \{0 \sim 1\}$  being the circle with one 0-cell and one 1-cell. Given a positive integer  $m$  we denote by  $X_m = [0, m] / \{0 \sim m\}$  the  $m$ -fold cover of  $X$  associated to the subgroup  $m\mathbb{Z}$  of  $\mathbb{Z}$  and equipped with the cell structure lifted from  $X$ .

Let  $T$  be a positive real number. The subset  $U \subset \text{Sub}_{\mathbb{Z}}^{fi}$  that consists of all the finite index subgroups  $N\mathbb{Z}$  in  $\mathbb{Z}$  with  $N \geq 4T$  is a  $\mathbb{Z}$ -Farber neighborhood. Pick some subgroup  $N\mathbb{Z}$  in  $U$ . We now explain how to construct a rebuilding of quality  $(T, O(1))$  of  $X_N$ . To do so, first pick some sequence of integers

$$0 = a_0 < a_1 \dots < a_m = N \quad \text{with} \quad T/2 \leq a_{i+1} - a_i \leq T.$$

For each  $t \in [0, N]$  let  $\iota(t)$  be the integer in  $[0, m-1]$  determined by  $t \in [a_{\iota(t)}, a_{\iota(t)+1})$ . The cellular maps  $\mathbf{g}: X_N \rightarrow X_m$  and  $\mathbf{h}: X_m \rightarrow X_N$  defined by

$$\mathbf{g}(t) = \iota(t) + \min\{t - a_{\iota(t)}, 1\} \quad \text{and} \quad \mathbf{h}(t) = a_{\lfloor t \rfloor} + (a_{\lfloor t \rfloor + 1} - a_{\lfloor t \rfloor})(t - \lfloor t \rfloor)$$

are homotopy inverses to each other. The explicit homotopy  $\mathbf{P}: X_N \times [0, 1] \rightarrow X_N$  between 1 and  $\mathbf{h} \circ \mathbf{g}$  is given by

$$\mathbf{P}(t, s) = \min\{a_{\iota(t)} + (t - a_{\iota(t)})(1 + s(a_{\iota(t)+1} - a_{\iota(t)} - 1)), a_{\iota(t)+1}\}.$$

One easily verifies that the  $\ell^2$ -operator norms of the induced chain maps satisfy

$$\|g\|, \|h\|, \|\rho\| = O(T + 1) \text{ and } \|\partial\|, \|\partial'\| \leq 2.$$

The number of cells in  $X_m$  is  $m \in [N/T, 2N/T]$  in dimension 0 and 1. Hence,  $\mathbb{Z}$  has the cheap  $\alpha$ -rebuilding property for any  $\alpha \in \mathbb{N}$ .  $\square$

**10.4. Applications.** Recall that a group  $\Gamma$  is called **polycyclic** if there is a sequence of subgroups

$$\Gamma = A_0 \triangleright A_1 \triangleright \cdots \triangleright A_n = \{1\}$$

such that each quotient group  $A_i/A_{i+1}$  is cyclic. As a corollary of Theorem 10.10 and Lemma 10.13 we obtain

**Corollary 10.14.** *Let  $\Gamma$  be a residually finite countable group. The following holds:*

- (1) *Let  $\Gamma' \leq \Gamma$  be a finite index subgroup. Then  $\Gamma$  has the cheap  $\alpha$ -rebuilding property if and only if  $\Gamma'$  does.*
- (2) *If  $\Gamma$  has an infinite normal subgroups  $N$  such that  $\Gamma/N$  is of type  $F_\alpha$  and  $N$  has the cheap  $\alpha$ -rebuilding property, then  $\Gamma$  has the cheap  $\alpha$ -rebuilding property.*
- (3)  *$\mathbb{Z}^m$  has the cheap  $\alpha$ -rebuilding property for every  $\alpha$ .*
- (4) *Infinite polycyclic groups have the cheap  $\alpha$ -rebuilding property for every  $\alpha$ .*

*Proof.* 1. If  $\Gamma$  has the cheap  $\alpha$ -rebuilding property, then the witnesses  $(X, \kappa, U(T))$  for  $\Gamma$  yield witnesses for  $\Gamma'$ . Conversely, if  $\Gamma'$  has the cheap  $\alpha$ -rebuilding property one can assume, up to passing to a further finite index subgroup, that it is normal in  $\Gamma$ . Then pick any  $(\alpha - 1)$ -connected CW-complex  $\Omega$  with a free action of  $\Gamma/\Gamma'$  that is finite in every dimension  $\leq \alpha$  and apply Theorem 10.10 to the  $\Gamma$ -action on  $\Omega$  defined through the quotient map  $\Gamma \rightarrow \Gamma/\Gamma'$ .

2. Let  $B$  be a classifying space (CW-complex)  $B$  for  $\Gamma/N$  with finite  $\alpha$ -skeleton. The group  $\Gamma$  acts on its universal cover  $\tilde{B}$  with cell-stabilizers all equal to  $N$  and one can apply Theorem 10.10 to this action.

3. and 4. finally follow from Lemma 10.13 and 2. by induction.  $\square$

**Example 10.15.** *Corollary 10.14, item (2) implies for instance that  $\mathrm{SL}_d(\mathbb{Z}) \rtimes \mathbb{Z}^d$  has the cheap  $\alpha$ -rebuilding property for every  $\alpha$ . The same is true for the standard braid groups since it has an infinite cyclic center.*

**10.5. Right-angled groups.** The following proposition connects the cheap 1-rebuilding property to the rewirings of right-angled groups considered in [AGN17].

**Proposition 10.16.** *Let  $\Gamma$  be a residually finite group that is **right-angled**, i.e., that admits a finite generating list  $\{\gamma_1, \dots, \gamma_m\}$  of elements of infinite order such that  $[\gamma_i, \gamma_{i+1}] = 1$  for all  $i \in \{1, \dots, m - 1\}$ . Then  $\Gamma$  has the cheap 1-rebuilding property.*

*Proof of Proposition 10.16.* According to Example 10.12 it suffices to produce a cocompact action  $\Gamma \curvearrowright \mathcal{G}$  on a connected graph such that

- vertex stabilizers have the 1-rebuilding property, and
- edge stabilizers are infinite.

We build the graph  $\mathcal{G}$  as follows: for each  $i \in \{1, \dots, m\}$ , let  $H_i$  be the (free abelian) infinite subgroup of  $\Gamma$  generated by  $\gamma_i$ . The vertex set of  $\mathcal{G}$  consists of all the conjugates of the  $H_i$  in  $\Gamma$ , and two distinct conjugates  $gH_i g^{-1}$  and  $hH_j h^{-1}$  represent an edge if and only if the subgroup they generate is abelian.

The conjugation action of  $\Gamma$  on the set of its subgroups induces an action of  $\Gamma$  on  $\mathcal{G}$ . Observe that, by hypothesis, the vertices corresponding to  $H_1, \dots, H_m$  belong to the same connected component of  $\mathcal{G}$ . It follows that for every  $g \in \Gamma$  the vertices  $gH_1g^{-1}, \dots, gH_mg^{-1}$  also belong to one connected component of  $\mathcal{G}$ . Now for any  $j \in \{1, \dots, m\}$  we have

$$gH_jg^{-1} = (g\gamma_j^{-1})H_j(g\gamma_j^{-1})^{-1},$$

and, since  $\{\gamma_1, \dots, \gamma_m\}$  is a generating set of  $\Gamma$ , we conclude that the graph  $\mathcal{G}$  is connected.

The stabilizer of each vertex, or each edge, contains a normal subgroup that is a free infinite abelian subgroup (a conjugate of some  $H_j$ ). These stabilizers therefore have the cheap 1-rebuilding property (by Corollary 10.14(2) and Lemma 10.13).

Finally, the quotient  $\Gamma \backslash \mathcal{G}$  is a finite graph (on  $m$  vertices). Theorem 10.10 applies to the action of  $\Gamma$  on  $\mathcal{G}$  and we conclude that  $\Gamma$  itself has the cheap 1-rebuilding property.  $\square$

The cheap  $\alpha$ -rebuilding property goes beyond right-angled groups, even for  $\alpha = 1$ . We now give an example.

**Example 10.17** (Non right-angled examples  $\Lambda \ltimes \mathbb{Z}^2$ ). *If a free group  $\Lambda \leq \mathrm{SL}_2(\mathbb{Z})$  does not contain any non-trivial unipotent element, then any embedding of  $\mathbb{Z}^2$  in the canonical semi-direct product  $\Lambda \ltimes \mathbb{Z}^2$  lies inside the obvious normal  $\mathbb{Z}^2$ : If  $(\lambda_1, a_2), (\lambda_2, a_2) \in \Lambda \ltimes \mathbb{Z}^2$  commute (and generate  $\mathbb{Z}^2$ ), then up to considering some power of them, one can assume  $\lambda_1 = \lambda_2$ . It follows that  $\lambda_1^{-1}a_2^{-1}a_1\lambda_1 = a_2^{-1}a_1$ , so that  $\lambda_1$  is unipotent, thus trivial.*

*In order to find such a subgroup  $\Lambda$  of  $\mathrm{SL}_2(\mathbb{Z})$ , it's enough to consider a finite cover of the modular orbifold  $\mathrm{SL}_2(\mathbb{Z}) \backslash \mathbb{H}^2$  that is a surface  $S$  of genus  $g \geq 2$  (with cusps). The fundamental group of the compact surface  $S'$  obtained by adding one point to each cusp of  $S$  surjects onto the free group  $\mathbf{F}_2$ . Finally pick a pull-back of this free group along the maps  $\mathrm{SL}_2(\mathbb{Z}) \geq \pi_1(S) \rightarrow \pi_1(S') \rightarrow \mathbf{F}_2$ . It does not contain any non-trivial unipotent element.*

*Such a semi-direct product  $\Gamma = \Lambda \ltimes \mathbb{Z}^2$  is thus not right-angled while it has the cheap  $\alpha$ -rebuilding property for all  $\alpha$  by Corollary 10.14 (2).*

**10.6. Artin groups.** Let  $I$  be a finite set. A **Coxeter matrix**  $M = (m_{ij})$  on  $I$  is an  $I$ -by- $I$  symmetric matrix  $M = (m_{ij})$  with entries in  $\mathbb{N} \cup \{\infty\}$  such that  $m_{ii} = 1$  for all  $i$  and such that whenever  $i \neq j$ ,  $m_{ij} \geq 2$ . Associated to  $M$  are

- a **Coxeter group**  $W_I$ , or simply  $W$ , given by the presentation

$$W = \langle s_i, i \in I \mid (s_i s_j)^{m_{ij}} = 1, (i, j) \in I^2 \rangle$$

with the convention that the relation is ignored if  $m_{ij} = \infty$ ,

- a **Artin group**  $A_I$ , or simply  $A$ , given by the presentation

$$A = \langle a_i, i \in I \mid \underbrace{a_i a_j a_i a_j a_i \cdots}_{m_{ij}} = \underbrace{a_j a_i a_j a_i a_j \cdots}_{m_{ij}}, (i, j) \in I^2 \rangle, \quad \text{and}$$

- a simplicial complex, called the **nerve** of  $M$ , that we define below.

If the corresponding Coxeter group  $W$  is finite we say that  $A$  is **spherical**. These Artin groups are sometimes also referred to as ‘Artin groups of finite type’ in the literature.

If  $J \subseteq I$ , let  $M_J$  denote the minor of  $M$  whose rows and columns are indexed by  $J$ , and let  $W_J$ , resp.  $A_J$ , be the corresponding Coxeter group, resp. Artin group. It is known [Bou81] that the natural map  $W_J \rightarrow W_I$ , resp.  $A_J \rightarrow A_I$ , is injective and hence  $W_J$ , resp.  $A_J$ , can be identified with the subgroup of  $W_I$ , resp.  $A_I$ , generated by  $\{s_i : i \in J\}$ , resp.  $\{a_i : i \in J\}$ .

The **nerve** of  $M$  is the simplicial complex  $L$  whose vertex set is  $I$  and a subset  $\sigma \subseteq I$  spans a simplex if and only if  $W_\sigma$  is finite.

Charney and Davis [CD95, Section 3] have associated to  $M$  a simple complex of groups  $\mathcal{C}$ , in the sense of [BH99, Definition 12.11]. The construction goes as follows: first consider the partially

ordered set (poset)

$$\mathcal{P} = \{J \subseteq I : W_J \text{ is finite}\}$$

ordered by inclusion. To any  $\sigma \in \mathcal{P}$  one associates the Artin group  $A_\sigma$ , and for each  $\tau \leq \sigma$  the associate homomorphism  $A_\tau \rightarrow A_\sigma$  is the natural inclusion. The geometric realization of  $\mathcal{C}$  is simply connected (in fact contractible), since the empty set is an initial object for the poset  $\mathcal{P}$ . In particular, the fundamental group of the complex of groups  $\mathcal{C}$  is  $A$  (the direct limit of the  $A_\sigma$ ). The complex of groups  $\mathcal{C}$  is developable, that is, it arises from the action of a group on a simplicial complex. More precisely, let

$$A\mathcal{P} = \{gA_J : g \in A, J \in \mathcal{P}\},$$

ordered by inclusion. The stabilizer of  $gA_J \in A\mathcal{P}$  is the subgroup  $gA_Jg^{-1}$ ; hence,  $A$  acts without inversion on  $A\mathcal{P}$ . Let

$$X = |A\mathcal{P}'|$$

be the geometric realization of the derived poset of  $A\mathcal{P}$ ; it is the simplicial complex, referred to as the ‘modified Deligne complex’ in [CD95], where  $k$ -simplices correspond to totally ordered chains of elements in  $A\mathcal{P}$ . The complex of groups  $\mathcal{C}$  explicitly arises from the action of  $A$  on  $X$ .

It is conjectured that  $X$  is contractible and this conjecture is equivalent to the celebrated  $K(\pi, 1)$  **conjecture** for Artin groups. The main theorem of [CD95] is that this conjecture holds when the nerve  $L$  is a **flag complex**, i.e., when the Coxeter group  $W_J$  associated to every clique  $J$  is finite. This holds for instance if  $A$  is either spherical or a right-angled Artin group. The conjecture also holds whenever  $L$  has dimension at most 1. Recently, Paolini and Salvetti proved that the  $K(\pi, 1)$  conjecture holds for all affine Artin Groups [PS21].

**Theorem 10.18** (Artin groups with  $(\alpha - 1)$ -connected nerve). *Let  $M$  be a Coxeter matrix whose nerve  $L$  is  $(\alpha - 1)$ -connected. Suppose that the  $K(\pi, 1)$ -conjecture holds for the associated Artin group  $A$  and that  $A$  is residually finite. Then  $A$  has the cheap  $\alpha$ -rebuilding property.*

*Proof.* We first consider the case where  $A$  is spherical, or equivalently where  $L$  is a simplex. In [Bes99, Main Theorem], Bestvina proves that  $A$  is then commensurable with  $G \times \mathbb{Z}$ , where  $G = A/\langle \Delta^2 \rangle$  is the quotient of  $A$  by some central element  $\Delta^2$ , and moreover  $G$  acts cocompactly on a contractible simplicial complex  $X$  and the action is transitive on the vertices with stabilizers  $\mathbb{Z}/2\mathbb{Z}$ . It follows that  $G$  is of type  $F_\alpha$  for every  $\alpha$  (see e.g., [Geo08, Theorem 7.3.1]). Applying Corollary 10.14 (2) we get that  $G \times \mathbb{Z}$  has the cheap  $\alpha$ -rebuilding property for every  $\alpha$ . By Corollary 10.14 (1), so has  $A$ .

Consider now the case where  $A$  is of general type. We would like to apply our Theorem 10.10 to the action of  $A$  on the modified Deligne complex  $X$ ; but the stabilizers of the vertices of the form  $A_J$  for  $J = \emptyset$  are trivial. We thus modify  $X$  as follows: Let  $\Omega \subset X$  be the sub-complex associated to the sub-poset  $\mathcal{P}_0$  of  $\mathcal{P}$  consisting of the non-empty subsets  $J \subseteq I$ . The  $A$ -action on  $X$  preserves  $\Omega$  and the cell stabilizers are now isomorphic to the non trivial spherical Artin groups  $A_J$ ,  $J \neq \emptyset$ . The complement  $X \setminus \Omega$  is the disjoint union of the open stars  $St(v)$  of the vertices  $v$  of  $X$  corresponding to the elements  $gA_\emptyset \in A\mathcal{P}$  ( $g \in A$ ). Each such star  $St(v)$  is the cone over the link of  $v$  in  $X$  and this link is isomorphic to the simplicial complex  $|\mathcal{P}'_0|$  associated to the poset of non-empty spherical subsets of  $I$ , i.e., the nerve  $L$  of  $M$ . By hypothesis the latter is  $(\alpha - 1)$ -connected and  $X$  is contractible, it therefore follows from Mayer-Vietoris and Hurewicz Theorems (and van Kampen Theorem in degree 1) that  $\Omega$  is  $(\alpha - 1)$ -connected. Then a direct application of Theorem 10.10 shows that the Artin group  $A$  has the cheap  $\alpha$ -rebuilding property as soon as it is residually finite.  $\square$

**Example 10.19.** *The direct product  $\Gamma = \mathbf{F}_2 \times \mathbf{F}_2 \times \cdots \times \mathbf{F}_2$  of  $k$  copies of the free group on 2 generators has the cheap  $(k - 1)$ -rebuilding property. This follows from Theorem 10.18 when  $k \geq 2$ . The nerve of these right-angled Artin groups has the homotopy type of a  $(k - 1)$ -dimensional sphere.*

*Remark 10.20.* In fact, to obtain Theorem 10.18, instead of the  $K(\pi, 1)$  conjecture, it's enough to assume that the modified Deligne complex  $X$  is  $(\alpha - 1)$ -connected: this is the condition in the proof for  $\Omega$  to be  $(\alpha - 1)$ -connected.

### 10.7. Homology growth.

**Theorem 10.21.** *Let  $\Gamma$  be a finitely presented residually finite group of type  $F_{\alpha+1}$  that has the cheap  $\alpha$ -rebuilding property for some non-negative integer  $\alpha$ . Then, for every Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$ , coefficient field  $K$  and  $0 \leq j \leq \alpha$  we have*

$$\lim_{n \rightarrow \infty} \frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\text{tors}}|}{[\Gamma : \Gamma_n]} = 0.$$

*Proof.* Let  $X$  be a  $K(\Gamma, 1)$  CW-complex with finite  $\alpha$ -skeleton. Let  $T \geq 1$ . Then there exists a  $\Gamma$ -Farber neighborhood  $U \subseteq \text{Sub}_\Gamma^{fi}$  such that for any finite index subgroup  $\Gamma_1 \in U$  there exists a  $\alpha$ -rebuilding  $(X_1 = \Gamma_1 \backslash \tilde{X}, X'_1)$  of quality  $(T, O(1))$ . In degree  $\leq \alpha$ , the complex  $X'_1$  has  $O([\Gamma : \Gamma_1]/T)$  cells of dimension and the norms of the boundary maps are  $O(T + 1)$ . It follows from (9.1) that for all  $j \leq \alpha$  we have

$$\dim_K H_j(\Gamma_n, K) = O([\Gamma : \Gamma_1]T^{-1}).$$

Letting  $T \rightarrow \infty$  this proves the first part of the theorem. By Proposition 9.1 we similarly get

$$\log |H_j(\Gamma_1, \mathbb{Z})_{\text{tors}}| \leq O([\Gamma : \Gamma_1]T^{-1} \log(T + 1)),$$

for  $j = 0, \dots, \alpha - 1$ . Letting  $T \rightarrow \infty$  this proves the second part of the theorem for all  $0 \leq j \leq \alpha - 1$ .

To handle the remaining case  $j = \alpha$ , we take advantage of the fact that in Proposition 9.1 the only information needed from dimension  $\alpha + 1$  is the norm of the boundary maps  $\partial_{\alpha+1}$  (but not the number of  $(\alpha + 1)$ -cells) and that the norm of the boundary maps is bounded when taking coverings. We proceed exactly as in the end of the proof of Theorem 8.1: by adding finitely many  $(\alpha + 1)$ -cells to  $X$ , we can make it  $\alpha$ -aspherical (see [Geo08, Theorem 8.2.1]). Write  $\Phi = \sqcup_I \mathbb{B}^{\alpha+1}$  for this collection of  $(\alpha + 1)$ -cells and let  $f: \partial\Phi = \sqcup_I \mathbb{S}^\alpha \rightarrow X^{(\alpha)}$  be the map that attaches the  $(\alpha + 1)$ -cells to  $X$ . Then  $X^+ := X \sqcup_f \Phi$  is a finite  $\alpha$ -aspherical CW-complex with fundamental group  $\Gamma$ .

Let  $T, U$  be as above and let  $\Gamma_1 \in U$  be a finite index subgroup. Let  $X_1 = \Gamma_1 \backslash \tilde{X}$  and let  $(X_1, X'_1, \mathbf{g}, \mathbf{h}, \mathbf{P})$  be an  $\alpha$ -rebuilding of quality  $(T, O(1))$  given by the rebuilding property of  $\Gamma$ . We also consider  $X_1^+ = \Gamma_1 \backslash \tilde{X}^+$ . The CW-complex  $X_1^+$  can be written as  $X_1^+ = X_1 \sqcup_{f_1} \Phi_1$  where  $\Phi_1 = \sqcup_{I_1} \mathbb{B}^{\alpha+1}$  is the preimage of  $\Phi$  in  $X_1^+$  and  $f_1: \partial\Phi_1 = \sqcup_{I_1} \mathbb{S}^\alpha \rightarrow X_1^{(\alpha)}$  is the lift of  $f$ . We have a diagram

$$\begin{array}{ccccc} \Phi_1 & \longleftarrow & \partial\Phi_1 & \xrightarrow{f_1} & X_1 \\ \text{id} \updownarrow \text{id} & & \text{id} \updownarrow \text{id} & & \mathbf{g} \updownarrow \mathbf{h} \\ \Phi_1 & \longleftarrow & \partial\Phi_1 & \xrightarrow{\varphi} & X'_1, \end{array}$$

with  $\varphi = \mathbf{g} \circ f_1$ . By Proposition 4.1 the space  $X'_1 \sqcup_\varphi \Phi_1$  is homotopy equivalent to

$$X_1 \sqcup_{f_1} \Phi_1 = X_1^+.$$

Therefore,  $H_\alpha(\Gamma_1, \mathbb{Z}) = H_\alpha(X'_1 \sqcup_\varphi \Phi_1)$ . The map  $f_1$  being of lift of  $f$ , its norm satisfies  $\|f_1\| \leq \|f\| = O(1)$ . The norm of the boundary map on  $X'_1 \sqcup_\varphi \Phi_1$  in degree  $\alpha + 1$  is bounded by  $\|\varphi\| \leq \|f_1\| \|g\| = O(T + 1)$ . By Proposition 9.1 we have

$$\log |H_\alpha(X'_1 \sqcup_\varphi \Phi_1, \mathbb{Z})_{\text{tors}}| \leq \|\partial_{\alpha+1}\|_{(X'_1 \sqcup_\varphi \Phi_1)} \times \#^{(\alpha)}(X'_1 \sqcup_\varphi \Phi_1) \leq O(\log(T + 1)[\Gamma : \Gamma_1]T^{-1}).$$

Since any Farber sequence eventually falls into  $U$ , we get the theorem by letting  $T \rightarrow \infty$ .  $\square$

We end this section with a question.

**Question 10.22.** *Does every residually finite amenable group of type  $F_\alpha$  have the cheap  $\alpha$ -rebuilding property?*

In this context, Kar, Kropholler and Nikolov [KKN17] prove the vanishing of the  $j$ -th torsion growth for  $j \leq \alpha$ .

## 11. LATTICES IN SEMI-SIMPLE LIE GROUPS (PROOF OF THEOREM B)

Let  $\mathbf{G}$  be an affine algebraic group defined over  $\mathbb{Q}$ . The **radical** of  $\mathbf{G}$  is the greatest connected normal solvable subgroup. The group  $\mathbf{G}$  is called **semi-simple** if its radical is trivial. In the following, we assume that  $\mathbf{G}$  is connected and semi-simple.

An algebraic group  $\mathbf{T}$  over  $\mathbb{Q}$  is an **algebraic torus** if the group of its complex points  $\mathbf{T}(\mathbb{C})$  is isomorphic to a product of  $\mathrm{GL}_1(\mathbb{C})$ . If  $\mathbf{T}$  is isomorphic to a product of  $\mathrm{GL}_1$  themselves over  $\mathbb{Q}$ , then  $\mathbf{T}$  is said to be **split** over  $\mathbb{Q}$ . The maximal dimension of a torus in  $\mathbf{G}$  that is split over  $\mathbb{Q}$  is called the **rational rank** of  $\mathbf{G}$ .

A Zariski-closed subgroup  $\mathbf{P}$  of  $\mathbf{G}$  is called **parabolic** if  $\mathbf{P}$  contains a connected solvable subgroup, i.e., a **Borel subgroup** of  $\mathbf{G}$ . If  $\mathbf{P}$  is defined over  $\mathbb{Q}$ , then  $\mathbf{P}$  is called a rational parabolic subgroup. The **unipotent radical** of  $\mathbf{P}$  is defined to be its largest normal subgroup consisting entirely of unipotent elements.

A  $\mathbb{Z}$ -structure of  $\mathbf{G}$  is given by a faithful  $\mathbb{Q}$ -embedding of  $\mathbf{G}$  in some  $\mathrm{GL}_n$ . We define  $\Gamma = \mathbf{G}_{\mathbb{Z}}$  to be the intersection of  $\mathbf{G}$  with  $\mathrm{GL}_n(\mathbb{Z})$  and the level  $N$  **principal congruence subgroup**  $\Gamma(N)$  to be the intersection of  $\mathbf{G}$  with the kernel of  $\mathrm{GL}_n(\mathbb{Z}) \rightarrow \mathrm{GL}_n(\mathbb{Z}/N\mathbb{Z})$ . Here  $N$  is some positive integer.

**Theorem 11.1.** *Let  $\mathbf{G}$  be a connected semi-simple affine algebraic group defined over  $\mathbb{Q}$  equipped with a  $\mathbb{Z}$ -structure. Let  $r$  be the rational rank of  $\mathbf{G}$ . Then there exists a constant  $\kappa = \kappa(\mathbf{G})$  such that for every **principal congruence subgroup**  $\Gamma(N) \leq \mathbf{G}_{\mathbb{Z}}$ , every  $j \leq r - 1$  and every coefficient field  $K$ , we have*

$$\dim_K |H_j(\Gamma(N), K)| = \kappa N^{(1-\delta)\dim \mathbf{G}} \quad \text{and} \quad \log |H_j(\Gamma(N), \mathbb{Z})_{\mathrm{tors}}| \leq \kappa N^{(1-\delta)\dim \mathbf{G}} \log N, \quad (11.1)$$

where

$$\delta = \min \left\{ \frac{\dim \mathbf{U}}{\dim \mathbf{G}} : \mathbf{U} \text{ is the unipotent radical of a parabolic subgroup of } \mathbf{G} \right\}.$$

*Remark 11.2.* In case  $\mathbf{G} = \mathrm{SL}_d$  we have  $r = d - 1$  and  $\dim \mathbf{G} = d^2 - 1$ . Any parabolic subgroup of  $\mathbf{G}$  is conjugated to a subgroup of block upper triangular matrices, and the unipotent radical of such is the subgroup where the block diagonal elements are the identity. It follows that

$$\delta = \frac{d-1}{d^2-1} = \frac{1}{d+1}.$$

Since there exists a universal constant  $C$  such that

$$1 > \frac{[\mathrm{SL}_d(\mathbb{Z}) : \Gamma(N)]}{N^{d^2-1}} > C > 0,$$

Theorem 11.1 implies Theorem B, as well as the congruence case of Theorem C, of the introduction.

*Proof of Theorem 11.1.* We first recall the spherical Tits building  $\Delta_{\mathbb{Q}}(\mathbf{G})$  associated with  $\mathbf{G}$  over  $\mathbb{Q}$  (see [Tit75, Tit74]); it is a simplicial set whose non-degenerate simplices are in bijection with proper rational parabolic subgroups of  $\mathbf{G}$ . Each proper maximal rational parabolic subgroup corresponds to a vertex of  $\Delta_{\mathbb{Q}}(\mathbf{G})$  and  $k+1$  proper parabolic subgroups  $\mathbf{P}_0, \dots, \mathbf{P}_k$  are the vertices of a  $k$ -simplex if and only if their intersection  $\mathbf{P}_0 \cap \dots \cap \mathbf{P}_k$  is a rational parabolic subgroup, and this simplex corresponds to the parabolic subgroup  $\mathbf{P}_0 \cap \dots \cap \mathbf{P}_k$ .

If the rational rank  $r$  of  $\mathbf{G}$  is equal to 1, then  $\Delta_{\mathbb{Q}}(\mathbf{G})$  is a countable collection of points. Otherwise  $\Delta_{\mathbb{Q}}(\mathbf{G})$  is a spherical building (see [Tit74, Theorem 5.2]). For any maximal  $\mathbb{Q}$ -split torus  $\mathbf{T}$ , all the

rational parabolic subgroups containing  $\mathbf{T}$  form an apartment in this building and each of these sub-complexes give a simplicial triangulation of the sphere of dimension  $\alpha = r - 1$  so that  $\Delta_{\mathbb{Q}}(\mathbf{G})$  has the homotopy type of a bouquet of  $\alpha$ -spheres (Solomon–Tits theorem [Sol69]); in particular it is  $(\alpha - 1)$ -connected. We refer to [Bro98, §V.5] for this and more about spherical buildings.

The rational points  $\mathbf{G}(\mathbb{Q})$  of  $\mathbf{G}$  act on the set of rational parabolic subgroups by conjugation and hence on  $\Delta_{\mathbb{Q}}(\mathbf{G})$ : for any  $g \in \mathbf{G}(\mathbb{Q})$  and any rational parabolic  $\mathbf{P}$ , the simplex associated to  $\mathbf{P}$  is mapped to the simplex associated to  $g\mathbf{P}g^{-1}$ . Let  $\Gamma \subset \mathbf{G}(\mathbb{Q})$  be an arithmetic subgroup of  $\mathbf{G}(\mathbb{Q})$ . Then by reduction theory (see e.g. [PR94, Theorem 4.15]) there are only finitely many  $\Gamma$ -conjugacy classes of rational parabolic subgroups. Therefore, the quotient  $\Gamma \backslash \Delta_{\mathbb{Q}}(\mathbf{G})$  is a finite simplicial complex.

From now on we let  $\Gamma = \mathbf{G}_{\mathbb{Z}}$ , we fix some principal congruence subgroup  $\Gamma(N)$  and let  $\Omega$  denote the rational Tits building  $\Delta_{\mathbb{Q}}(\mathbf{G})$ . Up to replacing  $\Omega$  by its barycentric subdivision we may furthermore assume that for every cell  $\sigma \subset \Omega$  the stabilizer  $\Gamma_{\sigma}$  acts trivially on  $\sigma$ .

The stabilizer  $\Gamma_{\sigma}$  of a simplex  $\sigma \subset \Omega$  associated to a rational parabolic subgroup  $\mathbf{P}$  contains the intersection  $\mathbf{P}_{\mathbb{Z}}$  of  $\mathbf{P}$  with  $\mathrm{GL}_d(\mathbb{Z})$ . In particular  $\Gamma_{\sigma}$  contains the  $\mathbb{Z}$ -points  $\mathbf{U}(\mathbb{Z})$  of the unipotent radical  $\mathbf{U}$  of  $\mathbf{P}$  as a normal subgroup. The group  $\mathbf{U}(\mathbb{Z})$  is finitely generated, torsion-free, nilpotent and its intersection with the stabilizer  $\Gamma(N)_{\sigma}$  is equal to

$$\ker(\mathbf{U}(\mathbb{Z}) \rightarrow \mathbf{U}(\mathbb{Z}/N\mathbb{Z}))$$

which is of index at least  $N^{\delta \dim \mathbf{G}}$  in  $\mathbf{U}(\mathbb{Z})$ . On the other hand it is well known that

$$[\Gamma : \Gamma(N)] \leq O(N^{\dim \mathbf{G}}),$$

see e.g. [BL12, bottom of page 3134]. It therefore follows from Theorem 8.1 (with  $\Gamma_1 = \Gamma(N)$ ) that there exists a constant  $c = c(\mathbf{G})$  that depends on  $\mathbf{G}$  but not on  $\Gamma(N)$  and a finite CW-complex  $\Sigma_N^+$  such that the following properties hold.

- (1) The CW-complex  $\Sigma_N^+$  has a  $\alpha$ -connected universal cover.
- (2) In each dimension  $\leq \alpha$ , the total number of cells of  $\Sigma_N^+$  is bounded by

$$cN^{(1-\delta) \dim \mathbf{G}}. \tag{11.2}$$

- (3) In each degree  $\leq \alpha + 1$  the norm of the boundary operator on the chain complex  $C_{\bullet}(\Sigma_N^+)$  is bounded by

$$cN^c. \tag{11.3}$$

We conclude by applying (9.1) and Proposition 9.1. Let  $\kappa = c^2 \log c$ . Then, in each degree  $j \leq \alpha$  we have both

$$\dim_K |H_j(\Gamma(N), K)| \leq \kappa N^{(1-\delta) \dim \mathbf{G}} \quad \text{and} \quad \log |H_j(\Gamma(N))_{\mathrm{tors}}| \leq \kappa N^{(1-\delta) \dim \mathbf{G}} \log N.$$

□

We used the fact that the stabilizers of cells in the action of  $\Gamma$  on  $\Omega$  have infinite normal unipotent subgroups. Hence, by Lemma 10.14 they all have the cheap  $\alpha$ -rebuilding property, for every  $\alpha$ . Upon applying Theorem 10.10 to the action of  $\Gamma$  on the rational Tits building  $\Omega$  we get

**Theorem 11.3.** *Let  $\mathbf{G}$  be a connected semi-simple affine algebraic group defined over  $\mathbb{Q}$  equipped with a  $\mathbb{Z}$ -structure. Let  $r$  be the rational rank of  $\mathbf{G}$ . Then  $\mathbf{G}_{\mathbb{Z}}$  has the cheap  $(r - 1)$ -rebuilding property.*

Theorem 10.21 now yields.

**Theorem 11.4.** *Let  $\mathbf{G}$  be a connected semi-simple affine algebraic group defined over  $\mathbb{Q}$  equipped with a  $\mathbb{Z}$ -structure. Let  $r$  be the rational rank of  $\mathbf{G}$ . Then for any Farber sequence  $(\Gamma_n)_{n \in \mathbb{N}}$  in  $\mathbf{G}_{\mathbb{Z}}$  and for any coefficient field  $K$  we have*

$$\lim_{n \rightarrow \infty} \frac{\dim_K H_j(\Gamma_n, K)}{[\Gamma : \Gamma_n]} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\log |H_j(\Gamma_n, \mathbb{Z})_{\mathrm{tors}}|}{[\mathbf{G}_{\mathbb{Z}} : \Gamma_n]} = 0,$$

for  $j = 0, \dots, r - 1$ .

This implies Theorem B and the first part of Theorem C of the introduction.

## 12. APPLICATION TO MAPPING CLASS GROUPS (PROOF OF THEOREM D)

Let  $S$  be a closed orientable surface of genus  $g$  with  $b$  connected boundary components. We assume that

$$\chi(S) = 2 - 2g - b < 0 \quad \text{and} \quad b \geq 4 \text{ if } g = 0.$$

We write  $\mathcal{MCG}(S)$  for the mapping class group of  $S$ . Let us recall the construction of the curve complex  $\mathcal{C}(S)$ .

The curve complex  $\mathcal{C}(S)$  is a combinatorial cell complex whose  $k$ -cells consist of collections of  $k + 1$  closed, disjoint, essential, pairwise non-homotopic simple closed curves on  $S$ , considered up to homotopy. In particular, a 0-dimensional cell corresponds to the homotopy class of a simple closed curve and top dimensional cells correspond to maximal families of pairwise non-homotopic essential closed curves in  $S$ . It is equipped with a co-compact  $\mathcal{MCG}(S)$ -action.

Under our hypotheses  $S$  admits a hyperbolic structure and the size of any maximal family of pairwise non-homotopic essential closed curves in  $S$  is finite, equal to  $3g - 3 + b$ . So the dimension of  $\mathcal{C}(S)$  is  $3g - 4 + b$ .

Now let  $\mathcal{F}$  be a finite collection of  $m$  closed, disjoint, essential, pairwise non-homotopic simple closed curves on  $S$ . The stabilizer of the homotopy class of  $\mathcal{F}$  in  $\mathcal{MCG}(S)$ , denoted  $\text{Stab } \mathcal{F}$ , fits into a short exact sequence:

$$1 \longrightarrow \mathbb{Z}^m \longrightarrow \text{Stab } \mathcal{F} \longrightarrow \mathcal{MCG}'(S \setminus \mathcal{F}) \longrightarrow 1,$$

where  $\mathcal{MCG}'(S \setminus \mathcal{F})$  is the subgroup of those mapping classes in  $\mathcal{MCG}(S \setminus \mathcal{F})$  that permute the boundary in such a way that it can still be glued back to obtain  $\mathcal{F}$  in  $S$ ; see e.g. [Min06, Equation (1)]. The group  $\mathcal{MCG}'(S \setminus \mathcal{F})$  has finite index in  $\mathcal{MCG}(S \setminus \mathcal{F})$ , which is of class  $F_k$  for every  $k \geq 0$  by [Iva01, Theorem 5.4.A]. Hence the stabilizers  $\text{Stab } \mathcal{F}$  are of class  $F_k$  for every  $k \geq 0$ .

In order to ensure the assumption (Cond 2) that the cell-stabilizers act freely on them, we consider instead the action of  $\Gamma = \mathcal{MCG}(S)$  on the barycentric subdivision  $\Omega$  of  $\mathcal{C}(S)$ . Now the stabilizer of any cell  $\sigma$  of  $\Omega$  is a finite index subgroup of some  $\text{Stab } \mathcal{F}$  as above. Therefore  $\Gamma_\sigma$  is of class  $F_\alpha$  for every  $\alpha \geq 0$  and it has a finite rank normal free abelian subgroup, so by Corollary 10.14 it has the cheap  $\alpha$ -rebuilding property for every  $\alpha$ .

The homotopy type of  $\mathcal{C}(S)$  (and therefore that of  $\Omega$ ) was identified by Harer [Har86, Thm 3.5]; it is homotopy equivalent to a bouquet of spheres of dimension  $\alpha(g, b)$  where

$$\alpha(g, b) := \begin{cases} 2g - 2 & \text{if } g \geq 2 \text{ and } b = 0; \\ 2g - 3 + b & \text{if } g \geq 1 \text{ and } b \geq 1; \\ b - 4 & \text{if } g = 0 \text{ and } b \geq 4. \end{cases}$$

Thus the space  $\Omega$  is  $(\alpha(g, b) - 1)$ -connected. By Theorem 10.10, we obtain:

**Theorem 12.1.** *The mapping class group  $\mathcal{MCG}(S)$  has the cheap  $\alpha(g, b)$ -rebuilding property.*

By [Iva01, Theorem 5.4.A],  $\mathcal{MCG}(S)$  is of class  $F_\alpha$  for every  $\alpha \geq 0$ . Applying Proposition 10.21 finishes the proof of Theorem D.  $\square$

Curiously we had a hard time trying to apply our method to  $\text{Out}(\mathbf{F}_n)$ , thus our question:

**Question 12.2.** *What is the range of  $\alpha$  for which  $\text{Out}(\mathbf{F}_n)$  has the cheap  $\alpha$ -rebuilding property?*



## REFERENCES

- [ABB<sup>+</sup>17] Miklos Abert, Nicolas Bergeron, Ian Biring, Tsachik Gelander, Nikolay Nikolov, Jean Raimbault, and Iddo Samet. On the growth of  $L^2$ -invariants for sequences of lattices in Lie groups. *Ann. Math. (2)*, 185(3):711–790, 2017. [4](#)
- [AGMY20] Avner Ash, Paul E. Gunnells, Mark McConnell, and Dan Yasaki. On the growth of torsion in the cohomology of arithmetic groups. *J. Inst. Math. Jussieu*, 19(2):537–569, 2020. [2](#), [3](#)
- [AGN17] Miklos Abert, Tsachik Gelander, and Nikolay Nikolov. Rank, combinatorial cost, and homology torsion growth in higher rank lattices. *Duke Math. J.*, 166(15):2925–2964, 2017. [3](#), [7](#), [41](#)
- [AOS20] Grigori Avramidi, Boris Okun, and Kevin Schreve. Mod  $p$  and torsion homology growth in nonpositive curvature, 2020. [7](#), [8](#)
- [Ati76] Michael F. Atiyah. Elliptic operators, discrete groups and von Neumann algebras. In *Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974)*, pages 43–72. Astérisque, No. 32–33. 1976. [1](#)
- [BD15] Jeffrey F. Brock and Nathan M. Dunfield. Injectivity radii of hyperbolic integer homology 3-spheres. *Geom. Topol.*, 19(1):497–523, 2015. [2](#), [8](#)
- [Ber18] Nicolas Bergeron. Torsion homology growth in arithmetic groups. In *European Congress of Mathematics*, pages 263–287. Eur. Math. Soc., Zürich, 2018. [2](#), [8](#)
- [Bes99] Mladen Bestvina. Non-positively curved aspects of Artin groups of finite type. *Geom. Topol.*, 3:269–302, 1999. [43](#)
- [BGS20] Uri Bader, Tsachik Gelander, and Roman Sauer. Homology and homotopy complexity in negative curvature. *J. Eur. Math. Soc. (JEMS)*, 22(8):2537–2571, 2020. [3](#)
- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999. [42](#)
- [BK17] Martin R. Bridson and Dessislava H. Kochloukova. Volume gradients and homology in towers of residually-free groups. *Math. Ann.*, 367(3-4):1007–1045, 2017. [3](#), [7](#)
- [BL12] Mikhail Belolipetsky and Alexander Lubotzky. Manifolds counting and class field towers. *Adv. Math.*, 229(6):3123–3146, 2012. [46](#)
- [BLLS14] Nicolas Bergeron, Peter Linnell, Wolfgang Lück, and Roman Sauer. On the growth of Betti numbers in  $p$ -adic analytic towers. *Groups Geom. Dyn.*, 8(2):311–329, 2014. [3](#)
- [Bou81] Nicolas Bourbaki. *Groupes et algèbres de Lie*. Chapitres 4, 5 et 6. Éléments de Mathématique. Paris etc.: Masson. 288 p. (1981)., 1981. [42](#)
- [Bro98] Kenneth S. Brown. *Buildings*. Springer Monographs in Mathematics. Springer-Verlag, New York, 1998. Reprint of the 1989 original. [46](#)
- [BS01] Itai Benjamini and Oded Schramm. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.*, 6:no. 23, 13, 2001. [34](#)
- [BŞV16] Nicolas Bergeron, Mehmet Haluk Şengün, and Akshay Venkatesh. Torsion homology growth and cycle complexity of arithmetic manifolds. *Duke Math. J.*, 165(9):1629–1693, 2016. [2](#), [8](#)
- [BV13] Nicolas Bergeron and Akshay Venkatesh. The asymptotic growth of torsion homology for arithmetic groups. *J. Inst. Math. Jussieu*, 12(2):391–447, 2013. [2](#), [3](#), [5](#), [33](#), [34](#)
- [Cal15] Frank Calegari. The stable homology of congruence subgroups. *Geom. Topol.*, 19(6):3149–3191, 2015. [3](#)
- [CD95] Ruth Charney and Michael W. Davis. The  $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups. *J. Am. Math. Soc.*, 8(3):597–627, 1995. [42](#), [43](#)
- [CE09] Frank Calegari and Matthew Emerton. Bounds for multiplicities of unitary representations of cohomological type in spaces of cusp forms. *Ann. of Math. (2)*, 170(3):1437–1446, 2009. [3](#)
- [CE11] Frank Calegari and Matthew Emerton. Mod- $p$  cohomology growth in  $p$ -adic analytic towers of 3-manifolds. *Groups Geom. Dyn.*, 5(2):355–366, 2011. [3](#)
- [CE16] Frank Calegari and Matthew Emerton. Homological stability for completed homology. *Math. Ann.*, 364(3-4):1025–1041, 2016. [3](#)
- [CGdLS19] Alessandro Carderi, Damien Gaboriau, and Mikael de la Salle. Non-standard limits of graphs and some orbit equivalence invariants, 2019. [3](#)
- [DD89] Warren Dicks and Martin J. Dunwoody. *Groups acting on graphs*. Cambridge University Press, Cambridge, 1989. [31](#)
- [Eme14] Vincent Emery. Torsion homology of arithmetic lattices and  $K_2$  of imaginary fields. *Math. Z.*, 277(3-4):1155–1164, 2014. [2](#)
- [Fra18] Mikolaj Fraczyk. Growth of mod  $-2$  homology in higher rank locally symmetric spaces. *arXiv preprint arXiv:1801.09283*, 2018. [3](#)
- [Gab00] Damien Gaboriau. Coût des relations d’équivalence et des groupes. *Invent. Math.*, 139(1):41–98, 2000. [3](#), [7](#)

- [Geo08] Ross Geoghegan. *Topological methods in group theory*, volume 243 of *Graduate Texts in Mathematics*. Springer, New York, 2008. [4](#), [5](#), [8](#), [9](#), [14](#), [17](#), [18](#), [25](#), [33](#), [43](#), [44](#)
- [Har86] John L. Harer. The virtual cohomological dimension of the mapping class group of an orientable surface. *Invent. Math.*, 84(1):157–176, 1986. [47](#)
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. [8](#), [23](#)
- [Iva01] Nikolai V. Ivanov. Chapter 12 - mapping class groups. In R.J. Daverman and R.B. Sher, editors, *Handbook of Geometric Topology*, pages 523 – 633. North-Holland, Amsterdam, 2001. [47](#)
- [KKN17] Aditi Kar, Peter Kropholler, and Nikolay Nikolov. On growth of homology torsion in amenable groups. *Math. Proc. Cambridge Philos. Soc.*, 162(2):337–351, 2017. [3](#), [45](#)
- [KN16] Aditi Kar and Nikolay Nikolov. On deficiency gradient of groups. *Int. Math. Res. Not. IMRN*, (3):696–716, 2016. [3](#), [7](#)
- [Lüc94] Wolfgang Lück. Approximating  $L^2$ -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.*, 4(4):455–481, 1994. [1](#), [7](#)
- [Mal49] Aaanatly I. Malcev. On a class of homogeneous spaces. *Izvestiya Akad. Nauk. SSSR. Ser. Mat.*, 13:9–32, 1949. [24](#)
- [Min06] Yair Minsky. Curve complexes, surfaces and 3-manifolds. In *International Congress of Mathematicians*, volume 2, pages 1001–1033. 2006. [47](#)
- [PR94] Vladimir Platonov and Andrei Rapinchuk. *Algebraic groups and number theory*, volume 139 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1994. Translated from the 1991 Russian original by Rachel Rowen. [46](#)
- [PS21] Giovanni Paolini and Mario Salvetti. Proof of the  $K(\pi, 1)$  conjecture for affine Artin groups. *Invent. Math.*, 224(2):487–572, 2021. [43](#)
- [Sau16] Roman Sauer. Volume and homology growth of aspherical manifolds. *Geom. Topol.*, 20(2):1035–1059, 2016. [3](#)
- [Sch15] Peter Scholze. On torsion in the cohomology of locally symmetric varieties. *Ann. Math. (2)*, 182(3):945–1066, 2015. [2](#)
- [Sol69] Louis Solomon. The Steinberg character of a finite group with  $BN$ -pair. pages 213–221, 1969. [46](#)
- [Sou99] Christophe Soulé. Perfect forms and the Vandiver conjecture. *J. Reine Angew. Math.*, 517:209–221, 1999. [2](#), [5](#), [33](#)
- [SZ94] Garrett Stuck and Robert J. Zimmer. Stabilizers for ergodic actions of higher rank semisimple groups. *Ann. Math. (2)*, 139(3):723–747, 1994. [4](#)
- [Tit74] Jacques Tits. *Buildings of spherical type and finite  $BN$ -pairs*. Lecture Notes in Mathematics, Vol. 386. Springer-Verlag, Berlin-New York, 1974. [45](#)
- [Tit75] Jacques Tits. On buildings and their applications. In *Proceedings of the International Congress of Mathematicians (Vancouver, B. C., 1974)*, Vol. 1, pages 209–220, 1975. [45](#)

M. A.: RENYI INSTITUTE OF MATHEMATICS, REALTANODA UTCA 13-15, 1053 BUDAPEST, HUNGARY  
*Email address:* [abert@renyi.hu](mailto:abert@renyi.hu)

N. B.: ENS / PSL UNIVERSITY, DÉPARTEMENT DE MATHÉMATIQUES ET APPLICATIONS, F-75005, PARIS, FRANCE  
*Email address:* [nicolas.bergeron@ens.fr](mailto:nicolas.bergeron@ens.fr)  
*URL:* <https://sites.google.com/view/nicolasbergeron/accueil>

M. F.: UNIVERSITY OF CHICAGO, DEPARTMENT OF MATHEMATICS, 5734 S UNIVERSITY AVE, CHICAGO, IL 60637, USA  
*Email address:* [mfraczyk@math.uchicago.edu](mailto:mfraczyk@math.uchicago.edu)

D. G.: ENS-LYON, UMR CNRS 5669 / UNIVERSITÉ DE LYON, UMPA, 69007 LYON, FRANCE  
*Email address:* [damien.gaboriau@ens-lyon.fr](mailto:damien.gaboriau@ens-lyon.fr)  
*URL:* <https://perso.ens-lyon.fr/gaboriau/>