



**HAL**  
open science

## Prevalence of hematozoan parasites in Red Kite nestlings from France

Michael Coeurdassier, Nicolas Bassin, Thibaut Powolny, Christophe Morin,  
Eve Afonso

► **To cite this version:**

Michael Coeurdassier, Nicolas Bassin, Thibaut Powolny, Christophe Morin, Eve Afonso. Prevalence of hematozoan parasites in Red Kite nestlings from France. *Wilson journal of ornithology*, 2021, 162 (2), pp.521-527. 10.1007/s10336-020-01838-w . hal-03278548

**HAL Id: hal-03278548**

**<https://hal.science/hal-03278548v1>**

Submitted on 5 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Michael Coeurdassier<sup>1</sup> . Nicolas Bassin<sup>1</sup> . Thibaut Powolny<sup>1</sup> . Christophe Morin<sup>2,3</sup> .  
2 Eve Afonso<sup>1</sup>

3

4 **Prevalence of hematozoan parasites in Red Kite nestlings**  
5 **from France**

6

7 <sup>1</sup> Department Chrono-environment – University of Franche-Comté, UMR/CNRS  
8 6249, 16 route de Gray, 25030 Besançon, France

9 <sup>2</sup> LPO Franche-Comté, 7 rue Voirin, 25000 Besançon, France

10 <sup>3</sup> 70120 Confracourt, France

11

12 Author for correspondence: Michael Coeurdassier [fcomte.fr](mailto:michael.coeurdassier@univ-<br/>13 fcomte.fr)

14

15 ORCID identifiers: Michael COEURDASSIER: [https://orcid.org/0000-0002-6260-  
17 311X](https://orcid.org/0000-0002-6260-<br/>16 311X) / Nicolas BASSIN: no ORCID Id / Thibaut POWOLNY: [https://orcid.org/0000-  
19 0001-6580-9853](https://orcid.org/0000-<br/>18 0001-6580-9853) / Christophe MORIN: no ORCID Id / Eve AFONSO:  
<https://orcid.org/0000-0001-5836-9909>

19

**Abstract**

21 Blood pathogens may be harmful for birds and thus, represent a potential threat for  
22 some populations. To date, only few data are available on the prevalence of  
23 hematozoans in the Red Kite, a raptor endemic to the Western Palearctic listed as  
24 Near Threatened within its global range. Here, we report the first screening of  
25 haemosporidian parasites in Red Kite nestlings from French breeding areas. The  
26 blood of 47 nestlings was sampled in three areas in Eastern France and the  
27 prevalence of three genera of blood pathogens, *Leucocytozoon*, *Haemoproteus* and  
28 *Plasmodium*, was measured based on PCR-assays. Thirteen nestlings were positive  
29 to *Leucocytozoon sp.* (n=13) and *Plasmodium relictum* (n=1), which corresponds to  
30 an overall prevalence of 28%. This prevalence is higher than those previously  
31 reported for the Red Kite in other countries. This is the first record of *P. relictum*, a  
32 potential pathogen for birds, in the Red Kite. No relationship was found between the  
33 body condition of nestlings and the occurrence of haemosporidians. Our findings  
34 suggest that hematozoans could be a limited threat for nestlings but prevalence in  
35 adults, a sensitive stage for the population dynamic of Red Kite, remains poorly  
36 known and should be further investigated at least in its main breeding areas.

37

38 Key words: Haemosporida, *Haemoproteus*, *Leucocytozoon*, *Plasmodium*, raptor

39 **Declarations**

40 Funding: This study was financially supported by the French Ministère de l'Écologie  
41 et du Développement Durable (Plan National d'Actions pour la conservation du  
42 Milan Royal), by the Région Franche-Comté (BIOSCOPE project, convention 2016-  
43 0059) and by the Long Term Ecological Research (LTER) site Zone Atelier Arc  
44 Jurassien.

45 Conflicts of interest/Competing interests: The authors declare that they have no  
46 interest or competing conflicts

47 Ethical approval: All the manipulations were performed with authorizations no.  
48 2015093-0007 and 2015093-0023 in the framework of the Red Kite national  
49 conservation plan.

50 All the authors consent to participate and for publication

51 Availability of data and material: The data presented in this article are referenced in  
52 the metadata platform dat@OSU proposed by the *Observatoire des Sciences de*  
53 *l'Univers Terre Homme Environnement Temps Astronomie* (OSU THETA)

54 Code availability: NA

55 Authors' contributions: Michael Coeurdassier designed the study. Christophe Morin  
56 and Michael Coeurdassier managed and carried out fieldwork and sampling. Eve  
57 Afonso and Nicolas Bassin conducted the pathogen analysis. Michael Coeurdassier,  
58 Eve Afonso, Thibaut Powolny and Nicolas Bassin analyzed and interpreted the data  
59 and wrote the manuscript. All co-authors commented on the manuscript and  
60 accepted the final version.

61 **Aknowledgements**

62 We are grateful to Damien Bertin, Violaine Champion, Natacha Leclerc, Geneviève  
63 Magnon, Dominique Michelat, Jacques Morel, François Rey-Demaneuf, François  
64 Richard, Michel Sauret, Bruno Tissot and all the volunteers for ensuring the  
65 monitoring of the studied Red Kite populations. We warmly thank the three  
66 anonymous referees for their contribution to improve this article. This study was  
67 financially supported by the French Ministère de l'Ecologie et du Développement  
68 Durable (Plan National d'Actions pour la conservation du Milan Royal), by the  
69 Région Franche-Comté (BIOSCOPE project, convention 2016-0059) and by the  
70 LTER site *Zone Atelier Arc Jurassien*. The molecular analyses were carried out on  
71 the PEA<sup>2t</sup> platform. The experiments comply with the current laws of France when  
72 they were performed.

73

## 74 Introduction

75 The Red Kite *Milvus milvus* is an Accipitridae endemic to the Western Palearctic. It  
76 breeds mainly in Germany, France and Spain, which together hold around 75% of  
77 the global population (Knott et al. 2010). In these countries, the populations have  
78 been estimated to be declining, which led the International Union for the  
79 Conservation of Nature to assess the Red Kite as Near threatened (BirdLife  
80 International 2020) within its global range while it is currently listed as vulnerable in  
81 France (UICN France et al. 2016). The main threats identified are poisoning by  
82 anticholinesterasic pesticides and rodenticides, decrease of livestock,  
83 homogenization of landscapes and potentially wind turbines (BirdLife International  
84 2020). In both Europe and France, the decline reported during the period 1990-2010  
85 has led to the implementation of a Conservation Plan in which the health status of  
86 the Red Kite was surveyed, notably on nestlings that were caught at nests for  
87 ringing and wing-marking. Among the health indicators monitored and considering  
88 that ecotoxicological issues were identified, chemical concentrations in blood were  
89 measured and we showed that 50% of nestlings were exposed to organic  
90 compounds and/or metals (Coourdassier et al. 2014; Powolny et al. 2020). If  
91 infection by pathogens has not been pointed out as a threat for the Red Kite, only  
92 few studies have focused on the prevalence of blood parasites and no data are  
93 available on the prevalence of hematozoan pathogens in Red Kite nestlings from  
94 France. Haemosporidians belonging to three genera of protozoan (*Haemoproteus*,  
95 *Leucocytozoon*, and *Plasmodium*) are intracellular blood parasites transmitted to  
96 hosts by bloodsucking dipteran vectors, they have been commonly detected in birds  
97 including raptors (Remple 2004). *Haemoproteus* and *Leucocytozoon* are often  
98 considered to have little or no pathogenic effects on birds but their presence in some  
99 raptor nestlings has been related to subclinical effects and even host death.  
100 *Plasmodium*, the causative agent of malaria in animals, can be a serious avian

101 pathogen, *Plasmodium relictum* being the species most commonly associated with  
102 virulent disease (Remple 2004). In raptors, haemosporidians may also negatively  
103 affect health such as body condition and life traits (Korpimaki et al. 1993; Hatchwell  
104 et al. 2001; Sol et al. 2003; Garvin et al. 2006). Here, we aim to measure the  
105 prevalence of common blood parasites in birds of the genera *Haemoproteus*,  
106 *Leucocytozoon* and *Plasmodium* in nestlings of three areas monitored in Eastern  
107 France. The relationship between prevalence and body condition of the nestlings  
108 was also checked to document whether such infection could affect individuals.

## 109 **Methods**

### 110 **Study areas and sample collection**

111 Fieldwork was carried out in 2015 from 4<sup>th</sup> to 25<sup>th</sup> June in Franche-Comté region,  
112 Eastern France. Three breeding populations of Red Kite, from three areas (called  
113 “Area” factor hereafter) were monitored: Sundgau Belfortain (47°380170’N,  
114 6°510460’E), Dugeon-Remoray basin (46°540220’N, 6°210170’E) and Besançon  
115 plateau (47°140350’N, 6°010190’E). Forty-seven chicks (23 males and 24 females)  
116 corresponding to 24 nests (called “Nest” factor hereafter) were retained for the  
117 study. The sampling was implemented 15 days prior to the expected fledging date,  
118 when the chicks were approximately 25–40 days old. Nestlings were removed from  
119 the nests by an experienced arborist climber and carried to the ground for data  
120 collection. First, the body mass (g) and the wing length (i.e., length from the  
121 extremity of the longest primary feather to the wrist of the right wing in mm) were  
122 measured. Blood (2 mL) was sampled by trained operators from the brachial vein  
123 using lithium-heparin coated syringes. As recommended (Fair et al. 2010), the  
124 volume of blood collected was <0.5% of the chick’s body mass. Blood samples were  
125 kept in the dark at 4°C and then stored at -20°C until processed in the laboratory.  
126 The sex of the chicks was determined according to a PCR method (Griffiths et al.

127 1998) as described in Powolny et al. (2020). All the manipulations were performed  
128 with governmental authorizations no. 2015093-0007 and 2015093-0023 in the  
129 framework of the Red Kite national conservation plan.

### 130 **Body condition estimate**

131 In the present study, we define body condition as a proxy of energy reserves, mainly  
132 fat mass that represents a broad indicator of the health status of nestlings.

133 According to Labocha and Hayes (2012), body mass of birds is « *as good or nearly*  
134 *as good an indicator of fat content, as any of the other condition indices* ». Because  
135 body mass can be easily and reliably measured, we used it as an estimate of body  
136 condition. As we studied growing individuals, difference in body mass could be due  
137 to slight differences in specimen age rather than in fat reserve. Mougeot and  
138 Bretagnolle (2006) showed that the wing length was closely correlated to the age of  
139 Red Kite nestlings. Moreover, the relationship they established is almost linear  
140 within the range of ages of the kite chicks we studied, thus, we used the wing length  
141 as an estimate of chick age. Body condition was calculated as the residuals of the  
142 linear model [body mass (in g) ~ wing length (in mm)], which corresponds to an  
143 assessment of the body mass following a statistical elimination (sensus Grafen and  
144 Hails 2002) of age differences. Positive residuals were related to chicks in relative  
145 good condition while negative residuals represented kites in poor condition.

### 146 **DNA extraction, PCR amplification, sequencing, and parasite detection**

147 DNA was extracted from blood samples using the DNeasy Blood & Tissue kit  
148 (Qiagen, France) and according to manufacturer's instructions (protocol "Purification  
149 of Total DNA from Animal Blood or Cells"). A nested PCR was conducted to amplify  
150 a fragment of cytochrome b gene (542 bp) from species of *Plasmodium*,  
151 *Haemoproteus* and *Leucocytozoon* as recommended by Pérez-Rodríguez et al.  
152 (2013). A first PCR was performed in total volumes of 12.5µL including 1 × Accustart



153 II PCR ToughMix (Quanta Biosciences, USA), 0.6 mM of each primer  
154 (Plas1F/HaemNR3), and 22 ng of total genomic DNA. The PCR program consisted  
155 in an activation step of 3 min at 94°C, followed by 20 cycles of 30 sec at 94°C, 20  
156 sec at 52°C, 30 sec at 72°C, and a final elongation of 5 min at 72°C. We then used  
157 PCR products from the first PCR to conduct the following reaction in a total volume  
158 of 25µL comprising 1 × Accustart II PCR ToughMix, 0.6 mM of each primer  
159 (3760F/HaemJR4) and 2.5µL of PCR products from the first PCR. The PCR  
160 program consisted in an activation step of 3 minutes at 94°C, followed by 35 cycles  
161 of 30 sec at 94°C, 20 sec at 52°C, 30 sec at 70°C, and a final elongation of 5 min at  
162 70°C. PCR products were then separated and visualized using a Qiaxcel device  
163 (Qiagen, France) and a DNA High Resolution kit (Qiagen). A negative control (PCR  
164 grade water instead of DNA extract) was introduced every ten samples. PCR  
165 products which yielded an approximately 590 bp fragment (including primers) were  
166 confirmed by sequencing. The amplified products were purified using the QIAquick  
167 PCR Purification Kit (Qiagen) according to manufacturer's instructions. Direct  
168 sequencing of the purified PCR products was performed with an automated  
169 sequencer (Applied Biosystems 3130 Genetic Analyzer) using the primers employed  
170 for the second PCR reactions. For each PCR-positive sample, we aligned forward  
171 and reverse DNA sequences to obtain a consensus of 479 bp (Pérez-Rodríguez et  
172 al. 2013). A homology search of the sequences generated in this study was then  
173 performed by conducting an online search with the NCBI Basic Local Alignment  
174 Search Tool for nucleotides (BLASTN) in the GenBank database  
175 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

176 Mixed infections were checked by examining electropherograms using the software  
177 BioEdit following the recommendations of Pérez-Tris and Bensch (2005). When  
178 double peaks were observed, we compared double peak patterns with the DNA  
179 sequences of parasite haplotypes already available on Genbank to identify which

180 parasite haplotypes were involved in mixed infections (Pérez-Rodríguez et al.,  
181 2013).

## 182 **Data analysis**

183 Difference of infection status (i.e., presence/absence of hematozoans) between sex  
184 was checked with a binomial generalized linear mixed effect model (GLMER) with  
185 'Sex' as fixed factors and 'Nest' nested into 'Area' (Area/Nest) as random effects  
186 [corresponding full model: Status ~ Sex + (1|Area/Nest)]. The relationship between  
187 the presence of at least one positive chick in the brood ('Positive brood', binomial  
188 variable) and the brood size was tested with a binomial generalized linear mixed  
189 effect model (GLMER) with 'Brood size' as fixed factors and 'Area' as random  
190 effects [corresponding full model: Positive brood ~ Brood size + (1|Area)].

191 Relationship between body condition and infection status of the individual was  
192 checked with a linear mixed effect model (LMER) with 'Wing length', 'Sex' (in  
193 interaction or not with 'Wing length') and 'Status' as fixed factors and 'Nest' nested  
194 into 'Area' (Area/Nest) as random effects [corresponding full model: Body mass ~  
195 Wing length + Sex + Status + (1|Area/Nest)] (Zuur et al., 2009). The null model was  
196 also tested and the most parsimonious model with the lowest Akaike information  
197 criterion (AICc) was selected, a difference of AICc > 2 was retained to evidence  
198 difference between 2 models (Burnham and Anderson, 2010). Normality of model  
199 residuals and variance homoscedasticity were verified. All statistical analyses were  
200 performed using R 3.4.2. with the packages *lme4*, *pgirmess* and *lmerTest*.

201

## 202 **Results**

203 Among the 47 chicks studied, 30 came from the Dugeon-Remoray basin (15  
204 females and 15 males), 14 from the Besançon plateau (eight females and six males)  
205 and three from the Sundgau belfortain (one female and two males). Haemosporidian

206 parasites were found in 13 chicks (i.e., 28%), *Leucocytozoon* was detected in eight  
207 males and five females from the three studied areas and *Plasmodium* in a single  
208 male also infected by *Leucocytozoon* of the Sundgau belfortain population (Fig. 1).  
209 No difference of *Leucocytozoon* prevalence was evidenced between sex (binomial  
210 GLMER,  $p = 0.33$ ) or between brood size (binomial GLMER,  $p = 0.66$ ). We found  
211 two *Leucocytozoon* haplotypes, which perfectly matched (i.e. with a 100% identity  
212 on 479bp) with L-MILVUS1 (Genbank accession number JN164716) and L-  
213 MILANS4 (JN164713) haplotypes described in Pérez-Rodríguez et al. (2013). The  
214 DNA sequence corresponding to L-MILVUS1 also closely matched with  
215 *Leucocytozoon toddi* (GenBank accession number DQ177272) with a 99% identity.  
216 The DNA sequence amplified in the only sample positive for *Plasmodium* matched  
217 with *Plasmodium relictum* (GenBank accession number LN835311) with a 100%  
218 identity over 462bp. Although *Leucocytozoon toddi* is a typical parasite of birds  
219 (Valkiūnas et al. 2010), we cannot exclude that we detected an unreferenced  
220 species genetically close, we thus retained *Leucocytozoon* ssp. hereafter, while we  
221 considered that DNA from *Plasmodium relictum* was detected in the corresponding  
222 positive nestling. The only PCR-positive sample for *Plasmodium relictum* was also  
223 positive for *Leucocytozoon* sp. (L-MILVUS1 haplotype) and we recorded three other  
224 individuals with a co-occurrence of both *Leucocytozoon* haplotypes L-MILVUS1 and  
225 L-MILANS4, among which two nestlings from the same brood. This suggests that  
226 co-infections can occur relatively frequently, both for interspecific (1/47 individual)  
227 and inter-haplotypes (3/47) co-occurrences.

228 At a same wing length, females are heavier than males (best LMER selected with  
229 'Sex' as factor,  $p < 0.0001$ , Table 1. Estimate for males = -84.8,  $df = 31.1$ ,  $t = -5.45$ ).  
230 Thus, the body condition of the nestlings is presented for both sex separately in Fig.  
231 1. No difference of body condition was detected between infected and non-infected  
232 kites whatever the sex ( $p = 0.15$ , Table 1).

233

234 **Discussion**

235 Our findings showed that nestlings of Red Kite are infected by haemosporidians,  
236 notably *Leucocytozoon ssp.* and *P. relictum*, in three French breeding areas. The  
237 prevalence of 28% we measured was higher than those commonly reported for  
238 *Leucocytozoon* in Red Kite chicks from German, Spanish, UK or Danish populations  
239 while *Plasmodium* was less frequently found (Table 2). Overall, if we aggregate all  
240 the studies reported in Table 2 except our and without distinction of age, the  
241 prevalences measured in the Red Kite (n=366) were 3.8%, 0.3% and 0.0% for  
242 *Leucocytozoon*, *Haemoproteus* and *Plasmodium*, respectively. Co-infection by two  
243 haplotypes of *Leucocytozoon* corresponding to some of those previously described  
244 in Spanish populations of Red Kite (Pérez-Rodríguez et al. 2013) were found in  
245 three nestlings of the French populations studied. We also point out that the  
246 detection of *P. relictum* in one single nestling of a French population is the first  
247 record for the Red Kite while several haplotypes of *Plasmodium* were detected in  
248 the Black Kite *Milvus migrans* (Pérez-Rodríguez et al. 2013; Nourani et al. 2020).  
249 More generally, *P. relictum* has already been reported in species belonging to  
250 Accipitridae and Falconidae families (Bennett et al. 1993). It is considered as a  
251 serious pathogen for birds and has been associated to virulent diseases and even  
252 death for translocated Gyrfalcon *Falco rusticolus* (Remple 2004). However, to our  
253 knowledge, no information is available on its harmfulness for Accipitridae and the  
254 individual infected by *P. relictum* we found did not show obvious health troubles.

255 Prevalence of blood parasites in diurnal raptors is highly variable and may reach  
256 50% or more notably in adults (e.g., Forrester et al. 1994; Castellucci et al. 1998),  
257 the genera *Haemoproteus* and *Leucocytozoon* being the most frequently detected  
258 while *Plasmodium* commonly has lower prevalence (Remple 2004). Overall,

259 nestlings have been reported to exhibit relatively low prevalence of Haemosporida in  
260 numerous studies (e.g., Gutiérrez-López et al. 2015; Hanel et al. 2016) even if  
261 notable exceptions exist (Toyne and Ashford, 1997, Jeffries et al. 2015). Such  
262 differences of prevalences between juveniles and adults have been explained by the  
263 higher probability of adults contracting an infection in successive year (Podmokła et  
264 al. 2014). For the Red Kite, available data does not allow to identify a difference of  
265 prevalence between adults and nestlings mainly because of the low number of  
266 positive individuals and unbalanced samples (Table 2).

267 The comparison of Haemosporida prevalence of Red Kite in Spain to those  
268 determined in other studies (Table 2) suggests that it could be higher in more  
269 northern areas such as France and potentially in Denmark, UK and Western Europe  
270 despite low sample sizes. The prevalence of haemosporidians in diurnal raptors has  
271 been explained by different factors. Some are related to populations such as the  
272 abundance of Diptera vectors within the environment (Sol et al. 2003, Sehgal 2015)  
273 while others rely more on species features, such as the world geographic range,  
274 breeding habitat and embryonic development (Tella et al. 1999). In the case of the  
275 Red Kite, abundance of vectors has never been investigated directly in the previous  
276 studies published but it was assumed that the low prevalence measured in Spain  
277 could be due to the dryness of the environment, making it unsuitable for the main  
278 vectors of haemosporidian parasites (Tella et al. 1999). In Franche-Comté, the  
279 climate is more wet (annual rainfall from 1000 to 1500 mm against 600 mm in Spain)  
280 and numerous ponds, lakes and streams composed the landscape of the studied  
281 areas, which could at least partly explain the relatively high prevalence of  
282 *Leucocytozoon*. Moreover, the world range of the Red Kite limited to Western  
283 Europe, the absence of inter-continental migration notably through Africa and the  
284 fact that it breeds in open-woodland habitat are all together factors that may explain

285 relatively low Haemosporida prevalence in raptors and more generally animals  
286 (Tella et al. 1999; Altizer et al. 2011).

287 Lower body condition has been related to infection status (i.e., infected or not) by  
288 haemosporidians in both passerines (Garvin et al. 2006, Hatchwell et al. 2000) and  
289 raptors (Dawson and Bortolotti 2000). We did not show any effect of infection of Red  
290 Kite by *Leucocytozoon* on body condition, which suggests that these blood parasites  
291 might have no or weak pathogenic consequences on nestlings and/or that the  
292 infection intensity is too low to impair condition. In the American kestrel *Falco*  
293 *sparverius*, Dawson and Bortolotti (2000) found that a reduced condition of males  
294 was associated with higher intensity of infections by *Haemoproteus* using blood  
295 smear as a diagnostic tool. Moreover, Schoenle et al. (2017) reported that  
296 measuring intensity infection rather than only presence/absence could be important  
297 when evaluating the role of infection in influencing hosts' behaviour, physiology or  
298 fitness. If the molecular methods we used are considered as relevant to diagnose  
299 infection status of Red Kite by blood parasites (Martinsen et al. 2006, Krone et al.  
300 2008), blood smears remain useful to measure the infection intensity in individuals  
301 and should be implemented complementarily to PCR screening in further studies.

302 Overall, both the low to medium prevalences reported in the different European  
303 populations studied and the absence of negative correlation with body condition we  
304 show suggest that hematozoans in Red Kite nestlings could be a limited threat for  
305 this life stage even if the detection of *P. relictum* in one chick questions about its  
306 occurrence at a larger scale and possible health effects on some individuals. In  
307 addition, recent findings showed that low-level chronic malaria infection reduced life  
308 span as well as the lifetime number and quality of offspring in a wild population of  
309 great reed warblers *Acrocephalus arundinaceus*, these delayed fitness effects of  
310 malaria being mediated by telomere degradation (Asghar et al. 2015). This stresses  
311 the need for further research on "subtle" hematozoan effect on individual in different

312 avian groups including raptors. Moreover, the prevalence in adults has been only  
313 assessed in Spain, a country where it is particularly low, while breeder's survival is a  
314 critical parameter in the dynamic of raptor populations (Saether and Bakke 2000).  
315 Further investigation associating both qualitative and quantitative methods on a  
316 significant number of Red Kites and including adults, from other European areas,  
317 notably in Germany and France, would be needed to confirm whether blood  
318 pathogens can be considered as negligible threat for the Red Kite population.

319

## 320 **References**

- 321 Altizer S, Bartel R, Han BA (2011) Animal migration and infectious disease risk.  
322 Science 331: 296–302
- 323 Asghar M, Hasselquist D, Hansson B, Zehtindjiev P, Westerdahl H, Bensch S (2015)  
324 Hidden costs of infection: Chronic malaria accelerates telomere degradation and  
325 senescence in wild birds. Science 347: 436–438
- 326 Bennett GF, Peirce MA, Ashford RW (1993) Avian haematozoa: mortality and  
327 pathogenicity. J Nat Hist 27: 993–1001
- 328 BirdLife International (2020) Species factsheet: *Milvus milvus*. <http://www.birdlife.org>.  
329 Accessed 03 March 2020
- 330 Burnham KP, Anderson DR (2010) Model selection and multimodel inference: a  
331 practical information-theoretic approach. 2nd edn. Springer Press, New York
- 332 Castellucci SA, Oplinger SB, Klucsarits JR (1998) Hematology and detection of  
333 hemoparasites in the American Kestrel (*Falco sparverius*) during summer nesting  
334 period. J Pa Acad Sci 72: 29–31

- 335 Coeurdassier M, Crini N, Amiot C, Fourel I, Berny P, Brand J, Scheifler R, Fritsch C,  
336 Faggio G, Mionnet A, Morin C, Riols R (2014) Exposure of nestlings Red Kite to  
337 rodenticides, PAHs and metals. 24th SETAC Europe Meeting, Basel
- 338 Dawson RD, Bortolotti GR (2000) Effects of hematozoan parasites on condition and  
339 return rates of American kestrels. *Auk* 117: 373–380
- 340 Fair J, Paul E, Jones J (2010) Guidelines to the use of wild birds in research. Retrieved  
341 from [http://sccp.ca/sites/default/files/species-](http://sccp.ca/sites/default/files/species-habitat/documents/Guidelines%20to%20the%20use%20of%20wild%20birds%20in%20researchornithological%20council_August2010.pdf)  
342 [habitat/documents/Guidelines%20to%20the%20use%20of%20wild%20birds%20in%20r](http://sccp.ca/sites/default/files/species-habitat/documents/Guidelines%20to%20the%20use%20of%20wild%20birds%20in%20researchornithological%20council_August2010.pdf)  
343 [esearchornithological%20council\\_August2010.pdf](http://sccp.ca/sites/default/files/species-habitat/documents/Guidelines%20to%20the%20use%20of%20wild%20birds%20in%20researchornithological%20council_August2010.pdf)
- 344 Forrester DJ, Telford SR, Foster GW, Bennett GF (1994) Blood parasites of raptors in  
345 Florida. *J Raptor Res* 28: 226–231
- 346 Garvin MC, Szell CC, Moore FR (2006) Blood parasites of nearctic–neotropical migrant  
347 passerine birds during spring trans-gulf migration: impact on host body condition. *J*  
348 *Parasitol* 92: 990–996
- 349 Grafen A, Hails R (2002) *Modern statistics for the life sciences*. Oxford University  
350 Press, Oxford
- 351 Griffiths R, Double MC, Orr K, Dawson RJ (1998) A DNA test to sex most birds. *Mol*  
352 *Ecol* 7: 1071–1075
- 353 Gutiérrez-López R, Gangoso L, Martínez-de la Puente J, Fric J, López-López P,  
354 Mailleux M, Muñoz J, Touati L, Samraoui B, Figuerola J (2015) Low prevalence of  
355 blood parasites in a long-distance migratory raptor: the importance of host habitat.  
356 *Parasite Vector* 8: 189. <https://doi.org/10.1186/s13071-015-0802-9>.
- 357 Hanel J, Doležalová J, Stehlíková Š, Modrý D, Chudoba J, Synek P, Votýpka J (2016)  
358 Blood parasites in Northern Goshawk (*Accipiter gentilis*) with an emphasis to  
359 *Leucocytozoon toddi*. *Parasitol Res* 115: 263–270




- 360 Hatchwell BJ, Wood MJ, Anwar MA, Chamberlain DE, Perrins CM (2001) The  
361 haematozoan parasites of common blackbirds *Turdus merula*: associations with host  
362 condition. *Ibis* 143: 420–426
- 363 Jeffries MI, Miller RA, Laskowski MD, Carlisle JD (2015) High prevalence of  
364 *Leucocytozoon* parasites in nestling Northern Goshawks (*Accipiter gentilis*) in the  
365 Northern Great Basin, U.S.A. *J Raptor Res* 49: 294–302
- 366 Korpimäki E, Hakkarainen H, Bennett GF (1993) Blood parasites and reproductive  
367 success of Tengmalm's Owls: detrimental effects on females but not on males? *Funct*  
368 *Ecol* 7: 420-424.
- 369 Krone O, Priemer J, Streich J, Sommer P, Langgemach T, Lessow O (2001)  
370 Haemosporida of birds of prey and owls from Germany. *Acta Protozool* 40: 281–290
- 371 Krone O, Waldenström J, Valkiūnas G, Lessow O, Müller K, Iezhova TA, Fickel J,  
372 Bensch S (2008) Haemosporidian blood parasites in European birds of prey and  
373 owls. *J Parasitol* 94: 709–715
- 374 Labocha MK, Hayes JP (2012) Morphometric indices of body condition in birds: a  
375 review. *J Ornithol* 153: 1–22
- 376 Laursen IM (2008) Environmental contamination, blood parasites, and home range of  
377 the Red Kite (*Milvus milvus*) in Denmark. Master of Science Dissertations, University  
378 of Copenhagen
- 379 Martinsen ES, Paperna I, Schall JJ (2006) Morphological versus molecular identification of  
380 avian Haemosporida: An exploration of three species concepts. *Parasitology* 133: 279–  
381 288
- 382 Mougeot F, Bretagnolle V (2006) Breeding biology of the Red Kite *Milvus milvus* in  
383 Corsica. *Ibis* 148: 436–448

- 384 Nourani L, Zakeri S, Djadid ND (2020) Dynamics of prevalence and distribution pattern  
385 of avian *Plasmodium* species and its vectors in diverse zoogeographical areas - A  
386 review. *Infect Genet Evol* 104244. <https://doi.org/10.1016/j.meegid.2020.104244>
- 387 Peirce MA (1981) Distribution and host-parasite check-list of the Haematozoa of birds  
388 in Western Europe. *J Nat Hist* 15: 419–458
- 389 Peirce MA, Greenwood AG, Cooper JE (1983) Haematozoa of raptors and other birds  
390 from Britain, Spain and the United Arab emirates. *Avian Pathol* 12: 443–446
- 391 Pérez-Rodríguez A, de la Puente J, Onrubia A, Pérez-Tris J (2013) Molecular  
392 characterization of haemosporidian parasites from kites of the genus *Milvus* (Aves:  
393 Accipitridae). *Int J Parasitol* 43: 381–387
- 394 Perez-Tris J, Bensch S (2005). Diagnosing genetically diverse avian malarial infections  
395 using mixed-sequence analysis and TA-cloning. *Parasitology* 131: 15-23
- 396 Podmokła E, Dubiec A, Drobniak SM, Arct A, Gustafsson L, Cichoń M (2014)  
397 Determinants of prevalence and intensity of infection with malaria parasites in the  
398 Blue Tit. *J Ornithol* 155: 721–727
- 399 Poulin R (1996) Helminth growth in vertebrate hosts: Does host sex matter? *Int J*  
400 *Parasitol* 26: 1311–1315
- 401 Powolny T, Bassin N, Crini N, Fourel I, Morin C, Pottinger TG, Massemin S, Zahn S,  
402 Coeurdassier M (2020) Corticosterone mediates telomere length in raptor chicks  
403 exposed to chemical mixture. *Sci Total Environ* 706: 135083
- 404 Remple JD (2004) Intracellular Hematozoa of raptors: a review and update. *J Avian*  
405 *Med Surg* 18: 75–88
- 406 Saether B-E, Bakke O (2000) Avian life history variation and contribution of  
407 demographic traits to the population growth rate. *Ecology* 81: 642–653

- 408 Schalk G, Forbes MR (1997) Male biases in parasitism of mammals: effects of study  
409 type, host age, and parasite taxon. *Oikos* 78: 67-74
- 410 Schoenle LA, Kernbach M, Haussmann MF, Bonier F, Moore IT (2017) An experimental  
411 test of the physiological consequences of avian malaria infection. *J Anim Ecol* 86: 1483-  
412 1496
- 413 Sehgal RNM (2015) Manifold habitat effects on the prevalence and diversity of avian  
414 blood parasites. *Int J Parasitol* 4: 421–430
- 415 Sol D, Jovani R, Torres J (2003) Parasite mediated mortality and host immune  
416 response explain age-related differences in blood parasitism in birds. *Oecologia* 135:  
417 542–547
- 418 Tella JL, Blanco G, Forero MG, Gajon A, Donazar JA, Hiraldo, F. (1999) Habitat, world  
419 geographic range, and embryonic development of hosts explain the prevalence of  
420 avian Hematozoa at small spatial and phylogenetic scales. *Proc Natl Acad Sci U.S.A*  
421 96: 1785–1789
- 422 Toyne EP, Ashford RW (1997) Blood parasites of nestling goshawks. *J Raptor Res* 31:  
423 81–83
- 424 UICN France, MNHN, LPO, SEOF, ONCFS (2016) La Liste rouge des espèces  
425 menacées en France - Oiseaux de France métropolitaine.  
426 [https://inpn.mnhn.fr/docs/LR\\_FCE/UICN-LR-Oiseaux-diffusion.pdf](https://inpn.mnhn.fr/docs/LR_FCE/UICN-LR-Oiseaux-diffusion.pdf). Accessed 11  
427 March 2020 (in French)
- 428 Valkiūnas G, Sehgal RNM, Iezhova TA, Hull AC (2010) Identification of *Leucocytozoon*  
429 *toddi* group (Haemosporida: Leucocytozoidae), with remarks on the species  
430 taxonomy of leucocytozoids. *J Parasitol* 96: 170–177
- 431 Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and  
432 extensions in ecology with R. Springer Press, New York

433 *Figure caption*

434 **Fig. 1** Body mass of the 47 Red Kite chicks studied standardised against wing  
 435 length taken as a proxy of age (Mougeot and Bretagnolle 2006) for females and  
 436 males separately. The individuals for which DNA from *Leucocytozoon* ssp. was  
 437 detected in blood were shown as a black circle in which the corresponding  
 438 haplotype 1 (L-MILANS4, Genbank accession number JN164713) and/or 2 (L-  
 439 MILVUS1, JN164716) is reported. The grey and black hatched dot corresponds to  
 440 the male infected by both *Leucocytozoon* (haplotype 2) and *Plasmodium relictum*.



441

442 **Table 1** Linear mixed-effect models tested to explain body mass as response. The best  
 443 model selected based on corrected Akaike information criteria is reported in bold (LL:  
 444 maximized log-likelihood; K: number of estimated parameters; AICc: Corrected Akaike's  
 445 information criterion; w\_ic: Akaike weight)

Model tested (Body mass ~ )	LL	K	AICc	$\Delta$ AICc	w_ic
Wing length + Sex + (1 Area/Nest)	<b>-258.9</b>	<b>6</b>	<b>532.0</b>	<b>0.00</b>	<b>0.39</b>
Wing length + Sex * Status + (1 Area/Nest)	-256.5	8	532.9	0.89	0.25
Wing length + Sex + Status + (1 Area/Nest)	-258.1	7	533.0	1.03	0.23
Wing length * Sex + (1 Area/Nest)	-258.7	7	534.2	2.20	0.13
Wing length + (1 Area/Nest)	-266.1	5	543.6	11.57	0.00

446

447 **Table 2** Summary of the data available on Red Kite infection by the haemosporidian  
 448 parasites *Leucocytozoon spp.* (*Leuco.*), *Haemoproteus spp.* (*Haemop.*) and  
 449 *Plasmodium spp.* (*Plasmo.*). (n = total number of screened individuals; *nr* = not  
 450 reported; *ns* = not screened)

Area/country	Age	n	Prevalence in % [ <i>n</i> of infected kites]			Reference
			<i>Leuco.</i>	<i>Haemop.</i>	<i>Plasmod.</i>	
Western Europe	<i>nr</i>	4	25 [1]	25 [1]	0 [0]	1
Germany	Nestling	13	15 [2]	0 [0]	0 [0]	2
	Juvenile	2	0 [0]	0 [0]	0 [0]	
	Adult	9	0 [0]	0 [0]	0 [0]	
	<i>nr</i>	2	0 [0]	0 [0]	0 [0]	3
UK	<i>nr</i>	1	0 [0]	0 [0]	0 [0]	4
Spain	<i>nr</i>	46	0 [0]	0 [0]	0 [0]	5
	Juvenile	164	4 [6]	0 [0]	0 [0]	6
	Adult	104	2 [2]	0 [0]	0 [0]	
Denmark	Nestling	17	18 [3]	0 [0]	0 [0]	7
	Juvenile	1	0 [0]	0 [0]	0 [0]	
	Adult	1	0 [0]	0 [0]	0 [0]	
France	Nestling	47	28 [13]	0 [0]	2 [1]	This study

451 1: Peirce (1981), 2: Krone *et al.* (2001), 3: Krone *et al.* (2008), 4: Peirce *et al.* (1983), 5: Tella *et al.* (1999),  
 452 6: Pérez-Rodríguez *et al.* (2013), 7: Laursen (2008).