



HAL
open science

The Quantitative Collapse of Concurrent Games with Symmetry

Pierre Clairambault, Hugo Paquet

► **To cite this version:**

Pierre Clairambault, Hugo Paquet. The Quantitative Collapse of Concurrent Games with Symmetry. 2023. hal-03280420v2

HAL Id: hal-03280420

<https://hal.science/hal-03280420v2>

Preprint submitted on 18 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE QUANTITATIVE COLLAPSE OF CONCURRENT GAMES WITH SYMMETRY

PIERRE CLAIRAMBAULT AND HUGO PAQUET

Univ Lyon, EnsL, UCBL, CNRS, LIP, F-69342, LYON Cedex 07, France
e-mail address: pierre.clairambault@ens-lyon.fr

University of Oxford
e-mail address: hugo.paquet@cs.ox.ac.uk

ABSTRACT. We explore links between the thin concurrent games of Castellan, Clairambault and Winskel, and the weighted relational models of linear logic studied by Laird, Manzonetto, McCusker and Pagani. More precisely, we show that there is an interpretation-preserving “collapse” functor from the former to the latter. On objects, the functor defines for each game a set of possible *execution states*. Defining the action on morphisms is more subtle, and this is the main contribution of the paper.

Given a strategy and an execution state, our functor needs to *count* the witnesses for this state within the strategy. Strategies in thin concurrent games describe non-linear behaviour explicitly, so in general each witness exists in countably many symmetric *copies*. The challenge is to define the right notion of witnesses, factoring out this infinity while matching the weighted relational model. Understanding how witnesses compose is particularly subtle and requires a delve into the combinatorics of witnesses and their symmetries.

In its basic form, this functor connects thin concurrent games and a relational model weighted by $\mathbb{N} \cup \{+\infty\}$. We will additionally consider a generalised setting where both models are weighted by elements of an arbitrary continuous semiring; this covers the probabilistic case, among others. Witnesses now additionally carry a value from the semiring, and our interpretation-preserving collapse functor extends to this setting.

1. INTRODUCTION

The *relational model* is one of the simplest model of linear logic. It naturally gives rise to a model of higher-order programming often described as *quantitative*, because aspects of computation such as the multiplicity of function calls are represented explicitly. The model assigns to every type a set known as its *web*, whose elements are thought of as (desequentialized) execution states, and to any term, a relation.

Relations are equivalently boolean-valued matrices, and a natural extension of the model consists in considering more general coefficients. This idea was first explored by Lamarche [Lam92] and developed in detail by Laird, Manzonetto, McCusker and Pagani [LMMP13, Lai20]. Their construction gives a family of *weighted relational models*, in which the interpretation of a term is a matrix assigning to each point of the web a *weight*, coming

Key words and phrases: denotational semantics, concurrent games, relational models.

from a continuous semiring \mathcal{R} , the *resource semiring*. This has an operational interpretation for a program M : while the relational model only indicates whether a given state α *can* be realized by an execution of M (*i.e.* do we have $\alpha \in \llbracket M \rrbracket?$), the weighted relational model aggregates information about all executions that realize this state. In the simplest case, $\mathcal{R} = \mathbb{N} \cup \{+\infty\}$, the model simply *counts* these executions. Using various semirings one can adequately represent probabilistic evaluation, best and worst case time analysis, etc.

Another well-established quantitative model – or rather, family of models – is game semantics [HO00, AJM00]. In game semantics, an execution is regarded as a play in a two-player game between Player (playing for the program), and Opponent (playing for the environment). Types are presented as games, whose rules specify the possible executions, and terms as strategies describing the interactive behaviour of the program under any evaluation context. The connections between game semantics and relational semantics have been thoroughly studied [BDER97, Mel05, Bou09, Ong17]. In particular, the family of *concurrent games* [RW11, CCW19] inherit from Melliès’ *asynchronous games* [Mel05] a particularly neat relationship with relational semantics. In this framework, both games and strategies are *event structures* [Win86], and as such admit a canonical notion of state/position: the *configurations*. As we will see, the web can be recovered as a subset of the configurations of the game. Then, we can “collapse” a strategy into a relation, by recording which of these configurations are reached, and forgetting the chronological history (see [Cla19] for a recent account). In an affine setting, *i.e.* without replication, this collapse operation can immediately be generalised to weighted relations with $\mathcal{R} = \mathbb{N} \cup \{+\infty\}$: if σ is a strategy on a game A , and x is (a point of the web corresponding to) a configuration of A , the collapse simply *counts* the distinct configurations of σ realizing x .

The difficulty arises in the non-affine setting, necessary for languages with duplication of resources. For this, the mature extension of concurrent games is *thin concurrent games with symmetry* [CCW19]. In thin concurrent games, infinite games arise from the construction $!A$, which creates countably many copies of A labelled with natural numbers, called *copy indices*. (This is similar to the situation in AJM games [AJM00].) Games are equipped with sets of bijections called *symmetries* (so they are *event structures with symmetry* [Win07]) which specify authorized reindexings. Additionally strategies must act uniformly with respect to these symmetries. The collapse to the relational model is relatively undisturbed by symmetry: points of the web now correspond to a subset of *symmetry classes* of configurations and, as before, we collapse a strategy to the set of symmetry classes it reaches (see *e.g.* [CC21]). However the extension to the *weighted* relational model is no longer obvious. We cannot simply count all concrete configurations of σ witnessing some symmetry class: there are infinitely many. This prompts the central question of this paper: how can we count configurations up to symmetry, in order to match coefficients of the weighted relational model? In other words: what does the weighted relational model count?

An answer to this question is our main contribution. For a symmetry class of configurations of the game, an apparent “obvious” solution is to consider the set of corresponding *symmetry classes* of configurations of the strategy. Surprisingly, the induced coefficient is wrong! Instead we are led to introduce a notion of *positive witnesses* for a given symmetry class of configurations of the game. We will show that counting positive witnesses yields an interpretation-preserving collapse to the relational model weighted by $\mathbb{N} \cup \{+\infty\}$. In proving this, the main challenge is functoriality of the collapse: whereas in the affine case, witnesses in a composite $\tau \odot \sigma$ cleanly correspond to pairs of witnesses in σ and τ , this fails with symmetry, and is only salvaged via a proper account of symmetries on both sides.

Finally, we also extend our results to any continuous semiring \mathcal{R} with a condition called *integer division*. We consider an extension of thin concurrent games where strategies carry valuations in \mathcal{R} , and show that this collapses into the \mathcal{R} -weighted relational model.

Related work. To our knowledge, the first quantitative collapse from games to relations is from *probabilistic* concurrent games to the relational model weighted by $\overline{\mathbb{R}}_+ = \mathbb{R} \cup \{+\infty\}$ in [CCPW18]. However, when working on an extension to a quantum language [CdV20], the first author discovered an error in [CCPW18]: the paper uses symmetry classes as witnesses, which —as we show here— is inadequate. The correct notion of witness and its validity *w.r.t.* composition was established by the first author in an unpublished report [Cla20]. Here we complete this to a full interpretation-preserving functor and to the \mathcal{R} -weighted case. In particular, Theorem 6.15 for $\mathcal{R} = \overline{\mathbb{R}}_+$ corrects the collapse theorem of [CCPW18].

Outline. In Section 2 we recall the \mathcal{R} -weighted relational model and the language of concern for most of the paper, a non-deterministic **PCF**. In Section 3, we recall *thin concurrent games* and the corresponding interpretation of non-deterministic **PCF**. In Section 4 we address the main challenge of the paper, the definition of positive witnesses and their compatibility with composition. In Section 5, we fix $\mathcal{R} = \mathbb{N} \cup \{+\infty\}$, show a number of properties ensuring that the interpretation is preserved, and prove our main result (Theorem 5.20). Finally, in Section 6 we generalize the result to an arbitrary \mathcal{R} (Theorem 6.15).

2. THE WEIGHTED RELATIONAL MODEL AND **nPCF**

Notations. If X is a set, we write $\mathcal{P}_f(X)$ for the finite subsets, and $\mathcal{M}_f(X)$ for finite multisets. For $x_1, \dots, x_n \in X$, $[x_1, \dots, x_n] \in \mathcal{M}_f(X)$ is the corresponding multiset. We use $\mu, \nu \in \mathcal{M}_f(X)$ to range over multisets; and write $\mu + \nu \in \mathcal{M}_f(X)$ for the sum of multisets, where $x \in X$ has multiplicity the sum of its multiplicities in X and Y . If $\mathcal{R} = (|\mathcal{R}|, +, \cdot, 0, 1)$ is a semiring and $x, y \in |\mathcal{R}|$, the *Kronecker symbol* $\delta_{x,y}$ means 1 if $x = y$, and 0 otherwise. If X is a set, we write $\sharp X \in \mathbb{N} \cup \{+\infty\}$ for its cardinality if X is finite, $+\infty$ otherwise.

We assume some familiarity with categorical logic, in particular Seely categories [Mel09].

2.1. Continuous semirings. We first recall the construction of the *\mathcal{R} -weighted relational model*, where \mathcal{R} is a continuous semiring of *resources*. Our presentation follows [LMMP13].

A **complete partial order (cpo)** is a poset (X, \leq) with a bottom and such that any directed subset $D \subseteq X$ has a sup $\vee D \in X$. For a cpo X , $F : X \rightarrow X$ is **continuous** if it is monotone and preserves all suprema of directed sets, *i.e.* $F(\vee D) = \vee(F(D))$. An n -ary function $X^n \rightarrow X$ is continuous if it is continuous in each of its parameters.

Definition 2.1. A **continuous semiring** \mathcal{R} is a semiring $(|\mathcal{R}|, +, \cdot, 0, 1)$ equipped with a partial order \leq such that $(|\mathcal{R}|, \leq)$ is a cpo with 0 as bottom, and $+$ and \cdot are continuous.

We often denote the carrier set $|\mathcal{R}|$ just by \mathcal{R} . The point of considering the ordered structure on \mathcal{R} , is that for any \mathcal{R} and possibly infinite subset $S \subseteq \mathcal{R}$, the indexed sum

$$\sum_{x \in S} x = \bigvee_{F \subseteq_f S} \left(\sum_{x \in F} x \right) \quad (2.1)$$

is always defined as the supremum of all the partial sums.

We impose two further conditions on continuous semirings. As in [LMMP13], they should be **commutative**: $r \cdot r' = r' \cdot r$ for all $r, r' \in \mathcal{R}$. Additionally, they should *have integer division*. If $x \in \mathcal{R}$ and $n \in \mathbb{N}$ is an integer, then one may define $n * x = x + \dots + x$ (with n occurrences of x). We say that \mathcal{R} has **integer division** if for all $n \geq 1$, for all $x, y \in \mathcal{R}$, if $n * x = n * y$ then $x = y$. Unlike commutativity, this condition is not required in [LMMP13]; nevertheless, all examples considered in [LMMP13] do have integer division. From now on, we assume all continuous semirings satisfy these two conditions.

Our core example of a continuous semiring is the following:

Definition 2.2. We write \mathcal{N} for the continuous semiring $(|\mathcal{N}|, +, \cdot, 0, 1)$ equipped with the standard order on \mathbb{N} extended with $x \leq +\infty$ for all $x \in \mathcal{N}$. To ensure continuity we take $+$ to be the usual sum extended with $(+\infty) + x = x + (+\infty) = +\infty$, and \cdot to be multiplication extended with $+\infty \cdot 0 = 0 \cdot +\infty = 0$, and $+\infty \cdot x = x \cdot +\infty = +\infty$ for any $x > 0$.

As described in [LMMP13], \mathcal{N} may be used to *count* operational reduction sequences in a non-deterministic language. There are other examples [LMMP13], including the completed non-negative reals $\overline{\mathbb{R}}_+$, which provide an adequate model for **PCF** with probabilistic choice. (\mathcal{N} has a canonical place among those examples, because it is an initial object in the category of continuous semirings and structure-preserving continuous maps.)

2.2. Weighted relations. We fix a continuous semiring \mathcal{R} and define the category $\mathcal{R}\text{-Rel}$ of \mathcal{R} -weighted relations. An **\mathcal{R} -relation** from a set X to a set Y is simply a function

$$\alpha : X \times Y \rightarrow \mathcal{R},$$

also written $\alpha : X \rightarrow Y$, regarded as a matrix with coefficients in \mathcal{R} . We usually write $\alpha_{x,y}$ for the coefficient $\alpha(x, y) \in \mathcal{R}$. For $\alpha : X \rightarrow Y$ and $\beta : Y \rightarrow Z$ and $x \in X, z \in Z$, we set

$$(\beta \circ \alpha)_{x,z} = \sum_{y \in Y} \alpha_{x,y} \cdot \beta_{y,z} \quad (2.2)$$

for the coefficients of the **composition** $\beta \circ \alpha : X \rightarrow Z$. For X a set, the **identity** on X has $(\text{id}_X)_{x,x'} = \delta_{x,x'}$; *i.e.* the diagonal matrix on X with only 1's as diagonal coefficients.

Proposition 2.3. *For any continuous semiring \mathcal{R} , there is a category $\mathcal{R}\text{-Rel}$ with sets as objects, and \mathcal{R} -relations from X to Y as morphisms.*

2.3. Categorical structure. $\mathcal{R}\text{-Rel}$ is a Seelye category: it is symmetric monoidal closed with finite products and a linear exponential comonad. We review this structure now. In fact, although this is not true in general for Seelye categories, $\mathcal{R}\text{-Rel}$ is *compact closed*.

2.3.1. Compact closed structure. The *tensor* $X \otimes Y$ of two sets X, Y , is simply their cartesian product $X \times Y$. The **tensor** of $\alpha_1 : X_1 \rightarrow Y_1$ and $\alpha_2 : X_2 \rightarrow Y_2$ has coefficients

$$(\alpha_1 \otimes \alpha_2)_{(x_1, x_2), (y_1, y_2)} = (\alpha_1)_{x_1, y_1} \cdot (\alpha_2)_{x_2, y_2}$$

for $(x_1, x_2) \in X_1 \otimes X_2$ and $(y_1, y_2) \in Y_1 \otimes Y_2$, yielding $\alpha_1 \otimes \alpha_2 : X_1 \otimes X_2 \rightarrow Y_1 \otimes Y_2$.

This operation yields a bifunctor $- \otimes - : \mathcal{R}\text{-Rel} \times \mathcal{R}\text{-Rel} \rightarrow \mathcal{R}\text{-Rel}$ completed with

$$\begin{aligned} \alpha_{X,Y,Z} &: (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z) \\ \lambda_X &: 1 \otimes X \rightarrow X \\ \rho_X &: X \otimes 1 \rightarrow X \\ s_{X,Y} &: X \otimes Y \rightarrow Y \otimes X \end{aligned}$$

defined as the obvious variants of the identity matrix, where $1 = \{\bullet\}$ is a singleton set. Those satisfy the necessary naturality and coherence properties, making $\mathcal{R}\text{-Rel}$ a symmetric monoidal category. Furthermore, any set X has a *dual* X^* defined simply as X itself, and

$$\eta_X : 1 \rightarrow X \times X, \quad \epsilon_X : X \times X \rightarrow 1$$

turn $\mathcal{R}\text{-Rel}$ into a *compact closed category*. In particular, it follows that $\mathcal{R}\text{-Rel}$ is automatically *symmetric monoidal closed*. For X and Y any two sets, this gives us a notion of *linear arrow* $X \multimap Y$, defined simply as $X \times Y$. We also get a **currying** bijection for X, Y, Z :

$$\Lambda : \mathcal{R}\text{-Rel}(X \otimes Y, Z) \simeq \mathcal{R}\text{-Rel}(X, Y \multimap Z)$$

with, for any $\alpha : X \otimes Y \rightarrow Z$, $\Lambda(\alpha)_{x,(y,z)} = \alpha_{(x,y),z}$. We also get an **evaluation** morphism: $\text{ev}_{X,Y} : (X \multimap Y) \otimes X \rightarrow Y$ defined as having coefficients $(\text{ev}_{X,Y})_{((x,y),x'),y'} = \delta_{x,x'} \cdot \delta_{y,y'}$.

2.3.2. Cartesian structure. Furthermore, $\mathcal{R}\text{-Rel}$ is cartesian. First, the empty set \emptyset is a terminal object, also written \top . If X, Y are sets, we define $X \& Y$ as $X + Y$ their **tagged disjoint union**, defined as $(\{1\} \times X) \uplus (\{2\} \times Y)$. Note that here and from now on, we use \uplus to denote the standard set-theoretic union, when it is known to be disjoint. We have

$$\pi_1 : X \& Y \rightarrow X, \quad \pi_2 : X \& Y \rightarrow Y,$$

the **projections** respectively defined as $(\pi_1)_{(i,x),y} = 1$ if $i = 1$ and $x = y$, and 0 otherwise – π_2 is defined symmetrically. For $\alpha : X \rightarrow Y$ and $\beta : X \rightarrow Z$, their **pairing** is

$$\langle \alpha, \beta \rangle : X \rightarrow Y \& Z,$$

defined with $\langle \alpha, \beta \rangle_{(1,x),z} = \alpha_{x,z}$ and $\langle \alpha, \beta \rangle_{(2,y),z} = \beta_{y,z}$. This makes $\mathcal{R}\text{-Rel}$ a cartesian category. One must keep in mind that $\mathcal{R}\text{-Rel}$ is *not* cartesian closed, as the closed structure is with respect to the tensor \otimes and not the cartesian product $\&$.

2.3.3. Linear exponential comonad. We define a comonad $!$ on $\mathcal{R}\text{-Rel}$. On objects the operation $X \mapsto !X$ constructs the free commutative comonoid: this is defined as $!X = \mathcal{M}_f(X)$. The action on morphisms is determined by the universal property of $!X$, but we give an explicit definition. For $\alpha : X \rightarrow Y$, we set

$$(!\alpha)_{\mu,[y_1,\dots,y_n]} = \sum_{\substack{(x_1,\dots,x_n), \text{ s.t. } 1 \leq i \leq n \\ \mu = [x_1,\dots,x_n]}} \prod \alpha_{x_i,y_i}.$$

Note that $!\alpha$ only has nonzero coefficients for pairs of multisets of the same size. Likewise, we define $\epsilon_X : !X \rightarrow X$, $\delta_X : !X \rightarrow !!X$, $m_{X,Y} : !X \otimes !Y \rightarrow !(X \& Y)$ with

$$\begin{aligned} (\epsilon_X)_{\mu,x} &= \delta_{\mu,[x]} \\ (\delta_X)_{\mu,[\nu_1,\dots,\nu_n]} &= \delta_{\mu,\nu_1+\dots+\nu_n} \\ (m_{X,Y})_{([x_1,\dots,x_n],[y_1,\dots,y_p]),\mu} &= \delta_{\mu,[(1,x_1),\dots,(1,x_n),(2,y_1),\dots,(2,y_p)]} \end{aligned}$$

$\overline{\Gamma \vdash \mathbf{tt} : \mathbb{B}}$	$\overline{\Gamma \vdash \mathbf{ff} : \mathbb{B}}$	$\overline{\Gamma \vdash n : \mathbb{N}}$	$\overline{\Gamma, x : A \vdash x : A}$	$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x^A. M : A \rightarrow B}$
$\frac{\Gamma \vdash M : A \rightarrow B \quad \Gamma \vdash N : A}{\Gamma \vdash MN : B}$		$\frac{\Gamma \vdash M : \mathbb{B} \quad \Gamma \vdash N_1 : \mathbb{X} \quad \Gamma \vdash N_2 : \mathbb{X}}{\Gamma \vdash \mathbf{if} M N_1 N_2 : \mathbb{X}}$		
$\frac{\Gamma \vdash M : \mathbb{N}}{\Gamma \vdash \mathbf{succ} M : \mathbb{N}}$	$\frac{\Gamma \vdash M : \mathbb{N}}{\Gamma \vdash \mathbf{pred} M : \mathbb{N}}$	$\frac{\Gamma \vdash M : \mathbb{N}}{\Gamma \vdash \mathbf{iszero} M : \mathbb{B}}$	$\frac{\Gamma \vdash M : A \rightarrow A}{\Gamma \vdash \mathcal{Y} M : A}$	

Figure 1: Typing rules for **PCF**

and $m^0 : 1 \rightarrow !\top$ is defined as 1 on its only point $(\bullet, [])$. We have defined all the structure of a Seely category [Mel09], and the necessary axioms can be verified. We obtain that the Kleisli category $\mathcal{R}\text{-Rel}_!$ is cartesian closed.

2.3.4. *Recursion.* For the interpretation of **nPCF** in $\mathcal{R}\text{-Rel}$ we must give structure for recursion. First, for any sets X, Y , we order the homset $\mathcal{R}\text{-Rel}(X, Y)$ pointwise, i.e.

$$\alpha \leq \beta \quad \Leftrightarrow \quad \forall x \in X, y \in Y, \alpha_{x,y} \leq_{\mathcal{R}} \beta_{x,y}.$$

It is straightforward that this defines a cpo, with bottom \perp the zero matrix. All operations on weighted relations involved in the Seely category structure (i.e. composition, tensor, pairing and $!(-)$) are continuous with respect to this order.

As all operations are continuous, we can define, for every set X , a continuous operator

$$\begin{aligned} F & : \mathcal{R}\text{-Rel}_!(\top, !(X \multimap X) \multimap X) \rightarrow \mathcal{R}\text{-Rel}_!(\top, !(X \multimap X) \multimap X) \\ \alpha & \mapsto \lambda f. f(\alpha f) \end{aligned}$$

where the λ -calculus notation is well-defined since $\mathcal{R}\text{-Rel}_!$ is cartesian closed. We then define $\mathcal{Y}_X \in \mathcal{R}\text{-Rel}_!(\top, !(X \multimap X) \multimap X)$ as usual with

$$\mathcal{Y}_X = \bigvee_{n \in \mathbb{N}} F^n(\perp) \in \mathcal{R}\text{-Rel}_!(\top, !(X \multimap X) \multimap X),$$

and with a context Y , we set $\mathcal{Y}_{Y,X} = \mathcal{Y}_X \circ_! e_Y$ where $\circ_!$ denotes Kleisli composition and $e_Y \in \mathcal{R}\text{-Rel}_!(Y, \top)$ the terminal morphism, yielding $\mathcal{Y}_{Y,X} \in \mathcal{R}\text{-Rel}_!(Y, !(X \multimap X) \multimap X)$.

2.4. **Interpretation of nPCF.** Now, we define **nPCF** and its interpretation.

2.4.1. *Non-deterministic PCF.* The **types** of **nPCF** are given by the following grammar:

$$A, B ::= \mathbb{B} \mid \mathbb{N} \mid A \rightarrow B$$

where \mathbb{B} and \mathbb{N} are respectively types for *booleans* and *natural numbers*. We refer to \mathbb{B} and \mathbb{N} as *ground types*, and use \mathbb{X}, \mathbb{Y} to range over those. **Typed terms** are defined via the typing rules of Figure 1 – in this paper, all terms are well-typed. **Contexts** are lists of typed variables $x_1 : A_1, \dots, x_n : A_n$. **Typing judgments** have the form $\Gamma \vdash M : A$, where Γ is a context and A is a type. In addition to the rules listed in Figure 1, the language has an explicit exchange rule for permuting variable declarations in contexts. Conditionals are

$$\begin{array}{ll}
\text{if} : \ !(\mathbb{B} \ \& \ \mathbb{X} \ \& \ \mathbb{X}) \ \multimap \ \mathbb{X} & \text{succ} : \ \! \mathbb{N} \ \multimap \ \mathbb{N} \\
\text{if}_{\gamma,v} = 1 & \text{if } \gamma = [(1, \mathbf{tt}), (2, v)] \\
& \text{or } \gamma = [(1, \mathbf{ff}), (3, v)], \\
\text{if}_{\gamma,v} = 0 & \text{otherwise.} \\
\text{pred} : \ \! \mathbb{N} \ \multimap \ \mathbb{N} & \text{iszero} : \ \! \mathbb{N} \ \multimap \ \mathbb{B} \\
\text{pred}_{\gamma,n} = 1 & \text{if } \gamma = [0] \text{ and } n = 0, \\
& \text{or } \gamma = [k+1] \text{ and } n = k, \\
\text{pred}_{\gamma,n} = 0 & \text{otherwise.} \\
& \text{iszero}_{\gamma,b} = 1 \text{ if } \gamma = [0] \text{ and } b = \mathbf{tt}, \\
& \text{or } \gamma = [k+1] \text{ and } b = \mathbf{ff}, \\
& \text{iszero}_{\gamma,b} = 0 \text{ otherwise.}
\end{array}$$

Figure 2: Interpretation of basic **PCF** combinators

restricted to the base type, but as usual in call-by-name general conditionals can be defined as syntactic sugar. Finally, **nPCF** also has a **non-deterministic choice** with typing rule:

$$\overline{\Gamma \vdash \mathbf{coin} : \mathbb{B}}$$

We omit the operational semantics [LMMP13], and only recall that we get for $\vdash M : \mathbb{X}$ and value $v : \mathbb{X}$ a *weight*, defined as the number of distinct reduction sequences evaluating M to v . We write $M \Downarrow^n v$ if there are exactly n reduction sequences from M to v .

2.4.2. *Interpretation.* We may now define the interpretation of **nPCF** in $\mathcal{N}\text{-Rel}_!$.

To every type A we associate a set $\langle A \rangle$, its **web**, defined by $\langle \mathbb{B} \rangle = \{\mathbf{tt}, \mathbf{ff}\}$ and $\langle \mathbb{N} \rangle = \mathbb{N}$ the set of natural numbers, extended to all types via $\langle A \rightarrow B \rangle = \langle A \rangle \multimap \langle B \rangle$. For contexts:

$$\langle x_1 : A_1, \dots, x_n : A_n \rangle = \&_{1 \leq i \leq n} \langle A_i \rangle,$$

and terms $\Gamma \vdash M : A$ are interpreted as morphisms $\langle M \rangle \in \mathcal{N}\text{-Rel}_!(\langle \Gamma \rangle, \langle A \rangle)$. We omit the standard definitions for the λ -calculus constructions. For **PCF** combinators, we set:

$$\begin{aligned}
\langle \Gamma \vdash v : \mathbb{X} \rangle_{\gamma,v'} &= \delta_{\gamma, \square} \cdot \delta_{v,v'} \\
\langle \Gamma \vdash \mathbf{if} \ M \ N_1 \ N_2 : \mathbb{X} \rangle &= \mathbf{if} \circ_! \langle \langle M \rangle, \langle N_1 \rangle, \langle N_2 \rangle \rangle \\
\langle \Gamma \vdash \mathbf{pred} \ M : \mathbb{N} \rangle &= \mathbf{pred} \circ_! \langle M \rangle \\
\langle \Gamma \vdash \mathbf{succ} \ M : \mathbb{N} \rangle &= \mathbf{succ} \circ_! \langle M \rangle \\
\langle \Gamma \vdash \mathbf{iszero} \ M : \mathbb{B} \rangle &= \mathbf{iszero} \circ_! \langle M \rangle \\
\langle \Gamma \vdash \mathcal{Y} \ M : A \rangle &= \mathcal{Y}_{\langle \Gamma \rangle, \langle A \rangle} \langle M \rangle
\end{aligned}$$

with the weighted relations in Figure 2. Finally, we set $\langle \Gamma \vdash \mathbf{coin} : \mathbb{B} \rangle_{\gamma,v} = \langle \mathbf{tt} \rangle_{\gamma,v} + \langle \mathbf{ff} \rangle_{\gamma,v}$, so that $\langle \Gamma \vdash \mathbf{coin} : \mathbb{B} \rangle : \langle \Gamma \rangle \multimap \langle \mathbb{B} \rangle$ as required, concluding the interpretation of **nPCF**.

The reader is referred to [LMMP13] for the proof of the following adequacy property:

Theorem 2.4. *For any $\vdash M : \mathbb{X}$, for any value $v : \mathbb{X}$ and $n \in \mathbb{N}$, $M \Downarrow^n v$ iff $\langle M \rangle_{\square,v} = n$.*

2.4.3. *What does the \mathcal{N} -weighted relational model count?* Theorem 2.4 shows that at ground types, the \mathcal{N} -weighted relational model counts the distinct reduction sequences to a value. But for a general type A , a term $\vdash M : A$, and $x \in \langle A \rangle$, the meaning of the coefficient $\langle M \rangle_x \in \mathbb{N} \uplus \{+\infty\}$ is more difficult to describe. Even at higher-order we expect it to be related to the cardinality of some set of concrete witnesses: but which one?

In this paper we give one answer to this question, in terms of *concurrent games*.

3. CONCURRENT GAME SEMANTICS OF **nPCF**

Our games model of **nPCF** is based on *thin concurrent games* [CCW19], to which we add an *exhaustivity* mechanism inspired by Melliès [Mel05]. Game semantics is naturally *affine* and the purpose of exhaustivity is to ensure strict linearity, in order to establish a tighter correspondence with \mathcal{R} -Rel. As we will see, the two models can be related by a functor preserving some of the Seely category structure.

3.1. Event structures with symmetry. We start with preliminaries on *event structures with symmetry* [Win07], the mathematical structure on which thin concurrent games rest.

3.1.1. Event structures. Specifically, we use prime event structures with binary conflict:

Definition 3.1. An **event structure (es)** is a triple $E = (|E|, \leq_E, \#_E)$, where $|E|$ is a countable set of **events**, \leq_E is a partial order called **causal dependency** and $\#_E$ is an irreflexive symmetric binary relation on $|E|$ called **conflict**, satisfying:

$$\begin{aligned} \text{finite causes: } & \forall e \in |E|, [e]_E = \{e' \in |E| \mid e' \leq_E e\} \text{ is finite,} \\ \text{conflict inheritance: } & \forall e_1 \#_E e_2, \forall e_2 \leq_E e'_2, e_1 \#_E e'_2. \end{aligned}$$

We write $e \rightarrow_E e'$ for **immediate causality**, *i.e.* $e <_E e'$ with no event in between. A notion of critical importance for working with event structures is that of *configurations*:

Definition 3.2. A (finite) **configuration** of event structure E is a finite $x \subseteq |E|$ which is

$$\begin{aligned} \text{down-closed: } & \forall e \in x, \forall e' \in |E|, e' \leq_E e \implies e' \in x. \\ \text{consistent: } & \forall e, e' \in x, \neg(e \#_E e'). \end{aligned}$$

We write $\mathcal{C}(E)$ for the set of finite configurations on E .

The set $\mathcal{C}(E)$ is naturally ordered by inclusion; it is the *domain of configurations*. Configurations are typically ranged over by variables x, y, z . For $x, y \in \mathcal{C}(E)$, we write $x \dashv\!\! \dashv y$ if x is immediately below y in the inclusion order, *i.e.* there is $e \in |E|$ such that $e \notin x$ and $y = x \cup \{e\}$ – in that case, we also write $x \vdash_E e$ and say that x **enables** e . Observe also that any $x \in \mathcal{C}(E)$ inherits a partial order \leq_x , the restriction of \leq_E to $x \times x$. We usually consider a configuration $x \in \mathcal{C}(E)$ as a partially ordered set.

Event structures are a so-called *truly concurrent* model: rather than presenting observable execution traces, they list computational events along with their causal dependence and independence. The causal order \leq_E is “conjunctive”: for an event to occur, *all* its dependencies must be met first. The conflict relation $\#_E$ represents an irreconcilable non-deterministic choice. Finally, *configurations* provide the adequate notion of *state*.

3.1.2. Symmetry. Plain event structures are not expressive enough for our purposes, notably to handle repetitions in games. Instead, we use event structures *with symmetry*:

Definition 3.3. An **isomorphism family** on event structure E is a set $\mathcal{S}(E)$ of bijections between configurations of E , satisfying the additional conditions:

- groupoid:* $\mathcal{S}(E)$ contains identity bijections; is closed under composition and inverse.
- restriction:* for all $\theta : x \simeq y \in \mathcal{S}(E)$ and $x \supseteq x' \in \mathcal{C}(E)$, there is a (necessarily) unique $\theta \supseteq \theta' \in \mathcal{S}(E)$ such that $\theta' : x' \simeq y'$.
- extension:* for all $\theta : x \simeq y \in \mathcal{S}(E)$, $x \subseteq x' \in \mathcal{C}(E)$, there is a (not necessarily unique) $\theta \subseteq \theta' \in \mathcal{S}(E)$ such that $\theta' : x' \simeq y'$.

The pair $(E, \mathcal{S}(E))$ is called an **event structure with symmetry (ess)**.

We regard isomorphism families as *proof-relevant* equivalence relations: they convey the information of which configurations are interchangeable, witnessed by an explicit bijection.

If E is an ess, we call the elements of $\mathcal{S}(E)$ **symmetries**. It is easy to prove that symmetries are automatically order-isos [Win07]. We write $\theta : x \cong_E y$ to mean that $\theta : x \simeq y$ is a bijection s.t. $\theta \in \mathcal{S}(E)$ with $x = \text{dom}(\theta)$ and $y = \text{cod}(\theta)$. We also write $x \cong_E y$ to mean that there is a symmetry θ s.t. $\theta : x \simeq_E y$. This induces an equivalence relation on configurations – we write $\mathcal{C}_{\cong}(E)$ for the set of equivalence classes, called **symmetry classes**, and use $x, y, z \in \mathcal{C}_{\cong}(E)$ to range over them. Symmetry classes are always non-empty sets of configurations, and the symmetry class of the empty configuration is always a singleton $\{\emptyset\}$. Abusing notation we write $\emptyset \in \mathcal{C}_{\cong}(E)$, which should cause no confusion.

In *thin concurrent games*, both games and strategies are certain ess.

3.2. Games. We introduce our games, and the corresponding constructions.

3.2.1. Definition. First, we recall *thin concurrent games* in the sense of [CCW19]:

Definition 3.4. A **thin concurrent game (tcg)** is an ess $A = (|A|, \leq_A, \#_A)$ with isomorphism families $\mathcal{S}(A), \mathcal{S}_+(A), \mathcal{S}_-(A)$ s.t. $\mathcal{S}_+(A) \subseteq \mathcal{S}(A)$, $\mathcal{S}_-(A) \subseteq \mathcal{S}(A)$, and

$$\text{pol}_A : |A| \rightarrow \{-, +\}$$

a **polarity function** preserved by symmetries, and additionally subject to the conditions:

- orthogonality:* for all $\theta \in \mathcal{S}(A)$, if $\theta \in \mathcal{S}_+(A) \cap \mathcal{S}_-(A)$, then $\theta = \text{id}_x$ for some $x \in \mathcal{C}(A)$,
- receptivity:* if $\theta \in \mathcal{S}_-(A)$ and $\theta \sqsubseteq^- \theta' \in \mathcal{S}(A)$, then $\theta' \in \mathcal{S}_-(A)$,
- +receptivity:* if $\theta \in \mathcal{S}_+(A)$ and $\theta \sqsubseteq^+ \theta' \in \mathcal{S}(A)$, then $\theta' \in \mathcal{S}_+(A)$,

where $\theta \sqsubseteq^p \theta'$ means that $\theta \subseteq \theta'$ adding only (pairs of) events of polarity p .

We shall see examples in Section 3.2.2, accompanying the constructions. Intuitively, negative events correspond to Opponent moves, i.e. actions of the execution environment, while events with positive polarity are Player moves, i.e. actions of the program under study. Symmetries correspond to changing the copy indices arising from $!(-)$. Positive symmetries reindex Player events, while negative symmetries reindex Opponent events.

For us in this paper, a *game* will be a tcg along with a *payoff function*:

Definition 3.5. A **game** is a tcg $A = (|A|, \leq_A, \#_A, \mathcal{S}(A), \mathcal{S}_+(A), \mathcal{S}_-(A), \text{pol}_A)$ with

$$\kappa_A : \mathcal{C}(A) \rightarrow \{-1, 0, +1\}$$



a **payoff function** satisfying the following conditions:

- invariant:* for all $\theta : x \cong_A y$, we have $\kappa_A(x) = \kappa_A(y)$,
- representable:* postponed until Section 4.2.2.

Writing $\min(A)$ for the minimal events of A , a **--game** must additionally satisfy:

- negative:* for all $a \in \min(A)$, $\text{pol}_A(a) = -$,
- initialized:* $\kappa_A(\emptyset) \geq 0$.

Finally, a **--game** A is **strict** if $\kappa_A(\emptyset) = 1$ and all its initial moves are in pairwise conflict. It is **well-opened** if it is strict with exactly one initial move.

Figure 3: The $--$ -game \mathbf{B} Figure 4: The $--$ -game \mathbf{N}

Contexts and types of \mathbf{nPCF} will be represented as strict $--$ -games, but more general games will arise during the model construction.

The condition *representable* is not necessary to get a model of \mathbf{nPCF} but only for the collapse theorem. We shall only introduce and describe it later on, when it becomes relevant. The payoff function κ_A assigns a value to each configuration. Configurations with payoff 0 are called **complete**: they correspond to *terminated* executions, which have reached an adequate stopping point. Otherwise, κ_A assigns a responsibility for why a configuration is non-complete. If $\kappa_A(x) = -1$ then Player is responsible, otherwise it is Opponent.

The payoff structure helps to manage the mismatch between game semantics, which are inherently *affine*, and relational semantics, which are inherently *linear*. Using payoff we will restrict to the strategies that behave linearly; then we can investigate the properties of our collapse at the level of Seely categories. This can also be achieved with other techniques; for example one make the weighted relational model affine by decomposing the comonad $!$ on $\mathcal{R}\text{-Rel}$ as the composition $!_{\text{contr}} \circ !_{\text{weak}}$ of a comonad allowing weakening, and one allowing arbitrary duplication. A similar construction appears in [Mel09, 8.10]. With either approach we obtain a cartesian closed functor between the respective Kleisli categories for $!$.

From *invariant*, all configurations in a symmetry class $x \in \mathcal{C}_{\cong}(A)$ have the same payoff, so we may write $\kappa_A(x)$ unambiguously. We now introduce constructions on games.

3.2.2. *Basic games.* Firstly we draw in Figures 3 and 4 the $--$ -games corresponding to the basic types \mathbb{B} and \mathbb{N} . The diagrams represent the event structures, read from top (the minimum) to bottom (maximal events). Events are annotated with their polarity, and the wiggly line indicates *conflict* – we adopt the convention that we only draw *minimal* conflict, *i.e.* we omit it when it can be deduced via *conflict inheritance*. In Figure 4, all positive events are assumed to be in pairwise conflict (this is not reflected in the diagram for readability). The isomorphism families are not represented, but for these games they are trivial and only comprise identity bijections between configurations. Finally, we have

$$\kappa_{\mathbf{X}}(\emptyset) = +1, \quad \kappa_{\mathbf{X}}(\{\mathbf{q}\}) = -1, \quad \kappa_{\mathbf{X}}(\{\mathbf{q}, v\}) = 0,$$

with $\mathbf{X} \in \{\mathbf{B}, \mathbf{N}\}$. This covers all possible configurations on \mathbf{B} and \mathbf{N} .

Although there are more configurations on \mathbf{B} and \mathbf{N} than points in (\mathbb{B}) and (\mathbb{N}) , the mismatch is resolved when considering *complete* configurations:

Lemma 3.6. *Writing $\mathcal{C}^0(A) = \{x \in \mathcal{C}(A) \mid \kappa_A(x) = 0\}$, we have two bijections:*

$$\begin{array}{ccc} \mathcal{C}^0(\mathbf{B}) & \simeq & (\mathbb{B}) \\ \{\mathbf{q}, b\} & \mapsto & b \end{array} \quad \begin{array}{ccc} \mathcal{C}^0(\mathbf{N}) & \simeq & (\mathbb{N}) \\ \{\mathbf{q}, n\} & \mapsto & n. \end{array}$$

So the web (\mathbb{X}) corresponds to complete configurations of \mathbf{X} . For affine or linear languages with no replication, this correspondence is preserved by all type constructors. But in the presence of replication, one must consider *complete symmetry classes* instead:

Lemma 3.7. *Writing $\mathcal{C}_{\cong}^0(A) = \{x \in \mathcal{C}_{\cong}(A) \mid \kappa_A(x) = 0\}$, we have two bijections:*

$$s^{\mathbb{B}} : \mathcal{C}_{\cong}^0(\mathbf{B}) \simeq (\mathbb{B}) \qquad s^{\mathbb{N}} : \mathcal{C}_{\cong}^0(\mathbf{N}) \simeq (\mathbb{N}).$$

This is not saying much: the symmetries on \mathbf{B} and \mathbf{N} are trivial, so symmetry classes are in one-to-one correspondence with configurations. This will not always be the case.

3.2.3. *Basic constructions on ess.* We start with some constructions on plain ess.

Definition 3.8. Consider E_1 and E_2 two event structures with symmetry.

Then, we define their **parallel composition** $E_1 \parallel E_2$ as comprising the components:

$$\begin{aligned} \text{events:} & & |E_1 \parallel E_2| & = & \{1\} \times |E_1| \uplus \{2\} \times |E_2| \\ \text{causality:} & & (i, e) \leq_{E_1 \parallel E_2} (j, e') & \Leftrightarrow & i = j \ \& \ e \leq_{E_i} e' \\ \text{conflict:} & & (i, e) \#_{E_1 \parallel E_2} (j, e') & \Leftrightarrow & i = j \ \& \ e \#_{E_i} e', \\ \text{symmetry:} & & \theta \in \mathcal{S}(E_1 \parallel E_2) & \Leftrightarrow & \exists \theta_1 \in \mathcal{S}(E_1), \theta_2 \in \mathcal{S}(E_2), \theta = \theta_1 \parallel \theta_2 \end{aligned}$$

where, if $\theta_i : x_i \cong_{E_i} y_i$, we set $(\theta_1 \parallel \theta_2)(i, e) = (i, \theta_i(e))$.

Note that any configuration $x \in \mathcal{C}(E_1 \parallel E_2)$ decomposes uniquely as $(\{1\} \times x_1) \uplus (\{2\} \times x_2)$, which we also write $x_1 \parallel x_2$. This is compatible with symmetry: if $\theta : x_1 \parallel x_2 \cong_{E_1 \parallel E_2} y_1 \parallel y_2$, then θ decomposes uniquely as $\theta_1 \parallel \theta_2$ with $\theta_i : x_i \cong_{E_i} y_i$. We may also observe that any symmetry class $x \in \mathcal{C}_{\cong}(E_1 \parallel E_2)$ has the form $x_1 \parallel x_2$, which is shorthand for $\{x_1 \parallel x_2 \mid x_1 \in x_1, x_2 \in x_2\}$ for $x_1 \in \mathcal{C}_{\cong}(E_1)$ and $x_2 \in \mathcal{C}_{\cong}(E_2)$.

We also use a variant of the above where components are in conflict:

Definition 3.9. Let E_1 and E_2 be two event structures with symmetry.

Then, we define their **sum** $E_1 + E_2$ as comprising the components:

$$\begin{aligned} \text{events:} & & |E_1 + E_2| & = & \{1\} \times |E_1| \uplus \{2\} \times |E_2| \\ \text{causality:} & & (i, e) \leq_{E_1 + E_2} (j, e') & \Leftrightarrow & i = j \ \& \ e \leq_{E_i} e' \\ \text{conflict:} & & (i, e) \#_{E_1 + E_2} (j, e') & \Leftrightarrow & i \neq j \ \vee \ e \#_{E_i} e', \\ \text{symmetry:} & & \theta \in \mathcal{S}(E_1 + E_2) & \Leftrightarrow & \exists \theta_1 \in \mathcal{S}(E_1), \theta_2 \in \mathcal{S}(E_2), \theta = \theta_1 \parallel \theta_2, \end{aligned}$$

where, necessarily, one of θ_1 or θ_2 must be empty.

Any non-empty $x \in \mathcal{C}(E_1 + E_2)$ may be written uniquely either as $\{1\} \times x_1$ for $x_1 \in \mathcal{C}(E_1)$, or as $\{2\} \times x_2$ for $x_2 \in \mathcal{C}(E_2)$. By convention, we write $(1, x_1) \in \mathcal{C}(E_1 + E_2)$ for the former and $(2, x_2) \in \mathcal{C}(E_1 + E_2)$ for the latter. Likewise, non-empty symmetries on $E_1 + E_2$ may be written uniquely either as $(1, \theta_1)$ or as $(2, \theta_2)$ for $\theta_i \in \mathcal{S}(E_i)$, defined in the obvious way. For $x_1 \in \mathcal{C}_{\cong}(E_1)$, we also write $(1, x_1) \in \mathcal{C}_{\cong}(E_1 \& E_2)$ as a shorthand for $\{(1, x_1) \mid x_1 \in x_1\}$ and likewise for $(2, x_2) \in \mathcal{C}_{\cong}(E_1 \& E_2)$ for $x_2 \in \mathcal{C}_{\cong}(E_2)$. Every non-empty symmetry class on $E_1 \& E_2$ may be written uniquely via one of these two shapes.

3.2.4. *Basic constructions on games.* Our first construction is the **dual** A^\perp of a game A . We set A^\perp as the same ess as A , changing only the components relative to polarities: more precisely, $\text{pol}_{A^\perp} = -\text{pol}_A$, $\mathcal{S}_+(A^\perp) = \mathcal{S}_-(A)$, $\mathcal{S}_-(A^\perp) = \mathcal{S}_+(A)$, and $\kappa_{A^\perp} = -\kappa_A$.

Parallel composition splits into *tensor* and *par*, which differ only in their payoff function:

\otimes	-1	0	+1
-1	-1	-1	-1
0	-1	0	+1
+1	-1	+1	+1

\wp	-1	0	+1
-1	-1	-1	+1
0	-1	0	+1
+1	+1	+1	+1

Figure 5: Payoff for \otimes and \wp

Definition 3.10. Consider two games A and B .

We define their **tensor** $A \otimes B$ as having ess $A \parallel B$, with the additional components:

$$\begin{aligned}
\text{polarities:} \quad & \text{pol}_{A \otimes B}(1, a) = \text{pol}_A(a) \\
& \text{pol}_{A \otimes B}(2, b) = \text{pol}_B(b) \\
\text{positive symmetries:} \quad & \theta_A \parallel \theta_B \in \mathcal{S}_+(A \otimes B) \Leftrightarrow \theta_A \in \mathcal{S}_+(A) \ \& \ \theta_B \in \mathcal{S}_+(B) \\
\text{negative symmetries:} \quad & \theta_A \parallel \theta_B \in \mathcal{S}_-(A \otimes B) \Leftrightarrow \theta_A \in \mathcal{S}_-(A) \ \& \ \theta_B \in \mathcal{S}_-(B) \\
\text{payoff:} \quad & \kappa_{A \otimes B}(x_A \parallel x_B) = \kappa_A(x_A) \otimes \kappa_B(x_B)
\end{aligned}$$

where the binary operation \otimes on $\{-1, 0, +1\}$ is defined in Figure 5. We also define the **par** $A \wp B$ with the same components, except for $\kappa_{A \wp B}(x_A \parallel x_B) = \kappa_A(x_A) \wp \kappa_B(x_B)$.

The tensor of two $--$ -games is again a $--$ -game. The par also preserves $--$ -games, but we will often use it on more general games: for $--$ -games A and B we will eventually define a strategy *from A to B* as a strategy on $A^\perp \wp B$. This is a game but not a $--$ -game.

Analogously to Lemma 3.7, we have:

Lemma 3.11. Consider A and B any games. Then, we have

$$\begin{aligned}
r_{A,B}^\otimes : \mathcal{C}(A \otimes B) &\simeq \mathcal{C}(A) \times \mathcal{C}(B) & s_{A,B}^\otimes : \mathcal{C}_{\leq}^0(A \otimes B) &\simeq \mathcal{C}_{\leq}^0(A) \times \mathcal{C}_{\leq}^0(B) \\
x_A \parallel x_B &\mapsto (x_A, x_B) & x_A \parallel x_B &\mapsto (x_A, x_B)
\end{aligned}$$

and likewise, $s_{A,B}^\wp : \mathcal{C}_{\leq}^0(A \wp B) \simeq \mathcal{C}_{\leq}^0(A) \times \mathcal{C}_{\leq}^0(B)$ with the same function.

The proof is straightforward, and uses that in Figure 5, for either \otimes and \wp , a configuration $x_A \parallel x_B$ has null payoff iff it has null payoff on both sides. This connects the tensor product of games with that in the relational model, which is defined as the cartesian product of sets. We move to another construction on games.

Definition 3.12. For two strict $--$ -games A_1 and A_2 , we define their **with** $A_1 \& A_2$ as having ess $A_1 + A_2$, with the additional components:

$$\begin{aligned}
\text{polarities:} \quad & \text{pol}_{A \& B}(1, a) = \text{pol}_A(a) \\
& \text{pol}_{A \& B}(2, b) = \text{pol}_B(b) \\
\text{positive symmetries:} \quad & (i, \theta) \in \mathcal{S}_+(A_1 \& A_2) \Leftrightarrow \theta \in \mathcal{S}_+(A_i) \\
\text{negative symmetries:} \quad & (i, \theta) \in \mathcal{S}_-(A_1 \& A_2) \Leftrightarrow \theta \in \mathcal{S}_-(A_i), \\
\text{payoff:} \quad & \kappa_{A_1 \& A_2}((i, x)) = \kappa_{A_i}(x), \quad (x \neq \emptyset) \\
& \kappa_{A_1 \& A_2}(\emptyset) = 1,
\end{aligned}$$

yielding a strict $--$ -game.

As we will see, this construction gives a cartesian product in our forthcoming category of strategies. It can also be applied to non-strict $--$ -games, but then it is not a product: if one of the A_i is not strict then the corresponding projection does not respect payoff (in the sense of Definition 3.25), because we have set $\kappa_{A_1 \& A_2}(\emptyset) = 1$. On the other hand

having $\kappa_{A_1 \& A_2}(\emptyset) = 0$ breaks the correspondence with the relational model, since the empty configuration does not correspond in a canonical way to one of the components.

On complete symmetry classes this matches the corresponding construction in \mathcal{R} -Rel:

Lemma 3.13. *Consider A and B any strict $--$ games. Then, we have*

$$\begin{aligned} r_{A,B}^{\&} &: \mathcal{C}^{\neq \emptyset}(A \& B) \simeq \mathcal{C}^{\neq \emptyset}(A) + \mathcal{C}^{\neq \emptyset}(B) \\ s_{A,B}^{\&} &: \mathcal{C}_{\simeq}^0(A \& B) \simeq \mathcal{C}_{\simeq}^0(A) + \mathcal{C}_{\simeq}^0(B). \end{aligned}$$

This generalizes directly to the n -ary case. The bijection follows our notation (i, x) for symmetry classes of non-empty configurations. As discussed above, the bijection $s_{A,B}^{\&}$ relies on strictness, which ensures that configurations with null payoff cannot be empty.

3.2.5. *Arrow.* Next we give the construction of a *linear* function space. The event structure $A \multimap B$ is easier to describe when B is well-opened, so we consider this case first.

Definition 3.14. Consider two games A and B , where B is well-opened and has unique initial move b_0 . The **linear function space** $A \multimap B$ has the following components:

$$\begin{aligned} \text{events:} & \quad |A \multimap B| = |A^\perp \parallel B| \\ \text{causality:} & \quad \leq_{A \multimap B} = \leq_{A^\perp \parallel B} \uplus \{((2, b_0), (1, a)) \mid a \in |A|\} \\ \text{conflict:} & \quad \#_{A \multimap B} = \#_{A^\perp \parallel B} \\ \text{symmetries:} & \quad \mathcal{S}(A \multimap B) = \{\theta : x \cong_{A^\perp \parallel B} y \mid x, y \in \mathcal{C}(A \multimap B)\} \\ \text{polarities:} & \quad \text{pol}_{A \multimap B} = \text{pol}_{A^\perp \parallel B} \\ \text{positive symmetries:} & \quad \mathcal{S}_+(A \multimap B) = \{\theta : x \cong_{A^\perp \parallel B}^+ y \mid x, y \in \mathcal{C}(A \multimap B)\} \\ \text{negative symmetries:} & \quad \mathcal{S}_-(A \multimap B) = \{\theta : x \cong_{A^\perp \parallel B}^- y \mid x, y \in \mathcal{C}(A \multimap B)\} \\ \text{payoff:} & \quad \kappa_{A \multimap B}(x_A \parallel x_B) = \kappa_{A^\perp}(x_A) \wp \kappa_B(x_B) \quad (x_B \neq \emptyset) \\ & \quad \kappa_{A \multimap B}(\emptyset) = 1. \end{aligned}$$

This is a well-opened $--$ game.

Again, symmetry classes of $A \multimap B$ relate with the corresponding construction in \mathcal{R} -Rel:

Lemma 3.15. *Consider two games A and B with B well-opened. Then, we have*

$$\begin{aligned} r_{A,B}^{\multimap} &: \mathcal{C}^{\neq \emptyset}(A \multimap B) \simeq \mathcal{C}(A) \times \mathcal{C}^{\neq \emptyset}(B) \\ s_{A,B}^{\multimap} &: \mathcal{C}_{\simeq}^0(A \multimap B) \simeq \mathcal{C}_{\simeq}^0(A) \times \mathcal{C}_{\simeq}^0(B) \end{aligned}$$

Again, the bijection $s_{A,B}^{\multimap}$ relies on B being *strict*: if we had $\kappa_B(\emptyset) = 0$, then there would be no symmetry class on $A \multimap B$ corresponding to a pair (x_A, \emptyset) with x_A non-empty.

It is easy to extend the construction of $A \multimap B$ to the case where B is *strict* but not well-opened: in that case, B has a canonical form $B \cong \&_{b \in \min(B)} B_b$ with B_b well-opened, for the obvious notion of isomorphism between games. This lets us set:

$$A \multimap B = \&_{b \in \min(B)} A \multimap B_b,$$

with in particular $A \multimap \top = \top$ for \top the empty game with $\kappa_\top(\emptyset) = 1$. We retain:

Lemma 3.16. *Consider two games A and B with B strict. Then, we have bijections*

$$\begin{aligned} r_{A,B}^{\multimap} &: \mathcal{C}^{\neq \emptyset}(A \multimap B) \simeq \mathcal{C}(A) \times \mathcal{C}^{\neq \emptyset}(B) \\ s_{A,B}^{\multimap} &: \mathcal{C}_{\simeq}^0(A \multimap B) \simeq \mathcal{C}_{\simeq}^0(A) \times \mathcal{C}_{\simeq}^0(B) \end{aligned}$$

Proof. Obtained by composition

$$\begin{aligned}
\mathcal{C}^{\neq\emptyset}(A \multimap B) &\simeq \sum_{b \in \min(B)} \mathcal{C}^{\neq\emptyset}(A \multimap B_b) \\
&\simeq \sum_{b \in \min(B)} \mathcal{C}(A) \times \mathcal{C}^{\neq\emptyset}(B_b) \\
&\simeq \mathcal{C}(A) \times \left(\sum_{b \in \min(B)} \mathcal{C}^{\neq\emptyset}(B_b) \right) \\
&\simeq \mathcal{C}(A) \times \mathcal{C}^{\neq\emptyset}(B)
\end{aligned}$$

using Lemmas 3.13 and 3.15, together with the distributivity of cartesian product over disjoint union. The exact same reasoning applies to complete symmetry classes. \square

For $x_A \in \mathcal{C}(A)$ and $x_B \in \mathcal{C}^{\neq\emptyset}(B)$, we write $x_A \multimap x_B = (r_{A,B}^-)^{-1}(x_A, x_B) \in \mathcal{C}^{\neq\emptyset}(A \multimap B)$, and likewise $\times_A \multimap \times_B \in \mathcal{C}_{\cong}^0(A \multimap B)$ for $\times_A \in \mathcal{C}_{\cong}^0(A)$ and $\times_B \in \mathcal{C}_{\cong}^0(B)$.

3.2.6. Exponentials. We define $!(-)$, which will eventually extend to a linear exponential comonad. It is the only source of non-trivial symmetries in the games used for **nPCF**. For a $--$ -game A , $!A$ is understood as an infinitary tensor of symmetric copies of A . The point is to allow for duplication and weakening, and so in particular $!A$ is *not* strict.

Definition 3.17. Consider A a $--$ -game. Then, we define the **bang** $!A$ with the components:

$$\begin{array}{ll}
\text{events:} & |!A| = \mathbb{N} \times |A| \\
\text{causality:} & (i, a_1) \leq_{!A} (j, a_2) \Leftrightarrow i = j \wedge a_1 \leq_A a_2 \\
\text{conflict:} & (i, a_1) \#_{!A} (j, a_2) \Leftrightarrow i = j \wedge a_1 \#_A a_2 \\
\text{symmetries:} & \theta \in \mathcal{S}(!A) \Leftrightarrow \exists \pi : \mathbb{N} \simeq \mathbb{N}, \exists (\theta_n)_{n \in \mathbb{N}} \in \mathcal{S}(A)^{\mathbb{N}} \\
& \forall (i, a) \in \text{dom}(\theta), \theta(i, a) = (\pi(i), \theta_i(a)) \\
\text{polarities:} & \text{pol}_{!A}(i, a) = \text{pol}_A(a) \\
\text{positive symmetries:} & \theta \in \mathcal{S}_+(!A) \Leftrightarrow \exists (\theta_n)_{n \in \mathbb{N}} \in \mathcal{S}_+(A)^{\mathbb{N}}, \\
& \forall (i, a) \in \text{dom}(\theta), \theta(i, a) = (i, \theta_i(a)) \\
\text{negative symmetries:} & \theta \in \mathcal{S}_-(!A) \Leftrightarrow \exists \pi : \mathbb{N} \simeq \mathbb{N}, \exists (\theta_n)_{n \in \mathbb{N}} \in \mathcal{S}_-(A)^{\mathbb{N}}, \\
& \forall (i, a) \in \text{dom}(\theta), \theta(i, a) = (\pi(i), \theta_i(a)) \\
\text{payoff:} & \kappa_{!A}(\|_{i \in I} x_i) = \bigotimes_{i \in I} \kappa_A(x_i) \quad (I \subseteq \mathbb{N}, \forall i \in I, x_i \neq \emptyset) \\
& \kappa_{!A}(\emptyset) = 0
\end{array}$$

where $\|_{i \in I} x_i = \biguplus_{i \in I} \{i\} \times x_i$. This yields a $--$ -game $!A$.

The definition of payoff uses implicitly that the tensor operation on $\{-1, 0, +1\}$ defined in Figure 5 is associative. The $--$ -game $!A$ is non-strict by intention: Opponent is free to open any number of copies, including zero.

Again, by considering symmetry classes we recover the matching construction in $\mathcal{R}\text{-Rel}$.

Lemma 3.18. Consider A a $--$ -game. Then, we have bijections

$$s_A^! : \mathcal{C}_{\cong}^0(!A) \simeq \mathcal{M}_f(\mathcal{C}_{\cong}^{0, \neq\emptyset}(A)) \quad s_A^{!, \neq\emptyset} : \mathcal{C}_{\cong}^{0, \neq\emptyset}(!A) \simeq \mathcal{M}_f^{\neq\emptyset}(\mathcal{C}_{\cong}^{0, \neq\emptyset}(A))$$

with $\mathcal{C}_{\cong}^{0, \neq\emptyset}(A)$ the non-empty complete classes, and $\mathcal{M}_f^{\neq\emptyset}(X)$ the non-empty finite multisets.

In particular, if A is a strict $--$ -game, the first bijection specializes to

$$s_A^! : \mathcal{C}_{\cong}^0(!A) \simeq \mathcal{M}_f(\mathcal{C}_{\cong}^0(A)).$$

Proof. We prove the first bijection. We first define $s_A^!$ on concrete configurations of $!A$:

$$\begin{aligned} s_A^! &: \mathcal{C}^0(!A) \rightarrow \mathcal{M}_f(\mathcal{C}_{\cong}^{0, \neq \emptyset}(A)) \\ \emptyset &\mapsto \square \\ \parallel_{i \in I} x_i &\mapsto [x_i \mid i \in I] \quad (\forall i \in I, x_i \neq \emptyset) \end{aligned}$$

where x_i denotes the symmetry class of x_i . It is straightforward that if $x \cong_{!A} y$ then $s_A^!(x) = s_A^!(y)$, so $s_A^!$ lifts to $s_A^! : \mathcal{C}_{\cong}^0(!A) \rightarrow \mathcal{M}_f(\mathcal{C}_{\cong}^{0, \neq \emptyset}(A))$, keeping the same notation.

Injective. Assume $s_A^!(\parallel_{i \in I} x_i) = s_A^!(\parallel_{j \in J} y_j)$ with every x_i and y_j non-empty. We have

$$[x_i \mid i \in I] = [y_j \mid j \in J],$$

meaning that there is a bijection $\pi : I \simeq J$ such that for all $i \in I$, we have $x_i = y_{\pi(i)}$. In turn, this means that for all $i \in I$, there is some $\theta_i : x_i \cong_A y_{\pi(i)}$. Now, completing π to $\pi' : \mathbb{N} \simeq \mathbb{N}$ arbitrarily yields $\theta : \parallel_{i \in I} x_i \cong_{!A} \parallel_{j \in J} y_j$ as required.

Surjective. Consider $\mu \in \mathcal{M}_f(\mathcal{C}_{\cong}^{0, \neq \emptyset}(!A))$. If $\mu = \square$, its pre-image is the empty complete symmetry class. Otherwise, write $\mu = [x_i \mid i \in I]$ with each x_i non-empty. For $i \in I$, fix some $x_i \in x_i$. *W.l.o.g.* we may assume $I \subseteq_f \mathbb{N}$, so that $\parallel_{i \in I} x_i$ gives the required pre-image.

Clearly, $s_A^!$ restricts to $s_A^{!, \neq \emptyset} : \mathcal{C}_{\cong}^{0, \neq \emptyset}(!A) \simeq \mathcal{M}_f^{\neq \emptyset}(\mathcal{C}_{\cong}^{0, \neq \emptyset}(A))$. \square

Note that unless A is strict, not all finite multisets in $\mathcal{M}_f(\mathcal{C}_{\cong}^0(A))$ correspond to a complete symmetry class on $!A$. For instance $[\emptyset]$, or any $[\emptyset, \dots, \emptyset]$ do not.

3.2.7. Interpretation of types and arenas. We give the complete interpretation of types. *Types* are interpreted as well-opened $--$ -games: for base types we use the well-opened games \mathbf{B} and \mathbf{N} defined in Section 3.2.2, and we set $\llbracket A \rightarrow B \rrbracket = !\llbracket A \rrbracket \multimap \llbracket B \rrbracket$. *Contexts* are interpreted as strict $--$ -games, with $\llbracket x_1 : A_1, \dots, x_n : A_n \rrbracket = \&_{1 \leq i \leq n} \llbracket A_i \rrbracket$.

Putting together Lemmas 3.7, 3.13, 3.15 and 3.18, we immediately get:

Lemma 3.19. *For any type A and context Γ , there are bijections:*

$$s_A^{\text{ty}} : \langle A \rangle \simeq \mathcal{C}_{\cong}^0(\llbracket A \rrbracket) \quad s_{\Gamma}^{\text{ctx}} : \mathcal{M}_f(\langle \Gamma \rangle) \simeq \mathcal{C}_{\cong}^0(!\llbracket \Gamma \rrbracket).$$

So the web of A may be regarded as the set of complete symmetry classes of $\llbracket A \rrbracket$. The games obtained as the interpretation of types have a particular shape:

Definition 3.20. An **arena** is a $--$ -game A satisfying:

$$\begin{aligned} \textit{alternating:} & \text{ if } a_1 \rightarrow_A a_2, \text{ pol}_A(a_1) \neq \text{pol}_A(a_2), \\ \textit{forestial:} & \text{ if } a_1 \leq_A a \text{ and } a_2 \leq_A a, \text{ then } a_1 \leq_A a_2 \text{ or } a_2 \leq_A a_1. \end{aligned}$$

For any type A and context Γ , it is straightforward that the $--$ -games $\llbracket A \rrbracket$ and $\llbracket \Gamma \rrbracket$ are arenas. Arenas do not quite characterize the games arising from the interpretation, but get close enough for the purposes of this paper. More precisely, arenas guarantee that moves have at most one causal immediate predecessor, hence recovering the notion of *justifier* familiar from more traditional game semantics [CC21]. This will play a very minor role in this paper: it is necessary for the deadlock-freeness property used in Section 4.1.1.

3.3. Strategies. Now that games are set, we introduce *strategies* used to interpret terms.

3.3.1. *Plain strategies.* We start by recalling the notion of *causal strategy* in its formulation in [CC21] – here, we only say *strategy* as it is the only sort of strategy we consider.

Definition 3.21. A **prestrategy** on game A comprises an ess $(|\sigma|, \leq_\sigma, \#_\sigma, \mathcal{S}(\sigma))$ with

$$\partial : |\sigma| \rightarrow |A|$$

a function called the **display map**, subject to the following conditions:

- rule-abiding:* for all $x \in \mathcal{C}(\sigma)$, $\partial(x) \in \mathcal{C}(A)$,
- locally injective:* for all $s_1, s_2 \in x \in \mathcal{C}(\sigma)$, if $\partial(s_1) = \partial(s_2)$ then $s_1 = s_2$,
- symmetry-preserving:* for all $\theta \in \mathcal{S}(\sigma)$, $\partial(\theta) = \{(\partial(s_1), \partial(s_2)) \mid (s_1, s_2) \in \theta\} \in \mathcal{S}(A)$,
- \sim -receptive:* for all $\theta : x \cong_\sigma y$, and extensions $x \vdash_\sigma s_1^-$, $\partial(\theta) \vdash_{\mathcal{S}(A)} (\partial(s_1^-), a_2^-)$, there is a unique $s_2^- \in |\sigma|$ s.t. $\theta \vdash_{\mathcal{S}(\sigma)} (s_1^-, s_2^-)$ and $\partial(s_2^-) = a_2^-$,
- thin:* for all $\theta : x \cong_\sigma y$, and extension $x \vdash_\sigma s_1^+$, there is a *unique* extension $y \vdash_\sigma s_2^+$ such that $\theta \vdash_{\mathcal{S}(\sigma)} (s_1^+, s_2^+)$.

Additionally, we say that σ is a **strategy** if it satisfies the further three conditions:

- negative:* for all $s \in |\sigma|$, if s is minimal then s is negative,
- courteous:* for all $s_1 \rightarrow_\sigma s_2$, if $\text{pol}(s_1) = +$ or $\text{pol}(s_2) = -$ then $\partial(s_1) \rightarrow_A \partial(s_2)$,
- receptive:* for all $x \in \mathcal{C}(\sigma)$, for all $\partial(x) \vdash_A a^-$, there is a unique $x \vdash_\sigma s^- \in \mathcal{C}(\sigma)$ such that $\partial(s) = a$,


We write $\sigma : A$ to mean that σ is a strategy on game A .

We disambiguate some notations used in the definition. First, σ implicitly comes with polarities, imported from A as $\text{pol}_\sigma(s) = \text{pol}_A(\partial(s))$. We often tag events to indicate their polarity as in s^-, s^+ ; the sign is not considered part of the variable name but conveys the polarity information. We also used the enabling relation on isomorphism families, defined by $\theta \vdash_{\mathcal{S}(A)} (a_1, a_2)$ iff $(a_1, a_2) \notin \theta$ and $\theta \cup \{(a_1, a_2)\} \in \mathcal{S}(A)$.

A strategy is a causal presentation, in one global object, of the entire computational behaviour of a program on the interface with its execution environment. While the *events* or *moves* in $|\sigma|$ are not technically moves of the game, the *display map* ∂_σ associates to any move of σ the corresponding move in the game, and is subject to adequate conditions ensuring compatibility with the structure. More precisely, conditions *rule-abiding*, *locally injective* and *symmetry-preserving* together amount to ∂ being a **map of event structures with symmetry** [Win07] – the adequate simulation maps between ess.

Conditions *courteous* and *receptive* are the usual conditions for concurrent strategies [RW11, CCRW17], expressing that the strategy should be invariant under asynchronous delay. The condition *\sim -receptive* forces strategies to consider as symmetric pairs of Opponent events symmetric in the game, and hence to treat them uniformly. Finally, *thin* is a minimality condition forcing strategies to pick *one* canonical representative up to symmetry for positive moves. For further explanations and discussions on those conditions, see [CCW19].

As an illustration, we show a strategy in Figure 6. We take advantage of this to introduce our convention for drawing strategies. Ignoring dotted lines, the diagram represents the event structure $(|\sigma|, \leq_\sigma, \#_\sigma)$, via its immediate causality relation \rightarrow . For convenience, each node is labeled with the corresponding move in the game as observed through ∂_σ – as each move in the game originates in one of the ground types appearing in the type, we attempt as much as possible to place moves in the diagram in the corresponding column. The immediate causality in the game is conveyed through the dotted lines. The grey subscripts correspond to the *copy indices*, *i.e.* the tags introduced in Definition 3.17 to address the

Figure 6: A strategy $\sigma : !(B \multimap B) \multimap B$

components of $!A$. For moves under several $!$, we write the copy indices in a sequence ranging from the outermost to the innermost $!$. The representation is symbolic: the actual strategy comprises branches as shown for all $i, j, k \in \mathbb{N}$. Finally, the one component of σ missing in this representation is the isomorphism family $\mathcal{S}(\sigma)$ which is too unwieldy to draw – in this particular case it consists of all order-isomorphisms changing only copy indices.

Only a few strategies are definable via **nPCF**: for instance, in [CC21] the same strategies are used to model a higher-order concurrent language with shared memory. One could, with an adequate notion of innocence, characterize exactly those strategies definable through **nPCF** – this is done in [Cas17]. For our present purposes this is not required; nevertheless we have to add two further conditions for the collapse functor to exist.

3.3.2. *Visible strategies.* *Visibility* is a locality property for the control flow. First, we define:

Definition 3.22. Consider E an event structure.

A **grounded causal chain (gcc)** in E is $\rho = \{\rho_1, \dots, \rho_n\} \subseteq |E|$ forming

$$\rho_1 \rightarrow_E \dots \rightarrow_E \rho_n$$

a chain with ρ_1 minimal with respect to \leq_E . We write $\text{gcc}(E)$ for the gccs in E .

If σ is forestial, then $\text{gcc}(\sigma)$ comprises simply its *finite branches*. But in general, a gcc $\rho \in \text{gcc}(\sigma)$ is not down-closed; it may be seen as an individual *thread*, and σ may be regarded as a collection of such threads with the information of their non-deterministic branchings, as well as the points where they causally fork or merge. Visibility states that each thread respects the local scope, *i.e.* it may not use resources introduced in another thread:

Definition 3.23. A strategy $\sigma : A$ is **visible** if for all $\rho \in \text{gcc}(\sigma)$, $\partial_\sigma(\rho) \in \mathcal{C}(A)$.

In this paper, we require visibility as it restricts the behaviour of strategies just enough so that their interactions never deadlock. Visibility is far from sufficient in the way of capturing the behaviour of **nPCF** terms – for those the causal structure is always forest-shaped, which

trivially entails visibility. We work with visibility rather than a forest-shaped causality as it costs us nothing, and makes our collapse theorem slightly more general.

Much more information on visibility may be found in [CC21].

3.3.3. Exhaustive strategies. *Exhaustivity* is an elegant mechanism due to Mellès [Mel05]. As discussed previously, it ensures that strategies are *linear* rather than *affine*. It also enforces a form of well-bracketing, forcing strategies to respect the call stack discipline.

First, we must define a notion of *stopping state* for a strategy.

Definition 3.24. Consider A a game, and $\sigma : A$ a strategy. A configuration $x \in \mathcal{C}(\sigma)$ is **+covered** if for all $m \in x$, if m is maximal in x (with respect to \leq_σ), then $\text{pol}_\sigma(m) = +$.

We write $\mathcal{C}^+(\sigma)$ for the set of all +covered configurations of σ .

In other words, a configuration is +covered if no Opponent move is left unresponded.

Definition 3.25. Consider A a game, and $\sigma : A$ a strategy. We set the condition:

$$\textit{exhaustive: } \text{ for all } x \in \mathcal{C}^+(\sigma), \kappa_A(\partial_\sigma(x)) \geq 0.$$

For example, $\sigma : \mathbf{B}_1 \multimap \mathbf{B}_2$ (indices for disambiguation) *cannot* reply on \mathbf{B}_2 without evaluating its argument, as $\kappa_{\mathbf{B}_1 \multimap \mathbf{B}_2}(\{\mathbf{q}_2^-, b_2^+\}) = \kappa_{\mathbf{B}^\perp}(\emptyset) \wp \kappa_{\mathbf{B}}(\{\mathbf{q}^-, b^+\}) = -1 \wp 0 = -1$.

In this paper we include all details of the compositionality of exhaustivity, which to the best of our knowledge do not appear in any published source.

3.4. Composition of plain strategies. Postponing for now the stability under composition of visibility and exhaustivity, we recall the composition of plain strategies.

For games A and B , a **strategy from A to B** is a strategy on $A^\perp \wp B$, also written $A \vdash B$. Fix from now on A, B and C , and $\sigma : A \vdash B$, $\tau : B \vdash C$. We aim to define $\tau \odot \sigma : A \vdash C$. The concrete definition of composition is covered at length elsewhere [CCW19]; instead we give a characterization in terms of its *states*.

3.4.1. Synchronization. Given configurations $x^\sigma \in \mathcal{C}(\sigma)$, $x^\tau \in \mathcal{C}(\tau)$, by convention we write

$$\partial_\sigma(x^\sigma) = x_A^\sigma \parallel x_B^\sigma \in \mathcal{C}(A \vdash B), \quad \partial_\tau(x^\tau) = x_B^\tau \parallel x_C^\tau \in \mathcal{C}(B \vdash C),$$

for the corresponding projections to the game. In defining composition, the first stage is to capture when such configurations $x^\sigma \in \mathcal{C}(\sigma)$ and $x^\tau \in \mathcal{C}(\tau)$ may successfully *synchronise*.

Definition 3.26. Consider two configurations $x^\sigma \in \mathcal{C}(\sigma)$ and $x^\tau \in \mathcal{C}(\tau)$. They are **causally compatible** if (1) **matching**: $x_B^\sigma = x_B^\tau = x_B$; and (2) if the bijection

$$\varphi_{x^\sigma, x^\tau} : x^\sigma \parallel x_C^\tau \xrightarrow{\partial_\sigma \parallel x_C^\tau} x_A^\sigma \parallel x_B \parallel x_C^\tau \xrightarrow{x_A^\sigma \parallel \partial_\tau^{-1}} x_A^\sigma \parallel x^\tau,$$

obtained by composition using local injectivity of ∂_σ and ∂_τ , is **secured**, *i.e.* the relation

$$(m, n) \triangleleft (m', n') \iff m <_{\sigma \parallel C} m' \vee n <_{A \parallel \tau} n',$$

defined on (the graph of) $\varphi_{x^\sigma, x^\tau}$ by importing causal constraints of σ and τ , is acyclic.

Securedness eliminates deadlocks: two matching $x^\sigma \in \mathcal{C}(\sigma)$, $x^\tau \in \mathcal{C}(\tau)$ agree on the final state in B , but σ and τ may impose incompatible constraints making it unreachable.

3.4.2. *Plain composition.* It turns out that up to an adequate isomorphism, there is a *unique* strategy $\tau \odot \sigma : A \vdash C$ whose configurations are such synchronizations, in a way compatible with symmetry. In order to state this we first disambiguate our notion of isomorphism.

Definition 3.27. Consider A a game, and $\sigma, \tau : A$ two strategies.

An **isomorphism** $\varphi : \sigma \cong \tau$ is an invertible map of ess such that $\partial_\tau \circ \varphi = \partial_\sigma$.

Before we introduce composition, we extend two earlier notions from configurations to symmetries. A symmetry $\theta \in \mathcal{S}(\sigma)$ is **+covered** if either of $\text{dom}(\theta)$ or $\text{cod}(\theta)$ is: as symmetries are order-isomorphisms and preserve polarities, the difference is immaterial. We write $\mathcal{S}^+(\sigma)$ for the set of +covered symmetries of σ . Likewise, $\theta^\sigma \in \mathcal{S}(\sigma)$ and $\theta^\tau \in \mathcal{S}(\tau)$ are **causally compatible** if they are *matching*, i.e. $\partial_\sigma \theta^\sigma = \theta_A^\sigma \parallel \theta_B^\sigma$ and $\partial_\tau \theta^\tau = \theta_B^\tau \parallel \theta_C^\tau$ with $\theta_B^\sigma = \theta_B^\tau$; and *secured*, i.e. either $\text{dom}(\theta^\sigma), \text{dom}(\theta^\tau)$ or $\text{cod}(\theta^\sigma), \text{cod}(\theta^\tau)$ are.

Now we state the following proposition, whose proof may be found in [CC21]:

Proposition 3.28. Consider $\sigma : A \vdash B$ and $\tau : B \vdash C$ two strategies.

Then, there is a strategy $\tau \odot \sigma : A \vdash C$, unique up to iso, such that there are order-isos:

$$\begin{aligned} (- \odot -) & : \{(x^\tau, x^\sigma) \in \mathcal{C}^+(\tau) \times \mathcal{C}^+(\sigma) \mid x^\sigma \text{ and } x^\tau \text{ causally compatible}\} \simeq \mathcal{C}^+(\tau \odot \sigma) \\ (- \odot -) & : \{(\theta^\tau, \theta^\sigma) \in \mathcal{S}^+(\tau) \times \mathcal{S}^+(\sigma) \mid \theta^\sigma \text{ and } \theta^\tau \text{ causally compatible}\} \simeq \mathcal{S}^+(\tau \odot \sigma) \end{aligned}$$

commuting with dom, cod , and s.t. for $\theta^\sigma \in \mathcal{S}^+(\sigma)$ and $\theta^\tau \in \mathcal{S}^+(\tau)$ causally compatible,

$$\partial_{\tau \odot \sigma}(\theta^\tau \odot \theta^\sigma) = \theta_A^\sigma \parallel \theta_C^\tau.$$

So $\tau \odot \sigma$ is the unique (up to iso) strategy whose configurations correspond to matching pairs for which the causal constraints imposed by σ and τ are compatible.

The idea of considering matching pairs of configurations brings us close to the composition of relations, spans, or profunctors. In general, however, the composition of strategies is more restricted, because of *causal compatibility*. This is a well-known feature of game semantics, which allows for the interpretation of more sophisticated language constructs. But for the strategies of this paper, matching configurations are always causally compatible. This follows from *visibility*, which we will exploit to prove that composition never *deadlocks*; see Section 4.1.1. This is crucial for constructing a functor to the relational model.

3.4.3. *Copycat.* As we have defined a notion of composition, it is natural to introduce here the accompanying identity, the *copycat strategy*. For any game A , the copycat strategy $\mathfrak{c}_A : A \vdash A$ is an asynchronous forwarder: any Opponent move on either side enables (i.e. is the unique causal dependency for) the corresponding event on the other side.

We only give the definition of copycat on arenas, as it is sufficient and slightly simpler:

Definition 3.29. For each arena A , the **copycat strategy** $\mathfrak{c}_A : A \vdash A$ comprises:

$$\begin{aligned} |\mathfrak{c}_A| & = |A \vdash A| \\ \partial_{\mathfrak{c}_A}(i, a) & = (i, a) \\ (i, a) \leq_{\mathfrak{c}_A} (j, a') & \Leftrightarrow a <_A a'; \text{ or } a = a', \text{ pol}_{A \vdash A}(i, a) = - \text{ and } \text{pol}_{A \vdash A}(j, a') = + \\ (i, a) \#_{\mathfrak{c}_A} (j, a') & \Leftrightarrow a \#_A a', \end{aligned}$$

with symmetries those bijections of the form $\theta_1 \parallel \theta_2 : x_1 \parallel x_2 \cong_{\mathfrak{c}_A} y_1 \parallel y_2$ such that

$$\theta_1 : x_1 \cong_A y_1, \quad \theta_2 : x_2 \cong_A y_2, \quad \text{and} \quad \theta_1 \cap \theta_2 : x_1 \cap x_2 \cong_A y_1 \cap y_2.$$

As for composition, we shall rely on a characterization of its +covered configurations:

Proposition 3.30. *Consider A any arena. Then, we have:*

$$\begin{aligned}\mathcal{C}^+(\mathfrak{c}_A) &= \{x_A \parallel x_A \in \mathcal{C}(A \parallel A) \mid x_A \in \mathcal{C}(A)\} \\ \mathcal{S}^+(\mathfrak{c}_A) &= \{\theta_A \parallel \theta_A \in \mathcal{S}(A \parallel A) \mid \theta_A \in \mathcal{S}(A)\}.\end{aligned}$$

In $+$ -covered configurations, any Opponent moves must have caused a Player move; so they must all have been forwarded. This makes $+$ -covered configurations completely balanced; once again it hints at the link with relational semantics as through its $+$ -covered configurations, \mathfrak{c}_A matches the identity relation on configurations.

3.4.4. Congruence. Arenas and strategies up to isomorphism form a bicategory, where the 2-cells are given by a notion of *maps between strategies*: these are defined as maps of event structures that commute with the display maps [CCW19]. For the purposes of semantics, we usually consider strategies up to isomorphism, and this gives a category. Unfortunately this category is not the right one: the requirement that isomorphisms commute with the display map is too strict. For $!$ to satisfy the comonad laws, we require a more permissive equivalence relation between strategies, which allows for the choice of copy indices to be different in each strategy. Formally, this weaker notion of isomorphism is given by a map of event structures which commutes with the display maps *up to positive symmetry*:

Definition 3.31. Consider $\sigma, \tau : A$ two causal strategies on arena A .

A **positive isomorphism** $\varphi : \sigma \approx \tau$ is an isomorphism of ess satisfying

$$\partial_\tau \circ \varphi \sim^+ \partial_\sigma,$$

i.e. for all $x \in \mathcal{C}(\sigma)$, $\{(\partial_\sigma(s), \partial_\tau \circ \varphi(s)) \mid s \in x\} \in \mathcal{S}_+(A)$. If there is a positive iso $\varphi : \sigma \approx \tau$ we say σ and τ are **positively isomorphic**, and write $\sigma \approx \tau$.

This means that σ and τ are the same up to renaming of events. Intuitively this renaming might cause a reindexing of positive events, but it must keep the copy indices of negative events unchanged. Crucially, the induced notion of equivalence is preserved by composition, *i.e.* it is a congruence.

Proposition 3.32. *Consider $\sigma, \sigma' : A \vdash B$, $\tau, \tau' : B \vdash C$ s.t. $\sigma \approx \sigma'$ and $\tau \approx \tau'$.*

Then, we have $\tau \odot \sigma \approx \tau' \odot \sigma'$.

The proof is fairly elaborate [CCW19]. Details are out of scope, however the main argument is reviewed later as Proposition 4.4: from two configurations able to synchronize *up to symmetry*, one can extract symmetric configurations synchronizing *on the nose*.

3.5. Composition of visible and exhaustive strategies. The fact that visible strategies compose is detailed in [CC21], and we do not repeat the details here. Note that for any arena A , the copycat strategy $\mathfrak{c}_A : A \vdash A$ is visible, and for arenas A, B and C and strategies $\sigma : A \vdash B$ and $\tau : B \vdash C$, if σ and τ are visible then so is $\tau \odot \sigma$. We emphasize that these results apply to arenas but not to arbitrary games; in particular, the proof exploits that every event has a unique immediate predecessor in the game [CC21].

The compositionality of exhaustivity does not appear anywhere in the literature, so we include the details, straightforward as they are. We start with copycat.

Proposition 3.33. *For any arena A , $\mathfrak{c}_A : A \vdash A$ is exhaustive.*

Proof. By Lemma 3.30, $+$ -covered configurations of copycat are $x_A \parallel x_A \in \mathcal{C}^+(\mathfrak{c}_A)$ for $x_A \in \mathcal{C}(A)$. But then $\kappa_{A \vdash A}(x_A \parallel x_A) = (-\kappa_A(x_A)) \wp \kappa_A(x_A)$, always non-negative. \square

We now prove that exhaustive strategies are stable under composition.

Proposition 3.34. *Consider A, B and C games, and $\sigma : A \vdash B$, $\tau : B \vdash C$ exhaustive.*

Then, $\tau \odot \sigma : A \vdash C$ is exhaustive.

Proof. By Proposition 3.28, any $+$ -covered configuration of $\tau \odot \sigma$ has the form $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$ for $x^\sigma \in \mathcal{C}^+(\sigma)$ and $x^\tau \in \mathcal{C}^+(\tau)$ causally compatible. We write projections as:

$$\partial_\sigma x^\sigma = x_A \parallel x_B, \quad \partial_\tau x^\tau = x_B \parallel x_C$$

and $\partial_{\tau \odot \sigma}(x^\tau \odot x^\sigma) = x_A \parallel x_C$. If $\kappa_A(x_A) = -1$ or $\kappa_C(x_C) = 1$, then $\kappa_{A \vdash C}(x_A \parallel x_C) = 1$ and we are done. Hence, assume $\kappa_A(x_A) \geq 0$ and $\kappa_C(x_C) \geq 0$. If $\kappa_A(x_A) = 1$, then we must have $\kappa_B(x_B) = 1$ as well since $\kappa_{A \vdash B}(x_A \parallel x_B) \geq 0$ since σ is exhaustive. But then, with the same reasoning by exhaustivity of τ , $\kappa_C(x_C) = 1$, contradiction. Symmetrically, if $\kappa_C(x_C) = -1$ we may deduce that $\kappa_A(x_A) = -1$, contradiction. The only case left has $\kappa_A(x_A) = 0$ and $\kappa_C(x_C) = 0$, so that $\kappa_{A \vdash C}(x_A \parallel x_C) = 0$ as needed. \square

3.6. Categorical structure. We now outline the categorical structure of our model, as required for **nPCF**, exploiting the various constructions on games introduced in 3.2. Due to the 2-dimensional nature of the model (isomorphisms between strategies play an important role), there are subtleties in the presentation of this structure, which we explain now.

Strategies on thin concurrent games form a bicategory, where associativity and unit laws for composition hold only up to iso. Above we introduced a weaker notion, positive isomorphism, which is useful for considering strategies up to a choice of copy indices. This gives rise to another bicategory with the same objects and morphisms, but more 2-cells.

A natural step could be to present the categorical structure at this level (as in [Pdq20]). This is mathematically important but technical, because many additional coherence laws involving 2-cells must be verified. For this paper this is not necessary as any 2-dimensional structure disappears in the collapse to $\mathcal{R}\text{-Rel}$. Thus we compromise and define a model consisting of objects (arenas), morphisms (exhaustive, visible strategies), and an equivalence relation on each hom-set (\approx), such that the laws for composition hold up to \approx . In other words, we record which strategies are positively isomorphic, but forget the isos.

We call this a \sim -category. There is a general theory of these, in which the usual coherence laws for morphisms hold only up to equivalence¹. In particular there are canonical notions of (\sim)-functor, (\sim)-comonad, monoidal \sim -category, and so on, which we use below.

3.6.1. \sim -categories. We start by defining \sim -categories:

Definition 3.35. A (small) \sim -category \mathcal{C} consists in a set of *objects* \mathcal{C}_0 ; for each A, B , a set of *morphisms* $\mathcal{C}(A, B)$ with an equivalence relation \sim ; a *composition* operation

$$(- \circ -) : \mathcal{C}(B, C) \times \mathcal{C}(A, B) \rightarrow \mathcal{C}(A, C)$$

for all $A, B, C \in \mathcal{C}_0$; an *identity* morphism $\text{id}_A \in \mathcal{C}(A, A)$ for all $A \in \mathcal{C}_0$, subject to:

- associativity:* for all $f \in \mathcal{C}(A, B), g \in \mathcal{C}(B, C), h \in \mathcal{C}(C, D)$, $(h \circ g) \circ f \sim h \circ (g \circ f)$,
- identity:* for all $f \in \mathcal{C}(A, B)$, $\text{id}_B \circ f \sim f \circ \text{id}_A \sim f$,
- congruence:* for all $f \sim f' \in \mathcal{C}(A, B), g \sim g' \in \mathcal{C}(B, C)$, $g' \circ f' \sim g \circ f$.

¹We note that a \sim -category is an enriched bicategory [GS16], where the enrichment is over a 2-category of sets with equivalence relations, equivalence-preserving maps, and equivalence between maps, with the latter defined pointwise. This gives a formal justification for our definitions.

In particular, the development above already gives us our main \sim -category of interest:

Proposition 3.36. *There is \mathbf{Strat} , a \sim -category with arenas as objects; morphisms from A to B the exhaustive, visible strategies on $A \vdash B$; and equivalence relation \approx .*

Proof. The required structure was provided above; it remains to prove that associativity of composition and neutrality of copycat hold up to \approx . In other words, we must provide

$$\begin{aligned} \alpha_{\sigma,\tau,\delta} &: (\delta \odot \tau) \odot \sigma \approx \delta \odot (\tau \odot \sigma) \\ \lambda_\sigma &: \mathbf{c}_B \odot \sigma \approx \sigma \\ \rho_\sigma &: \sigma \odot \mathbf{c}_A \approx \sigma, \end{aligned}$$

for all $\sigma : A \vdash B$, $\tau : B \vdash C$ and $\delta : C \vdash D$. On $+$ -covered configurations, they are:

$$\begin{aligned} \alpha_{\sigma,\tau,\delta} &: \mathcal{C}^+((\delta \odot \tau) \odot \sigma) \cong \mathcal{C}^+(\delta \odot (\tau \odot \sigma)) & \lambda_\sigma &: \mathcal{C}^+(\mathbf{c}_B \odot \sigma) \cong \mathcal{C}^+(\sigma) \\ (x^\delta \odot x^\tau) \odot x^\sigma &\mapsto x^\delta \odot (x^\tau \odot x^\sigma) & (x_B \parallel x_B) \odot x^\sigma &\mapsto x^\sigma \end{aligned}$$

and symmetrically for ρ_σ ; using Propositions 3.28 and 3.30. In fact, these bijections between configurations are sufficient to define the required isomorphisms of strategies (provided one checks that they are order-isomorphisms, compatible with symmetry, and that they commute with display maps) – see [CC21]. In this paper, we only need the above characterization of their action on configurations. Congruence is Proposition 3.32. \square

Functors between \sim -categories must preserve \sim , and preserve composition and identities up to \sim . Of course, any \sim -category quotients to a category whose morphisms are equivalence classes. But it is preferable to refrain from quotienting: this way, the interpretation yields a concrete strategy rather than an equivalence class. This is particularly relevant for recursion which involves an complete partial order on concrete strategies, whereas it is unknown if positive isomorphism classes satisfy the adequate completeness properties. Note that this subtlety is present in all game semantics involving explicit copy indices, including in AJM games [AJM00], though it is usually handled implicitly.

3.6.2. Relative Seely categories. Following Section 3.2.4, some constructions are only available for *strict* games: $A \& B$ is only defined when A and B are strict (Definition 3.12), and $A \multimap B$ is only defined when B is strict (Definition 3.14). As we explain in more details below, this means that some of the Seely category structure in \mathbf{Strat} only exists *relative* to the inclusion functor $\mathbf{Strat}_s \hookrightarrow \mathbf{Strat}$, where \mathbf{Strat}_s is the full sub- \sim -category of *strict* arenas. We introduce a notion of *relative Seely category*, which generalises Seely categories, in which some constructions ($\&$ and \multimap) are only available for a sub-family of objects.

Additionally, while $!$ is always available in \mathbf{Strat} (there is a comonad $! : \mathbf{Strat} \rightarrow \mathbf{Strat}$), the collapse of Section 6 only preserves $!$ on strict arenas. So to precisely capture the logical content of this collapse operation, it makes sense to consider $!$ as a *relative comonad* $\mathbf{Strat}_s \hookrightarrow \mathbf{Strat}$, in the sense of [ACU10]. The axioms for relative Seely categories ensure that the induced Kleisli category (with objects are the strict ones), is cartesian closed.

Relative Seely categories model the fragment of Intuitionistic Linear Logic with

$$\begin{aligned} S, T &::= \top \mid S \& T \mid A \multimap S \\ A, B &::= 1 \mid A \otimes B \mid S \mid !S \end{aligned}$$

as formulas, separated into *strict* S, T and *general* A, B formulas. This corresponds to the logical structure preserved by our collapse; note that this covers all formulas involved in Girard’s call-by-name translation for **nPCF** types.

In the following we make use of the standard notions of relative adjunctions and relative comonads. For the reader unfamiliar with these, the definition also contains explicit data. For full details see Appendix A.1.

Definition 3.37. A **relative Seely category** is a symmetric monoidal category $(\mathcal{C}, \otimes, 1)$ equipped with a full subcategory \mathcal{C}_s together with the following data and axioms:

- \mathcal{C}_s has finite products $(\&, \top)$ preserved by the inclusion functor $J : \mathcal{C}_s \hookrightarrow \mathcal{C}$.
- For every $B \in \mathcal{C}$ there is a functor $B \multimap - : \mathcal{C}_s \rightarrow \mathcal{C}_s$, such that there is a natural bijection

$$\Lambda(-) : \mathcal{C}(A \otimes B, S) \simeq \mathcal{C}(A, B \multimap S).$$

for every $A \in \mathcal{C}$ and $S \in \mathcal{C}_s$. In other words, the functors

$$- \otimes B : \mathcal{C} \rightarrow \mathcal{C} \quad J(B \multimap -) : \mathcal{C}_s \rightarrow \mathcal{C}$$

form a J -relative adjunction $- \otimes B \dashv_J J(- \multimap B)$.

- There is a J -relative comonad $! : \mathcal{C}_s \rightarrow \mathcal{C}$. Concretely we have, for every $S \in \mathcal{C}_s$, an object $!S \in \mathcal{C}$ and a morphism $\epsilon_S : !S \rightarrow S$, and for every $\sigma : !S \rightarrow T$, a **promotion** $\sigma^\dagger : !S \rightarrow !T$, subject to three axioms [ACU10]:

$$\begin{aligned} \epsilon_T \circ \sigma^\dagger &= \sigma & (\sigma : !S \rightarrow T) \\ \epsilon_S^\dagger &= \text{id}_{!S} & (S \in \mathcal{C}_s) \\ (\tau \circ \sigma^\dagger)^\dagger &= \tau^\dagger \circ \sigma^\dagger & (\sigma : !S \rightarrow T, \tau : T \rightarrow U), \end{aligned}$$

which make $! : \mathcal{C}_s \rightarrow \mathcal{C}$ a functor, via $!\sigma = (\sigma \circ \epsilon_S)^\dagger$ for $\sigma : S \rightarrow T$.

- The functor $! : \mathcal{C}_s \rightarrow \mathcal{C}$ is symmetric strong monoidal $(\mathcal{C}_s, \&, \top) \rightarrow (\mathcal{C}, \otimes, 1)$, so

$$m_0 : 1 \rightarrow !\top \quad m_{S,T} : !S \otimes !T \rightarrow !(S \& T)$$

are natural isos, additionally compatible with promotion in the sense that the diagram

$$\begin{array}{ccc} !\Gamma & \xrightarrow{\langle f, g \rangle^\dagger} & !(S \& T) \\ \langle \epsilon, \epsilon \rangle^\dagger \downarrow & & \downarrow m^{-1} \\ !(\Gamma \& \Gamma) & & !S \otimes !T \\ & \searrow m^{-1} \quad \nearrow f^\dagger \otimes g^\dagger & \\ & !\Gamma \otimes !\Gamma & \end{array}$$

commutes for all $\Gamma, S, T \in \mathcal{C}_s$, $f \in \mathcal{C}(!\Gamma, S)$, $g \in \mathcal{C}(!\Gamma, T)$.

Note that any Seely category is canonically a relative Seely category.

For any relative Seely category, the Kleisli category associated with the relative comonad $!$ is cartesian closed. This category, denoted $\mathcal{C}_!$, has objects those of \mathcal{C}_s , and $\mathcal{C}_!(S, T) = \mathcal{C}(!S, T)$. The proof is essentially as for Seely categories; details are in Appendix A.2.

Lemma 3.38. *For a relative Seely category \mathcal{C} , the Kleisli category $\mathcal{C}_!$ is cartesian closed with finite products given as in \mathcal{C}_s , and function space $S \Rightarrow T = !S \multimap T$.*

For **Strat** these definitions must be taken in \sim -categorical form, this is what we do next. The generalisation is very straightforward, but we give the main definitions along the way.

3.6.3. *Symmetric monoidal structure.* We first extend the symmetric monoidal structure.

Symmetric monoidal \sim -categories are defined as expected; where the usual data required to preserve \sim and satisfy laws up to \sim . For \sim -categories \mathcal{C} and \mathcal{D} , their *product* $\mathcal{C} \times \mathcal{D}$ has objects $\mathcal{C}_0 \times \mathcal{D}_0$, morphisms $(A, B) \rightarrow (C, D)$ the pairs $(f, g) \in \mathcal{C}(A, C) \times \mathcal{D}(B, D)$, with obvious identity and compositions, and $(f, g) \sim (f', g')$ if $f \sim f'$ and $g \sim g'$.

Definition 3.39. A **symmetric monoidal \sim -category** is a \sim -category \mathcal{C} equipped with an object $1 \in \mathcal{C}_0$, a \sim -functor

$$(- \otimes -) : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C},$$

and, for all $A, B, C \in \mathcal{C}_0$, morphisms

$$\begin{aligned} \alpha_{A,B,C} &: (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C) \\ \lambda_A &: 1 \otimes A \rightarrow A \\ \rho_A &: A \otimes 1 \rightarrow A \\ s_{A,B} &: A \otimes B \rightarrow B \otimes A \end{aligned}$$

invertible up to \sim , and such that the usual naturality and coherence conditions for a symmetric monoidal category [ML13] hold up to \sim .

For Strat, on arenas the tensor is an instance of the tensor of games given in Definition 3.10. The unit 1 is the empty arena where the empty configuration has payoff 0. This operation extends to a tensor product of strategies, defined using parallel composition of event structures. Rather than giving the concrete construction we state a characterization in terms of $+$ -covered configurations:

Proposition 3.40. *Consider A, B, C, D arenas, and $\sigma : A \vdash B$, $\tau : C \vdash D$ strategies.*

Then, there is a strategy $\sigma \otimes \tau : A \otimes C \vdash B \otimes D$, unique up to iso, s.t. there are

$$\begin{aligned} (- \otimes -) &: \mathcal{C}^+(\sigma) \times \mathcal{C}^+(\tau) \simeq \mathcal{C}^+(\sigma \otimes \tau) \\ (- \otimes -) &: \mathcal{S}^+(\sigma) \times \mathcal{S}^+(\tau) \simeq \mathcal{S}^+(\sigma \otimes \tau) \end{aligned}$$

order-isos commuting with dom, cod, and s.t. for all $\theta^\sigma \in \mathcal{S}^+(\sigma)$ and $\theta^\tau \in \mathcal{S}^+(\tau)$,

$$\partial_{\sigma \otimes \tau}(\theta^\sigma \otimes \theta^\tau) = (\theta_A^\sigma \parallel \theta_C^\tau) \parallel (\theta_B^\sigma \parallel \theta_D^\tau).$$

This is clear from the construction of the tensor in [CCW19] with uniqueness coming from Lemma 4.11 of [CC21]. Bifactoriality up to isomorphism is proved in [CCW19]. The components of the symmetric monoidal structure are also given in [CCW19]; they are immediate variants of the copycat strategy of Definition 3.29. They also satisfy obvious characterizations of their $+$ -covered configurations, with for instance for the associator:

$$((x_A \parallel x_B) \parallel x_C) \parallel (x_A \parallel (x_B \parallel x_C)) \in \mathcal{C}^+(\alpha_{A,B,C})$$

for $x_A \in \mathcal{C}(A)$, $x_B \in \mathcal{C}(B)$ and $x_C \in \mathcal{C}(C)$ and $\mathcal{C}^+(\alpha_{A,B,C})$ containing exactly those, and similarly for λ_A, ρ_A and $s_{A,B}$. For this paper, the only new things to check are that the structural isomorphisms are exhaustive, and that the tensor preserves exhaustivity. The former is straightforward via characterization of configurations; we detail the latter.

Proposition 3.41. *If $\sigma : A \vdash B$ and $\tau : C \vdash D$ are exhaustive, so is $\sigma \otimes \tau : A \otimes C \vdash B \otimes D$.*

Proof. Consider $x^\sigma \otimes x^\tau \in \mathcal{C}^+(\sigma \otimes \tau)$, i.e. $x^\sigma \in \mathcal{C}^+(\sigma)$ and $x^\tau \in \mathcal{C}^+(\tau)$. If $\kappa_A(x_A^\sigma) = -1$ or $\kappa_C(x_C^\tau) = -1$, then we are done; thus assume $\kappa_A(x_A^\sigma) \geq 0$ and $\kappa_C(x_C^\tau) \geq 0$. If $\kappa_{B \otimes D}(x_B^\sigma \parallel x_D^\tau) = 1$ then we are done; thus assume $\kappa_{B \otimes D}(x_B^\sigma \parallel x_D^\tau) \leq 0$. If $\kappa_{B \otimes D}(x_B^\sigma \parallel x_D^\tau) = -1$, then $\kappa_B(x_B^\sigma) = -1$ or $\kappa_D(x_D^\tau) = -1$, say the former. Since σ is exhaustive and $x^\sigma \in \mathcal{C}^+(\sigma)$, we have $\kappa_A(x_A^\sigma) = -1$, contradiction – and likewise, $\kappa_D(x_D^\tau) = -1$ yields a contradiction.

So, $\kappa_{B \otimes D}(x_B^\sigma \parallel x_D^\tau) = 0$, *i.e.* $\kappa_B(x_B^\sigma) = 0$ and $\kappa_D(x_D^\tau) = 0$. Finally, assume $\kappa_A(x_A^\sigma) = 1$. Then, by exhaustivity of σ , $\kappa_B(x_B^\sigma) = 1$ as well, contradiction. So $\kappa_A(x_A^\sigma) = 0$, and likewise $\kappa_C(x_C^\tau) = 0$. Summing up, the overall payoff of $\partial_{\sigma \otimes \tau}(x^\sigma \otimes x^\tau)$ is 0, as required. \square

Overall, we have completed the symmetric monoidal structure of **Strat**:

Proposition 3.42. *Equipped with the above, **Strat** is a symmetric monoidal \sim -category.*

3.6.4. *Closed structure.* The (relative) closed structure is easily derived, using the linear function space construction of Section 3.2.5. For A, B, C arenas with C strict, the currying bijection $\Lambda(-) : \mathbf{Strat}(A \otimes B, C) \rightarrow \mathbf{Strat}(A, B \multimap C)$ leaves the ess unchanged and only affects the display map: for $\sigma : A \otimes B \rightarrow C$ and $x^\sigma \in \mathcal{C}^{\neq \emptyset}(\sigma)$ with $\partial_\sigma(x^\sigma) = (x_A^\sigma \parallel x_B^\sigma) \parallel x_C^\sigma$,

$$\partial_{\Lambda(\sigma)}(x^\sigma) = x_A^\sigma \parallel (x_B^\sigma \multimap x_C^\sigma)$$

using the notation introduced in Section 3.2.5.

The evaluation morphism $\mathbf{ev}_{B,C} : (B \multimap C) \otimes B \rightarrow C$ is a copycat-like strategy having

$$((x_B \multimap x_C) \parallel x_B) \parallel x_C \in \mathcal{C}^+(\mathbf{ev}_{B,C})$$

as $+$ -covered non-empty configurations, for any $x_B \in \mathcal{C}(B)$ and $x_C \in \mathcal{C}^{\neq \emptyset}(C)$.

3.6.5. *Cartesian products.* We now introduce the cartesian structure.

Definition 3.43. A \sim -category \mathcal{C} has **binary products** if for any $A, B \in \mathcal{C}$, there exists an object $A \& B \in \mathcal{C}$, and **projections** $\pi_A \in \mathcal{C}(A \& B, A)$ and $\pi_B \in \mathcal{C}(A \& B, B)$, such that for every $\Gamma \in \mathcal{C}$, $\sigma \in \mathcal{C}(\Gamma, A)$, and $\tau \in \mathcal{C}(\Gamma, B)$, there exists $\langle \sigma, \tau \rangle$, unique up to \sim , such that

$$\sigma \sim \pi_A \circ \langle \sigma, \tau \rangle \quad \tau \sim \pi_B \circ \langle \sigma, \tau \rangle.$$

An object $\top \in \mathcal{C}$ is **terminal** if for any $\Gamma \in \mathcal{C}$ there is $e_\Gamma \in \mathcal{C}(\Gamma, \top)$, unique up to \sim .

We prove that \mathbf{Strat}_s has binary products and a terminal object (so it has all *finite* products), and these are preserved by the inclusion functor $\mathbf{Strat}_s \hookrightarrow \mathbf{Strat}$.

Clearly, the arena \top with no events and $\kappa_\top(\emptyset) = 1$ is strict, and terminal in **Strat**. For any A and B strict arenas, their product is given by the construction $A \& B$ of Definition 3.12. We give a characterisation of the pairing construction $\langle -, = \rangle$.

Proposition 3.44. *For arenas Γ, A, B , with A, B strict, and strategies $\sigma : \Gamma \vdash A, \tau : \Gamma \vdash B$, there is a strategy $\langle \sigma, \tau \rangle : \Gamma \vdash A \& B$, unique up to iso, s.t. there are order-isos:*

$$\begin{aligned} \mathcal{C}^+(\sigma) + \mathcal{C}^+(\tau) &\simeq \mathcal{C}^+(\langle \sigma, \tau \rangle) \\ \mathcal{S}^+(\sigma) + \mathcal{S}^+(\tau) &\simeq \mathcal{S}^+(\langle \sigma, \tau \rangle) \end{aligned}$$

commuting with **dom**, **cod**, and such that for all $\theta^\sigma \in \mathcal{S}^+(\sigma)$ and $\theta^\tau \in \mathcal{S}^+(\tau)$, we have

$$\partial_{\langle \sigma, \tau \rangle}(\mathbf{inj}_\sigma(\theta^\sigma)) = \theta_\Gamma^\sigma \parallel (\theta_A^\sigma \parallel \emptyset), \quad \partial_{\langle \sigma, \tau \rangle}(\mathbf{inj}_\tau(\theta^\tau)) = \theta_\Gamma^\tau \parallel (\emptyset \parallel \theta_B^\tau)$$

writing $\mathbf{inj}_\sigma : \mathcal{C}^+(\sigma) \rightarrow \mathcal{C}^+(\langle \sigma, \tau \rangle)$ and $\mathbf{inj}_\tau : \mathcal{C}^+(\tau) \rightarrow \mathcal{C}^+(\langle \sigma, \tau \rangle)$ for the induced injections.

This result and the universal property of Definition 3.43 follow easily from the construction of the cartesian product of arenas given in [CC21]. Projections are the obvious copycat strategies, with $+$ -covered configurations

$$(x_A \parallel \emptyset) \parallel x_A \in \mathcal{C}^+(\pi_A), \quad (\emptyset \parallel x_B) \parallel x_B \in \mathcal{C}^+(\pi_B)$$

for $x_A \in \mathcal{C}(A), x_B \in \mathcal{C}(B)$. Their exhaustivity is easy; note however that this uses the fact that A and B are strict. We detail the proof that pairing preserves exhaustivity:

Proposition 3.45. *Consider Γ, A, B arenas with A, B strict, and $\sigma : \Gamma \vdash A, \tau : \Gamma \vdash B$.*

If σ and τ are exhaustive, then so is $\langle \sigma, \tau \rangle : \Gamma \vdash A \& B$.

Proof. Consider a $+$ -covered configuration of $\langle \sigma, \tau \rangle$, say for instance $\text{inj}_\sigma(x^\sigma) \in \mathcal{C}^+(\langle \sigma, \tau \rangle)$ for $x^\sigma \in \mathcal{C}^+(\sigma)$; with $\partial_{\langle \sigma, \tau \rangle} \text{inj}_\sigma(x^\sigma) = x_\Gamma^\sigma \parallel (x_A^\sigma \parallel \emptyset) \parallel x_A^\sigma$. If $\kappa_{A\&B}(x_A^\sigma \parallel \emptyset) = 1$, then we are done; assume $\kappa_{A\&B}(x_A^\sigma \parallel \emptyset) \leq 0$. Likewise, assume $\kappa_\Gamma(x_\Gamma^\sigma) \geq 0$. If $\kappa_\Gamma(x_\Gamma^\sigma) = 1$, then we must have $\kappa_A(x_A^\sigma) = 1$ as well since σ is exhaustive – but this implies $\kappa_{A\&B}(x_A^\sigma \parallel \emptyset) = 1$ as well, contradiction. Likewise, if $\kappa_{A\&B}(x_A^\sigma \parallel \emptyset) = -1$, then $\kappa_A(x_A^\sigma) = -1$ and so $\kappa_\Gamma(x_\Gamma^\sigma) = -1$ since σ is exhaustive, contradiction. The only case left has $\kappa_\Gamma(x_\Gamma^\sigma) = 0$ and $\kappa_{A\&B}(x_A^\sigma \parallel \emptyset) = 0$, so the display of $\text{inj}_\sigma(x^\sigma)$ has payoff 0. The symmetric reasoning applies to $+$ -covered configurations of the form $\text{inj}_\tau(x^\tau)$ for $x^\tau \in \mathcal{C}^+(\tau)$, which concludes the proof. \square

In summary, we have the following:

Proposition 3.46. *The full sub- \sim -category \mathbf{Strat}_s of strict arenas has finite products, preserved by the inclusion functor $\mathbf{Strat}_s \hookrightarrow \mathbf{Strat}$.*

3.6.6. *Exponential.* We define a \sim -comonad $! : \mathbf{Strat} \rightarrow \mathbf{Strat}$. By a general result [ACU10] the restriction of $!$ to the sub- \sim -category \mathbf{Strat}_s gives a relative comonad $! : \mathbf{Strat}_s \rightarrow \mathbf{Strat}$, and below we will show that this satisfies the axioms of Definition 3.37.

For an object $A \in \mathbf{Strat}$, $!A$ is given by Definition 3.17. Following the same pattern as for the other structure we introduce the functorial action of $!$ as follows:

Proposition 3.47. *Consider A, B arenas, and $\sigma : A \vdash B$ a strategy.*

Then there is a strategy $!\sigma : !A \vdash !B$ where the ess is obtained as in Definition 3.17 (without the clauses on positive and negative symmetries); there is an order-iso

$$[-] : \mathbf{Fam} \left(\mathcal{C}^{+, \neq \emptyset}(\sigma) \right) \simeq \mathcal{C}^+(\! \sigma)$$

with $\mathbf{Fam}(X)$ the set of families of elements of X indexed by finite subsets of \mathbb{N} , and where $\mathcal{C}^{+, \neq \emptyset}(\sigma)$ denotes the set of non-empty $+$ -covered configurations of σ . Moreover,

$$\partial_\sigma \left([(x^i)_{i \in I}] \right) = (\parallel_{i \in I} x_A^i) \parallel (\parallel_{i \in I} x_B^i)$$

where for all $i \in I$, $\partial_\sigma(x^i) = x_A^i \parallel x_B^i$.

The proof is direct from the definition of $!\sigma$ (see [CC21]). Though the symmetries of $!\sigma$ may also be described in a similar style (as for earlier operations), we refrain from doing so as it is slightly more heavy notationally: indeed, whereas for tensor and pairing the symmetries are constructed exactly as for configurations, in $!\sigma$ symmetries span components as they may freely exchange copy indices as stated in Definition 3.17.

The \sim -comonad structure of $!$ is given by natural transformations $\epsilon_A : !A \rightarrow A$ and $\delta_A : !A \rightarrow !!A$ whose components are, as usual, relabeled copycat strategies, characterized by the shape of their $+$ -covered configurations:

$$\begin{array}{ccc} \left(\parallel_{\langle i,j \rangle \in \mathbb{N}} x_A^{i,j} \right) & \parallel & \left(\parallel_{i \in \mathbb{N}} \parallel_{j \in \mathbb{N}} x_A^{i,j} \right) \in \mathcal{C}^+(\delta_A) \\ \{0\} \times x_A & \parallel & x_A \in \mathcal{C}^+(\epsilon_A) \end{array}$$

whenever these sets are finite, $x_A, x_A^{i,j} \in \mathcal{C}(A)$, and using a fixed bijection $\langle -, - \rangle : \mathbb{N}^2 \simeq \mathbb{N}$.

It is routine that these are exhaustive – for ϵ_A , the case of the empty set uses condition *initialized*. We must also prove that the functorial operation preserves exhaustivity:

Proposition 3.48. *Consider A, B arenas, and $\sigma : A \vdash B$ a strategy.*

If σ is exhaustive, then so is $!\sigma : !A \vdash !B$.

Proof. Consider $[(x^i)_{i \in I}] \in \mathcal{C}^+(!\sigma)$, where $I \subseteq_f \mathbb{N}$ and $x^i \in \mathcal{C}^{+, \neq \emptyset}(\sigma)$ for all $i \in I$. We have

$$\partial_\sigma \left([(x^i)_{i \in I}] \right) = \left(\parallel_{i \in I} x_A^i \right) \parallel \left(\parallel_{i \in I} x_B^i \right)$$

where $\partial_\sigma(x^i) = x_A^i \parallel x_B^i \in \mathcal{C}(A \vdash B)$. If $\kappa_{!B}(\parallel_{i \in I} x_B^i) = 1$ then we are done, so assume $\kappa_{!B}(\parallel_{i \in I} x_B^i) \leq 0$. Symmetrically, assume $\kappa_{!A}(\parallel_{i \in I} x_A^i) \geq 0$. If $\kappa_{!B}(\parallel_{i \in I} x_B^i) = -1$, then there must be some $i_0 \in \mathbb{N}$ such that $\kappa_B(x_B^{i_0}) = -1$. But because σ is exhaustive, this implies that $\kappa_A(x_A^{i_0}) = -1$ as well; but then $\kappa_{!A}(\parallel_{i \in I} x_A^i) = -1$, contradiction. So, $\kappa_{!B}(\parallel_{i \in I} x_B^i) = 0$, which entails that for all $i \in I$, $\kappa_B(x_B^i) = 0$ (note that as $x^i \in \mathcal{C}^{+, \neq \emptyset}(\sigma)$, all x_B^i are non-empty). Now, if $\kappa_{!A}(\parallel_{i \in I} x_A^i) = 1$, then there must be some $i_0 \in I$ such that $\kappa_A(x_A^{i_0}) = 1$. But as σ is exhaustive, this entails that $\kappa_B(x_B^{i_0}) = 1$ as well, contradiction. \square

The functor $!$ preserves \approx and satisfies all necessary axioms up to \approx [CC21], so that:

Lemma 3.49. *The functor $! : \mathbf{Strat} \rightarrow \mathbf{Strat}$ is a \sim -comonad, which restricts to a J -relative comonad $\mathbf{Strat}_s \rightarrow \mathbf{Strat}$, for J the inclusion functor.*

Finally we have, for strict B, C , a strategy $m_{B,C} : !B \otimes !C \rightarrow !(B \& C)$ characterized by

$$\left(\left(\parallel_{i \in \mathbb{N}} x_B^i \right) \parallel \left(\parallel_{j \in \mathbb{N}} x_C^j \right) \right) \parallel \left(\begin{array}{c} \uplus_{j \in \mathbb{N}} \{2j+1\} \times (\emptyset \parallel x_C^j) \\ \uplus \uplus_{i \in \mathbb{N}} \{2i\} \times (x_B^i \parallel \emptyset) \end{array} \right) \in \mathcal{C}^+(m_{B,C})$$

for $x_B^i \in \mathcal{C}(B), x_C^j \in \mathcal{C}(C)$. Likewise, the strategy $m_0 : !\top \rightarrow 1$ with only the empty configuration. The axioms can be verified, and we finally obtain:

Proposition 3.50. *The \sim -category \mathbf{Strat} , equipped with the full sub- \sim -category \mathbf{Strat}_s of strict arenas, and all components outlined above, is a relative Seely \sim -category.*

In particular, the Kleisli \sim -category $\mathbf{Strat}_!$, with objects strict arenas, is cartesian closed.

3.7. Interpretation of nPCF. Combinators of the simply-typed λ -calculus are handled in the standard way following the cartesian closed structure of $\mathbf{Strat}_!$ [LS88], the only structure left to finalize the interpretation is the interpretation of **nPCF** primitives, and recursion.

$$\begin{array}{ccc}
\text{if} : !(\mathbf{B} \ \& \ \mathbf{X} \ \& \ \mathbf{X}) \vdash \mathbf{X} & \text{if} : !(\mathbf{B} \ \& \ \mathbf{X} \ \& \ \mathbf{X}) \vdash \mathbf{X} & \text{succ} : !\mathbf{N} \vdash \mathbf{N} \\
\mathbf{q}_0^+ & \mathbf{q}_0^+ & \mathbf{q}_0^+ \\
\mathbf{tt}_0^- & \mathbf{ff}_0^- & n_0^- \\
\mathbf{q}_1^+ & \mathbf{q}_1^+ & (n+1)^+ \\
v_1^- & v_1^- & \\
v^+ & v^+ & \\
\end{array}$$

$$\begin{array}{ccc}
\text{iszero} : !\mathbf{N} \vdash \mathbf{B} & \text{iszero} : !\mathbf{N} \vdash \mathbf{B} & \text{pred} : !\mathbf{N} \vdash \mathbf{N} & \text{pred} : !\mathbf{N} \vdash \mathbf{N} \\
\mathbf{q}_0^+ & \mathbf{q}_0^+ & \mathbf{q}_0^+ & \mathbf{q}_0^+ \\
0_0^- & (n+1)_0^- & 0_0^- & (n+1)_0^- \\
\mathbf{tt}^+ & \mathbf{ff}^+ & 0^+ & n^+
\end{array}$$

Figure 7: Strategies for basic **PCF** combinators

$$\begin{aligned}
\llbracket \text{if } M \ N_1 \ N_2 \rrbracket &= \text{if} \odot! \langle \llbracket M \rrbracket, \llbracket N_1 \rrbracket, \llbracket N_2 \rrbracket \rangle \\
\llbracket \text{succ } M \rrbracket &= \text{succ} \odot! \llbracket M \rrbracket \\
\llbracket \text{pred } M \rrbracket &= \text{pred} \odot! \llbracket M \rrbracket \\
\llbracket \text{iszero } M \rrbracket &= \text{iszero} \odot! \llbracket M \rrbracket
\end{aligned}$$




Figure 8: Interpretation of basic combinators

Figure 9: Interpretation of **coin**

3.7.1. *Base types and primitives.* The arenas for the base types are given in Section 3.2.2.

For the **PCF** primitives, the constants $\Gamma \vdash \mathbf{tt}, \mathbf{ff} : \mathbb{B}$ and $\Gamma \vdash n : \mathbb{N}$ are interpreted by the obvious strategies which reply accordingly to the initial move. Conditionals along with the primitives **iszero**, **pred**, and **succ** are interpreted following the clauses of Figure 8, using the strategies shown in Figure 7 – note that we omit arrows to avoid clutter.

3.7.2. *Recursion.* The fixpoint combinator is defined as the least upper bound of its finite approximants, leveraging completeness properties of the following partial order:

Definition 3.51. Consider A a game, and $\sigma, \tau : A$ strategies.

We write $\sigma \preceq \tau$ if $\mathcal{C}(\sigma) \subseteq \mathcal{C}(\tau)$ – so in particular $|\sigma| \subseteq |\tau|$, and additionally:

- (1) for all $s_1, s_2 \in |\sigma|$, $s_1 \leq_\sigma s_2$ iff $s_1 \leq_\tau s_2$,
- (2) for all $s_1, s_2 \in |\sigma|$, $s_1 \#_\sigma s_2$ iff $s_1 \#_\tau s_2$,
- (3) for all $x, y \in \mathcal{C}(\sigma)$ and bijection $\theta : x \simeq y$, we have $\theta \in \mathcal{S}(\sigma)$ iff $\theta \in \mathcal{S}(\tau)$,
- (4) for all $s \in |\sigma|$, $\partial_\sigma(s) = \partial_\tau(s)$,

i.e. all components compatible with the inclusion.

Strategies on A , ordered by \preceq , form a *directed complete partial order*, with respect to which all operations on strategies are easily shown to be continuous. We have:

Proposition 3.52. Consider A a game, and D a directed set of strategies on A . Then:

$$\mathcal{C}^+(\vee D) = \bigcup_{\sigma \in D} \mathcal{C}^+(\sigma), \quad \mathcal{S}^+(\vee D) = \bigcup_{\sigma \in D} \mathcal{S}^+(\sigma).$$

Moreover, if every $\sigma \in D$ is exhaustive, then so is $\vee D : A$.

The supremum also preserves visibility; so for any A, B the homset $\text{Strat}(A, B)$ is a dcpo. Before defining the recursion operator via the usual fixpoint formula, we must deal with the minor inconvenience that the dcpo of strategies on A does not have a least element: strategies minimal for \triangleleft still have – by receptivity – events matching the minimal events of A , but they are free to *name* those arbitrarily. We solve this as in [CCW19]: we choose one minimal $\perp_A : A$. For any $\sigma : A$, we pick an isomorphic $\sigma \cong \sigma^b : A$ s.t. $\perp_A \triangleleft \sigma^b$, obtained by renaming minimal events. We write \mathcal{D}_A for the pointed dcpo of strategies above \perp_A .

As all operations on strategies examined so far are continuous, for any arena A we have

$$F : \mathcal{D}_{!\top \vdash (A \rightarrow A) \rightarrow A} \rightarrow \mathcal{D}_{!\top \vdash (A \rightarrow A) \rightarrow A} \\ \sigma \mapsto (\lambda f^{A \rightarrow A}. f(\sigma f))^b,$$

written in λ -calculus syntax relying on the constructions on strategies corresponding to the cartesian closed structure of $\text{Strat}_!$, continuous. By Kleene's fixpoint theorem

$$\mathcal{Y}_A = \bigvee_{n \in \mathbb{N}} F^n(\perp) \in \text{Strat}_!(\top, (A \rightarrow A) \rightarrow A)$$

is a least fixpoint of F . Finally, in the presence of a context Γ , we set $\mathcal{Y}_{\Gamma, A} = \mathcal{Y}_A \odot_! e_\Gamma$ where $e_\Gamma \in \text{Strat}_!(\Gamma, \top)$ is the terminal morphism, yielding $\mathcal{Y}_{\Gamma, A} \in \text{Strat}_!(\Gamma, !(A \multimap A) \multimap A)$.

This concludes the interpretation of **nPCF** in Strat . From [CC21], it is adequate with respect to may-convergence – however this statement will play no role here. Most of the rest of the paper will study the collapse of Strat (and its interpretation of **nPCF**) onto $\mathcal{N}\text{-Rel}$.

3.8. Collapsing Strat to $\mathcal{N}\text{-Rel}$. Back to the question asked in Section 2.4.3: what does the \mathcal{N} -weighted relational model count, on arbitrary higher-order types?

We shall answer this by providing a *collapse* interpretation-preserving functor, written

$$f(-) : \text{Strat} \rightarrow \mathcal{N}\text{-Rel}.$$

At first this seems simple: recall from Lemma 3.19 that for any type A , there is $s_A^{\text{ty}} : \langle A \rangle \simeq \mathcal{C}_{\cong}^0(\llbracket A \rrbracket)$ a bijection through which we regard $\langle A \rangle$ – the **web** of A , as it is usually called – as the set of symmetry classes of null payoff of $\llbracket A \rrbracket$. Given $\sigma : A$ and $x_A \in \mathcal{C}_{\cong}^0(\llbracket A \rrbracket)$, we must associate a *weight* $(f(\sigma))_{x_A} \in \mathbb{N} \cup \{+\infty\}$. It seems natural for this weight to be

$$(f(\sigma))_{x_A} = \sharp(\text{wit}_\sigma(x_A))$$

the cardinality of an adequately chosen set of *witnesses* of x_A . On ground types this seems rather clear: for instance, considering $\vdash M : \mathbb{B}$ defined as $M = \mathbf{if\ coin\ tt\ tt}$, we have

$$\llbracket M \rrbracket = \begin{array}{c} \mathbf{q}^- \\ \swarrow \quad \searrow \\ \mathbf{tt}^+ \text{---} \text{---} \text{---} \mathbf{tt}^+ \end{array} \quad \begin{array}{l} \llbracket M \rrbracket_{\mathbf{tt}} = 2 \\ \llbracket M \rrbracket_{\mathbf{ff}} = 0 \end{array}$$

so one may guess that $\text{wit}(x_A) = \{x^\sigma \in \mathcal{C}(\sigma) \mid \partial_\sigma(x^\sigma) \in x_A\}$ – however, this is inadequate beyond ground types: if $x^\sigma \in \mathcal{C}(\sigma)$ includes any copyable Opponent move, then x^σ automatically has countably many symmetric copies prompted by copies of that Opponent move, displayed to the same configuration of the game up to symmetry. So this notion of witnesses is useless save in the most simple cases, as it makes no account of symmetry.

So what is the right set of witnesses up to symmetry? *How to count configurations of a strategy up to symmetry?* This question, the crux of the paper, is investigated next.

4. WITNESSES AND COMPOSITION

To set up our collapse, the most challenging proof obligation – by far – is preservation of composition. Here we present the right definition of collapse, along with the proof that it preserves composition. But rather than arriving there directly, we take a more indirect route, showing some of the subtleties constraining the solution.

4.1. Finding the Right Witnesses. A fitting starting point for this discussion is to recall

$$(\beta \circ \alpha)_{x,z} = \sum_{y \in Y} \alpha_{x,y} \cdot \beta_{y,z} \quad (4.1)$$

the composition of $\alpha \in \mathcal{N}\text{-Rel}(X, Y)$ and $\beta \in \mathcal{N}\text{-Rel}(Y, Z)$, which we must relate to the composition of strategies. As explained above, for $\sigma \in \text{Strat}(A, B)$ and $x_A \in \mathcal{C}_{\cong}^0(A), x_B \in \mathcal{C}_{\cong}^0(B)$, we expect $(f(\sigma))_{x_A, x_B}$ to be the cardinality of a well-chosen set $\text{wit}_{\sigma}(x_A, x_B)$ of *witnesses* for $x_A \parallel x_B \in \mathcal{C}_{\cong}^0(A \vdash B)$. Thus, (4.1) strongly hints at a bijection

$$\text{wit}_{\tau \circ \sigma}(x_A, x_C) \simeq \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} \text{wit}_{\sigma}(x_A, x_B) \times \text{wit}_{\tau}(x_B, x_C) \quad (4.2)$$

for all $\sigma \in \text{Strat}(A, B), \tau \in \text{Strat}(B, C), x_A \in \mathcal{C}_{\cong}^0(A)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$.

4.1.1. Witnesses without symmetry. Let us start by ignoring symmetry, and first count witnesses for plain complete configurations $x_A \in \mathcal{C}^0(A)$. It seems natural to define:

$$\text{wit}_{\sigma}(x_A, x_B) = \{x^{\sigma} \in \mathcal{C}^+(\sigma) \mid \partial_{\sigma}(x^{\sigma}) = x_A \parallel x_B\}.$$

for all $\sigma \in \text{Strat}(A, B), x_A \in \mathcal{C}^0(A)$ and $x_B \in \mathcal{C}^0(B)$.

Working with $+$ -covered configurations seems reasonable, as Proposition 3.28 entails:

$$(- \odot -) : \{(x^{\tau}, x^{\sigma}) \in \mathcal{C}^+(\tau) \times \mathcal{C}^+(\sigma) \mid x^{\sigma} \text{ and } x^{\tau} \text{ causally compatible}\} \simeq \mathcal{C}^+(\tau \odot \sigma)$$

preserving the display maps for all $\sigma : A \vdash B$ and $\tau : B \vdash C$, which seems close to (4.2). The two obstacles to (4.2) are that: (1) this bijection has an additional requirement that synchronized $x^{\sigma} \in \mathcal{C}^+(\sigma)$ and $x^{\tau} \in \mathcal{C}^+(\tau)$ should be *causally compatible* – a constraint which does not appear in (4.2); and (2) we must be sure that for all $x^{\sigma} \in \mathcal{C}^+(\sigma)$ and $x^{\tau} \in \mathcal{C}^+(\tau)$ synchronizable, if $x_A \in \mathcal{C}^0(A)$ and $x_C \in \mathcal{C}^0(C)$, then $x_B^{\sigma} = x_B^{\tau} \in \mathcal{C}^0(B)$.

For (1), we invoke the “*deadlock-free lemma*” from [CC21]:

Lemma 4.1. *Consider A, B, C arenas, $\sigma : A \vdash B$ and $\tau : B \vdash C$ visible strategies, $x^{\sigma} \in \mathcal{C}(\sigma)$ and $x^{\tau} \in \mathcal{C}(\tau)$ with a symmetry $\theta : x_B^{\sigma} \cong_B x_B^{\tau}$. Then, the composite bijection*

$$\varphi : x^{\sigma} \parallel x_C^{\tau} \xrightarrow{\partial_{\sigma} \parallel x_C^{\tau}} x_A^{\sigma} \parallel x_B^{\sigma} \parallel x_C^{\tau} \xrightarrow{x_A^{\sigma} \parallel \theta \parallel x_C^{\tau}} x_A^{\sigma} \parallel x_B^{\tau} \parallel x_C^{\tau} \xrightarrow{x_A^{\sigma} \parallel \partial_{\tau}^{-1}} x_A^{\sigma} \parallel x^{\tau},$$

is secured, in the sense that the relation \triangleleft , defined on the graph of φ with

$$(l, r) \triangleleft (l', r')$$

whenever $l \langle_{\sigma} \parallel \langle_C \rangle l'$ or $r \langle_A \parallel \langle_{\tau} \rangle r'$, is acyclic².

²For θ an identity, this exactly means that x^{σ} and x^{τ} satisfy the *secured* condition of Definition 3.26.

The quite subtle proof is out of the scope of the paper, the interested reader is referred to [CC21]. It is one of the main properties of *visibility* that composition never deadlocks, so that when dealing with visible strategies, the *causally compatible* requirement of Proposition 3.28 is redundant – the lemma above also covers the case of synchronization through a symmetry θ , which will be necessary later on in the paper.

Next, for (2), we prove the following property:

Lemma 4.2. *Consider games A, B and C ; and exhaustive $\sigma : A \vdash B$ and $\tau : B \vdash C$.*

For all $x^\sigma \in \mathcal{C}^+(\sigma)$ and $x^\tau \in \mathcal{C}^+(\tau)$, if $\kappa_A(x_A^\sigma) = 0$ and $\kappa_C(x_C^\tau) = 0$, then $\kappa_B(x_B) = 0$.

Proof. Seeking a contradiction, assume $\kappa_B(x_B) = 1$. Then $\kappa_{B \vdash C}(x_B \parallel x_C^\tau) = -1$, contradicting that τ is exhaustive. Symmetrically, $\kappa_B(x_B) = -1$ contradicts that σ is exhaustive. \square

From these two statements, we may now deduce as claimed:

Corollary 4.3. *Consider A, B, C arenas, $\sigma \in \text{Strat}(A, B)$, $\tau \in \text{Strat}(B, C)$. Then, we have*

$$\text{wit}_{\tau \circ \sigma}(x_A, x_C) \simeq \sum_{x_B \in \mathcal{C}^0(B)} \text{wit}_\sigma(x_A, x_B) \times \text{wit}_\tau(x_B, x_C) \quad (4.3)$$

for all $x_A \in \mathcal{C}^0(A)$ and $x_C \in \mathcal{C}^0(B)$.

Proof. We construct the bijection by building functions in both directions.

Take $y \in \text{wit}_{\tau \circ \sigma}(x_A, x_C)$, i.e. $y \in \mathcal{C}^+(\tau \circ \sigma)$ s.t. $\partial_{\tau \circ \sigma}(y) = x_A \parallel x_C$. By Proposition 3.28, $y = x^\tau \circ x^\sigma$ s.t. $x^\sigma \in \mathcal{C}^+(\sigma)$ and $x^\tau \in \mathcal{C}^+(\tau)$ are causally compatible, and with

$$\partial_\sigma(x^\sigma) = x_A \parallel x_B \quad \partial_\tau(x^\tau) = x_B \parallel x_C$$

for some $x_B \in \mathcal{C}(B)$. But by Lemma 4.2 we have $\kappa_B(x_B) = 0$, so we return (x_B, x^σ, x^τ) .

Now, take $x_B \in \mathcal{C}^0(B)$, $x^\sigma \in \text{wit}_\sigma(x_A, x_B)$ and $x^\tau \in \text{wit}_\tau(x_B, x_C)$. By Lemma 4.1, x^σ and x^τ are causally compatible, so $x^\tau \circ x^\sigma \in \mathcal{C}^+(\tau \circ \sigma)$ s.t. $\partial_{\tau \circ \sigma}(x^\tau \circ x^\sigma) = x_A \parallel x_C$.

That these constructions are inverses of one another follows from Proposition 3.28. \square

This is a good starting point, which we must now extend to deal with symmetry.

4.1.2. *Witnesses as symmetry classes.* How shall we extend (4.3) to account for symmetry?

Since points of the web correspond to symmetry classes of configurations of the game, it would seem sensible for witnesses to be symmetry classes of configurations of the strategy:

$$\text{wit}_\sigma^{\cong}(x_A, x_B) = \{x^\sigma \in \mathcal{C}_\cong^+(\sigma) \mid \partial_\sigma(x^\sigma) = x_A^\sigma \parallel x_B^\sigma\}$$

for $\sigma : A \vdash B$, $x_A \in \mathcal{C}_\cong^0(A)$ and $x_B \in \mathcal{C}_\cong^0(B)$; where $\mathcal{C}_\cong^+(\sigma)$ is the set of symmetry classes of $+$ -covered configurations of σ ; and where $\partial_\sigma(x^\sigma) \in \mathcal{S}(A \vdash B)$ is well-defined since ∂_σ preserves symmetry. With this definition, are we able to extend the bijection (4.3)?


Consider first $y \in \text{wit}_{\tau \circ \sigma}^{\cong}(x_A, x_C)$. Take any representative $x^\tau \circ x^\sigma \in y$. The configurations x^σ and x^τ synchronize in $x_B^\sigma = x_B^\tau = x_B$, with symmetry class $x_B \in \mathcal{C}_\cong^0(B)$. So taking x^σ and x^τ the respective symmetry classes of x^σ and x^τ , we obtain

$$x_B \in \mathcal{C}_\cong^0(B), \quad x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B), \quad x^\tau \in \text{wit}_\tau^{\cong}(x_B, x_C)$$

as required. From the characterization of the symmetries of composition in Proposition 3.28, one deduces that this does not depend on the choice of the representative $x^\tau \circ x^\sigma \in y$.

What about the other direction? Consider now $x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B)$ and $x^\tau \in \text{wit}_\tau^{\cong}(x_B, x_C)$ for some $x_B \in \mathcal{C}_\cong^0(B)$, and pick representatives $x^\sigma \in x^\sigma$ and $x^\tau \in x^\tau$. Displaying them as

$$\partial_\sigma(x^\sigma) = x_A^\sigma \parallel x_B^\sigma \quad \partial_\tau(x^\tau) = x_B^\tau \parallel x_C^\tau,$$

Figure 10: Composition of σ and τ

we must compute their synchronization. But we may not have $x_B^\sigma = x_B^\tau$, indeed we only know that $x_B^\sigma, x_B^\tau \in \mathbf{x}_B$ so that there must be some $\theta : x_B^\sigma \cong_B x_B^\tau$. Fortunately, thin concurrent games come with tools to compute such synchronizations up to symmetry – in particular, at the heart of the proof of Proposition 3.32 is the following property:

Proposition 4.4. *Consider $\sigma : A \vdash B$ and $\tau : B \vdash C$ two strategies.*

For $x^\sigma \in \mathcal{C}^+(\sigma)$, $x^\tau \in \mathcal{C}^+(\tau)$ and $\theta : x_B^\sigma \cong_B x_B^\tau$ s.t. the composite bijection is secured:

$$x^\sigma \parallel x_C^\tau \stackrel{\partial_\sigma \parallel C}{\cong} x_A^\sigma \parallel x_B^\sigma \parallel x_C^\tau \stackrel{A \parallel \theta \parallel C}{\cong} x_A^\sigma \parallel x_B^\tau \parallel x_C^\tau \stackrel{A \parallel \partial_\tau^{-1}}{\cong} x_A^\sigma \parallel x^\tau,$$

then there are (necessarily unique) $y^\sigma \in \mathcal{C}^+(\sigma)$ and $y^\tau \in \mathcal{C}^+(\tau)$ causally compatible, and

$$\varphi^\sigma : x^\sigma \cong_\sigma y^\sigma, \quad \varphi^\tau : x^\tau \cong_\tau y^\tau,$$

such that we have $\varphi_A^\sigma \in \mathcal{S}_-(A)$, $\varphi_C^\tau \in \mathcal{S}_+(C)$, and $\varphi_B^\tau \circ \theta = \varphi_B^\sigma$.

See Appendix A.3. Intuitively, we play $\mathcal{S}(\sigma)$ and $\mathcal{S}(\tau)$ against each other. By \sim -receptivity and extension they adjust their copy indices interactively until they reach an agreement, *i.e.* pairs of configurations matching on the nose.

In the situation at hand the securedness assumption is automatic by Lemma 4.1, so that we get indeed $x^\sigma \cong_\sigma y^\sigma$ and $x^\tau \cong_\tau y^\tau$ matching on the nose, synchronizing to

$$y^\tau \in y^\sigma \in \mathcal{C}^+(\tau \circ \sigma)$$

whose symmetry class yields $y \in \text{wit}_{\tau \circ \sigma}^{\cong}(\mathbf{x}_A, \mathbf{x}_C)$ as needed.

Having given constructions in both directions, it might seem that we are essentially done, with a version of (4.2) essentially following the case without symmetry. But remember that above we started from witnesses $x^\sigma \in \text{wit}_\sigma^{\cong}(\mathbf{x}_A, \mathbf{x}_B)$ and $x^\tau \in \text{wit}_\tau^{\cong}(\mathbf{x}_B, \mathbf{x}_C)$, for which we first chose representatives $x^\sigma \in \mathcal{C}^+(\sigma)$, $x^\tau \in \mathcal{C}^+(\tau)$ and a symmetry $\theta : x_B^\sigma \cong_B x_B^\tau$. So one should not overlook the proof obligation that the construction is invariant under the choice of these representatives. Unfortunately, the symmetry class $y \in \text{wit}_{\tau \circ \sigma}^{\cong}(\mathbf{x}_A, \mathbf{x}_C)$ *does* depend on the choice of x^σ , x^τ , and θ – indeed, this was the error made in [CCPW18].

In fact, (4.2) does *not* hold for this notion of witness.

Example 4.5. Consider the arena α with one event \mathbf{q}^- , $\kappa_\alpha(\emptyset) = 1$ and $\kappa_\alpha(\{\mathbf{q}^-\}) = 0$, and

$$\sigma : !(!\alpha \multimap \alpha) \vdash !\alpha \multimap \alpha, \quad \tau : !(!\alpha \multimap \alpha)$$

two strategies as pictured in Figure 10. Their assignment of copy indices uses functions

$$f : \mathbb{N} \rightarrow \mathbb{N}, \quad g : \mathbb{N} \rightarrow \mathbb{N}, \quad h : \mathbb{N} \rightarrow \mathbb{N}, \quad k : \mathbb{N}^2 \rightarrow \mathbb{N},$$

whose precise identity has no impact on the discussion. We are interested in their composition, which unfolds as in Figure 10, yielding four pairwise conflicting, non-symmetric positive moves. Enriching **nPCF** with a type α with no constant, the substitution

$$(\lambda x^\alpha. f (f x)) [(\lambda y^\alpha. \mathbf{if} \text{ coin } y y)/f]$$

gives a perfect syntactic counterpart to the composition in Figure 10. Because there are two calls to the non-deterministic choice, this reduces to $\lambda x^\alpha. x$, but in four different ways. Observe that in the copy indices of each positive move of $\tau \odot \sigma$, one can read back the way the two non-deterministic choices were resolved: the upper row corresponds to the first call yielding $\mathbf{q}_{i,f(i)}^+$, the leftmost column to the second call yielding $\mathbf{q}_{i,f(i)}^+$, and so on.

From the composition, $\tau \odot \sigma$ has four witnesses for $\begin{pmatrix} \mathbf{q}^- \\ \vdots \\ \mathbf{q}^+ \end{pmatrix} \in \mathcal{C}_{\cong}^0(!\alpha \multimap \alpha)$, while

$$\sharp \text{wit}_{\sigma}^{\cong} \left(\begin{pmatrix} \mathbf{q}^- & \mathbf{q}^- & \mathbf{q}^- \\ \vdots & \vdots & \vdots \\ \mathbf{q}^+ & \mathbf{q}^+ & \mathbf{q}^+ \end{pmatrix} \right) = 1, \quad \sharp \text{wit}_{\tau}^{\cong} \left(\begin{pmatrix} \mathbf{q}^- & \mathbf{q}^- \\ \vdots & \vdots \\ \mathbf{q}^+ & \mathbf{q}^+ \end{pmatrix} \right) = 3,$$

as σ and τ cannot synchronize on any other symmetry class, this contradicts (4.2).

Indeed, up to symmetry, there are exactly *three* configurations of τ corresponding to two calls: (1) both choices may be resolved with $\mathbf{q}_{i,f(i)}$; (2) both choices may be resolved with $\mathbf{q}_{i,g(i)}$; and (3) we may have one of each. The point is that there is a symmetry

$$\begin{pmatrix} \mathbf{q}^- & \mathbf{q}^- \\ \vdots & \vdots \\ \mathbf{q}_{i,f(i)}^+ & \mathbf{q}_{j,g(j)}^+ \end{pmatrix} \cong_{!(\alpha \multimap \alpha)} \begin{pmatrix} \mathbf{q}^- & \mathbf{q}^- \\ \vdots & \vdots \\ \mathbf{q}_{j,f(j)}^+ & \mathbf{q}_{i,g(i)}^+ \end{pmatrix} \quad (4.4)$$

swapping the two calls, even though they do give rise to separate configurations in $\tau \odot \sigma$.

So, symmetry classes of $+$ -covered configurations are not the right witness: they count only once symmetry classes that should intuitively weight more, as they admit endo-symmetries that may affect the result. Indeed one can correct this accounting by appropriately weighting groups of endo-symmetries of symmetry classes – see Appendix C.1. This suggests links with generalized species of structures [FGHW08]; but our original question regarding what *concrete objects* the weighted relational model *counts* remains open.

4.1.3. *Concrete witnesses.* We now provide an alternative, more concrete notion of witness.

Intuitively, we must find a refinement of symmetry still letting us consider strategies up to their specific choice of copy indices, but nevertheless keeping the two configurations of (4.4) separate. Concretely, one may observe that the *swap* of (4.4) is only possible if Opponent changes their copy indices, exchanging i and j . Here we use a fundamental property of our setting: being *thin concurrent games* (Definition 3.4), arenas have sets of *positive* and *negative* symmetries $\mathcal{S}_+(A)$ and $\mathcal{S}_-(A)$; so we may consider configurations of strategies up to *positive* symmetry only. This has a very strong consequence:

Lemma 4.6. *Consider A a game, $\sigma : A$ a strategy on A , and $\theta \in \mathcal{S}(\sigma)$.*

If $\partial_\sigma(\theta) \in \mathcal{S}_+(A)$, then, $x = y$ and $\theta = \text{id}_x$.

Proof. A quite direct consequence of *thin*, which prevents Player from imposing symmetries not prompted by a prior Opponent exchange. See Lemma 3.28 in [CCW19]. \square




Figure 11: Witnesses for non-canonical representatives




Figure 12: Witnesses for canonical representatives

In other words, sub-groupoids of $\mathcal{S}(\sigma)$ mapping to positive symmetries of the game are all reduced to identities. Consider now fixed, for any arena A , the choice of a representative $\underline{x}_A \in \underline{x}_A$ for any $x_A \in \mathcal{C}_{\cong}^0(A)$. This invites the definition of **positive witnesses**:

$$\text{wit}_\sigma^+(\underline{x}_A, \underline{x}_B) = \{x^\sigma \in \mathcal{C}^+(\sigma) \mid x_A^\sigma \cong_A^- \underline{x}_A \ \& \ x_B^\sigma \cong_B^+ \underline{x}_B\}, \quad (4.5)$$

for $x_A \in \mathcal{C}_{\cong}^0(A)$ and $x_B \in \mathcal{C}_{\cong}^0(B)$, which will indeed turn out to be the right one. Notice the pleasant fact that this ranges over actual configurations of σ rather than symmetry classes³.

4.2. Representability. We introduce our last technical ingredient, *representability*.

4.2.1. Canonical representatives. The definition of wit^+ in (4.5) includes a significant subtlety: the dependency on the choice of the representatives $\underline{x}_A \in \underline{x}_A$, $\underline{x}_B \in \underline{x}_B$. Unfortunately, not only the set $\text{wit}_\sigma^+(\underline{x}_A, \underline{x}_B)$ depends on the choice of \underline{x}_A and \underline{x}_B , but even its *cardinality*:

Example 4.7. Consider $A = \text{!}(\text{!}\alpha \multimap \alpha)$, $B = \text{!}\alpha \multimap \alpha$, $\underline{x}_A = \begin{pmatrix} \mathbf{q}_0^- & \mathbf{q}_1^- \\ \vdots & \vdots \\ \mathbf{q}_0^+ & \mathbf{q}_1^+ \end{pmatrix}$ and $\underline{x}_B = \begin{pmatrix} \mathbf{q}_0^- \\ \vdots \\ \mathbf{q}_0^+ \end{pmatrix}$.

We show in Figures 11 and 12 the sets $\text{wit}_\sigma^+(\underline{x}_A, \underline{x}_B)$ for $\sigma : A$ the left hand side strategy in Figure 10, with the choice of representatives as shown. Up to positive symmetry, Player is free to associate \mathbf{q}_0^+ to either minimal event of \underline{x}_A . But as the symmetry is positive, the copy index of the subsequent Opponent move is forced by \underline{x}_A and this association. For Figure 12 the two choices make no difference as the bottom events of \underline{x}_A have the same index 0. In contrast, in Figure 11 these indices differ, and so yield distinct concrete witnesses.

To explain this mismatch, it is helpful to explicitly factor in the positive symmetries by

$$\sim^+ \text{wit}_\sigma(\underline{x}_A, \underline{x}_B) = \{(\theta_A, x^\sigma, \theta_B) \mid x^\sigma \in \mathcal{C}^+(\sigma), \theta_A : \underline{x}_A \cong_A^- x_A^\sigma, \theta_B : x_B^\sigma \cong_B^+ \underline{x}_B\} \quad (4.6)$$

the set of \sim^+ -witnesses, allowing us to prove the following property:

³An early sign that wit^+ is better behaved is that unlike wit^\cong , it does not depend on the choice of the symmetry for σ – recall from Section A.1.2 in [CCW19] that the symmetry is *not* unique.

Proposition 4.8. Consider A, B arenas, $\sigma : A \vdash B$, and $x_A \in \mathcal{C}_{\cong}^0(A), x_B \in \mathcal{C}_{\cong}^0(B)$.

Then, the cardinality of $\sim^{\pm}\text{-wit}_{\sigma}^+(x_A, x_B)$ does not depend on $\underline{x}_A \in x_A$ and $\underline{x}_B \in x_B$.

Proof. See Appendix A.5. □

For instance, in Figure 11, each positive witness admits exactly *one* pair of symmetries making it a \sim^+ -witness, whereas for Figure 12, the unique witness yields *two* \sim^+ -witnesses.

So which of Figure 11 and 12 is right, if any? Letting the weighted relational model be the judge, Figure 12 is better: indeed σ is the strategy of a pure λ -term to which the weighted relational model associates only weights 0 and 1. Tracking down the pathological behaviour in Figure 11, the crux of the issue is the inability, in the representative \underline{x}_A from Figure 11, to exchange the minimal moves while staying in \underline{x}_A . The only negative symmetry

$$\begin{array}{ccc} \mathbf{q}_0^- & \mathbf{q}_1^- & \cong_A^- \\ \vdots & \vdots & \\ \mathbf{q}_{0,0}^+ & \mathbf{q}_{1,1}^+ & \cong_A^+ \end{array} \begin{array}{ccc} \mathbf{q}_1^- & \mathbf{q}_0^- & \\ \vdots & \vdots & \\ \mathbf{q}_{1,0}^+ & \mathbf{q}_{0,1}^+ & \end{array}$$

(signified here by moves having the same position in the diagram) with domain \underline{x}_A exchanging \mathbf{q}_0^- and \mathbf{q}_1^- has codomain distinct from \underline{x}_A , intuitively causing the extra witness. In contrast, the analogous swap is an endosymmetry of the representative \underline{x}_A for Figure 12.

The next definition aims to capture the representatives for which wit^+ is well-behaved:

Definition 4.9. Consider A a game, and $x \in \mathcal{C}(A)$.

We say that x is **canonical** iff any $\theta : x \cong_A x$ factors uniquely as

$$x \cong_A^- x \cong_A^+ x,$$

with in particular x in the middle.

This definition extends a basic property of thin concurrent games:

Lemma 4.10. Consider A a thin concurrent game, and $\theta : x \cong_A y$ any symmetry.

Then, there exist unique $z \in \mathcal{C}(A)$, $\theta^- : x \cong_A z$ and $\theta^+ : z \cong_A y$ such that $\theta = \theta^+ \circ \theta^-$.

Proof. See Lemma 3.19 in [CCW19]. □

Likewise, any $\theta : x \cong_A y$ factors uniquely as $\theta^- \circ \theta^+$ for some $\theta^- \in \mathcal{S}_-(A)$, $\theta^+ \in \mathcal{S}_+(A)$.

The representative \underline{x}_A of Figure 12 is canonical, while that of Figure 11 is not. In our collapse, we will need to ensure that we compute witnesses only on canonical representatives.

4.2.2. Representability. This asks two questions: (1) does there always exist a canonical representative for any symmetry class?; and (2) is the cardinality of witnesses now invariant under the choice of a canonical representative?

For (1), for tcgs as in Definition 3.4 or games as in Definition 3.5, the answer is no – see Appendix B. For arenas as in Definition 3.20, we do not know. Likewise, though it is not hard to prove (2) for games arising from **PCF** types, we do not have an answer in general. So we must instead ask games to carry an explicit choice of canonical representatives:

Definition 4.11. A game A is **representable** when it comes equipped with a function

$$\underline{(-)}_A : \mathcal{C}_{\cong}^0(A) \rightarrow \mathcal{C}^0(A)$$

such that for all $x_A \in \mathcal{C}_{\cong}^0(A)$, $\underline{x}_A \in x_A$ is canonical.

$$\begin{array}{lll}
\underline{x}_A^+ = \underline{x}_A & \underline{(1, x)}_{A \& B} = (1, \underline{x}_A) & \underline{x} \parallel \underline{y}_{A \wp B} = \underline{x}_A \wp \underline{x}_B \\
\underline{x} \parallel \underline{y}_{A \otimes B} = \underline{x}_A \parallel \underline{x}_B & \underline{(2, x)}_{A \& B} = (2, \underline{x}_B) & \underline{x} \multimap \underline{y}_{A \multimap B} = \underline{x}_A \multimap \underline{y}_B
\end{array}$$

Figure 13: Representation functions for symmetry-free game constructions

This provides the condition left missing in Definition 3.5; but leaves us with the proof obligation of constructing the representation function for all game constructions. For basic games $1, \top, \mathbf{B}, \mathbf{N}, \alpha$ which have trivial symmetry, the representation function is obvious. For the game constructions only propagating symmetry (*i.e.* dual, tensor, par, with, and linear arrow), we set the representation function as specified in Figure 13 – it is direct that it preserves canonicity. Most importantly, for the bang construction, we set

$$[\underline{x}^1, \dots, \underline{x}^n]_{!A} = \parallel_{1 \leq i \leq n} \underline{x}_A^i \in \mathcal{C}^0(!A)$$

relying on Lemma 3.18, assuming chosen a sequential writing $[x_1, \dots, x_n]$ for every multiset.

This definition indeed always yield a canonical representative:

Lemma 4.12. *Consider A a representable $--$ game.*

Then $!A$, equipped with the function above, is representable.

Proof. We must show that for all $\underline{x} = [\underline{x}^1, \dots, \underline{x}^n] \in \mathcal{C}_{\cong}^0(!A)$, $\underline{x}_{!A}$ is canonical. Consider

$$\theta : \parallel_{1 \leq i \leq n} \underline{x}_A^i \cong_{!A} \parallel_{1 \leq i \leq n} \underline{x}_A^i$$

any symmetry. By definition, there is $\pi : \mathbb{N} \simeq \mathbb{N}$ a permutation, and a family $(\theta_i)_{i \in \mathbb{N}} \in \mathcal{S}(A)^{\mathbb{N}}$ s.t. for all $(i, a) \in \underline{x}_{!A}$, we have $\theta(i, a) = (\pi(i), \theta_i(a))$. But then, we have $\theta_i : \underline{x}_A^i \cong_A \underline{x}_A^{\pi(i)}$ which means that $\underline{x}^i = \underline{x}^{\pi(i)}$, so that $\underline{x}_A^i = \underline{x}_A^{\pi(i)}$ – hence θ_i is an endo-symmetry.

Now, we use that \underline{x}_A^i is canonical, which entails that θ_i factors as

$$\underline{x}_A^i \cong_A^- \underline{x}_A^i \cong_A^+ \underline{x}_A^i$$

using which we may finally factor θ as $\theta^+ \circ \theta^-$ where $\theta^+(i, a) = (i, \theta_{\pi^{-1}(i)}^+(a))$ and $\theta^-(i, a) = (\pi(i), \theta_i^-(a))$ – it is direct by definition that $\theta^+ : \underline{x}_{!A} \cong_{!A}^+ \underline{x}_{!A}$ and $\theta^- : \underline{x}_{!A} \cong_{!A}^- \underline{x}_{!A}$. \square

From now on, all games come equipped with a representation. As stated above, not every tcg admits a representation (see Appendix B for a counter-example). However, non-representable games seem to lie outside of the interpretation of any reasonable type.

4.3. Preservation of Composition. For A, B arenas and $\sigma : A \vdash B$, we may finally set:

$$(f(\sigma))_{\underline{x}_A, \underline{x}_B} = \sharp \text{wit}_{\sigma}^+(\underline{x}_A, \underline{x}_B), \quad (4.7)$$

for any $\underline{x}_A \in \mathcal{C}_{\cong}^0(A)$, $\underline{x}_B \in \mathcal{C}_{\cong}^0(B)$, and where $\text{wit}_{\sigma}^+(\underline{x}_A, \underline{x}_B)$ is defined as in (4.5) using the canonical representatives. Our aim is to prove the following equality:

$$(f(\tau \circ \sigma))_{\underline{x}_A, \underline{x}_C} = \sum_{\underline{x}_B \in \mathcal{C}_{\cong}^0(B)} (f(\sigma))_{\underline{x}_A, \underline{x}_B} \times (f(\tau))_{\underline{x}_B, \underline{x}_C},$$

and the natural route seems to be by setting up a bijection

$$\text{wit}_{\tau \circ \sigma}^+(\underline{x}_A, \underline{x}_C) \simeq \sum_{\underline{x}_B \in \mathcal{C}_{\cong}^0(B)} \text{wit}_{\sigma}^+(\underline{x}_A, \underline{x}_B) \times \text{wit}_{\tau}^+(\underline{x}_B, \underline{x}_C). \quad (4.8)$$

However, while it must hold, there does not seem to be any simple way of constructing this bijection explicitly. We use an indirect route, considering witnesses with symmetry.

4.3.1. Interaction witnesses with symmetry. In (4.8) above, going from right to left is problematic as we get triples (x_B, x^σ, x^τ) where, in general, there is no reason to have $x_B^\sigma = x_B^\tau$. We do have $x_B^\sigma \cong_B^+ x_B$ and $x_B \cong_B^- x_B^\tau$, but with no specified symmetries.

So to approach (4.8), we shall start by studying synchronizable pairs of

$$(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^\pm \text{wit}_\sigma(x_A, x_B), \quad (\Omega_B^-, x^\tau, \Omega_C^+) \in \sim^\pm \text{wit}_\tau(x_B, x_C),$$

witnesses with symmetry as in (4.6). So we have $x^\sigma \in \mathcal{C}^+(\sigma)$, $x^\tau \in \mathcal{C}^+(\tau)$ and symmetries

$$x_A \xleftarrow{\theta_A^-} x_A^\sigma \quad x_B^\sigma \xrightarrow{\theta_B^+} x_B \xleftarrow{\Omega_B^-} x_B^\tau \quad x_C^\tau \xrightarrow{\Omega_C^+} x_C$$

allowing us to synchronize x^σ and x^τ using Proposition 4.4.

We shall compare those with witnesses in the composition, starting by defining:

Definition 4.13. Consider A, B and C arenas; $\sigma \in \text{Strat}(A, B)$ and $\tau \in \text{Strat}(B, C)$ strategies; and $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$.

The **interaction witnesses** on x_A, x_B, x_C is the set of all $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$ s.t.

$$x_A^\sigma \cong_A^- x_A, \quad x_B^\sigma = x_B^\tau \in x_B, \quad x_C^\tau \cong_C^+ x_C,$$

we write $\text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C)$ for this set. Likewise, the **\sim^+ -interaction witnesses** comprise

$$\theta_A^- : x_A^\sigma \cong_A^- x_A, \quad x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma), \quad \theta_C^+ : x_C^\tau \cong_C^+ x_C,$$

with $x_B^\sigma = x_B^\tau \in x_B$. We write $\sim^\pm \text{wit}_{\sigma, \tau}(x_A, x_B, x_C)$ for this set.

Note that we obviously have, for any $x_A \in \mathcal{C}_{\cong}^0(A)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$

$$\text{wit}_{\tau \odot \sigma}^+(x_A, x_C) \simeq \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} \text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C) \quad (4.9)$$

as interaction witnesses are exactly witnesses of the composition with a specified symmetry class $x_B \in \mathcal{C}_{\cong}^0(B)$ in the middle. To establish (4.8), we shall now study a connection between \sim^+ -interaction witnesses and synchronizable pairs of \sim^+ -witnesses.

4.3.2. Synchronization up to symmetry. The property we shall prove is a quantitative elaboration on Proposition 4.4, so we start with an explicit reformulation of Proposition 4.4.

Lemma 4.14. Consider A, B, C arenas, $\sigma \in \text{Strat}(A, B)$ and $\tau \in \text{Strat}(B, C)$.

Then, for any $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$, for any pair of \sim^+ -witnesses

$$(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^\pm \text{wit}_\sigma^+(x_A, x_B), \quad (\Omega_B^-, x^\tau, \Omega_C^+) \in \sim^\pm \text{wit}_\tau^+(x_B, x_C),$$

there are unique $\omega^\sigma : x^\sigma \cong_\sigma y^\sigma$, $\nu^\tau : x^\tau \cong_\tau y^\tau$, $\Theta_B : x_B \cong_B y_B$ and \sim^+ -interaction witness

$$(\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+) \in \sim^\pm \text{wit}_{\sigma, \tau}(x_A, x_B, x_C)$$

with $y_B^\sigma = y_B^\tau = y_B$, such that the following diagrams commute:

$$\begin{array}{ccccc}
& \theta_A^- & x_A^\sigma & & x_B^\sigma \xrightarrow{\theta_B^+} \underline{x}_B \xleftarrow{\Omega_B^-} x_B^\tau & & x_C^\tau & \Omega_C^+ \\
& \swarrow & \downarrow \omega_A^\sigma & \omega_B^\sigma \downarrow & \downarrow \Theta_B & \downarrow \nu_B^\tau & \nu_C^\tau \downarrow & \searrow \\
\underline{x}_A & & y_A^\sigma & & y_B^\sigma = y_B = y_B^\tau & & y_C^\tau & \psi_C^+ \\
& \swarrow & \downarrow \psi_A^- & & & & \downarrow \psi_C^+ & \searrow \\
& & & & & & & \underline{x}_C
\end{array}$$

Proof. Existence. First, by Lemma 4.1, the bijection induced by $(\Omega_B^-)^{-1} \circ \theta_B^+ : x_B^\sigma \cong_B x_B^\tau$ is secured. Thus we can apply Proposition 4.4, yielding $y^\tau \odot y^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$ along with

$$\omega^\sigma : y^\sigma \cong_\sigma x^\sigma, \quad \nu^\tau : y^\tau \cong_\tau y^\tau,$$

such that $\nu_B^\tau \circ ((\Omega_B^-)^{-1} \circ \theta_B^+) = \omega_B^\sigma$. Furthermore we may set $\psi_A^- = \theta_A^- \circ (\omega_A^\sigma)^{-1}$ and $\psi_C^+ = \Omega_C^+ \circ (\nu_C^\tau)^{-1}$ – overall, the following diagrams commute

$$\begin{array}{ccccc}
& \theta_A^- & x_A^\sigma & & x_B^\sigma \xrightarrow{\theta_B^+} \underline{x}_B \xrightarrow{(\Omega_B^-)^{-1}} x_B^\tau & & x_C^\tau & \Omega_C^+ \\
& \swarrow & \downarrow \omega_A^\sigma & \omega_B^\sigma \downarrow & \downarrow \nu_B^\tau & & \nu_C^\tau \downarrow & \searrow \\
\underline{x}_A & & y_A^\sigma & & y_B^\sigma = y_B = y_B^\tau & & y_C^\tau & \psi_C^+ \\
& \swarrow & \downarrow \psi_A^- & & & & \downarrow \psi_C^+ & \searrow \\
& & & & & & & \underline{x}_C
\end{array}$$

which we complete by setting $\Theta_B : \underline{x}_B \rightarrow y_B$ as either path around the center diagram.

Uniqueness. First, $y^\tau \odot y^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$, ω^σ, ν^τ are unique by the uniqueness clause in Proposition 4.4. It follows that ψ_A^- and ψ_C^+ and Θ_B are determined by the diagram. \square

In particular, to $(\theta_A^-, x^\sigma, \theta_B^+) \in \mathcal{W}_\sigma^+(\underline{x}_A, \underline{x}_B)$ and $(\Omega_B^-, x^\tau, \Omega_C^+) \in \mathcal{W}_\tau^+(\underline{x}_B, \underline{x}_C)$ we have associated a \sim^+ -interaction witness $(\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+) \in \mathcal{W}_{\sigma, \tau}^+(\underline{x}_A, \underline{x}_B, \underline{x}_C)$. Counting-wise, it seems \sim^+ -interaction witnesses have fewer degrees of liberty than pairs of \sim^+ -witnesses as the latter have no symmetry on B ; in the lemma above this is mitigated by the fact that the construction also extracts Θ_B . The next step is to reverse this: from

$$(\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+) \in \mathcal{W}_{\sigma, \tau}^+(\underline{x}_A, \underline{x}_B, \underline{x}_C)$$

and $\Theta_B : \underline{x}_B \cong_B y_B$, we must make Θ_B act on y^σ and y^τ to recover the original x^σ and x^τ .

4.3.3. Negative symmetries acting on strategies. In thin concurrent games, strategies can always adjust their copy indices to match a change in Opponent's copy indices:

Lemma 4.15. *Consider A a game, $\sigma : A$ a strategy, $x^\sigma \in \mathcal{C}(\sigma)$ and $\theta^- : x_A^\sigma \cong_A^- y_A$.*

Then, there are unique $\varphi : x^\sigma \cong_\sigma y^\sigma$ and $\theta^+ : y_A \cong_A^+ y_A^\sigma$ such that

$$\partial_\sigma \varphi = \theta^+ \circ \theta^- : x_A^\sigma \cong_A y_A^\sigma.$$

Proof. See Lemma B.4 in [CCW19]. \square

Opponent changes their copy indices by applying the negative symmetry $\theta^- : x_A^\sigma \cong_A^- y_A$, and Player adapts by applying the unique $\varphi : x^\sigma \cong_\sigma y^\sigma$. The resulting configuration y_A^σ might not be equal to y_A , but it is *positively symmetric*, reflecting Player's adjusted indices.

4.3.4. *Symmetries acting on \sim^+ -interaction witnesses.* We use this to reverse Lemma 4.14.

Lemma 4.16. *Consider A, B, C arenas, $\sigma \in \text{Strat}(A, B)$ and $\tau \in \text{Strat}(B, C)$.*

Then, for any $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$, \sim^+ -interaction witness

$$(\psi_A^-, y^\tau \circ y^\sigma, \psi_C^+) \in \sim^{\pm}\text{-wit}_{\sigma, \tau}(x_A, x_B, x_C)$$

with $y_B^\sigma = y_B^\tau = y_B$, any $\Theta_B : x_B \cong_B y_B$, there are unique $\omega^\sigma : x^\sigma \cong_\sigma y^\sigma$, $\nu^\tau : x^\tau \cong_\tau y^\tau$ and

$$(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^{\pm}\text{-wit}_\sigma^+(x_A, x_B), \quad (\Omega_B^-, x^\tau, \Omega_C^+) \in \sim^{\pm}\text{-wit}_\tau^+(x_B, x_C),$$

a pair of \sim^+ -witnesses, such that the following diagram commutes:

$$\begin{array}{ccccc} & & x_B^\sigma & \xrightarrow{\theta_B^+} & x_B & \xleftarrow{\Omega_B^-} & x_B^\tau & & & & x_C^\tau & & & \Omega_C^+ \\ & \theta_A^- & \downarrow \omega_A^\sigma & & \downarrow \omega_B^\sigma & & \downarrow \nu_B^\tau & & & & \downarrow \nu_C^\tau & & & \downarrow \psi_C^+ \\ x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & = & y_B & = & y_B^\tau & & y_C^\tau & \xrightarrow{\psi_C^+} & x_C & \\ & & \downarrow \omega_A^\sigma & & \downarrow \Theta_B & & \downarrow \nu_B^\tau & & & & & & & \end{array}$$

Proof. The first step is to factor Θ_B^{-1} in two ways, as in the diagram

$$\begin{array}{ccccccc} & & z_B^1 & \xrightarrow{\Phi_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\ & & \uparrow \Phi_B^- & & \uparrow \Theta_B^{-1} & & \uparrow \Psi_B^+ \\ x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & = & y_B^\tau & \xrightarrow{\psi_C^+} & x_C \\ & & \downarrow \omega_A^\sigma & & \downarrow \omega_B^\sigma & & \downarrow \nu_B^\tau & & \end{array}$$

following Lemma 4.10. By Lemma 4.15 we can make Φ_B^- act on x^σ . This yields

$$\lambda_A^- : x_A^\sigma \cong_A^- y_A^\sigma, \quad \omega^\sigma : x^\sigma \cong_\sigma y^\sigma, \quad \Delta_B^+ : x_B^\sigma \cong_B^+ z_B^1,$$

unique such that the following diagram commutes:

$$\begin{array}{ccccccc} & & z_B^1 & \xrightarrow{\Phi_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\ & & \uparrow \Phi_B^- & & \uparrow \Theta_B^{-1} & & \uparrow \Psi_B^+ \\ x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & = & y_B^\tau & \xrightarrow{\psi_C^+} & x_C \\ & & \downarrow \omega_A^\sigma & & \downarrow \omega_B^\sigma & & \downarrow \nu_B^\tau & & \end{array}$$

leaving in grey the irrelevant parts of the full diagram for context. Setting $\theta_A^- = \psi_A^- \circ \lambda_A^-$ and $\theta_B^+ = \Phi_B^+ \circ \Delta_B^+$, we have found data making the following diagram commute:

$$\begin{array}{ccccccc} & & z_B^1 & \xrightarrow{\Phi_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\ & & \uparrow \Phi_B^- & & \uparrow \Theta_B^{-1} & & \uparrow \Psi_B^+ \\ x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & = & y_B^\tau & \xrightarrow{\psi_C^+} & x_C \\ & & \downarrow \omega_A^\sigma & & \downarrow \omega_B^\sigma & & \downarrow \nu_B^\tau & & \end{array}$$

We shall now prove uniqueness of this data. Assume that we have other symmetries $\gamma_A^- : u_A^\sigma \cong_A^- x_A$, $\varpi^\sigma : u^\sigma \cong_S y^\sigma$ and $\gamma_B^+ : u_B^\sigma \cong_B^+ x_B$ making the following diagram commute:

$$\begin{array}{ccccccc}
& & \gamma_A^- & u_A^\sigma & u_B^\sigma & \xrightarrow{\gamma_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\
& & \swarrow & \downarrow \varpi_A^\sigma & \downarrow \varpi_B^\sigma & & \downarrow \Theta_B & \nearrow \Psi_B^+ & \\
x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & \xlongequal{\quad} & y_B & \xlongequal{\quad} & y_C^\tau & \xrightarrow{\psi_C^+} & x_C
\end{array}$$

Then, it follows that the following diagram also commutes:

$$\begin{array}{ccccccc}
& & (\psi_A^-)^{-1} \circ \gamma_A^- & u_A^\sigma & u_B^\sigma & \xrightarrow{(\Phi_B^+)^{-1} \circ \gamma_B^+} & z_B^1 & \xrightarrow{\Phi_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\
& & \swarrow & \downarrow \varpi_A^\sigma & \downarrow \varpi_B^\sigma & & \downarrow \Phi_B^- & \nearrow \Theta_B^{-1} & \nearrow \Psi_B^+ & & \\
x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & \xlongequal{\quad} & y_B & \xlongequal{\quad} & y_C^\tau & \xrightarrow{\psi_C^+} & x_C
\end{array}$$

By uniqueness for Lemma 4.15, it follows that $u^\sigma = x^\sigma$, $\omega^\sigma = \varpi^\sigma$, $\lambda_A^- = (\psi_A^-)^{-1} \circ \gamma_A^-$ so $\gamma_A^- = \theta_A^-$, and $(\Phi_B^+)^{-1} \circ \gamma_B^+ = \Delta_B^+$ so $\gamma_B^+ = \theta_B^+$. Altogether, we have proved that there are

$$\theta_A^- : x_A^\sigma \cong_\sigma x_A, \quad \omega^\sigma : x^\sigma \cong_\sigma y^\sigma, \quad \theta_B^+ : x_B^\sigma \cong_B^+ x_B,$$

unique making the following diagram commute:

$$\begin{array}{ccccccc}
& & \theta_A^- & x_A^\sigma & x_B^\sigma & \xrightarrow{\theta_B^+} & x_B & \xleftarrow{\Psi_B^-} & z_B^2 \\
& & \swarrow & \downarrow \omega_A^\sigma & \downarrow \omega_B^\sigma & & \downarrow \Theta_B & \nearrow \Psi_B^+ & \\
x_A & \xleftarrow{\psi_A^-} & y_A^\sigma & & y_B^\sigma & \xlongequal{\quad} & y_B & \xlongequal{\quad} & y_C^\tau & \xrightarrow{\psi_C^+} & x_C
\end{array}$$

The lemma follows by performing the exact same reasoning on the right hand side. \square

4.3.5. *The interaction bijection.* If B is a game and $x_B \in \mathcal{C}_{\cong}^0(B)$, let us write $\mathcal{S}(x_B)$ for the set of endosymmetries on x_B , i.e. symmetries $\theta_B : x_B \cong_B x_B$. We shall use the fact that for any $x, y \in x_B$, there are exactly as many symmetries $x \cong_B y$ as in $\mathcal{S}(x_B)$. Indeed let us fix, for any $x \in x_B$, a symmetry $\kappa_x : x \cong_B x_B$. For any $x, y \in x_B$, we then have

$$\begin{aligned}
(-)[x, y] : \mathcal{S}(x_B) &\rightarrow \mathcal{S}(B)(x, y) \\
\theta &\mapsto \kappa_y^{-1} \circ \theta \circ \kappa_x
\end{aligned}$$

writing $\mathcal{S}(B)(x, y)$ for the set of all $\theta : x \cong_B y$. It is elementary that this is a bijection.

Using this, we finally have, for any A, B, C games and $\sigma \in \text{Strat}(A, B)$, $\tau \in \text{Strat}(B, C)$:

Corollary 4.17. *Fix $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$. Then, there is a bijection*

$$\Upsilon : \sim^\pm\text{-wit}_\sigma(x_A, x_B) \times \sim^\pm\text{-wit}_\tau(x_B, x_C) \simeq \sim^\pm\text{-wit}_{\sigma, \tau}(x_A, x_B, x_C) \times \mathcal{S}(x_B)$$

such that for any $\Upsilon(x^\sigma, x^\tau) = (y^\tau \odot y^\sigma, \Theta)$, we have $x^\sigma \cong_\sigma y^\sigma$ and $x^\tau \cong_\tau y^\tau$.

Proof. Given $(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^\pm\text{-wit}_\sigma(x_A, x_B)$ and $(\Omega_B^-, x^\tau, \Omega_C^+) \in \sim^\pm\text{-wit}_\tau(x_B, x_C)$, we get

$$(\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+) \in \sim^\pm\text{-wit}_{\sigma, \tau}(x_A, x_B, x_C)$$

and $\Theta_B : x_B \cong_B y_B$ by Lemma 4.14; so we set $\Upsilon((\theta_A^-, x^\sigma, \theta_B^+), (\Omega_B^-, x^\tau, \Omega_C^+))$ as:

$$((\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+), (\kappa_{y_B} \circ \Theta_B)).$$

Reciprocally, given $(\psi_A^-, y^\tau \odot y^\sigma, \psi_C^+) \in \sim^\pm \text{wit}_{\sigma, \tau}(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C)$ and $\Xi_B \in \mathcal{S}(\mathbf{x}_B)$, we set $\Theta_B = \kappa_{y_B}^{-1} \circ \Xi_B$ and apply Lemma 4.16 to get back two \sim^+ -witnesses:

$$\begin{aligned} (\theta_A^-, x^\sigma, \theta_B^+) &\in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A, \mathbf{x}_B) \\ (\Omega_B^-, x^\tau, \Omega_C^+) &\in \sim^\pm \text{wit}_\tau(\mathbf{x}_B, \mathbf{x}_C) \end{aligned}$$

That these constructions are inverses of each other is an immediate consequence from the uniqueness properties in Lemmas 4.14 and 4.16. \square

4.3.6. Preservation of composition. Extending earlier notations, for any $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$, we write $\mathcal{S}_+(\mathbf{x}_A)$ for the group of positive endo-symmetries on $\underline{\mathbf{x}}_A$, and likewise for $\mathcal{S}_-(\mathbf{x}_A)$. As for general symmetries, if $x, y \in \mathbf{x}_A$ s.t. $x \cong_A^+ \underline{\mathbf{x}}_A$ and $y \cong_A^+ \underline{\mathbf{x}}_A$, then there is a bijection

$$\begin{aligned} (-)[x, y] : \mathcal{S}_+(\mathbf{x}_A) &\simeq \mathcal{S}_+(A)(x, y) \\ \theta^+ &\mapsto (\kappa_y^+)^{-1} \circ \theta^+ \circ \kappa_x^+ \end{aligned}$$

for $\mathcal{S}_+(A)(x, y)$ the set of $\theta^+ : x \cong_A^+ y$; and having chosen a positive symmetry $\kappa_x^+ : x \cong_A^+ \underline{\mathbf{x}}_A$ for all x positively symmetric to $\underline{\mathbf{x}}_A$ – the same hold for negative symmetries. Thus:

Lemma 4.18. *Consider strategies $\sigma \in \text{Strat}(A, B)$, $\tau \in \text{Strat}(B, C)$, and symmetry classes $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$, $\mathbf{x}_B \in \mathcal{C}_{\cong}^0(B)$ and $\mathbf{x}_C \in \mathcal{C}_{\cong}^0(C)$. Then we have bijections*

$$\begin{aligned} \Psi &: \sim^\pm \text{wit}_\sigma(\mathbf{x}_A, \mathbf{x}_B) \simeq \mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}_+(\mathbf{x}_B) \times \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B) \\ \Xi &: \sim^\pm \text{wit}_{\sigma, \tau}(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) \simeq \mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}_+(\mathbf{x}_C) \times \text{wit}_{\sigma, \tau}^+(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) \end{aligned}$$

s.t. for all $(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A, \mathbf{x}_B)$, writing $\Psi(\theta_A^-, x^\sigma, \theta_B^+) = (\psi_A^-, \psi_B^+, y^\sigma)$, $x^\sigma = y^\sigma$; and likewise, writing $\Xi(\theta_A^-, x^\tau \odot x^\sigma, \theta_C^+) = (\psi_A^-, \psi_C^+, y^\tau \odot y^\sigma)$, then $x^\sigma = y^\sigma$ and $x^\tau = y^\tau$.

Proof. To $(\theta_A^-, x^\sigma, \theta_B^+) \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A, \mathbf{x}_B)$, simply associate

$$\Psi(\theta_A^-, x^\sigma, \theta_B^+) = (\theta_A^- \circ (\kappa_{x_A}^-)^{-1}, \theta_B^+ \circ (\kappa_{x_B}^+)^{-1}, x^\sigma) \in \mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}_+(\mathbf{x}_B) \times \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B),$$

it is straightforward that this is a bijection. The proof for Ξ is the same. \square

We now compose these bijections, to obtain:

Lemma 4.19. *Consider strategies $\sigma \in \text{Strat}(A, B)$, $\tau \in \text{Strat}(B, C)$, and symmetry classes $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$, $\mathbf{x}_B \in \mathcal{C}_{\cong}^0(B)$ and $\mathbf{x}_C \in \mathcal{C}_{\cong}^0(C)$. Then we have a bijection:*

$$\begin{aligned} \Phi &: \mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}(\mathbf{x}_B) \times \mathcal{S}_+(\mathbf{x}_C) \times \text{wit}_{\sigma, \tau}^+(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) \\ &\simeq \mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}(\mathbf{x}_B) \times \mathcal{S}_+(\mathbf{x}_C) \times \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B) \times \text{wit}_\tau^+(\mathbf{x}_B, \mathbf{x}_C) \end{aligned}$$

such that writing $\Phi(\theta_A^-, \theta_B, \theta_C^+, x^\tau \odot x^\sigma) = (\varphi_A^-, \varphi_B, \varphi_C^+, y^\sigma, y^\tau)$, $x^\sigma \cong_\sigma y^\sigma$ and $x^\tau \cong_\tau y^\tau$.

Proof. The bijection is obtained through the following composition:

$$\begin{aligned} &\mathcal{S}_-(\mathbf{x}_A) \times \mathcal{S}(\mathbf{x}_B) \times \mathcal{S}_+(\mathbf{x}_C) \times \text{wit}_{\sigma, \tau}^+(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) \\ &\simeq \mathcal{S}(\mathbf{x}_B) \times \sim^\pm \text{wit}_{\sigma, \tau}(\mathbf{x}_A, \mathbf{x}_B, \mathbf{x}_C) \\ &\simeq \sim^\pm \text{wit}_\sigma(\mathbf{x}_A, \mathbf{x}_B) \times \sim^\pm \text{wit}_\tau(\mathbf{x}_B, \mathbf{x}_C) \\ &\simeq \mathcal{S}_-(\mathbf{x}_A) \times \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B) \times \mathcal{S}_+(\mathbf{x}_B) \times \mathcal{S}_-(\mathbf{x}_B) \times \text{wit}_\tau^+(\mathbf{x}_B, \mathbf{x}_C) \times \mathcal{S}_+(\mathbf{x}_C) \\ &\simeq \mathcal{S}_-(\mathbf{x}_B) \times \mathcal{S}(\mathbf{x}_B) \times \mathcal{S}_+(\mathbf{x}_C) \times \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B) \times \text{wit}_\tau^+(\mathbf{x}_B, \mathbf{x}_C). \end{aligned}$$

using Ξ in Lemma 4.18, then Υ^{-1} in Corollary 4.17, then Ψ from Lemma 4.18 for σ and τ , and finally the bijection $\mathcal{S}(\mathbf{x}_B) \simeq \mathcal{S}_+(\mathbf{x}_B) \times \mathcal{S}_-(\mathbf{x}_B)$ coming from the canonicity of $\underline{\mathbf{x}}_B$. That Φ preserves symmetry classes in σ and τ is an immediate verification. \square

From Φ , it immediately follows for $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$ we have

$$\sharp\text{wit}_{\sigma,\tau}^+(x_A, x_B, x_C) = \sharp\text{wit}_{\sigma}^+(x_A, x_B) \times \sharp\text{wit}_{\tau}^+(x_B, x_C), \quad (4.10)$$

however there is no clear way to realize the corresponding bijection directly, without invoking symmetries. Of course the bijection must exist for cardinality reasons, but then it may not preserve symmetry classes in σ and τ – which is necessary for the quantitative generalization in Section 6. Nevertheless, (4.10) allows us to conclude the core result of the paper:

Corollary 4.20. *Consider $\sigma \in \text{Strat}(A, B)$ and $\tau \in \text{Strat}(B, C)$. Then,*

$$f(\tau \odot \sigma)_{x_A, x_C} = \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} f(\sigma)_{x_A, x_B} \times f(\tau)_{x_B, x_C}$$

for all $x_A \in \mathcal{C}_{\cong}^0(A)$ and $x_C \in \mathcal{C}_{\cong}^0(C)$.

Proof. We perform the following direct computation, using (4.9) and (4.10).

$$\begin{aligned} \sharp\text{wit}_{\tau \odot \sigma}^+(x_A, x_C) &= \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} \sharp\text{wit}_{\sigma,\tau}^+(x_A, x_B, x_C) \\ &= \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} \sharp\text{wit}_{\sigma}^+(x_A, x_B) \times \text{wit}_{\tau}^+(x_B, x_C). \quad \square \end{aligned}$$

5. PRESERVATION OF THE INTERPRETATION

Now that preservation of composition is clear, we deal with the rest of the interpretation.

5.1. Structure-preserving functors. We first set up the categorical machinery.

5.1.1. Cartesian closed functors. We start with cartesian closed functors, the appropriate notion of morphisms between cartesian closed categories, preserving the interpretation of the simply-typed λ -calculus. This can be straightforwardly adapted to \sim -categories.

Definition 5.1. Let \mathcal{C}, \mathcal{D} be cartesian closed \sim -categories. A \sim -functor

$$F : \mathcal{C} \rightarrow \mathcal{D}$$

is **cartesian closed** if it comes equipped with for any $A, B \in \mathcal{C}_0$, maps

$$\begin{aligned} k^\top &: \top \rightarrow F\top \\ k_{A,B}^\& &: FA \& FB \rightarrow F(A \& B) \\ k_{A,B}^\Rightarrow &: FA \Rightarrow FB \rightarrow F(A \Rightarrow B) \end{aligned}$$

invertible up to \sim ; and such that the following diagrams commute up to \sim :

$$\begin{array}{ccc} & FA & \\ \pi_1 \nearrow & & \nwarrow F(\pi_1) \\ FA \& FB & \xrightarrow{k_{A,B}^\&} F(A \& B) \\ \pi_2 \searrow & & \nearrow F(\pi_2) \\ & FB & \end{array} \qquad \begin{array}{ccc} & & F(A \Rightarrow B) \& FA \xrightarrow{k_{A \Rightarrow B, A}^\&} F((A \Rightarrow B) \& A) \\ & & \uparrow k_{A, B \& FA}^\Rightarrow \\ (FA \Rightarrow FB) \& FA & \xrightarrow{\text{ev}_{FA, FB}} FB \end{array} \begin{array}{c} \downarrow F(\text{ev}_{A, B}) \end{array}$$

We use notions of cartesian closed categories with explicit structure, which must be preserved up to isomorphism. It is not necessary to require that $t^{\&}$ and t^{\rightarrow} are natural (up to \sim); this automatically follows. Likewise, preservation of projections and evaluation suffice to ensure that pairing and currying are also preserved.

In fact, cartesian closed \sim -functors ensure preservation up to isomorphism of the interpretation of the simply-typed λ -calculus, in the following sense: assume chosen

$$k^\alpha : \llbracket \alpha \rrbracket_{\mathcal{D}} \rightarrow F(\llbracket \alpha \rrbracket_{\mathcal{C}})$$

an isomorphism for any base type α . Then, by induction on types one can form isos

$$\begin{aligned} k_A^{\text{ty}} &: \llbracket A \rrbracket_{\mathcal{D}} \rightarrow F(\llbracket A \rrbracket_{\mathcal{C}}) \\ k_\Gamma^{\text{ctx}} &: \llbracket \Gamma \rrbracket_{\mathcal{D}} \rightarrow F(\llbracket \Gamma \rrbracket_{\mathcal{D}}) \end{aligned}$$

in \mathcal{D} for every type A and context Γ ; it is then a lengthy exercise to prove that

$$\begin{array}{ccc} \llbracket \Gamma \rrbracket_{\mathcal{D}} & \xrightarrow{\llbracket M \rrbracket_{\mathcal{D}}} & \llbracket A \rrbracket_{\mathcal{C}} \\ k_\Gamma^{\text{ctx}} \downarrow & \sim & \downarrow k_A^{\text{ty}} \\ F(\llbracket \Gamma \rrbracket_{\mathcal{C}}) & \xrightarrow{F(\llbracket M \rrbracket_{\mathcal{C}})} & F(\llbracket A \rrbracket_{\mathcal{C}}) \end{array}$$

for every simply-typed λ -term $\Gamma \vdash M : A$; *i.e.* F preserves the interpretation up to iso.

5.1.2. Relative Seely \sim -functors.

Definition 5.2. Let \mathcal{C}, \mathcal{D} be relative Seely \sim -categories. A **relative Seely (\sim -)functor** $\mathcal{C} \rightarrow \mathcal{D}$ is a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ which restricts to $F : \mathcal{C}_s \rightarrow \mathcal{D}_s$, equipped with:

- For every $A, B \in \mathcal{C}$, morphisms

$$\begin{aligned} t_{A,B}^{\otimes} &: FA \otimes FB \rightarrow F(A \otimes B) \\ t^1 &: 1 \rightarrow F1; \end{aligned}$$

making (F, t^{\otimes}, t^1) a symmetric monoidal functor $(\mathcal{C}, \otimes, 1) \rightarrow (\mathcal{D}, \otimes, 1)$;

- for every $S, T \in \mathcal{C}_s$, morphisms

$$\begin{aligned} t_{S,T}^{\&} &: FS \& FT \rightarrow F(S \& T) \\ t^\top &: \top \rightarrow F\top; \end{aligned}$$

- For every $A \in \mathcal{C}$ and $S \in \mathcal{C}_s$, a morphism

$$t_{A,S}^{\circ} : FA \multimap FS \rightarrow F(A \multimap S);$$

- For every $S \in \mathcal{C}_s$, a morphism

$$t_S^! : !FS \rightarrow F!S;$$

all of which are invertible up to \sim and satisfy the coherence axioms of Figure 14 up to \sim , such that for every $S, T \in \mathcal{C}_s$ and $f : !S \rightarrow T$, the diagram

$$\begin{array}{ccc} !FS & \xrightarrow{(Ff \circ t_S^!)^!} & !FT \\ t_S^! \downarrow & & \downarrow t_T^! \\ F!S & \xrightarrow{F(f^!)} & F(!T) \end{array} \quad (5.1)$$

commutes up to \sim .




Figure 14: Coherence diagrams for relative Seely functors

In this paper, we only use the following property of relative Seely functors:

Proposition 5.3. *A relative Seely \sim -functor $F : \mathcal{C} \rightarrow \mathcal{D}$ induces a cartesian closed \sim -functor $F_! : \mathcal{C}_! \rightarrow \mathcal{D}_!$ defined by $F_!(S) = F(S)$ for all $S \in \mathcal{C}_!$,*

$$F_!(f) = (Ff) \circ t_S^! \in \mathcal{D}_!(FS, FT)$$

for all $f \in \mathcal{C}_!(S, T)$, and equipped with the following structural isomorphisms:

$$\begin{aligned} k^\top &= t^\top \circ \epsilon_\top && \in \mathcal{D}_!(\top, F\top) \\ k_{S,T}^\& = t_{S,T}^\& \circ \epsilon_{FS \& FT} && \in \mathcal{D}_!(FS \& FT, F(S \& T)) \\ k_{S,T}^{\Rightarrow} &= t_{FS,T}^{\Rightarrow} \circ ((t_S^!)^{-1} \multimap FT) \circ \epsilon_{FS \Rightarrow FT} && \in \mathcal{D}_!(FS \Rightarrow FT, F(S \Rightarrow T)). \end{aligned}$$

Proof. A lengthy but direct diagram chase. \square

This sets most of the proof obligations for proving soundness of the collapse from **Strat** to **\mathcal{N} -Rel**: we must show that $f(-)$ yields a relative Seely \sim -functor from **Strat** to **\mathcal{N} -Rel**.

5.2. A symmetric monoidal \sim -functor. We define the functor $f(-)$ and equip it with relative Seely structure, starting with symmetric monoidal structure.

5.2.1. A \sim -functor. As expected, on arenas we set $f(A) = \mathcal{C}_{\cong}^0(A)$. From Corollary 4.20, we already have an operation preserving composition

$$f(-) : \mathbf{Strat} \rightarrow \mathcal{N}\text{-Rel}.$$

To get a \sim -functor, it remains to check that $f(-)$ preserves identities and \sim .

Proposition 5.4. *Consider A an arena, and $x_A, y_A \in \mathcal{C}_{\cong}^0(A)$. Then, $f(\mathbf{c}_A)_{x_A, y_A} = \delta_{x_A, y_A}$.*

Proof. First, assume $x_A \neq y_A$ and, seeking a contradiction, consider $z_A \parallel z_A \in \mathbf{wit}_{\mathbf{c}_A}^+(x_A, y_A)$, relying on Proposition 3.30 for the shape of $+$ -covered configurations. So there are

$$\theta_A^- : z_A \cong_A^- x_A, \quad \theta_A^+ : z_A \cong_A^+ y_A,$$

and so $x_A \cong_A y_A$ by composition, contradiction.

Next we must show that for all $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$, $\text{wit}_{\mathbf{x}_A}^+(\mathbf{x}_A, \mathbf{x}_A)$ has exactly one element. First, it is immediate by Proposition 3.30 that $\underline{x}_A \parallel \underline{x}_A \in \text{wit}_{\mathbf{x}_A}^+(\mathbf{x}_A, \mathbf{x}_A)$. For uniqueness, consider $x_A \parallel x_A \in \text{wit}_{\mathbf{x}_A}^+(\mathbf{x}_A, \mathbf{x}_A)$. By definition of \sim^+ -witnesses, there are symmetries

$$\theta_A^- : x_A \cong_A^- \underline{x}_A, \quad \theta_A^+ : x_A \cong_A^+ \underline{x}_A,$$

but because \underline{x}_A is canonical this entails that $x_A = \underline{x}_A$. \square

Next we prove that $f(-)$ preserves \sim . As in the target category the equivalence relation \sim is the identity, this amounts to $f(-)$ being invariant under \approx .

Proposition 5.5. *Consider A, B arenas, and $\sigma, \tau \in \text{Strat}(A, B)$ such that $\sigma \approx \tau$.*

Then, for all $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$ and $\mathbf{x}_B \in \mathcal{C}_{\cong}^0(B)$, $f(\sigma)_{\mathbf{x}_A, \mathbf{x}_B} = f(\tau)_{\mathbf{x}_A, \mathbf{x}_B}$.

Proof. By Definition 3.31, there is a positive isomorphism $\varphi : \sigma \approx \tau$. Recall that this means

$$\{(\partial_\sigma(s), \partial_\tau \circ \varphi(s)) \mid s \in x\} \in \mathcal{S}_+(A \vdash B)$$

for all $x \in \mathcal{C}(\sigma)$, with φ an isomorphism of ess – we write $\psi_A^x \parallel \psi_B^x$ for this symmetry, satisfying by construction $\psi_A^x : x_A^\sigma \cong_A^- y_A^\tau$ and $\psi_B^x : x_B^\sigma \cong_B^+ y_B^\tau$ writing $y^\tau = \varphi x^\sigma \in \mathcal{C}^+(\tau)$.

Now, for $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$ and $\mathbf{x}_B \in \mathcal{C}_{\cong}^0(B)$, we construct a bijection

$$\begin{aligned} \varphi & : \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B) \simeq \text{wit}_\tau^+(\mathbf{x}_A, \mathbf{x}_B) \\ & \quad x^\sigma \mapsto \varphi(x^\sigma). \end{aligned}$$

Indeed, consider $x^\sigma \in \text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B)$. By definition, there are $\theta_A^- : x_A^\sigma \cong_A^- \underline{x}_A$ and $\theta_B^+ : x_B^\sigma \cong_B^+ \underline{x}_B$. Now, $\varphi(x^\sigma) \in \mathcal{C}^+(\tau)$ as φ is an order-isomorphism preserving polarities. Furthermore, we have $\theta_A^- \circ (\psi_A^x)^{-1} : y_A^\tau \cong_A^- \underline{x}_A$ and $\theta_B^+ \circ (\psi_B^x) : y_B^\tau \cong_B^+ \underline{x}_B$; which entails $y^\tau \in \text{wit}_\tau^+(\mathbf{x}_A, \mathbf{x}_B)$ as required. By the symmetrical reasoning φ^{-1} sends $\text{wit}_\tau^+(\mathbf{x}_A, \mathbf{x}_B)$ to $\text{wit}_\sigma^+(\mathbf{x}_A, \mathbf{x}_B)$ and they are clearly mutual inverses, which concludes the proof. \square

5.2.2. *Preservation of monoidal structure.* Next, $f(-)$ is a symmetric monoidal \sim -functor.

For A, B any arenas, we provide the components:

$$\begin{aligned} t_{A,B}^\otimes & : f(A) \times f(B) \xrightarrow{\mathcal{N}\text{-Rel}} f(A \otimes B) \\ t^1 & : 1 \xrightarrow{\mathcal{N}\text{-Rel}} f(1) \end{aligned}$$

defined by $(t_{A,B}^\otimes)_{(\mathbf{x}_A, \mathbf{x}_B), y} = \delta_{y, \mathbf{x}_A \parallel \mathbf{x}_B}$ for every $\mathbf{x}_A \in \mathcal{C}_{\cong}^0(A)$ and $\mathbf{x}_B \in \mathcal{C}_{\cong}^0(B)$; and $(t^1)_{\bullet, \emptyset} = 1$.

Proposition 5.6. *We have $(f(-), t^\otimes, t^1)$ a symmetric monoidal \sim -functor.*

Proof. The crux is the naturality of t^\otimes , corresponding to the fact that the tensor operation on morphisms for Strat and $\mathcal{N}\text{-Rel}$ agree, *i.e.* the following diagram commutes in $\mathcal{N}\text{-Rel}$

$$\begin{array}{ccc} f(A) \times f(B) & \xrightarrow{t_{A,B}^\otimes} & f(A \otimes B) \\ f(\sigma) \otimes f(\tau) \downarrow & & \downarrow f(\sigma \otimes \tau) \\ f(A') \times f(B') & \xrightarrow{t_{A',B'}^\otimes} & f(A' \otimes B') \end{array}$$

for all A, B, A', B' arenas, and $\sigma \in \text{Strat}(A, A'), \tau \in \text{Strat}(B, B')$. To prove this, we invoke the characterizing property of the tensor of strategies in Proposition 3.40 – we have

$$(- \otimes -) : \mathcal{C}^+(\sigma) \times \mathcal{C}^+(\tau) \simeq \mathcal{C}^+(\sigma \otimes \tau)$$

such that $\partial_{\sigma \otimes \tau}(x^\sigma \otimes x^\tau) = (x_A^\sigma \parallel x_B^\tau) \parallel (x_{A'}^\sigma \parallel x_{B'}^\tau)$ for all $x^\sigma \in \mathcal{C}^+(\sigma)$ and $x^\tau \in \mathcal{C}^+(\tau)$. For all $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$, $y_{A'} \in \mathcal{C}_{\cong}^0(A')$ and $y_{B'} \in \mathcal{C}_{\cong}^0(B')$, this immediately restricts to

$$(- \otimes -) : \mathbf{wit}_\sigma^+(x_A, y_{A'}) \times \mathbf{wit}_\tau^+(x_B, y_{B'}) \rightarrow \mathbf{wit}_{\sigma \otimes \tau}^+(x_A \parallel x_B, y_{A'} \parallel y_{B'}).$$

Via this bijection, both paths around the diagram compute to the quantity:

$$\sharp(\mathbf{wit}_\sigma^+(x_A, y_{A'}) \times \mathbf{wit}_\tau^+(x_B, y_{B'}))$$

for all $(x_A, x_B) \in f(A) \times f(B)$ and $y_{A'} \parallel y_{B'} \in f(A \otimes B)$; as required.

The further coherence conditions, expressing that the associators, unitors and symmetries agree in both categories, are all immediate verifications relying on the characterization of the $+$ -covered configurations of the corresponding strategies. \square

5.3. A relative Seely \sim -functor. Next we study the preservation of the modality $!(-)$, which is the most challenging. Then we will deal with \multimap and $\&$.

5.3.1. Preservation of the action of $!$ on morphisms. Inspecting the requirements for relative Seely functors (Definition 5.2) we must first show that $f(-)$ preserves strict objects; this is immediate since every object is strict in $\mathcal{N}\text{-Rel}$. We must then exhibit $t_C^! : !f(C) \rightarrow f(!C)$ for every strict C , and show commutation of the diagram (5.1) up to \sim . However, both in \mathbf{Strat} and $\mathcal{R}\text{-Rel}$, the relative comonad $!$ is in fact a proper comonad; this means that we have a concrete presentation of promotion: for every $f : !C \rightarrow D$, $f^\dagger = !f \circ \delta_C$. Thus the diagram (5.1) amounts to the following, for every $f : !C \rightarrow D$ with C, D strict:

$$\begin{array}{ccccccc} !FC & \xrightarrow{\delta_C} & !!FC & \xrightarrow{!t_C} & !F!C & \xrightarrow{!Ff} & !FD \\ & \searrow^{t_C} & & & & & \swarrow^{t_D} \\ & & F!C & \xrightarrow{F\delta_C} & F!!C & \xrightarrow{F!f} & F!D \end{array} \quad (5.2)$$

For this, we will need to understand how the functorial action of $!$ in \mathbf{Strat} relates to that of $!$ in $\mathcal{N}\text{-Rel}$. We study this now, before giving the definition of the maps $t_C^!$.

The comparison is subtle and it seems a good idea to first recall the definition in $\mathcal{N}\text{-Rel}$:

$$(!\alpha)_{\mu, [y_1, \dots, y_n]} = \sum_{\substack{(x_1, \dots, x_n) \text{ s.t. } 1 \leq i \leq n \\ \mu = [x_1, \dots, x_n]}} \prod \alpha_{x_i, y_i} \quad (5.3)$$

for any weighted relation $\alpha : X \dashrightarrow Y$. Something tricky is going on here. It looks like we are summing over all permutations of $\{1, \dots, n\}$, but no: permutations that lead to the same tuple are counted only once (and the rest of the term is invariant under permutations yielding the same tuple). We must understand how this arises in game semantics.

We recall the game semantical definition that we must match against (5.3). Consider A, B arenas, $\sigma \in \mathbf{Strat}(A, B)$, $x_A \in \mathcal{C}_{\cong}^0(!A)$ and $y_B \in \mathcal{C}_{\cong}^0(!B)$, respectively with

$$\underline{x}_A = \parallel_{1 \leq i \leq p}^{\neq \emptyset} x_A^i, \quad \underline{y}_B = \parallel_{1 \leq i \leq n}^{\neq \emptyset} y_B^i$$

where, and from now on, we label these parallel compositions with “ $\neq \emptyset$ ” to emphasize that each component is non-empty. By definition, $f(!\sigma)_{x_A, y_B} = \sharp \mathbf{wit}_{! \sigma}^+(x_A, y_B)$, where we have

$$\mathbf{wit}_{! \sigma}^+(x_A, y_B) \simeq \sum_{x \in \text{Sym}_A^-(\underline{x}_A)} \sum_{y \in \text{Sym}_B^+(\underline{y}_B)} \mathbf{wit}_{! \sigma}(x, y)$$

with $\text{Sym}_A^-(\underline{x}_{!A}) = \{x \in \mathcal{C}(!A) \mid x \cong_{!A}^- \underline{x}_{!A}\}$ and $\text{Sym}_B^+(\underline{y}_{!B}) = \{y \in \mathcal{C}(!B) \mid y \cong_{!B}^+ \underline{y}_{!B}\}$.

Our task is to link this sum to (5.3), which will require us to gradually decompose further its elements. First the sum over all $y \in \text{Sym}_{!B}^+(\underline{y}_{!B})$ may be described very simply:

Lemma 5.7. *There is a bijection:*

$$\begin{aligned} \text{Sym}_{!B}^+(\underline{y}_{!B}) &\simeq \prod_{1 \leq i \leq n} \text{Sym}_B^+(y_B^i) \\ \|\neq \emptyset\|_{1 \leq i \leq n} y_B^i &\mapsto (y_B^1, \dots, y_B^n) \end{aligned}$$

Proof. Obvious by definition of positive symmetries of $!B$ in Definition 3.17. \square

In contrast, the set $\text{Sym}_{!A}^-(\underline{x}_{!A})$ is much wilder, as negative symmetries on A are free to change copy indices at will. Yet, the data of some $x \cong_{!A}^- \underline{x}_{!A}$ may be witnessed by distinct symmetries; in fact even the action of the symmetry on copy indices is not uniquely defined.

To help reason on $\text{Sym}_{!A}^-(\underline{x}_{!A})$ we need more structure. For $x \in \text{Sym}_{!A}^-(\underline{x}_{!A})$, we write

$$x = \|\neq \emptyset\|_{k \in K_x} x_A^k$$

where $x_A^k \in \mathcal{C}(A)$ for $k \in K_x$. We choose, for each $x \in \text{Sym}_{!A}^-(\underline{x}_{!A})$ a bijection $\pi_x : K_x \simeq \{1, \dots, p\}$ such that for all $k \in K_x$, $x_A^k \cong_A^- \underline{x}_A^{\pi_x(k)}$. If π is a permutation on $\{1, \dots, p\}$, we say it is an **isotropy** of $\underline{x}_{!A}$ if for all $1 \leq i \leq p$ we have $\underline{x}_A^i \cong_A^- \underline{x}_A^{\pi(i)}$, i.e. $\underline{x}_A^i = \underline{x}_A^{\pi(i)}$. Isotropies of $\underline{x}_{!A}$ form a group $\mathfrak{m}(\underline{x}_{!A})$, the **isotropy group** of $\underline{x}_{!A}$. Now, we prove:

Lemma 5.8. *We have the following bijection:*

$$\begin{aligned} \mathfrak{m}(\underline{x}_{!A}) \times \text{Sym}_{!A}^-(\underline{x}_{!A}) &\simeq \sum_{K \subseteq_f \mathbb{N}} \sum_{\pi: K \simeq \{1, \dots, p\}} \prod_{k \in K} \text{Sym}_A^-(\underline{x}_A^{\pi(k)}) \\ (\vartheta, x) &\mapsto (K_x, \vartheta \circ \pi_x, (x_A^k)_{k \in K_x}) \end{aligned}$$

Proof. We first check that this map is well-defined. Consider $\vartheta \in \mathfrak{m}(\underline{x}_{!A})$ and $x \cong_{!A}^- \underline{x}_{!A}$. We must show that for all $k \in K_x$, we have $x_A^k \cong_A^- \underline{x}_A^{\vartheta \circ \pi_x(k)}$. We know that $x_A^k \cong_A^- \underline{x}_A^{\pi_x(k)}$. Moreover, by definition of $\mathfrak{m}(\underline{x}_{!A})$, we have $\underline{x}_A^{\pi_x(k)} = \underline{x}_A^{\vartheta \circ \pi_x(k)}$; so $x_A^k \cong_A^- \underline{x}_A^{\vartheta \circ \pi_x(k)}$.

We define its inverse. To $K \subseteq_f \mathbb{N}$, $\pi : K \simeq \{1, \dots, p\}$, and $(x_A^k)_{k \in K}$, we associate

$$(\pi \circ \pi_x^{-1}, x) \in \mathfrak{m}(\underline{x}_{!A}) \times \text{Sym}_{!A}^-(\underline{x}_{!A})$$

where $x = \|\neq \emptyset\|_{k \in K} x_A^k \cong_{!A}^- \underline{x}_{!A}$ as required. It is clear that the two are inverses. \square

Relying on this bijection, we may start the following computation:

$$\begin{aligned} \mathfrak{m}(\underline{x}_{!A}) \times \text{wit}_{! \sigma}^+(\underline{x}_{!A}, \underline{y}_{!B}) &\simeq \sum_{\vartheta \in \mathfrak{m}(\underline{x}_{!A})} \sum_{(\|\neq \emptyset\|_{k \in K} x_A^k) \in \text{Sym}_{!A}^-(\underline{x}_{!A})} \sum_{y \in \text{Sym}_{!B}^+(\underline{y}_{!B})} \text{wit}_{! \sigma}(\|\neq \emptyset\|_{k \in K} x_A^k, y) \\ &\simeq \sum_{K \subseteq_f \mathbb{N}} \sum_{\pi: K \simeq \{1, \dots, p\}} \sum_{(x_A^k)_{k \in K}} \sum_{(y_B^i)_{1 \leq i \leq n}} \text{wit}_{! \sigma}(\|\neq \emptyset\|_{k \in K} x_A^k, \|\neq \emptyset\|_{1 \leq i \leq n} y_B^i) \end{aligned}$$

where $(x_A^k)_{k \in K}$ ranges over $\prod_{k \in K} \text{Sym}_A^-(\underline{x}_A^{\pi(k)})$ and $(y_B^i)_{1 \leq i \leq n}$ over $\prod_{1 \leq i \leq n} \text{Sym}_B^+(\underline{y}_B^i)$.

Now, let us recall that Proposition 3.47 gives us an order-iso

$$[-] : \text{Fam}(\mathcal{C}^{+, \neq \emptyset}(\sigma)) \simeq \mathcal{C}^+(\! \sigma)$$

with $\text{Fam}(X)$ the set of families of elements of X indexed by finite subsets of \mathbb{N} , such that

$$\partial_{! \sigma}([\!(x^i)_{i \in I}\!]) = (\|\!_{i \in I} x_A^i\| \|\!_{i \in I} x_B^i\|)$$

where for all $i \in I$, $\partial_\sigma(x^i) = x_A^i \parallel x_B^i$. In particular, this entails that the set above can be non-empty only if $K \subseteq \{1, \dots, n\}$; so it is in bijection with

$$\simeq \sum_{K \subseteq \{1, \dots, n\}} \sum_{\pi: K \simeq \{1, \dots, p\}} \sum_{(x_A^k)_{k \in K}} \sum_{(y_B^i)_{1 \leq i \leq n}} \text{wit}_{!_\sigma} \left(\parallel_{k \in K}^{\neq \emptyset} x_A^k, \parallel_{1 \leq i \leq n}^{\neq \emptyset} y_B^i \right) \quad (5.4)$$

To simplify the sum further we shall need the next lemma. It is a variant of Lemma 5.8, but also dealing with the fact that we might have fewer non-empty configurations on A than on B , and introducing a sum over sequences of symmetry classes akin to (5.3).

Lemma 5.9. *We have the following bijection:*

$$\sum_{K \subseteq \{1, \dots, n\}} \sum_{\pi: K \simeq \{1, \dots, p\}} \prod_{k \in K} \text{Sym}_A^-(\underline{x}_A^{\pi(k)}) \simeq \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} z_A^i}} \sum_{\vartheta \in \mathfrak{m}(x_{!A})} \prod_{1 \leq i \leq n} \text{Sym}_A^-(z_A^i)$$

Proof. As for (5.3), the sum on the right hand side ranges over all tuples. Fix in advance, for all $\vec{z} = (z_A^1, \dots, z_A^n)$ such that $\underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} z_A^i$, an injection $\kappa_{\vec{z}}: \{1, \dots, p\} \rightarrow \{1, \dots, n\}$ s.t. for all $1 \leq i \leq p$, $\underline{x}_A^i \cong_A \underline{z}_A^{\kappa_{\vec{z}}(i)}$. Necessarily, z_A^i is empty for all $i \notin \text{cod}(\kappa_{\vec{z}})$.

Given $K \subseteq \{1, \dots, n\}$, $\pi: K \simeq \{1, \dots, p\}$ and $(x_A^k)_{k \in K}$, we set $\vec{z} = (z_A^1, \dots, z_A^n)$ with

$$z_A^i = \begin{cases} x_A^{\pi(i)} & \text{if } i \in K \\ \emptyset & \text{otherwise} \end{cases},$$

by construction we have $\underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} z_A^i$. We set $\vartheta = \pi \circ \kappa_{\vec{z}} \in \mathfrak{m}(x_{!A})$. Finally, we set

$$x_A^i = \begin{cases} x_A^i & \text{if } i \in K \\ \emptyset & \text{otherwise} \end{cases};$$

if $i \notin K$, $x_A^i = z_A^i = \emptyset$; if $i \in K$, $x_A^i \cong_A^- \underline{x}_A^{\pi(i)}$ by hypothesis and $z_A^i = \underline{x}_A^{\pi(i)}$ by construction.

Reciprocally, consider $\vec{z} = (z_A^1, \dots, z_A^n)$, $\vartheta \in \mathfrak{m}(x_{!A})$ and $(x_A^i)_{1 \leq i \leq n}$. We set K as the subset of all $k \in \{1, \dots, n\}$ such that z_A^k is non-empty. We set the bijection

$$\begin{aligned} \pi &: K \simeq \{1, \dots, p\} \\ k &\mapsto \vartheta \circ \kappa_{\vec{z}}^{-1}(k) \end{aligned}$$

which is well-defined as K is exactly the codomain of $\kappa_{\vec{z}}$. For every $k \in K$, we set $(x_A^k)_{k \in K}$ simply as the restriction of the family $(x_A^i)_{1 \leq i \leq n}$ to K – and we do indeed have

$$x_K^k \cong_A^- z_A^k \cong_A^- \underline{x}_A^{\kappa_{\vec{z}}^{-1}(k)} \cong_A^- \underline{x}_A^{\vartheta \circ \kappa_{\vec{z}}^{-1}(k)} = \underline{x}_A^{\pi(k)}.$$

Finally, it is a direct verification that these constructions are inverses. \square

We start again computing from (5.4). Substituting the bijection of the lemma above:

$$\begin{aligned} &\simeq \sum_{K \subseteq \{1, \dots, n\}} \sum_{\pi: K \simeq \{1, \dots, p\}} \sum_{(x_A^k)_{k \in K}} \sum_{(y_B^i)_{1 \leq i \leq n}} \text{wit}_{!_\sigma} \left(\parallel_{k \in K}^{\neq \emptyset} x_A^k, \parallel_{1 \leq i \leq n}^{\neq \emptyset} y_B^i \right) \\ &\simeq \sum_{\vartheta \in \mathfrak{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} z_A^i}} \sum_{(x_A^i)_{1 \leq i \leq n}} \sum_{(y_B^i)_{1 \leq i \leq n}} \text{wit}_{!_\sigma} \left(\parallel_{1 \leq i \leq n} x_A^i, \parallel_{1 \leq i \leq n}^{\neq \emptyset} y_B^i \right) \end{aligned}$$

where now $(x_A^i)_{1 \leq i \leq n}$ ranges over $\prod_{1 \leq i \leq n} \text{Sym}_A^-(z_A^i)$ and $(y_B^i)_{1 \leq i \leq n}$ over $\prod_{1 \leq i \leq n} \text{Sym}_B^+(y_B^i)$.

Some of the x_A^i may now be empty, but both parallel compositions range over the same indices. Thanks to this we may apply Proposition 3.47, which directly yields:

$$\simeq \sum_{\vartheta \in \mathfrak{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} \underline{z}_A^i}} \sum_{(x_A^i)_{1 \leq i \leq n}} \sum_{(y_B^i)_{1 \leq i \leq n}} \prod_{1 \leq i \leq n} \text{wit}_\sigma(x_A^i, y_B^i)$$

We may now complete the computation, with:

$$\begin{aligned} &\simeq \sum_{\vartheta \in \mathfrak{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} \underline{z}_A^i}} \prod_{1 \leq i \leq n} \sum_{x_A^i \cong_A^- \underline{z}_A^i} \sum_{y_B^i \cong_B^+ \underline{x}_B^i} \text{wit}_\sigma(x_A^i, y_B^i) \\ &\simeq \sum_{\vartheta \in \mathfrak{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} \underline{z}_A^i}} \prod_{1 \leq i \leq n} \text{wit}_\sigma^+(z_A^i, y_B^i), \end{aligned}$$

which concludes the construction of the following bijection:

Lemma 5.10. *For A, B arenas, strategy $\sigma \in \text{Strat}(A, B)$, and symmetry classes $x_{!A} \in \mathcal{C}_{\cong}^0(!A)$, $y_{!B} \in \mathcal{C}_{\cong}^0(!B)$ with $y_{!B} = [y_B^1, \dots, y_B^n]$ with each y_B^i non-empty, we have a bijection*

$$U : \sum_{\pi \in \mathfrak{m}(x_{!A})} \text{wit}_{! \sigma}^+(x_{!A}, y_{!B}) \simeq \sum_{\varpi \in \mathfrak{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} \text{wit}_\sigma^+(z_A^i, y_B^i)$$

such that for all $K(\pi, x^{! \sigma}) = (\varpi, ((z_A^1, \dots, z_A^n), (x^i)_{1 \leq i \leq n}))$, we have $x^{! \sigma} = [(x^i)_{1 \leq i \leq n}]$.

Proof. Note $\underline{x}_{!A} \cong_A \parallel_{1 \leq i \leq n} \underline{z}_A^i$ iff $\underline{x}_{!A} = [z_A^i | z_A^i \neq \emptyset]$ by Lemma 3.18 – complete symmetry classes of $!A$ match finite multisets of *non-empty* complete symmetry classes of A . \square

We may finally deduce the desired equality:

Corollary 5.11. *For A, B arenas, strategy $\sigma \in \text{Strat}(A, B)$, and symmetry classes $x_{!A} \in \mathcal{C}_{\cong}^0(!A)$, $y_{!B} \in \mathcal{C}_{\cong}^0(!B)$ with $y_{!B} = [y_B^1, \dots, y_B^n]$ and each y_B^i non-empty, we have:*

$$\# \text{wit}_{! \sigma}^+(x_{!A}, y_{!B}) = \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} \# \text{wit}_\sigma^+(z_A^i, y_B^i) \quad (5.5)$$

Proof. By Lemma 5.10, taking the cardinalities we have the equality:

$$\# \mathfrak{m}(x_{!A}) \times \# \text{wit}_{! \sigma}^+(x_{!A}, y_{!B}) = \# \mathfrak{m}(x_{!A}) \times \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ \underline{x}_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} \# \text{wit}_\sigma^+(z_A^i, y_B^i)$$

from which the result follows by dividing by $\# \mathfrak{m}(x_{!A})$ (which we can do as it is finite). \square

As for composition, we must pad the desired identity with further symmetry groups in order to realize it. The equation (5.5) is very much like (5.4), and we will use this result to show that $f(-)$ has the appropriate preservation properties for $!$. First we explain why only the relative comonad structure is preserved, and not the full comonad structure.

5.3.2. *Non-preservation of the comonad !.* For C a strict arena, we define

$$t_C^! : !f(C) \rightarrow f(!C)$$

where $(t_C^!)_{\mu, x_{!C}} = \delta_{\mu, s_C^!(x_{!C})}$, via the bijection of Lemma 3.18. For strict C, D , we deduce from Corollary 5.11 that for $\sigma \in \text{Strat}(C, D)$, the following diagram commutes in $\mathcal{N}\text{-Rel}$:

$$\begin{array}{ccc} !f(C) & \xrightarrow{t_C^!} & f(!C) \\ !f(\sigma) \downarrow & & \downarrow f(!\sigma) \\ !f(D) & \xrightarrow{t_D^!} & f(!D) \end{array}$$

So $t^!$ is a natural transformation $! \circ f \rightarrow f \circ ! : \text{Strat}_s \rightarrow \mathcal{N}\text{-Rel}$. However, having this for strict C and D is not sufficient, because the construction of the Kleisli category relies crucially on promotion. We must therefore consider strategies of the form $\sigma : !C \rightarrow D$, where of course here $!C$ is not strict and the property above does not directly apply.

The issue is that there *is* a difference between $!f(A) = \mathcal{M}_f(f(A))$ and $f(!A)$ for A non-strict: the latter has only one empty configuration, whereas the former distinguishes between elements $[\emptyset, \dots, \emptyset]$ containing n occurrences of \emptyset , for every n .

Consequently, the naturality square above fails for any reasonable extension of $t_A^!$ to non-strict A . For instance, considering $\sigma \in \text{Strat}(1, \mathbf{B})$ that immediately answers \mathbf{tt} ,

$$(!f(\sigma))_{\emptyset^n, \mathbf{tt}^p} = \delta_{n,p}$$

where $\emptyset^n, \mathbf{tt}^p$ are the obvious multisets. In other words, the relational model remembers how many times σ “does not call” its argument. In contrast, we have $f(!\sigma)_{\emptyset, \mathbf{tt}^p} = 1$ for all $p \in \mathbb{N}$ – $\mathcal{C}_{\cong}^0(!1)$ is a singleton set. Fortunately, this mismatch disappears for promotion.

5.3.3. *Preservation of promotion.* We verify the necessary diagram (5.2).

Proposition 5.12. *Consider C, D strict arenas, and $\sigma \in \text{Strat}(!C, D)$.*

Then, promotion is preserved, i.e. the following diagram commutes in $\mathcal{N}\text{-Rel}$:

$$\begin{array}{ccccccc} \mathcal{M}_f(\mathcal{C}_{\cong}^0(C)) & \xrightarrow{\delta_C} & \mathcal{M}_f(\mathcal{M}_f(\mathcal{C}_{\cong}^0(C))) & \xrightarrow{t_C^!} & \mathcal{M}_f(\mathcal{C}_{\cong}^0(!C)) & \xrightarrow{!f(\sigma)} & \mathcal{M}_f(\mathcal{C}_{\cong}^0(D)) \\ & \searrow^{t_C^!} & \mathcal{C}_{\cong}^0(!C) & \xrightarrow{f(\delta_C)} & \mathcal{C}_{\cong}^0(!!C) & \xrightarrow{f(!\sigma)} & \mathcal{C}_{\cong}^0(!D) & \swarrow^{t_D^!} \end{array}$$

Proof. For $\mu \in \mathcal{M}_f(\mathcal{C}_{\cong}^0(C))$ and $y_{!D} \in \mathcal{C}_{\cong}^0(!D)$, the upper-right path evaluates to:

$$\sum_{\substack{(x_{!C}^1, \dots, x_{!C}^n) \text{ s.t.} \\ x_{!C}^1 + \dots + x_{!C}^n = \mu}} \prod_{1 \leq i \leq n} f(\sigma)_{x_{!C}^i, y_D^i} \quad (5.6)$$

writing $y_{!D} = [y_D^1, \dots, y_D^n]$, inlining $s_D^!$ and $s_C^!$. This is by (5.3) and direct computation.

For the other path, first note that for any $x_{!C} \in \mathcal{C}_{\cong}^0(!C)$ and $y_{!!C} \in \mathcal{C}_{\cong}^0(!!C)$ we have

$$f(\delta_C)_{x_{!C}, y_{!!C}} = \delta_{x_{!C}, y_{!C}^1 + \dots + y_{!C}^n}$$

where $y_{!!C} = [y_{!C}^1, \dots, y_{!C}^n]$ with each $y_{!C}^i$ non-empty – this is proved by a direct elaboration of Proposition 5.4. Relying on this and Corollary 5.11, the bottom-left path evaluates to

$$\sum_{\substack{(x_{!C}^1, \dots, x_{!C}^n) \text{ s.t.} \\ \sum [x_{!C}^i | x_{!C}^i \neq \emptyset] = \mu}} \prod_{1 \leq i \leq n} f(\sigma)_{x_{!C}^i, y_D^i}$$

which is almost (5.6), except for the side-condition. But fortunately, $\sum[x_{iC}^i \mid 1 \leq i \leq n] = \sum[x_{iC}^i \mid x_{iC}^i \neq \emptyset]$ as the empty symmetry class corresponds to the empty multiset. \square

We see that the mismatch causing the failure of naturality disappears with the promotion, as the junk enumeration of multisets in the relational model is erased by the sum.

5.3.4. *A relative Seely \sim -functor.* To wrap up, we introduce the missing components

$$\begin{aligned} t^\top &: \emptyset \rightarrow f(\top) \\ t_{C,D}^{\&} &: f(C) + f(D) \rightarrow f(C \& D) \\ t_{A,C}^{\circ} &: f(A) \times f(C) \rightarrow f(A \multimap C) \end{aligned}$$

for A, B, C, D arenas with C, D strict, defined by t^\top with empty domain, and

$$(t_{C,D}^{\&})_{x,x} = \delta_{x, s_{C,D}^{\&}(x)} \quad (t_{A,C}^{\circ})_{x,x} = \delta_{x, s_{A,C}^{\circ}(x)}.$$

The missing five coherence diagrams of Figure 14 are direct, from an analysis of the symmetry classes reached by the component strategies. As for copycat in Proposition 5.4, this follows from the description of the $+$ -covered configurations of projections in Section 3.6.5, dereliction and monoidality in Section 3.6.6, and evaluation in Section 3.6.4. Altogether:

Corollary 5.13. *We have a relative Seely \sim -functor $f(-) : \text{Strat} \rightarrow \mathcal{N}\text{-Rel}$.*

So by Proposition 5.3 we have a cartesian closed \sim -functor $f_!(-) : \text{Strat}_! \rightarrow \mathcal{N}\text{-Rel}_!$.

5.4. **Preservation of the Interpretation.** The above covers the simply-typed λ -calculus; it remains to address constants and primitives, and recursion.

5.4.1. *Mediating isomorphisms.* By Proposition 5.3, we have isos in $\mathcal{N}\text{-Rel}_!$ for A, B strict

$$\begin{aligned} k^\top &: \top \rightarrow f(\top) \\ k_{A,B}^{\&} &: f(A) \& f(B) \rightarrow f(A \& B) \\ k_{A,B}^{\Rightarrow} &: !f(A) \multimap f(B) \rightarrow f(A \Rightarrow B) \end{aligned}$$

with $A \Rightarrow B = !A \multimap B$. To these, for ground \mathbb{X} we add $k^{\mathbb{X}} : (\mathbb{X}) \rightarrow f([\mathbb{X}])$ defined as $t^{\mathbb{X}} \circ \epsilon_{(\mathbb{X})}$, with $t^{\mathbb{X}} \in \mathcal{N}\text{-Rel}([\mathbb{X}], f([\mathbb{X}]))$ set as $(t^{\mathbb{X}})_{x,x} = \delta_{x, s^{\mathbb{X}}(x)}$ with $s^{\mathbb{X}}$ from Lemma 3.7.

We generalize these mediating isos to all types, by defining isomorphisms $k_\Gamma^{\text{ctx}} : (\Gamma) \rightarrow f([\Gamma])$ and $k_A^{\text{ty}} : (A) \rightarrow f([A])$ in $\mathcal{N}\text{-Rel}_!$ inductively, as follows:

$$\begin{aligned} k_\emptyset^{\text{ctx}} &= k^\top &: \top &\rightarrow f(\top) \\ k_{\Gamma, x:A}^{\text{ctx}} &= k_{[\Gamma], [A]}^{\&} \odot! (k_\Gamma^{\text{ctx}} \&! k_A^{\text{ty}}) &: (\Gamma, x:A) &\rightarrow f([\Gamma, x:A]) \\ k_{A \rightarrow B}^{\text{ty}} &= k_{[A], [B]}^{\Rightarrow} \odot! ((k_A^{\text{ty}})^{-1} \Rightarrow k_B^{\text{ty}}) &: (A \rightarrow B) &\rightarrow f([A \rightarrow B]) \end{aligned}$$

where $\&_!$ is the functorial action of the cartesian product in $\mathcal{N}\text{-Rel}_!$.

These isos may be described more directly in the linear category $\mathcal{N}\text{-Rel}$. First we set:

$$\begin{aligned} t_\emptyset^{\text{ctx}} &= t^\top &: \top &\rightarrow f(\top) \\ t_{\Gamma, x:A}^{\text{ctx}} &= t_{[\Gamma], [A]}^{\&} \odot (t_\Gamma^{\text{ctx}} \& t_A^{\text{ty}}) &: (\Gamma, x:A) &\rightarrow f([\Gamma, x:A]) \\ t_\Gamma^{\text{lctx}} &= t_{[\Gamma]}^! \odot! (t_\Gamma^{\text{ctx}}) &: !(\Gamma) &\rightarrow f(![\Gamma]) \\ t_{A \rightarrow B}^{\text{ty}} &= t_{[A], [B]}^{\circ} \odot ((t_A^{\text{ty}})^{-1} \multimap t_B^{\text{ty}}) &: [A \rightarrow B] &\rightarrow f([A \rightarrow B]) \\ t_A^{\text{ty}} &= t_{[A]}^! \odot! (t_A^{\text{ty}}) &: !(A) &\rightarrow f(![A]) \end{aligned}$$

$$\begin{array}{ccc}
\begin{array}{ccc}
(\Gamma) & \xrightarrow{(\nu)} & (\mathbb{X}) \\
k_{\Gamma}^{\text{ctx}} \downarrow & & \downarrow k_{\mathbb{X}}^{\text{ty}} \\
f([\Gamma]) & \xrightarrow{f_1([\nu])} & f([\mathbb{X}])
\end{array} &
\begin{array}{ccc}
(\mathbb{B}) \& (\mathbb{X}) \& (\mathbb{X}) & \xrightarrow{\text{if}} & (\mathbb{X}) \\
k \downarrow & & \downarrow k^{\mathbb{X}} \\
f(\mathbf{B} \& \mathbf{X} \& \mathbf{X}) & \xrightarrow{f_1(\text{if})} & f([\mathbb{X}])
\end{array} &
\begin{array}{ccc}
(\mathbb{N}) & \xrightarrow{\text{succ}} & (\mathbb{N}) \\
k^{\mathbb{N}} \downarrow & & \downarrow k^{\mathbb{N}} \\
f(\mathbf{N}) & \xrightarrow{f_1(\text{succ})} & f(\mathbf{N})
\end{array} \\
\\
\begin{array}{ccc}
(\mathbb{N}) & \xrightarrow{\text{pred}} & (\mathbb{N}) \\
k^{\mathbb{N}} \downarrow & & \downarrow k^{\mathbb{N}} \\
f(\mathbf{N}) & \xrightarrow{f_1(\text{pred})} & f(\mathbf{N})
\end{array} &
\begin{array}{ccc}
(\mathbb{N}) & \xrightarrow{\text{iszero}} & (\mathbb{B}) \\
k^{\mathbb{N}} \downarrow & & \downarrow k^{\mathbb{B}} \\
f(\mathbf{N}) & \xrightarrow{f_1(\text{iszero})} & f(\mathbf{B})
\end{array} &
\begin{array}{ccc}
(\Gamma) & \xrightarrow{(\text{coin})} & (\mathbb{B}) \\
k_{\Gamma}^{\text{ctx}} \downarrow & & \downarrow k^{\mathbb{B}} \\
f([\Gamma]) & \xrightarrow{f_1([\text{coin}])} & f(\mathbf{B})
\end{array}
\end{array}$$

Figure 15: Preservation of basic primitives

and then we may prove the following lemma:

Lemma 5.14. *For any context Γ and type A , we have $k_{\Gamma}^{\text{ctx}} = t_{\Gamma}^{\text{ctx}} \circ \epsilon_{[\Gamma]}$ and $k_A^{\text{ty}} = t_A^{\text{ty}} \circ \epsilon_{[A]}$.*

Proof. A direct diagram chase. \square

Finally, we also give the following concrete characterization of the linear mediating isos:

Lemma 5.15. *For Γ a context, A a type, $\gamma \in \mathcal{M}_f([\Gamma])$, $x_{\Gamma} \in \mathcal{C}_{\cong}^0([\Gamma])$, $a \in [A]$, $x_A \in \mathcal{C}_{\cong}^0(A)$,*

$$(t_{\Gamma}^{\text{ctx}})_{\gamma, x_{\Gamma}} = \delta_{s_{\Gamma}^{\text{ctx}}(\gamma), x_{\Gamma}} \quad (t_A^{\text{ty}})_{a, x_A} = \delta_{s_A^{\text{ty}}(a), x_A}.$$

Proof. A direct computation. \square

We must prove that the two interpretations match up to these mediating isos. For constants and primitives this is the following lemma, which holds by immediate inspection:

Lemma 5.16. *The diagrams of Figure 15 commute in $\mathcal{N}\text{-Rel}_!$, for any context Γ , ground type \mathbb{X} , and with $k : (\mathbb{B}) \& (\mathbb{X}) \& (\mathbb{X}) \rightarrow f(\mathbf{B} \& \mathbf{X} \& \mathbf{X})$ the obvious isomorphism.*

5.4.2. *Recursion.* Preservation of the recursion combinator boils down to:

Proposition 5.17. *Consider A, B arenas. Then, the collapse function is continuous:*

$$f(-) : \text{Strat}(A, B) \rightarrow \mathcal{N}\text{-Rel}(f(A), f(B))$$

Proof. Consider directed $D \subseteq \text{Strat}(A, B)$, and $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$. We have

$$\text{wit}_{\sqrt{D}}^+(x_A, x_B) = \bigcup_{\sigma \in D} \text{wit}_{\sigma}^+(x_A, x_B)$$

directly by Proposition 3.52. But additionally, we have

$$\mathcal{P}_f(\text{wit}_{\sqrt{D}}^+(x_A, x_B)) = \bigcup_{\sigma \in D} \mathcal{P}_f(\text{wit}_{\sigma}^+(x_A, x_B)). \quad (5.7)$$

Indeed, if $X \subseteq_f \text{wit}_{\sqrt{D}}^+(x_A, x_B)$ then there is a finite $Y \subseteq_f D$ such that

$$X \subseteq \bigcup_{\sigma \in Y} \text{wit}_{\sigma}^+(x_A, x_B),$$

but as D is directed, there is some $\tau \in D$ such that for all $\sigma \in Y$, we have $\sigma \leq \tau$. It immediately follows that $X \subseteq \text{wit}_\tau^+(x_A, x_B)$ as well. Reciprocally, if $X \subseteq_f \text{wit}_\sigma^+(x_A, x_B)$ for some $\sigma \in D$, then clearly $X \subseteq_f \text{wit}_{\vee D}^+(x_A, x_B)$ as well since $\sigma \leq \vee D$.

From there, we can calculate:

$$f(\vee D)_{x_A, x_B} = \bigvee_{X \subseteq_f \text{wit}_{\vee D}^+(x_A, x_B)} \#X = \bigvee_{\sigma \in D} \bigvee_{X \subseteq_f \text{wit}_\sigma^+(x_A, x_B)} \#X = \bigvee_{\sigma \in D} f(\sigma)_{x_A, x_B}$$

using the definition of $f_\sigma(x_A, x_B)$ as a limit of finite sums. \square

From there, it is easy to deduce preservation of the recursion combinator:

Proposition 5.18. *Consider Γ a context and A a type. Then, the diagram*

$$\begin{array}{ccc} \langle \Gamma \rangle & \xrightarrow{\mathcal{Y}_{\langle \Gamma \rangle, \langle A \rangle}} & \langle (A \rightarrow A) \rightarrow A \rangle \\ k_\Gamma^{\text{ctx}} \downarrow & & \downarrow k_{(A \rightarrow A) \rightarrow A}^{\text{ty}} \\ f(\llbracket \Gamma \rrbracket) & \xrightarrow{f_!(\mathcal{Y}_{\llbracket \Gamma \rrbracket, \llbracket A \rrbracket})} & f(\llbracket (A \rightarrow A) \rightarrow A \rrbracket) \end{array}$$

commutes in $\mathcal{N}\text{-Rel}_!$.

Proof. First, the following diagram commutes in $\mathcal{N}\text{-Rel}_!$ for all $n \in \mathbb{N}$:

$$\begin{array}{ccc} \top & \xrightarrow{\mathcal{Y}_{\langle A \rangle}^n} & \langle (A \rightarrow A) \rightarrow A \rangle \\ k^\top \downarrow & & \downarrow k_{(A \rightarrow A) \rightarrow A}^{\text{ty}} \\ f(\top) & \xrightarrow{f_!(\mathcal{Y}_{\llbracket A \rrbracket}^n)} & f(\llbracket (A \rightarrow A) \rightarrow A \rrbracket) \end{array}$$

as follows directly by induction on n , by direct application of the preservation of the cartesian closed structure. By Proposition 5.17, we may take the supremum and get that

$$\begin{array}{ccc} \top & \xrightarrow{\mathcal{Y}_{\langle A \rangle}} & \langle (A \rightarrow A) \rightarrow A \rangle \\ k^\top \downarrow & & \downarrow k_{(A \rightarrow A) \rightarrow A}^{\text{ty}} \\ f(\top) & \xrightarrow{f_!(\mathcal{Y}_{\llbracket A \rrbracket})} & f(\llbracket (A \rightarrow A) \rightarrow A \rrbracket) \end{array}$$

commutes in $\mathcal{N}\text{-Rel}_!$. The proposition follows by preservation of the terminal object. \square

5.4.3. *Preservation of the interpretation.* Finally, we may conclude our main theorem.

Theorem 5.19. *Consider $\Gamma \vdash M : A$ any term of **nPCF**. Then,*

$$\begin{array}{ccc} \langle \Gamma \rangle & \xrightarrow{\langle M \rangle} & \langle A \rangle \\ k_\Gamma^{\text{ctx}} \downarrow & & \downarrow k_A^{\text{ty}} \\ f(\llbracket \Gamma \rrbracket) & \xrightarrow{f_!(\llbracket M \rrbracket)} & f(\llbracket A \rrbracket) \end{array}$$

commutes in $\mathcal{N}\text{-Rel}_!$.

Proof. By induction on the typing derivation $\Gamma \vdash M : A$. The combinators of the simply-typed λ -calculus follow by preservation of the cartesian closed structure. For constants, it follows from Lemma 5.16. For **if**, **pred**, **succ**, **iszero** and **coin**, it follows from Lemma 5.16 and the preservation of the cartesian structure. Recursion is by Proposition 5.18. \square

A direct reformulation of this theorem is the following:

Theorem 5.20. *Consider any term $\Gamma \vdash M : A$ of **nPCF**, and $\gamma \in \llbracket \Gamma \rrbracket, a \in \llbracket A \rrbracket$. Then,*

$$\llbracket M \rrbracket_{\gamma, a} = \# \text{wit}_{\llbracket M \rrbracket}^+(s_{\Gamma}^{\text{ctx}}(\gamma), s_A^{\text{ty}}(a)).$$

Proof. By Theorem 5.19, we have

$$\begin{array}{ccc} \llbracket \Gamma \rrbracket & \xrightarrow{\llbracket M \rrbracket} & \llbracket A \rrbracket \\ k_{\Gamma}^{\text{ctx}} \downarrow & & \downarrow k_A^{\text{ty}} \\ J(\llbracket \Gamma \rrbracket) & \xrightarrow{f_{\llbracket M \rrbracket}} & J(\llbracket A \rrbracket) \end{array}$$

in $\mathcal{N}\text{-Rel}$. Simplifying Kleisli composition and using Lemma 5.14, we have

$$\begin{array}{ccc} \mathcal{M}_f(\llbracket \Gamma \rrbracket) & \xrightarrow{\llbracket M \rrbracket} & \llbracket A \rrbracket \\ \downarrow !(t_{\Gamma}^{\text{ctx}}) & & \downarrow t_A^{\text{ty}} \\ !J(\llbracket \Gamma \rrbracket) & \xrightarrow{t_{\llbracket \Gamma \rrbracket}^!} J(!\llbracket \Gamma \rrbracket) \xrightarrow{J\llbracket M \rrbracket} & J(\llbracket A \rrbracket) \end{array}$$

in $\mathcal{N}\text{-Rel}$; but by definition of t_{Γ}^{ctx} , this amounts exactly to

$$\begin{array}{ccc} \mathcal{M}_f(\llbracket \Gamma \rrbracket) & \xrightarrow{\llbracket M \rrbracket} & \llbracket A \rrbracket \\ t_{\Gamma}^{\text{ctx}} \downarrow & & \downarrow t_A^{\text{ty}} \\ J(!\llbracket \Gamma \rrbracket) & \xrightarrow{J\llbracket M \rrbracket} & J(\llbracket A \rrbracket) \end{array}$$

in $\mathcal{N}\text{-Rel}$. The theorem follows by Lemma 5.15 and direct computation. \square

This answers our original question: at higher-order types, the weighted relational model counts witnesses in the concurrent game semantics, up to positive symmetry.

6. COLLAPSE OF \mathcal{R} -WEIGHTED STRATEGIES

In this last technical section, we show how all the results above generalize to the collapse of strategies whose configurations are labelled with elements of a continuous semiring \mathcal{R} .

6.1. \mathcal{R} -strategies. We build a relative Seely \sim -category $\mathcal{R}\text{-Strat}$, for any \mathcal{R} .

6.1.1. Basic definition. As for Strat , the objects of $\mathcal{R}\text{-Strat}$ are all arenas. We define:

Definition 6.1. Consider A a game. An \mathcal{R} -strategy on A is a strategy $\sigma : A$, with

$$\mathcal{V}_{\sigma} : \mathcal{C}^+(\sigma) \rightarrow \mathcal{R}$$

a **valuation**, *invariant under symmetry*: for all $x \cong_{\sigma} y$, $\mathcal{V}_{\sigma}(x) = \mathcal{V}_{\sigma}(y)$.

For instance, using $\mathcal{R} = \overline{\mathbb{R}}_+$, we may adjoin to the strategy $\mathbf{coin} : \mathbf{B}$ a valuation $\mathcal{V} : \mathcal{C}^+(\mathbf{coin}) \rightarrow \overline{\mathbb{R}}_+$ with $\mathcal{V}(\{\mathbf{q}^-, \mathbf{t}^+\}) = \mathcal{V}(\{\mathbf{q}^-, \mathbf{f}^+\}) = \frac{1}{2}$; representing a fair coin toss.

For A and B arenas, the homset $\mathcal{R}\text{-Strat}(A, B)$ comprises *visible, exhaustive* \mathcal{R} -strategies on $A \vdash B$ – visibility and exhaustivity are undisturbed by the presence of the valuation.

6.1.2. *Basic strategies and operations.* For any arena A and any $x_A \parallel x_A \in \mathcal{C}^+(\mathfrak{C}_A)$, we set

$$\mathcal{V}_{\mathfrak{C}_A}(x_A \parallel x_A) = 1,$$

as by Proposition 3.30 all $+$ -covered configurations of copycat have this form. All copycat strategies involved in the relative Seely \sim -category structure are made into \mathcal{R} -strategies similarly, by setting their valuation to be 1 everywhere – this covers associators, unitors, projections, evaluation, dereliction, digging, and Seely isomorphisms. All operations on strategies involved in the relative Seely category structure are lifted to \mathcal{R} -strategies with:

$$\begin{aligned} \mathcal{V}_{\tau \odot \sigma}(x^\tau \odot x^\sigma) &= \mathcal{V}_\sigma(x^\sigma) \cdot \mathcal{V}_\tau(x^\tau) & \mathcal{V}_{\langle \sigma, \tau \rangle}(\text{inj}_\tau(x^\tau)) &= \mathcal{V}_\tau(x^\tau) \\ \mathcal{V}_{\sigma \otimes \tau}(x^\sigma \otimes x^\tau) &= \mathcal{V}_\sigma(x^\sigma) \cdot \mathcal{V}_\tau(x^\tau) & \mathcal{V}_{!(\sigma)}(\prod_{i \in I} (x^i)) &= \prod_{i \in I} (\mathcal{V}_\sigma(x^i)) \\ \mathcal{V}_{\langle \sigma, \tau \rangle}(\text{inj}_\sigma(x^\sigma)) &= \mathcal{V}_\sigma(x^\sigma) \end{aligned}$$

leveraging the characterizations of $+$ -covered configurations for these operations, respectively found in Propositions 3.28, 3.40, 3.44, and 3.47. In the last case, \prod denotes the iterated product (\cdot) of \mathcal{R} . One must ensure that the resulting valuation is invariant under symmetry, which is immediate (in the last case, using that the product \cdot is commutative).

Likewise, the partial order \leq is extended to \mathcal{R} -strategies by setting $\sigma \leq \tau$ if it holds for the underlying strategies, and $\mathcal{V}_\sigma(x) = \mathcal{V}_\tau(x)$ for all $x \in \mathcal{C}^+(\sigma)$. It is clear that \leq retains the same completeness properties, and that all operations on \mathcal{R} -strategies are continuous.

In particular, for any strict arenas Γ and A , we may define the recursion combinator

$$\mathcal{Y}_{\Gamma, A} \in \mathcal{R}\text{-Strat}_!(\Gamma, !(A \multimap A) \multimap A)$$

exactly as in Section 2.3.4 – the same strategy, with valuation again set to 1 everywhere.

6.1.3. *Positive isomorphisms.* Finally, we must adapt the equivalence relation on strategies.

Definition 6.2. Consider A a game, and $\sigma, \tau : A$ two \mathcal{R} -strategies.

A **positive isomorphism** $\varphi : \sigma \approx \tau$ is a positive isomorphism between the underlying strategies, such that for all $x \in \mathcal{C}^+(\sigma)$, we have $\mathcal{V}_\tau(\varphi(x)) = \mathcal{V}_\sigma(x)$.

We say that $\sigma, \tau : A$ are **positively isomorphic**, written $\sigma \approx \tau$, if there exists a positive isomorphism $\varphi : \sigma \approx \tau$. We must ensure that this valuation-aware equivalence relation is still preserved under all operations on strategies – it is evident for all, save composition.

For composition, we need more information on how positive isos are propagated:

Proposition 6.3. Let $\sigma, \sigma' \in \text{Strat}(A, B)$; $\tau, \tau' \in \text{Strat}(B, C)$; $\varphi : \sigma \approx \sigma'$, $\psi : \tau \approx \tau'$.

Then, there exists a positive isomorphism $\psi \odot \varphi : \tau \odot \sigma \approx \tau' \odot \sigma'$ such that for all $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$, writing $y^{\tau'} \odot y^{\sigma'} = (\psi \odot \varphi)(x^\tau \odot x^\sigma) \in \mathcal{C}^+(\tau' \odot \sigma')$, we have

$$\varphi(x^\sigma) \cong_{\sigma'} y^{\sigma'}, \quad \psi(x^\tau) \cong_{\tau'} y^{\tau'}.$$

See Appendix A.4 for the proof. From that, we may deduce:

Proposition 6.4. Let $\sigma, \sigma' \in \mathcal{R}\text{-Strat}(A, B)$; $\tau, \tau' \in \mathcal{R}\text{-Strat}(B, C)$; $\varphi : \sigma \approx \sigma'$, $\psi : \tau \approx \tau'$.

Then, $\psi \odot \varphi : \tau \odot \sigma \approx \tau' \odot \sigma'$ is a positive iso between \mathcal{R} -strategies.

Proof. For $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$, writing $y^{\tau'} \odot y^{\sigma'} = (\psi \odot \varphi)(x^\tau \odot x^\sigma)$, we calculate

$$\begin{aligned} \mathcal{V}_{\tau' \odot \sigma'}((\psi \odot \varphi)(x^\tau \odot x^\sigma)) &= \mathcal{V}_{\sigma'}(y^{\sigma'}) \cdot \mathcal{V}_{\tau'}(y^{\tau'}) \\ &= \mathcal{V}_{\sigma'}(\varphi(x^\sigma)) \cdot \mathcal{V}_{\tau'}(\psi(x^\tau)) \\ &= \mathcal{V}_\sigma(x^\sigma) \cdot \mathcal{V}_\tau(x^\tau) \\ &= \mathcal{V}_{\tau \odot \sigma}(x^\tau \odot x^\sigma), \end{aligned}$$

where $\mathcal{V}_{\sigma'}(y^{\sigma'}) = \mathcal{V}_{\sigma'}(\varphi(x^\sigma))$ as $y^{\sigma'} \cong_{\sigma'} \varphi(x^\sigma)$ and likewise for τ . \square

Corollary 6.5. *There is a relative Seely \sim -category $\mathcal{R}\text{-Strat}$.*

Proof. It remains to establish the required positive isomorphisms, *i.e.* to show that the corresponding positive isomorphisms for **Strat** preserve valuations. As an illustration, recall from Proposition 3.36 that associativity is realized with the positive isomorphism

$$\alpha_{\sigma,\tau,\delta} : (\delta \odot \tau) \odot \sigma \approx \delta \odot (\tau \odot \sigma)$$

such that $\alpha_{\sigma,\tau,\delta}((x^\delta \odot x^\tau) \odot x^\sigma) = x^\delta \odot (x^\tau \odot x^\sigma)$ for all $(x^\delta \odot x^\tau) \odot x^\sigma \in \mathcal{C}^+(\delta \odot (\tau \odot \sigma))$. Clearly, this preserves valuations by associativity of \cdot . Other cases are similar. \square

6.1.4. *Interpretation of $\mathcal{R}\text{-PCF}$.* All basic strategies for **nPCF** primitives have valuation set to 1 everywhere, completing the interpretation of **nPCF**. But valuations remain trivial:

Proposition 6.6. *Consider $\Gamma \vdash M : A$ a term of **nPCF**.*

Then, for all $x \in \mathcal{C}^+(\llbracket M \rrbracket)$, we have $\mathcal{V}_{\llbracket M \rrbracket}(x) = 1$.

This is obvious: all basic strategies have all valuations 1, and the operations on strategies only involve the product \cdot of \mathcal{R} , never the sum. To explain that, recall that in $\mathcal{R}\text{-Rel}$ the sum serves to aggregate weights for all executions made distinct by non-deterministic choices. But $\mathcal{R}\text{-Strat}$ maintains explicit branching information, and each witness represents only one individual execution – so it makes sense that coefficients should remain 1.

So as to better illustrate the model of \mathcal{R} -strategies, we add a new primitive:

$$\frac{\Gamma \vdash M : A}{\Gamma \vdash r \cdot M : A}$$

for all $r \in \mathcal{R}$ – we refer to $\mathcal{R}\text{-PCF}$ for the enriched language. There is a matching operation:

Definition 6.7. Consider A a game, and $\sigma : A$ a \mathcal{R} -strategy.

We set $r \cdot \sigma : A$ with strategy σ and valuation $\mathcal{V}_{r \cdot \sigma}(x) = r \cdot \mathcal{V}_\sigma(x)$ for all $x \in \mathcal{C}^+(\sigma)$.

Altogether, this yields an interpretation of $\mathcal{R}\text{-PCF}$ into $\mathcal{R}\text{-Strat}_!$, sending a term $\Gamma \vdash M : A$ to $\llbracket M \rrbracket \in \mathcal{R}\text{-Strat}_!(\llbracket \Gamma \rrbracket, \llbracket A \rrbracket)$. We must also set the interpretation of $\mathcal{R}\text{-PCF}$ in $\mathcal{R}\text{-Rel}$, set with the exact same clauses as for the interpretation in $\mathcal{N}\text{-Rel}$, except for:

$$(\Gamma \vdash r \cdot M : A)_{\mu,a} = r \cdot (\Gamma \vdash M : A)_{\mu,a}.$$

This completes the interpretation of any $\Gamma \vdash M : A$ as $(\llbracket M \rrbracket) \in \mathcal{R}\text{-Rel}_!(\llbracket \Gamma \rrbracket, \llbracket A \rrbracket)$, which we must now compare with $\llbracket M \rrbracket \in \mathcal{R}\text{-Strat}_!(\llbracket \Gamma \rrbracket, \llbracket A \rrbracket)$.

6.2. **A relative Seely \sim -functor.** Next, we show how Corollary 5.13 extends in the presence of quantitative valuations. With the earlier developments of this paper this is mostly a formality: as all earlier compatibility results are realized by explicit bijections between sets of witnesses, we must only exploit that these bijections preserve valuations.

First, we define the quantitative collapse as follows. For any $\sigma \in \mathcal{R}\text{-Strat}(A, B)$, we set:

$$f(\sigma)_{x_A, y_B} = \sum_{x \in \text{wit}_\sigma^+(x_A, x_B)} \mathcal{V}_\sigma(x) \quad (6.1)$$

for all $x_A \in \mathcal{C}_{\cong}^0(A)$ and $x_B \in \mathcal{C}_{\cong}^0(B)$. It is a clear generalization of (4.7), with all witnesses weighted according to their valuation. It is a conservative extension of (4.7): for strategies arising from terms without $r \cdot -$, by Proposition 6.6 it is equivalent to (4.7).

6.2.1. *Composition.* First, we show that (6.1) is compatible with composition. Fortunately, it suffices to exploit the bijections introduced in Section 4.3, along with *integer division*:

Proposition 6.8. *For $\sigma \in \mathcal{R}\text{-Strat}(A, B)$, $\tau \in \mathcal{R}\text{-Strat}(B, C)$, $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_C \in \mathcal{C}_{\cong}^0(C)$:*

$$f(\tau \odot \sigma)_{x_A, x_C} = \sum_{x_B \in \mathcal{C}_{\cong}^0(B)} f(\sigma)_{x_A, x_B} \cdot f(\tau)_{x_B, x_C}.$$

Proof. Let us first fix some $x_B \in \mathcal{C}_{\cong}^0(B)$. We then perform the computation in \mathcal{R} :

$$\begin{aligned} & (\#\mathcal{S}_-(x_A)) * (\#\mathcal{S}(x_B)) * (\#\mathcal{S}_+(x_C)) * \sum_{x^\tau \odot x^\sigma \in \text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C)} \mathcal{V}_{\tau \odot \sigma}(x^\tau \odot x^\sigma) \\ = & \sum_{\theta_A^- \in \mathcal{S}_-(x_A)} \sum_{\theta_B \in \mathcal{S}(x_B)} \sum_{\theta_C^+ \in \mathcal{S}_+(x_C)} \sum_{x^\tau \odot x^\sigma \in \text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C)} \mathcal{V}_\sigma(x^\sigma) \cdot \mathcal{V}_\tau(x^\tau) \\ = & \sum_{\varphi_A^- \in \mathcal{S}_-(x_A)} \sum_{\varphi_B \in \mathcal{S}(x_B)} \sum_{\varphi_C^+ \in \mathcal{S}_+(x_C)} \sum_{y^\sigma \in \text{wit}_\sigma^+(x_A, x_B)} \sum_{y^\tau \in \text{wit}_\tau^+(x_B, x_C)} \mathcal{V}_\sigma(y^\sigma) \cdot \mathcal{V}_\tau(y^\tau) \\ = & (\#\mathcal{S}_-(x_A)) * (\#\mathcal{S}(x_B)) * (\#\mathcal{S}_+(x_C)) * \left[\left(\sum_{y^\sigma \in \text{wit}_\sigma^+(x_A, x_B)} \mathcal{V}_\sigma(y^\sigma) \right) \cdot \left(\sum_{y^\tau \in \text{wit}_\tau^+(x_B, x_C)} \mathcal{V}_\tau(y^\tau) \right) \right] \\ = & (\#\mathcal{S}_-(x_A)) * (\#\mathcal{S}(x_B)) * (\#\mathcal{S}_+(x_C)) * (f(\sigma)_{x_A, x_B} \cdot f(\tau)_{x_B, x_C}) \end{aligned}$$

using the definition of integer multiplication in \mathcal{R} and of the valuation of $\tau \odot \sigma$; then substituting by Φ of Lemma 4.19 and using that the valuation is invariant under symmetry; and using distributivity of \cdot over $+$ in \mathcal{R} and the definition of integer multiplication.

Now, since \mathcal{R} satisfies *integer division*, we may deduce the equality in \mathcal{R} :

$$\sum_{x^\tau \odot x^\sigma \in \text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C)} \mathcal{V}_{\tau \odot \sigma}(x^\tau \odot x^\sigma) = f(\sigma)_{x_A, x_B} \cdot f(\tau)_{x_B, x_C},$$

by dividing each side by $(\#\mathcal{S}_-(x_A))$, $(\#\mathcal{S}(x_B))$, $(\#\mathcal{S}_+(x_C))$. By (4.9), summing both sides over all $x_B \in \mathcal{C}_{\cong}^0(B)$ concludes the proof of the desired equation. \square

We do not know if integer division is really needed. One could avoid it by extracting from the bijection Φ in Lemma 4.19 a direct bijection preserving symmetry classes

$$\text{wit}_{\sigma, \tau}^+(x_A, x_B, x_C) \simeq \text{wit}_\sigma^+(x_A, x_B) \times \text{wit}_\tau^+(x_B, x_C),$$

but it is not immediately clear how to do that. Assuming integer division does not remove any interesting example of continuous semiring; so we did not push this. We obtain:

Proposition 6.9. *The operation $f(-) : \mathcal{R}\text{-Strat} \rightarrow \mathcal{R}\text{-Rel}$ is a \sim -functor.*

Proof. It remains to prove that $f(-)$ preserves the identities and the equivalence relation. For identities the proof of Proposition 5.4 applies, using that valuations for \mathfrak{a}_A are 1.

For the equivalence relation, given $\sigma, \tau \in \mathcal{R}\text{-Strat}(A, B)$ and $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$, given $\varphi : \sigma \approx \tau$ we proved in Proposition 5.5 that φ specializes to a bijection

$$\varphi : \text{wit}_\sigma^+(x_A, x_B) \simeq \text{wit}_\tau^+(x_A, x_B),$$

but as φ is now required to preserve valuations, we have $f(\sigma)_{\times_A, \times_B} = f(\tau)_{\times_A, \times_B}$ as needed. \square

6.2.2. *Preservation of symmetric monoidal structure.* For the tensor, it is straightforward:

Proposition 6.10. *The operation $f(-)$ is a symmetric monoidal \sim -functor.*

Proof. The structural isomorphisms involved are the \mathcal{R} -weighted relations defined with the same formulas as in Section 5.2.2. All coherence laws follow. For naturality of $t_{A,B}^\otimes$, we build on the proof of Proposition 5.6 by noting that the bijection

$$(- \otimes -) : \text{wit}_\sigma^+(\times_A, \times_{A'}) \times \text{wit}_\tau^+(\times_B, \times_{B'}) \rightarrow \text{wit}_{\sigma \otimes \tau}^+(\times_A \parallel \times_B, \times_{A'} \parallel \times_{B'})$$

is such that $\mathcal{V}_{\sigma \otimes \tau}(x^\sigma \otimes x^\tau) = \mathcal{V}_\sigma(x^\sigma) \cdot \mathcal{V}_\tau(x^\tau)$ by definition of the valuation for tensor. From this, the naturality of $t_{A,B}^\otimes$ follows by an immediate calculation. \square

6.2.3. *Preservation of promotion.* As expected, for C a strict arena we set $t_C^!$ as the \mathcal{R} -weighted relation defined with the same formula as in Section 5.3.2. We prove:

Proposition 6.11. *For A, B arenas, strategy $\sigma \in \text{Strat}(A, B)$, and symmetry classes $x_{!A} \in \mathcal{C}_{\cong}^0(!A)$, $y_{!B} \in \mathcal{C}_{\cong}^0(!B)$ with $y_{!B} = [y_B^1, \dots, y_B^n]$ with each y_B^i non-empty, we have:*

$$f(!\sigma)_{x_{!A}, y_{!B}} = \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} f(\sigma)_{z_A^i, y_B^i}$$

Proof. We perform the computation:

$$\begin{aligned} \sharp \mathbf{m}(x_{!A}) * f(!\sigma)_{x_{!A}, y_{!B}} &= \sum_{\pi \in \mathbf{m}(x_{!A})} \sum_{x^{! \sigma} \in \text{wit}_\sigma^+(x_{!A}, y_{!B})} \mathcal{V}_{! \sigma}(x^{! \sigma}) \\ &= \sum_{\varpi \in \mathbf{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \sum_{(x^i) \in \prod_{1 \leq i \leq n} \text{wit}_\sigma^+(z_A^i, y_B^i)} \mathcal{V}_{! \sigma}([x^i]_{1 \leq i \leq n}) \\ &= \sum_{\varpi \in \mathbf{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \sum_{(x^i) \in \prod_{1 \leq i \leq n} \text{wit}_\sigma^+(z_A^i, y_B^i)} \prod_{1 \leq i \leq n} \mathcal{V}_\sigma(x^i) \\ &= \sum_{\varpi \in \mathbf{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} \sum_{x^i \in \text{wit}_\sigma^+(z_A^i, y_B^i)} \mathcal{V}_\sigma(x^i) \\ &= \sum_{\varpi \in \mathbf{m}(x_{!A})} \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} f(\sigma)_{z_A^i, y_B^i} \\ &= \sharp \mathbf{m}(x_{!A}) * \sum_{\substack{(z_A^1, \dots, z_A^n) \text{ s.t.} \\ x_{!A} = [z_A^i | z_A^i \neq \emptyset]}} \prod_{1 \leq i \leq n} f(\sigma)_{z_A^i, y_B^i} \end{aligned}$$

using the definition of integer multiplication; the bijection U in Lemma 5.10; the definition of the valuation for $! \sigma$; distributivity of \cdot over sum; definition of $f(\sigma)$; and again definition of integer multiplication. Finally, the desired equality follows by integer division. \square

From this, it follows – with the same proof – that promotion is preserved as in Proposition 5.12. As for composition, it is not clear whether one can avoid integer division here.

From this point, we can conclude the preservation of the relative Seely structure.

Corollary 6.12. *We have a relative Seely \sim -functor $f(-) : \mathcal{R}\text{-Strat} \rightarrow \mathcal{R}\text{-Rel}$.*

Proof. It remains to define t^\top , $t_{C,D}^\&$ and $t_{A,C}^\circ$ for A, C, D arenas with C, D strict – those are defined with the same formulas as in Section 5.3.4. The coherence laws follow likewise. \square

6.3. Preservation of the Interpretation. While Corollary 6.12 does the heavy lifting, there remain a few things to check. First, preservation of recursion boils down to:

Proposition 6.13. *Consider A, B arenas. Then, the collapse function is continuous:*

$$f(-) : \mathcal{R}\text{-Strat}(A, B) \rightarrow \mathcal{R}\text{-Rel}(f(A), f(B))$$

Proof. Consider directed $D \subseteq \mathcal{R}\text{-Strat}(A, B)$, and $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$. We compute:

$$\begin{aligned} f(\vee D)_{x_A, x_B} &= \sum_{x^\sigma \in \text{wit}_{\vee D}^+(x_A, x_B)} \mathcal{V}_\sigma(x^\sigma) \\ &= \bigvee_{X \subseteq_f \text{wit}_{\vee D}^+(x_A, x_B)} \sum_{x^\sigma \in X} \mathcal{V}_\sigma(x^\sigma) \\ &= \bigvee_{\sigma \in D} \bigvee_{Y \subseteq_f \text{wit}_\sigma^+(x_A, x_B)} \sum_{x^\sigma \in Y} \mathcal{V}_\sigma(x^\sigma) \end{aligned}$$

which is $\bigvee_{\sigma \in D} f(\sigma)_{x_A, x_B}$ by definition – we used the definition of infinite sums, and (5.7). \square

It follows that the recursion combinator is preserved, with the same proof as Proposition 5.18. Likewise, all the diagrams in Figure 15 immediately hold. Finally, the interpretations of $r \cdot -$ trivially agree with each other as well. To conclude, we have:

Theorem 6.14. *Consider $\Gamma \vdash M : A$ any term of $\mathcal{R}\text{-PCF}$. Then,*

$$\begin{array}{ccc} (\Gamma) & \xrightarrow{\langle M \rangle} & (A) \\ k_\Gamma^{\text{ctx}} \downarrow & & \downarrow k_A^{\text{by}} \\ f(\llbracket \Gamma \rrbracket) & \xrightarrow{f(\llbracket M \rrbracket)} & f(\llbracket A \rrbracket) \end{array}$$

commutes in $\mathcal{R}\text{-Rel}_!$.

Proof. As for Theorem 5.19 with the ingredients introduced in this section. \square

As in the earlier case, we also provide a more concrete statement:

Theorem 6.15. *Consider any term $\Gamma \vdash M : A$ of $\mathcal{R}\text{-PCF}$, and $\gamma \in (\Gamma)$, $a \in (A)$. Then,*

$$(M)_{\gamma, a} = \sum_{x \in \text{wit}_{\llbracket M \rrbracket}^+(s_\Gamma^{\text{ctx}}(\gamma), s_A^{\text{by}}(a))} \mathcal{V}_{\llbracket M \rrbracket}(x).$$

Proof. Direct from Theorem 6.14, with the same proof as for Theorem 5.20. \square

From this, one obtains game semantics for various continuous semirings, inheriting adequacy properties from [LMMP13]. Details are out of scope of the paper.

The weighted relational model is inherently infinite, because the sum (2.2) involved in the composition of weighted relations has no reason to be finite. This infinitary nature is sometimes criticized; for instance probabilistic coherence spaces [EPT11] consist in enriching the weighted relational model with a biorthogonality construction ensuring (among other things) that all coefficients remain finite. So it is noteworthy that no infinity arises in \mathcal{R} -Strat: the construction unfolds just fine with only a plain semiring – or in fact, only a *monoid* $(|\mathcal{R}|, \cdot, 1)$! Indeed, as it stands, the sum only arises when collapsing to \mathcal{R} -Rel.

7. CONCLUSION

As a rough approximation, there are essentially two families of denotational models in the legacy of linear logic: on the one hand the *web-based* semantics such as relational models, coherence spaces and their weighted counterparts, arising from Girard’s quantitative semantics [Gir88]; and on the other hand the *interactive semantics* drawing inspiration, among others, from Girard’s geometry of interaction [Gir89]. The two families are great for different things: the former family has had impressive achievements in modeling quantitative aspects of programming, with notably the recent full abstraction result for probabilistic **PCF** due to Ehrhard, Pagani and Tasson [EPT18]; while the latter has proved particularly powerful in capturing effectful programming languages [MT16]. It is certainly puzzling that these families, though sharing such a close genesis, have remained almost separated!

We believe the results presented here are an important step towards bringing these two families together, aiming towards a unified landscape of quantitative denotational models of programming languages. We proved this for **PCF**, but there is no doubt that this extends to other languages or evaluation strategies – in fact, the first author and de Visme proved a similar collapse theorem for the (call-by-value) *quantum λ -calculus* [CdV20] (this relies on some of the constructions of this paper, first appearing in an unpublished technical report by the first author [Cla20]). Of course, much remains to be done: notably, we would like to understand better the links between thin concurrent games and generalized species of structure [FGHW08]. Much of the present development is also reminiscent of issues related to rigid resource terms and the Taylor development of λ -terms [OA20].

Acknowledgments. We would like to thank Marc de Visme for lively discussions during the development of this work, and the counter-example to representability (Appendix B).

This work was supported by the ANR project DyVerSe (ANR-19-CE48-0010-01); and by the Labex MiLyon (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007), operated by the French National Research Agency (ANR).

REFERENCES

- [ACU10] Thorsten Altenkirch, James Chapman, and Tarmo Uustalu. Monads need not be endofunctors. In *International Conference on Foundations of Software Science and Computational Structures*, pages 297–311. Springer, 2010.
- [AJM00] Samson Abramsky, Radha Jagadeesan, and Pasquale Malacaria. Full abstraction for PCF. *Inf. Comput.*, 163(2):409–470, 2000.

- [BDER97] Patrick Baillot, Vincent Danos, Thomas Ehrhard, and Laurent Regnier. Timeless games. In *Computer Science Logic, 11th International Workshop, CSL '97, Annual Conference of the EACSL, Aarhus, Denmark, August 23-29, 1997, Selected Papers*, pages 56–77, 1997.
- [Bou09] Pierre Boudes. Thick subtrees, games and experiments. In *Typed Lambda Calculi and Applications, 9th International Conference, TLCA 2009, Brasilia, Brazil, July 1-3, 2009. Proceedings*, pages 65–79, 2009.
- [Cas17] Simon Castellan. *Concurrent structures in game semantics. (Structures concurrentes en sémantique des jeux)*. PhD thesis, University of Lyon, France, 2017.
- [CC21] Simon Castellan and Pierre Clairambault. Disentangling parallelism and interference in game semantics. 2021. Submitted.
- [CCPW18] Simon Castellan, Pierre Clairambault, Hugo Paquet, and Glynn Winskel. The concurrent game semantics of probabilistic PCF. In Anuj Dawar and Erich Grädel, editors, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018, Oxford, UK, July 09-12, 2018*, pages 215–224. ACM, 2018.
- [CCRW17] Simon Castellan, Pierre Clairambault, Silvain Rideau, and Glynn Winskel. Games and strategies as event structures. *Log. Methods Comput. Sci.*, 13(3), 2017.
- [CCW19] Simon Castellan, Pierre Clairambault, and Glynn Winskel. Thin games with symmetry and concurrent hyland-ong games. *Log. Methods Comput. Sci.*, 15(1), 2019.
- [CdV20] Pierre Clairambault and Marc de Visme. Full abstraction for the quantum lambda-calculus. *Proc. ACM Program. Lang.*, 4(POPL):63:1–63:28, 2020.
- [Cla19] Pierre Clairambault. A tale of additives and concurrency in game semantics. 2019. Submitted.
- [Cla20] Pierre Clairambault. Learning to count up to symmetry. *CoRR*, abs/2006.05080, 2020.
- [EPT11] Thomas Ehrhard, Michele Pagani, and Christine Tasson. The computational meaning of probabilistic coherence spaces. In *Proceedings of the 26th Annual IEEE Symposium on Logic in Computer Science, LICS 2011, June 21-24, 2011, Toronto, Ontario, Canada*, pages 87–96. IEEE Computer Society, 2011.
- [EPT18] Thomas Ehrhard, Michele Pagani, and Christine Tasson. Full abstraction for probabilistic PCF. *J. ACM*, 65(4):23:1–23:44, 2018.
- [FGHW08] Marcelo Fiore, Nicola Gambino, Martin Hyland, and Glynn Winskel. The cartesian closed bicategory of generalised species of structures. *Journal of the London Mathematical Society*, 77(1):203–220, 2008.
- [Gir88] Jean-Yves Girard. Normal functors, power series and λ -calculus. *Ann. Pure Appl. Log.*, 37(2):129–177, 1988.
- [Gir89] Jean-Yves Girard. Geometry of interaction 1: Interpretation of system f. In *Studies in Logic and the Foundations of Mathematics*, volume 127, pages 221–260. Elsevier, 1989.
- [GS16] Richard Garner and Michael Shulman. Enriched categories as a free cocompletion. *Advances in Mathematics*, 289:1–94, 2016.
- [HO00] J. M. E. Hyland and C.-H. Luke Ong. On full abstraction for PCF: I, II, and III. *Inf. Comput.*, 163(2):285–408, 2000.
- [Lai20] James Laird. Weighted models for higher-order computation. *Inf. Comput.*, 275:104645, 2020.
- [Lam92] François Lamarche. Quantitative domains and infinitary algebras. *Theor. Comput. Sci.*, 94(1):37–62, 1992.
- [LMMP13] Jim Laird, Giulio Manzonetto, Guy McCusker, and Michele Pagani. Weighted relational models of typed lambda-calculi. In *28th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2013, New Orleans, LA, USA, June 25-28, 2013*, pages 301–310. IEEE Computer Society, 2013.
- [LS88] Joachim Lambek and Philip J Scott. *Introduction to higher-order categorical logic*, volume 7. Cambridge University Press, 1988.
- [Mel05] Paul-André Melliès. Asynchronous games 4: A fully complete model of propositional linear logic. In *20th IEEE Symposium on Logic in Computer Science (LICS 2005), 26-29 June 2005, Chicago, IL, USA, Proceedings*, pages 386–395. IEEE Computer Society, 2005.
- [Mel09] Paul-André Melliès. Categorical semantics of linear logic. Panoramas et Synthèses, 2009.
- [ML13] Saunders Mac Lane. *Categories for the working mathematician*, volume 5. Springer Science & Business Media, 2013.

- [MT16] Andrzej S. Murawski and Nikos Tzevelekos. An invitation to game semantics. *ACM SIGLOG News*, 3(2):56–67, 2016.
- [OA20] Federico Olimpieri and Lionel Vaux Auclair. On the taylor expansion of λ -terms and the groupoid structure of their rigid approximants. *CoRR*, abs/2008.02665, 2020.
- [Ong17] C.-H. Luke Ong. Quantitative semantics of the lambda calculus: Some generalisations of the relational model. In *32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017, Reykjavik, Iceland, June 20-23, 2017*, pages 1–12. IEEE Computer Society, 2017.
- [Paq20] Hugo Paquet. *Probabilistic concurrent game semantics*. PhD thesis, PhD thesis, University of Cambridge, 2020.
- [RW11] Silvain Rideau and Glynn Winskel. Concurrent strategies. In *Proceedings of the 26th Annual IEEE Symposium on Logic in Computer Science, LICS 2011, June 21-24, 2011, Toronto, Ontario, Canada*, pages 409–418. IEEE Computer Society, 2011.
- [Ulm68] Friedrich Ulmer. Properties of dense and relative adjoint functors. *Journal of Algebra*, 8(1):77–95, 1968.
- [Win86] Glynn Winskel. Event structures. In *Advances in Petri Nets*, pages 325–392, 1986.
- [Win07] Glynn Winskel. Event structures with symmetry. *Electron. Notes Theor. Comput. Sci.*, 172:611–652, 2007.

APPENDIX A. POSTPONED PROOFS AND CONSTRUCTIONS

A.1. Theory of relative adjunctions and comonads. We first recall the basic theory of relative adjunctions, and relative comonads. These are defined with respect to any functor $J : \mathcal{C} \rightarrow \mathcal{D}$, but we only give the special case where \mathcal{C} is a full subcategory of \mathcal{D} , and $J : \mathcal{C} \hookrightarrow \mathcal{D}$ is the inclusion functor. The general definitions can be found in e.g. [ACU10].

Relative adjunctions. If $F : \mathcal{C} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{D}$, we say that F is a **J -relative left adjoint** to G if for every $C \in \mathcal{C}$ and $B \in \mathcal{B}$ there is a natural bijection

$$\mathcal{B}(F(C), B) \cong \mathcal{D}(C, G(B)).$$

We say that F is a **J -relative right adjoint** to G if for all $C \in \mathcal{C}$ and $B \in \mathcal{B}$ there is

$$\mathcal{B}(B, F(C)) \cong \mathcal{D}(G(B), C).$$

a natural bijection. These two situations are respectively pictured as the diagrams below:

$$\begin{array}{ccc}
 & \mathcal{B} & \\
 F \nearrow & \dashv & \searrow G \\
 \mathcal{C} & \xrightarrow{\quad} & \mathcal{D}
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \mathcal{B} & \\
 F \nearrow & \vdash & \searrow G \\
 \mathcal{C} & \xrightarrow{\quad} & \mathcal{D}
 \end{array}
 \tag{A.1}$$

Note that this definition is asymmetric: if F is a right adjoint to G relative to J , then it does *not* make sense to say that G is a J -left adjoint to F .

Relative comonads. A **J -relative comonad** consists of: (1) for every $C \in \mathcal{C}$, an object $!C \in \mathcal{D}$; (2) for every $C \in \mathcal{C}$, a morphism $\epsilon_C : !C \rightarrow C$; and (3) for every $B, C \in \mathcal{C}$ and $f : !B \rightarrow C$, a morphism $f^\dagger : !B \rightarrow !C$, such that for $A, B, C \in \mathcal{C}$,

- (1) if $f : !B \rightarrow C$, then $f = \epsilon_C \circ f^\dagger$,
- (2) $\epsilon_C^\dagger = \text{id}_{!C}$,
- (3) if $f : !A \rightarrow B$ and $g : !B \rightarrow C$, then $(g \circ f^\dagger)^\dagger = g^\dagger \circ f^\dagger$.

The axioms ensure that $!$ can be extended to a functor, sending $f : B \rightarrow C$ to $(f \circ \epsilon_B)^\dagger$.

The Kleisli category of a J -relative comonad. The relationship between adjunctions and comonads extends to the J -relative setting. For F and G as in the right-hand diagram in (A.1), their composite GF is a relative comonad. Conversely, any relative comonad has an associated **Kleisli category** $\mathcal{C}_!$ which can be used to construct a relative adjunction: it has objects those of \mathcal{C} , and homsets given by $\mathcal{C}_!(B, C) = \mathcal{D}(!B, C)$. We have a situation

$$\begin{array}{ccc} & \mathcal{C}_! & \\ \nearrow & \vdash & \searrow \\ \mathcal{C} & \xrightarrow{\quad} & \mathcal{D} \end{array}$$

where the right adjoint is identity-on-objects and maps $f : B \rightarrow C$ to $f \circ \epsilon_B$, and the left adjoint maps $C \in \mathcal{C}_!$ to $!C \in \mathcal{D}$ and $f \in \mathcal{C}_!(B, C)$ to $f^! \in \mathcal{D}(!B, !C)$.

A.2. The Kleisli category of a relative Seely category. For a relative Seely category \mathcal{C} as in Definition 3.37, the Kleisli category for the relative comonad $!$ is cartesian closed. We give some details of the proof. Recall that the situation is the following:

$$\begin{array}{ccc} & \mathcal{C}_! & \\ \nearrow F & \vdash & \searrow G \\ \mathcal{C}_s & \xrightarrow{\quad} & \mathcal{C} \end{array}$$

where F and G are defined in the last paragraph of the previous section.

Products. It is easy to show that J -relative right adjoints preserve the limits that J preserves, see e.g. [Ulm68]. Since J preserves products, $\mathcal{C}_!$ has all finite products constructed as in \mathcal{C}_s , with projections $\pi_A \circ \epsilon_{A \& B} \in \mathcal{C}_!(A \& B, A)$ and $\pi_B \circ \epsilon_{A \& B} \in \mathcal{C}_!(A \& B, B)$.

Cartesian closure. For $A, B \in \mathcal{C}_s$, we define the function space $A \Rightarrow B = !A \multimap B$; we know this is an object of \mathcal{C}_s since by definition the functor $!A \multimap -$ has type $\mathcal{C}_s \rightarrow \mathcal{C}_s$. We use the (relative) closed structure of \mathcal{C} to derive the required bijection. For $A, B, C \in \mathcal{C}_s$,

$$\begin{aligned} \mathcal{C}_!(A \& B, C) &= \mathcal{C}(! (A \& B), C) \\ &\cong \mathcal{C}(!A \otimes !B, C) && \text{(using } m_{A,B} \text{)} \\ &\cong \mathcal{C}(!A, !B \multimap C) = \mathcal{C}_!(A, B \Rightarrow C) \end{aligned}$$

and this is natural. The evaluation map $\text{ev}_{A,B} \in \mathcal{C}_!((A \Rightarrow B) \& A, B)$ is given by

$$!((A \Rightarrow B) \& A) \xrightarrow{m_{A,B}^{-1}} !(A \Rightarrow B) \otimes !A \xrightarrow{\epsilon_{A \Rightarrow B} \otimes !A} !A \multimap B \otimes !A \xrightarrow{\text{ev}_{!A,B}} B.$$

A.3. Synchronization up to Symmetry. First we include the proof of:

Proposition A.1. *Consider $\sigma : A \vdash B$ and $\tau : B \vdash C$ two strategies.*

For $x^\sigma \in \mathcal{C}^+(\sigma)$, $x^\tau \in \mathcal{C}^+(\tau)$ and $\theta : x_B^\sigma \cong_B x_B^\tau$ s.t. the composite bijection is secured:

$$x^\sigma \parallel x_C^\tau \xrightarrow{\partial_\sigma \parallel C} x_A^\sigma \parallel x_B^\sigma \parallel x_C^\tau \xrightarrow{A \parallel \theta} x_A^\sigma \parallel x_B^\tau \parallel x_C^\tau \xrightarrow{A \parallel \partial_\tau^{-1}} x_A^\sigma \parallel x^\tau,$$

then there are (necessarily unique) $y^\sigma \in \mathcal{C}^+(\sigma)$ and $y^\tau \in \mathcal{C}^+(\tau)$ causally compatible, and

$$\varphi^\sigma : x^\sigma \cong_\sigma y^\sigma, \quad \varphi^\tau : x^\tau \cong_\tau y^\tau,$$

such that we have $\varphi_A^\sigma \in \mathcal{S}_-(A)$, $\varphi_C^\tau \in \mathcal{S}_+(C)$, and $\varphi_B^\tau \circ \theta = \varphi_B^\sigma$.

Proof. Let us write A_- for A with symmetry replaced with $\mathcal{S}(A_-) = \mathcal{S}_-(A)$, $\mathcal{S}_-(A_-) = \mathcal{S}_-(A)$, and $\mathcal{S}_+(A_-)$ reduced to identity bijections. Likewise, C_+ is C with isomorphism families restricted to positive symmetries. Then, then we consider the maps of essp

$$(\partial_\sigma \parallel \text{id}_C) : \sigma \parallel C_+ \rightarrow A^\perp \parallel B \parallel C^\perp \quad (\text{id}_A \parallel \partial_\tau) : A_- \parallel \tau \rightarrow A \parallel B^\perp \parallel C$$

are dual *pre-~-strategies* in the sense of [CCW19]. *Existence* then follows directly from an application of Lemma 3.23 of [CCW19] to these two. For *uniqueness*, consider $z^\sigma \in \mathcal{C}^+(\sigma)$ and $z^\tau \in \mathcal{C}^+(\tau)$ causally compatible together with $\psi^\sigma : z^\sigma \cong_\sigma x^\sigma$ and $\psi^\tau : z^\tau \cong_\tau x^\tau$ with $\psi_A^\sigma \in \mathcal{S}_-(A)$, $\psi_C^\tau \in \mathcal{S}_+(C)$ and $\psi_B^\tau \circ \theta = \psi_B^\sigma$. Then,

$$\partial_\sigma(\psi^\sigma \circ (\varphi^\sigma)^{-1}) = \theta_A^- \parallel (\psi_B^\sigma \circ (\varphi_B^\sigma)^{-1}) \quad \partial_\tau(\psi^\tau \circ (\varphi^\tau)^{-1}) = (\psi_B^\tau \circ (\varphi_B^\tau)^{-1}) \parallel \theta_C^+$$

where $\psi_B^\sigma \circ (\varphi_B^\sigma)^{-1} = \psi_B^\tau \circ \theta \circ \theta^{-1} \circ (\varphi_B^\tau)^{-1} = \psi_B^\tau \circ (\varphi_B^\tau)^{-1}$, so by Proposition 3.28,

$$\omega = (\psi^\sigma \circ (\varphi^\sigma)^{-1}) \odot (\psi^\tau \circ (\varphi^\tau)^{-1}) \in \mathcal{S}^+(\tau \odot \sigma)$$

but its image by $\partial_{\tau \odot \sigma}$ is a positive symmetry, so ω is an identity symmetry by Lemma 3.28 of [CCW19]. It follows easily from Proposition 3.28 that $\psi^\sigma = (\varphi^\sigma)^{-1}$ and $\psi^\tau = (\varphi^\tau)^{-1}$. \square

We can also prove the same property on *symmetries* rather than configurations. For this, we use *higher symmetries* on ess: if $\theta, \theta' \in \mathcal{S}(E)$, we write $\Theta : \theta \cong_E \theta'$ for a bijection between their graphs, such that writing $\text{dom}(\Theta) = \{(a_1, b_1) \mid ((a_1, a_2), (b_1, b_2)) \in \Theta\}$,

$$\text{dom}(\Theta) : \text{dom}(\theta) \cong_E \text{dom}(\theta')$$

and likewise for $\text{cod}(\Theta)$.

Proposition A.2. *Consider $\sigma : A \vdash B$ and $\tau : B \vdash C$ two strategies.*

For $\theta^\sigma \in \mathcal{S}^+(\sigma)$, $\theta^\tau \in \mathcal{S}^+(\tau)$ and $\Theta : \theta_B^\sigma \cong_B \theta_B^\tau$ s.t. the composite bijection is secured:

$$\theta^\sigma \parallel \theta_C^\tau \xrightarrow{\partial_\sigma \parallel C} \theta_A^\sigma \parallel \theta_B^\sigma \parallel \theta_C^\tau \xrightarrow{A \parallel \Theta \parallel C} \theta_A^\sigma \parallel \theta_B^\tau \parallel \theta_C^\tau \xrightarrow{A \parallel \partial_\tau^{-1}} \theta_A^\sigma \parallel \theta^\tau,$$

then there are (necessarily unique) $\vartheta^\sigma \in \mathcal{S}^+(\sigma)$ and $\vartheta^\tau \in \mathcal{S}^+(\tau)$ causally compatible, and

$$\Phi^\sigma : \theta^\sigma \cong_\sigma \vartheta^\sigma, \quad \Phi^\tau : \theta^\tau \cong_\tau \vartheta^\tau,$$

s.t. Φ_A^σ is negative (i.e. $\text{dom}(\Phi_A^\sigma)$ and $\text{cod}(\Phi_A^\sigma)$ negative), Φ_C^τ is positive, and $\Phi_B^\tau \circ \Theta = \Phi_B^\sigma$.

Proof. Let us write $\theta^\sigma : x^\sigma \cong_\sigma y^\sigma$ and $\theta^\tau : x^\tau \cong_\tau y^\tau$. Applying Proposition 4.4, we get

$$\varphi^\sigma : x^\sigma \cong_\sigma u^\sigma \quad \varphi^\tau : x^\tau \cong_\tau u^\tau \quad \psi^\sigma : y^\sigma \cong_\sigma v^\sigma \quad \psi^\tau : y^\tau \cong_\tau v^\tau$$

where u^σ, u^τ causally compatible, v^σ, v^τ causally compatible, and satisfying additional properties not listed here. We may then define ϑ^σ and ϑ^τ as the missing sides of:

$$\begin{array}{ccc} x^\sigma & \xrightarrow{\varphi^\sigma} & u^\sigma \\ \theta^\sigma \downarrow & & \\ y^\sigma & \xrightarrow{\psi^\sigma} & v^\sigma \end{array} \quad \begin{array}{ccc} x^\tau & \xrightarrow{\varphi^\tau} & u^\tau \\ \theta^\tau \downarrow & & \\ y^\tau & \xrightarrow{\psi^\tau} & v^\tau \end{array}$$

and $\Phi^\sigma : \theta^\sigma \cong_\sigma \vartheta^\sigma$ and $\Phi^\tau : \theta^\tau \cong_\tau \vartheta^\tau$ induced by those commuting diagrams. It is then a simple diagram chasing that the additional properties are satisfied. \square

A.4. Horizontal Composition of Positive Isomorphisms. Next we detail the proof of:

Proposition A.3. *Let $\sigma, \sigma' \in \text{Strat}(A, B)$; $\tau, \tau' \in \text{Strat}(B, C)$; $\varphi : \sigma \approx \sigma'$, $\psi : \tau \approx \tau'$.*

Then, there exists a positive isomorphism $\psi \odot \varphi : \tau \odot \sigma \approx \tau' \odot \sigma'$ such that for all $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$, writing $y^{\tau'} \odot y^{\sigma'} = (\psi \odot \varphi)(x^\tau \odot x^\sigma) \in \mathcal{C}^+(\tau' \odot \sigma')$, we have

$$\varphi(x^\sigma) \cong_{\sigma'} y^{\sigma'}, \quad \psi(x^\tau) \cong_{\tau'} y^{\tau'}.$$

Proof. We define $\psi \odot \varphi$ on $+$ -covered configurations. Take $x^\tau \odot x^\sigma \in \mathcal{C}^+(\tau \odot \sigma)$. There is, of course, no reason why $\varphi(x^\sigma)$ and $\psi(x^\tau)$ would be compatible. However, since φ and ψ are positive isomorphisms, there are (unique) symmetries $\theta_A^-, \theta_B^+, \theta_B^-, \theta_C^+$ such that

$$\begin{array}{ccc} x^\sigma & \xrightarrow{\partial_\sigma} & x_A^\sigma \parallel x_B^\sigma \\ \varphi \downarrow & & \downarrow \theta_A^- \parallel \theta_B^+ \\ \varphi(x^\sigma) & \xrightarrow{\partial_{\sigma'}} & \varphi(x^\sigma)_A \parallel \varphi(x^\sigma)_B \end{array} \quad \begin{array}{ccc} x^\tau & \xrightarrow{\partial_\tau} & x_B^\tau \parallel x_C^\tau \\ \psi \downarrow & & \downarrow \theta_B^- \parallel \theta_C^+ \\ \psi(x^\tau) & \xrightarrow{\partial_{\tau'}} & \psi(x^\tau)_B \parallel \psi(x^\tau)_C \end{array}$$

commute. We show that there are unique $y^{\tau'} \odot y^{\sigma'} \in \mathcal{C}^+(\tau' \odot \sigma')$ and symmetries

$$\omega : \varphi(x^\sigma) \cong_{\sigma'} y^{\sigma'} \quad \nu : \psi(x^\tau) \cong_{\tau'} y^{\tau'}$$

such that $\omega_A \in \mathcal{S}_-(A)$, $\nu_C \in \mathcal{S}_+(C)$ and $\omega_B \circ \theta_B^+ = \nu_B \circ \theta_B^-$.

Existence. We get $\theta_B = \theta_B^- \circ (\theta_B^+)^{-1} : \varphi(x^\sigma)_B \cong_B \psi(x^\tau)_B$ a mediating symmetry between $\varphi(x^\sigma)$ and $\psi(x^\tau)$ and from the two diagrams above we easily deduce that

$$\begin{array}{ccccccc} \varphi \parallel \theta_C^+ \swarrow & x^\sigma \parallel x_C^\tau & \xrightarrow{\partial_\sigma \parallel x_C^\tau} & x_A^\sigma \parallel x_B \parallel x_C^\tau & \xrightarrow{x_A^\sigma \parallel \partial_\tau^{-1}} & x_A^\sigma \parallel x^\tau & \xrightarrow{\theta_A^- \parallel \psi} \\ z^{\sigma'} \parallel z_C^{\tau'} & \xrightarrow{\partial_{\sigma'} \parallel z_C^{\tau'}} & z_A^{\sigma'} \parallel z_B^{\sigma'} \parallel z_C^{\tau'} & \xrightarrow{z_A^{\sigma'} \parallel \theta_B \parallel z_C^{\tau'}} & z_A^{\sigma'} \parallel z_B^{\tau'} \parallel z_C^{\tau'} & \xrightarrow{z_A^{\sigma'} \parallel \partial_\tau^{-1}} & z_A^{\sigma'} \parallel z^{\tau'} \end{array}$$

commutes, writing $z^{\sigma'} = \varphi(x^\sigma)$ and $z^{\tau'} = \psi(x^\tau)$. As the (bijection induced by) the top row is secured and φ, ψ are order-isomorphisms, it follows that the (bijection induced by) the bottom row is also secured. Therefore, applying Proposition 4.4, it follows that there are

$$y^{\tau'} \odot y^{\sigma'} \in \mathcal{C}^+(\tau' \odot \sigma') \quad \omega : z^{\sigma'} \cong_{\sigma'} y^{\sigma'} \quad \nu : z^{\tau'} \cong_{\tau'} y^{\tau'}$$

such that $\omega_A \in \mathcal{S}_-(A)$, $\nu_C \in \mathcal{S}_+(C)$ and $\nu_B \circ \theta_B = \omega_B$ as required. For *uniqueness*, if

$$u^{\tau'} \odot u^{\sigma'} \in \mathcal{C}^+(\tau' \odot \sigma') \quad \mu : z^{\sigma'} \cong_{\sigma'} u^{\sigma'} \quad \gamma : z^{\tau'} \cong_{\tau'} u^{\tau'}$$

then $(\mu \circ \omega^{-1}) \odot (\gamma \circ \nu^{-1}) \in \mathcal{S}(\tau' \odot \sigma')$ displays to a positive symmetry of $A \vdash C$, so is an identity by Lemma 3.28 of [CCW19]. By Proposition 3.28, $\mu = \omega$ and $\gamma = \nu$.

Now, we may set $(\psi \odot \varphi)(x^\tau \odot x^\sigma) = y^{\tau'} \odot y^{\sigma'}$. To prove preservation of symmetry, we perform the exact same construction on symmetries, using Proposition A.2, which commutes with domain and codomain. The inverse $(\psi \odot \varphi)^{-1}$ is constructed similarly. The fact that these are inverses and their monotonicity are direct consequences of the uniqueness of the construction above. Finally, any order-isomorphism preserving symmetry between ess is generated by a unique isomorphism of ess, see *e.g.* Lemma D.4 from [CC21]. \square

A.5. Invariance of \sim^+ -witnesses. We show that the cardinality of \sim^+ -witnesses do not depend on the choice of representative.

Proposition A.4. *Consider A, B arenas, $\sigma : A \vdash B$, and $x_A \in \mathcal{C}_{\cong}^0(A), x_B \in \mathcal{C}_{\cong}^0(B)$.*

Then, the cardinality of $\sim^{\pm}\text{-wit}_{\sigma}^{\pm}(x_A, x_B)$ does not depend on $x_A \in \mathbf{x}_A$ and $x_B \in \mathbf{x}_B$.

Proof. We show the following. Consider A a game, $\sigma : A$ a strategy. For any $x_A \in \mathcal{C}(A)$, set $\sim^{\pm}\text{-wit}_{\sigma}(x_A) = \{(x^{\sigma}, \psi^{\pm}) \mid \psi^{\pm} : x_A^{\sigma} \cong_A x_A\}$. Then, for any $x_A \cong_A y_A$, we have a bijection $\sim^{\pm}\text{-wit}_{\sigma}(x_A) \simeq \sim^{\pm}\text{-wit}_{\sigma}(y_A)$. Indeed, fix $\theta : x_A \cong_A y_A$, which can be factored uniquely as $\theta^+ \circ \theta^-$ and as $\vartheta^- \circ \vartheta^+$ by Lemma 4.10. Now, given $(x^{\sigma}, \psi^{\pm}) \in \sim^{\pm}\text{-wit}_{\sigma}(x_A)$, then there are unique $y^{\sigma} \in \mathcal{C}(\sigma)$, $\varphi^{\sigma} : x^{\sigma} \cong_{\sigma} y^{\sigma}$, $\omega^+ : y_A^{\sigma} \cong_A^+ z_A$ s.t. the following diagram commutes.

$$\begin{array}{ccccc} x_A^{\sigma} & \xrightarrow{\psi^+} & x_A & \xrightarrow{\vartheta^+} & z_A' \\ \varphi_A^{\sigma} \downarrow & & \downarrow \theta^- & & \downarrow \vartheta^- \\ y_A^{\sigma} & \xrightarrow{\omega^+} & z_A & \xrightarrow{\theta^+} & y_A \end{array}$$

For *existence*, refactor $\theta^- \circ \psi^+ = \Theta^+ \circ \Theta^-$ by Lemma 4.10, say $\Theta^- : x_A^{\sigma} \cong_A^- u_A$ and $\Theta^+ : u_A \cong_A^+ z_A$. By Lemma 4.15, there are unique $\varphi^{\sigma} : x^{\sigma} \cong_{\sigma} y^{\sigma}$ and $\Omega^+ : y_A^{\sigma} \cong_A^+ u_A$ such that $\Omega^+ \circ \varphi_A^{\sigma} = \Theta^-$. Setting $\omega^+ = \Theta^+ \circ \Omega^+$ satisfies our constraints.


Uniqueness. If we also have $y^{\sigma'}, \varphi^{\sigma'}$, and $\omega^{+'}$ satisfying those constraints, then $\varphi^{\sigma} \circ (\varphi^{\sigma'})^{-1} \in \mathcal{S}(\sigma)$ maps to the identity which is a positive symmetry, so must be an identity by Lemma 4.6. It follows that $y^{\sigma} = y^{\sigma'}$, $\varphi^{\sigma} = \varphi^{\sigma'}$, and by necessity $\omega^+ = \omega^{+'}$ as well.

This yields a construction from $\sim^{\pm}\text{-wit}_{\sigma}(x_A)$ to $\sim^{\pm}\text{-wit}_{\sigma}(y_A)$. Note that the construction is symmetric and may be applied from $\sim^{\pm}\text{-wit}_{\sigma}(y_A)$ to $\sim^{\pm}\text{-wit}_{\sigma}(x_A)$ as well via θ^{-1} . That the two constructions are inverse follows immediately from the uniqueness property. \square

APPENDIX B. NOT EVERY GAME IS REPRESENTABLE

The following counter-example is due to Marc de Visme.

Example B.1. Consider the tcg A , with events, polarities, and causality and follows:



Its symmetry comprises all order-isomorphisms between configurations. The negative symmetry has all order-isomorphisms included in one of the two maximal bijections

$$\begin{array}{ccc} \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_1 & \oplus_2 \end{array} & \cong_A^- & \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_1 & \oplus_2 \end{array} & \cong_A^- & \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_2 & \oplus_1 \end{array} \end{array}$$

where again, the bijection matches those events in the corresponding position of the diagram. Likewise, the positive symmetry has all order-isomorphisms included in one of:

$$\begin{array}{ccc} \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_1 & \oplus_2 \end{array} & \cong_A^+ & \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_1 & \oplus_2 \end{array} & \cong_A^+ & \begin{array}{cc} \ominus_1 & \ominus_2 \\ \vdots & \vdots \\ \oplus_2 & \oplus_1 \end{array} \end{array}$$

forming, altogether, a tcg. Then, the endosymmetry

$$\begin{array}{ccc} \ominus_1 & & \ominus_2 \\ \vdots & & \vdots \\ & \oplus_1 & \\ & \cong_A & \\ & & \oplus_1 \\ & & \ominus_2 & & \ominus_1 \end{array}$$

which is neither positive nor negative, uniquely factors as

$$\begin{array}{ccc} \ominus_1 & & \ominus_2 \\ \vdots & & \vdots \\ & \oplus_1 & \\ & \cong_A^- & \\ & & \oplus_2 & & \cong_A^+ & \\ & & & & & \oplus_1 \\ & & & & & \ominus_2 & & \ominus_1 \end{array}$$

which is not formed of endosymmetries. So this configuration is not canonical, but its only symmetric $\{\ominus_1, \ominus_2, \oplus_2\}$ is not canonical either, for the same reason.

APPENDIX C. FURTHER CONTENT ON GROUPOIDS OF STRATEGIES

C.1. On Weights of Symmetry Classes. How should one correct the sum, if one is to count symmetry classes instead of positive witnesses? Let us fix A a game, $\sigma : A$ any strategy, and $x_A \in \mathcal{C}_{\cong}^0(A)$. We show how negative symmetries act on \sim^+ -witnesses of x_A .

Proposition C.1. *For any $(x^\sigma, \theta^+) \in \sim^{\pm}\text{-wit}_\sigma(x_A)$ and $\varphi^- \in \mathcal{S}_-(x_A)$ there are unique $(y^\sigma, \psi^+) \in \sim^{\pm}\text{-wit}_\sigma(x_A)$ and $\phi : x^\sigma \cong_\sigma y^\sigma$ such that the following diagram commutes:*

$$\begin{array}{ccc} x_A^\sigma & \xrightarrow{\theta^+} & x_A \\ \phi_A^\sigma \downarrow & & \downarrow \varphi^- \\ y_A^\sigma & \xrightarrow{\psi^+} & x_A \end{array}$$

Proof. Consider $(x^\sigma, \theta^+) \in \sim^{\pm}\text{-wit}(x_A)$ and $\varphi^- \in \mathcal{S}_-(x_A)$. We show that there is unique $\phi^\sigma : x^\sigma \cong_\sigma y^\sigma$ and $\psi^+ : y_A^\sigma \cong_A^+ x_A$ making the following diagram commute:

$$\begin{array}{ccc} x_A^\sigma & \xrightarrow{\theta^+} & x_A \\ \phi_A^\sigma \downarrow & & \downarrow \varphi^- \\ y_A^\sigma & \xrightarrow{\psi^+} & x_A \end{array}$$

For existence, by Lemma 4.10, $\varphi^- \circ \theta^+ : x_A^\sigma \cong_A x_A$ factors uniquely as $\Xi^+ \circ \Xi^- : x_A^\sigma \cong_A x_A$. Next, by Lemma 4.15, there is $\phi^\sigma : x^\sigma \cong_\sigma y^\sigma$ such that we have

$$\phi_A^\sigma = \Omega^+ \circ \Xi^- : x_A^\sigma \cong_A y_A^\sigma$$

for some $\Omega^+ : y_A \cong_A^+ y_A^\sigma$. We then form $\psi^+ = \Xi^+ \circ (\Omega^+)^{-1}$ to conclude.

For uniqueness, if we have $\varphi_1 : x^\sigma \cong_\sigma y^\sigma$ and $\varphi_2 : x^\sigma \cong_\sigma z^\sigma$ satisfying the requirements,

$$\begin{array}{ccccc} y_A^\sigma & \xrightarrow{(\sigma\varphi_1)^{-1}} & x_A^\sigma & \xrightarrow{\sigma\varphi_2} & z_A^\sigma \\ +\downarrow & & +\downarrow & & \downarrow + \\ x_A & \xrightarrow{(\varphi^-)^{-1}} & x_A & \xrightarrow{\varphi^-} & x_A \end{array}$$

commutes, so $(\sigma\varphi_2) \circ (\sigma\varphi_1)^{-1} = \sigma(\varphi_2 \circ \varphi_1^{-1})$ is positive, so by Lemma 3.28 of [CCW19] we have $\varphi_2 \circ \varphi_1^{-1} = \text{id}$, so $\varphi_1 = \varphi_2$. \square

It follows easily that there is a group action

$$(- \curvearrowright -) : \mathcal{S}_-(\mathbf{x}_A) \times \sim^\pm \text{wit}(\mathbf{x}_A) \rightarrow \sim^\pm \text{wit}(\mathbf{x}_A),$$

though we shall not use this specifically.

Next, we show that representatives of symmetry classes of configurations in σ can always be chosen to be positively symmetric to the chosen representative in the game.

Lemma C.2. *Consider $x^\sigma \in \text{wit}_\sigma^\cong(\mathbf{x}_A)$. Then, there exists $x^\sigma \in \mathbf{x}^\sigma$ such that $\partial_\sigma x^\sigma \cong_A^+ \underline{\mathbf{x}}_A$.*

Proof. By hypothesis, $\partial_\sigma x^\sigma = x_A^\sigma \in \mathbf{x}_A$, so there exists

$$x_A^\sigma \cong_A^\theta \underline{\mathbf{x}}_A$$

which factors uniquely as $x_A^\sigma \cong_A^{\theta^-} y_A \cong_A^{\theta^+} \underline{\mathbf{x}}_A$ by Lemma 4.10. But then, by Lemma 4.15, there is $\varphi : x^\sigma \cong_\sigma y^\sigma$ and $\psi^+ : y_A \cong_A^+ y_A^\sigma$ such that

$$\partial_\sigma \varphi = \psi^+ \circ \theta^- : x_A^\sigma \cong_A y_A^\sigma,$$

so that in particular $y^\sigma \in \mathbf{x}^\sigma$ and $y_A^\sigma \cong_A^+ y_A \cong_A^+ \underline{\mathbf{x}}_A$. \square

So, for each $x^\sigma \in \text{wit}_\sigma^\cong(\mathbf{x}_A)$ we fix a representative \underline{x}^σ such that $\underline{x}_A^\sigma \cong_A^+ \underline{\mathbf{x}}_A$. We also choose a reference $\theta_{\underline{x}^\sigma}^+ : \underline{x}_A^\sigma \cong_A^+ \underline{\mathbf{x}}_A$. Finally, for every $x^\sigma \in \mathbf{x}^\sigma$ we choose $\kappa_{x^\sigma} : \underline{x}^\sigma \cong_\sigma x^\sigma$.

Our aim is, for every symmetry class $\mathbf{x}^\sigma \in \text{wit}_\sigma^\cong(\mathbf{x}_A)$, count the number of concrete witnesses in \mathbf{x}^σ . We introduce some notations for this set – let us write

$$\begin{aligned} \text{wit}_\sigma^+[\mathbf{x}^\sigma] &= \{x^\sigma \in \text{wit}_\sigma^+(\mathbf{x}_A) \mid x^\sigma \in \mathbf{x}^\sigma\} \\ \sim^\pm \text{wit}_\sigma[\mathbf{x}^\sigma] &= \{x^\sigma \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A) \mid x^\sigma \in \mathbf{x}^\sigma\} \end{aligned}$$

for the concrete witnesses (resp. \sim^+ -witnesses) within a symmetry class $\mathbf{x}^\sigma \in \text{wit}_\sigma^\cong(\mathbf{x}_A)$.

Then, we prove the following bijection:

Proposition C.3. *There is a bijection $\sim^\pm \text{wit}_\sigma[\mathbf{x}^\sigma] \times \mathcal{S}(\mathbf{x}^\sigma) \simeq \mathcal{S}(\mathbf{x}_A)$.*

Proof. First we show that for every $(x^\sigma, \theta^+) \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A)$ and $\varphi \in \mathcal{S}(\mathbf{x}^\sigma)$, there is a unique $\psi \in \mathcal{S}(\mathbf{x}_A)$ such that the following diagram commutes:

$$\begin{array}{ccc} \underline{\mathbf{x}}_A^\sigma & \xrightarrow{\theta_{\underline{x}^\sigma}^+} & \underline{\mathbf{x}}_A \\ (\kappa_{x^\sigma} \circ \varphi)_A \downarrow & & \downarrow \psi \\ x_A^\sigma & \xrightarrow{\theta^+} & \underline{\mathbf{x}}_A \end{array}$$

but this is obvious, as ψ is determined by composition from the other components.

Reciprocally, we show that for all $\psi \in \mathcal{S}(\mathbf{x}_A)$, there are unique $(x^\sigma, \theta^+) \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A)$ and $\varphi \in \mathcal{S}(\mathbf{x}^\sigma)$ such that the same diagram above commutes. First, by canonicity of $\underline{\mathbf{x}}_A$, $\psi = \psi^+ \circ \psi^-$ for $\psi^- \in \mathcal{S}_-(\mathbf{x}_A)$ and $\psi^+ \in \mathcal{S}_+(\mathbf{x}_A)$. By Proposition C.1, there are unique $(y^\sigma, \omega^+) \in \sim^\pm \text{wit}_\sigma(\mathbf{x}_A)$ and $\phi^\sigma : \underline{\mathbf{x}}^\sigma \cong_\sigma y^\sigma$ such that the following diagram commutes:

$$\begin{array}{ccc} \underline{\mathbf{x}}_A^\sigma & \xrightarrow{\theta_{\underline{x}^\sigma}^+} & \underline{\mathbf{x}}_A \\ \phi_A^\sigma \downarrow & & \downarrow \psi^- \\ y_A^\sigma & \xrightarrow{\omega^+} & \underline{\mathbf{x}}_A \end{array}$$

We may then define $x^\sigma := y^\sigma$, $\theta^+ := \psi^+ \circ \omega^+$, and $\varphi := (\kappa_{y^\sigma})^{-1} \circ \phi^\sigma$ and the diagram is obviously satisfied. It remains to prove uniqueness, so assume we have $(z^\sigma, \vartheta^+) \in \sim^{\pm} \text{wit}_\sigma(x_A)$ and $\nu \in \mathcal{S}(x^\sigma)$ such that the following diagram commutes:

$$\begin{array}{ccc} \underline{x}_A^\sigma & \xrightarrow{\theta_{x^\sigma}^+} & \underline{x}_A \\ (\kappa_{z^\sigma \circ \nu})_A \downarrow & & \downarrow \psi \\ z_A^\sigma & \xrightarrow{\vartheta^+} & \underline{x}_A \end{array}$$

But then $(\kappa_{z^\sigma \circ \nu}) \circ (\kappa_{x^\sigma} \circ \varphi^{-1}) \in \mathcal{S}(\sigma)$ displays to a positive symmetry, so must be an identity by Lemma 4.6. Thus $x^\sigma = y^\sigma$, so $\nu = \varphi$, and so $\theta^+ = \vartheta^+$ as it is uniquely determined from the other components by the diagram. This gives constructions in both directions, and that they are inverses follows directly from the uniqueness properties. \square

From that bijection, we may conclude the following result:

Theorem C.4. *Consider A a game, $\sigma : A$ and $x_A \in \mathcal{C}_{\cong}^0(A)$, and $x^\sigma \in \text{wit}_\sigma^{\cong}(x_A)$. Then,*

$$\#\text{wit}_\sigma^+[x^\sigma] = \frac{\#\mathcal{S}_-(x_A)}{\#\mathcal{S}(x^\sigma)}.$$

Proof. By Proposition C.3, we have $\#\sim^{\pm} \text{wit}_\sigma[x^\sigma] \times \#\mathcal{S}(x^\sigma) = \#\mathcal{S}(x_A)$, so we have

$$\#\mathcal{S}_+(x_A) \times \#\text{wit}_\sigma^+[x^\sigma] \times \#\mathcal{S}(x^\sigma) = \#\mathcal{S}_+(x_A) \times \#\mathcal{S}_-(x_A)$$

using Lemma 4.18 and canonicity of \underline{x}_A . The identity follows. \square

This finally lets us state the collapse formula for symmetry classes – below we use that by invariance under symmetry, any the valuation of any \mathcal{R} -strategy lifts canonically to symmetry classes. So we can finally reformulate (6.1) as:

Theorem C.5. *Consider $\sigma \in \mathcal{R}\text{-Strat}(A, B)$ and $x_A \in \mathcal{C}_{\cong}^0(A)$, $x_B \in \mathcal{C}_{\cong}^0(B)$. Then,*

$$f(\sigma)_{x_A, x_B} = \sum_{x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B)} \frac{\#\mathcal{S}_-(x_A)}{\#\mathcal{S}(x^\sigma)} * \mathcal{V}_\sigma(x^\sigma)$$

Proof. We calculate:

$$\begin{aligned} f(\sigma)_{x_A, x_B} &= \sum_{x^\sigma \in \text{wit}_\sigma^+(x_A, x_B)} \mathcal{V}_\sigma(x^\sigma) \\ &= \sum_{x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B)} \sum_{x^\sigma \in \text{wit}_\sigma^+[x^\sigma]} \mathcal{V}_\sigma(x^\sigma) \\ &= \sum_{x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B)} \#\text{wit}_\sigma^+[x^\sigma] * \mathcal{V}_\sigma(x^\sigma) \\ &= \sum_{x^\sigma \in \text{wit}_\sigma^{\cong}(x_A, x_B)} \frac{\#\mathcal{S}_-(x_A)}{\#\mathcal{S}(x^\sigma)} * \mathcal{V}_\sigma(x^\sigma) \end{aligned}$$

\square

C.2. Absorption of Symmetries. As final contribution, we include a property which, though not used for the main results of this paper, was required for the quantum collapse of [CdV20]. As such, we believe it fits with the present development.

For A a game, $\sigma : A$ a strategy, and $x_A \in \mathcal{C}_{\cong}^0(A)$, a variant of the \sim^+ -witnesses is

$$\sim\text{-wit}_{\sigma}^+(x_A) = \{(x^{\sigma}, \theta) \mid x^{\sigma} \in \text{wit}_{\sigma}^+(x_A), \theta : x_A^{\sigma} \cong_A x_A\},$$

so we still consider witnesses $x^{\sigma} \in \mathcal{C}^+(\sigma)$ such that $x_A^{\sigma} \cong_A^+ x_A$ still, but associated with all possible *symmetries*, not only positive symmetries as in $\sim^{\pm}\text{-wit}_{\sigma}(x_A)$. Our last contribution consists in counting $\sim\text{-wit}_{\sigma}^+(x_A)$, compared to $\sim^{\pm}\text{-wit}_{\sigma}(x_A)$. First we need:

Lemma C.6. *Consider A a tcg, $x_A \in \mathcal{C}_{\cong}(A)$, and $x \in \mathcal{C}(A)$ s.t. $x \cong_A^+ x_A$.*

Then, any $\theta : x \cong_A x_A$ factors uniquely as $\theta^- \circ \theta^+$, where $\theta^+ : x \cong_A^+ x_A$, $\theta^- \in \mathcal{S}_-(x_A)$.

Proof. Fix some $\varphi : x \cong_A^+ x_A$. Now, take $\theta : x \cong_A x_A$. By Lemma 4.10, θ factors uniquely as $\theta^- \circ \theta^+$, where $\theta^+ : x \cong_A^+ z$ and $\theta^- : z \cong_A^- x_A$ for some $z \in \mathcal{C}(A)$. But then,

$$\varphi \circ \theta^{-1} : x_A \cong_A x_A$$

factors via $(\varphi \circ (\theta^+)^{-1}) : z \cong_A^+ x_A$ and $(\theta^-)^{-1} : x_A \cong_A^- z$, so $x_A = z$ since x_A is canonical. \square

Corollary C.7. *There is a bijection $\sim\text{-wit}_{\sigma}^+(x_A) \simeq \mathcal{S}_-(x_A) \times \sim^{\pm}\text{-wit}_{\sigma}(x_A)$.*

Proof. First, we show that for all $(x^{\sigma}, \theta) \in \sim\text{-wit}_{\sigma}^+(x_A)$ there are unique $y^{\sigma} \in \mathcal{C}^+(\sigma)$, $\varphi^{\sigma} \in \mathcal{S}(\sigma)$, $\theta^- \in \mathcal{S}_-(A)$, $\theta^+ \in \mathcal{S}_+(A)$ and $\psi^+ \in \mathcal{S}_+(A)$, such that the diagram commutes:

$$\begin{array}{ccc} x_A^{\sigma} & \xrightarrow{\theta^+} & x_A \\ \varphi_A^{\sigma} \downarrow & \searrow \theta & \downarrow \theta^- \\ y_A^{\sigma} & \xrightarrow{\psi^+} & x_A \end{array}$$

By Lemma C.6, θ factors uniquely as claimed. But then, the other components and their uniqueness follows from Proposition C.1. Reciprocally, we show that for all $\theta^- \in \mathcal{S}_-(x_A)$ and $(y^{\sigma}, \psi^+) \in \sim^{\pm}\text{-wit}_{\sigma}(x_A)$, there are unique $x^{\sigma} \in \mathcal{C}(\sigma)$, $\theta \in \mathcal{S}(A)$, $\theta^+ \in \mathcal{S}_+(A)$ and $\varphi^{\sigma} \in \mathcal{S}(\sigma)$ such that the diagram above commutes – but this is again Proposition C.1.

These two constructions immediately provide the two sides of the bijection, and that they are inverses immediately follows from the uniqueness. \square