



**HAL**  
open science

# Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean

Viviane Bout-roumazeilles, N Combourieu-Nebout, S Desprat, G Siani, J-L Turon

## ► To cite this version:

Viviane Bout-roumazeilles, N Combourieu-Nebout, S Desprat, G Siani, J-L Turon. Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean. 2021. hal-03280600

**HAL Id: hal-03280600**

**<https://hal.science/hal-03280600>**

Preprint submitted on 7 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).  
Please refer to the corresponding final paper in CP if available.

# Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean

V. Bout-Roumazielles<sup>1</sup>, N. Combourieu-Nebout<sup>2</sup>, S. Desprat<sup>3</sup>, G. Siani<sup>4</sup>, and J.-L. Turon<sup>5</sup>

<sup>1</sup>Géosystèmes, UMR8217, CNRS – Université Lille 1, 59655 Villeneuve d’Ascq, France

<sup>2</sup>Laboratoire des Sciences du Climat et de l’Environnement, laboratoire Mixte CNRS-CEA-UVSQ, Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France

<sup>3</sup>Ecole Pratique des Hautes Etudes, Environnements et Paléoenvironnements Océaniques, UMR5808, CNRS – EPOC, Université Bordeaux I, 33405 Talence, France

<sup>4</sup>Laboratoire des Interactions et Dynamique des Environnements de Surface (IDES), UMR8148, CNRS – Université de Paris-Sud, Bât 504, 91405 Orsay Cedex, France

<sup>5</sup>Environnements et Paléoenvironnements Océaniques, UMR5808, CNRS – EPOC, Université Bordeaux I, 33405 Talence, France

Received: 12 July 2012 – Accepted: 17 July 2012 – Published: 27 July 2012

Correspondence to: V. Bout-Roumazielles (viviane.bout@univ-lille1.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Atmospheric supply

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

The objectives were to retrace the eolian and fluvial terrigenous supplies in a sediment core from the Sicilian-Tunisian Strait by coupling mineralogical, grain-size and geochemical approaches, in order to get informations on the atmospheric versus riverine contributions to sedimentation on the southern side of central Mediterranean since the last glacial. The eolian supply is dominant over the whole interval, excepted during the sapropel S1 when riverine contribution apparently became significant, and particles provenance has been modified since Last Glacial. Saharan contribution increased during the Bølling-Allerød, evidencing the persistence of aridity over North Africa although the northern Mediterranean already experienced moister and warmer conditions. The Younger Dryas is marked by proximal dust inputs highlighting intense regional eolian activity. A southward migration of dust provenance toward Sahel occurred at the onset of the Holocene, likely resulting from a southward position of the Inter Tropical Convergence Zone, probably associated with a large-scale atmospheric reorganization. Finally, a peculiar high terrigenous flux associated with drastic modifications of the mineralogical and geochemical sediment signature occurred during the sapropel S1, suggesting the propagation of fine-particles derived from major floodings of the Nile River – resulting from enhanced rainfall on northeastern Africa – and their transportation across the Sicilian-Tunisian Strait by intermediate water-masses.

## 1 Introduction

The Mediterranean is a transitional area where northern and southern climatic influences tightly interact (e.g. Magny et al., 2009). Previous studies revealed that moist conditions developed in the Mediterranean during the early Holocene while a progressive orbitally-driven trend to aridification characterized the mid and late Holocene. This climatic evolution displays a contrasting regional pattern, with an abrupt transition at 5.7 kyr on the western Mediterranean and a more gradual transition in the eastern

CPD

8, 2921–2968, 2012

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mediterranean (Cheddadi et al., 1991; Ariztegui et al., 2000; Magny et al., 2002, 2007a; Frigola et al., 2007; Tzedakis, 2007; Sadori et al., 2008; Roberts et al., 2008, 2011; Jalut et al., 2009; Peyron et al., 2011). Precipitation estimations based on lake levels, fire and pollen association, and on speleothem and isotope records, also provide evidences of contrasting seasonality across the Mediterranean during the Holocene (Magny et al., 2003, 2007, 2009; Zanchetta et al., 2007; Tzedakis, 2007; Roberts et al., 2008; Peyron et al., 2011; Vanni re et al., 2011), but estimating the respective atmospheric and oceanic control on Mediterranean climatic evolution through their impact on eolian and fluvial systems is still complex. A multiproxy study of the terrigenous supply would help retracing the variability of both eolian and fluvial systems.

The nature and provenance of fine-grained terrigenous particles in the Mediterranean is mainly controlled by the balance between riverine supplies driven by precipitation regime on the surrounding continents and eolian supplies from the Sahara (Bergametti et al., 1989b; Matthewson et al., 1995; Guerzoni and Chester, 1996; Foucault and M li res, 2000; Goudie and Middleton, 2001). Several studies used deep-sea clay mineral associations as tracers of source regions and as indicators of water mass fluctuations (Chamley, 1975; Petschick et al., 1996; Fagel et al., 1997, 2006; Gingele et al., 2001; Liu et al., 2003; Boulay et al., 2005; Colin et al., 2010). Indeed, the mineralogical nature of sediments, which depends on the petrographic characteristics of their source areas (e.g., Bout-Roumazeilles et al., 1999; Sionneau et al., 2008), has been used to retrace detrital particles provenance in the Mediterranean and thus to assess the respective eolian and riverine contributions to deep-sea sedimentation and provide valuable information on the sediment propagation pathways (Caquineau et al., 1998, 2002; Bout-Roumazeilles et al., 2007; Erhmann et al., 2007a; Hamann et al., 2009; Kandler et al., 2009; Formenti et al., 2011a, 2011b). The clay mineral fraction also provides information on climatic conditions, such as precipitation and runoff patterns over the adjacent continents (Chamley, 1989; Montero et al., 2009, 2010), as well as on the dynamics of river inputs (Pinsak and Murray, 1960). The geochemical signature of clays would complementary help characterizing the main sources and evidencing

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



specific transportation patterns or transfer processes (Haug et al., 2001; Kandler et al., 2009; Formenti et al., 2011a, b). Coupling clay mineral and grain-size studies would help investigating the sedimentary transfer regime because sediment grain-size distribution, which is primarily driven by sedimentary processes, reflects transportation conditions (Ehrmann et al., 2007; Montero et al., 2009; Sionneau et al., 2010). In this frame, combining clay mineralogy with grain-size analyses and geochemical tracers would allow retracing significant variations of detrital supply in the Sicilian-Tunisian strait and inferring any major modifications of both atmospheric and oceanic terrigenous transfers patterns since the last deglaciation.

## 2 Geographical settings

### 2.1 Oceanic circulation

The Mediterranean Sea is a concentration basin where evaporation exceeds precipitation plus freshwater discharge. The surface Atlantic water entering from Gibraltar Strait transforms into Modified Atlantic Water (MAW) as it flows eastward (Fig. 1). The eastern Mediterranean surface water displays a counterclock wise circulation (Pickard and Emery, 1982; Pinardi and Masetti, 2000). The Levantine Intermediate Water (LIW) forms in the northwest Levantine basin (Fig. 1) through convection cells at mid-depth during winter (Lascaratos et al., 1998). The LIW flows westward in sub-surface layers (~ 150 to 600–800 m) towards the northernmost Adriatic Sea and western Mediterranean (Fig. 1) via the relatively shallow Siculo-Tunisian Strait. The dense Eastern Mediterranean Deep Water (EMDW) – formed in the Adriatic Sea (Fig. 1) and occasionally in the Aegean Sea – and Western Mediterranean Deep Water (WMDW) – formed in the Gulf of Lions – fill the deep basin (< 800 m depth) (Wüst, 1961; Pickard and Emery, 1982; Malanotte-Rizzoli and Hodt, 1988; Klein et al., 1999). The Mediterranean Intermediate Water represents up to 80 % of the Mediterranean Outflow (MOW) into the Atlantic Ocean (Bryden and Stommel, 1984). This general pattern is highly dependent

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on environmental conditions including eolian activity and precipitations distribution and recent alteration of the ocean/atmosphere coupling has resulted in enhanced deep-water formation in the Aegean Sea (Malanotte-Rizzoli et al., 1999).

## 2.2 Present-day river supplies

At present day, a major part of detrital clays is supplied to the Mediterranean via rivers (Fig. 1), the most important being the Nile River (from  $120$  to  $230 \times 10^6 \text{ tyr}^{-1}$ ) discharging in the eastern Mediterranean. The Po River, discharging into the Adriatic Sea ( $17 \times 10^6 \text{ tyr}^{-1}$ ) and southeastern European rivers associated with Turkish rivers, providing about respectively  $30 \times 10^6 \text{ tyr}^{-1}$  and  $17 \times 10^6 \text{ tyr}^{-1}$  to the Aegean Sea, are some major contributors to sedimentation into the central Mediterranean (Holeman, 1968; Milliman and Syvitski, 1992; Stanley et al., 1992; Erhmann et al., 2007b; Garzanti et al., 2006; Hamann et al., 2009). The detrital supply through the Dardanelles Strait is reduced ( $0.9 \times 10^6 \text{ tyr}^{-1}$ ) because most sediment is trapped within the Black Sea and Marmara Sea (Erhmann et al., 2007a). Central Mediterranean is not affected by detrital supply from the Rhône River nor from the Ebro River which discharge into the western Mediterranean. Finally riverine contribution to the central Mediterranean from areas bordering the southern shores is negligible due to the narrow drainage basin and sparse rainfall (Martin and Milliman, 1997).

## 2.3 Eolian supply

The estimation of the present-day eolian contribution to deep-sea sedimentation varies from  $3.9 \times 10^6 \text{ tyr}^{-1}$  to  $120 \times 10^6 \text{ tyr}^{-1}$  (Bergametti et al., 1989c; Matthewson et al., 1995; Guernozi and Chester, 1996; Goudie and Middleton, 2001). This wide range of estimation reflects intermittent and seasonal variations of dust outbreaks toward Europe (D'Almedia, 1986; Guerzoni et al., 1997). Satellite imagery, back trajectories and observations indicate that fine particles originating from North Africa (Fig. 1) are transported by southerly/southwesterly winds (Scirocco, Ghibbli) toward the central

CPD

8, 2921–2968, 2012

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mediterranean (Ganor et al., 2000; Goudie and Middleton, 2001; Washington et al., 2005; Engelstaeder et al., 2006; Israelevich et al., 2012). Arid and semi-arid regions of North Africa (Fig. 1) are the main areas of production of dust all-year long (Coudé-Gaussen, 1982; Pye, 1987): Tunisia and northern Algeria, Morocco and western Sahara, the South Algeria-Mali region, the Bodélé depression and the southern Egypt-northern Sudan (Brooks and Legrand, 2000; Caquineau et al., 2002; Israelevich et al., 2002; Prospero et al., 2002; Goudie, 2003; Formenti et al., 2011a). Most of the northward aerosols transportation across the Mediterranean is linked to the seasonal displacement of cyclones over the Mediterranean (Folger, 1970; Guerzoni et al., 1997; Moulin et al., 1997; Rodriguez et al., 2001). Maximum aerosol is observed over central and eastern Mediterranean in spring (Luck and Ben Othman, 2002; O'Hara et al., 2006) and summer when anticyclonic conditions initiates drought over the area (Prospero et al., 2002; Koren et al., 2006; Roberts, 2008; Israelevich et al., 2012). Although Saharan fine particles are mainly produced during warm and hydrolysis period, it is transported from paleosols and little consolidated formations during dry intervals when vegetation is reduced and eolian erosion more efficient (Brooks and Legrand, 2000). During summer, the Saharan depression favors the transport of dust over the western basin, progressively moving toward central Mediterranean (toward Corsica and Italy) at the end of summer (Bergametti et al., 1989a; Moulin et al., 1997; Barry and Chorley, 1998).

### 3 Materials and methods

#### 3.1 Materials

The core MD04-2797CQ (36°57' N-11°40'E) was taken during the PRIVILEGE-PRIMAROSA cruise in 2004 at 771 m water-depth in the Sicilian-Tunisian Strait (Fig. 1). According to its geographical setting, the sedimentation of the core is likely influenced by both marine and eolian supplies. The scarcity of rivers along the North African

margin, and their non-permanent characters prevents the area from any major and perennial fluvial contribution (O'Hara et al., 2006). By contrast, the area is submitted to intense wind-driven supply all year long, carrying along with the dominant south-southwestern low-altitudes dusty winds (Scirocco) (Barry and Chorley, 1998) which reach the Adriatic Sea by crossing the Mediterranean (Fig. 1). According to the main wind system, the proximal source of dust for the Sicilian-Tunisian strait is likely the Tunisia-northern Algeria area (Coudé-Gaussen, 1987). The Mali-Algeria frontier or the Moroccan Atlas (Brooks and Legrand, 2000; Caquineau et al., 2002; Prospero et al., 2002; Goudie, 2003) could also contribute to sedimentation either during intense dust episode or when specific and/or seasonal atmospheric configuration affect the main wind directions (Bergametti et al., 1989a, 1989b; Ganor et al., 1991; Pye, 1992; Avila et al., 1997; Goudie and Middleton, 2001; Prospero et al., 2002; Washington et al., 2005).

### 3.2 Chronostratigraphy

The age model is based on 13 AMS  $^{14}\text{C}$  datations (Rouis-Zargouni et al., 2010, 2012). The ages were corrected using reservoir ages of 400 yr during the Holocene, the Younger Dryas and the late glacial, 560 yr during the Bølling-Allerød and 800 yr during the Heinrich event 1 and Older Dryas (H1-OD), following recommendations of Siani et al. (2001) (Table 1). The top of the core is dated at 668 ka cal. BP, suggesting that the sea-sediment interface was not preserved during coring. The Terrigenous Mass Accumulation rates (MART) were calculated as follow:  $\text{MART} (\text{g kyr}^{-1} \text{cm}^{-2}) = \text{Linear Sedimentation Rates} (\text{cm kyr}^{-1}) \cdot \text{dry density} (\text{g cm}^{-3}) \cdot (100 - (\% \text{opal} + \% \text{CaCO}_3))$ .

### 3.3 Clay minerals

The analyses were performed according to the protocol described in Bout-Roumazielles et al. (1999). All samples were first decalcified with 0.2 N hydrochloric acid. The excess acid was removed by repeated centrifugations. The clay-sized fraction

## Atmospheric supply

V. Bout-Roumazielles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



( $< 2 \mu\text{m}$ ) was isolated by settling, and oriented on glass slides (oriented mounts). Three XRD (X-ray diffraction) determinations were performed using a Bruker D4 Endeavor coupled with Lynxeye detector: (a) untreated sample; (b) glycolated sample (after saturation for 12 h in ethylene glycol); (c) sample heated at  $490^\circ\text{C}$  for two hours. Each clay mineral is then characterized by its layer plus interlayer interval as revealed by XRD analysis. Smectite is characterized by a peak at  $14 \text{ \AA}$  on the untreated sample test, which expands to  $17 \text{ \AA}$  after saturation in ethylene glycol and retracts to  $10 \text{ \AA}$  after heating. Illite presents a basal peak at  $10 \text{ \AA}$  on the three tests (natural, glycolated, and heated). The illite crystallinity, or Kübler Index is based on the expression of the width of the illite peak at  $10 \text{ \AA}$  and allows identifying the anchizone, the limit of diagenesis and the onset of the epizone. Chlorite is characterized by peaks at  $14 \text{ \AA}$ ,  $7 \text{ \AA}$ ,  $4.72 \text{ \AA}$  and  $3.53 \text{ \AA}$  on the three tests. Kaolinite is characterized by peaks at  $7 \text{ \AA}$  and  $3.57 \text{ \AA}$  on the untreated sample and after saturation in ethylene glycol. Both peaks disappear or are strongly reduced after heating. Palygorskite presents a basal peak at  $10.34 \text{ \AA}$ , accompanied by a weaker peak at  $6.44 \text{ \AA}$ , on both untreated and glycolated tests. The  $10.34 \text{ \AA}$  peak collapses at  $10 \text{ \AA}$  after heating (Brindley and Brown, 1980). The presence of palygorskite has been confirmed by MET observations of the palygorskite-rich samples. Semi-quantitative estimation of clay mineral abundances, based on the pseudo-voigt deconvolution for the doublets illite-palygorskite ( $10 \text{ \AA}$ – $10.34 \text{ \AA}$ ) and kaolinite-chlorite ( $3.57 \text{ \AA}$ – $3.53 \text{ \AA}$ ), was performed using the software MacDiff developed by Petschick (2001).

**3.4 Grain-size**

Grain-size analyses were performed on the carbonate- and opal-free fraction of the sediment using a Malvern Mastersizer 2000 laser ( $0.02$ – $2000 \mu\text{m}$ ) following protocols described in details in Montero et al. (2009). After defloculation, an aliquot of the sample was measured using 10 % ultrasonication in the Hydro S dispersion cell, once beam obscuration ranges between 12 and 15 %. The mode (i.e. most frequent grain-size in

µm), the percentage of clay (< 2µm), cohesive silt (2–10 µm), sortable-silt (10–63 µm) and sand (> 63 µm) are reported.

### 3.5 XRF-scanning

Elemental abundances were measured on U-channels using the Avaatech core scanner from the EPOC laboratory at the University of Bordeaux. The sediment was protected with Ultralene® X-ray transmission foil in order to avoid contamination. The data were acquired at a 30 s count time, using 10 kV voltage and 400 mA intensity. The results are expressed in counts per second, with a 2 % precision according to standard samples. Trace elements are normalized to Al (Tribovillard et al., 2008), in order to avoid dilution effect by carbonates.

## 4 Results

### 4.1 Clay mineralogy

The clay mineral fraction is composed of kaolinite (45 %), smectite (25 %) and illite (15 %). Chlorite and palygorskite are secondary components with 10 and 6 % of the clay content respectively (Table S1). The composition of the clay mineral assemblage displays some variations over the studied interval (Fig. 2). The kaolinite content remains slightly below 45 % between 18.5 and 11.7 ka. After what it increases and represents 50 % between 10.5 and 8.5 ka. The percentage of kaolinite decreases at 8.5 ka and remains low between 8 and 5.8 ka. It then increases progressively between 5.8 and 2 ka and reached its maximum (> 50 %) around 2 ka. The smectite remains stable between 18.5 and 8.5 ka excepted two maxima around 12 ka and 11 ka (Fig. 2). The content in smectite increases up to 38 % of the clay mineral fraction between 8.5 and 5.8 ka as the kaolinite content is minimum. This smectite-rich interval is interrupted by a slightly depleted level around 7.2 ka (Fig. 2). The smectite content decreases progressively since 5.8 ka down to 15 % at the top of the core. The illite content is maximum

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

between 18.5 and 11.7 ka, then it drops down and remains low ( $< 15\%$ ) between 11.7 and 5 ka. Then illite increases slightly and remains stable around 15% over the more recent part of the record (Fig. 2). The Kübler Index is low ( $< 0.4^\circ 2\theta$ ) indicating high structural order when the content in illite is maximum. By contrast, the Kübler Index is higher ( $< 0.5^\circ 2\theta$ ) – low ordering- between 11.7 and 5 ka when the percentage of illite is reduced. The illite to kaolinite ratio (I/K) is characterized by a drastic change at 11.7 ka from values around 0.35 to values as low as 0.25. The chlorite content does not correlate with other clay species and shows slight variations, being higher than average between 18.5 and 15.5 ka. The palygorskite is less abundant between 8.5 and 5.8 ka, while smectite is maximum (Fig. 2).

## 4.2 Grain-size

The grain-size varies between 4  $\mu\text{m}$  and 10  $\mu\text{m}$ , indicating the detrital fraction is mainly composed of cohesive silt-size particles. Cohesive particles indeed represent 50 to 70% of the total terrigenous fraction, being slightly less abundant between 18.5 and 11.7 ka and between 7.5 and 5.5 ka (Fig. 3). By contrast, the clay-size fraction is generally low (around 11%), excepted between 8.5 and 5.5 ka when it represents up to 25% of the detrital material. The sortable silt (10  $\mu\text{m}$  to 63  $\mu\text{m}$ ) composes 25% of the detrital fraction, being more abundant between 13 and 12 ka where it represents 45%. The sand-size fraction is rare, but shows two maxima between 14 and 12 ka and between 8.5 and 7.5 ka (Fig. 3).

## 4.3 Elemental geochemistry

The Al content displays a decreasing trend from 18.5 to 14 ka (Fig. 4). The Ca to Fe ratio is used to distinguish areas dominated by carbonates (Sahara) from areas where hydrolysis is intense (alterites). The Ca/Fe ratio is characterized by large variations, with maximum values between 15 and 13 ka, and between 5 and 3 ka, while the ratio is minimum before 15 ka and between 2 and 1 ka. Some intervals are characterized

by higher than average K/Al ratio, between 14 and 11.7, between 5.8 and 5.5 ka and between 2.5 and 1 ka. The two latter intervals are also characterized by the highest values of the Ti/Al ration (Fig. 4). The Zr/Al ratio displays a peculiar behavior, peaking at 13.4 ka, 7 ka, 6.4 ka, 1.3 ka and being high between 12.8 and 12.3 ka.

## 5 Discussion

### 5.1 Mineralogical and geochemical characterization of particle provenance and transport patterns

Illite and chlorite are abundant in sediments from the Ionian Sea (Fig. 1), where they mainly derive from the southern European rivers flowing into the Adriatic (Po River) and Aegean Seas (southeastern European Rivers) (Venkatarathnam and Ryan, 1971; Dominik and Stoffers, 1978; Chamley, 1989; Alonso and Maldonado, 1990; Tomadin, 2000). Smectite predominates in the eastern basin, being abundant in the Marmara Sea (Ergin et al., 2012; Armynot et al., 2012), in the Levantine basin (Hamann et al., 2009) and in the southeastern Aegean Sea (Erhmann et al., 2007a). Smectite characterizes the Nile River suspended loads, deriving from Cenozoic volcanic provinces of the Ethiopian Highlands (Ukstins et al., 2002; Hamann et al., 2009), and is further redistributed as suspended clay particles within the Mediterranean surface water (Venkatarathnam and Ryan, 1971; Foucault and Mélières, 2000; Hamann et al., 2009). Palygorskite has been shown to be abundant in tunisian dunes and is used as a main eolian tracer, because it may be transported over long-range distance, crossing the Mediterranean through meridian transfers (Coudé-Gaussen et al., 1982; Robert et al., 1984; Molinaroli, 1996). Illite is also a dominant component of eolian dust originating from Sahara when associated with palygorskite (Coudé-Gaussen and Blanc, 1985; Guerzoni et al., 1999; Foucault and Mélières, 2000; Goudie and Middleton, 2001), whereas illite associated with abundant kaolinite originate from southeastern areas of North Africa (Chester et al., 1977; Stanley and Wingerath, 1996; Foucault

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Mélières, 2000; Goudie and Middleton, 2001; O'Hara et al., 2006; Hamann et al., 2009). The illite to kaolinite ratio (I/K) is a useful tracer of Saharan versus Sahelian dust provenance (Schütz and Seibert, 1987; Caquineau et al., 1998, 2002; Formenti et al., 2011a), which is thought to be conservative after long-range transport (Chester et al., 1972; Caquineau et al., 1998, 2002; Kandler et al., 2009). Highest ratio characterizes the northwestern areas (Fig. 1), whereas low ratio retraces the Sahelian belt: the lowest ratios are observed in dust collected in Sahel (0.1 to 0.3), intermediate values characterize south and central Sahara (0.3 to 0.7), moderate to high values are observed in Tunisia and northern Algeria (1 to 2) whereas highest ratios (1.3 to 2.6 up to 6) occur in northwestern Sahara (Fig. 1) and in Morocco (Chester et al., 1972; Glaccum and Prospero, 1980; Paquet et al., 1984; Coudé-Gaussen, 1991; Kiefert et al., 1996; Avila et al., 1997; Caquineau et al., 1998, 2002; Kandler et al., 2009). The I/K ratio also displays a longitudinal gradient (Fig. 1) with ratio decreasing from northwestern Africa (I/K = 2 to 1.1) toward northeastern Africa (I/K = 0.2 to 0.7) (Gomes et al., 1990; Ganor and Forer, 1996; Caquineau et al., 1998).

Geochemical data are used to pinpoint the origin of the terrigenous fraction and to retrace transportation processes. North and central Sahara are characterized by carbonate-rich soils, whereas the Sahelian area submitted to intense hydrolysis is depleted in Ca and enriched in Fe (Kandler et al., 2009; Formenti et al., 2011a, b). As a result, the Ca/Fe ratio is used to distinguish the Saharan and Sahelian provenance, complementary with the I/K ratio: Tunisia-northern Algeria source is characterized by high carbonate (up to 50 %) and low Fe contents (Chester et al., 1984; Bergametti et al., 1989), Moroccan Atlas has a high carbonate content (up to 70 %) and a rather high Fe content (Bergametti et al., 1989; Avila et al., 1997; Kandler et al., 2009), Libya displays intermediate Fe and carbonate contents (Guieu et al., 2002; O'Hara et al., 2006) whereas southern Algeria shows intermediate Fe content associated with low carbonate content (< 25 %) (Paquet et al., 1984; Bergametti et al., 1989; Alastuey et al., 2005).

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The grain-size of dust displays strong seasonal variations (Guerzoni et al., 1997), with a fine population (2 to 10  $\mu\text{m}$ , mode around 2.7  $\mu\text{m}$ ) characterizing long-distance and high latitude transport and a coarser population (5 to 50  $\mu\text{m}$ , mode around 20  $\mu\text{m}$ ) typical of dust storms carried over few km, being restricted to adjacent continental and marine areas (Prospero, 1981; Torres-Padron et al., 2002). Comparison between Ti/Al and Zr/Al ratios and grain-size distribution are thus used to identify specific eolian vs. riverine transport processes, associated with the development of sand dunes during the Holocene, or with remote fine-grained eolian outbreaks or coarse-grained fluvial supply (Haug et al., 2001; Martinez-Ruiz et al., 2003; Erhmann et al., 2007b; Hamann et al., 2009; Montero et al., 2009; Skonieczny et al., 2011).

## 5.2 Bølling-Allerød

The Bølling-Allerød (B-A) is marked by enhanced terrigenous supply starting at 14.6 ka, associated with increased contributions of both illite and palygorskite while kaolinite and chlorite decrease (Fig. 2). The relationship between illite content and sortable silt-size fraction (14  $\mu\text{m}$ ) indicate that illite is mostly transported through eolian processes. The concomitant increase of the Ca/Fe and I/K ratio suggests that the increased eolian supply is associated with a modification of the main detrital source (Fig. 5a). Indeed, increased proportion of Ca vs. Fe during the B-A may basically be interpreted as reflecting higher productivity but this hypothesis is not supported by the dinoflagellate record from core MD04-2797 (Rouis-Zargouni et al., 2010). The increased proportion of illite may indicate a northward migration of the main clay provenance over Libya, as illite has been shown to be more abundant in dust collected over northern Libya (O'Hara et al., 2006). Nevertheless the I/K ratio range – varying between 0.35 and 0.4 – during the B-A is rather low compared with observed values for Libya and does not further support this hypothesis (Fig. 5a). The observed sedimentological characteristics tend to indicate Tunisia-Algeria as the main dust provenance (Formenti et al., 2011a). According to previously published data, the mineralogical composition is indeed consistent with a source from northern or central Algeria (Fig. 7a), but the low content in

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



palygorskite ruled out any major contribution from tunisian loess (Bout-Roumazeilles et al., 2007). The development of humid conditions during the B-A is evidenced by an increased contribution of river vs. eolian supply in the Aegean Sea (core SL128, Fig. 1) (Hamann et al., 2008), by pollen association from the Alboran (ODP976, Fig. 1) and Adriatic Sea (Combourieu-Nebout et al., 1998, 2002; Fletcher and Sanchez-Goñi, 2008) and by a speleothem record from the eastern Mediterranean (Bar-Matthew et al., 2003), contrasting with the clay mineral record from the Sicilian-Tunisian Strait. The Sea Surface Temperatures (SST, Fig. 3) – as reconstructed from planktonic *foraminifera* at site MD04-2797 – are high during the Bølling-Allerød (Ellassami et al., 2007), consistently with general warm oceanic conditions, relatively high humidity and temperatures compared with present-day in Sicily and central Mediterranean (Ramrath et al., 2000; Allen et al., 2002; Zielhofer et al., 2008; Incarbona et al., 2010). The dominance of semi-desert plants at site MD04-2797 (Desprat et al., 2012) confirms the persistence of dry continental conditions over Africa during the B-A. These results tend to confirm contrasting climatic evolution (e.g. Roberts et al., 2008) with humid conditions on the northeastern and eastern Mediterranean whereas aridity still persisted on the southern Mediterranean.

### 5.3 Younger Dryas

A major peak in the Zr/Al ratio is observed between around 12.8 ka, associated with an increase in the abundance of both sortable-silt and sand-size particles (Fig. 5b). The clay association does not exhibit major modification but smectite is more abundant at the expense of kaolinite, displaying a peak at 12 ka (Fig. 2). These characteristics indicate the supply of coarser particles that may either be associated to a volcanic supply, or to river supply or to an intense dust episode (Martinez-Ruiz et al., 2003; Erhmann et al., 2007b; Hamann et al., 2009; Skonieczny et al., 2011). Smectite peaking at 12 ka supports the hypothesis of a tephra (Fig. 5d). Moreover, when compared with the chronology of tephras (e.g. Zanchetta et al., 2011), the increase of Zr may be associated with the Agrano Pomici Principali (12.8–12.2 ka) tephra (DiVito et al.,

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





1999; Siani et al., 2004). Nevertheless, optical observations of the sediment do not give any support to this hypothesis. The hypothesis of enhanced river supply is also unlikely considering the general arid climatic conditions that prevailed during the Younger Dryas (YD) (Lézine et al., 1995; Gasse et al., 2000). This specific level could thus alternatively reflect intense Sahara eolian activity that has previously been reported in sediment cores from the northeastern Tropical Atlantic (De Menocal et al., 2000; Cole et al., 2009; Skonieczny et al., 2012).

Cold intervals are generally associated with intense dust deposition/eolian activity resulting from reduced vegetation promoting the erosion of soils, low moisture availability and enhanced atmospheric circulation. The North Atlantic Oscillation (NAO), which is responsible for some part of the Mediterranean climate variability since its impact on the wind activity and precipitation balance (Pittalwala and Hameed, 1991; Moulin et al., 1997; Sanchez-Goñi et al., 2002), has been shown to be positive during the YD (Kim et al., 2007). A positive phase of the NAO is thought to be responsible for the development of dry conditions over southern Europe and North Africa and for promoting northward intrusions of Saharan air masses toward the western Mediterranean (ODP976, Fig. 1) during cold climatic events of the last climatic cycle (Combourieu-Nebout et al., 2002; Bout-Roumazielles et al., 2007). These cold events were characterized by increased palygorskite supply associated with the dominance of arid steppe (*Artemisia*) and the occurrence of endemic pollen (*Argania*) pinpointing a dust source from western Morocco being transported along the North African coast (Bout-Roumazielles et al., 2007). The absence of specific enrichment in palygorskite during YD in the Sicilian-Tunisian Strait indicates that the main long-range transportation pattern of fine Saharan dust is not responsible for the peculiar characteristic of sediments deposited in the Sicilian-Tunisian Strait. This is in agreement with grain-size and geochemical data, which rather point to medium to short-range transportation of coarse particles from proximal continental areas (Pye and Tsoar, 1987; McTainsh et al., 1997; O'Hara et al., 2006). The occurrence of such a coarse-grain proximal supply during the YD, consistent with the presence of loess deposits and coastal dunes in Tunisia (Coudé-Gaussen et al., 1987;

**Atmospheric supply**

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Grousset et al., 1992; Crouvi et al., 2010) and further supported by model suggesting that Libyan desert and coastal areas are major coarse-grain dust sources for central and eastern Mediterranean (Callot et al., 2000), may thus reflect local eolian activity forced by ocean-atmosphere linkage at global scale (Fig. 7b).

## 5.4 Onset of the Holocene

The end of the Younger Dryas (11.7 ka) is marked by a drastic change in the clay assemblage (Figs. 5a, 6a and b) and a decrease of the sortable silt proportion. The kaolinite increases while illite diminishes, resulting in a progressive decrease of the I/K ratio. This shift, associated to a modification of the illite structural composition, evidences a major change of provenance of the clay fraction, interpreted as an increase contribution of the Sahelian area (Fig. 6a). This observation is supported by the synchronous changes in geochemical signature (Fig. 6c) confirming enhanced contribution of strongly hydrolyzed domains. Low I/K indicates enhanced contribution from the southern Algeria-Mali border region (Paquet et al., 1984; Bergametti et al., 1989b; Alastuey et al., 2005; Bout-Roumzeilles et al., 2007), but the clay association suggests a provenance from southern Algeria rather than from northern Mali, because kaolinite is more abundant southward (e.g. Bout-Roumzeilles et al., 2007), becoming dominant in southern Algeria whereas the Mali border is characterized by smectite-rich assemblages (Fig. 7c). The abundance of semi-desert pollen taxa exhibits a sharp decrease at 12.25 ka (Desprat et al., 2012), before the mineralogical transition, suggesting that the modification of vegetation, i.e. a major change in steppe composition and a development of scarce deciduous oak woodlands, might prevent soils from weathering, displacing southward the areas of eolian erosion. The end of the YD is marked by a similar modification of the clay mineral fraction in a core from the northeastern Tropical Atlantic off Senegal (Skonieczny et al., 2012) and interpreted as recording a southward shift of the Inter Tropical Convergence Zone (ITCZ), which strongly affect the transport of Saharan dust toward the Atlantic Ocean. Moreover, the onset of the Holocene is also marked by a decrease of the I/K ratio off Portugal (Stumpf et al.,

## Atmospheric supply

V. Bout-Roumzeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2011). The synchronicity of the clay modification at these sites, submitted to distinct wind regimes, signify that the end of the YD is marked by a large-scale atmospheric re-organization over the North African continent, with the development of dry conditions during the early Holocene, affecting both the high-latitude long-range transportation and the regional low-latitude wind regime over the Mediterranean.

## 5.5 Sapropel S1

The time interval between 8.6 and 5.5 ka is marked by the highest terrigenous flux and smectite content, with a significant grain-size fining (Figs. 2 and 5d). The associated increase in particulate organic carbon (POC) during the whole interval suggests presapropelic conditions (Essallami et al., 2007). Both timing and peculiar characteristics suggest this interval to be related to the most recent organic-rich sapropel S1 (Kidd et al., 1978) deposited during the mid-Holocene in the eastern Mediterranean (Emeis et al., 1996; Emeis et al., 2000; De Lange et al., 2008). During the sapropel, increased precipitation on the surrounding continents provided freshwater discharge and promoted strong stratification of the water column and enhanced nutrient supply, resulting in increased productivity and improved preservation due to oxygen depletion of the water column (Rossignol-Strick et al., 1982; Rohling, 1994; Abu-Zied et al., 2008). Increased precipitations are thought to be linked to either orbitally-forced enhanced monsoon activity (Rossignol-Strick et al., 1982; Hilgen, 1991; Lourens et al., 1996; Emeis et al., 2000) or to increased rainfall over the Mediterranean region (Kallel et al., 1997; Magny et al., 2002). Indeed, although some data appear to be consistent over the Mediterranean basin (Sadori et al., 2011), rainfall records display seasonal modulation (Sadori et al., 2004, 2008; Frisia et al., 2006; Magny et al., 2007, 2012; Kotthoff et al., 2008; Jalut et al., 2009; Colonese et al., 2010; Giraudi et al., 2011; Peyron et al., 2011). Speleothems indicate increased moisture availability in both western (Corchia Cave, Zanchetta et al., 2007), and eastern Mediterranean (Soreq Cave, Bar-Matthews et al., 2000) consistent with lakes records from East Anatolia (Stevens et al., 2001; e.g. Roberts et al., 2008) but also evidence contrasting moisture originating either from

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Atlantic (Bard et al., 2002) or from the Mediterranean (Arz et al., 2003). Moreover rainfall patterns also display slightly different chronology in Italian lakes (Roberts et al., 2008) and stalagmite records (Frisia et al., 2006; Zanchetta et al., 2007). Consequently, most studies focused on the precipitation distribution and moisture availability that prevailed on the continent at the time of S1.

But investigations of the marine propagation of the sapropelic layer from eastern toward western Mediterranean evidenced the absence of true sapropels and/or synchronous events (Mercone et al., 2000) in the western basin, suggesting lower export of production fluxes into the basin (Martinez-Ruiz et al., 2003) or different hydrographic conditions (Weldeab et al., 2003). Smectite provenance during the sapropel in the Sicilian-Tunisian Strait would provide additional information on the modifications of oceanic environmental conditions. Indeed, central Mediterranean is depleted in smectite (Bout-Roumazielles et al., 2007), which is rare in southern Algeria and absent from northern and central Algeria. By contrast, the southern Algeria-Mali border (Fig. 1) is locally characterized by a high content in smectite (Bout-Roumazielles et al., 2007). But, at present-day, this area is submitted to trade-winds, promoting a dominant low-altitude westward transportation and ruling out any significant contribution of this area to sedimentation in the central Mediterranean. A volcanic origin of smectite from Sicilia is not supported by optical observations or by the chronology of the main tephra (Siani et al., 2004). The enhanced supply of smectite may originate from tunisian loess (Fig. 1) (e.g. Bout-Roumazielles et al., 2007), but the low content of palygorskite during the smectite-rich interval does not further support this interpretation. Moreover, the enhanced moisture availability over that period, associated with pollen association – i.e. open oak forest with heath underbrush or maquis and Asteraceae-Poaceae-Cyperaceae steppe (Desprat et al., 2012) preventing soils from erosion – and the geochemical signal give little support to significant eolian activity. These observations highlight that smectite may alternatively be supplied toward the Mediterranean by the Nile River and transported as suspended particles within the Mediterranean surface water (Ventakatarathnam and Ryan, 1971; Stanley and Wingerath, 1996; Foucault

**Atmospheric supply**

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Mélières, 2000; Hamann et al., 2009). This smectite-rich interval is indeed characterized by an increase abundance of palladium content, suggesting the presence of Pd-bearing smectite, which are generally associated with mafic and ultramafic rocks (Sawlowicz, 1993) and may indicate a specific contribution from the volcanic and ultramafic materials outcropping in Ethiopia (Fig. 7d). This interpretation is consistent with an increased contribution of the Blue Nile draining the Cenozoic volcanic provinces of Ethiopian Highlands (Ukstins et al., 2002), during the African Humid period (Revel et al., 2010). In the Levantine basin (SL112, Fig. 1), the terrigenous fraction, deriving from the Nile River (Krom et al., 2002; Schilman et al., 2001; Hamann et al., 2008), reveals an increase in cohesive fine-grained particles during the Sapropel S1 (Hamann et al., 2010), interpreted as enhanced pluvial period in Egypt (Revel et al., 2010) during the African Humid Period, AHP (Cole et al., 2009; De Menocal et al., 2000).

Considering uncertainties on the age models, the enhanced contribution of fine particles is synchronous in the Levantine Sea and in the Sicilian-Tunisian Strait, supporting a provenance of the fine smectite-rich particles from the Eastern Mediterranean basin. The abundance of fine particles may thus result from slightly slower oceanic circulation during the Sapropel S1 promoting the deposition of the finest fraction through settling processes (Cram and O'Sullivan, 1999). The circulation pattern is consistent with this hypothesis since the Levantine Intermediate Water (200–600 m water-depth) overflows toward the western Mediterranean basin across the Sicilian-Tunisian Strait. Moreover, cold-water corals and benthic *foraminifera* assemblages evidenced the impact of sapropelic conditions in shallow water (600 m water-depth) from the Ionian Sea (Fink et al., 2012), giving additional support to a westward propagation of sapropelic conditions through intermediate water-masses (Fig. 7d). The occurrence of a smectite-rich interval in the Alboran Sea (Bout-Roumzeilles et al., 2007), and on the Atlantic side of the Gibraltar Strait during the AHP, suggest a potential transport of smectite through the Mediterranean Overflow Water (Stumpf et al., 2010).

**Atmospheric supply**

V. Bout-Roumzeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 5.6 Late Holocene

The upper part of the record is marked by an increase of the Ti/Al and Zr/Al ratios (Fig. 5c), associated with a high supply in quartz, both suggesting enhanced wind activity. The dominance of silt-size particles and the stable MART (Fig. 5c) suggest long-range transportation rather than enhanced eolian supply. The increases in kaolinite and palygorskite abundances (Fig. 5c) as well as the low Ca/Fe and I/K ratios indicate a provenance from the Sahelian area and/or the southwest Morocco (Fig. 7e). This event is synchronous with the occurrence of cool/arid events in the Mediterranean (Rapid Climate Coolings-RCC; e.g. ~ 3.5–2.5 kyr BP), suggesting relationships with high-latitude climatic phenomena and altered runoff regime (Mayewski et al., 2004). Previous studies evidenced a climate change around 4.5 ka cal. BP, with distinct regional late Holocene climatic evolution in the Mediterranean (Roberts et al., 2008; Sadori et al., 2008; Peyron et al., 2011; Sadori et al., 2011; Vanni re et al., 2011; Magny et al., 2012). Southern sites experienced a drastic decrease in summer precipitations (Magny et al., 2012) while northern sites were submitted to increased precipitation. According to present-day atmospheric pattern, the late Holocene mineralogical and geochemical records are consistent with a mean southward position of the ITCZ which may also explain the low-latitude aridity associated to the RCC (Haug et al., 2001; Hodell et al., 2001). A southward position of the ITCZ would indeed explain the geographically contrasted Mediterranean climatic evolution by promoting the aridification of North Africa while increased summer precipitation occurred in northern Mediterranean (Vanni re et al., 2011) as well as enhanced high-altitude export of dust (Skonieczny et al., 2011).

CPD

8, 2921–2968, 2012

### Atmospheric supply

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 6 Conclusions

The clay mineral, grain-size and geochemical studies of sediment deposited on the Sicilian-Tunisian Strait help retracing atmospheric vs. riverine terrigenous supplies variability since the last glacial in central Mediterranean. Although the eolian supply is dominant at the studied site – excepted during the sapropel S1 – both flux and the main provenance of particles display strong variations, related to aridity/moisture balance and vegetation cover, driven by large-scale atmospheric reorganization.

The Bølling-Allerød is marked by increased terrigenous flux while both illite and palygorskite became main components of the clay mineral fraction. The concomitant dominance of silt-size particles indicates an eolian origin for this enhanced detrital supply. The geochemical and mineralogical data indicate a dominant Saharan contribution, pinpointing central and northern Algeria as a main provenance. This increased eolian contribution from Sahara highlights contrasting regional climatic evolution, with the development of moist conditions over the North and eastern Mediterranean during the Bølling-Allerød, while glacial aridity persisted on the southern Mediterranean.

A short-term coarse-grained and Zr-rich interval characterized the Younger Dryas, as arid and cold climatic conditions promoted soil erosion. Similar events characterize the western Mediterranean and the northeastern Tropical Atlantic Ocean, although clay mineralogy clearly indicates different provenances. These results suggest local increase of wind activity driven by wide ocean-atmosphere interactions.

The onset of the Holocene is marked by a major change of clay mineralogy and crystallinity corresponding to a southward migration of the main clay provenance toward the Sahelian belt. This change is associated with a progressive development of the Mediterranean vegetation and the southward shift of the ITCZ. Similar signals are recorded in the northeastern tropical Atlantic Ocean and off Portugal, suggesting a large-scale atmospheric reorganization.

High terrigenous flux associated with the dominance of very fine-grained Pd-rich smectite characterized the early to mid-Holocene in central Mediterranean, while the

CPD

8, 2921–2968, 2012

### Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



organic content indicates pre-sapropelic conditions. The sedimentological characteristics rule out any eolian supply but rather suggest enhanced riverine contribution from remote area during the sapropel S1. The peculiar signature of this event in the central Mediterranean and in the Levantine basin suggests the propagation through intermediate water-masses of fine-grained smectite originating from the Nile River during the sapropel S1, resulting from enhanced precipitations on northeast Africa, coincident with the late phase of the AHP.

Finally, an increase supply of dust originating from northern Sahel or southern Morocco characterizes the last 3 kyr, indicating intense eolian activity linked to modification of the rainfall regime.

**Supplementary material related to this article is available online at:**  
**<http://www.clim-past-discuss.net/8/2921/2012/cpd-8-2921-2012-supplement.pdf>.**

*Acknowledgements.* This study was funded by the ANR LAMA. Philippe Martinez (EPOC – University of Bordeaux) is thanked for producing XRF analysis of the core. Léa-Marie Bernard Emaile, Deny Malengros and Romain Abraham are thanked for technical assistance for XRD analysis and laser grain-size measurements.



The publication of this article is financed by CNRS-INSU.

CPD

8, 2921–2968, 2012

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## References

- Abu-Zied, R. H., Rohling, E. J., Jorissen, F. J., Fontanier, C., Casford, J. S. L., and Cooke, S.: Benthic *foraminiferal* response to changes in bottom-water oxygenation and organic carbon flux in the eastern Mediterranean during LGM to Recent times, *Mar. Micropaleontol.*, 67, 46–68, 2008.
- Alastuey, A., Querol, X., Castillo, S., Escudero, M., Avila, A., Cuevas, E., Torres, C., Romero, P.-M., Exposito, F., Garcia, O., Pedro Diaz, J., Dingenen, R. V., and Putaud, J. P.: Characterization of TSP and PM<sub>2.5</sub> at Izana and Sta. Cruz de Tenerife (Canary Islands, Spain) during a Saharan Dust Episode (July 2002), *Atmos. Environ.*, 39, 4715–4728, 2005.
- Allen, J. R. M., Watts, W. A., McGee, E., and Huntley, B.: Holocene environmental variability – The record from Lago Grande di Monticchio, Italy, *Quaternary. Int.*, 88, 69–80, 2002.
- Alonso, B. and Maldonado, A.: Late Quaternary sedimentation patterns of the Ebro turbidite systems (northwestern Mediterranean): Two styles of deep-sea deposition, *Mar. Geol.*, 95, 353–377, 1990.
- Ariztegui, D., Asioli, A., Lowe, J. J., Trincardi, F., Vigliotti, L., Tamburini, F., Chondrogianni, C., Accorsi, C. A., Bandini Mazzanti, M., Mercuri, A. M., Van Der Kaars, S., Mckenzie, J. A., and Oldfield, F.: Palaeoclimate and the formation of sapropel S1: inferences from Late Quaternary lacustrine and marine sequences in the central Mediterranean region, *Paleogeogr. Palaeocl.*, 158, 215–240, 2000.
- Armynot du Châtelet, E., Bout-Roumazeilles, V., Coccioni, R., Frontalini, F., Guillot, F., Kaminiski, M. A., Recourt, P., Riboulleau, A., Trentesaux, A., Tribouvillard, N., and Ventalon, S.: Environmental control on shell structure and composition of agglutinated *foraminifera* in the Marmara Sea, *Mar. Geol.*, submitted, 2012.
- Arz, H. W., Lamy, F., Pätzold, J., Müller, P. J., and Prins, M.: Mediterranean moisture source for an early-Holocene humid period in the Northern Red Sea, *Science*, 300, 118–121, 2003.
- Avila, A., Queralt-Mitjans, I., and Alarcón, M.: Mineralogical composition of African dust delivered by red rains over northeastern Spain, *J. Geophys. Res.*, 102, 21977–21996, 1997.
- Bar-Matthews, M., Ayalon, A., and Kaufman, A.: Timing and hydrological conditions of Sapropel events in the eastern Mediterranean, as evident from speleothems, Soreq cave, Israel, *Chem. Geol.*, 169, 145–156, 2000.
- Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., and Hawkesworth, C.J.: Sea-land oxygen isotopic relationship from planktonic *foraminifera* and speleothems in the East-

CPD

8, 2921–2968, 2012

## Atmospheric supply

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ern Mediterranean region and their implication for paleorainfall during interglacial intervals, *Geochim. Cosmochim. Ac.*, 67, 3181–3199, 2003.

Bard, E., Delaygue, G., Rostek, F., Antonioli, F., Silenzi, S., and Schrag, D.: Hydrological conditions in the western Mediterranean basin during the deposition of Sapropel 6 (ca. 175 kyr), *Earth Planet. Sc. Lett.*, 202, 481–494, 2002.

Barry, R. G. and Chorley, R. J.: *Atmosphere, Weather and Climate*, 7th Edn., Routledge, London, 409 pp., 1998.

Bergametti, G., Dutot, A. L., Buat-Menard, P., Losno, R., and Rémondaki, E.: Seasonal variability of the elemental composition of atmospheric aerosol particles over the northwestern Mediterranean, *Tellus*, 41, 353–361, 1989a.

Bergametti, G., Gomes, L., Coudé-Gaussen, G., Rognon, P., and Le Coustumer, M. N.: African dust observed over Canary Islands: Source-regions identification and transport pattern for some summer situations, *J. Geophys. Res.*, 94, 14855–14864, 1989b.

Bergametti, G., Gomes, L., Remoudaki, E., Desbois, M., Martin, D., and Buat-Menard, P.: Present transport and deposition patterns of African dusts to the North-Western Mediterranean, in: *Paleoclimatology and Paleometeorology: Modern and past patterns of global atmospheric transport*, edited by: Leinen, M. and Sarnthein, M., Dordrecht, Kluwer Academic, 282, 227–251, 1989c.

Boulay, S., Colin, C., Trentesaux, A., Frank, N., and Liu, Z.: Sediment sources and East Asian monsoon intensity over the last 450 ky. Mineralogical and geochemical investigations on South China Sea sediments, *Paleogeogr. Palaeocl.*, 228, 260–277, 2005.

Bout-Roumazeilles, V., Cortijo, E., Labeyrie, L., and Debrabant, P.: Clay mineral evidence of nepheloid layer contribution to the Heinrich layers in the Northwest Atlantic, *Paleogeogr. Palaeocl.*, 146, 211–228, 1999.

Bout-Roumazeilles, V., Nebout, N. C., Peyron, O., Cortijo, E., Landais, A., and Masson-Delmotte, V.: Connection between South Mediterranean climate and North African atmospheric circulation during the last 50,000 yr BP North Atlantic cold events, *Quaternary Sci. Rev.*, 26, 3197–3215, 2007.

Breed, C. S. and Breed, W. J.: Dunes and other windforms of Central Australia, and a comparison with linear dunes on the Moenkopi Plateau, Arizona, Apollo-Souyez Test Project Summary Science Report, v.2, *Earth Observation and Photography*, edited by: El-Baz, F. and Warner, D. M., National Aeronautics and Space Administration Special Publication, 412, 319–358, 1979.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Brindley, G. W. and Brown, G.: Crystal Structures of Clay Minerals and their X-ray identification, Mineralogical Society, London, 1980.
- Brooks, N. and Legrand, M.: Dust variability over northern Africa and rainfall in the Sahel, in: Linking Climate Change to Landsurface Change, edited by: McLaren, S. J. and Kniveton, D., Kluwer Academic, Dordrecht, 1–25, 2000.
- 5 Bryden, H. L. and Stommel, H. M.: Limiting processes that determine basic features of the circulation in the Mediterranean Sea, *Oceanol. Acta*, 7, 289–296, 1984.
- Callot, Y., Marticorena, B., and Bergametti, G.: Geomorphologic approach for modelling the surface features of arid environments in a model of dust emissions: application to the Sahara desert, *Geodin. Acta*, 13, 245–270, 2000.
- 10 Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M.-C., and Chatenet, B.: Saharan dust: clay ratio as a relevant tracer to assess the origin of soil-derived aerosols, *Geophys. Res. Lett.*, 25, 983–986, 1998.
- Caquineau, S., Gaudichet, A., Gomes, L., and Legrand, M.: Mineralogy of Saharan dust transported over northwestern tropical Atlantic Ocean in relation to source regions, *J. Geophys. Res.*, 107, 4251, doi:10.1029/2000JD000247, 2002.
- 15 Chamley, H.: Sédimentation argileuse en mer Ionienne au Plio-Pléistocène d'après l'étude des forages 125 DSDP, *B. Soc. Geol. Fr.*, 7, 1131–1143, 1975.
- Chamley, H.: Clay sedimentology, Berlin, Spinger-Verlag, 623 pp., 1989.
- 20 Cheddadi, R., Rossignol-Strick, M., and Fontugne, M.: Eastern Mediterranean paleoclimates from 26 to 5 kaBP by pollen and isotopic analysis of a core in the anoxic Bannock Basin, *Mar. Geol.*, 100, 53–66, 1991.
- Chester, R., Elderfield, H., Griffin, J. J., Johnson, L. R., and Padgham, R. C.: Eolian dust along the eastern margins of the Atlantic Ocean, *Mar. Geol.*, 13, 91–105, 1972.
- 25 Chester, R., Baxter, G., Behairy, A. K. A., Connor, K., Cross, D., Elderfield, H., and Padgham, R. C.: Soil-sized eolian dusts from the lower troposphere of the eastern Mediterranean Sea, *Mar. Geol.*, 24, 201–217, 1977.
- Chester, R., Sharples, E. J., Sanders, G. S., and Saydam, A. C.: Saharan dust incursion over the Tyrrhenian Sea, *Atmos. Environ.*, 18, 929–935, 1984.
- 30 Cole, J. M., Goldstein, S. L., Demenocal, P. B., Hemming, S. R., and Grousset, F. E.: Contrasting compositions of Saharan dust in the eastern Atlantic Ocean during the last deglaciation and African Humid Period, *Earth Planet. Sc. Lett.*, 278, 257–266, 2009.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Colin, C., Siani, G., Sicre, M.-A., and Liu, Z.: Impact of the east Asian monsoon rainfall changes on the erosion of the Mekong River basin over the past 25,000 yr, *Mar. Geol.*, 271, 84–92, 2010.
- 5 Colonese, A. C., Zanchetta, G., Dotsika, E., Drysdale, R. N., Fallick, A. E., Grifoni Cremonesi, R., and Manganelli, G.: Early-Middle Holocene land snail shell stable isotope record from Grotta di Latronico 3 (Southern Italy), *J. Quaternary Sci.*, 25, 1347–1359, 2010.
- Combourieu-Nebout, N., Paterne, M., Turon, J. L., and Siani, G.: A high-resolution record of the last deglaciation on the central Mediterranean Sea: Palaeovegetation and palaeohydrological evolution, *Quaternary Sci. Rev.*, 17, 303–317, 1998.
- 10 Combourieu Nebout, N., Turon, J. L., Zahn, R., Capotondi, L., Londeix, L., and Pahnke, K.: Enhanced aridity and atmospheric high-pressure stability over the western Mediterranean during the North Atlantic cold events of the past 50 k.y, *Geology*, 30, 863–866, 2002.
- Coudé-Gaussen, G.: Les poussières éoliennes sahariennes. Mise au point, *Revue de géomorphologie dynamique*, 31, 49–69, 1982.
- 15 Coudé-Gaussen, G.: Saharan dust, *J. Geophys. Res.*, 92, 9753–9771, 1987.
- Coudé-Gaussen, G.: Les poussières sahariennes, John Libbey Euro Text, 390 pp., 1991.
- Coudé-Gaussen, G. and Blanc, P.: Présence de grains éolisés de palygorskite dans les poussières actuelles et les sédiments récents d'origine désertique, *B. Soc. Geol. Fr.*, 1, 571–579, 1985.
- 20 Coudé-Gaussen, G., Hillaire-Marcel, C., and Rognon, P.: Origine et évolution pédologique des fractions carbonatées dans les loess des Matmata (Sud-Tunisien) d'après leurs teneurs en  $^{13}\text{C}$  et  $^{18}\text{O}$ , *C. R. Acad. Sc. Paris*, 295, 939–942, 1982.
- Coudé-Gaussen, G., Rognon, P., Rapp, A., and Nihlén, T.: Dating of peridesert loess in Matmata, south Tunisia, by radiocarbon and thermoluminescence methods, *Z. Geomorp. N. F.*, 31, 129–144, 1987.
- 25 Cram, A. and O'Sullivan, G.: Neogene sapropels in the Mediterranean: a review, *Mar. Geol.*, 153, 11–28, 1999.
- Crouvi, O., Amit, R., Enzel, Y., and Gillespie, A. R.: Active sand seas and the formation of desert loess, *Quaternary Sci. Rev.*, 29, 2087–2098, 2010.
- 30 D'Almeida, G. A.: A model for Saharan dust transport, *J. Clim. Appl. Meteorol.*, 25, 903–916, 1986.

**Atmospheric supply**

V. Bout-Roumazielles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- De Lange, G. J., Thomson, J., Reitz, A., Slomp, C. P., Principato, M. S., Erba, E., and Corselli, C.: Synchronous basin-wide formation and redox-controlled preservation of a Mediterranean sapropel, *Nat. Geosci.*, 1, 606–610, 2008.
- deMenocal, P. B., Ortiz, J., Guilderson, T., Adkins, J., Sarnthein, M., Baker, L., and Yarusinsky, M.: Abrupt onset and termination of the African Humid Period: rapid climatic responses to gradual insolation forcing, *Quaternary Sci. Rev.*, 19, 347–361, 2000.
- Desprat, S., Rouis Zargouni, I., Esselami, L., Turon, J. L., Combourieu Nebout, N., Sicre, M. A., Bout Roumazielles, V., Dewilde, F., and Siani, G.: Holocene vegetation and climatic changes at the southernmost tip of the Central Mediterranean from a direct land-sea correlation, in prep., 2012.
- Di Vito, M. A., Isaia, R., Orsi, G., Southon, J., de Vita, S., D'Antonio, M., Pappalardo, L., and Piocchi, M.: Volcanism and deformation since 12,000 years at the Campi Flegrei caldera (Italy), *J. Volcanol. Geoth. Res.*, 91, 221–246, 1999.
- Dominik, J. and Stoffers, P.: The influence of Late Quaternary stagnations on clay sedimentation in the eastern Mediterranean Sea, *Geol. Rundsch.*, 68, 302–317, 1978.
- Ehrmann, W., Schmiedl, G., Hamann, Y., and Kuhnt, T.: Distribution of clay minerals in surface sediments of the Aegean Sea: a compilation, *Int. J. Earth Sci.*, 96, 769–780, 2007a.
- Ehrmann, W., Schmiedl, G., Hamann, Y., Kuhnt, T., Hemleben, C., and Siebel, W.: Clay minerals in late glacial and Holocene sediments of the northern and southern Aegean Sea, *Paleogeogr. Palaeocl.*, 249, 36–57, 2007b.
- Emeis, K. C., Robertson, A. E. S., Richter, C., and shipboard scientific party: Proc. ODP Init. Repts., 160, Ocean Drilling Program, College Station, TX, doi:10.2973/odp.proc.ir.160.103.1996, 1996.
- Emeis, K.-C., Struck, U., Schulz, H.-M., Bernasconi, S., Sakamoto, T., and Martinez-Ruiz, F.: Temperature and salinity of Mediterranean Sea surface waters over the last 16,000 years: constraints on the physical environment of S1 sapropel formation based on stable oxygen isotopes and alkenone unsaturation ratios, *Paleogeogr. Palaeocl.*, 158, 259–280, 2000.
- Engelstaedter, S., Tegen, I., and Washington, R.: North African dust emissions and transport, *Earth Sci. Rev.*, 79, 73–100, doi:10.1016/j.earsci.2006.06.004, 2006.
- Ergin, M., Bayhan, E., and Temel, A.: Clay mineral distribution in last Glacial-Holocene sediment cores from the eastern Marmara Sea (Cinarcik Basin-Irzmir Gulf Transition), NW-Turkey: Multisources and transport paths in a two-way flow system, *Quaternary Int.*, 261, 53–74, 2012.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Essallami, I., Sicre, M. A., Kallel, N., Labeyrie, L., and Siani, G.: Hydrological changes in the Mediterranean Sea over the last 30,000 years, *Geochem. Geophys. Geos.*, 8, Q07002, doi:10.1029/2007GC001587, 2007.
- 5 Fagel, N. and Hillaire-Marcel, C.: Glacial/interglacial instabilities of the Western Boundary Under Current during the last 365 kyr from Sm/Nd ratios of the sedimentary clay-size fractions at ODP site 646 (Labrador Sea), *Mar. Geol.*, 232, 87–99, 2006.
- Fagel, N., Hillaire-Marcel, C., and Robert, C.: Changes in the Western Boundary Undercurrent outflow since the Last Glacial Maximum, from smectite/illite ratios in deep Labrador Sea sediments, *Paleoceanography*, 12, 79–96, 1997.
- 10 Fink, H. G., Wienberg, C., Hebbeln, D., McGregor, H. V., Schmiedl, G., Taviani, M., and Freiwald, A.: Oxygen control on Holocene cold-water coral development in the eastern Mediterranean Sea, *Deep-Sea Res. Pt. I*, 62, 89–96, 2012.
- Fletcher, W. I. and Sanchez-Goñi, M. F.: Orbital- and sub-orbital-scale climate impacts on vegetation of the western Mediterranean basin over the last 48,000 yr, *Quaternary Res.*, 70, 451–464, 2008.
- 15 Folger, D. W.: Wind transport of land derived mineral, biogenic and industrial matter in the Atlantic, *Deep-Sea Res.*, 17, 433–459, 1970.
- Formenti, P., Rajot, J. L., Desboeufs, K., Saïd, F., Grand, N., Chevaillier, S., and Schmechtig, C.: Airborne observations of mineral dust over western Africa in the summer Monsoon season: spatial and vertical variability of physico-chemical and optical properties, *Atmos. Chem. Phys.*, 11, 6387–6410, doi:10.5194/acp-11-6387-2011, 2011a.
- 20 Formenti, P., Schütz, L., Balkanski, Y., Desboeufs, K., Ebert, M., Kandler, K., Petzold, A., Scheuven, D., Weinbruch, S., and Zhang, D.: Recent progress in understanding physical and chemical properties of African and Asian mineral dust, *Atmos. Chem. Phys.*, 11, 8231–8256, doi:10.5194/acp-11-8231-2011, 2011b.
- Foucault, A. and Mélières, F.: Palaeoclimatic cyclicity in central Mediterranean Pliocene sediments: the mineralogical signal, *Paleogeogr. Palaeocl.*, 158, 311–323, 2000.
- 25 Frigola, J., Moreno, A., Cacho, I., Canals, M., Sierro, F. J., Flores, J. A., Grimalt, J. O., Hodell, D. A., and Curtis, J. H.: Holocene climate variability in the western Mediterranean region from a deepwater sediment record, *Paleoceanography*, 22, PA2209, doi:10.1029/2006PA001307, 2007.
- 30

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Frisia, S., Borsato, A., Mangini, A., Spötl, C., and Madonna, G.: Holocene climate variability from a discontinuous stalagmite record and the Mesolithic to Neolithic transition, *Quaternary Res.*, 66, 388–400, 2006.
- 5 Ganor, E. and Foner, H. A.: The mineralogical and chemical properties and behaviour of aeolian Sahara dust over Israel, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., 163–172, Kluwer Academic, Norwell, Mass, 1996.
- Ganor, E., Foner, H., Brenner, S., Neeman, E., and Lavi, N.: Chemical composition of aerosols settling in Israel following dust storms, *Atmos. Environ.*, 25, 2665–2670, 1991.
- 10 Ganor, E., Deutsch, Y., and Foner, H. A.: Mineralogical composition and sources of airborne settling particles on Lake Kinneret (the sea of Galilee) in Israel, *Water Air Soil Poll.*, 118, 245–262, doi:10.1023/A:1005167230795, 2000.
- Garzanti, E., Ando, S., Vezzoli G., Abdel Megid, A. A., and El Kammar, A.: Petrology of Nile River sands (Ethiopia and Sudan): sediment budgets and erosion patterns, *Earth Planet. Sc. Lett.*, 252, 327–341, 2006.
- 15 Gasse, F.: Hydrological changes in the African tropics since the Last Glacial Maximum, *Quaternary Sci. Rev.*, 19, 189–211, 2000.
- Gingele, F., De Decker, P., and Hillenbrand, C.-D.: Clay mineral distribution in surface sediments between Indonesia and NW Australia – source and transport by ocean currents, *Mar. Geol.*, 179, 135–146, 2001.
- 20 Giraudi, C.: Eolian sand in peridesert northwestern Libya and implications for Late Pleistocene and Holocene Sahara expansions, *Paleogeogr. Palaeocl.*, 218, 161–173, 2005.
- Giraudi, C., Magny, M., Zanchetta, G., and Drysdale, R. N.: The Holocene climatic evolution of the Mediterranean Italy: A review of the continental geological data, *Holocene*, 21, 105–115, 2011.
- 25 Glaccum, R. A. and Prospero, J. M.: Saharan aerosols over the tropical north Atlantic – Mineralogy, *Mar. Geol.*, 37, 295–321, 1980.
- Gomes, L., Bergametti, G., Coudé-Gaussen, G., and Rognon, P.: Submicron desert dust: A sandblasting process, *J. Geophys. Res.*, 95, 927–935, 1990.
- Goudie, A. S.: *Great warm deserts of the world*, Oxford University Press, Oxford, 2003.
- 30 Goudie, A. S. and Middleton, N. J.: Saharan dust storms: nature and consequences, *Earth Sci. Rev.*, 56, 179–204, 2001.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Grousset, F. E., Rognon, P., Coudé-Gaussen, G., and Pédemay, P.: Origins of peri-Saharan dust deposits traced by their Nd and Sr isotopic composition, *Paleogeogr. Palaeoclimatol.*, 93, 203–212, 1992.

Guerzoni, S. and Chester, R.: The impact of Desert Dust across the Mediterranean, Dordrecht, Kluwer Academic, 1996.

Guerzoni, S., Molinaroli, E., and Chester, R.: Saharan dust inputs to the Western Mediterranean Sea: depositional patterns, geochemistry and sedimentological implications, *Deep-Sea Res. Pt. II*, 44, 631–654, 1997.

Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loyé-Pilot, M.-D., Measures, C., Migon, C., Molinaroli, E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, *Prog. Oceanogr.*, 44, 147–190, 1999.

Guieu, C., Loyé-Pilot, M. D., Ridame, C., and Thomas, C.: Chemical characterization of the Saharan dust end-member: Some biogeochemical implications for the western Mediterranean Sea, *J. Geophys. Res.*, 107, 4258, doi:10.1029/2001jd000582, 2002.

Hamann, Y., Ehrmann, W., Schmiedl, G., Krüger, S., Stuetz, J.-B., and Kuhnt, T.: Sedimentation processes in the Eastern Mediterranean Sea during the Late Glacial and Holocene revealed by end-member modelling of the terrigenous fraction in marine sediments, *Mar. Geol.*, 248, 97–114, 2008.

Hamann, Y., Ehrmann, W., Schmiedl, G., and Kuhnt, T.: Modern and late Quaternary clay mineral distribution in the area of the SE Mediterranean Sea, *Quaternary Res.*, 71, 453–464, 2009.

Hamann, Y., Bowen, S. W., Ersoy, O., Ehrmann, W., and Aydar, E.: First evidence of a distal early Holocene ash layer in Eastern Mediterranean deep-sea sediments derived from the Anatolian volcanic province, *Quaternary Res.*, 73, 497–506, 2010.

Haug, G., Hughen, K. A., Sigman, D. M., Peterson, L. C., and Röhl, U.: Southward migration of the Intertropical Convergence Zone through the Holocene, *Science*, 293, 1304–1308, 2001.

Hilgen, H. J.: Astronomical calibration of Gauss to Matuyama sapropels in the Mediterranean and implication for the geomagnetic polarity time scale, *Earth Planet. Sc. Lett.*, 104, 226–244, 1991.

Holeman, J. N.: The sediment yield of major rivers of the world, *Water Resour. Res.*, 4, 737–744, 1968.



**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Incarbona, A., Martrat, B., Di Stefano, E., Grimalt, J. O., Pelosi, N., Patti, B., and Tranchida, G.: Primary productivity variability on the Atlantic Iberian margin over the last 70,000 years: evidence from coccolithophores and fossil organic compounds, *Paleoceanography*, 25, PA2218, doi:10.1029/2008PA001709, 2010.
- 5 Israelevich, P. L., Levin, Z., Joseph, J. H., and Ganor, E.: Desert aerosol transport in the Mediterranean region as inferred from TOMS aerosol index, *J. Geophys. Res.*, 107, 4572, doi:10.1029/2001jd002011, 2002.
- Israelevich, P., Ganor, E., Alpert, P., Kishcha, P., and Stupp, A.: Predominant transport paths of Saharan dust over the Mediterranean Sea to Europe, *J. Geophys. Res.*, 117, D02205, doi:10.1029/2011JD016482, 2012.
- 10 Jalut, G., Deboubat, J. J., Fontugne, M., and Otto, T.: Holocene circum-Mediterranean vegetation changes: climate forcing and human impact, *Quaternary Int.*, 200, 4–18, 2009.
- Jones, G. A.: Advective transport of clay minerals in the region of the Rio Grande rise, *Mar. Geol.*, 58, 187–212, 1984.
- 15 Kallel, N., Paterne, J. C., Duplessy, C., Vergnaud-Grazzini, C., Labeyrie, L., Arnold, M., Fontugne, M., and Pierre, C.: Enhanced rainfall in the Mediterranean region during the last sapropel event, *Oceanol. Acta*, 20, 697–712, 1997.
- Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knipertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUN 2006, *Tellus B*, 61, 32–50, 2009.
- 20 Kidd, R. B., Cita, M. B., and Ryan, W. B. F.: Stratigraphy of the Eastern Mediterranean Sapropel Sequences Recovered during DSDP LEG 42A and Their Paleoenvironmental Significance, edited by: Hsü, K. J., Montadert, L., Garrison, R. E., Fabricius, F. H., Kidd, R. B., Müller, C., Cita, M. B., Bizon, G., Wright, R. C., Erickson, A. J., Bernoulli, D., and Mélières, F., *Initial Rep. Deep Sea 42*, US Govt. Printing Office, Washington, DC, 421–443, 1978.
- 25 Kiefert, L., McTainsh, G. H., and Nickling, W. G.: Sedimentological characteristics of Saharan and Australian Dusts, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic Publishers, Dordrecht, 183–190, 1996.
- 30 Kim, J., Jung, C. H., Choi, B.-C., Oh, S.-N., Brechtel, F. J., Yoon, S.-C., and Kim, S.-W.: Number size distribution of atmospheric aerosols during ACE-Asia dust and precipitation events, *Atmos. Environ.*, 41, 4841–4855, 2007.



**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Klein, B., Roether, W., Manca, B. B., Bregant, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: The large deep water transient in the Eastern Mediterranean, *Deep-Sea Res. Pt. I*, 46, 371–414, 1999.
- 5 Koren, I., Kaufman, Y. J., Washington, R., Todd, M. C., Rudich, Y., Vanderlei Martins, J., and Rosenfeld, D.: The Bodélé depression: A single spot in the Sahara that provides most mineral dust to the Amazon forest, *Environ. Res. Lett.*, 1, 014005, doi:10.1088/1748-9326/1/1/014005, 2006.
- Kotthoff, U., Pross, J., Müller, U. C., Peyron, O., Schmiedl, G., Schulz, H., and Bordon, A.: Climate dynamics in the borderlands of the Aegean Sea during formation of sapropel S1 deduced from a marine pollen record, *Quaternary Sci. Rev.*, 27, 832–845, 2008.
- 10 Krom, M. D., Stanley, J. D., Cliff, R. A., and Woodwad, J. C.: Nile River sediment fluctuations over the past 7000 yr and their key role in sapropel development, *Geology*, 30, 71–74, 2002.
- Lascaratos, A. and Nittis, K.: A high-resolution three-dimensional numerical study of intermediate water formation in the Levantine Sea, *J. Geophys. Res.*, 103, 18497–18511, 1998.
- 15 Lézine, A.-M., Turon, J. L., and Buchet, G.: Pollen analyses off Senegal: evolution of the coastal environment during the last deglaciation, *J. Quaternary Sci.*, 10, 95–105, 1995.
- Liu, Z., Trentesaux, A., Clemens, S. C., Colin, C., Wang, P., Huang, B., and Boulay, S.: Clay mineral assemblages in the northern South China Sea: implications for East Asian monsoon evolution over the past 2 million years, *Mar. Geol.*, 201, 133–146, 2003.
- 20 Lourens, L. J., Antonarakou, F. J., Hilgen, F. J., Van Hoof, A. A. M., Vergnaud Grazzini, C., and Zachariasse, W. J.: Evaluation of the Plio-Pleistocene astronomical time-scale, *Paleoceanography*, 11, 391–431, 1996.
- Luck, J. M. and Ben Othman, D.: Trace element and Pb isotope variability during rainy events in the NW Mediterranean: constraints on anthropogenic and natural sources, *Chem. Geol.*, 182, 443–460, 2002.
- 25 Magny, M., Miramont, C., and Sivan, O.: Assessment of the impact of climate and anthropogenic factors on Holocene Mediterranean vegetation in Europe on the basis of palaeohydrological records, *Paleogeogr. Palaeoclimatol.*, 186, 47–59, 2002.
- Magny, M., Bégeot, C., Guiot, J., and Peyron, O.: Contrasting patterns of hydrological changes in Europe in response to Holocene climate cooling phases, *Quaternary Sci. Rev.*, 22, 1589–30 1596, 2003.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Magny, M., Vanni re, B., de Beaulieu, J.-L., B geot, C., Heiri, O., Millet, L., Peyron, O., and Walter-Simonnet, A. V.: Early-Holocene climatic oscillations recorded by lake-level fluctuations in west-central Europe and in central Italy, *Quaternary Sci. Rev.*, 26, 1951–1964, 2007.
- 5 Magny, M., Vanni re, B., Zanchetta, G., Fouache, E., Touchais, G., Petrika, L., Coussot, C., Walter-Simonnet, A. V., and Arnaud, F.: Possible complexity of the climatic event around 4300–3800 cal BP in the central and western Mediterranean, *Holocene*, 19, 823–833, 2009.
- Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vanni re, B., and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the south- and north-central Mediterranean, *J. Quaternary Sci.*, 27, 290–296, 2012.
- 10 Malanotte-Rizzoli, P. and Hecht, A.: Large-scale properties of the Eastern Mediterranean: a review, *Oceanol. Acta*, 11, 323–335, 1988.
- Malanotte-Rizzoli, P., Manca, B. B., Ribera, D., Alcal , M., Theocharis, A., Brenner, S., Budillon, G., and Ozsoy, E.: The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations, *Dynam. Atmos. Oceans*, 29, 365–395, 1999.
- 15 Martin, J. M. and Milliman, J. D.: EROS 2000 (European Rivers Ocean System). The western Mediterranean Sea: an introduction, *Deep-Sea Res.*, 44, 521–529, 1997.
- Martinez-Ruiz, F., Paytan, A., Kastner, M., Gonzalez-Donoso, J. M., Linares, D., Bernasconi, S., and Jimenez-Espejo, F. J.: A comparative study of the geochemical and mineralogical characteristics of the S1 sapropel in the western and eastern Mediterranean, *Palaeogeogr. Palaeoclimatol.*, 190, 23–37, 2003.
- 20 Matthewson, A. P., Shimmield, G., Kroon, D., and Fallick, A. E.: A 300 kyr high-resolution aridity record of the North African continent, *Paleoceanography*, 10, 677–692, 1995.
- Mayewski, P. A., Rohling, E. J., Stager, J. C., Karlen, W., Maasch, K. A., Meeker, L. D., Meyerson, E. A., Gasse, F., Van Kreveld, S., Holmgren, K., Lee-Thorp, J., Rosqvist, G., Rack, F., Staubwasser, M., Schneider, R. R., and Steig, E. J.: Holocene climate variability, *Quaternary Res.*, 62, 243–255, 2004.
- 25 McTainsh, G. H., Nickling, W. G., and Lynch, A. W.: Dust deposition and particle size in Mali, West Africa, *Catena*, 29, 307–322, 1997.
- Mercone, D., Thomson, J., Croudace, I. W., Siani, G., Paterne, M., and Troelstra, S.: Duration of S1, the most recent sapropel in the eastern Mediterranean Sea, as indicated by accelerator mass spectrometry radiocarbon and geochemical evidence, *Paleoceanography*, 15, 336–347, 2000.
- 30

**Atmospheric supply**

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Milliman, J. D. and Syvitski, J. P. M.: Geomorphic/Tectonic control of sediment discharge to the ocean: The importance of small mountainous rivers, *J. Geol.*, 100, 520–540, 1992.
- Molinarioli, E.: Mineralogical characterization of Saharan dust with a review to its final destination in Mediterranean sediments, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic, 153–162, 1996.
- 5 Montero-Serrano, J. C., Bout-Roumazielles, V., Tribovillard, N., Sionneau, T., Riboulleau, A., Bory, A., and Flower, B.: Sedimentary evidence of deglacial megafloods in the northern Gulf of Mexico (Pigmy Basin), *Quaternary Sci. Rev.*, 28, 3333–3347, 2009.
- 10 Montero-Serrano, J. C., Bout-Roumazielles, V., Sionneau, T., Tribovillard, N., Bory, A., Flower, B. P., Riboulleau, A., Martinez, P., and Billy, I.: Changes in precipitation regimes over North America during the Holocene as recorded by mineralogy and geochemistry of Gulf of Mexico sediments, *Global Planet. Change*, 74, 132–143, 2010.
- Moulin, C., Lambert, C. E., Dulac, F., and Dayan, U.: Control of atmospheric export of dust from North Africa by the North Atlantic Oscillation, *Nature*, 387, 691–694, 1997.
- 15 O'Hara, S. L., Clarke, M. L., and Elatrash, M. S.: Field measurements of desert dust deposition in Libya, *Atmos. Environ.*, 40, 3881–3897, 2006.
- Paquet, H., Coudé-Gaussen, G., and Rognon, P.: Etude minéralogique de poussières sahariennes le long d'un itinéraire entre 19° et 35° de latitude nord, *Rev. Geol. Dyn. Geogr.*, 25, 257–265, 1984.
- 20 Petschick, R.: MacDiff v4.2.5 (Free Geological Software). Geologisch-Paläontologisches Institut, Universität Frankfurt/Main, available at: <http://www.geol-pal.uni-frankfurt.de/Staff/Homepages/Petschick/classicsoftware.html#MacDiff> (last access: 25 July 2012), 2001.
- Petschick, R., Kuhn, G., and Gingele, F.: Clay mineral distribution in surface sediments of the South Atlantic: Sources, transport, and relation to oceanography, *Mar. Geol.*, 130, 203–230, 1996.
- 25 Peyron, O., Goring, S., Dormoy, I., Kothoff, U., Porss, J., de Beaulieu, J. L., La Mantia, T., Schneider, R., and Magny, M.: Holocene seasonality changes in the central Mediterranean region reconstructed from the pollen sequences of Lake Accessa (Italy) and Tenaghi Philippon (Greece), *Holocene*, 21, 131–146, 2011.
- 30 Pickard, G. L. and Emery, W. J.: *Descriptive Physical Oceanography – An Introduction*, Pergamon Press, San Diego, 249 pp., 1982.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: a review, *Palaeogeogr. Palaeoclimatol.*, 158, 153–174, 2000.
- Pinsak, A. P. and Murray, H. H.: Regional clay mineral patterns in the Gulf of Mexico, *Clays and Clay Minerals*, 7th National conference, 162–177, 1960.
- Pittalwala, I. and Hameed, I.: Simulation of the North Atlantic Oscillation in a general circulation model, *Geophys. Res. Lett.*, 18, 841–844, 1991.
- Prospero, J. M.: Arid regions as sources of minerals aerosols in the marine atmosphere, *Geol. Soc. Am. S.*, 186, 71–86, 1981.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the nimbus 7 total ozone mapping spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40, 1002, doi:10.1029/2000RG000095, 2002.
- Pye, K.: *Aeolian dust and dust deposits*, New-York, London, Academic Press, 334 pp., 1987.
- Pye, K.: Aeolian dust transport and deposition over Crete and adjacent parts of the Mediterranean Sea, *Earth Surf. Proc. Land.*, 17, 271–288, 1992.
- Pye, K. and Tsoar, H.: The mechanics and geological implications of dust transport and deposition in deserts with particular reference to loess formation and dune sand diagenesis in the northern Negev, Israel, in: *Desert Sediments Ancient and Modern*, edited by: Frostick, L. E. and Reid, I., *Geol. Soc. Spec. Publ.*, 35, 139–156, 1987.
- Ramrath, A., Sadori, L., and Negendank, J. F. W.: Sediments from Lago di Mezzano, central Italy: A record of Lateglacial/Holocene climatic variations and anthropogenic impact, *Holocene*, 10, 87–95, 2000.
- Revel, M., Ducassou, E., Grousset, F. E., Bernasconi, S. M., Migeon, S., Revillon, S., Mascle, J., Murat, A., Zaragosi, S., and Bosch, D.: 100,000 years of African monsoon variability recorded in sediments of the Nile margin, *Quaternary Sci. Rev.*, 29, 1342–1362, 2010.
- Robert, C., Gauthier, A., and Chamley, H.: Origine autochtone et allochtone des argiles récentes de haute altitude en Corse, *Géologie de la Méditerranée*, XI, 243–253, 1984.
- Roberts, N., Jones, M. D., Benkaddour, A., Eastwood, W. J., Filippi, M. L., Frogley, M. R., Lamb, H. F., Leng, M. J., Reed, J. M., Stein, M., Stevens, L., Valero-Garces, B., and Zanchetta, G.: Stable isotope records of Late Quaternary climate and hydrology from Mediterranean lakes: the ISOMED synthesis, *Quaternary Sci. Rev.*, 27, 2426–2441, 2008.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Roberts, N., Brayshaw, D., Kuzucuoglu, C., Perez, R., and Sadori, L.: The mid-Holocene climatic transition in the Mediterranean: causes and consequences, *Holocene*, 21, 3–14, 2011.
- Rodriguez, S., Querol, X., Alastuey, A., Kallos, G., and Kakaliagou, O.: Saharan dust contributions to PM10 and TSP levels in Southern and Eastern Spain, *Atmos. Environ.*, 35, 2433–2447, 2001.
- 5 Rohling, E. J.: Review and new aspects concerning the formation of eastern Mediterranean sapropels, *Mar. Geol.*, 122, 1–28, 1994.
- Rossignol-Strick, M.: Mediterranean Quaternary sapropels, an immediate response to the African monsoon to variation of insolation, *Paleogeogr. Palaeocl.*, 49, 237–263, 1985.
- 10 Rossignol-Strick, M., Nesteroff, W., Olive, P., and Vernaud-Grazzini, C.: After the deluge: Mediterranean stagnation and sapropel formation, *Nature*, 295, 105–110, 1982.
- Rouis-Zargouni, I., Turon, J.-L., Londeix, L., Essallami, L., Kallel, N., and Sicre, M.-A.: Environmental and climatic changes in the central Mediterranean Sea (Siculo-Tunisian Strait) during the last 30 ka based on dinoflagellate cyst and planktonic *foraminifera* assemblages, *Paleogeogr. Palaeocl.*, 285, 17–29, 2010.
- 15 Rouis-Zargouni, I., Tron, J. L., Londeix, L., Kallel, N., and Essallami, L.: The last glacial-interglacial transition and dinoflagellate cysts in the western Mediterranean sea, *C. R. Geosci.*, 344, 99–109, doi:10.1016/j.crte.2012.01.002, 2012.
- Sadori, L., Giraudi, C., Pettitti, P., and Ramrath, A.: Human impact at Lago di Mezzano (central Italy) during the Bronze Age: a multidisciplinary approach, *Quaternary Int.*, 113, 5–17, 2004.
- 20 Sadori, L., Zanchetta, G., and Giardini, M.: Last Glacial to Holocene palaeoenvironmental evolution at Lago di Pergusa (Sicily, Southern Italy) as inferred by pollen, microcharcoal, and stable isotopes, *Quaternary Int.*, 181, 4–14, 2008.
- Sadori, L., Jahns, S., and Peyron, O.: Mid-Holocene vegetation history of the central Mediterranean, *Holocene*, 21, 117–129, 2011.
- 25 Saliot, A.: The Mediterranean Sea. The Handbook of Environmental Chemistry, v.5, Water Pollution, Springer-Verlag, Berlin, 408 pp., 2005.
- Sánchez-Goñi, M. F., Cacho, I., Turon, J. L., Guiot, J., Sierro, F. J., Peyrouquet, J.-P., Grimalt, J. O., and Shackleton, N.: Synchronicity between marine and terrestrial responses to millennial scale climatic variability during the last glacial period in the Mediterranean region, *Clim. Dynam.*, 19, 95–105, 2002.
- 30 Sawlowicz, Z.: Iridium and other platinum-group elements as geochemical markers in sedimentary environments, *Paleogeogr. Palaeocl.*, 104, 253–270, 1993.

**Atmospheric supply**

V. Bout-Roumazeilles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schilman, B., Bar-Matthews, M., Almogi-Labin, A., and Luz, B.: Global climate instability reflected by Eastern Mediterranean marine records during the late Holocene, *Paleogeogr. Palaeocl.*, 176, 157–176, 2001.

Schütz, L. and Sebert, M.: Mineral aerosols and source identification, *J. Aerosol Sci.*, 18, 1–10, 1987.

Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., and Haddad, G.: Mediterranean sea-surface radiocarbon reservoir age changes since the last glacial maximum, *Science*, 294, 1917–1920, 2001.

Siani, G., Sulpizio, R., Paterne, M., and Sbrana, A.: Tephrostratigraphy study for the last 18,000 <sup>14</sup>C years in a deep-sea sediment sequence for the South Adriatic, *Quaternary Sci. Rev.*, 23, 2485–2500, 2004.

Sionneau, T., Bout-Roumazeilles, V., Biscaye, P. E., Van Vliet-Lanoe, B., and Bory, A.: Clay mineral distributions in and around the Mississippi River watershed and Northern Gulf of Mexico: sources and transport patterns, *Quaternary Sci. Rev.*, 27, 1740–1751, 2008.

Sionneau, T., Bout-Roumazeilles, V., Flower, B. P., Bory, A., Tribouvillard, N., Kissel, C., Van Vliet-Lanoe, B., and Montero Serrano, J. C.: Provenance of freshwater pulses in the Gulf of Mexico during the last deglaciation, *Quaternary Res.*, 74, 235–245, 2010.

Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Stuut, J.-B., Meyer, I., Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., and Ndiaye, T.: The 7–13 March 2006 major Saharan outbreak: Multi-proxy characterization of mineral dust deposited on the West African margin, *J. Geophys. Res.*, 116, D18210, doi:10.1029/2011JD016173, 2011.

Stanley, D. J. and Wingerath, J. G.: Clay mineral distributions to interpret Nile cell provenance and dispersal, I. Lower River Nile to delta sector, *J. Coastal Res.*, 12, 911–929, 1996.

Stanley, D. J., Warne, A. G., Davis, H. R., Bernasconi, M. P., and Chen, Z.: Nile Delta, *Nat. Geogr. Res. Expl.*, 8, 22–51, 1992.

Stevens, L. R., Wright Jr., H. E., and Ito, E.: Proposed changes in seasonality of climate during the late-glacial and Holocene at Lake Zeribar, Iran, *Holocene*, 11, 747–756, 2001.

Stumpf, R., Frank, M., Schönfeld, J., and Haley, B. A.: Late Quaternary variability of Mediterranean Outflow Water from radiogenic Nd and Pb isotopes, *Quaternary Sci. Rev.*, 29, 2462–2572, doi:10.1016/j.quascirev.2010.06.021, 2010.

- Stumpf, R., Frank, M., Schönfeld, J., and Haley, B. A.: Climatically driven changes in sediment supply on the SW Iberian shelf since the last Glacial Maximum, *Earth Planet. Sc. Lett.*, 312, 80–90, 2011.
- Tomadin, L.: Sedimentary fluxes and different dispersion mechanisms of the clay sediments in the Adriatic Basin, *Rend. Fis. Acc. Lincei*, 9, 161–174, 2000.
- Torres-Padrón, M. E., Gelado-Caballero, M. D., Collado-Sánchez, C., Siruela-Matos, V. F., Cardona-Castellano, P. J., and Hernández-Brito, J. J.: Variability of dust inputs to the CANIGO zone, *Deep-Sea Res. Pt. II*, 49, 3455–3464, 2002.
- Tribouvillard, N., Bout-Roumazielles, V., Algeo, T., Lyons, T. W., Sionneau, T., Montero-Serrano, J. C., Riboulleau, A., and Baudin, F.: Paleodepositional conditions in the Orca Basin as inferred from organic matter and trace metal contents, *Mar. Geol.*, 254, 62–72, 2008.
- Tzedakis, P. C.: Seven ambiguities in the Mediterranean palaeoenvironmental narrative, *Quaternary Sci. Rev.*, 26, 2042–2066, 2007.
- Ukstins, I. A., Renne, P. R., Wolfenden, E., Baker, J. A., Ayalew, D., and Menzeis, M.: Matching conjugate volcanic rifted margins: 40Ar/39Ar chrono-stratigraphy of pre-and syn-rift bimodal flood volcanism in Ethiopia and Yemen, *Earth Planet. Sc. Lett.*, 198, 289–306, 2002.
- Vannière, B., Power, M. J., Roberts, N., Tinner, W., Carrion, J., Magny, M., and Bartlein, P.: Circum-Mediterranean fire activity and climate changes during the mid Holocene environmental transition (8500–2500 cal yr BP), *Holocene*, 21, 53–73, 2011.
- Venkatarathnam, K. and Ryan, W. B. F.: Dispersal patterns of clay minerals in the sediments of the Eastern Mediterranean Sea, *Mar. Geol.*, 11, 261–282, 1971.
- Washington, R. and Todd, M. C.: Atmospheric controls on mineral dust emission from the Bodélé depression, Chad: the role of the low level jet, *Geophys. Res. Lett.*, 32, L17701, doi:10.1029/2005GL023597, 2005.
- Weldeab, S., Siebel, W., Wehausen, R., Emeis, K.-C., Schmiedl, G., and Hemleben, C.: Late Pleistocene sedimentation in the Western Mediterranean Sea: implications for productivity changes and climatic conditions in the catchment areas, *Paleogeogr. Palaeocl.*, 190, 121–137, 2003.
- Wüst, G.: On the vertical circulation on the Mediterranean Sea, *J. Geophys. Res.*, 66, 3261–3271, 1961.
- Zanchetta, G., Drysdale, R. N., Hellstrom, J. C., Fallick, A. E., Isola, I., Gagan, M. K., and Pareschi, M. T.: Enhanced rainfall in the Western Mediterranean during deposition of sapro-

**Atmospheric supply**

V. Bout-Roumazielles et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





pel S1: Stalagmite evidence from Corchia cave (Central Italy), *Quaternary Sci. Rev.*, 26, 279–286, 2007.

Zanchetta, G., Sulpizio, R., Roberts, N., Cioni, R., Eastwood, W. J., Siani, G., Caon, B., Pat-  
terne, M., and Santacroce, R.: Tephrostratigraphy, chronology and climatic events of the  
5 Mediterranean basin during the Holocene: An overview, *Holocene*, 21, 33–52, 2011.

Zielhofer, C. and Faust, D.: Mid- and Late Holocene fluvial chronology of Tunisia, *Quaternary  
Sci. Rev.*, 27, 580–588, 2008.

## Atmospheric supply

V. Bout-Roumazeilles et  
al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Atmospheric supply

V. Bout-Roumazeilles et al.

**Table 1.** Radiocarbon ages performed on core MD04-2797. The ages were corrected using reservoir ages of 400 yr during the Holocene, the Younger Dryas and late glacial, 560 yr during the Bølling-Allerød and 800 yr during the Heinrich event H1 and the Older Dryas, following recommendations of Siani et al. (2001).

Depth (cm)	Species	<sup>14</sup> C age (yr BP)	Error ±1σ	Corrected age (yr BP)	Error ±1σ	Calibrated age (cal yr BP)	Error ±1σ	Laboratory
0	<i>G. inflata</i>	1105*	20	705	20	668	9	ARTEMIS
80	<i>foraminifera</i>	5496	95	5093	95	5827	144	ARTEMIS
160	<i>foraminifera</i>	6700	85	6300	85	7241	120	ARTEMIS
199	<i>foraminifera</i>	7523	81	7123	81	7967	71	ARTEMIS
240	<i>foraminifera</i>	8113	81	7713	81	8488	98	ARTEMIS
330	<i>G. ruber</i>	8888	110	8488	110	9477	112	ARTEMIS
410	<i>foraminifera</i>	10 863	32	10 463	32	12 493	49	ARTEMIS
470	<i>G. inflata</i>	12 728	173	12 168	172	14 016	35	ARTEMIS
510	<i>G. ruber</i>	13 900	141	13 100	141	15 934	621	ARTEMIS
610	<i>G. ruber</i>	15 590	50	15 190	50	18 725	370	ARTEMIS
700	<i>foraminifera</i>	17 660	70	17 260	70	20 414	180	ARTEMIS
940	<i>foraminifera</i>	23 415	163	23 015	163	27 879	331	ARTEMIS
1030	<i>foraminifera</i>	26 095	7	25 695	7	30 490	156	ARTEMIS

\* Data from Rouis-Zargouni et al. (2010).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion








## Atmospheric supply

V. Bout-Roumazielles et al.



**Fig. 2.** Clay mineralogy of core MD04-2797: kaolinite (%), smectite (%), illite (%), chlorite (%), palygorskite (%), I/K : illite to kaolinite ratio, Kübler index (crystallinity), (MART) Terrigenous Mass Accumulation Rates ( $\text{g kyr}^{-1} \text{cm}^{-2}$ ) and quartz content (counts/s) for the last 20 000 yr. B-A = Bølling-Allerød, YD = Younger Dryas, S1 = sapropel S1 (Mercone et al., 2000).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Atmospheric supply

V. Bout-Roumazeilles et al.



**Fig. 3.** Grain size analysis of core MD04-2797: mode ( $\mu\text{m}$ ), % of cohesive silt ( $2 < > 10 \mu\text{m}$ ), % of sortable silt ( $10 < > 63 \mu\text{m}$ ), % of clay ( $< 2 \mu\text{m}$ ) and % of sand ( $> 63 \mu\text{m}$ ). MART from Fig. 1. Sea Surface Temperatures (SST) for May–June and October–November based on *foraminifera* association (Essallami et al., 2007),  $\delta^{18}\text{O}$  (‰) and  $\delta^{13}\text{C}$  from planktonic *foraminifera* *G. bullioides* (Essallami et al., 2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Atmospheric supply

V. Bout-Roumazielles et al.



**Fig. 4.** Geochemical data for core MD04-2797, Al content (counts/s), Ti/Al ratio, Zr/Al ratio, Ca/Fe ratio and Pd content (counts/s). MART from Fig. 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Atmospheric supply

V. Bout-Roumazeilles et al.



**Fig. 5.** Comparison of multiproxy data, **(a):** I/K, Ca/Fe and their interpretation on particles provenance for the Bølling-Allerød, **(b):** grain-size compared with Zr/Al during the Younger Dryas, **(c):** quartz content (counts/s), Ti/Al ratio and kaolinite % over the last 3000 yr, **(d):** Smectite (%), characterizing the sapropel S1, % of clay, content in Pd (counts/s) and MART.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Fig. 6.** Comparison of the mineralogical and geochemical signatures before and after 11.7 kyr, **(a)**: kaolinite versus illite (XRD counts/s) evidencing contrasting provenance between 18.5 and 11.7 ka (red squares) and between 11.7 ka and top of the core (blue squares); **(b)**: illite to kaolinite ratio as a function of illite crystallinity (Kübler Index), diagenesis, epizone and anchizone limits, **(c)**: Ca content versus Fe content (counts XRF).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Atmospheric supply

V. Bout-Roumazeilles et al.



**Fig. 7.** Main provenance (dashed areas) and potential eolian – low level (red arrows) or altitude winds (black arrows) – and riverine (green arrows) transportation patterns of clay-mineral particles for 5 timeslices, **(a)** Bølling-Allerød; **(b)** Younger Dryas; **(c)** early Holocene; **(d)** Sapropel S1; **(e)** late Holocene.