



HAL
open science

Deglacial and Holocene vegetation and climatic changes at the southernmost tip of the Central Mediterranean from a direct land-sea correlation

S Desprat, N Combourieu-Nebout, L Essallami, M A Sicre, I Dormoy, O Peyron, G Siani, Viviane Bout-roumazeilles, J L Turon

► To cite this version:

S Desprat, N Combourieu-Nebout, L Essallami, M A Sicre, I Dormoy, et al.. Deglacial and Holocene vegetation and climatic changes at the southernmost tip of the Central Mediterranean from a direct land-sea correlation. 2021. hal-03280601

HAL Id: hal-03280601

<https://hal.science/hal-03280601>

Preprint submitted on 7 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

Deglacial and Holocene vegetation and climatic changes at the southernmost tip of the Central Mediterranean from a direct land-sea correlation

S. Desprat¹, N. Combourieu-Nebout², L. Essallami³, M. A. Sicre², I. Dormoy⁴,
O. Peyron⁴, G. Siani⁵, V. Bout Roumazeilles⁶, and J. L. Turon⁷

¹EPHE, Environnements et Paléoenvironnements Océaniques et Continentaux (EPOC),
UMR5805, CNRS – Université Bordeaux 1, 33405 Talence, France

²LSCE, UMR1572, CNRS/CEA/UVSQ, 91198 Gif-sur-Yvette Cedex, France

³GEOGLOB, Sfax Faculty of Sciences, 3038 Sfax, Tunisia

⁴Chrono-Environnement, UMR6249, CNRS – Université de Franche-Comté,
25030 Besançon, France

⁵IDES, Earth Sciences Department, Université Paris XI, 91405 Orsay, France

⁶Géosystèmes, UMR8217, CNRS – Université Lille 1, 59655 Villeneuve d'Ascq, France

⁷EPOC, UMR5805, CNRS – Université Bordeaux 1, 33405 Talence, France

5687

Received: 5 November 2012 – Accepted: 9 November 2012 – Published: 21 November 2012

Correspondence to: S. Desprat (s.desprat@epoc.u-bordeaux1.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

Despite a large number of studies, the long-term and millennial to centennial-scale climatic variability in the Mediterranean region during the last deglaciation and the Holocene is still debated, in particular in the Southern Central Mediterranean. In this paper, we present a new marine pollen sequence (MD04-2797CQ) from the Siculo-Tunisian Strait documenting the regional vegetation and climatic changes in the Southern Central Mediterranean during the last deglaciation and the Holocene.

The MD04-2797CQ marine pollen sequence shows that semi-desert plants dominated the vegetal cover in the Southern Central Mediterranean between 18 and 12.3 kyr BP indicating prevailing dry conditions during the deglaciation, even during the Greenland Interstadial (GI)-1. Such arid conditions likely restricted the expansion of the trees and shrubs despite the GI-1 climatic amelioration. Across the transition Greenland Stadial (GS)-1 – Holocene, Asteraceae-Poaceae steppe became dominant till 10.1 kyr. This record underlines with no chronological ambiguity that even though temperatures increased, deficiency in moisture availability persisted into the Early Holocene. Temperate trees and shrubs with heaths as oak forest understorey or heath maquis expanded between 10.1 and 6.6 kyr, while Mediterranean plants only developed from 6.6 kyr onwards. These changes in vegetal cover show that the regional climate in Southern Central Mediterranean was wetter during Sapropel 1 (S1) and became drier during the Mid- to Late Holocene. Wetter conditions during S1 were likely due to increased winter precipitation while summers remained dry. We suggest, in agreement with published modelling experiments, that the increased melting of the Laurentide Ice Sheet between 10 to 6.8 kyr in conjunction with weak winter insolation played a major role in the development of winter precipitation maxima in the Mediterranean region in controlling the strength and position of the North Atlantic storm track.

Finally, our data provide evidences of centennial-scale vegetation and climatic changes in the Southern Central Mediterranean. During the wet Early Holocene, alkenones-derived cooling episodes are synchronous to herbaceous composition

5689

changes that indicate muted changes in precipitation. In contrast, enhanced aridity episodes, as detected by strong reduction in trees and shrubs, are recorded during the Mid- to Late Holocene. We show that the impact of the Holocene cooling events depend on the baseline climate states insolation and ice sheet volume, shaping the response of the mid-latitude atmospheric circulation.

1 Introduction

The Mediterranean region received increasing attention during the past decade when it was identified as a major “climate change hot spot” in future climate modeling projections as a result of the global increase in greenhouse gases (Giorgi, 2006; IPCC, 2007). Sensitivity of the Mediterranean climate and ecosystems to global climate change during glacial and interglacial periods has been demonstrated by a numerous paleoclimatic studies. Holocene Mediterranean records show that the long-term Holocene climatic evolution in response to orbital forcing can be described by three phases: a humid phase from 11.5 to 7 kyr, a transition phase from 7 to 5.5 kyr and an aridification phase from 5.5 kyr to present-day (Jalut et al., 2009; Finné et al., 2011; Pérez-Obiol et al., 2011). However, Holocene environmental conditions in the Mediterranean region remain a matter of debate. Davis and Brewer (2009) even hypothesized that higher Early Holocene moisture availability was not related to increased precipitation but decreased temperature. The respective contribution of climate and human activities to environmental changes, the contrasting and complex information provided by different proxy records and the heterogeneity and complexity of this region contribute to the existing uncertainties on the Holocene climate evolution in the different regions of the Mediterranean (Tzedakis, 2007; Roberts et al., 2011a). This is particularly true for the Southern Central Mediterranean as shown by an increasing number of lake level, pollen and speleothem records depicting contrasting signals. Some paleorecords from Sicily show wetter conditions (Sadori and Narcisi, 2001; Magny et al., 2011b; Peyron et al., 2012b) while others indicate drier conditions during the Early Holocene (Tinner

5690

et al., 2009; Calò et al., 2012). Note that “Early Holocene” is defined here as the 11.7 to 6.5 kyr interval and “Mid- to Late Holocene” the interval from 6.5 kyr to present-day. On the southern rim of the Mediterranean Sea, peat sequences from Kroumirie (Ben Tiba and Reille, 1982; Ben Tiba, 1995; Stambouli-Essassi et al., 2007) give a synthetic view of the Holocene vegetation history in the humid mountainous NW Tunisia. More detailed sequences in this area and in the semi-arid plains are needed to get a complete description of the Holocene climatic changes in Northern Tunisia. In addition, delayed forest expansion appears to be recorded by few sites from the southernmost areas of the Mediterranean region suggesting that dry conditions may have persisted into the Holocene (Tzedakis, 2007). A clearer picture of the long-term climatic changes in the Southern Central Mediterranean is a crucial step towards a mechanistic understanding of the regional hydroclimatic changes in response to large-scale orbital climatic changes.

Deglacial vegetation changes in the Central Mediterranean are based on few pollen sequences mainly located in Central Italy and Adriatic Sea (Watts et al., 1996; Combourieu-Nebout et al., 1998; Magri, 1999; Magri and Sadori, 1999). These records clearly show the afforestation associated to the climatic improvement of the last deglaciation as well as the response of the Mediterranean forest to millennial-scale climatic changes. However, the low temporal resolution of available records (Ben Tiba and Reille, 1982; Brun, 1983) does not allow to evaluate the impact of the millennial-scale variability during the last deglaciation. Since the identification of abrupt changes in the North Atlantic during the Holocene (Bond et al., 1997), a number of publications have shown that such changes affected the Mediterranean Sea and borderlands (e.g. synthesis papers and references therein: Magny et al., 2007a; Fletcher and Zielhofer, 2011; Peyron et al., 2011; Roberts et al., 2011b; Sadori et al., 2011). The complexity of the Mediterranean climate as well as the interpretation of proxies and chronological uncertainties remains a major limitation to our understanding of abrupt climatic changes. Records from Central Mediterranean provide evidences of increased dryness during the Holocene cooling episodes which contrasts with the increase in humidity in

5691

the European mid-latitudes, above 40° N (Magny et al., 2011b). Recently, a change in nature and tempo of centennial-scale variability during the mid-Holocene has been identified in the SW Mediterranean (Fletcher et al., 2012). However, key questions remain about the impact of Holocene cooling events on the Mediterranean climate and ecosystems and the processes responsible for the propagation and modulation of the North Atlantic variability into the Mediterranean region.

In this paper, we present pollen data from a marine record retrieved from the Siculo-Tunisian strait providing a regional picture of climate and vegetation changes in the southernmost Central Mediterranean during the last deglaciation and the Holocene. Our record based on a direct land-sea correlation will bring new insight into regional long-term trend and millennial to centennial-scale climatic changes. We compare our results with other records from Central and Western Mediterranean to show a coherent pattern between records. Finally, we discuss the potential links between the Mediterranean climate evolution and the low and mid-latitude atmospheric circulation features in relation with insolation and ice sheet volume forcings.

2 Environmental setting and potential pollen source

Core MD04-2797CQ (36°57' N 11°40' E, 771 m water-depth) was retrieved from the Siculo-Tunisian Strait at ~ 50 km east off Cape Bon at the tip of the North-Eastern Tunisian coast and at ~ 110 km off Sicily (Fig. 1). The strait bottom morphology is characterized by a sill and two troughs separating the Sicilian and Tunisian continental shelf areas. The upper water column is occupied by the Modified Atlantic Water (MAW). This water mass is fed by the Atlantic Tunisian Current (ATC) which is the southern branch of the MAW flowing eastward along the Tunisian coast through the strait while the northern branch flows along the Sicilian coast (Fig. 1) (Béranger et al., 2004). The upper water mass above the core site is influenced by the ATC. At depth, waters originating from the Eastern Mediterranean flowing westward across the Siculo-Tunisian Strait are composed of Levantine Intermediate Water and transitional Eastern Mediterranean

5692

water (Astraldi et al., 2002). The surface ocean circulation is driven by the atmospheric circulation dominated by westerly to north-westerly winds (Pinardi et al., 2005).

The present-day climate in Northern Tunisia and Sicily is of Mediterranean-type, characterized by warm and dry summers and cool and wet winters. The orography and maritime influences modulate the climate in both regions. An west to east humidity gradient exists in Northern Tunisia between the humid to sub-humid northwestern mountains and the semi-arid northeastern lowlands while in Southern Tunisia arid conditions prevail (INRF, 1976). The Sicily climate ranges from semi-arid to sub-humid at the highest elevations of the island (Drago, 2005). The autumn-winter-spring seasonal precipitation mainly derives from moisture advection from the Atlantic Ocean. The Western Mediterranean Sea is another source of moisture for the surrounding landmasses through evaporation and atmospheric circulation. Summer dryness results from the subsidence development associated with the northward displacement of the Hadley cell (Lionello et al., 2006).

The vegetation distribution in the Central Mediterranean is controlled by temperature and precipitation gradients and seasonality. The potential vegetation cover of the Medjerda watershed is dominated by *Olea-lentiscus* maquis in the eastern semi-arid areas and by holm oak-aleppo pine forest in the southern upper semi-arid High Tell and Tunisian Dorsal. In the most North-Western Kroumirie region of the Medjerda watershed, cork oak with *Erica arborea* underbrush and high altitude zeen oak forest prevail due to humid and sub-humid conditions in the mountains. Cape Bon considered as the “backbone” of the Tunisian Dorsal is the closest landmass to our core site where moisture from maritime influence leads to semi-arid to sub-humid conditions. Vegetation of Cape Bon is dominated by *Oleo-lentiscus* maquis, kermes oak (*Quercus coccifera*) and Thuya (Gausson and Vernet, 1958; Posner, 1988; El Euch, 1995; Boussaid et al., 1999).

In addition, numerous permanent or temporary wetlands are located in Tunisia (Posner, 1988) mostly in the humid, sub-humid and upper semi-arid bioclimatic zones (INRF, 1976). Wetlands from Northern Tunisia such as Garaet el Ichkeul are

5693

characterized by the dominance of Cyperaceae in the marsh zone although recently replaced by Chenopodiaceae due to decreasing freshwater inflow as a consequence of dams and pipeline constructions (Ramdani et al., 2001). In temporary wetlands such as Sebket Ariana located in the semi-arid region, north-east of Tunis, vegetation is dominated by halophyte plants in particular Chenopodiaceae (Posner, 1988).

In Sicily, vegetation is distributed along humidity and temperature gradients resulting from the orography. *Oleo-ceratonion* maquis dominates the lowlands in the warmest and driest areas of the SE Sicily. At higher elevations, the potential vegetation has been assigned to mesophilous Mediterranean evergreen woods (*Quercetum ilicis*). Increased humidity in the submontane areas allows the development of deciduous woodlands (*Quercetalia pubescenti-petraeae*) while at highest altitudes of Sicily, *Geranio versicoloris-Fagion* association is favoured (Morinello, 1999).

A number of studies have shown that marine pollen signal provides an integrated picture of the vegetation from the nearby continents (Heusser and Balsam, 1977; Naughton et al., 2007). Studies on present-day pollen deposition in marine sediments show that pollen grains are fluvial and wind-transported from the adjacent continent and that agglomeration, flocculation and zooplankton feeding processes enable rapid sinking of pollen through the water column (Heusser and Balsam, 1977; Heusser, 1978; Chmura et al., 1999; Dupont and Wyputta, 2003; Mudie and McCarthy, 2006). Further, cores such as MD04-2797 CQ located near continental regions with well-developed hydrographic basins and prevailing offshore winds, mainly recruit pollen from rivers (Heusser, 1978; Turon, 1984; Dupont and Wyputta, 2003). Pollen grains belonging to the fine-particle fraction have a similar behaviour as fine sediments during the sedimentary processes (Muller, 1959; Chmura and Eisma, 1995). Medjerda and Miliane oueds are the two most important and permanent water streams in Tunisia discharging into the Gulf of Tunis and providing sediments to the shelf area (Fig. 1). The Medjerda oued is the main Tunisian stream flowing SW–NE between the Northern Tell and High Tell mountains, from Eastern Algeria to the Gulf of Tunis. Floods are recurrent in Northern Tunisia which receives most rainfall in winter and spring. Floods enhance material

5694

transport and contributed to the recent filling in of the Gulf of Tunis (Zahar et al., 2008). The riverine system of Sicily is not well developed and climate varies from semi-arid in the low land to sub-humid upland. Wind-driven sediment input to the shelf might be important in semi-arid regions. Given the dominant Westerly winds and surface ocean circulation, the bathymetry of the strait and large river systems of Northern Tunisia, the Southern Central Mediterranean and most probably also Northern Tunisia are likely to be main source of pollens to our core site. Therefore, even though long-range transport of pollen cannot be ruled out, we interpret the pollen signal of core MD04-2797CQ as reflecting vegetation changes in this region.

3 Material and methods

Sediment core MD04-2797CQ was subsampled for pollen analysis every 2 to 10 cm over the first 590 cm. The sample preparation technique followed the procedure described by de Vernal et al. (1996). Exotic pollen (*Lycopodium*) was added to each sample to determine pollen concentrations. After chemical and physical treatments (cold HCl, cold HF, cold KOH and sieving through 10 μ m nylon mesh screens), the residue was mounted unstained in glycerol. Pollen was counted using a light microscope at 400 and 1000 (oil immersion) magnifications. 100 to 160 pollen grains without *Pinus* and a minimum of 20 taxa were counted in each of sample. 10 out of 94 samples analysed were considered as sterile due to extremely low pollen concentration, in particular in the upper 80 cm of the core. Pollen percentages for each taxon are based on the main pollen sum excluding *Pinus* because of its over-representation in marine deposits (Heusser and Balsam, 1977; Turon, 1984). *Pinus* percentages were calculated from the main pollen sum including *Pinus*. Spores and aquatic pollen percentages were obtained from the total sum that includes all pollen grains plus aquatic plant and indeterminate pollen grains and Pteridophyta spores.

Quantitative climatic reconstruction based on the Modern Analogue Technique (MAT) was performed from pollen data. The MAT has been used to reconstruct here, annual,

5695

winter and summer temperatures and precipitation by considering the proportion of pollen assemblages (Guiot, 1990). Methods and results are presented and discussed in details in Peyron et al. (2012).

Direct land-sea correlation is obtained in comparing pollen to marine proxies from MD04-2797CQ sediments. Dinocyst and planktic foraminifera assemblages and derived SSTs, planktic foraminifera $\delta^{18}\text{O}$ and SST obtained from the C_{37} alkenones unsaturation index have been published earlier (Essallami et al., 2007; Rouis-Zargouni et al., 2010). In this paper, we compare our pollen data with the new high resolution alkenone-derived SST record (Sicre et al., 2012).

4 Results and interpretations

4.1 Chronological framework and pollen analysis results

Core MD04-2797CQ chronology was established from 13 radiocarbon dates among which 6 are from the upper 4 m of the core corresponding to the Holocene interval (Table 1). Except for the date at 0 cm, all radiocarbon dates have been replicated, thus increasing the reliability of the age-depth model. Averages of the ^{14}C date duplicates were corrected for a reservoir age of 400 yr for the Holocene, GS-1 and LGM intervals, of 560 yr for GI-1 (level 470 cm) and of 800 yr for Heinrich Stadial 1 (HS1) (level 510 cm), according to Siani et al. (2001). Corrected dates were calibrated using the calibration curve INTCAL 09 (Reimer et al., 2009). The age-depth model was obtained using linear interpolation between dated levels (Fig. 2). The upper 80 cm represents the last 5800 yr and is only chronologically constrained by two radiocarbon dates (5830 and 670 yr cal BP). A strong change in sedimentation rate occurred in between the two dated levels increasing age uncertainties for this interval. Here after, ka and kyr will be used for ages in cal BP.

According to the age-depth model (Fig. 2), the MD04-2797CQ pollen record encompasses the last 18 000 yr. Pollen analyses were performed at high temporal resolution

5696

generally considered as a cold tolerant-moisture demanding conifer. Colder conditions and moisture availability at mid to high altitudes during HS 1 and GS-1 could have favoured cedar expansion at the expense of deciduous oak in NW Africa, from the Moroccan Rif to the High Tell. The Mediterranean region presents a wide range of habitats due to highly variable topography and local climate conditions which enables refugia zones during the last glacial period (Médail and Diadema, 2009). This work supports the idea that tree populations maintained in the mountainous region of North Africa even during the most severe episodes of the last deglaciation.

During GI-1, SSTs increased, semi-desert plants and cedar reduced while some deciduous and evergreen oak woodlands and Mediterranean shrubs (*Olea*) developed (PZ-2; Fig. 4). This finding is indicative of relatively warmer and wetter climatic conditions than during HS 1 and GS-1. Despite the temperature rise, the semi-desert plants remained dominant and tree and shrubs expansion limited compared to terrestrial pollen records from Italy (Magri and Sadori, 1999; Allen et al., 2002) or Western Mediterranean Sea pollen records (Fletcher and Sánchez Goñi, 2008; Combourieu-Nebout et al., 2009). This result can be explained by lower water availability in the semi-arid areas of Southern Central Mediterranean. Terrigenous supply to the Siculo-Tunisian Strait remained characterized by a high content of aeolian-driven particles, such as illite and palygorskite, which also indicate prevailing dry conditions during GI-1 (Bout-Roumazeilles et al., 2012). Strikingly, our pollen data do not indicate a tree and shrub expansion maximum at the beginning of the GI-1 but rather slightly higher values at the end of the interstadial at ~ 13.4 kyr following a long HS1 to GI-1 transition. This feature contrasts with the sharp temperature rise and early GI-1 optimum in Greenland ice cores (NGRIP members, 2004) (Fig. 4) and has already been identified from speleothem records from Northern Tunisia, Southern France and Southern Germany (Genty et al., 2006) as well as from pollen records from the Alboran Sea and Southern Alps (Combourieu-Nebout et al., 2009; Finsinger et al., 2011). This North–South contrast may be linked to the North Atlantic oceanic and atmospheric circulations and sea ice cover (von Grafenstein et al., 1999; Genty et al., 2006). In the Southern Alps,

5699

the gradual expansion of *Quercus* have been attributed to increasing summer temperatures and longer growing season probably related to insolation (Finsinger et al., 2011). Stacked temperature records over the latitudinal bands 30–0° S and 0–30° N (Shakun et al., 2012) indicate lower temperature during GI-1 than during the Holocene and display the same peculiar increasing temperatures over the course of the interstadial. Since tropical ocean temperature and associated evaporation regulate the atmospheric moisture content feeding the North Atlantic storm track (Braconnot et al., 2007), stronger inland moisture transport toward the end of GI-1 may be associated with the tropical climate in relation with orbital forcing.

4.3 Holocene long-term vegetation and climate changes

4.3.1 Vegetation and climate in and off Tunisia

In core MD04-2797CQ, the GS-1–Holocene transition is marked by a sharp decrease in semi-desert plants at ~ 12.3 kyr while alkenone-SSTs and summer SSTs derived from planktonic foraminifera assemblages increased by 9 °C (Essallami et al., 2007) (PZ-4; Figs. 3 and 4). These vegetation and SST changes seem to precede by 600 yr the sharp rise in temperature recorded in Greenland (Rasmussen et al., 2006). One can attribute this time difference to age uncertainties of the records. However, this difference is quite large and may suggest earlier changes in temperature and precipitation in the Southern Mediterranean region compared to the northern latitudes. Open vegetation with Asteraceae, Poaceae and some semi-desert plants (mainly Chenopodiaceae and *Ephedra fragilis*-type) remained dominant until 10.1 kyr when the open oak forest expands. Although Mediterranean and temperate trees and shrubs slightly increased at 12.3 kyr, they remained restricted at some woodlands/scrubs despite high alkenone-SSTs (Fig. 4). This vegetation change indicates a shift to warmer conditions and increased humidity during the Early Holocene but moisture availability remained too weak before 10.1 kyr to enable a forest development. In the Southern Mediterranean, temperature influences forest composition but moisture availability is overall

5700

critical for forest development (Quezel, 2002). During this interval, decreasing semi-desert plants and increasing Cyperaceae and Ericaceae values suggest a gradual increase in moisture availability (Fig. 4). Therefore, the abrupt tree and shrub expansion observed at 10.1 kyr would likely result from crossing ecological thresholds during gradual climatic change. The gradual decrease in aridity before 10 kyr is supported by mineralogical and geochemical data found at core site MD04-2797CQ (Bout-Roumazeilles et al., 2012).

From 10.1 to 6.6 kyr, open deciduous oak forest with heath understorey and maquis developed along with a maximal sedge expansion (PZ-5). Mediterranean plants (mainly *Pistacia*, evergreen *Quercus* and *Olea*) remained limited over this interval and only start expanding after 6.6 kyr at the expense of heath maquis (PZ-6). At present-day, in the Central Mediterranean and in particular in Tunisia, Ericaceae are mainly represented by *Erica arborea*, *E. scoparia* and *E. multiflora* (Posner, 1988; Ojeda et al., 1998). Although *E. arborea* and *E. scoparia* have a wide range of distribution, they are related to humid climate and particularly atmospheric humid conditions (Jalut et al., 2000). This is well illustrated by the Ericaceae distribution in Northern Tunisia. Both species are widely distributed in the sub-humid zones of NW Tunisia and Cape Bon as understorey of *Quercus canariensis* and *Quercus suber* or as maquis of heather, while *E. multiflora* mainly grows in the *Olea-lentiscus* maquis or with aleppo pine and holm oak association in the semi-arid areas of Northern Tunisia (Posner, 1988). Therefore, the development of Ericaceae along with deciduous trees suggests that the most humid conditions of the Holocene occurred during this phase. Another line of evidence supporting this interpretation is the concurrent maximal development of sedges. Cyperaceae is a family comprising numerous species that can grow under a wide range of ecological conditions although most of them develop in humid area (Brun, 1985). Therefore, increased precipitation can potentially favour their contribution in the Asteraceae-Poaceae steppes. Sedge expansion can also reflect enhanced humid habitats, such as inland or coastal wetlands that are common features in Northern Tunisia. For instance, extensive sedge marshland surrounded Garaet el Ichkeul lake

5701

until dam constructions dramatically decreased water supply (Ramdani et al., 2001). Part of the strong sedge representation in our record might also result from coastal environment modifications related to post-glacial sea-level rise up to 6.6 kyr. The highest sedge percentages are also seen in MD95-2043 and ODP 976 sites during the same period, although lower than in our sequence. In these Alboran Sea records, sedges are associated with maximum forest development sustained by the most humid conditions of the Holocene (Fletcher and Sánchez Goñi, 2008; Combourieu-Nebout et al., 2009). Our pollen record therefore suggests that the 10–6.6 kyr interval is the wettest Holocene phase. However, in NE Tunisia water availability was never high enough, even during the most humid phase to sustain the regional development of a closed forest. Decreasing Ericaceae and Cyperaceae between 6.6 and 5 kyr (PZ-6) suggest a transition toward drier conditions. After 5 kyr, the expansion of Mediterranean plants such as evergreen *Quercus* and *Olea* suggests lower available summer moisture and warmer winter.

Precipitation estimates derived from our pollen record support the earlier interpretation of highest rainfall between 10 and 6.6 kyr, occurring mainly in winter, with mean annual precipitation of ~670 mm (Fig. 5, Peyron et al., 2012a). This period coincides with the highest pollen concentrations (Fig. 4) possibly due to both strong terrigenous inputs and increase biomass. After 5 kyr, while dryness increased on land, the SSTs show moderate warming, although different time-scale variability resulting from different temporal resolution may introduce bias. This warming seems however to be confirmed by increasing abundances of warm water dinocysts (Rouis-Zargouni et al., 2010).

Interpretation of our pollen record may appear contradictory regarding the gradual increase of temperate trees and shrubs all along the Holocene. Changes in post-glacial sea level may have influenced our marine pollen signal by modifying the dominant pollen transport agent from fluvial to aeolian. Mediterranean post-glacial sea-level reconstructions suggest that Cape Bon coastline and Medjerda river mouth were closer to our core site until 6.5 kyr when the Laurentide ice sheets (LIS) melting was completed (Lambeck et al., 2004). This could have led to enhance the representation of

5702

lowland relative to upland vegetation up to 6.5 kyr and that of the upland open forest afterwards. However, a shift to drier climate could have promoted aeolian transport. Although pollen transport changes may have modified the pollen signal, they do not alter the main vegetation changes. In particular, changes in the lowland shrub communities such as the replacement of heath maquis by *oleo-lentiscus* scrubs remain reliable to evaluate the main traits of the Holocene vegetation and climate history. Sedimentological analyses of core MD04-2797CQ agree with our interpretation of the pollen record. High sedimentation rates, decrease of aeolian-driven clay particles (palygorskite and illite), enhanced contribution of smectite likely reflecting fluvial supply and higher Zr/Al ratio indicate increased sediment supply to the shelf. Higher riverine discharge are likely sustained by more precipitation between 6 and 9.5 kyr (Bout-Roumazeilles et al., 2012).

4.3.2 Climatic or human-induced vegetation changes

The respective role of climatic change and anthropogenic pressure since the Neolithic has been a long standing debate in our understanding of the Mediterranean vegetation changes recorded by pollen sequences (De Beaulieu et al., 2005; Jalut et al., 2009). In Sicily, human impact appears to be detected in paleorecords since the Bronze Age and not since the Neolithic (Sadori and Giardini, 2007), and intense land use began with the Greek and Roman periods (Tinner et al., 2009). Human impact on the Tunisian environments became noticeable from the Phoenician and Carthaginian epochs (from XIIth to IIth century BC) with a strong agricultural development essentially based on cereal cultivations (Brun, 1983). However, our pollen record does not show this cereal signal. It is known that the dispersal of the large sized cereal pollen grains is poor (Vuorela, 1973), making difficult to get a reliable representation of cultivated Poaceae in marine pollen records. The Late Holocene *Olea* expansion is clearly detected in our record at around 3.2 kyr that may result from human activities during the Bronze Age. Olive tree expansion is also detected at that time in sequences from the Adriatic Sea, Italy and Sicily (Sadori and Narcisi, 2001; Di Rita and Magri, 2009;

5703

Combourieu-Nebout, 2012). Management of olive tree populations for fruit production started as early as the Bronze Age in the Western Mediterranean (Terral et al., 2004). However, as shown by our record, the expansion of *Olea* in Southern Italy is accompanied by a widespread increase of other Mediterranean trees and shrubs that may thus reveal a concomitant vegetation response to climate during the Bronze Age (Di Rita and Magri, 2009). Because of the low chronological constraints in our sequence between 600 and 5800 yr BP, *Olea* increase may also have occurred during Roman times. Expansion of olive trees at around 2000 yr cal BP during Roman occupation is detected in Southern Italy and Sicily (Terral et al., 2004; Di Rita and Magri, 2009). Roman developed an intensive olive tree cultivation for fruits and oil in the Mediterranean region (Terral et al., 2004). Therefore, human influence cannot be discarded since the Bronze Age but beforehand our pollen record primarily reflects climatic conditions.

4.3.3 Major climatic changes in the Southern Central Mediterranean

Land-sea correlation from core MD04-2797CQ indicates without chronological ambiguity that Holocene expansion of tree populations in Southern Central Mediterranean lagged temperature increase by two millennia. Delayed or slow afforestation due to moisture deficiency in the southernmost areas of the Mediterranean has been previously suggested from few terrestrial records (Tzedakis, 2007; Tinner et al., 2009). Dryness in Sicily before 10 kyr is supported by lake level and pollen data (Fig. 6g–i) (Sadori and Narcisi, 2001; Tinner et al., 2009; Magny et al., 2011b). Close examination of the Tunisian pollen records from Kroumirie (Stambouli-Essassi et al., 2007) and from Gulf of Gabes (Brun, 1979, 1985) shows that the deciduous oak forest expansion in NW Tunisia and of *Olea-Pistacia* scrubs in central eastern appears delayed by at least one millennium with respect to the beginning of the Holocene (Fig. 6c). Despite loose chronological framework and temporal resolution of the Tunisian pollen records, dry conditions of the Early Holocene derived from our sequence appear to be a robust feature in Northern Tunisia. In addition, the pollen and terrigenous records from core MD04-2797CQ parallel the $\delta^{13}\text{C}$ signal of the Tunisian speleothem record from

5704

La Mine cave (Fig. 6d) (Genty et al., 2006). Both records suggest a gradual climatic improvement and vegetation development in Southern Central Mediterranean immediately after the GS-1 cooling up to the climate optimum around 10 kyr. This is in contrast with Lake Preola record showing an abrupt rather than gradual change in lake level (Magny et al., 2011b). This contrasting feature could be season-related with a gradual change in winter precipitation from GS-1 to 10 kyr and an abrupt change in summer precipitation at 10 kyr.

Increased moisture availability during the Early Holocene, between 10 kyr and 6.6 kyr, is also consistent with available Tunisian and Sicilian records (Fig. 6). This accounts for example for the expansion of forest upland at Lake Pergusa and Majen M'Hida (Sadori and Narcisi, 2001; Stambouli-Essassi et al., 2007) and of *Pistacia* scrubs at the Western Sicilian coastal sites Lake Preola and Gorgo Basso (Tinner et al., 2009; Calò et al., 2012). After 6.6 kyr, our data and most of the Sicilian and Northern Tunisian records show similar long-term climate changes. Pollen and sedimentary $\delta^{18}\text{O}$ records and pollen-derived precipitation estimates from Lake Pergusa, lake level record from Lake Preola and stalagmite record from Grotta di Carburangeli suggest a transitional period of decreasing humidity or high amplitude variability in rainfall in Sicily between the wet 10 to 7 kyr interval and the pronounced drier period from 5–4 kyr to present-day (Frisia et al., 2006; Sadori et al., 2008; Magny et al., 2011a; Peyron et al., 2012b). Similarly, increasing dry conditions by 4–5 kyr is reflected by the replacement of the zeen oak forest by a cork oak forest with heath understorey at Dar Fatma in the humid NW Tunisia (Ben Tiba and Reille, 1982). Alluvial records from the Medjerda floodplain (Fig. 6e) showing increased fluvial dynamics with recurrent floods since 5 kyr also suggest prevailing arid conditions over the end of the Holocene (Zielhofer and Faust, 2008).

In contrast, the pollen records from the Sicilian coastal lakes Gorgo Basso, Lake Preola and Biviera di Gela also show a pronounced vegetation change at approx. 7–6.5 kyr. This change has been interpreted as a second step of humidity increase caused by a decrease in summer dryness (Noti et al., 2009; Tinner et al., 2009; Calò

5705

et al., 2012). In addition, in Western Sicily, no shift is recorded by 5 kyr, the evergreen forest declines only around 2.8–2.2 kyr likely because of intensified anthropic pressure on Mediterranean ecosystems (Tinner et al., 2009; Calò et al., 2012). However, the long-term climatic variations derived from vegetation changes at Biviera di Gela are not unequivocal and may as well be consistent with our results. Although maximum Mediterranean forest expansion between 7 and 5 kyr is interpreted as a response to moister climate at Biviera di Gela (Noti et al., 2009), the reduction of mesophilous trees over this interval might also be indicative of a rather contrasting drying trend. Furthermore, the partial shift from evergreen forest to *Pistacia*-dominated shrublands at this site became pronounced by 5 to 4 kyr, concurring with a Mid to Late Holocene aridification reinforced by increasing human impact (Noti et al., 2009). A number of hypotheses, although not entirely satisfactory, have been proposed to reconcile these coastal Sicilian pollen records with Lake Preola lake-level record and Lake Pergusa pollen sequence, such as spatial heterogeneity of landscape and climate (Tinner et al., 2009) or changes in seasonal and interannual variability of drought (Magny et al., 2011b). Possible mis-assignment of *Pistacia* pollen taxa to the evergreen species *P. lentiscus*, while pollen grains of the evergreen and deciduous species (*P. terebinthus* and *P. atlantica*) are indistinguishable is another plausible explanation (Sadori et al., 2011). Recently, Calò et al. (2012) also suggested that discrepancies between pollen and lake level records at the Sicilian coastal sites would come from seasonal biases of each proxy.

Rainfall seasonality in response to orbital forcing is an important issue. Lower or higher spring/summer dryness in Southern Central Mediterranean between 10 and 7 kyr than during the Late Holocene has been deduced from continental records (Magny et al., 2011a; Calò et al., 2012) while our marine record shows that summer precipitation (Psum) at regional scale remained low (50 to 100 mm) all over the Holocene, with no significant change (Fig. 5; Peyron et al., 2012a). However, even when Psum are estimated higher, they remained < 100 mm (Magny et al., 2011a). This denotes that dry summers remained a characteristic of the Holocene climate in that region. Subtle variations in Psum from semi-arid regions are particularly difficult to assess

5706

from pollen records since summer moisture availability for plants not only depend on summer precipitations but also on groundwater recharge during fall and winter that is critical to sustain vegetation during summer drought (Quezel, 2002; Fletcher et al., 2012), on continental evaporation related to summer temperatures and cloudiness, atmospheric humidity, storminess or soil absorption/retention. In contrast, Tunisian and Sicilian pollen and speleothem records (Frisia et al., 2006; Magny et al., 2011a) agrees with our observations (Fig. 5; Peyron et al., 2012a) underlying that the strongest winter and annual rainfall in the Southern Central Mediterranean occurred during the 10.1–6.6 kyr interval, followed by a decrease toward the Late Holocene.

4.3.4 Relationships with large-scale climatic changes

The Mediterranean climate results from complex interactions between low and mid-latitude atmospheric circulations and local features such as orography or land sea-contrast (Brayshaw et al., 2011). The winter Mediterranean hydroclimate depends on the Mediterranean storm activity (and associated precipitation) which is strongly influenced by the position and strength of the mid-latitude North Atlantic (NA) storm track (Brayshaw et al., 2010), while summer dryness is tightly linked to the summer extension and strength of the Hadley cell and associated subtropical subsidence (Lionello et al., 2006). Holocene changes of both NA storm track and tropical convection in response to insolation forcing are therefore critical factors to explain the long-term climatic evolution in Central Mediterranean and seasonal precipitation changes.

The drier conditions persisting into the Early Holocene up to 10.1 kyr are also detected in the SW and NE Mediterranean with a West to East gradient (Dormoy et al., 2009). The level of annual moisture availability always remained higher in the SW probably due to the stronger Atlantic influence, allowing Mediterranean plants to expand during the first millennia of the Holocene (Fletcher and Sánchez Goñi, 2008; Combourieu-Nebout et al., 2009). Such drier conditions are potentially due to the boreal summer insolation maximum of the Early Holocene (Tzedakis, 2007) resulting in deficiency in effective moisture availability due to strong evaporation (Renssen and

5707

Isarin, 2001). However, limited winter rainfall due to reduced influence of westerly flow-induced cyclones on the Mediterranean region is also probably involved (Kotthoff et al., 2008). The gradual shift from the GS-1 dry-cold situation to the Holocene warm-humid conditions might also be the result of the progressive reorganisation of the global atmospheric circulation associated with the retreat of Northern Hemisphere ice sheets. The LIS which disappeared around 6.8 kyr (Carlson et al., 2008) was large enough to impact on the atmospheric circulation (Montero-Serrano et al., 2010). The Fennoscandian Ice Sheet (FIS) which persisted up to 9 kyr could have influenced the response of the European hydroclimate to insolation forcing in maintaining the NA storm track in a relatively southern position (Magny et al., 2003). However, the NA storm track remained narrow enough not to bring high humidity in the Mediterranean.

The period comprised between 10.1 and 6.6 kyr encompasses to the Saproel 1 (Mercone et al., 2000). Increased precipitation in Southern Central Mediterranean is coeval with maximum monsoon activity as shown by tropical African paleohydrological records (Lézine et al., 2011). However, a number of evidences clearly show that the African monsoon never reached the Mediterranean region (Tzedakis, 2007). Our data thus confirm these earlier results since the estimated increase in annual precipitation remains limited on average to 700 mm yr^{-1} in our southern region and mainly due to a change in winter/fall precipitation (Fig. 5). In contrast, monsoonal intensification during summer could have played a predominant role on the Mediterranean summer dry season. Limited summer rainfall in Southern Central Mediterranean has been attributed to the development and persistence of subtropical anticyclones as a result of the reinforcement of the Hadley cell related to the African monsoon (Tinner et al., 2009; Gaetani et al., 2011). According to (Harrison et al., 1992), the summer northern position and higher strength of the North Atlantic Subtropical High in response to high summer insolation could have blocked the eastward convection and even generate an offshore flow in Southern Europe. This pattern would explain summer dryness over the whole Central Mediterranean region and the northern progression of summer dryness

as observed in pollen and lake level records (Magny et al., 2007a; Finsinger et al., 2010; Peyron et al., 2011).

The winter rainfall maximum from 10.1 to 6.6 kyr, observed in the Southern Central Mediterranean, appears to be a broad feature of the climate in the Western, Central and North-Eastern Mediterranean regions (Fletcher and Sánchez Goñi, 2008; Kotthoff et al., 2008; Combourieu-Nebout et al., 2009; Dormoy et al., 2009; Jalut et al., 2009; Pérez-Obiol et al., 2011; Peyron et al., 2011). The spatial extent and relative simultaneity of maximum winter suggests common mechanisms related to remote controlling factors. Simulations using global and regional models HadSM3 and HadRM3, suggest that increased winter precipitation during the Early Holocene would be linked to shifting NA storm track in response to orbital forcing (Brayshaw et al., 2010, 2011). The Early-Mid Holocene orbital configuration imposes a weaker winter latitudinal insolation gradient that would drive the northern tropical Hadley cell to be narrower and in a more southern position in winter. Consequently, the NA storm-track would weaken and shift to the south, thus enhancing Mediterranean cyclogenesis and winter precipitation (Brayshaw et al., 2010).

A contrasting hypothesis involving the Arctic Oscillation/North Atlantic situation (AO/NAO) has been proposed by Davis and Brewer (2009). According to these authors, increase in moisture availability in the Mediterranean region would be caused by lower evaporation due to cooler conditions (in contrast with our data from the Central Mediterranean). Cool and dry southern conditions would be explained by a more dominant positive AO (Davis and Brewer, 2009) and more northern NA storm track. Bonfils et al. (2004) showed that NAO does not support the winter temperatures and precipitation distribution during the mid-Holocene as shown by European paleodata. They suggest instead an insolation-induced southern entry of the NA storm track in the European mid-latitudes than during the Late Holocene.

The Early Holocene deglaciation of the LIS may have also contributed to increase winter precipitation in the Mediterranean region. The rapid retreat and increased melting of the LIS notably occurred during the wettest phase of the Mediterranean Holocene

5709

climate, between 10 and 6.8 kyr (Carlson et al., 2008). The above mentioned modelling experiments performed by Brayshaw et al. (2010) do not show a significant impact of the LIS on the NA storm track. Increased cyclogenesis is simulated only in the Western North Atlantic due to the increased temperature gradient resulting from the LIS cooling effect. Another modelling experiment with the fully coupled AOGCM ModelE-R has been performed to simulate the 9 kyr-climate with a more realistic LIS topography and appropriated freshwater routing (Carlson et al., 2008). ModelE-R outputs showed a stronger impact of the LIS on the ocean and atmospheric circulation over the North Atlantic region. Suppression of the Labrador Sea Water (LSW) formation, substantial reduction of the AMOC (by 15 %), and SST cooling in the NW Atlantic and temperature decrease in NE North America are simulated in agreement with paleodata (Carlson et al., 2008). If effective AMOC reduction is debatable (Renssen et al., 2005; Hoogakker et al., 2011), oceanic circulation reorganisation occurred when LIS retreat had completed. At ~6.5 kyr, deep water convection in the Nordic Seas reduced and LSW formation reactivated (Hoogakker et al., 2011). According to the ModelE-R experiment, the impact of the LIS melting on atmospheric and North Atlantic circulation induced a southern and stronger NA storm track providing slightly more winter precipitation to the mid and Southern Europe (Carlson et al., 2008, SI: Fig. 4).

We propose that during the Early Holocene, insolation and ice sheets, in particular the LIS decay, modulated winter precipitation in the Western to Central Mediterranean regions by altering oceanic and atmospheric circulation patterns of the North Atlantic, and strengthening the Mediterranean cyclogenesis. After the final demise of the LIS, increase in winter insolation and latitudinal insolation gradient probably contributed to the long-term decrease in winter precipitation and seasonality in the Western and Central Mediterranean by gradually shifting northward the NA storm track, which thus affected less frequently the Mediterranean region.

5710

4.4 Impact of the Holocene centennial-scale climatic changes on the temperature and hydroclimate in the Southern Central Mediterranean

The pollen record from core MD047-2497CQ displays a series of eight vegetation centennial-scale changes superimposed to the Holocene long-term evolution (11.5–11 kyr, 10.3–9.9 kyr, 9.5–9.1 kyr, 8.5–7.9 kyr, 7.3–6.7 kyr, 6.4–6 kyr, and ~5.5 kyr and 4.5 kyr).

During the Early Holocene dry phase, two centennial-scale dry and cold events at ~11.2 kyr and 10.1 kyr are clearly marked by increases of semi-desert plants with lower tree and shrubs percentages. During the first event, a large amplitude SST cooling is synchronous to vegetation changes, while no SST change is detected during the second one. Further, three cooling events of the SST record at ~9.3 kyr, 8.2 kyr, 7.0 kyr punctuated the wettest Holocene phase (10.1–6.6 kyr) with the last one at ~6.2 kyr, at the onset of the transition toward drier Mid- to Late Holocene. All these events are coeval with a change in herbaceous composition as shown by the strong reductions of Cyperaceae, most of the time replaced by Asteraceae (*Taraxacum*). Strikingly, they do not coincide with trees and shrubs reductions as we would have expected (Fig. 7). For instance, the Mediterranean forest percentages remain stable during the ~9.3 kyr and 6.2 kyr events while they decline at the end of the 7.0 kyr event. Different SST and Mediterranean forest variations have also been reported from the Alboran Sea record MD95-2043 (Fletcher et al., 2012) underlying the possible asynchronism between Mediterranean SST millennial-scale cooling and tree development. A seasonal bias towards summer for SST derived from alkenones and winter for the Mediterranean forest is proposed to explain this divergent pattern (Fletcher et al., 2012). However, alkenone-SSTs from core MD04-2797CQ during the interval 10.1–6.6 kyr do not parallel those found in core MD95-2043, suggesting that they might be controlled by different processes. We will see hereafter that centennial SST coolings from the Siculo-Tunisian Strait are associated to drier conditions, as revealed by the herbaceous layer changes.

5711

Centennial-scale variations in sedge percentages may have different causes. They could result from variations in the extent of coastal humid areas due to minor sea level changes. Such centennial-scale sea level changes during the Early Holocene may be produced by deglacial pulses, sea-water temperature changes through thermal water expansion or AMOC variations (Levermann et al., 2005; Lombard et al., 2005; Tornqvist and Hijma, 2012). Flooding can also favour sedge development in estuary zones and pollen export to the ocean (Bernhardt et al., 2012). However, high sedge percentages do not coincide with increased periods of flooding as recorded in the fluvial floodplains of Northern Tunisia (Zielhofer and Faust, 2008). Regarding the whole pollen assemblage, the most straightforward interpretation is restricted humid areas due to drier conditions. The concomitant *Taraxacum* increase may witness low moisture availability although relatively weak because semi-desert plants did not expand. In modern pollen assemblages, the highest *Taraxacum* percentages are found in dry environments of Greece and Morocco, where mean annual temperatures span from 12 and 17 °C and precipitation from 0 to 250 mm (Leroy, 1997). Therefore, a weak decrease in precipitation and slight cooling as shown by SSTs remaining above 16 °C would explain why oak populations were not affected by abrupt climate changes during that period. In contrast, over the last 6 kyr, our pollen record shows two strong abrupt forest reductions around 5.5 kyr and 4.5 kyr likely resulting from enhanced dryness, even though no SST decreases are recorded.

All the events detected before 6.6 kyr parallel the widely documented abrupt coolings punctuating the Early Holocene: the Preboreal Oscillation (PBO) at ~11.2 kyr (Björck, 1996), the Boreal Oscillation at ~10.1 kyr (Björck et al., 2001; Boch et al., 2009), the 9.3 kyr event (Rasmussen et al., 2007), the “8.2-event” (Alley et al., 1997) and a final event at ~7.4 kyr (Bond et al., 2001) which likely corresponds to the drier episode detected at ~7.0 kyr bearing in mind the chronological uncertainties (Fig. 7). Our pollen record confirms a decrease in precipitation in the Mediterranean region during the Early Holocene cooling episodes (Combourieu-Nebout et al., 2009; Dormoy et al., 2009; Pross et al., 2009; Fletcher et al., 2010). In addition, the centennial drier events in our

5712

record match the higher lake level episodes in the European mid-latitudes (Magny and Bégeot, 2004) (Fig. 7). This finding supports the Early Holocene contrasting pattern of the hydrological changes between the mid-latitudes and Southern Europe (Magny and Bégeot, 2004; Fletcher et al., 2012). These cooling events have been associated with
 5 deglacial freshwater outbursts in the North Atlantic from proglacial lakes such as Lake Agassiz (Teller et al., 2002), Lake Labrador-Ungava (Jansson and Kleman, 2004), Lake Superior (Yu et al., 2010) or Baltic ice Lake (Nesje et al., 2004), and with solar activity minima (Bond et al., 2001; Magny and Bégeot, 2004). Early Holocene meltwater pulses and solar forcing have been shown to alter the AMOC and modify the atmospheric
 10 circulation in the mid-latitudes but also in the tropics (Bond et al., 2001; Magny and Bégeot, 2004; Marchitto et al., 2010). A southern entry point of the Atlantic jet and associated cyclones in Europe and a stronger zonal flow due to enhanced temperature gradient between low and high latitudes, are suggested from the European hydrological reconstructions (Magny et al., 2003, 2007b; Magny and Bégeot, 2004).

15 Abrupt forest reductions around 5.5 kyr and 4.5 kyr may be attributed to major widespread cooling events dating to 5.6–5 kyr and ~4.2 kyr, respectively (Magny and Haas, 2004; Drysdale et al., 2006; Magny et al., 2009). This coincidence might be fortuitous because the age control and time resolution of our sequence is low over the last 5.8 kyr. Nevertheless, drier conditions in the Southern Mediterranean at ~5.5 and
 20 4.2 kyr are also suggested by lake-level decreases (Magny et al., 2007a, 2011b). An episode of arboreal biomass reduction at ~4.2 kyr is recorded in the Central Mediterranean. It has been associated with climate change despite the possible human impact at the beginning of the Bronze Age (Magri and Parra, 2002; Sadori et al., 2011). Both events at ~5.5 and 4.2 kyr are associated with higher lake levels at mid-latitudes in
 25 Europe and in Central Italy in response to North Atlantic cooling and decrease in solar activity (Magny, 2004; Magny et al., 2007a) (Fig. 7). However, hydrological changes at 4.2 and 5.5 kyr are complex (Magny et al., 2006, 2009, 2011b). For instance, the short-lived dry event in Central Italy probably coeval to the “4.2 kyr event” occurred during a cooler and wetter interval between 4.5 and 3.8 kyr (Magny et al., 2009). In contrast,

5713

forest reduction episodes in Western Mediterranean at ~5.4–4.5 kyr and 3.7–2.9 kyr do not match those events (Fig. 7). They have been associated to dry conditions in the mid-latitudes and enhanced wind strength in the northern latitudes (Fletcher et al.,
 5 2012). However, it is not clear if these forest reduction episodes actually correspond to warm or cold events in the North Atlantic because of a different timing and periodicity of the centennial scale variations recorded in the Alboran Sea pollen record and in the North Atlantic IRD record (Fletcher W. J., personal communication, 2012).

Therefore, given the limitations of the chronology of our core, forest reductions reconstructed from our record may be linked either (1) to higher lake level in the mid-
 10 European latitude and lower lake level at Lake Preola, or (2) to the forest reduction in the Western Mediterranean. Answer to this question is crucial to determine the factors controlling the centennial-scale variability during the Mid- to Late Holocene and their impact on the Mediterranean region. From 6 kyr onwards, dry episodes in the Southern Central Mediterranean could involve (1) wetter conditions above 40° N due to a southern and stronger Atlantic jet like during the Early Holocene cooling events (Magny et al.,
 15 2004) or (2) drier conditions in the Mediterranean and Mid European latitudes due to an intensification and northward shift of the NA storm track and subsequent limited penetration of storms in the Mediterranean region (Fletcher et al., 2012).

Apart from this, our record shows that changes on land are of weaker amplitude during the Early Holocene, in particular during Sapropel 1. Weaker changes in Mediterranean forest in the SW Mediterranean also characterize the Early Holocene as compared to the Mid- to Late Holocene as shown by the MD95-2043 pollen sequence (Fig. 7). The notable weak expression of the 8.2 event in the Central Mediterranean has already been emphasized from the Accesa or Preola sequences (Magny et al., 2011b;
 20 Peyron et al., 2011). This event is even missing in Moroccan and Tunisian alluvial sequences (Zielhofer et al., 2008). This finding emphasizes that impact of cooling events in the Early Holocene was relatively weak on the Western to Central Mediterranean hydroclimate while it appears to have been stronger during the Mid to Late Holocene. Baseline climate states, such as insolation and ice sheet volume, exert a strong control

5714

- Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichet, Th., Hewitt, C. D., Kageyama, M., Kitoh, A., Lâiné, A., Loutre, M.-F., Marti, O., Merkel, U., Ramstein, G., Valdes, P., Weber, S. L., Yu, Y., and Zhao, Y.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum – Part 1: experiments and large-scale features, *Clim. Past*, 3, 261–277, doi:10.5194/cp-3-261-2007, 2007.
- Brayshaw, D. J., Hoskins, B., and Black, E.: Some physical drivers of changes in the winter storm tracks over the North Atlantic and Mediterranean during the Holocene, *Philos. T. Roy. Soc. A*, 368, 5185–5223, 2010.
- Brayshaw, D. J., Rambeau, C. M. C., and Smith, S. J.: Changes in mediterranean climate during the holocene: insights from global and regional climate modelling, *Holocene*, 21, 15–31, 2011.
- Broecker, W. S.: Was the Younger Dryas triggered by a flood?, *Science*, 312, 1146–1148, 2006.
- Brun, A.: Recherches palynologiques sur les sédiments du Golfe de Gabès: résultats préliminaires, *Géologie méditerranéenne: la Mer Pélagienne*, Marseille, 1979,
- Brun, A.: Etude palynologique des sédiments marins Holocènes de 5000 BP à l'actuel dans le Golfe de Gabès (Mer Pélagienne), *Pollen Spores*, 25, 437–460, 1983.
- Brun, A.: La couverture steppique en Tunisie au Quaternaire supérieur, *CR Acad. Sci. II*, 14, 1085–1090, 1985.
- Cacho, I., Grimalt, J. O., Canals, M., Sbaiffi, L., Shackleton, N. J., Schönfeld, J., and Zahn, R.: Variability of the Western Mediterranean sea surface temperature during the last 25,000 years and its connection with the Northern Hemisphere climatic changes, *Paleoceanography*, 16, 40–52, 2001.
- Calò, C., Henne, P. D., Curry, B., Magny, M., Vescovi, E., La Mantia, T., Pasta, S., Vannièrè, B., and Tinner, W.: Spatio-temporal patterns of Holocene environmental change in Southern Sicily, *Palaeogeogr. Palaeocl.*, 323–325, 110–122, doi:10.1016/j.palaeo.2012.01.038, 2012.
- Carlson, A. E., Legrande, A. N., Oppo, D. W., Came, R. E., Schmidt, G. A., Anslow, F. S., Licciardi, J. M., and Obbink, E. A.: Rapid early Holocene deglaciation of the Laurentide ice sheet, *Nat. Geosci.*, 1, 620–624, 2008.
- Chmura, G. L. and Eisma, D.: A palynological study of surface and suspended sediments on a tidal flat: implications for pollen transport and deposition in coastal waters, *Mar. Geol.*, 128, 183–200, 1995.

5719

- Chmura, G. L., Smirnov, A., and Campbell, I. D.: Pollen transport through distributaries and depositional patterns in coastal waters, *Palaeogeogr. Palaeocl.*, 149, 257–270, 1999.
- Combourieu-Nebout, N., Peyron, O., Sadori, L., Dormoy, I., and Joannin, S.: Central Mediterranean vegetation and climate changes during the Holocene through pollen records around the Adriatic Sea, *Clim. Past Discuss.*, in preparation, 2012.
- Combourieu-Nebout, N., Paterne, M., Turon, J.-L., and Siani, G.: A high-resolution record of the Last Deglaciation in the Central Mediterranean sea: palaeovegetation and palaeohydrological evolution, *Quaternary Sci. Rev.*, 17, 303–332, 1998.
- Combourieu-Nebout, N., Peyron, O., Dormoy, I., Desprat, S., Beaudouin, C., Kotthoff, U., and Marret, F.: Rapid climatic variability in the west Mediterranean during the last 25 000 years from high resolution pollen data, *Clim. Past*, 5, 503–521, doi:10.5194/cp-5-503-2009, 2009.
- Correa-Metrio, A., Urrego, D. H., Cabrera, K., and Bush, M.: paleoMAS: paleoecological analysis. R package version 2.0, The Comprehensive R Archive Network, 2011.
- Davis, B. A. S. and Brewer, S.: Orbital forcing and role of the latitudinal insolation/temperature gradient, *Clim. Dynam.*, 32, 143–165, 2009.
- De Beaulieu, J. L., Miras, Y., Andrieu-Ponel, V., and Guiter, F.: Vegetation dynamics in North-Western Mediterranean regions: instability of the Mediterranean bioclimate, *Plant Biosyst.*, 139, 114–126, 2005.
- de Vernal, A., Henry, M., and Bilodeau, G.: Techniques de préparation et d'analyse en micropaléontologie, *Les cahiers du GEOTOP*, 3, 16–27, 1996.
- Di Rita, F. and Magri, D.: Holocene drought, deforestation and evergreen vegetation development in the Central Mediterranean: a 5500 years record from Lago Alimini Piccolo, Apulia, Southeast Italy, *Holocene*, 19, 295–306, 2009.
- Dormoy, I., Peyron, O., Combourieu-Nebout, N., Goring, S., Kotthoff, U., Magny, M., and Pross, J.: Terrestrial climate variability and seasonality changes in the Mediterranean region between 15 000 and 4000 years BP deduced from marine pollen records, *Clim. Past*, 5, 615–632, doi:10.5194/cp-5-615-2009, 2009.
- Drago, A.: Atlante climatologico della Sicilia – seconda edizione, *Riv. Ital. Agrometeorol.*, 2, 67–83, 2005.
- Drysdale, R., Zanchetta, G., Hellstrom, J., Maas, R., Fallick, A., Pickett, M., Cartwright, I., and Piccini, L.: Late Holocene drought responsible for the collapse of Old World civilizations is recorded in an Italian cave flowstone, *Geology*, 34, 101–104, 2006.

5720

- Dupont, L. and Wyputta, U.: Reconstructing pathways of aeolian pollen transport to the marine sediments along the coastline of SW Africa, *Quaternary Sci. Rev.*, 22, 157–174, 2003.
- El Euch, F.: Le sylvo pastoralisme en Tunisie, in: *Sylvo pastoral Systems. Environmental, Agricultural and Economic Sustainability, Cahiers Options Méditerranéennes, CIHEAM, Zaragoza*, 1995.
- 5 Essallami, L., Sicre, M. A., Kallel, N., Labeyrie, L., and Siani, G.: Hydrological changes in the Mediterranean Sea over the last 30,000 years, *Geochem. Geophys. Geosyst.*, 8, Q07002, doi:10.1029/2007gc001587, 2007.
- Finné, M., Holmgren, K., Sundqvist, H. S., Weiberg, E., and Lindblom, M.: Climate in the Eastern Mediterranean, and adjacent regions, during the past 6000 years – a review, *J. Archaeol. Sci.*, 38, 3153–3173, doi:10.1016/j.jas.2011.05.007, 2011.
- 10 Finsinger, W., Colombaroli, D., De Beaulieu, J. L., Valsecchi, V., Vannière, B., Vescovi, E., Chapron, E., Lotter, A. F., Magny, M., and Tinner, W.: Early to mid-Holocene climate change at Lago dell'Accesa (Central Italy): climate signal or anthropogenic bias?, *J. Quaternary Sci.*, 25, 1239–1247, 2010.
- 15 Finsinger, W., Lane, C. S., van Den Brand, G. J., Wagner-Cremer, F., Blockley, S. P. E., and Lotter, A. F.: The lateglacial *Quercus* expansion in the Southern European Alps: rapid vegetation response to a late Allerød climate warming?, *J. Quaternary Sci.*, 26, 694–702, 2011.
- Fletcher, W. J. and Sánchez Goñi, M. F.: Orbital- and sub-orbital-scale climate impacts on vegetation of the Western Mediterranean basin over the last 48,000 years, *Quaternary Res.*, 20 70, 451–464, 2008.
- Fletcher, W. J. and Zielhofer, C.: *Fragility of Western Mediterranean landscapes during Holocene Rapid Climate Changes*, Catena, 2011.
- Fletcher, W. J., Sanchez Goñi, M. F., Peyron, O., and Dormoy, I.: Abrupt climate changes of the last deglaciation detected in a Western Mediterranean forest record, *Clim. Past*, 6, 245–264, doi:10.5194/cp-6-245-2010, 2010.
- 25 Fletcher, W. J., Debret, M., and Sanchez Goñi, M. F.: Mid-Holocene emergence of a low-frequency millennial oscillation in western Mediterranean climate: Implications for past dynamics of the North Atlantic atmospheric westerlies, *The Holocene*, in press, doi:10.1177/0959683612460783, 2012.
- 30 Frisia, S., Borsato, A., Mangini, A., Spötl, C., Madonia, G., and Sauro, U.: Holocene climate variability in Sicily from a discontinuous stalagmite record and the Mesolithic to Neolithic transition, *Quaternary Res.*, 66, 388–400, 2006.

5721

- Gaetani, M., Pohl, B., Douville, H., and Fontaine, B.: West African Monsoon influence on the summer Euro-Atlantic circulation, *Geophys. Res. Lett.*, 38, L09705, doi:10.1029/2011GL047150, 2011.
- Gaussen, H. and Vernet, A.: *Carte Internationale du Tapis Végétal. Tunis-Sfax*, Gouvernement Tunisien, Institut Géographique National, Paris, 1958.
- 5 Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K., Chkir, N., Hellstrom, J., Wainer, K., and Bourges, F.: Timing and dynamics of the last deglaciation from European and North African $\delta^{13}\text{C}$ stalagmite profiles – comparison with Chinese and South Hemisphere stalagmites, *Quaternary Sci. Rev.*, 25, 2118–2142, doi:10.1016/j.quascirev.2006.01.030, 2006.
- 10 Giorgi, F.: Climate change hot-spots, *Geophys. Res. Lett.*, 33, L08707, doi:10.1029/2006GL025734, 2006.
- Guiot, J.: Methodology of the last climatic cycle reconstruction from pollen data, *Palaeogeogr. Palaeoclimatol.*, 80, 49–69, 1990.
- 15 Harrison, S. P., Prentice, I. C., and Bartlein, P. J.: Influence of insolation and glaciation on atmospheric circulation in the North Atlantic sector: implications of general circulation model experiments for the Late Quaternary climatology of Europe, *Quaternary Sci. Rev.*, 11, 283–299, 1992.
- Heusser, L.: Spores and pollen in the marine realm, in: *Introduction to Marine Micropaleontology*, edited by: Haq, B. U. and Boersma, A., Elsevier, New York, 327–339, 1978.
- 20 Heusser, L. E. and Balsam, W. L.: Pollen distribution in the NE Pacific ocean, *Quaternary Res.*, 7, 45–62, 1977.
- Hoogakker, B. A. A., Chapman, M. R., McCave, I. N., Hillaire-Marcel, C., Ellison, C. R. W., Hall, I. R., and Telford, R. J.: Dynamics of North Atlantic deep water masses during the Holocene, *Paleoceanography*, 26, PA4214, doi:10.1029/2011PA002155, 2011.
- 25 INRF: *Carte Bioclimatique de la Tunisie selon la classification d'Emberger. Etages et Variantes*, Institut National de Recherches Forestières, République Tunisienne, 1976.
- IPCC: *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Geneva, Switzerland, 104 pp., 2007.
- 30 Jalut, G., Esteban Amat, A., Bonnet, L., Gauquelin, T., and Fontugne, M.: Holocene climatic changes in the Western Mediterranean, from South-East France to South-East Spain, *Palaeogeogr. Palaeoclimatol.*, 160, 255–290, 2000.

5722

- Jalut, G., Dedoubat, J. J., Fontugne, M., and Otto, T.: Holocene circum-Mediterranean vegetation changes: climate forcing and human impact, *Quaternary Int.*, 200, 4–18, doi:10.1016/j.quaint.2008.03.012, 2009.
- Jansson, K. N. and Kleman, J.: Early Holocene glacial lake meltwater injections into the Labrador Sea and Ungava Bay, *Paleoceanography*, 19, PA1001, 1001–1012, 2004.
- Johnsen, S. J., Clausen, H. B., Dansgaard, W., Fuhrer, K., Gundestrup, N., Hammer, C. U., Iversen, P., Jouzel, J., Stauffer, B., and Steffensen, J. P.: Irregular glacial interstadials recorded in a new Greenland ice core, *Nature*, 359, 311–313, 1992.
- Juggins, S.: *Package "rioja" – Analysis of Quaternary Science Data*, The Comprehensive R Archive Network, 2009.
- Kotthoff, U., Pross, J., Müller, U. C., Peyron, O., Schmiedl, G., Schulz, H., and Bordon, A.: Climate dynamics in the borderlands of the Aegean Sea during formation of sapropel S1 deduced from a marine pollen record, *Quaternary Sci. Rev.*, 27, 832–845, doi:10.1016/j.quascirev.2007.12.001, 2008.
- Lambeck, K., Antonioli, F., Purcell, A., and Silenzi, S.: Sea-level change along the Italian coast for the past 10,000 years, *Quaternary Sci. Rev.*, 23, 1567–1598, doi:10.1016/j.quascirev.2004.02.009, 2004.
- Leroy, S. A. G.: Climatic and non-climatic lake-level changes inferred from a Plio-Pleistocene lacustrine complex of Catalonia (Spain): palynology of the tres pins sequences, *J. Paleolimnol.*, 17, 347–367, 1997.
- Levermann, A., Griesel, A., Hofmann, M., Montoya, M., and Rahmstorf, S.: Dynamic sea level changes following changes in the thermohaline circulation, *Clim. Dynam.*, 24, 347–354, doi:10.1007/s00382-004-0505-y, 2005.
- Lézine, A. M., Hély, C., Grenier, C., Braconnot, P., and Krinner, G.: Sahara and Sahel vulnerability to climate changes, lessons from Holocene hydrological data, *Quaternary Sci. Rev.*, 30, 3001–3012, 2011.
- Lombard, A., Cazenave, A., DoMinh, K., Cabanes, C., and Nerem, R. S.: Thermosteric sea level rise for the past 50 years; comparison with tide gauges and inference on water mass contribution, *Global Planet. Change*, 48, 303–312, doi:10.1016/j.gloplacha.2005.02.007, 2005.
- Lundqvist, J. and Saarnisto, M.: Summary of project IGCP-253, *Quaternary Int.*, 28, 9–18, 1995.
- MacAyeal, D. R.: Binge/purge oscillations of the Laurentide ice sheet as a cause of the North Atlantic's Heinrich events, *Paleoceanography*, 8, 775–785, 1993.

5723

- Magny, M.: Holocene climate variability as reflected by mid-European lake-level fluctuations and its probable impact on prehistoric human settlements, *Quaternary Int.*, 113, 65–79, 2004.
- Magny, M. and Bégeot, C.: Hydrological changes in the European midlatitudes associated with freshwater outbursts from Lake Agassiz during the Younger Dryas event and the early Holocene, *Quaternary Res.*, 61, 181–192, 2004.
- Magny, M. and Haas, J. N.: A major widespread climatic change around 5300 cal yr BP at the time of the Alpine Iceman, *J. Quaternary Sci.*, 19, 423–430, 2004.
- Magny, M., Bégeot, C., Guiot, J., and Peyron, O.: Contrasting patterns of hydrological changes in Europe in response to Holocene climate cooling phases, *Quaternary Sci. Rev.*, 22, 1589–1596, 2003.
- Magny, M., Leuzinger, U., Bortenschlager, S., and Haas, J. N.: Tripartite climate reversal in Central Europe 5600–5300 years ago, *Quaternary Res.*, 65, 3–19, 2006.
- Magny, M., de Beaulieu, J. L., Drescher-Schneider, R., Vannièrè, B., Walter-Simonnet, A. V., Miras, Y., Millet, L., Bossuet, G., Peyron, O., Brugiapaglia, E., and Leroux, A.: Holocene climate changes in the Central Mediterranean as recorded by lake-level fluctuations at Lake Accesa (Tuscany, Italy), *Quaternary Sci. Rev.*, 26, 1736–1758, 2007a.
- Magny, M., Vannièrè, B., de Beaulieu, J. L., Bégeot, C., Heiri, O., Millet, L., Peyron, O., and Walter-Simonnet, A. V.: Early-Holocene climatic oscillations recorded by lake-level fluctuations in West-Central Europe and in Central Italy, *Quaternary Sci. Rev.*, 26, 1951–1964, 2007b.
- Magny, M., Vannièrè, B., Zanchetta, G., Fouache, E., Touchais, G., Petrika, L., Coussot, C., Walter-Simonnet, A. V., and Arnaud, F.: Possible complexity of the climatic event around 4300–3800 cal BP in the Central and Western Mediterranean, *Holocene*, 19, 823–833, 2009.
- Magny, M., Peyron, O., Sadori, L., Ortu, E., Zanchetta, G., Vannièrè, B., and Tinner, W.: Contrasting patterns of precipitation seasonality during the Holocene in the South- and North-Central Mediterranean, *J. Quaternary Sci.*, 27, 290–296, 2011a.
- Magny, M., Vannièrè, B., Calo, C., Millet, L., Leroux, A., Peyron, O., Zanchetta, G., La Mantia, T., and Tinner, W.: Holocene hydrological changes in South-Western Mediterranean as recorded by lake-level fluctuations at Lago Preola, a coastal lake in Southern Sicily, Italy, *Quaternary Sci. Rev.*, 30, 2459–2475, doi:10.1016/j.quascirev.2011.05.018, 2011b.
- Magri, D.: Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (Central Italy), *Rev. Palaeobot. Palyno.*, 106, 171–208, 1999.

5724

- Magri, D. and Parra, I.: Late Quaternary Western Mediterranean pollen records and African winds, *Earth Planet. Sc. Lett.*, 200, 401–408, 2002.
- Magri, D. and Sadori, L.: Late Pleistocene and Holocene pollen stratigraphy at Lago di Vico, Central Italy, *Veg. Hist. Archaeobot.*, 8, 247–260, 1999.
- 5 Marchitto, T. M., Muscheler, R., Ortiz, J. D., Carriquiry, J. D., and Van Geen, A.: Dynamical response of the tropical pacific ocean to solar forcing during the early holocene, *Science*, 330, 1378–1381, 2010.
- McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D., and Brown-Leger, S.: Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes, *Nature*, 428, 834–837, 2004.
- 10 Médail, F. and Diadema, K.: Glacial refugia influence plant diversity patterns in the Mediterranean Basin, *J. Biogeogr.*, 36, 1333–1345, 2009.
- Mercone, D., Thomson, J., Croudace, I. W., Siani, G., Paterne, M., and Troelstra, S.: Duration of S1, the most recent sapropel in the Eastern Mediterranean Sea, as indicated by accelerator mass spectrometry radiocarbon and geochemical evidence, *Paleoceanography*, 15, 336–347, 2000.
- 15 Montero-Serrano, J. C., Bout-Roumazielles, V., Sionneau, T., Tribovillard, N., Bory, A., Flower, B. P., Riboulleau, A., Martinez, P., and Billy, I.: Changes in precipitation regimes over North America during the Holocene as recorded by mineralogy and geochemistry of Gulf of Mexico sediments, *Global Planet. Change*, 74, 132–143, 2010.
- 20 Mudie, P. J. and McCarthy, F. M. G.: Marine palynology: potentials for onshore-offshore correlation of Pleistocene–Holocene records, *T. Roy. Soc. S. Afr.*, 61, 139–157, 2006.
- Muller, J.: Palynology of recent Orinoco delta and shelf sediments, *Micropaleontology*, 5, 1–32, 1959.
- 25 Naughton, F., Sanchez Goni, M. F., Desprat, S., Turon, J.-L., Duprat, J., Malaize, B., Joli, C., Cortijo, E., Drago, T., and Freitas, M. C.: Present-day and past (last 25,000 years) marine pollen signal off Western Iberia, *Mar. Micropaleontol.*, 62, 91–114, 2007.
- Nesje, A., Dahl, S. O., and Bakke, J.: Were abrupt Lateglacial and early-Holocene climatic changes in Northwest Europe linked to freshwater outbursts to the North Atlantic and Arctic Oceans?, *Holocene*, 14, 299–310, 2004.
- 30 NGRIP members: High-resolution record of Northern Hemisphere climate extending into the last interglacial period, *Nature*, 431, 147–151, 2004.

5725

- Noti, R., van Leeuwen, J. F. N., Colombaroli, D., Vescovi, E., Pasta, S., la Mantia, T., and Tinner, W.: Mid- and late-holocene vegetation and fire history at Biviere di Gela, a coastal lake in Southern Sicily, Italy, *Veg. Hist. Archaeobot.* 18, 371–387, 2009.
- Ojeda, F., Arroyo, J., and Marañón, T.: The phytogeography of European and Mediterranean heath species (Ericoideae, Ericaceae): a quantitative analysis, *J. Biogeogr.*, 25, 165–178, 1998.
- 5 Pérez-Obiol, R., Jalut, G., Julià, R., Pèlachs, A., Iriarte, M. J., Otto, T., and Hernández-Beloqui, B.: Mid-holocene vegetation and climatic history of the iberian peninsula, *Holocene*, 21, 75–93, 2011.
- 10 Peterson, L. C., Haug, G. H., Hughen, K. A., and Rohl, U.: Rapid changes in the hydrologic cycle of the tropical Atlantic during the last glacial, *Science*, 290, 1947–1951, 2000.
- Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., de Beaulieu, J. L., Drescher-Schneider, R., Vanniére, B., and Magny, M.: Holocene seasonality changes in the central mediterranean region reconstructed from the pollen sequences of Lake Accessa (Italy) and Tenaghi Philippon (Greece), *Holocene*, 21, 131–146, 2011.
- 15 Peyron, O., Combourieu-Nebout, N., Magny, M., Goring, S., Joannin, S., Dormoy, I., Brayshaw, D., de Beaulieu, J.-L., Brugiapaglia, E., Desprat, S., Kouli, K., Kotthoff, U., Pross, J., and Sadori, L.: Holocene climate changes in Mediterranean area: a model-data comparison, *Clim. Past Discuss.*, in preparation, 2012a.
- 20 Peyron, O., Magny, M., Gorin, S., Joannin, S., de Beaulieu, J.-L., Brugiapaglia, E., Sadori, L., Garfi, G., Kouli, K., and Combourieu-Nebout, N.: Contrasting patterns of climatic changes during the Holocene in Central Mediterranean area (Italy) reconstructed from pollen data, *Clim. Past Discuss.*, in preparation, 2012b.
- Pinardi, N., Zavatarelli, M., Arneri, E., Crise, A., and Ravaioli, M.: The Physical sedimentary and ecological structure and variability of shelf areas in the Mediterranean Sea, in: *The Global Coastal Ocean – Interdisciplinary Regional Studies and Syntheses*, edited by: Robinson, A. R. and Brink, K. H., Harvard University Press, Cambridge, MA and London, 1245–1331, 2005.
- 25 Posner, S. D.: Biological diversity and tropical forests in Tunisia, Agency for International Development, 206 pp., 1988.
- 30 Pross, J., Kotthoff, U., Müller, U. C., Peyron, O., Dormoy, I., Schmiedl, G., Kalaitzidis, S., and Smith, A. M.: Massive perturbation in terrestrial ecosystems of the Eastern Mediterranean region associated with the 8.2 kyr BP climatic event, *Geology*, 37, 887–890, 2009.

5726

- Quezel, P.: Réflexions sur l'évolution de la flore et de la végétation au Maghreb méditerranéen, Ibis Press, 2002.
- Ramdani, M., Flower, R. J., Elkhiaiti, N., Kraïem, M. M., Fathi, A. A., Birks, H. H., and Patrick, S. T.: North African wetland lakes: characterization of nine sites included in the CAS-SARINA Project, *Aquat. Ecol.*, 35, 281–302, 2001.
- 5 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M.-L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *J. Geophys. Res.*, 111, D06102, doi:10.1029/2005JD006079, 2006.
- 10 Rasmussen, S. O., Vinther, B. M., Clausen, H. B., and Andersen, K. K.: Early Holocene climate oscillations recorded in three Greenland ice cores, *Quaternary Sci. Rev.*, 26, 1907–1914, 2007.
- Rasmussen, S. O., Seierstad, I. K., Andersen, K. K., Bigler, M., Dahl-Jensen, D., and Johnsen, S. J.: Synchronization of the NGRIP, GRIP, and GISP2 ice cores across MIS 2 and palaeoclimatic implications, *Quaternary Sci. Rev.*, 27, 18–28, 2008.
- Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Burr, G., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., McCormac, F. G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J., Turney, C. S. M., van der Plicht, J., and Weyhenmeyer, C.: IntCal09 and Marine09 radiocarbon age calibration curves, 0–50 000 yr cal BP, *Radiocarbon*, 51, 1111–1150, 2009.
- Renssen, H. and Isarin, R. F. B.: The two major warming phases of the last deglaciation at ~14.7 and ~11.5 kyr cal BP in Europe: climate reconstructions and AGCM experiments, *Global Planet. Change*, 30, 117–153, 2001.
- 25 Renssen, H., Goosse, H., and Fichefet, T.: Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene, *Geophys. Res. Lett.*, 32, 1–4, 2005.
- Roberts, N., Brayshaw, D., Kuzucuoglu, C., Perez, R., and Sadori, L.: The mid-Holocene climatic transition in the Mediterranean: causes and consequences, *Holocene*, 21, 3–13, 2011a.

5727

- Roberts, N., Eastwood, W. J., Kuzucuoglu, C., Fiorentino, G., and Caracuta, V.: Climatic, vegetation and cultural change in the eastern mediterranean during the mid-holocene environmental transition, *Holocene*, 21, 147–162, 2011b.
- 5 Rouis-Zargouni, I., Turon, J.-L., Londeix, L., Essallami, L., Kallel, N., and Sicre, M.-A.: Environmental and climatic changes in the Central Mediterranean Sea (Siculo-Tunisian Strait) during the last 30 ka based on dinoflagellate cyst and planktonic foraminifera assemblages, *Palaeogeogr. Palaeoclimatol.*, 285, 17–29, 2010.
- Sadori, L. and Giardini, M.: Charcoal analysis, a method to study vegetation and climate of the Holocene: The case of Lago di Pergusa (Sicily, Italy), *Geobios*, 40, 173–180, 2007.
- 10 Sadori, L. and Narcisi, B.: The Postglacial record of environmental history from Lago di Pergusa, Sicily, *Holocene*, 11, 655–670, 2001.
- Sadori, L., Zanchetta, G., and Giardini, M.: Last Glacial to Holocene palaeoenvironmental evolution at Lago di Pergusa (Sicily, Southern Italy) as inferred by pollen, microcharcoal, and stable isotopes, *Quaternary Int.*, 181, 4–14, 2008.
- 15 Sadori, L., Jahns, S., and Peyron, O.: Mid-Holocene vegetation history of the Central Mediterranean, *Holocene*, 21, 117–129, 2011.
- Sanchez Goñi, M. F. and Harrison, S. P.: Millennial-scale climate variability and vegetation changes during the Last Glacial: concepts and terminology, *Quaternary Sci. Rev.*, 29, 2823–2827, doi:10.1016/j.quascirev.2009.11.014, 2010.
- 20 Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B., Schmitner, A., and Bard, E.: Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation, *Nature*, 484, 49–54, doi:10.1038/nature10915, 2012.
- Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., and Haddad, G.: Mediterranean Sea surface radiocarbon reservoir age changes since the Last Glacial Maximum, *Science*, 294, 1917–1920, doi:10.1126/science.1063649, 2001.
- 25 Sicre, M. A., Siani, G., Genty, D., Kallel, N., and Essallami, L.: North-South SST evolution across the Central Mediterranean basin during the last deglacial, *Clim. Past Discuss.*, in preparation, 2012.
- Stambouli-Essassi, S., Roche, E., and Bouzid, S.: Evolution of vegetation and climatic changes in North-Western Tunisia during the last 40 millennia, *Geo. Eco. Trop.*, 31, 171–214, 2007.
- 30 Teller, J. T., Leverington, D. W., and Mann, J. D.: Freshwater outbursts to the oceans from glacial Lake Agassiz and their role in climate change during the last deglaciation, *Quaternary Sci. Rev.*, 21, 879–887, 2002.

5728




Fig. 1. Map of Central Mediterranean with location of the studied core MD04-2797CQ and the pollen sequences mentioned in the text from the Southern Central Mediterranean. DF: Dar Fatma (Ben Tiba and Reille, 1982), MH: Majen Ben M'Hida (Stambouli-Essassi et al., 2007), GB: Gorgo Basso (Tinner et al., 2009), LP: Lake Pergusa (Sadori and Narcisi, 2001), BG: Biviera di Gela (Noti et al., 2009).

5733




Fig. 2. Age-depth model for core MD04-2797CQ.

5734




Fig. 3. Percentages of major pollen taxa versus depth, Core MD04-2797CQ, Siculo-Tunisian Strait. Boundaries of pollen zones are based on visual inspection of data and application of clustering analysis (see text for details).

5735




Fig. 4. Caption on next page.

5736

Fig. 4. Vegetation and climatic changes on land over the last 18 000 yr. From the bottom to the top: tree and shrubs (grey) and total (white) pollen concentrations, pollen percentages of selected pollen taxa and ecological groups, SST derived from alkenones (Sicre et al., 2012) from core MD04-2797CQ, $\delta^{18}\text{O}$ from NGRIP ice core on GGC05 time-scale (converted to BP) (Rasmussen et al., 2006; Vinther et al., 2006). Long-term Holocene vegetation evolution has been detected by applying a smoothing cubic spline method to selected pollen taxa data. Pollen zones (bottom) and stratigraphical framework (top) are indicated. Note that we will name the Younger Dryas as Greenland Stadial 1 (GS-1) following the INITIMATE recommendations (Lowe et al., 2008) but we will keep using HS 1 in the text (as defined by Sánchez Goñi and Harrison, 2010) because the isotopic changes during GS-2 are difficult to reconcile from one Greenland ice core to another (Rasmussen et al., 2008) and the correlation between GS-2a and HS 1 has to be confirmed.

5737




Fig. 5. Box-Whisker plots of estimated annual, summer and winter temperatures (TANN, MTWA, MTCO) and precipitation (Pann, Psum, Pwin) derived from the pollen data of core MD04-2797CQ (Peyron et al., 2012a).

5738




Fig. 7. Comparison of the centennial-scale variability recorded in core MD04-2797CQ with changes detected in the Western Mediterranean, middle latitudes of Europe and North Atlantic. From left to right: Mediterranean forest pollen percentages from core MD95-2043 (Fletcher and Sánchez Goñi, 2008): 3-points moving average (bold green line) with forest reduction episodes indicated by the grey bars as in Fletcher et al. (2012); MD04-2797CQ pollen percentages and alkenone-SSTs (bold lines: cubic smoothing splines with a degree of freedom adjusted to highlight the centennial-scale changes) with abrupt vegetation and SST changes as described in the text; Mid-European lake level (LL) changes (Magny, 2004); Stacked Hematite Stained Grains (HSG) from North Atlantic records (Bond et al., 2001).