



HAL
open science

Molards as an analogue for ejecta-ice interactions on Mars

Costanza Morino, Susan J. Conway, Coralie Peignaux, Antoine Lucas, Kristian Svennevig, Frances Butcher, Gioachino Roberti, Meven Philippe, Jake Collins-May

► **To cite this version:**

Costanza Morino, Susan J. Conway, Coralie Peignaux, Antoine Lucas, Kristian Svennevig, et al.. Molards as an analogue for ejecta-ice interactions on Mars. EGU General Assembly 2021, Apr 2021, Online, Austria. 10.5194/egusphere-egu21-9628 . hal-03280950v2

HAL Id: hal-03280950

<https://hal.science/hal-03280950v2>

Submitted on 3 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



This is a repository copy of *Molards as an analogue for ejecta-ice interactions on Mars*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/174163/>

Version: Published Version

Proceedings Paper:

Morino, C., Conway, S., Peignaux, C. et al. (6 more authors) (2021) Molards as an analogue for ejecta-ice interactions on Mars. In: EGU General Assembly 2021 Abstracts. vEGU21: Engage • Discover • Inspire, 19-30 Apr 2021, Virtual conference. European Geosciences Union (EGU) .

<https://doi.org/10.5194/egusphere-egu21-9628>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

EGU21-9628

<https://doi.org/10.5194/egusphere-egu21-9628>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Molards as an analogue for ejecta-ice interactions on Mars

Costanza Morino¹, Susan Conway², Coralie Peignaux², Antoine Lucas³, Kristian Svennevig⁴, Frances Butcher⁵, Gioachino Roberti⁶, Meven Philippe², and Jake Collins-May⁷

¹CNRS UMR 5204 Laboratoire Environnements Dynamiques et Territoires De La Montagne, Le Bourget-du-Lac, France (costanza.morino@gmail.com)

²CNRS UMR6112 Laboratoire de Planétologie et Géodynamique, Nantes Cedex 3, France

³Université de Paris, Institut de physique du globe de Paris, CNRS, F-75005, Paris, France

⁴Geological Survey of Denmark and Greenland, Copenhagen, Denmark

⁵Department of Geography, The University of Sheffield, Sheffield, UK

⁶Minerva Intelligence Inc., Vancouver, British Columbia, Canada

⁷The School of Geography, Politics and Sociology, Newcastle University, Newcastle, UK

The 125-km-diameter Hale impact crater is located in the southern hemisphere of Mars and has been dated to 1 Ga (Early to Middle Amazonian; Jones et al., 2011). It is thought to have penetrated the martian cryosphere, because it hosts landforms indicating volatile mobilisation post-impact: its ejecta are lobate and bear channels, and the interior is pervasively pitted and hosts alluvial fans (Collins-May et al. 2020; El-Maarry et al., 2013; Jones et al., 2011; Tornabene et al., 2012). Here, we test the hypothesis that conical mounds found within the ejecta are “molards” by comparing them to terrestrial analogues. Molards are conical mounds of debris resulting from the degradation of blocks of ice-rich material which have been mobilised by a landslide and are found in periglacial environments (Morino et al., 2019).

Our study area (240x180 km) is in the South-East part of the Hale impact crater ejecta (36°–39°S, 36°–31°W). We analyse the spatial and topographic distribution of the conical mounds using orbital images from 25 cm/pixel to 15 m/pixel and measure their height, width and slope using 1 m/pixel elevation data. We then compare them to conical mounds on the deposits of the 2010 Mount Meager debris avalanche, Canada (Roberti et al. 2017) and of the 2000 Paatuut landslide in western Greenland (Dahl-Jensen et al. 2004).

The conical mounds of the Hale impact crater are located at the distal boundary of the thickest part of the ejecta blanket, which reflects the spatial distribution of mounds along the distal parts of the terminal lobe of the Mount Meager debris avalanche. Furthermore, mounds in the Hale impact crater have comparable shapes and flank slopes to molards in the Mount Meager and Paatuut case studies, but are one order of magnitude bigger. This size difference is consistent with the flow-depth that transported the blocks also being one order of magnitude bigger than on Earth.

We infer that conical mounds near the Hale impact crater are a result of fragmented blocks of ice-

cemented regolith produced by the impact and transported by the ejecta flows, and finally degraded into cones of debris (molards) by the loss of interstitial ice. Our interpretation supports the prevailing hypothesis that the Hale impact event penetrated the martian cryosphere and further provides important constraints on the rheology of martian ejecta deposits that can be tested by future studies and in other locations on Mars.

We acknowledge financial support for the PERMOLARDS project from French National Research Agency (ANR-19-CE01-0010).