



HAL
open science

Deer exclusion unveils abiotic filtering in forest understory plant assemblages

Simon Chollet, Christophe Baltzinger, Morgane Maillard, Jean-Louis Martin

► **To cite this version:**

Simon Chollet, Christophe Baltzinger, Morgane Maillard, Jean-Louis Martin. Deer exclusion unveils abiotic filtering in forest understory plant assemblages. *Annals of Botany*, 2021, 128 (3), pp.371-381. 10.1093/aob/mcab079 . hal-03281826

HAL Id: hal-03281826

<https://hal.science/hal-03281826v1>

Submitted on 29 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Deer exclusion unveils abiotic filtering in forest understory plant
assemblages**

Simon Chollet^{1*}, Christophe Baltzinger², Morgane Maillard³, Jean-Louis Martin³

¹ *University of Rennes, CNRS, ECOBIO [Ecosystèmes, biodiversité, évolution] – UMR 6553, Rennes, France ;* ² *INRAE Val de Loire, Research Unit Forest Ecosystems, Domaine des Barres, 45290, Nogent-sur-Vernisson, France ;* ³ *CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France*

* *For correspondence. E-mail simon.chollet@univ-rennes1.fr*

Accepted Manuscript

- **Background and Aims** The role of deer (family Cervidae) in ecosystem functioning has traditionally been neglected by forest ecologists due to the animal's scarcity in most parts of the northern hemisphere. However, the dramatic rebound in deer populations throughout the 20th century has brought deer browsing to the forefront of forest ecological questioning. Today there is ample evidence that deer affect tree regeneration, understory plant and animal diversity and even litter decomposition. However, the mechanisms underlying the effects of deer on forest ecosystems remain unclear. Among others, the relative role of abiotic factors versus biotic interactions (e.g. herbivory) in shaping plant assemblages remains largely unknown.
- **Methods** We used a large-scale experiment with exclosures distributed along abiotic gradients to understand the role of black-tailed deer (*Odocoileus hemionus sitchensis*) on forest understory on the Haida Gwaii archipelago (western Canada), a unique context where most of the key ecological effects of deer presence had already been intensively studied.
- **Key Results** Our results demonstrate that 20 years of deer exclusion resulted in a clear increase in vascular plant richness, diversity and cover, and caused a decline in bryophyte cover. Exclusion also unveiled abiotic (i.e. soil water availability and fertility) filtering of plant assemblages that would otherwise have been masked by the impact of abundant deer populations. However, deer exclusion did not lead to an increase in beta diversity, probably because some remnant species had a competitive edge to regrow after decades of over-browsing.

- **Conclusions** We demonstrated that long-term herbivory by deer can be a dominant factor structuring understory plant communities that overwhelms abiotic factors. However, while exclosures prove useful to assess overall effects of large herbivores, the results from our studies at broader scales on the archipelago, suggest that enclosure experiments should be used cautiously when inferring the mechanisms at work.

Key words: abiotic factors, assembly rules, biotic interactions, bryophyte, legacy effects, overbrowsing, Sitka black-tailed deer, vascular plant.

Accepted Manuscript

INTRODUCTION

Growing evidence indicates that deer influence biodiversity and functioning of temperate and boreal forest ecosystems much more than previously thought. During the second part of the 20th century, most cervid (family Cervidae, hereafter deer) populations increased dramatically in many forest ecosystems of the northern hemisphere. The reasons for this recovery from severe historical population depletions are multiple. They vary among regions and include hunting regulations, restocking programs, changes in forestry and eradication of predator (Alverson *et al.* 1988; Crête and Daigle 1999; Warren 2011; Martin *et al.* 2020). As a result, deer populations became increasingly perceived as “over-abundant” in many countries (Côté *et al.* 2004; Takatsuki 2009; Carpio *et al.* 2021). Evidence indicates that abundant deer cause tree regeneration failure (e.g. Gill 1992), modify plant communities (e.g. Wiegmann and Waller, 2006), change litter decomposition (e.g. Chollet *et al.* 2021) and trigger cascading effects on animals that depend on the understory (e.g. Chollet and Martin 2013).

The studies of deer effects on forest ecosystems used a variety of methods including enclosure and enclosure experiments, experimental reduction of deer abundance through culls, comparison of sites varying in deer abundance (in a synchronic or diachronic way), the study of sites inaccessible to herbivores (rocky refuges, uncolonized islands) or simulated herbivory (i.e. clipping) (Waller 2014). However, most of the results came from enclosure experiments, with a large number of studies located in North America, Japan and Europe (Bernes *et al.* 2018). The limits of enclosure studies have been pointed out repeatedly. They include the fact that they only allow the comparison of one level of browsing with a complete lack of browsing. Furthermore, their results usually only focus on the effect of defoliation by deer and neglect their effect through trampling, soil litter disturbance/scraping, urine and faeces deposition or seed dispersal (Hester *et al.* 2000; Bergström and Edenius 2003; Waller

2014). In addition, as Mladenoff and Stearns (1993) pointed out, extrapolating the results of enclosure experiments from the stand scale to regional trends could be meaningless, because the small size of enclosures prevents information on factors (e.g. edaphic factors) other than those related to deer effect on the vegetation. In fact light availability, water stress and nutrient deficiencies are primary factors limiting plant long-term persistence and, consequently, influence community composition in most forest understories (Muller 2014; Neufeld and Young 2014). In a modelling exercise Mladenoff and Stearns (1993) suggested that the lack of hemlock (*Tsuga canadensis*) regeneration was not solely controlled by deer, and that a population reduction alone would not ensure successful regeneration. This study, among others, raised the question of the hierarchy between biotic interactions and abiotic factors in shaping plant assemblages.

In grass dominated ecosystems, large herbivores are predicted to reduce the dominance of the more competitive species and therefore to favour spatial plant coexistence (Olf and Ritchie 1998; Adler *et al.* 2001). One of the mechanisms behind this heterogeneity is that the reduction of competition due to herbivory allows a stronger control of community composition by abiotic filters. However, in boreal and temperate forests, where understories are dominated by slow growing perennial forbs and shrubs, the prediction of intense deer browsing effect on spatial vegetation heterogeneity may be quite different (Roberts and Gilliam 2014; Waller 2014). In fact, in these ecosystems, one could predict that high deer abundance would lead to a decrease in species richness and diversity, including beta diversity (i.e. measure of community heterogeneity), as understory plants may have more difficulty to withstand severe defoliation, and/or may be more palatable (Bardgett and Wardle 2003; Nuttle *et al.* 2014; Forbes *et al.* 2019). In such ecosystems, the relative role of biotic interactions and abiotic factors may be reversed, with stronger biotic than abiotic filtering. Despite such expected differences among ecosystem types, the effect of deer on vegetation

spatial heterogeneity in forest ecosystems have been largely neglected, compared to grassland or savannah ecosystems (Gao and Carmel 2020). When effects of large herbivores in these systems were actually tested, some results suggested that strong abiotic gradients could actually affect the response of the forest vegetation to herbivory (e.g. Piazza *et al.*, 2016; Randall and Walters, 2011; Rhodes *et al.*, 2017 but see Suzuki *et al.* 2013). For example, Piazza *et al.* (2016) demonstrated that the impact of livestock on the composition and diversity of understorey vegetation increased along a precipitation gradient in *Nothofagus* forests of Argentina, with the wetter sites being more affected than the drier. In another study evaluating impacts of elk (*Cervus canadensis*), mule deer (*Odocoileus hemionus*) and cattle (*Bos taurus*) on aspen (*Populus tremuloides*) regeneration, Rhodes *et al.* (2017) highlighted a combined effect of herbivore density and abiotic factors (precipitation, elevation and stand composition).

In the present study we explore the effects of Sitka black-tailed deer (*Odocoileus hemionus sitchensis*) on the understory vegetation along abiotic gradients with the intention to evaluate the spatial heterogeneity of deer impact and vegetation response before and after deer exclusion. Specifically, our goal was to understand the relative role of abiotic and biotic filtering in the assemblage of understory plant communities. In order to achieve this, we designed a large scale enclosure experiment in a context where the ecological effects of deer presence had been intensively studied, the archipelago of Haida Gwaii in western Canada. Studies there documented a dramatic effect of deer on tree regeneration (Martin and Baltzinger 2002; Stroh *et al.* 2008), on understory and shoreline plant communities composition (Stockton *et al.* 2005; Martin *et al.* 2010; Chollet, Baltzinger, Ostermann, *et al.* 2013), on bryophyte communities (Chollet, Baltzinger, Le Saout, *et al.* 2013). They also showed a cascade of effects on insects and birds (Allombert, Stockton, *et al.* 2005; Allombert, Gaston, *et al.* 2005; Chollet *et al.* 2015), and the possibility of plant and bird

communities recovery after a drastic reduction in deer density (Chollet *et al.* 2016). We used 20 exclosures placed in pairs in ten sites throughout the main island of the archipelago, Graham, where deer were initially introduced in the late 19th century resulting in the longest deer presence on Haida Gwaii (Fig. 1). Our intention here is to assess plant community recovery after twenty years of deer exclusion. Based on our previous results throughout the archipelago, we expected that after 20 years of recovery the vegetation should be more abundant, more diverse and more heterogeneous (higher beta diversity) within the deer exclosures than outside of them. In addition, a comparison of these results to those we obtained with different approaches on Haida Gwaii (Martin and Baltzinger 2002; Stockton *et al.* 2005; Chollet, Baltzinger, Ostermann, *et al.* 2013; Chollet *et al.* 2016) should give us a unique opportunity to critically assess the relative contribution of results from exclosures to the overall question of deer effects on vegetation.

MATERIALS AND METHODS

Study area and experimental design

The study took place on Haida Gwaii (formerly known as Queen Charlotte Islands, British Columbia, Canada), the largest and most isolated archipelago on the west coast of Canada. The islands (more than 150, including small islets) are largely formed of volcanic and sedimentary rock, together with intrusions of granitic rock (Sutherland Brown 1968) and are mainly covered with temperate rain forest classified in the Coastal Western Hemlock Zone (Banner *et al.* 2014). In these forests, the dominating canopy trees are Western redcedar (*Thuja plicata*), Western hemlock (*Tsuga heterophylla*) and locally Sitka spruce (*Picea sitchensis*). The shrub layer is usually dominated by regeneration of Western hemlock and shrubs like Salal (*Gaultheria shallon*), red huckleberry (*Vaccinium parvifolium*), blueberries (*Vaccinium alaskaense* and *V. ovalifolium*) and false azalea (*Menziesia ferruginea*) (Banner

et al. 2014). Typical herb layer species are Deer fern (*Blechnum spicant*), false lily-of-the-valley (*Maianthemum dilatatum*), twayblades (*Listera ovata* and *L. cordata*), single delight (*Moneses uniflora*) and five-leaved bramble (*Rubus pedatus*) (Banner *et al.* 2014). Overall, the flora is similar to the one found in South-eastern Alaska and Northern British Columbia, but strongly impoverished due to insularity (i.e. 655 vascular plant species versus 2300, Banner *et al.* 2014). The climate is cool and oceanic, with a strong precipitation gradient. Mean annual precipitation ranges from 1000 mm on the east side of the islands to 7000 mm on their west side (Banner *et al.* 2014). Sitka black-tailed deer were introduced in the north of the largest island (Graham) in the late 19th century. In absence of their predators the population rapidly increased to reach densities higher than those found on the mainland (Martin and Baltzinger 2002). The average population density on the islands was estimated to range from 13 deer to 37 deer per km² (Engelstoft 2001; Stockton *et al.* 2005).

We used our enclosure experiment to characterize understory vegetation change following 20 years of deer exclusion on Graham. The experiment consisted of 20 square enclosures (5 x 5 m, 25m²) set up in 1997 in 10 sites (two enclosures per site, Fig.1). We placed all enclosures in remnant old-growth forest patches dominated by Western redcedar. Each enclosure was paired with an unfenced area considered as a control plot. We chose site locations that reflected at best the forest island diversity in terms of precipitation, light availability and soil fertility. In each enclosure and associated unfenced control plot we recorded the % cover of all vascular plant species present in the 0 to 4m layer, as well as overall bryophyte cover (all species grouped) in the initial year, 1997 and in the final year, 2017. During the entire period, we checked and maintained fences regularly. Only one enclosure was destroyed in the winter 2016/2017 by tree-fall, leaving 19 intact enclosures for the 2017 survey. For each plot (inside and outside enclosure) we sampled soil in five randomly selected location in 2017 to evaluate soil fertility. From the mixed subsample we

measured total carbon and nitrogen content (g / g dry soil) of freeze-dried soil using an Elementar Vario El Cube Analyzer (Elementar, Langenselbold, Germany).

Statistical analyses

Because skilled but different observers sampled the vegetation during the initial (1997) and the final (2017) monitoring sessions, we decided to minimize biases due to observer effect in species detection and in plant cover estimation by comparing characteristics of plant communities inside and outside enclosure only within a given year (i.e. in 1997 and in 2017).

For each year we compared alpha species richness, total species cover, species diversity (measured by Shannon index), beta diversity, indicator species (using IndVal, Dufrene and Legendre 1997) and species composition (with Non-Metric Multidimensional Scaling, NMDS) for vascular plant species inside and outside deer enclosures. In addition, we compared total bryophyte cover inside and outside enclosures. Concerning beta diversity, as demonstrated by Baselga (2010), measures of dissimilarity are the resultant of two components, species turnover and nestedness, that could be partitioned. In the former the differences come from a replacement of species between assemblages but in the latter the poorest site is only a subset of the richest site (absence of replacement of species). In both case species assemblages are obviously dissimilar but not for the same reason. In addition the dissimilarity could be defined based on a qualitative (i.e. species presence) or quantitative (i.e. species abundance) dimension, which could also be partitioned (Baselga 2013). We studied beta diversity by partitioning the two components of a quantitative dissimilarity index (i.e. Bray-Curtis).

In order to study changes of assembly rules resulting from deer exclusion we evaluated the affinity of plant assemblages for soil moisture and nutrient regimes inside and outside enclosures using null models (Gotelli and Graves 1996). Species assemblage affinity

was calculated as the mean value of each species affinity index present in the plot. We extracted semi-quantitative estimations of species affinity for moisture and nutrient from Klinkenberg (2020). These are equivalent to Ellenberg's indicator values in the European flora. We compared the mean species assemblage value of each plot (inside and outside) to 4,999 random assemblages drawn from the species pool (combination of all observed species). For a species richness S_i in plot i , our null model randomly sampled S_i species, each with a probability equal to its frequency in the species pool. For each plot we compared the observed value to the distribution generated by the 4,999 runs of the null model by calculating Standardized Effect Size (SES) as:

$$SES = \frac{(I_{obs} - \mu I_{null})}{\sigma I_{null}}$$

with I_{obs} the observed metric, μI_{null} and σI_{null} respectively the mean and standard deviation of the plot null distribution.

We compared treatment (inside vs. outside enclosure) and site effects on vascular species richness, Shannon index, beta diversity, total species cover and bryophyte cover with two-way ANOVAs. We studied the effect of precipitation (i.e. longitude as a proxy of the gradient, increasing from east to west), light availability (i.e. canopy cover as a proxy) and soil fertility (i.e. total soil C : N as a proxy) on vascular species richness, diversity, cover, bryophyte cover and SES with multiple regression models (including stepwise selection of explanatory variables) in 2017. To complement NMDS we tested the difference in species composition with Permutational Multivariate Analysis of Variance (PERMANOVA).

All statistical analyzes were performed in the R environment (R Core Team 2020) using the *vegan* (Oksanen *et al.* 2019), *betapart* (Baselga *et al.* 2018) and *Indicspecies* (Cáceres *et al.* 2020) packages.

RESULTS

At the outset of the study vascular species richness, Shannon diversity index, vascular plant cover and bryophytes cover were similar inside and outside exclosures, but varied among sites (Fig. 2 Supplementary data Table S1). After 20 years of deer exclusion the three metrics assessing vascular species were significantly higher inside than outside. Differences among sites remained significant (Fig. 2, Supplementary data Table S1). At the end of the period bryophyte cover had significantly decreased inside exclosures with an absence of differences among sites (Fig. 2, Supplementary data Table S1).

Multiple regression models indicated that at the end of the study variation in soil fertility (C : N) explained the among site variation in species richness, vascular plant cover, diversity (Shannon) and to a lesser extent moss cover variations in the exclosures (Fig. 3, Supplementary data Table S2). Conversely, outside of the exclosures, only species richness could be explained by C : N and this to only a limited extent (Fig. 3, Supplementary data Table S2). For vascular plants, the higher the soil fertility the higher the three studied community metrics, whereas the opposite pattern was observed for moss cover.

Our null models indicate non-random filtering for mean soil moisture and nutrient affinity index of plant assemblages inside exclosures (Fig. 4). Assemblages located on the west coast (i.e. higher precipitation record) were composed of species with a higher need for soil humidity (Fig. 4A) and species assemblages of more fertile sites were composed of species with a higher affinity for soil nutrient (Fig. 4B). These non-random trends in community assembly we observed inside the exclosures were lacking or weaker in plots placed outside of the exclosures (Fig. 4).

We found that only eight of the 44 species were significantly associated with a treatment, and it was only for the inside of exclosures in 2017 (Table 1). Among these species

were two trees (*Chamaecyparis nootkatensis*, *Thuja plicata*), three shrubs (*Gaultheria shallon*, *Menziesia ferruginea*, *Vaccinium parvifolium*), two forbs (*Cornus canadensis*, *Rubus pedatus*) and one fern (*Blechnum spicant*).

In 1997 beta diversity (as measured with Bray-Curtis index) did not differ between the inside and outside of exclosures ($F= 0.44$, $p>0.05$, Fig. 5A), and there was no significant difference among sites ($F= 1.04$, $p>0.05$). Twenty years later beta diversity was significantly lower for plots sampled inside the exclosures than outside ($F= 20.37$, $p<0.001$, Fig. 5B) but again without any difference among sites ($F= 0.28$, $p>0.05$). The higher dissimilarity observed in 2017 outside of the exclosures mainly reflected a difference in species abundance, as some individuals were lost from one site to the other but without substitution [significant difference in nestedness (Bray-Curtis gra component, Fig 5F) but not in turnover (Bray-Curtis bal component, Fig 5D)].

The Non Metric Multidimensional Scaling shows that the composition of plant communities were totally overlapping at the beginning of the study (in 1997) but that this was no more the case at the end (in 2017, Fig. 6, Supplementary data Table S3). This result is confirmed by the PERMANOVA where we found no significant differences between inside and outside plots in 1997 ($R^2=0.007$, $F=0.56$, $p>0.05$) but a difference in 2017 ($R^2=0.20$, $F=11.7$, $p<0.001$). The site effect was important in 1997 ($R^2=0.63$, $F=5.07$, $p<0.001$) as well as in 2017 ($R^2=0.35$, $F=2.23$, $p<0.001$). In 2017, the heterogeneity among plots was lower inside than outside exclosures, indicating increased similarity after 20 years of deer exclusion (Fig. 6).

DISCUSSION

Deer as dominant biotic filter

Plant community assembly is classically considered as the result of the interplay of dispersal, and of abiotic and biotic hierarchical filtering (Keddy 1992; Lortie *et al.* 2004). The key result in this study is that abundant deer presence can act as a strong biotic filter overwhelming abiotic factors in the control of the vegetation. While the presence of deer resulted in low vegetation cover outside of the exclosures, the increases in species richness, diversity and plant cover inside the exclosures depended not only on deer exclusion for 20 years, but were also strongly dependent on soil fertility. The most fertile sites (i.e. low C : N) benefited the more from deer exclusion, whereas sites with low fertility showed weaker response to deer exclusion. This lack of change in plots with the lower amount of nutrient is key because it shows that recovery from deer overabundance will only be possible, at least in the medium term, on relatively richer sites. Outside of the exclosures, only species richness was influenced by soil fertility, and to a lesser extent to what we observed inside of the exclosures. These differences in plant community response to soil fertility when deer are abundant suggest that they act as a strong biotic filter in the assembly of plant communities, overwhelming abiotic filters. This interpretation is confirmed by our results testing the change in plant assemblage affinity for soil moisture and nutrients (Fig. 4). In fact, we found that in absence of deer, assemblages were non-random, with wet and fertile sites harbouring communities characterized by species with higher affinity for wet and rich soils respectively. However, in presence of deer (i.e. outside of exclosures) these trends fade away.

Deer differentially affect community characteristics

More generally, our results confirm that abundant deer can be a dominant factor controlling understorey vegetation in Haida Gwaii forests. Twenty years of deer exclusion resulted in a clear increase of vascular plant cover and diversity. This increase of understorey vegetation was concomitant with a reduction in bryophyte cover, probably resulting from increased competition for light and for growing space because of the developing vascular plant cover (Chollet *et al.* 2013; Chollet *et al.* 2016). Interestingly we documented only a relatively small increase in alpha species richness, suggesting that deer, although strongly controlling plant abundance, have a comparatively smaller effect on species presence. However, this interpretation could be challenged by previous results we obtained at broader scales on the archipelago (Stockton *et al.* 2005; Chollet, Baltzinger, Ostermann, *et al.* 2013). These showed a positive effect of deer absence on plant species richness on islands never colonized by deer (Stockton *et al.* 2005), on refuges inaccessible to deer (Chollet, Baltzinger, Ostermann, *et al.* 2013) or at the landscape scale after a severe experimental reduction in deer density (Chollet *et al.* 2016). We hypothesize that the relatively small increase in species richness observed in our small-size and isolated exclosures after twenty years of deer exclusion may reflect a deficit in colonization. Woodland specialist-plant species, in particular, are indeed known to be poor colonists (Bossuyt *et al.* 1999; Rooney and Waller 2003; Verheyen and Hermy 2004; Waller 2014). Nevertheless, the changes we observed in the vegetation are consistent with the results obtained throughout the northern hemisphere where deer population greatly increased during the 20th century (Rooney and Waller 2003; Côté *et al.* 2004; Takatsuki 2009; Habeck and Schultz 2015; Ramirez *et al.* 2018) or in places where deer were introduced (Tanentzap *et al.* 2009; Nuñez *et al.* 2010).

Deer effects are selective

The species which benefited the more from deer exclusion (as revealed by the IndVal analyses) were mainly shrubs and tree species, all considered as typical from the Coastal Western Hemlock Zone and highly palatable to deer (Pojar *et al.* 1980). One of this species, Western redcedar, is of particular interest as it represents an essential element of Haida culture (Turner 2004). The negative effect of deer on Yellow cedar (*Chamaecyparis nootkatensis*) is also important because this species has been recently found to be also negatively affected by global warming on Haida Gwaii (Comeau *et al.* 2019) as well as on the mainland (Hennon *et al.* 2012), questioning the future of this important species on the islands. The three shrub species which benefited from deer exclusion (*Gaultheria shallon*, *Menziesia ferruginea*, *Vaccinium parvifolium*) are also widely used by first nations as food or medicine and are considered as diagnostic elements of the Coastal Western Hemlock Zone forest type (Turner 2004; Banner *et al.* 2014).

Deer exclusion leads to local biotic homogenization

Twenty years after deer exclusion, vegetation heterogeneity was lower inside than outside of the exclosures (Fig. 5 and 6), suggesting a local biotic homogenization (Olden and Rooney 2006). This pattern of dominance of some species inside exclosures could reflect the fact that large herbivores, by reducing the dominance of the more competitive species (browsing effect), or by creating microhabitat (trampling effect), or by moving seeds (dispersal effect), create heterogeneity that could lead to increased beta diversity. This has often been documented in grass dominated ecosystems (Olf and Ritchie 1998). An alternative hypothesis is that the dominance of competitive species observed inside the exclosures was partly an artifact linked to past overbrowsing (i.e. legacy effect). Under this hypothesis, species dominance could be understood as the result of the better recovery of a

minority of species that had a higher ability to withstand the prolonged presence of deer. This would confer them an initial advantage once protected, and limit the increase of poor colonizing species initially absent or more suppressed. Although we cannot fully affirm which of these non-exclusive hypotheses prevails, our results on beta diversity partitioning support the second hypothesis. In presence of deer, the higher beta diversity is explained by heterogeneity in abundance, and not by species replacement (Fig. 5). In other words, species present outside of the exclosures were also present inside, but with less variation in their abundance among sites. This suggests that herbivory, trampling or dispersal are not creating heterogeneity, but that, after deer exclusion, the abundance of the species already present increases and converges to similar cover values among sites. This understory homogenization inside the exclosures could be illustrated by the strong increase in cover of competitive and relatively fast growing species like *Gaultheria shallon* (Fraser *et al.* 1995; Bennett *et al.* 2011). That dominance of a small number of competitive species may disappear in the long term and at larger scale. This is actually what we observed in the already mentioned landscape-scale experiment of deer density reduction which, ultimately, triggered an increase in both vegetation and bird community heterogeneity at the broader scale (Chollet *et al.* 2016).

CONCLUSION

Our study demonstrates that deer, when overabundant, could overwhelm abiotic factors controlling plant communities (here soil water availability and fertility). In addition, deer reduced alpha species richness, diversity and the cover of understory plants typical from the studied costal rainforest. Despite consistency with other studies on the archipelago using different methodologies, this study based on a twenty-year exclosure experiment provided results not entirely in agreement with those observed at more broader and more realistic

spatial and/or temporal scales. The less dramatic increase in species richness or the decrease in beta diversity we observed after (only) 20 years of excluding deer from small sections of forests (25 m²) could be patterns sensitive to study length or scale. We actually highlighted the key importance of addressing questions at the appropriate temporal scale in a research conducted in the same study system on deer effects on soil physicochemical properties and bacterial communities (Maillard *et al.* 2021). This called caution when using even 20 year-long enclosure experiments to assess the interactions between deer and vegetation or other ecosystem components and properties. We could overcome the difficulty by combining the enclosure experiment with other approaches involving islands with deer and islands never colonized by deer, or the vegetation in natural refuges protected from deer browsing, or the experimental reduction of deer density. But this was only made possible, on one hand, by the rare and unplanned experimental situation created by the introduction of deer to Haida Gwaii and, on the other hand, by the also rare ability to thoroughly study this system for decades.

Accepted Manuscript

ACKNOWLEDGEMENT

This work was made possible through the bringing together of data, results and contributions that span well over two decades. We thank Lisa Blant, Isabel Buttler, Julie Gresser, Carolyn Hesseltine, Rob Kelly, Soizic Le Saout, Annie McKenzie, Juliane Schörghuber and Bruno Vila for their help in the field, Barb Rowsell and Janet Gray for the continuous monitoring of the enclosures over the years and Noémie Stroh for exploratory analyses. We also thank two anonymous reviewers for their useful comments on a previous version of the manuscript. Author contributions: SC, CB and JLM conceived the study, all authors collected the data, SC led the analysis and wrote the first draft, all authors commented the manuscript.

FUNDING

This work was supported by the British Columbia South Moresby Forest Replacement Account [contract # SMFRA99DQC-002, SMFRA Project 24.2], the Forest Renewal British Columbia [PA97335-BRE] and by the Agence Nationale de la Recherche [ANR BLAN 171801 Bambi], the French Ministry of Foreign affairs and the France Canada Research Fund (Grant to Morgane Maillard). *Cemagref* funded CB field missions in 2005 and 2009. The Research Group on Introduced Species (RGIS), The Canadian Wildlife Service (CWS), the Laskeek Bay Conservation Society (LBCS), Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and Haida Heritage Site, and the BC Ministry of Forests, Lands and Natural Resource Operation (Haida Gwaii Natural Resource District) provided logistical support.

LITERATURE CITED

- Adler P, Raff D, Lauenroth W. 2001.** The Effect of Grazing on the Spatial Heterogeneity of Vegetation. *Oecologia* **128**: 465–479.
- Allombert S, Gaston AJ, Martin J-L. 2005.** A natural experiment on the impact of overabundant deer on songbird populations. *Biological Conservation* **126**: 1–13.
- Allombert S, Stockton S, Martin J-L. 2005.** A Natural Experiment on the Impact of Overabundant Deer on Forest Invertebrates. *Conservation Biology* **19**: 1917–1929.
- Alverson WS, Waller DM, Solheim SL. 1988.** Forests Too Deer: Edge Effects in Northern Wisconsin. *Conservation Biology* **2**: 348–358.
- Banner A, MacKenzie WH, Pojar J, MacKinnon A, Saunders SC, Klassen H. 2014.** *A Field Guide to Ecosystem Classification and Identification for Haida Gwaii.*
- Bardgett RD, Wardle DA. 2003.** Herbivore-Mediated Linkages Between Aboveground and Belowground Communities. *Ecology* **84**: 2258–2268.
- Baselga A. 2010.** Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography* **19**: 134–143.
- Baselga A. 2013.** Separating the two components of abundance-based dissimilarity: balanced changes in abundance vs. abundance gradients. *Methods in Ecology and Evolution* **4**: 552–557.
- Baselga A, Orme D, Vileger S, et al. 2018.** *betapart: Partitioning Beta Diversity into Turnover and Nestedness Components.*
- Bennett JN, Blevins LL, Barker JE, Blevins DP, Prescott CE. 2011.** Increases in tree growth and nutrient supply still apparent 10 to 13 years following fertilization and vegetation control of salal-dominated cedar-hemlock stands on Vancouver Island. *Canadian Journal of Forest Research* **33**: 1516–1524.
- Bergström R, Edenius L. 2003.** From twigs to landscapes – methods for studying ecological effects of forest ungulates. *Journal for Nature Conservation* **10**: 203–211.
- Bernes C, Macura B, Jonsson BG, et al. 2018.** Manipulating ungulate herbivory in temperate and boreal forests: effects on vegetation and invertebrates. A systematic review. *Environmental Evidence* **7**: 13.
- Bossuyt B, Hermy M, Deckers J. 1999.** Migration of herbaceous plant species across ancient–recent forest ecotones in central Belgium. *Journal of Ecology* **87**: 629–638.
- Cáceres MD, Jansen F, Dell N. 2020.** *indicspecies: Relationship Between Species and Groups of Sites.*
- Carpio AJ, Apollonio M, Acevedo P. 2021.** Wild ungulate overabundance in Europe: contexts, causes, monitoring and management recommendations. *Mammal Review* **51**: 95–108.

Chollet S, Baltzinger C, Ostermann L, Saint-André F, Martin J-L. 2013. Importance for forest plant communities of refuges protecting from deer browsing. *Forest Ecology and Management* **289**: 470–477.

Chollet S, Baltzinger C, Saout SL, Martin J-L. 2013. A better world for bryophytes? A rare and overlooked case of positive community-wide effects of browsing by overabundant deer. *Écoscience* **20**: 352–360.

Chollet S, Bergman C, Gaston AJ, Martin J-L. 2015. Long-term consequences of invasive deer on songbird communities: Going from bad to worse? *Biological Invasions* **17**: 777–790.

Chollet S, Maillard M, Schörghuber J, Grayston SJ, Martin J-L. 2021. Deer slow down litter decomposition by reducing litter quality in a temperate forest. *Ecology* **102**: e03235.

Chollet S, Martin J-L. 2013. Declining woodland birds in North America: should we blame Bambi? *Diversity and Distributions* **19**: 481–483.

Chollet S, Padié S, Stockton S, Allombert S, Gaston AJ, Martin J-L. 2016. Positive plant and bird diversity response to experimental deer population reduction after decades of uncontrolled browsing. *Diversity and Distributions* **22**: 274–287.

Comeau VM, Daniels LD, Knochenmus G, Chavardès RD, Zeglen S. 2019. Tree-Rings Reveal Accelerated Yellow-Cedar Decline with Changes to Winter Climate after 1980. *Forests* **10**: 1085.

Côté SD, Rooney TP, Tremblay J-P, Dussault C, Waller DM. 2004. Ecological Impacts of Deer Overabundance. *Annual Review of Ecology, Evolution, and Systematics* **35**: 113–147.

Crête M, Daigle C. 1999. Management of indigenous North American deer at the end of the 20th century in relation to large predators and primary production. *Acta Veterinaria Hungarica* **47**: 1–16.

Dufrêne M, Legendre P. 1997. Species Assemblages and Indicator Species: the Need for a Flexible Asymmetrical Approach. *Ecological Monographs* **67**: 345–366.

Engelstoft C. 2001. Effects of Sitka black-tailed deer (*Odocoileus hemionus sitkensis*) on understory in old-growth forests on Haida Gwaii (Queen Charlotte Islands), British Columbia.

Forbes ES, Cushman JH, Burkepille DE, Young TP, Klope M, Young HS. 2019. Synthesizing the effects of large, wild herbivore exclusion on ecosystem function. *Functional Ecology* **33**: 1597–1610.

Fraser LH, Chanway CP, Turkington R. 1995. The competitive role of *Gaultheria shallon* on planted western hemlock and western red cedar saplings on northern Vancouver Island. *Forest Ecology and Management* **75**: 27–39.

Gao J, Carmel Y. 2020. Can the intermediate disturbance hypothesis explain grazing–diversity relations at a global scale? *Oikos* **129**: 493–502.

Gill RMA. 1992. A Review of Damage by Mammals in North Temperate Forests: 3. Impact on Trees and Forests. *Forestry: An International Journal of Forest Research* **65**: 363–388.

Gotelli NJ, Graves GR. 1996. *Null Models in Ecology*.

Habeck CW, Schultz AK. 2015. Community-level impacts of white-tailed deer on understory plants in North American forests: a meta-analysis. *AoB PLANTS* **7**.

Hennon PE, D'Amore DV, Schaberg PG, Wittwer DT, Shanley CS. 2012. Shifting Climate, Altered Niche, and a Dynamic Conservation Strategy for Yellow-Cedar in the North Pacific Coastal Rainforest. *BioScience* **62**: 147–158.

Hester AJ, Edenius L, Buttenschøn RM, Kuiters AT. 2000. Interactions between forests and herbivores: the role of controlled grazing experiments. *Forestry: An International Journal of Forest Research* **73**: 381–391.

Keddy PA. 1992. Assembly and response rules: two goals for predictive community ecology. *Journal of Vegetation Science* **3**: 157–164.

Klinkenberg B. 2020. E-Flora BC: Electronic Atlas of the Flora of British Columbia [eflora.bc.ca]. Lab for Advanced Spatial Analysis, Department of Geography, University of British Columbia, Vancouver.

Lortie CJ, Brooker RW, Choler P, et al. 2004. Rethinking plant community theory. *Oikos* **107**: 433–438.

Maillard M, Martin J-L, Chollet S, Catomeris C, Simon L, Grayston SJ. 2021. Belowground effects of deer in a temperate forest are time-dependent. *Forest Ecology and Management* **493**: 119228.

Martin J-L, Baltzinger C. 2002. Interaction among deer browsing, hunting, and tree regeneration. *Canadian Journal of Forest Research* **32**: 1254–1264.

Martin J-L, Chamailé- Jammes S, Waller DM. 2020. Deer, wolves, and people: costs, benefits and challenges of living together. *Biological Reviews* **95**: 782–801.

Martin J-L, Stockton SA, Allombert S, Gaston AJ. 2010. Top-down and bottom-up consequences of unchecked ungulate browsing on plant and animal diversity in temperate forests: lessons from a deer introduction. *Biological Invasions* **12**: 353–371.

Mladenoff DJ, Stearns F. 1993. Eastern Hemlock Regeneration and Deer Browsing in the Northern Great Lakes Region: A Re-examination and Model Simulation. *Conservation Biology* **7**: 889–900.

Muller RN. 2014. Nutrient Relations of the Herbaceous Layer in Deciduous Forest Ecosystems In: *The Herbaceous Layer in Forests of Eastern North America*. Oxford University Press, 14–34.

Neufeld HS, Young DR. 2014. Ecophysiology of the Herbaceous Layer in Temperate Deciduous Forests In: *The Herbaceous Layer in Forests of Eastern North America*. Oxford University Press, 35–95.

- Núñez MA, Bailey JK, Schweitzer JA. 2010.** Population, community and ecosystem effects of exotic herbivores: A growing global concern. *Biological Invasions* **12**: 297–301.
- Nuttle T, Ristau TE, Royo AA. 2014.** Long-term biological legacies of herbivore density in a landscape-scale experiment: forest understoreys reflect past deer density treatments for at least 20 years. *Journal of Ecology* **102**: 221–228.
- Oksanen J, Blanchet FG, Friendly M, et al. 2019.** *vegan: Community Ecology Package*.
- Olden JD, Rooney TP. 2006.** On defining and quantifying biotic homogenization. *Global Ecology and Biogeography* **15**: 113–120.
- Olf H, Ritchie ME. 1998.** Effects of herbivores on grassland plant diversity. *Trends in Ecology and Evolution* **13**: 261–265.
- Piazza M-V, Garibaldi LA, Kitzberger T, Chaneton EJ. 2016.** Impact of introduced herbivores on understory vegetation along a regional moisture gradient in Patagonian beech forests. *Forest Ecology and Management* **366**: 11–22.
- Pojar J, Lewis T, Roemer H, Wildford JD. 1980.** *Relationships between introduced black-tailed deer and the plant life of the Queen Charlotte Islands, British Columbia*. Smithers: Ministry of Forests.
- R Core Team. 2020.** R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ramirez JI, Jansen PA, Poorter L. 2018.** Effects of wild ungulates on the regeneration, structure and functioning of temperate forests: A semi-quantitative review. *Forest Ecology and Management* **424**: 406–419.
- Randall JA, Walters MB. 2011.** Deer density effects on vegetation in aspen forest understoreys over site productivity and stand age gradients. *Forest Ecology and Management* **261**: 408–415.
- Rhodes AC, Wan HY, St. Clair SB. 2017.** Herbivory impacts of elk, deer and cattle on aspen forest recruitment along gradients of stand composition, topography and climate. *Forest Ecology and Management* **397**: 39–47.
- Roberts MR, Gilliam FS. 2014.** Response of the Herbaceous Layer to Disturbance in Eastern Forests In: *The Herbaceous Layer in Forests of Eastern North America*. Oxford University Press, 302–320.
- Rooney TP, Waller DM. 2003.** Direct and indirect effects of white-tailed deer in forest ecosystems. *Forest Ecology and Management* **181**: 165–176.
- Stockton SA, Allombert S, Gaston AJ, Martin J-L. 2005.** A natural experiment on the effects of high deer densities on the native flora of coastal temperate rain forests. *Biological Conservation* **126**: 118–128.

Stroh N, Baltzinger C, Martin J-L. 2008. Deer prevent western redcedar (*Thuja plicata*) regeneration in old-growth forests of Haida Gwaii: Is there a potential for recovery? *Forest Ecology and Management*.

Sutherland Brown A. 1968. *Geology of the Queen Charlotte Islands, British Columbia.*

Suzuki M, Miyashita T, Kabaya H, Ochiai K, Asada M, Kikvidze Z. 2013. Deer herbivory as an important driver of divergence of ground vegetation communities in temperate forests. *Oikos* **122**: 104–110.

Takatsuki S. 2009. Effects of sika deer on vegetation in Japan: A review. *Biological Conservation* **142**: 1922–1929.

Tanentzap AJ, Burrows LE, Lee WG, Nugent G, Maxwell JM, Coomes DA. 2009. Landscape-level vegetation recovery from herbivory: progress after four decades of invasive red deer control. *Journal of Applied Ecology* **46**: 1064–1072.

Turner NJ. 2004. *Plants of Haida Gwaii: Xaadaa Gwaay gud gina k'Aws.* Winlaw, B.C: Sono Nis Press.

Verheyen K, Hermy M. 2004. Recruitment and growth of herb-layer species with different colonizing capacities in ancient and recent forests. *Journal of Vegetation Science* **15**: 125–134.

Waller D. 2014. Effects of Deer on Forest Herb Layers In: *The Herbaceous Layer in Forests of Eastern North America.* Oxford University Press, 369–399.

Waller DM. *Effects of Deer on Forest Herb Layers.* Oxford University Press.

Warren RJ. 2011. Deer overabundance in the USA: recent advances in population control. *Animal Production Science* **51**: 259–266.

Wiegmann SM, Waller DM. 2006. Fifty years of change in northern upland forest understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation* **129**: 109–123.

Tables

Table 1. Indicator species (measured by Indicator Value, Dufrêne and Legendre 1997) per treatment (Inside/Outside exclosures) per year (1997: onset; 2017: last monitoring).

	Inside	Outside
1997	No species associated to this treatment	No species associated to this treatment
2017	<i>Chamaecyparis nootkatensis</i> <i>Thuja plicata</i> <i>Gaultheria shallon</i> <i>Menziesia ferruginea</i> <i>Vaccinium parvifolium</i> <i>Cornus canadensis</i> <i>Rubus pedatus</i> <i>Blechnum spicant</i>	No species associated to this treatment

Accepted Manuscript

Figure captions

Figure 1. Study area and study sites

Figure 2. Comparison of vascular species richness, diversity (Shannon index), vascular plant cover and moss cover inside and outside deer exclosures in 1997 (left) and 2017 (right).

Statistical differences were tested with two way ANOVAs. *:p-value<0.05, ***: p-value<0.001.


Figure 3. Linear model inside (black dots) and outside (open triangles) deer exclosures in 2017 for variables selected in stepwise regression (see Supplementary data Table S2 for statistical models). Statistical significance was tested with Linear Regression Models. (*): p-value<0.1, *: p-value<0.05, ***: p-value<0.001.

Figure 4. Standardized Effect Sized (SES) of the mean soil moisture index (A) and soil nutrient index (B) as a function of respectively precipitation gradient (Distance to West Coast as a proxy) and soil fertility (soil C : N as a proxy) in 2017. Black dots correspond to plots inside deer exclosures and open triangles to plots outside deer exclosures. Statistical significance was tested with Linear Regression Models. (*): p-value<0.1, *: p-value<0.05.

Figure 5. Beta diversity measure with Bray-Curtis index (A, B) and partitioning in its two components: 1) balanced variation in abundance (Bray-Curtis bal) representing substitution of species (C, D), and 2) abundance gradients (Bray-Curtis gra) representing the change in abundance of the same species between plot (E, F) inside and outside deer exclosures in 1997 (left) and 2017 (right). Statistical differences were test with one-way ANOVA. ***: p<0.001.

Figure 6. NMDS analysis in 1997 (stress =0.21) and 2017 (stress =0.15). Open triangles are plots outside exclosures and black dots are plots inside exclosures.

Figure 1



Acces

scrip

Figure 2






Figure 3




Accepted Manuscript

Figure 4




Accepted Manuscript

Figure 5



A

Figure 6



Accepted Manuscript