



HAL
open science

Cadaverine regulates biotin synthesis to modulate primary root growth in Arabidopsis

Nicole Gibbs, Shih-Heng Su, Samuel Lopez-Nieves, Stéphane Mann, Claude Alban, Hiroshi Maeda, Patrick Masson

► **To cite this version:**

Nicole Gibbs, Shih-Heng Su, Samuel Lopez-Nieves, Stéphane Mann, Claude Alban, et al.. Cadaverine regulates biotin synthesis to modulate primary root growth in Arabidopsis. *The Plant Journal*, 2021, 107 (5), pp.1283-1298. 10.1111/tpj.15417 . hal-03291375

HAL Id: hal-03291375

<https://hal.science/hal-03291375v1>

Submitted on 20 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DR NICOLE GIBBS (Orcid ID : 0000-0002-2166-731X)

DR HIROSHI A MAEDA (Orcid ID : 0000-0003-0246-694X)

Article type : Original Article

Title: Cadaverine regulates biotin synthesis to modulate primary root growth in Arabidopsis

Authors: Nicole M. Gibbs^{1,#}, Shih-Heng Su¹, Samuel Lopez-Nieves², Stéphane Mann³, Claude Alban⁴, Hiroshi A. Maeda² and Patrick H. Masson^{1*}

Affiliations:

¹Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI 53706, USA.

²Department of Botany, University of Wisconsin-Madison, Madison, WI 53706, USA.

³ Muséum National d'Histoire Naturelle, UMR 7245, CNRS, MNHN, Molécules de Communication et Adaptation des Micro-organismes, CP 54, 57 Rue Cuvier, 75005 Paris, France.

⁴Univ. Grenoble Alpes, INRAE, CEA, CNRS, IRIG, LPCV, 38000 Grenoble, France.

#Current Address: Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological Studies, La Jolla, CA 92037, USA

Running Title: Cadaverine inhibits biotin synthesis

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/TPJ.15417](https://doi.org/10.1111/TPJ.15417)

This article is protected by copyright. All rights reserved

Keywords: Polyamines, Cadaverine, Biotin, Root architecture, *Arabidopsis thaliana*

*Corresponding author: email: phmasson@wisc.edu; address: Laboratory of Genetics, 425-g Henry Mall, Madison, WI 53706, USA.

Summary

Cadaverine, a polyamine, has been linked to modification of root growth architecture and response to environmental stresses in plants. However, the molecular mechanisms that govern the regulation of root-growth by cadaverine are largely unexplored. Here we conducted a forward genetic screen and isolated a mutation, *cadaverine hypersensitive 3 (cdh3)*, which resulted in increased root-growth sensitivity to cadaverine, but not other polyamines. This mutation affects the *BIO3-BIO1* biotin biosynthesis gene. Exogenous supply of biotin and a pathway intermediate downstream of BIO1, 7,8-diaminopelargonic acid, suppressed this cadaverine sensitivity phenotype. An *in vitro* enzyme assay showed cadaverine inhibits the BIO3-BIO1 activity. Furthermore, cadaverine-treated seedlings displayed reduced biotinylation of Biotin Carboxyl Carrier Protein 1 (BCCP1) of the Acetyl-CoA Carboxylase complex involved in *de novo* fatty acid biosynthesis, resulting in decreased accumulation of triacylglycerides. Taken together, these results revealed an unexpected role of cadaverine in the regulation of biotin biosynthesis, which leads to modulation of primary root growth of plants.

Introduction

Cadaverine (1,5-Diaminopentane) is a polyamine associated with environmental perception and modulation of biological processes in prokaryotic and eukaryotic organisms. In plants, alterations in cadaverine levels induce changes in root architecture (Gamarnik and Frydman, 1991; Niemi *et al.*, 2002; Liu *et al.*, 2014; Strohm *et al.*, 2015), fresh weight (Cassán, 2009; Tomar *et al.*, 2013), and stress response (Aziz *et al.*, 1998; Liu *et al.*, 2000; Shevyakova *et al.*, 2001; Kuznetsov *et al.*, 2007; Cassán, 2009; Sziderics *et al.*, 2010) with implications in crop yield and environmental stress resistance (Reviewed in (Jancewicz *et al.*, 2016).

Cadaverine is produced by decarboxylation of lysine in plastids (Bunsupa *et al.*, 2012), a process that has been shown to be upregulated by environmental stresses, including heat, drought and salt stress, in ice plant (Kuznetsov *et al.*, 2007; Shevyakova *et al.*, 2001), pepper (Sziderics *et al.*, 2010), and maize (Simon-Sarkadi *et al.*, 2014). On the other hand, free cadaverine levels are below the limit of detection in *Arabidopsis thaliana* (Bunsupa *et al.*,

2012; Shimizu *et al.*, 2019). However, all plants can accumulate cadaverine directly from their environment, in the form of molecules secreted by microbes from the phyllosphere, rhizosphere or endosphere, or diffusing from decomposing matter (Cassán, 2009). Importantly, both endogenous cadaverine and exogenously applied cadaverine can modulate plant growth and environment-related stress resistance (Reviewed in Jancewicz *et al.*, 2016). *Arabidopsis thaliana* seedlings exposed to exogenous cadaverine display altered root growth behavior, clearly indicating a role for this compound in root-growth regulation (Strohm *et al.*, 2015).

Similar to other polyamines, cadaverine catabolism produces hydrogen peroxide and other derived compounds, which serve as transducers in a variety of physiological, developmental, and environmental stress responses in animals, plants, and microbes. In *Leguminosae*, cadaverine is a precursor for quinolizidine alkaloids, which act to deter insects (Bunsupa *et al.*, 2012). Much of the polyamine content within plant cells has been shown to be in the conjugated form (Torrighiani *et al.*, 1987). Cadaverine can conjugate to lignin precursors, though other acceptors of conjugation are possible (Negrel *et al.*, 1992).

Despite these important roles of cadaverine in a broad aspect of plant development and physiology, very little is known about the molecular mechanisms that modulate plant responses to cadaverine. To date, three cadaverine-response genes have been identified in *Arabidopsis thaliana*: *OCT1*, *PAO4* and *SPMS* (Liu *et al.*, 2014; Strohm *et al.*, 2015). *OCT1* encodes an organic cation transporter previously implicated in carnitine transport (Lelandais-Brière *et al.*, 2007; Strohm *et al.*, 2015). *PAO4*, on the other hand, encodes a polyamine oxidase that functions in the back-conversion of spermine into spermidine, and spermidine into putrescine (Liu *et al.*, 2014). *SPMS* encodes a spermine synthase.

To better understand this process, we carried out a forward genetic screen in *Arabidopsis thaliana*, seeking mutations that affect root-growth responses to exogenous cadaverine. This screen revealed biotin as a possible target for the cadaverine-induced pathway. Also named vitamin B7 or H, biotin is an essential cofactor that functions in carboxylation reactions associated with *de novo* fatty acid biosynthesis and amino-acid homeostasis (Alban *et al.*, 2000).

In both plants and fungi, biotin synthesis starts with the conversion of pimeloyl-CoA to 7-keto-8-aminoperlarginic acid (KAPA), which is catalyzed by the cytosol- and peroxisome-localized KAPA synthase enzyme (encoded by the *BIO4* gene in *Arabidopsis*) (Pinon, 2005; Tanabe *et al.*, 2011). Subsequent reactions occur in the mitochondria, and produce 7,8-diaminoperlarginic acid (DAPA), dethiobiotin (DTB), and finally biotin. These reactions are catalyzed by the DAPA synthase (BIO1), DTB synthase (BIO3) and biotin synthase (BIO2) enzymes, respectively (Shellhammer and Meinke, 1990; Picciocchi *et al.*, 2001; Muralla *et al.*, 2007; Cobessi *et al.*, 2012). In plants and fungi, BIO1 and BIO3 form a bifunctional, homodimeric enzyme that is encoded by a single gene (*Arabidopsis BIO3-BIO1*, *At5G57590*) (Muralla *et al.*, 2007).

In *Arabidopsis*, five carboxylases have been shown to use biotin as a cofactor: Acetyl CoA carboxylase (ACCase, which catalyzes the first step in *de novo* fatty acid biosynthesis) (Sasaki *et al.*, 1993; Yanai *et al.*, 1995), 3-methylcrotonyl CoA carboxylase (MCCA, which contributes to leucine catabolism) (Alban *et al.*, 1993; Song *et al.*, 1994), propionyl CoA carboxylase (an enzyme that functions at the intersection between fatty-acid, amino-acid and sugar metabolism pathways) (Wurtele and Nikolau, 1990), geranyl-CoA carboxylase (Guan *et al.*, 1999), and pyruvate carboxylase (a key contributor to gluconeogenesis and lipogenesis) (Wurtele and Nikolau, 1990); (Reviewed in (Nikolau *et al.*, 2003)). A role for biotin beyond its function as a cofactor in carboxylases has also been suggested (Che, 2003), as has its potential contribution to the regulation of expression of defense-related genes (Li *et al.*, 2012). Furthermore, an uncharacterized nuclear-localized biotinylated protein has also been identified as potential regulator of target gene expression (Li *et al.*, 2012).

In this work, we show that cadaverine modulates primary root growth, mainly, by altering the biotin metabolic pathway in *Arabidopsis thaliana*. Cadaverine-induced inhibition of root growth can be rescued with supplementation of biotin and biotin-pathway intermediates downstream of BIO1. Furthermore, we observed that decreased biotinylation of ACCase subunit BCCP1, as a result of cadaverine treatment, leads to a reduction of fatty

acid synthesis. This study will contribute toward understanding how biotin limitation affects plant growth, and the role of cadaverine as a natural biotin inhibitor.

Results

The *cdh3* mutation enhances primary root growth sensitivity to cadaverine in *Arabidopsis*.

To identify molecular candidates in cadaverine response in plants, we carried out a forward genetic screen using a population of ethyl methanesulfonate (EMS)-mutagenized *Arabidopsis thaliana* plants in the Col-0 background. We screened M2 seedlings for altered root growth on media containing 200 μ M cadaverine. One of the cadaverine-hypersensitive mutants identified in this screen, named *cadaverine hypersensitive 3* (*cdh3*), showed a significant inhibition of primary root growth on cadaverine-containing media relative to wild type (Figure 1(a) and S1). This mutant is the main subject of investigation in this manuscript.

Cadaverine decreases primary root growth earlier in the *cdh3* mutant than the wild type.

To further characterize *cdh3*, we first plated wild type and *cdh3* mutant seeds on media containing cadaverine, ranging from 2 μ M to 2 mM, and quantified primary root growth under these conditions. The results demonstrated a strong inhibitory effect of cadaverine at concentrations as low as 2 μ M for *cdh3*, whereas a significant effect of cadaverine on wild-type root growth was observed only at concentrations of 20 μ M and higher (Figure 1(b)). At 20 μ M *cdh3* primary roots were 58.4% shorter than wild type. On control media lacking cadaverine, *cdh3* seedlings showed only a minor decrease in primary root growth compared to wild type (Figure 1(a)).

In previous work involving wild-type *Ler* seedlings, Strohm et al. showed that exogenous cadaverine inhibits primary root growth mostly by affecting cell elongation (Strohm et al., 2015). They also demonstrated the existence of substantial natural variation between *Arabidopsis* accessions in their primary root growth responses to exogenous cadaverine (Strohm et al., 2015). Therefore, we tested the effects of exogenous cadaverine

on wild-type Col-0, analyzing cell elongation and apical root meristem size. 8-day-old seedlings were grown on 200 μ M cadaverine or control media, and roots were treated with propidium iodide, a cell-wall stain (Strohm *et al.*, 2015), to determine cell sizes and zone boundaries in the root tip. The decrease in primary root growth was found to be mostly associated with alterations in cell size within the elongation zone (Figure S2), whereas the number of cells in the root apical meristem was not substantially affected in the wild type (Figure 1(c)-(e)). *cdh3*, however, showed a significantly smaller root meristem upon cadaverine treatment, while *cdh3* grown on control media showed an equal number of cells compared to wild type. In addition to decreased cell division and cell expansion in *cdh3*, propidium iodide permeated a number of cells within the elongation zone, suggesting cell death (Figure 1(c)).

We next wanted to determine the time at which cell elongation is modified following cadaverine treatment by measuring changes in tip growth over time. Five-day-old seedlings were transferred onto split agar plates containing control or 200 μ M cadaverine media and imaged for 4 days using a high-resolution camera. Wild type roots began to show a significant decrease in tip growth after 96 hours of cadaverine treatment. *cdh3*, on the other hand, showed a decrease in root tip growth already 40 hours after onset of cadaverine treatment (Figure 1(f)). Therefore, the *cdh3* mutation substantially increases the speed of root-tip response to exogenous cadaverine.

Because arginine-derived polyamines, such as putrescine and putrescine-derived spermidine and spermine, were previously shown to also affect primary root growth in young *Arabidopsis* seedlings, we tested *cdh3*'s ability to respond to these PAs as well (Liu *et al.*, 2014; Strohm *et al.*, 2015). The results demonstrated that *cdh3* and wild-type show similar root-growth response to putrescine and spermine, with *cdh3* showing only a mild resistance to 500 μ M spermidine (Figure 1(g)). Hence, the *cdh3* mutation increases root-growth sensitivity to cadaverine, and it has little to no effect on root responses to putrescine and putrescine-derived polyamines.

Taken together, our data indicate that *cdh3* alters a mechanism involved in the control of primary root growth in response to cadaverine, a process that targets the regulation of anisotropic cell expansion occurring in the main elongation zone as well as the meristem.

cdh3*'s enhanced cadaverine response is the result of a mutation in *BIO3-BIO1

To evaluate the genetic makeup of *cdh3*, we backcrossed it to Col-0, and self-pollinated the corresponding F1. Segregating F2 progeny were investigated for root-growth sensitivity to cadaverine. Results showed 22.6% seedlings (12 out of 53) exhibiting hypersensitive cadaverine response, consistent with a single, recessive mutation (Chi-Square value of 0.1179; $p \sim 0.3$). To clone *CDH3*, we generated a pool of 162 cadaverine-hypersensitive *cdh3* F2 seedlings from this segregating population, extracted DNA from this pool, and subjected it to whole genome Next-Generation Sequencing. The candidate mutation was mapped to a 4.6Mb region of chromosome 5, using the approach described in (Schneeberger *et al.*, 2009). Within this region, a non-synonymous mutation was found within the *BIO3-BIO1* gene (*At5G57590*), which encodes a bifunctional enzyme involved in biotin synthesis (Cobessi *et al.*, 2012). This mutation resulted in an A662T amino-acid change within the BIO1 catalytic pocket of the protein (Figure 2(a), 2(b)), which functions as an aminotransferase that catalyzes the conversion of KAPA into DAPA within the pathway. The A662 amino acid is located in close proximity to the PLP-cofactor binding site, potentially affecting enzymatic activity (Figure 2(c)).

To confirm association of the cadaverine-hypersensitive phenotype of *cdh3* with the A662T mutation in *BIO3-BIO1*, we transformed a wild-type *BIO3-BIO1* transgene under the control of its native promoter into *cdh3*. Four independent, homozygous transgenic lines (*cdh3* [*BIO3-BIO1pro::BIO3-BIO1*]) were recovered and screened on cadaverine-containing media. Results shown in Figure 2(d) demonstrated rescue of the cadaverine-hypersensitive phenotype of *cdh3* by the wildtype *BIO3-BIO1pro::BIO3-BIO1* transgene in all four independent lines. Furthermore, the transgene also rescued the decreased root-growth phenotype displayed by *cdh3* on control media, suggesting the corresponding A662T

mutation in *BIO3-BIO1* was also responsible for *cdh3* root-growth restriction on control media (Figure 1(a)).

While performing these transformation-rescue experiments, we also noted that three of the four transformation-rescue lines (lines 1, 3 and 4) displayed increased root-growth resistance to cadaverine, a phenotype that is opposite to the hypersensitive response displayed by *cdh3*. To determine if the transformation-rescue lines displayed increased transgene expression level, *BIO3-BIO1* expression was quantified using RT-qPCR on 8-day old seedlings grown on control media. While no significant expression alterations were found between *cdh3* and wild-type seedlings, transformation-rescue lines displayed 4.5- to 14.8-fold increases in *BIO3-BIO1* expression relative to wild type (Figure 2(e)). The level of root-growth insensitivity to cadaverine correlated with increased expression in resistant transformation-rescue lines ($R^2=0.68$). Taken together, these results suggest dosage of *BIO3-BIO1* affects root-growth response to cadaverine.

Cadaverine response is chemically suppressed by biotin

Initial studies with the embryo-lethal *bio1-1* mutant showed that knockout of biotin synthesis in plants can be rescued by application of exogenous biotin (Schneider *et al.*, 1989). To determine if *cdh3*'s hypersensitive root-growth response to cadaverine is the result of decreased biotin, we supplemented 200 μ M cadaverine-containing media with biotin at concentrations ranging from 10 nM to 100 nM (Schneider *et al.*, 1989). The cadaverine-hypersensitive phenotype displayed by *cdh3* roots was rescued in a dose-dependent manner to near wild type root growth by addition of biotin (Figure 3(a)). In addition, the inhibitory effect of cadaverine on wild-type root growth was also suppressed by addition of 80 nM biotin to the medium (Figure 3(a)). These data are compatible with cadaverine targeting the biotin pathway or modulating biotin availability or function. Alternatively, cadaverine and biotin could function through parallel pathways to modify primary root growth.

To identify a step in the biotin synthesis pathway that may be potentially affected by cadaverine, we tested intermediate compounds of the biotin biosynthesis pathway for their abilities to suppress root-growth response to cadaverine. Figure 3(b) shows that addition of

1 μM DAPA, the product of BIO1 activity, resulted in complete suppression of cadaverine's effects on root growth, whereas addition of 1 μM KAPA, the BIO1 substrate, did not. These results suggest that BIO2, BIO3, and BIO4 functions are not impaired by cadaverine. Rather, BIO1 enzymatic activity, channeling of DAPA to the BIO3 active site, or KAPA's availability or transport, is/are inhibited by cadaverine.

Since root-growth response to cadaverine can be alleviated by biotin supplementation in both *cdh3* and wild-type seedlings, we tested if we could identify cadaverine-hypersensitive lines that cannot be rescued by biotin, thereby identifying downstream regulators in the cadaverine-response pathway. *ORGANIC CATION TRANSPORTER 1* mutant, *oct1-1*, and *POLYAMINE OXIDASE 4* mutant, *pao4-1*, have been shown to be hypersensitive to cadaverine (Liu *et al.*, 2014; Strohm *et al.*, 2015). *oct1* and *pao4* mutants were tested for chemical rescue on biotin- and cadaverine-containing media. As shown in Figure 3(c) and 3(d) *oct1* and *pao4* mutant roots no longer displayed cadaverine hypersensitivity in the presence of biotin. Hence, biotin appears to function downstream of, or parallel to, *OCT1* and *PAO4* in the cadaverine-response pathway.

Cadaverine directly modulates biotin synthesis by altering BIO3-BIO1 enzymatic activity

As the results of the chemical rescue experiment suggest that cadaverine may target DAPA synthase, we expressed wild type BIO3-BIO1 recombinant protein *in vitro* to test whether its enzymatic activity was fully functional upon cadaverine treatment. Previous *in vitro* studies by Cobessi *et al.*, have shown that the product of the BIO1 reaction, DAPA, is channeled from the BIO1 active site to the BIO3 active site, and thus not released into the solution. Therefore, we chose to detect the product of BIO3, dethiobiotin (DTB) using a FRET based competition assay, in which DTB will displace the lower affinity 4'-hydroxyazobenzene-2-carboxylic acid (HABA) quencher from fluorescently tagged avidin (Batchelor *et al.*, 2007). We set up *in vitro* reactions in the presence or absence of cadaverine to determine whether cadaverine alters the enzyme activity. The result shown in Figure 4(a) indicates less formation of DTB in reactions containing cadaverine compared to

control. The data suggests that cadaverine affects the flow of molecules through the biotin synthesis pathway at the level of DAPA or DTB synthesis, leading to decreased biotin synthesis and altered root growth. Attempts to express the *cdh3* mutant protein *in vitro* were unsuccessful, suggesting the A662T mutation may affect BIO3-BIO1 protein stability.

BIO3-BIO1 transcript and protein are unaffected by cadaverine treatment

To determine if cadaverine also affects the regulation of *BIO3-BIO1* expression, we next quantified the levels of *BIO3-BIO1* transcripts in control and cadaverine-treated wild type and *cdh3* seedlings. RNA was extracted from whole seedlings germinated on cadaverine-containing media, or treated with cadaverine for 24, 48 or 72 hours, along with controls grown on cadaverine-free medium. No significant differences were identified between cadaverine-treated and control samples, or between *cdh3* and wild-type seedlings (Figure 4(b)). Additionally, cadaverine did not influence the expression of other biotin synthesis genes, *BIO4* (*At5G04620*) and *BIO2* (*At2G43360*) (Figure S3). This result contrasts with an earlier report that showed increased *BIO2* expression upon biotin depletion (Patton *et al.*, 1996). Together, these data suggest that cadaverine-induced inhibition of biotin synthesis under our conditions is not sufficiently severe to result in increased *BIO2* expression.

To investigate a potential effect of cadaverine on BIO3-BIO1 protein abundance, wild type, *cdh3*, and *cdh3* [*BIO3-BIO1pro::BIO3-BIO1*] rescue line #2 were germinated and grown on either control or cadaverine-containing media for 8 days. Proteins were extracted from dissected root and shoot tissues, separated by SDS-polyacrylamide gel electrophoresis, and subjected to western blot analysis, probing with a BIO3-BIO1-specific antibody (Cobessi *et al.*, 2012). The *cdh3* [*BIO3-BIO1pro::BIO3-BIO1*] rescue line expressed increased amounts of BIO3-BIO1 protein relative to wild type, irrespective of cadaverine treatment (Figure 4(c)), consistent with its increased *BIO3-BIO1* transcripts (Figure 2(e)). Minimal differences in relative BIO3-BIO1 protein abundance were observed between cadaverine-treated and control samples for all genotypes and tissues tested (Figure 4(c)), suggesting

that cadaverine has limited impacts on overall expression of the BIO3-BIO1 protein in *Arabidopsis*.

Cadaverine treatment decreases the biotinylation of biotin carboxyl carrier proteins in *planta*

Considering that cadaverine inhibits biotin synthesis, we hypothesized that treating *Arabidopsis* seedlings with cadaverine should result in decreased biotinylation of target proteins in plant tissues. Previous studies have shown that probing western blot membranes with a streptavidin probe allows the detection of a minimum of three biotinylated plant proteins, including methylcrotonoyl-CoA carboxylase (MCCA), involved in amino acid metabolism, and the biotin carboxyl carrier protein 1 and 2 subunits (BCCP1 and BCCP2) of chloroplastic, heterodimeric acetyl-CoA carboxylase (ACCase), involved in fatty acid metabolism (Aubert *et al.*, 1996; Che, 2003; Li *et al.*, 2012). BCCP2 is primarily expressed in flowers and developing seeds, but scarcely detectable in seedling tissues (Thelen *et al.*, 2001). We used a similar Western blot analysis to determine whether cadaverine treatment is sufficient to induce differences in MCAA and/or BCCP proteins biotinylation in dissected root and/or shoot tissues.

Wild type, *cdh3* and *cdh3* [*BIO3-BIO1pro::BIO3-BIO1*] #2 rescue seedlings were germinated and grown on control or cadaverine-containing media for 8 days. Roots and shoots were then dissected. Proteins extracted from these samples were subjected to western blot analysis using a streptavidin probe. The amounts of biotinylated MCCA protein showed little change upon cadaverine treatment in the root (Figure 5(a), 5(c)) or shoot (Figure 5(b) and 5(d)). On the other hand, a decrease in biotinylated BCCP1 protein was observed in both roots (Figure 5(a), 5(e)) and shoots (Figure 5(b), 5(f)), with a larger magnitude of decrease occurring in the roots.

Cadaverine-induced changes in biotinylated protein abundance could be due to decreased levels of target protein biotinylation, and/or relate to changes in abundance of the corresponding proteins. To investigate these possibilities, we used MCCA- and BCCP1-specific antibodies to analyze the abundance of these proteins in control and cadaverine-

treated seedlings. Results shown in Figure 5(c) demonstrated increased MCCA levels in the root upon cadaverine treatment only in *cdh3*, while no significant changes were observed in the shoot (Figure 5(d)). BCCP1 protein abundance in cadaverine-treated samples relative to controls, showed no significant changes in the root or shoot (Figure 5(e), 5(f)).

qRT-PCR analysis of *MCCA* and *BCCP1* expression showed no significant transcriptional alteration in *cdh3* and Col-0 upon cadaverine treatment, whether applied since germination, or for 24, 48 or 72 hours (Figure S4). Taken together, our data are consistent with cadaverine treatment leading to decreased biotin synthesis, thereby resulting in decreased biotinylation in BCCP1 protein in both shoots and roots. They are also consistent with decreased biotinylation level of MCCA protein in *cdh3* mutant roots, as increased MCCA apoprotein abundance is not accompanied by increased levels of biotinylated MCCA (Figure 5(c)).

In summary, cadaverine treatment significantly affects the amount of biotinylated BCCP1 protein in both shoots and roots.

Lipid composition is altered by cadaverine

Acetyl CoA Carboxylase requires biotinylation of the BCCP1 subunit to generate malonyl-CoA from acetyl-CoA at the first step of fatty acid biosynthesis. The cadaverine-induced reduction in biotinylated BCCP1 protein in plant tissues is expected to decrease fatty acid production. To test this hypothesis, we determined the impact of cadaverine treatment on the lipid profile of Arabidopsis seedlings. Wild type Col-0 seeds were germinated and grown on control media for 5 days, then transferred to either control media or media containing 200 μ M cadaverine, and allowed to grow for 72 more hours, a time previously determined to be sufficient to allow primary root growth inhibition by cadaverine (Figure 1). At that time, lipids were extracted from these samples, and subjected to lipidomics analyses by MS/MS (Shiva *et al.*, 2013). Total polar lipids as well as diacylglycerol (DAG) and triacylglycerol (TAG) lipids, were analyzed with five biological replicates.

The results showed that total triacylglycerols (TAG) decreased in abundance upon cadaverine treatment, with 6 out of 31 TAG species showing significant decreases (Figure 6),

all of which were TAGs having 18:3 acyls. On the other hand, TAGs containing short-chain fatty acids, 16:0 and 16:1, had little changes with the treatment (Figure 6). The reduced TAG synthesis along with the higher percentage of short-chain fatty acid compositions in TAGs were also previously observed in Arabidopsis plants with limited biotin availability (Pommerrenig *et al.*, 2012). Therefore, these results are consistent with the negative effect of cadaverine on biotin availability and TAG production.

We observed similar levels of total polar lipid species between cadaverine-treated and control seedlings, with slight decreases observed in total phosphatidic acid (PA) and phosphatidylinositol (PI), and an increase in phosphatidylcholine (PC) (Table S1). Although total diacylglycerol (DAG) levels were not altered by cadaverine treatment, specific DAGs changed in abundance in response to cadaverine, including increases in DAG 16:0/16:1, DAG 18:0/16:2 and DAG 18:3/dnOPDA, and decreases in DAG 18:3/18:3 content (Table S1). In addition, we have also seen modification of phospholipids, albeit below the significance threshold, with PI and PA decreasing with cadaverine treatment.

Discussion

We have shown that cadaverine functions to inhibit primary root growth by modulating biotin synthesis through inhibition of the BIO3-BIO1 enzyme. Cadaverine response can be chemically rescued with biotin, DAPA, but not KAPA. While KAPA could potentially be excluded from transport into seedlings, the results of our *in vitro* reaction show cadaverine to inhibit BIO3-BIO1 activity. Furthermore, cadaverine did not alter expression of BIO3-BIO1 at the protein level in wildtype or *cdh3*, and overexpression of *BIO3-BIO1* from our *cdh3* rescue line resulted in increased BIO3-BIO1 protein and resistance to cadaverine, in agreement with an effect on BIO3-BIO1 enzymatic function.

Analysis of the *BIO3-BIO1* mutant, *cdh3*, reported here shows biotin deficiency results in decreased primary root growth, which is similar to the phenotype observed in the hypomorphic mutation in *BIO4* (Li *et al.*, 2012). Furthermore, analysis of *cdh3* root growth on cadaverine-containing media also shows evidence of cell death in the elongation zone upon staining with propidium iodide (Figure 1(c)). Evidence of cell death was also reported in

shoots of a hypomorphic mutant of *BIO4* (Li *et al.*, 2012), indicating that decreases in biotin cause cell death in both roots and shoots. The localization of cell death to the elongation zone of cadaverine-treated *cdh3* may reflect an increased requirement for biotin in tissues that undergo rapid cell growth and expansion. Alternatively, deficiencies in fatty acid synthesis have been shown to result in an accumulation of reactive oxygen species in the mitochondria, along with cell death (Wu *et al.*, 2015). The similar phenotypes observed in cadaverine-treated *cdh3* mutant seedlings (this study) and biotin-deficient mutants (Li *et al.*, 2012; Wu *et al.*, 2015) thus suggests a deficiency in biotin synthesis may underlie the developmental impact of cadaverine.

Inhibition of biotin synthesis has long been a target for herbicides and antibiotics. While numerous compounds have been identified to inhibit biotin biosynthesis, to date only one natural product, amiclennomycin, had been identified to inhibit DAPA synthase in bacteria, through formation of a covalent adduct with PLP (Sandmark *et al.*, 2002). Our results indicate cadaverine is a natural DAPA synthase inhibitor in *Arabidopsis*. Cadaverine may function to inhibit BIO3-BIO1 through formation of a Schiff base with PLP, as observed with amiclennomycin, or through direct interaction with the binding site. BIO3-BIO1 inhibition by cadaverine cannot be mimicked by treatments with putrescine and/or putrescine-derived polyamines. The different inhibitory activities displayed by distinct polyamines may be determined by charge distribution of the amine groups or subcellular compartmentalization. Given the proximity of A662 to PLP, the *cdh3* mutant's hypersensitive response to cadaverine could be explained by an altered conformation of the mutant protein resulting in decreased reaction efficiency. However, multiple attempts at producing recombinant *cdh3* mutant enzyme failed, preventing us from directly testing this hypothesis. Future work will be aimed at characterizing cadaverine's inhibitory effect on DAPA synthase activity in plants.

As we have shown exogenous biotin rescues the cadaverine response phenotype, the effect of cadaverine treatment on primary root growth is in part associated with induced alterations in lipid composition, including decreased TAG content, and/or signaling phospholipids. Analysis of *SUC5*, required for biotin transport into the developing embryo from the maternal plant, shows the *suc5* and *bio1* or *bio2* double mutants rescued with

exogenous biotin have an increased proportion of short chain fatty acids (16:0 and 16:1), in accordance with our observation of cadaverine treatment (Figure 6; Table S1) (Pommerrenig *et al.*, 2012). The decrease in PI and PA could be due to increased activity through the phosphoinositide pathway to generate DAG to compensate for decreases in TAG. Alternatively, cadaverine may function to affect the components of the phosphoinositide pathway, independently from its effect on biotin, contributing to the root growth phenotype we see in response to cadaverine.

Biotin is also involved in amino acid homeostasis, gluconeogenesis, and other aspects of central metabolism. MCCAse has previously been shown to accumulate in response to biotin depletion or starvation to allow for recycling of amino acids (Aubert *et al.*, 1996; Che, 2003; Ding *et al.*, 2012). The increase of MCCA apoprotein in *cdh3* (Figure 5(a), 5(c)) is compatible with a mechanism that allows modulation of target MCCA protein abundance by biotin availability, in agreement with previous studies (Che, 2003; Li *et al.*, 2012). The absence of such changes in cadaverine-treated wild type plants may simply reflect a lower level of biotin limitation in wild type roots.

While root growth inhibition of *cdh3* was not observed with putrescine-derived polyamines (Figure 1(g)), the *pao4* phenotype and subsequent rescue by biotin (Figure 3(d)) suggests a connection between cadaverine and the putrescine-derived polyamines. Polyamine oxidases, such as PAO4, oxidize polyamines spermidine and spermine, thereby contributing to putrescine-derived polyamines homeostasis (Kamada-Nobusada *et al.*, 2008; Liu *et al.*, 2014). *pao4* mutants show increased spermine content, decreased spermidine and hydrogen peroxide, and changes in primary metabolism (Sequera-Mutiozabal *et al.*, 2016). Altered primary metabolism associated with the *pao4* mutation may reflect an indirect role for the putrescine-derived polyamines or their catabolic products in biotin synthesis or biotin's downstream effects.

The putrescine-derived polyamines are essential components for growth and development; for example, thermospermine is required for vasculature development in *Arabidopsis* (Kakehi *et al.*, 2008) and spermidine is required for eukaryotic translation (Park *et al.*, 1982). Studies on the functions of putrescine-derived polyamines have been

hampered by the embryo lethality associated with spermidine depletion (Imai *et al.*, 2004). Therefore, it is possible that putrescine-derived polyamines provide other essential roles that have yet to be uncovered.

OCT1 has been characterized as a carnitine transporter and also proposed to transport cations, such as polyamines (Lelandais-Brière *et al.*, 2007; Strohm *et al.*, 2015). In mammals the carnitine cycle allows acyl transport into the mitochondria for Beta-oxidation. An analogous mechanism has been proposed in plants after identification of a carnitine acyl carrier-like protein with homology to mammalian carnitine acyl carrier, but the role of carnitine in plants has yet to be further elucidated (Lawand *et al.*, 2002). *oct1* mutants could be sensitive to cadaverine because of altered polyamines homeostasis and/or localization. Alternatively, cadaverine-induced lipid deficiency could be exacerbated by altered acyl transport capabilities in the *oct1* mutants.

In some plants, cadaverine abundance has been found to increase in response to environmental stress, a process that is believed to contribute to stress mitigation. Furthermore, microbes from the phyllosphere, rhizosphere or endosphere have also been shown to provide plants with cadaverine, as does decomposing matter in proximity of a root system (Cassán, 2009). Both endogenous and exogenously applied cadaverine can modulate plant growth and environment-related stress resistance (Reviewed in Jancewicz *et al.*, 2016). We present here that cadaverine inhibits biotin biosynthesis and primary metabolism in *Arabidopsis* through direct inhibition of DAPA synthase, thereby reducing the accumulation of TAGs. If the physiological responses to cadaverine are adaptive, the presence of cadaverine, either through association with microbes (in most plants including *Arabidopsis thaliana*) or through stress-induced synthesis, where observed, may serve to divert a plant's resources from growth to defense responses. Future investigations addressing the conservation of cadaverine-response pathways in plants, and the development of methods aimed at better controlling the effects of cadaverine on biotin synthesis, its impact on primary metabolism within the biomass, and on root-system architecture, should yield important new insights, facilitating the development of new crop

varieties with improved productivity, enhanced energy content of the biomass, and increased adaptability to the stresses associated with climate change or growth on marginal lands.

Experimental Procedures

Seedling root growth measurements

Seedlings were grown on plates containing 1.5% agar, 0.5X Linsmaier and Skoog (LS) media pH 5.7 with 1.5% sucrose unless otherwise noted. After 48 hours of stratification, plates were moved to a 22°C growth chamber under long day conditions, 60 $\mu\text{mol}/\text{m}^2/\text{s}$ white light (cool-white fluorescent bulbs) and the wave assay was carried out as described in (Rutherford and Masson, 1996). Day lengths were 16 hours. After 6 days of growth, the position of the root tip was marked and seedlings were grown for 4 more days. After 10 days, plates were scanned using a document scanner and the growth from Day 6 to Day 10 was measured using the Neuron J plugin of Image J. Root growth was standardized to the control of each mutant line.

To determine if the response was specific to cadaverine, mutant lines were screened on media supplemented with putrescine (5 mM, 10 mM, and 15 mM), spermidine (300 μM , 500 μM and 700 μM), and spermine (200 μM , 300 μM and 400 μM), and root growth measurements were carried out as described above.

Meristem morphology and cell size

Zeiss confocal LSM780 microscope was used to characterize cell size and morphology of cadaverine-treated roots. 8 day-old seedlings were germinated on control media or 200 μM cadaverine-containing media. Seedlings were grown as described above. After 8 days in the growth chamber, seedlings were stained with 1.0 mg/mL propidium iodide for approximately 2 seconds and then rinsed with water. Roots were imaged using a 561 nm laser, 20X objective. Images were analyzed using Image J. Meristem was defined from the quiescent center to the first elongated cell in the cortex. The number of cells in the meristem was counted from the quiescent center along files of cortical cells to the first elongated cell (Perilli *et al.*, 2010). The cell length in the mature zone was quantified by measuring cells in the epidermis using Image J.

Root Tip Growth Measurement

Seedlings were germinated on control media, as above, and grown vertically to reduce waving. 3 days after germination, seedlings were transferred to control or 200 μM cadaverine-containing media and grown under 60 $\mu\text{mol}/\text{m}^2/\text{s}$ of continuous light. A high-resolution raspberry pi camera was used to image seedlings every 8 hours. Imaging began within 10 minutes after transfer. Image J was used to measure tip growth every 8 hours. Statistical difference in root tip growth between cadaverine and control treatment was tested using a mixed effects model with an F-test for the interaction between time and treatment, corrected with Tukey's adjustment for multiple comparisons in SAS Studio Version 3.8 (Gueorguieva and Krystal, 2004).

Forward Genetic Screen

To generate an EMS population, approximately 0.19 g of Col-0 seeds (M0) were treated with 50 mLs of 0.2% EMS for 15 hours, then rinsed with water 8 times (Weigel and Glazebrook, 2002). On the last wash, seeds were soaked in water for an hour. Seeds were then resuspended in 0.1% agar and pipetted onto soil. After EMS application M0 seeds are referred to as M1. M1 plants were self-pollinated and M2 seeds were collected in individual pools for each plant. 1,064 M2 lines were screened on plates containing LS media supplemented with 200 μM cadaverine. Seedlings showing a hypersensitive response, a decrease in root growth compared to the wild type on cadaverine media, were self-pollinated and the M3 lines were rescreened on 200 μM cadaverine and control media. Root growth of these M3 lines was quantified as described above. Student's T-test was done against Col-0 to determine if the mutant line was hypersensitive to cadaverine (Figure S1).

Next Generation Sequencing

A mapping population of *cadaverine hypersensitive 3* (*cdh3*) was generated by backcrossing *cdh3* to wild type Col-0. The segregating F2 population was screened on 50 μM cadaverine plates as described above, omitting sucrose. 162 cadaverine hypersensitive seedlings were pooled and frozen in liquid nitrogen, then homogenized with a 4.5 mm steel bead using Mixmill. DNA was extracted using the QIAGEN DNeasy plant mini kit. Next generation sequencing was done using Illumina HiSeq 2000 (UW-Biotech Facility) with 100 base pair,

paired-end reads. DNASTAR software (version 12.3) was used for alignment and variant identification. Reads with a Q value below 30 were removed from analysis. Variants present in multiple EMS mutant lines and Col-0 were subtracted from *cdh3* variant list. The location of the mutation was identified by looking at the percent SNP along the position in the genome (Zuryn *et al.*, 2010).

Chemical Rescue

Seedlings were germinated on LS media with or without 200 μ M cadaverine and 10 nM, 20 nM, 40 nM, 60 nM, 80 nM, or 100 nM biotin. Biotin was prepared as described in (Schneider *et al.*, 1989). Seedling root growth was quantified as described above. Student's T-test was done to determine significance against Col-0 using p-value of 0.05 as a cut off.

Transformation Rescue

Primers containing adapters for *Asc1* and *Pac1* restriction sites were used to amplify Col-0 wild-type *BIO3-BIO1* genomic DNA (*At5G57590*), from 1,233 base pairs upstream from the transcriptional start site to 382 base pairs 3' of the stop codon, yielding a 5,706 base pair product. The PCR reaction was restriction-digested with *Asc1* and *Pac1* and ligated into pMDC99 (Curtis, 2003). The ligated product was transformed into *E. coli* DUO competent cells (Lucigen, Middleton WI) and selected on 50 μ g/mL kanamycin. Transformation rescue was done on the *cdh3* mutant using the floral dip method as described (Clough and Bent, 1998) using *Agrobacteria* strain GV3101. T1 seeds were screened on 50 μ g/mL hygromycin to identify positive transformants.

Protein Expression and Isolation

Protein expression of pET-mBIO3-BIO1 was carried out as described previously (Cobessi *et al.*, 2012). The plasmid was introduced into Rosetta-2 *E. coli* competent cells (Novagen, Madison WI, USA) and cultured overnight at 37 °C, 200 r.p.m. in 10 mL LB medium containing 50 μ g/mL kanamycin. After overnight growth the culture was transferred into 500 mLs LB with appropriate antibiotic and supplemented with 10 μ M pyridoxine and 15 μ M thiamin (Cobessi *et al.*, 2012). The culture was incubated at 37°C until the OD₆₀₀ reached 0.3. At this point, 0.4 mM isopropylthio- β -D-galactoside (IPTG) was added, and the culture was grown for 16 hours at 18°C, shaking at 200 rpm. Cells were then harvested by

centrifuged at 4°C for 20 minutes at 4000g. Pellet was resuspend in lysis buffer with additional 1x protease inhibitor cocktail (Roche, Basel, Switzerland) and 500 µM PLP. Cell lysate was centrifuged at 10,000g for 30 mins at 4°C. Supernatant was collected and subjected to QIAexpress Ni-NTA column purification (Qiagen, Hilden, Germany). Eluted protein was desalted using Amicon Ultra centrifugal filter units, Ultra-15 MWCO 50 kDa (Millipore Sigma, Burlington, MA). Protein was stored and diluted in 100 mM HEPPS pH 8.6.

BIO3-BIO1 *in vitro* enzymatic assay

The assay was modified from the method developed by Cobessi et al., 2012. Reaction buffer contained the following: 100 mM HEPPS pH 8.6, 1 mM DTT, 100 mM NaCl, 5 mM MgCl₂, 3mM AdoMet, 10 mM NaHCO₃, and 0.3 mM ATP. Each reaction included various concentrations of racemic hydrochloride-KAPA (synthesized according to the procedure described by (Nudelman *et al.*, 2004) as substrate (0 µM or 20 µM), along with 0 or 50 µM Cadaverine dihydrochloride. BIO3-BIO1 enzyme at a final concentration of 6.9 µM was added to start the reactions. Reactions were run at 25°C for 8 minutes. At the end of each reaction, the enzyme was inactivated by addition of 20% acetic acid. Inactivated enzyme was removed from the reaction using a 10 kDa column (Pall NanoSep). Dethiobiotin (DTB) was detected using FluoReporter Biotin Quantitation Assay Kit (ThermoFisher F30751, Waltham MA). 25 µLs of reaction product was mixed with 25 µLs 2X Biotective Green Reagent and incubated for 5 minutes in the dark before detection using a Tecan Infinite M1000 Pro microplate reader with 495 nm (10nm bandwidth) excitation and 530 nm (10 nm bandwidth) emission. A standard curve of DTB was generated from 0 pmol to 80 pmol and used to determine the amount of DTB per sample.

Seedling growth for Gene Expression Analysis

Seedlings were germinated on control media and transferred to plates containing control or 200 µM cadaverine media for 24, 48 or 72 hours prior to collecting tissue at 8 days after germination. An additional subset of seedlings was germinated directly on control or 200 uM cadaverine media. To harvest, seedlings were flash frozen in liquid nitrogen and homogenized using a 4.5 mm steel bead and a Mixmill.

Gene Expression Analysis

RNA extraction was done using TRIzol reagent and Direct-zol RNA MiniPrep Kit, with a chloroform phase separation, according to manufacturer specifications (ZymoResearch). cDNA synthesis was done using SuperScript iii. qRT-PCR was done using Bullseye EvaGreen qPCR Mastermix (MidSci) and run on a LightCycler 480 II Instrument (Roche). Gene expression was calculated by delta (delta Ct) method and standardized to PP2A (At1G59830) reference gene (Ramakers *et al.*, 2003; Czechowski *et al.*, 2005). Fold-change of expression for all samples was then set relative to a control sample. Statistical significance was identified by ANOVA with Tukey's HSD correction (SAS version 9.3).

Protein Analysis

Seedling growth is as described above, with seedlings germinated on 1.5% agar, 0.5X Linsmaier and Skoog (LS) media pH 5.7 with 1.5% sucrose with or without or 200 μ M cadaverine. Tissue was frozen using dry ice after 8 days of growth. Frozen samples were homogenized using MixMill. Protein was extracted in the equivalent of 1 mL of protein extraction buffer (20mM MOPS (pH 7.5), 5 mM MgCl₂, 5 mM KCl, 50 mM Sucrose, Roche inhibitor cocktail tablet, 1 mM DTT, 7% glycerol (adapted from (Li *et al.*, 2012)) per 250 mg of tissue to. in. The homogenate was gently mixed at 4°C for 30 minutes, then centrifuged at 4°C for 30 minutes, 14,000 rpm to pellet the debris. Total protein was quantified and standardized for each gel using the Roche BCA assay. Approximately 150 μ g of protein was loaded onto a 12% Tris-Bis NuPAGE gel (Invitrogen) and run using the Invitrogen X-cell SureLock Mini-Cell Electrophoresis System. After electroblotting, the Immobilon-FL PVDF membrane was blocked for 1 hour at 4°C with a 5% milk/TBS solution (25 mM Tris-HCl, 140 mM NaCl, 3.0 mM KCl). Anti-RPT4A, a proteasome subunit-specific antibody generated in rabbit, was used as a loading control at 1:4,000 dilution. BIO3-BIO1 antibody was generated against guinea pig by Claude Alban and used at 1:25,000 (Cobessi *et al.*, 2012). BCCP1 and MCCA antibodies were generated against rabbit by Basil Nikolau and obtained from the Plant Antibody Facility (PAF, Ohio State) (Che, 2003; Li *et al.*, 2011). MCCA and BCCP1 antibodies were used at a concentration of 1:10,000. LiCor CW800, secondary antibodies were used according to manufacturer's instructions at 1:10,000 dilution. Membranes were

scanned using LICOR Odyssey. Image studio software Version 5.2.5 and Adobe Photoshop was used to quantify band intensity.

Biotin Containing Protein Analysis

Biotin containing proteins were quantified following protocol from Li et al., 2012. RPT4A loading control was used as described above. Membranes were blocked in TBST with 1% Tween20 prior to treating with Streptavidin. Streptavidin CW800 was used to detect biotinylated proteins at a concentration of 0.5 ug per mL in 1% milk TBST.

Lipid Profiling

Col-0 wild-type seedlings were grown on LS media as described above. After 5 days in the growth chamber, seedlings were transferred to plates containing control or 200 μ M cadaverine media, and allowed to grow for an additional 72 hours. Fresh tissue was submerged immediately in 75°C isopropanol with 0.01% butylated hydroxytoluene (BHT). Lipid extraction procedure was done according to (Devaiah et al., 2006). Samples were dried under nitrogen gas and shipped to Kansas State Lipidomics Research Center for analysis (Vu et al., 2014). Lipids were standardized to dry weight for each sample. Total acyl lipids, DAG and TAG were analyzed on a Waters Xevo TQS mass spectrometer. Samples contained 200 seedlings per biological replica, with a total of 5 biological repeats.

Accession Numbers

Sequence data from this article can be found in the EMBL/GenBank data libraries under accession numbers: BIO3-BIO1 (AT5G57590), BIO2 (AT2G43360), BIO4 (AT5G04620), MCCA (AT1G03090), BCCP1 (AT5G16390), PAO4 (AT1G65840), OCT1 (AT1G73220), PP2A (AT1G69960), RPT4A (AT5G43010).

Data Availability Statement

Data from this study can be provided by the corresponding author upon request.

Acknowledgements

This work was funded by HATCH award #WIS10338 from the College of Agriculture and Life Sciences, Fall Competition Award #MSN226688 from Office of the Vice Chancellor for

Research and Graduate Education, University of Wisconsin-Madison, and grant #80NSSC19K1483 from the National Aeronautics and Space Administration. We would like to thank David Meinke and Joe Shellhammer for technical advice regarding biotin quantification, Benjamin Minkoff and Thao Nygen for help troubleshooting protein purification, and David Cobessi for help with BIO3-BIO1 structure analysis. We are grateful for Sarah Swanson at the Newcomb Imaging Center, Department of Botany, UW-Madison, for confocal imaging. Tecan Infinite M1000 Pro data were obtained at the University of Wisconsin-Madison Biophysics Instrumentation Facility, which was established with support from the University of Wisconsin-Madison and grants BIR-9512577 (NSF) and S10RR13790 (NIH). The lipid analyses described in this work were performed at the Kansas Lipidomics Research Center Analytical Laboratory. Instrument acquisition and lipidomics method development was supported by National Science Foundation (EPS 0236913, MCB 1413036, DBI 0521587, DBI1228622), Kansas Technology Enterprise Corporation, K-IDeA Networks of Biomedical Research Excellence (INBRE) of National Institute of Health (P20GM103418), and Kansas State University.

Author Contributions

Designed research, N.M.G., H.A.M. and P.H.M.; Performed research, N.M.G., S-H.S., S.L-N., S.M., C.A.; Analyzed data, N.M.G., S-H.S., S.L-N., H.A.M., and P.H.M.; Wrote paper, N.M.G. and P.H.M.

Declaration of Interests

The authors declare no competing interests.

Supporting Information

Figure S1: Thirteen cadaverine hypersensitive mutants (*cdh1* to *cdh13*) were identified by screening for alterations in root responses to cadaverine.

Figure S2: Cadaverine decreases cell elongation.

Figure S3: The expression of genes involved in biotin synthesis is unaffected by cadaverine treatment.

Figure S4: *MCCA* and *BCCP1* gene expression is unaffected by cadaverine treatment.

Table S1: Lipid Analysis of 72-hour cadaverine treated seedlings.

References

- Alban, C., Baldet, P., Axiotis, S. and Douce, R.** (1993) Purification and Characterization of 3-Methylcrotonyl-Coenzyme A Carboxylase from Higher Plant Mitochondria. *Plant Physiology*, **102**, 957–965.
- Alban, C., Job, D. and Douce, R.** (2000) BIOTIN METABOLISM IN PLANTS. *Annual Review of Plant Physiology and Plant Molecular Biology*, **51**, 17–47.
- Aubert, S., Alban, C., Bligny, R. and Douce, R.** (1996) Induction of beta-methylcrotonyl-coenzyme A carboxylase in higher plant cells during carbohydrate starvation: evidence for a role of MCCase in leucine catabolism. *FEBS Letters*, **383**, 175–180.
- Aziz, A., Martin-Tanguy, J. and Larher, F.** (1998) Stress-induced changes in polyamine and tyramine levels can regulate proline accumulation in tomato leaf discs treated with sodium chloride. *Physiologia plantarum*, **104**, 195–202. Available at: https://www.researchgate.net/profile/Aziz_Aziz3/publication/225274396_Stress-induced_changes_in_polyamine_and_tyramine_levels_can_regulate_proline_accumulation_in_tomato_leaf_discs_treated_with_sodium_chloride_Physiol_Plant/links/0fcfd51152e5d86d26000000.pdf.
- Batchelor, R., Sarkez, A., Cox, W.G. and Johnson, I.** (2007) Fluorometric assay for quantitation of biotin covalently attached to proteins and nucleic acids. *BioTechniques*, **43**, 503–507. Available at: https://www.biotechniques.com/multimedia/archive/00001/BTN_A_000112564_O_1354a.pdf.
- Bunsupa, S., Katayama, K., Ikeura, E., Oikawa, A., Toyooka, K., Saito, K. and Yamazaki, M.** (2012) Lysine Decarboxylase Catalyzes the First Step of Quinolizidine Alkaloid Biosynthesis and Coevolved with Alkaloid Production in Leguminosae. *The Plant Cell*, **24**, 1202–1216.

Cassán, F. (2009) Cadaverine production by *Azospirillum brasilense* and its possible role in plant growth promotion and osmotic stress mitigation. *European Journal of Soil Biology*, **45**, 12–19.

Che, P. (2003) The Role of Biotin in Regulating 3-Methylcrotonyl-Coenzyme A Carboxylase Expression in *Arabidopsis*. *Plant Physiology*, **131**, 1479–1486.

Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant journal : for cell and molecular biology*, **16**, 735–743.

Cobessi, D., Dumas, R., Pautre, V., Meinguet, C., Ferrer, J.-L. and Alban, C. (2012) Biochemical and Structural Characterization of the *Arabidopsis* Bifunctional Enzyme Dethiobiotin Synthetase–Diaminopelargonic Acid Aminotransferase: Evidence for Substrate Channeling in Biotin Synthesis. *The Plant Cell*, **24**, 1608–1625.

Curtis, M.D. (2003) A Gateway Cloning Vector Set for High-Throughput Functional Analysis of Genes in *Planta*. *Plant Physiology*, **133**, 462–469. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC523872/pdf/1330462.pdf>.

Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K. and Scheible, W.-R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiology*, **139**, 5–17.

Devaiah, S.P., Roth, M.R., Baughman, E., Li, M., Tamura, P., Jeannotte, R., Welti, R. and Wang, X. (2006) Quantitative profiling of polar glycerolipid species from organs of wild-type *Arabidopsis* and a PHOSPHOLIPASE $\text{D}\alpha 1$ knockout mutant. *Phytochemistry*, **67**, 1907–1924.

Ding, G., Che, P., Ilarslan, H., Wurtele, E.S. and Nikolau, B.J. (2012) Genetic dissection of methylcrotonyl CoA carboxylase indicates a complex role for mitochondrial leucine catabolism during seed development and germination. *The Plant Journal*, **70**, 562–577. Available at: <http://onlinelibrary.wiley.com/store/10.1111/j.1365-313X.2011.04893.x/asset/j.1365-313X.2011.04893.x.pdf?v=1&t=jdkya03t&s=40aee41d5766d399aa4e720305ad6f85dafc7a4f>.

Gamarnik, A.A. and Frydman, R.B.R. (1991) Cadaverine, an Essential Diamine for the Normal Root Development of Germinating Soybean (*Glycine max*) Seeds. *Plant Physiology*, **97**, 778–785.

Guan, X., Diez, T., Prasad, T.K., Nikolau, B.J. and Wurtele, E.S. (1999) Geranoyl-CoA Carboxylase: A Novel Biotin-Containing Enzyme in Plants. *Arch Biochem Biophys*, **362**, 12–21.

Gueorguieva, R. and Krystal, J.H. (2004) Move Over ANOVA: Progress in Analyzing Repeated-Measures Data and Its Reflection in Papers Published in the Archives of General Psychiatry. *Arch Gen Psychiat*, **61**, 310.

Imai, A., Matsuyama, T., Hanzawa, Y., et al. (2004) Spermidine Synthase Genes Are Essential for Survival of Arabidopsis. *Plant Physiol*, **135**, 1565–1573.

Jancewicz, A.L., Gibbs, N.M. and Masson, P.H. (2016) Cadaverine's Functional Role in Plant Development and Environmental Response. *Front Plant Sci*, **7**, 870.

Takehi, J.I., Kuwashiro, Y., Niitsu, M. and Takahashi, T. (2008) Thermospermine is Required for Stem Elongation in Arabidopsis thaliana. *Plant and Cell Physiology*, **49**, 1342–1349.

Kamada-Nobusada, T., Hayashi, M., Fukazawa, M., Sakakibara, H. and Nishimura, M. (2008) A Putative Peroxisomal Polyamine Oxidase, AtPAO4, is Involved in Polyamine Catabolism in Arabidopsis thaliana. *Plant Cell Physiol*, **49**, 1272–1282.

Kuznetsov, V., Shorina, M., Aronova, E., Stetsenko, L., Rakitin, V. and Shevyakova, N. (2007) NaCl- and ethylene-dependent cadaverine accumulation and its possible protective role in the adaptation of the common ice plant to salt stress. *Plant Science*, **172**, 363–370.

Lawand, S., Dorne, A.-J., Long, D., Coupland, G., Mache, R. and Carol, P. (2002) Arabidopsis A BOUT DE SOUFFLE, Which Is Homologous with Mammalian Carnitine Acyl Carrier, Is Required for Postembryonic Growth in the Light. *Plant Cell*, **14**, 2161–2173.

- Lelandais-Brière, C., Jovanovic, M., Torres, G.A.M., Perrin, Y., Lemoine, R., Corre-Menguy, F. and Hartmann, C.** (2007) Disruption of AtOCT1, an organic cation transporter gene, affects root development and carnitine-related responses in Arabidopsis. *The Plant Journal*, **51**, 154–164.
- Li, J., Brader, G., Helenius, E., Kariola, T. and Palva, E.T.** (2012) Biotin deficiency causes spontaneous cell death and activation of defense signaling. *The Plant Journal*, **70**, 315–326.
- Li, X., Ilarslan, H., Brachova, L., Qian, H.-R., Li, L., Che, P., Wurtele, E.S. and Nikolau, B.J.** (2011) Reverse-genetic analysis of the two biotin-containing subunit genes of the heteromeric acetyl-coenzyme A carboxylase in Arabidopsis indicates a unidirectional functional redundancy. *Plant Physiology*, **155**, 293–314.
- Liu, K., Fu, H., Bei, Q. and Luan, S.** (2000) Inward potassium channel in guard cells as a target for polyamine regulation of stomatal movements. *Plant Physiology*, **124**, 1315–1326.
- Liu, T., Dobashi, H., Kim, D.W., Sagor, G.H.M., Niitsu, M., Berberich, T. and Kusano, T.** (2014) Arabidopsis mutant plants with diverse defects in polyamine metabolism show unequal sensitivity to exogenous cadaverine probably based on their spermine content. *Physiology and molecular biology of plants : an international journal of functional plant biology*, **20**, 151–159.
- Muralla, R., Chen, E., Sweeney, C., Gray, J.A., Dickerman, A., Nikolau, B.J. and Meinke, D.** (2007) A Bifunctional Locus (BIO3-BIO1) Required for Biotin Biosynthesis in Arabidopsis. *Plant Physiology*, **146**, 60–73.
- Negrel, J., Paynot, M. and Javelle, F.** (1992) Purification and Properties of Putrescine Hydroxycinnamoyl Transferase from Tobacco (*Nicotiana tabacum*) Cell Suspensions. *Plant Physiology*, **98**, 1264–1269.
- Niemi, K., Häggman, H. and Sarjala, T.** (2002) Effects of exogenous diamines on the interaction between ectomycorrhizal fungi and adventitious root formation in Scots pine in vitro. *Tree Physiology*, 1–9. Available at: <http://treephys.oxfordjournals.org/content/22/6/373.full.pdf>.

Nikolau, B.J., Ohlrogge, J.B. and Wurtele, E.S. (2003) Plant biotin-containing carboxylases. *Archives of biochemistry and biophysics*, **414**, 211–222.

Nudelman, Ayelet, Marcovici-Mizrahi, D., Nudelman, Abraham, Flint, D. and Wittenbach, V. (2004) Inhibitors of biotin biosynthesis as potential herbicides. *Tetrahedron*, **60**, 1731–1748.

Park, M.H., Cooper, H.L. and Folk, J.E. (1982) The biosynthesis of protein-bound hypusine (N epsilon -(4-amino-2-hydroxybutyl)lysine). Lysine as the amino acid precursor and the intermediate role of deoxyhypusine (N epsilon -(4-aminobutyl)lysine). *J Biological Chem*, **257**, 7217–22.

Patton, D.A., Volrath, S. and Ward, E.R. (1996) Complementation of an Arabidopsis thaliana biotin auxotroph with an Escherichia coli biotin biosynthetic gene. *Molecular & general genetics : MGG*, **251**, 261–266.

Perilli, S., Biology, S.S.P.D. and 2010 (2010) Analysis of root meristem size development. *Springer*, **655**, 177–187.

Piccocchi, A., Douce, R. and Alban, C. (2001) Biochemical Characterization of the Arabidopsis Biotin Synthase Reaction. The Importance of Mitochondria in Biotin Synthesis. *Plant Physiology*, **127**, 1224–1233. Available at: <http://www.plantphysiol.org/content/plantphysiol/127/3/1224.full.pdf>.

Pinon, V. (2005) Biotin Synthesis in Plants. The First Committed Step of the Pathway Is Catalyzed by a Cytosolic 7-Keto-8-Aminopelargonic Acid Synthase. *Plant Physiology*, **139**, 1666–1676.

Pommerrenig, B., Popko, J., Heilmann, M., Schulmeister, S., Dietel, K., Schmitt, B., Stadler, R., Feussner, I. and Sauer, N. (2012) SUCROSE TRANSPORTER 5 supplies Arabidopsis embryos with biotin and affects triacylglycerol accumulation. *The Plant Journal*, **73**, 392–404.

Ramakers, C., Ruijter, J.M., Deprez, R.H.L. and Moorman, A.F.M. (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neuroscience Letters*, **339**, 62–66. Available at: <https://pdf.sciencedirectassets.com/271072/1-s2.0-S0304394000X08555/1->

s2.0-S0304394002014234/main.pdf?x-amz-security-

token=AgoJb3JpZ2luX2VjEGYyaCXVzLWVhc3QtMSJGMEQCIFFINRf34MawRzYSCeQyswzut775Ti4tLz2LfTm84nkOAiBuTifBjO53kAHUtOvuTk41dQsktCXF5s617BidgIsalCraAwgfEAIaDDA1OTAwMzU0Njg2NSIMqwiIC1qPnlMny1ZyKrcD9IS%2FoXpTuGKrPJXa%2B5UfjJ%2BU798fBMrmStTBQcdreISW%2FRpRR%2F720hDKSYKWz1r6EQNdEoR4USU0SnKP23LatQL6e3M7fsORwdH9xCIUPduJSN6rHj5mcEOroLJG4OFZu2Lxd4qwgDDn4E6%2FqWDe7nC1QTMcW7Jh%2FJC2azM4p2u0HTidjWdZJ1yHdgLPcBX%2BchJXKmgp%2BfuW2vazx50BXs%2FWfpNwDZPNwVNFk3oAnMqJ8CXgGqBZEDufGHEvKKD%2BSehSkmW5yRX53ArQVzPYs%2FId0%2FRXbbqCcB%2FcdskrFxxlBAj9imZibdtVQmikR%2BL4EwLIyBLPIovbI4z%2BIw2f29IP2Ep5AIIATHUsgVzxXD1NdDdUdLavGni4Cfu7DPfpb1dWa%2FTY8UfSzu24e0RN7zhgAIU1bycNHwYVGKsl9cdQQ%2BzfgobKMmU6lx0wGt%2F0QJtati9qAqFzuuIhuu1O33bAD9L%2FvXrZMVp%2Bmccyy%2FS0BH7S888IWFLPAd2z47jKQmzExLGVYg9zdNwO1r3qP1ogB%2Fge72YbR1P1J3Pr1%2BTkKYZeQRaF3cCBvOmWTa9rmbwIN0EWezDg1%2B%2FkBTq1AfNwlxVYJiLaKmS%2Bs9nlbliYNcaH6CzRMSkOmmcOt6Vq9Zo.

Rutherford, R. and Masson, P.H. (1996) Arabidopsis thaliana sku mutant seedlings show exaggerated surface-dependent alteration in root growth vector. *Plant Physiology*, **111**, 987–998.

Sandmark, J., Mann, S., Marquet, A. and Schneider, G. (2002) Structural Basis for the Inhibition of the Biosynthesis of Biotin by the Antibiotic Amiclenomycin. *J Biol Chem*, **277**, 43352–43358.

Sasaki, Y., Hakamada, K., Suama, Y., Nagano, Y., Furusawa, I. and Matsuno, R. (1993) Chloroplast-encoded protein as a subunit of acetyl-CoA carboxylase in pea plant. *J Biological Chem*, **268**, 25118–23.

Schneeberger, K., Ossowski, S., Lanz, C., Juul, T., Petersen, A.H., Nielsen, K.L., Jørgensen, J.-E., Weigel, D. and Andersen, S.U. (2009) SHOREmap: simultaneous mapping and mutation identification by deep sequencing. *Nature Publishing Group*, **6**, 550–551.

Schneider, T., Dinkins, R., Robinson, K., Shellhammer, J. and Meinke, D.W. (1989) An embryolethal mutant of Arabidopsis thaliana is a biotin auxotroph. *Developmental biology*, **131**, 161–167.

Sequera-Mutiozabal, M.I., Erban, A., Kopka, J., Atanasov, K.E., Bastida, J., Fotopoulos, V., Alcázar, R. and Tiburcio, A.F. (2016) Global Metabolic Profiling of Arabidopsis Polyamine Oxidase 4 (AtPAO4) Loss-of-Function Mutants Exhibiting Delayed Dark-Induced Senescence. *Frontiers in plant science*, **7**, 1237–13. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4757743/pdf/fpls-07-00173.pdf>.

Shellhammer, J. and Meinke, D. (1990) Arrested Embryos from the bio1 Auxotroph of Arabidopsis thaliana Contain Reduced Levels of Biotin. *Plant Physiology*, **93**, 1162–1167.

Shevyakova, N.I., Rakitin, V.Y., Duong, D.B., Science, N.S.P. and 2001 (2001) Heat shock-induced cadaverine accumulation and translocation throughout the plant. *Elsevier*.

Shimizu, Y., Rai, A., Okawa, Y., Tomatsu, H., Sato, M., Kera, K., Suzuki, H., Saito, K. and Yamazaki, M. (2019) Metabolic diversification of nitrogen-containing metabolites by the expression of a heterologous lysine decarboxylase gene in Arabidopsis. *Plant J*, **100**, 505–521.

Shiva, S., Vu, H.S., Roth, M.R., Zhou, Z., Marepally, S.R., Nune, D.S., Lushington, G.H., Visvanathan, M. and Welti, R. (2013) Lipidomic analysis of plant membrane lipids by direct infusion tandem mass spectrometry. *Methods Mol Biology Clifton N J*, **1009**, 79–91.

Simon-Sarkadi, L., Ludidi, N. and Kocsy, G. (2014) Modification of cadaverine content by NO in salt-stressed maize. *Plant signaling & behavior*, **9**, e27598.

Song, J., Wurtele, E.S. and Nikolau, B.J. (1994) Molecular cloning and characterization of the cDNA coding for the biotin-containing subunit of 3-methylcrotonoyl-CoA carboxylase: identification of the biotin carboxylase and biotin-carrier domains. *Proceedings of the National Academy of Sciences*, **91**, 5779–5783.

Strohm, A.K., Vaughn, L.M. and Masson, P.H. (2015) Natural variation in the expression of ORGANIC CATION TRANSPORTER 1 affects root length responses to cadaverine in Arabidopsis. *Journal of Experimental Botany*, **66**, 853–862. Available at: <http://jxb.oxfordjournals.org/content/66/3/853>.

Sziderics, A.H., Oufir, M., Trognitz, F., Kopecky, D., Matušíková, I., Hausman, J.-F. and Wilhelm, E. (2010) Organ-specific defence strategies of pepper (*Capsicum annuum* L.) during early phase of water deficit. *Plant Cell Reports*, **29**, 295–305.

Tanabe, Y., Maruyama, J., Yamaoka, S., Yahagi, D., Matsuo, I., Tsutsumi, N. and Kitamoto, K. (2011) Peroxisomes are involved in biotin biosynthesis in *Aspergillus* and *Arabidopsis*. *Journal of Biological Chemistry*, **286**, 30455–30461.

Thelen, J.J., Mekhedov, S. and Ohlrogge, J.B. (2001) Brassicaceae express multiple isoforms of biotin carboxyl carrier protein in a tissue-specific manner. *Plant Physiology*, **125**, 2016–2028.

Tomar, P.C., Lakra, N. and Mishra, S.N. (2013) Effect of cadaverine on *Brassica juncea* (L.) under multiple stress. *Indian journal of experimental biology*, **51**, 758–763.

Torrigiani, P., Serafini-Fracassini, D. and Bagni, N. (1987) Polyamine Biosynthesis and Effect of Dicyclohexylamine during the Cell Cycle of *Helianthus tuberosus* Tuber. *Plant Physiol*, **84**, 148–152.

Vu, H.S., Shiva, S., Roth, M.R., et al. (2014) Lipid changes after leaf wounding in *Arabidopsis thaliana*: expanded lipidomic data form the basis for lipid co-occurrence analysis. *The Plant Journal*, **80**, 728–743.

Weigel, D.G. and Glazebrook, J. (2002) *Arabidopsis: A Laboratory Manual*,.

Wu, J., Sun, Y., Zhao, Y., et al. (2015) Deficient plastidic fatty acid synthesis triggers cell death by modulating mitochondrial reactive oxygen species. *Nature Publishing Group*, **25**, 621–633.

Wurtele, E.S. and Nikolau, B.J. (1990) Plants contain multiple biotin enzymes: discovery of 3-methylcrotonyl-CoA carboxylase, propionyl-CoA carboxylase and pyruvate carboxylase in the plant kingdom. *Archives of biochemistry and biophysics*, **278**, 179–186.

Accepted Article
Yanai, Y., Kawasaki, T., Shimada, H., Wurtele, E.S., Nikolau, B.J. and Ichikawa, N. (1995)

Genomic Organization of 251 kDa Acetyl-CoA Carboxylase Genes in Arabidopsis: Tandem Gene Duplication has Made Two Differentially Expressed Isozymes. *Plant Cell Physiol*, **36**, 779–787.

Zuryn, S., Gras, S.L., Jamet, K. and Jarriault, S. (2010) A Strategy for Direct Mapping and Identification of Mutations by Whole-Genome Sequencing. *Genetics*, **186**, 427–430.

Figures Legends

Figure 1: *cdh3* is hypersensitive to cadaverine. (a) 10 day old Col-0 and *cdh3* seedlings germinated on vertical plates containing 200 μ M cadaverine or control media. **(b)** Dose

Accepted Article

response of seedlings germinated on cadaverine media. Root growth between day 6 to day 10 is quantified. Root growth on control media (Average +/- standard error): Col 2.29 cm +/- 0.045 cm; *cdh3* 1.94 cm +/- 0.096 cm. Whiskers represent minimum to maximum values. **(c)** Seedlings were germinated on 200 μ M cadaverine or control media and treated with propidium iodide to stain the cell walls. Stained roots were then analyzed using confocal microscopy. Five seedlings were analyzed per treatment group. White stars indicate the quiescent center. White arrowheads indicate the end of the root meristem. **(d)** Length of the meristem was quantified by identifying cells in the cortex whose length is greater than the width, and measuring the distance from the distal wall this cell to the distal wall of the quiescent center. Bars represent standard error, with dots showing data points. **(e)** The numbers of cells in the meristems were counted from the quiescent center to the end of the meristem along the cortex. Bars represent standard error, with dots showing data points. **(f)** Root tip growth following transfer of 5 day old seedlings from control to split agar plates containing either control media, or media supplemented with 200 μ M cadaverine. Root tip growth was recorded using a high-resolution camera for 96 hours and quantified using Image J. Data points represent the three seedlings per treatment. Bars indicate standard error. * indicates p-value less than 0.05, ** indicates p-value less than 0.01, *** indicates p-value less than 0.001 analyzed by a mixed effects model with an F-test for the interaction between time and treatment with Tukey's adjustment for multiple comparisons. **(g)** *cdh3* and wild type root growths were measured from day 6 to day 10 on control or media supplemented with the indicated concentrations of putrescine, spermidine and spermine. Root growths were standardized to growth on control media for each genotype. Whiskers show minimum to maximum values. ANOVA with Tukey's HSD correction was used to determine significance.

Figure 2: *cdh3* contains a mutation in *BIO3-BIO1*. (a) Next generation sequencing revealed a peak on chromosome 5 with a high proportion of SNPs compared to the reference sequence. The table in Figure 2(a) shows mutations within the 4.6 Mb peak region containing nonsynonymous SNPs in protein coding genes. (b) The gene structure for *BIO3-BIO1* is shown indicating the position of the SNP in *cdh3*. (c) 3D structure of the catalytic site of *BIO1*, deduced from the crystal structure resolved in (Cobessi et al., 2012). This structure, imaged with PyMOL software, shows the substrate (KAPA, in pink) and the cofactor (PLP, in blue) bound to L666. The position of Ala662, which is converted into a threonine in *cdh3*, is indicated in yellow. (d) Quantification of standardized root growth on control and 200 μ M cadaverine-containing media from day 6 to day 10 of wild type, *cdh3*, or *cdh3* transformation rescue lines (*cdh3[BIO3-BIO1pro:BIO3-BIO1]*) with the wild type *BIO3-BIO1* transgene under the control of its native promoter. *BIO3-BIO1* rescue lines are at the T3 generation. Root growth on control media (Average +/- standard error): Col 2.54 cm +/- 0.045; *cdh3* 2.05 cm +/- 0.073 cm; *cdh3[BIO3-BIO1pro:BIO3-BIO1 1* 2.73 cm +/- 0.075; *cdh3[BIO3-BIO1pro:BIO3-BIO1 2* 2.37 cm +/- 0.104 cm; *cdh3[BIO3-BIO1pro:BIO3-BIO1 3* 2.36 cm +/- 0.063 cm; *cdh3[BIO3-BIO1pro:BIO3-BIO1 4* 2.53 cm +/- 0.073 cm. Whiskers represent minimum to maximum. (e) qRT-PCR analysis of *BIO3-BIO1* in 8 day old wild type, *cdh3* and *cdh3[BIO3-BIO1pro:BIO3-BIO1]* seedlings grown on control media. Bars represent standard error, with dots showing biological replicates. In (d) and (e), significant differences between

groups detected using ANOVA with Tukey's HSD correction ($p < 0.05$), are indicated with distinct letters.

Figure 3: Biotin suppresses wild-type primary root growth response to cadaverine and rescues *cdh3*'s hypersensitive root growth response. (a) Standardized root growth of wild type and *cdh3* mutant seedlings on 200 μM cadaverine-containing media in the presence of 0 nM to 100 nM biotin. *** indicates p-value less than 0.001 using Student's T-test. Root growth on control media (Average \pm standard error): Col-0 3.505 cm \pm 0.0925 cm ; *cdh3* 2.605 cm \pm 0.326 cm. Whiskers show minimum to maximum values. (b) Wild type and *cdh3* mutant root growth from day 6 to day 10 on media with or without 200 μM cadaverine, in the presence of 1 μM biotin, KAPA or DAPA. A simplified biotin synthesis pathway is shown on the right of this graph. Root growth on control media (Average \pm standard error): Col-0 3.452 cm \pm 0.582 cm; *cdh3* 2.985 cm \pm 0.115 cm. Minimum to maximum values are represented by whiskers. (c) Root growth of *oct1* mutants on media without or with 200 μM cadaverine and 1 μM biotin. *oct1-1* is in the WS accession background, and *oct1-2* is in the Col-0 background. Root growth on control media (Average \pm standard error): WS 1.663 cm \pm 0.0322 cm; *oct1-1* 1.569 cm \pm 0.066 cm; Col-0 1.523


cm +/- 0.174 cm; *oct1-2* 1.44 cm +/- 0.033 cm. Whiskers show minimum to maximum values. (d) Standardized root growth of wild type Col-0 and *pao4-1* and *pao4-2* mutant seedlings on media with or without 200 μ M cadaverine, in the presence or absence of 1 μ M biotin. Root growth on control media (Average +/- standard error): Col 1.938 cm +/- 0.457 cm; *pao4-1* 2.230 cm +/-0.046 cm; *pao4-2* 2.361 cm +/- 0.043 cm. In panels (b)-(d), significance was determined using ANOVA with Tukey's HSD correction. Whiskers show minimum to maximum values.

Figure 4: Cadaverine inhibits BIO3-BIO1 enzymatic activity. (a) BIO3-BIO1 enzyme was expressed in *E. coli* and affinity-purified. *In vitro* enzymatic assays contained 50 μ M cadaverine and 20 μ M KAPA. A negative control assay was carried out with 0 μ M KAPA with or without 50 μ M cadaverine, to control for background fluorescence (see experimental procedures). Reactions were run for 8 minutes before heat-treating to stop the reaction. * indicates a p-value less than 0.05 based on Student's T-test. Bars represent standard error, with individual data points showing biological replicates. (b) *BIO3-BIO1* expression is unaltered by cadaverine treatment. *BIO3-BIO1* expression was quantified using qRT-PCR


on cDNA prepared from RNA extracted from 8 day old seedlings germinated on 200 μ M cadaverine or control media, or transferred to cadaverine for 24, 48 or 72 hours. Expression is standardized to *PP2A* reference gene. No statistically significant differences were observed using ANOVA. Bars represent standard error with individual data points showing biological replicates. **(c)** BIO3-BIO1 protein expression was quantified using a western-blot approach. 8 day old seedlings were grown on cadaverine-containing media for 72 hours. Proteins were extracted from whole seedlings and run on a SDS-PAGE gel. Following electro-transfer, the PVDF membrane was exposed to a BIO3-BIO1-specific antibody to determine protein level. An antibody directed against a proteasomal protein, RPT4A was also used as a loading control. The blot shown here is representative of three biological replicates that gave similar results. In each blot, band intensities were quantified and standardized to RPT4A for each lane. The results of these quantifications are plotted in the graph shown to the right of the blot, with data points showing biological replicates. Bars indicate standard error. ** indicates p-value less than 0.01 using Student's T-test against Col-0 on control condition.

Figure 5: The levels of biotinylated proteins are altered following cadaverine treatment. (a,b): Seed were germinated on control or cadaverine-containing media. Following 8 days of treatment, roots **(a)** and shoots **(b)** were dissected and protein was extracted. 150 µg of protein were loaded on SDS polyacrylamide gel and electrophoresed. After electro-transfer onto a PVDF membrane, biotinylated proteins were detected with Licor IRDye CW800-labeled streptavidin or anti-BCCP1 and MCCA antibodies. In all cases, anti-RPT4A antibodies were used to control for loading differences between samples. The blots shown in panels **(a)** and **(b)** are representative of three biological replicates showing similar results. Quantifications of band intensities from these blots are summarized in panels **(c-f)** for all three biological replicates. **(c,d,e,f):** Protein amount and biotinylation were quantified based on band intensity, standardized to the RPT4A signals, and then standardized again to Col-0. The expression results shown in the graphs represent three biological replicates, with dots indicating the value of each replicate. Bars indicate standard error. * indicates a Student's T-test p-value below 0.05. ** indicates a p-value below 0.01, compared to Col-0 control treatment. **(c)** MCCA total protein and MCCA biotinylation in roots. **(d)** MCCA total protein and biotinylation in shoots. **(e)** BCCP1 total protein and BCCP1 biotinylation in roots. **(f)** BCCP1 total protein and BCCP1 biotinylation in shoots.


Figure 6: TAG content following 72 hours of cadaverine treatment. Col-0 seedlings were germinated on control media and transferred to either control or 200 μ M cadaverine-containing media after 5 days. 72 hours post-treatment, whole seedlings were harvested and used for lipid extraction and analysis using MS/MS. The average of five biological replicates are shown containing approximately 200 seedlings per replicate, with dots representing each replicate. Bars indicate standard error. Student's T-test was done to determine significance between cadaverine treatment and control. * indicates a p-value below 0.05. ** indicates a p-value below 0.01.




tpj_15417_f1.png




tpj_15417_f2.png




tpj_15417_f3.png



tpj_15417_f4.png



tpj_15417_f5.png



tpj_15417_f6.png