



**HAL**  
open science

## The climatic debt is growing in the understorey of temperate forests: Stand characteristics matter

Benoit Richard, Jean-luc Dupouey, Emmanuel Corcket, Didier Alard, Frédéric Archaux, Michaël Aubert, Vincent Boulanger, François Gillet, Estelle Langlois, Sébastien Macé, et al.

### ► To cite this version:

Benoit Richard, Jean-luc Dupouey, Emmanuel Corcket, Didier Alard, Frédéric Archaux, et al.. The climatic debt is growing in the understorey of temperate forests: Stand characteristics matter. *Global Ecology and Biogeography*, 2021, 30 (7), pp.1474-1487. 10.1111/geb.13312 . hal-03293479

**HAL Id: hal-03293479**

**<https://hal.science/hal-03293479v1>**

Submitted on 21 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 The climatic debt is growing in the understory of 2 temperate forests: **stand characteristics matter**

## 3 *Running title:*

- 4 • **Explaining** the climatic debt of temperate forests

## 5 **Authors**

6 Benoit Richard<sup>1,2</sup> | Jean-Luc Dupouey<sup>3</sup> | Emmanuel Corcket<sup>4</sup> | Didier Alard<sup>4</sup> |  
7 Frédéric Archaux<sup>5</sup> | Michaël Aubert<sup>2</sup> | Vincent Boulanger<sup>6</sup> | François Gillet<sup>7</sup> |  
8 Estelle Langlois<sup>2</sup> | Sébastien Macé<sup>6</sup> | Pierre Montpied<sup>3</sup> | Thérèse Beaufile<sup>7</sup> | Carole  
9 Begeot<sup>7</sup> | Patrick Behr<sup>3</sup> | Jean-Michel Boissier<sup>8</sup> | Sylvaine Camaret<sup>9</sup> | Richard  
10 Chevalier<sup>5</sup> | Guillaume Decocq<sup>1</sup> | Yann Dumas<sup>5</sup> | Richard Eynard-Machet<sup>10</sup> | Jean-  
11 Claude Gégout<sup>3</sup> | Sandrine Huet<sup>11</sup> | Valéry Malécot<sup>11</sup> | Pierre Margerie<sup>2</sup> | Arnaud  
12 Mouly<sup>7</sup> | Thierry Paul<sup>3</sup> | Benoît Renaux<sup>12</sup> | Pascale Ruffaldi<sup>7</sup> | Fabien Spicher<sup>1</sup> |  
13 Erwin Thirion<sup>3</sup> | Erwin Ulrich<sup>6</sup> | Manuel Nicolas<sup>6</sup> | Jonathan Lenoir<sup>1\*</sup>

## 14 **Authors' affiliations**

15 <sup>1</sup>UMR CNRS 7058 « Ecologie et Dynamique des Systèmes Anthropisés » (EDYSAN),  
16 Université de Picardie Jules Verne, Amiens, France

17 <sup>2</sup>Normandie Université, UNIROUEN, INRAE, ECODIV, Rouen, France

18 <sup>3</sup>Université de Lorraine, AgroParisTech, INRAE, SILVA, Nancy, France

19 <sup>4</sup>Université de Bordeaux, INRAE, BIOGECO, Pessac, France

20 <sup>5</sup>INRAE, UR EFNO, Domaine des Barres, Nogent-sur-Vernisson, France

21 <sup>6</sup>Office National des Forêts, Département RDI, Fontainebleau, France

22 <sup>7</sup>UMR CNRS 6249 Chrono-Environnement, Université Bourgogne Franche-Comté,  
23 Besançon, France

24 <sup>8</sup>Ecosylve, Les Déserts, France

25 <sup>9</sup>UFR LLSH, Université Savoie Mont Blanc, Domaine Universitaire de Jacob-  
26 Bellecombette, Chambéry, France

27 <sup>10</sup>e-Sciences, Cognin, France

28 <sup>11</sup>Agrocampus-Ouest, INRAE, Université d'Angers, IRHS, Beaucouzé, France

29 <sup>12</sup>Conservatoire botanique national du Massif central, Chavaniac-Lafayette, France

# The climatic debt is growing in the understory of temperate forests: **stand characteristics matter**

## *Running title:*

- **Explaining** the climatic debt of temperate forests

## **Abstract**

### **Aim**

Climate warming reshuffles biological assemblages towards less cold-adapted but more warm-adapted species, a process coined thermophilization. However, the velocity at which this process happens is generally lagging behind the velocity of climate change, generating a climatic debt which temporal dynamic remains misunderstood. Relying on high-resolution time series of vegetation data from a long-term monitoring network of permanent forest plots, we aim at quantifying the temporal dynamic – up to a yearly resolution – of the climatic debt in the understory of temperate forests before identifying the key determinants allowing to modulate it.

### **Location**

France

### **Time period**

1995-2017

### **Taxa studied**

Vascular plants

## 21 **Methods**

22 We used the community temperature index (CTI) to produce a time series of  
23 understory plant community thermophilization, which we subsequently compared to  
24 a time series of mean annual temperature changes over the same period and for the  
25 same sites. The direction and magnitude of the difference (i.e. the climatic debt) was  
26 finally analyzed using linear mixed-effect models to assess the relative contribution  
27 of abiotic and biotic determinants, including forest **stand characteristics**.

## 28 **Results**

29 We found a significant increase in CTI values over time (0.08–0.09°C/decade),  
30 **whereas** the velocity of mean annual temperature changes was three times higher  
31 over the same period (0.22–0.28°C/decade). Hence, the climatic debt increased over  
32 time and was greater in forest stands with higher basal area or older trees **as well as**  
33 **under warmer macroclimate**. By contrasts, a greater frequency of anthropogenic  
34 disturbances decreased the climatic debt, **while natural disturbances and herbivory**  
35 **had no impact**.

## 36 **Conclusions**

37 Although often overlooked in understanding the climatic debt of forest biodiversity,  
38 changes in forest stand characteristics may modulate the climatic debt by locally  
39 modifying microclimatic conditions. **Notably, the buffering effect** of the upper canopy  
40 layer **implies microclimate dynamics that** may provide more time for understory  
41 plant communities to locally adapt.

1  
2  
3 42 *Keywords*  
4  
5

6 43 Biodiversity, climate change, climatic debt, anthropogenic disturbances, long-term  
7  
8  
9 44 monitoring, plant community, thermophilization  
10  
11

12 45  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## 46 Introduction

47 Climate change induces a global redistribution of biodiversity at various spatial  
48 scales, with profound consequences on ecosystem functioning and the provision of  
49 ecosystem services (Hooper *et al.*, 2012; Pecl *et al.*, 2017). Typical biodiversity  
50 redistribution consists of species shifting their geographical range poleward in  
51 latitude and/or upslope in elevation (Lenoir & Svenning, 2015), tracking shifting  
52 isotherms through colonization and local extinction processes (Jump & Peñuelas,  
53 2005; Dullinger *et al.*, 2012). **Locally, changes in species' abundance and population  
54 dynamics, preceding species range shifts, are leading to important re-assemblages in  
55 community composition** (Hillebrand *et al.*, 2010; Walther, 2010). For example, some  
56 studies reported that local species assemblages are increasingly dominated by  
57 warm-adapted species (i.e. species from lower elevations and/or latitudes) at the  
58 expense of cold-adapted species (i.e. species from higher elevations and/or latitudes),  
59 a process referred to as “thermophilization” of communities (Gottfried *et al.*, 2012;  
60 De Frenne *et al.*, 2013; Zellweger *et al.*, 2020). However, species and communities  
61 may respond asynchronously and most likely with time lags relative to the forcing  
62 environmental changes, so that plant species assemblages shifting from one  
63 equilibrium state to another may not reflect environmental conditions prevailing at  
64 the time of the survey but may rather reflect former environmental conditions  
65 (Davis, 1984, 1989; Blonder *et al.*, 2017). Such discrepancies between the observed  
66 velocity of climate change and the slower velocities at which species and  
67 communities are actually responding have been termed “climatic debt” (i.e. the fact

1  
2  
3 68 that biotic responses are lagging behind climate change) (Menéndez *et al.*, 2006;  
4  
5 69 Bertrand *et al.*, 2011, 2016; Devictor *et al.*, 2012; Zellweger *et al.*, 2020).  
6  
7

8  
9 70 To assess the magnitude of the climatic debt or lag between the velocity of  
10  
11 71 mean annual temperature (MAT) changes and the velocity of thermophilization of  
12  
13 72 communities, one needs first to quantify the degree of thermophilization of local  
14  
15 73 species assemblages. The most straightforward approach is to compute the  
16  
17 74 community temperature index (CTI) (Devictor *et al.*, 2008; Zellweger *et al.*, 2020).  
18  
19 75 The CTI is the average value of either individual species' thermal optima or species-  
20  
21 76 indicator values for temperature across all species co-occurring in a focal  
22  
23 77 community. How CTI changes over time (i.e. the proportion of cold- and warm-  
24  
25 78 adapted species appearing or disappearing from the focal community) likely reflects  
26  
27 79 how communities are responding to climate changes (Devictor *et al.*, 2008; Bertrand  
28  
29 80 *et al.*, 2011). This index thus provides an intuitive proxy to estimate the velocity of  
30  
31 81 thermophilization of communities, and can be subsequently compared to the velocity  
32  
33 82 of observed temperature changes (e.g. both expressed in units of degrees Celsius per  
34  
35 83 decade; Devictor *et al.*, 2008). Changes in CTI (or analogous index) over time have  
36  
37 84 already been used to document thermophilization and delayed biotic responses of a  
38  
39 85 wide range of taxonomic groups (e.g. plants, butterflies, fishes and birds; Devictor *et*  
40  
41 86 *al.*, 2012; Bertrand *et al.*, 2011; Cheung *et al.*, 2013; Gaüzère *et al.*, 2017). However,  
42  
43 87 most of these studies relied on a relatively restricted number of observations over  
44  
45 88 time, usually a baseline survey and one or two resurveys at best, to compute  
46  
47 89 changes in CTI values over time (Savage & Vellend, 2015; Ash *et al.*, 2017; Fadrique  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 90 *et al.*, 2018). Some previous studies even relied on non-permanent or quasi-  
4  
5 91 permanent plots to assess changes in CTI values over time (Bertrand *et al.*, 2011; De  
6  
7  
8 92 Frenne *et al.*, 2013; Becker-Scarpitta *et al.*, 2019; Zellweger *et al.*, 2020).  
9  
10 93 Unfortunately, such studies relying on non-permanent plots tend to infer temporal  
11  
12 94 changes blurred by space-for-time substitutions, which are known to overestimate  
13  
14  
15 95 the magnitude of thermophilization of communities in response to contemporary  
16  
17 96 climate warming (Elmendorf *et al.*, 2015). Even when studies rely on strictly  
18  
19  
20 97 permanent plots, it requires more than one or two resurveys to build a detailed time  
21  
22 98 series of community composition changes from long-term monitoring programs.  
23  
24  
25 99 Strictly permanent monitoring programs relying on frequent resurveys (e.g. from  
26  
27 100 one-year to five-year intervals) during a sufficiently long time period (e.g. at least  
28  
29  
30 101 two decades) are needed to capture a temporal dynamic in the velocity of community  
31  
32 102 thermophilization and confirm linearity over time. Although such long-term time  
33  
34 103 series of biodiversity monitoring already exist (Dornelas *et al.*, 2018), we still lack a  
35  
36  
37 104 good knowledge and understanding of the temporal dynamic of the climatic debt.

38  
39  
40 105         Whether the climatic debt of plant communities is steadily growing over time  
41  
42 106 and what are the main underlying determinants of this dynamic remain open  
43  
44  
45 107 questions in global change ecology (Essl *et al.*, 2015; Bertrand *et al.*, 2016; Zellweger  
46  
47 108 *et al.*, 2020). For instance, a previous study focusing on temperate forests has  
48  
49  
50 109 suggested an increasing climatic debt in the response of understory plant  
51  
52 110 communities from the lowlands (Bertrand *et al.*, 2011). A follow-up study has  
53  
54 111 revealed that this growing climatic debt involved a complex combination of abiotic  
55  
56  
57  
58  
59  
60



1  
2  
3 112 and biotic factors with the greatest effect size captured by environmental conditions  
4  
5 113 during the baseline surveys (i.e. a greater climatic debt for plant communities  
6  
7  
8 114 occurring within initially warmer conditions) (Bertrand *et al.*, 2016). Noteworthy,  
9  
10 115 the same authors also found that **anthropogenic** disturbances contributed very little  
11  
12 116 and unexpectedly to the climatic debt. Yet, previous and more recent findings  
13  
14  
15 117 suggest the opposite with local disturbances amplifying forest community responses  
16  
17 118 to climate change (De Frenne *et al.*, 2013; Stevens *et al.*, 2015; Perring *et al.*, 2018;  
18  
19  
20 119 Brice *et al.*, 2019). A more recent study has even demonstrated that the climatic lag  
21  
22 120 in forest plant communities is primarily controlled by forest microclimate dynamics  
23  
24  
25 121 suggesting **forest stand characteristics** to be a strong determinant of the climatic  
26  
27 122 debt (Zellweger *et al.*, 2020). These discrepancies call for more in-depth  
28  
29  
30 123 investigations on the temporal dynamic of the climatic debt in forest understory  
31  
32 124 plant communities and the relative contribution of **anthropogenic and natural**  
33  
34 125 disturbances.

36  
37  
38 126 In addition to baseline environmental conditions and stand characteristics  
39  
40 127 modulated by **anthropogenic and natural disturbances**, ungulates directly impact  
41  
42 128 vegetation dynamics (Post & Pedersen, 2008; Olofsson *et al.*, 2009; Boulanger *et al.*,  
43  
44  
45 129 2018). **On the one hand**, Olofsson *et al.* (2009) have shown that herbivores inhibit  
46  
47 130 the climate-driven shrub expansion towards high latitudes, **indicating that**  
48  
49  
50 131 **herbivores could contribute to the climatic debt. On the other hand**, Post & Pedersen  
51  
52 132 (2008) have shown that plant community composition (e.g. dwarf birch, willow,  
53  
54 133 graminoids, forbs and mosses) on warmed plots (open-top chambers) did not differ  
55  
56  
57  
58  
59  
60

1  
2  
3 134 from that on ambient plots where herbivores grazed, **suggesting no effect of**  
4  
5 135 **herbivory on the climatic debt**. Altogether, these findings **show contrasting impacts**  
6  
7  
8 136 **of herbivory on** the climatic debt.  
9

10  
11 137 Here, we rely on a long-term (1995-2017) national (France) monitoring  
12  
13 138 network of permanent forest plots organized in a paired design of enclosure (fenced  
14  
15 139 area excluding ungulates) *vs.* control plots – the RENECOFOR monitoring program  
16  
17  
18 140 – with multiple resurveys (every year to every five years) to generate a high-  
19  
20 141 resolution and unprecedented time series of CTI values for understory plant  
21  
22 142 communities inside and outside the fenced area. Using this information-rich time  
23  
24 143 series, we aimed to: (i) compare the rates of CTI changes to the velocity of MAT  
25  
26 144 changes during the period 1995-2017; (ii) test whether the climatic debt of  
27  
28 145 understory plant communities increased steadily over time; and (iii) assess the  
29  
30 146 relative contribution of several abiotic and biotic factors known to be influential  
31  
32 147 from the scientific literature, including baseline environmental conditions, stand  
33  
34 148 characteristics that are influenced by **anthropogenic and natural disturbances**, and  
35  
36 149 biotic disturbances from large herbivores.  
37  
38  
39  
40  
41  
42

## 43 150 **Materials and Methods**

### 44 151 *A long-term national monitoring network of forest ecosystems*

45  
46  
47 152 To assess changes in understory plant communities, we used vegetation data  
48  
49 153 collected every five years, and even every year for some sites, throughout a long-  
50  
51 154 term French monitoring network of forest ecosystems (RENECOFOR,  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 155 <http://www1.onf.fr/renecofor>). The RENECOFOR network belongs to the European  
4  
5 156 ICP Forests Level II monitoring program (ICP Forests, <http://www.icp-forests.org>)  
6  
7  
8 157 and aims at monitoring the health and functioning of forest ecosystems. This  
9  
10 158 monitoring network was established in 1992 by the French National Forest Service  
11  
12 159 (ONF, Office National des Forêts). It includes 102 permanent plots distributed in  
13  
14  
15 160 state and communal forests throughout the whole metropolitan French territory  
16  
17 161 ([www1.onf.fr/renecofor/sommaire/sites](http://www1.onf.fr/renecofor/sommaire/sites)) (Ulrich, 1995 ; Nicolas *et al.*, 2014). All plots  
18  
19 162 are located in mature and even-aged pure stands, and cover an area of 2 ha with a  
20  
21 163 central fenced zone of 0.5 ha to exclude large herbivores (**Fig. 1**) (for more details,  
22  
23 164 see Boulanger *et al.*, 2018).  
24  
25  
26  
27

28 165         Since 1995, vegetation surveys have been conducted every five years (baseline  
29  
30 166 survey in 1995 and resurveys in 2000, 2005, 2010 and 2015). From the total pool of  
31  
32 167 102 permanent plots, we discarded 58 plots which suffered from major data gaps  
33  
34 168 throughout the studied time period (e.g. the resurvey of 2010 was systematically  
35  
36 169 missing for 53 plots due to a shortage in funding). This yielded a total of 44 plots for  
37  
38 170 which a full time series, at a 5-yr interval, was available between 1995 and 2015,  
39  
40 171 hereafter referred to as the five-year dataset (**Fig. 1**). Among these 44 plots, 8 plots  
41  
42 172 were monitored annually without discontinuity between 1995 and 2017, hereafter  
43  
44 173 referred to as the yearly dataset (**Fig. 1**). The whole workflow of analyses was  
45  
46 174 carried out separately for both the five-year and yearly datasets, as these two  
47  
48 175 datasets are very complementary (*Supporting Information App. Fig. 1*). Although  
49  
50 176 the five-year dataset covers a large spatial extent representative of all temperate  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 177 forests in France, its temporal resolution is rather coarse. *Per contra*, the yearly  
4  
5 178 dataset covers a restricted spatial extent but provides a very detailed time series at  
6  
7  
8 179 high temporal resolution. More information on the spatial location and dominant  
9  
10 180 tree species for each of 44 study plots are provided in *Supporting Information App.*  
11  
12  
13 181 *Table 1.*  
14  
15

16 182         During each monitoring campaign, the vegetation was surveyed by expert  
17  
18 183 botanists according to standardized protocols under a quality assurance and quality  
19  
20 184 control procedure. The quality assurance and quality control procedure involved  
21  
22 185 intercalibration exercises organized the year before each monitoring campaign with  
23  
24 186 all expert botanists surveying the same plots in the same forest (Camaret &  
25  
26 187 Brêthes, 2004; Archaux *et al.*, 2009). During each monitoring campaign, all plots  
27  
28 188 were surveyed both at spring and in summer/autumn, to cover the entire growing  
29  
30 189 season. All vascular plants were recorded in eight 2 m × 50 m subplots: four inside  
31  
32 190 and four outside the fenced area of 0.5 ha (**Fig. 1**). The cover-abundance of all taxa  
33  
34 191 occurring in four vegetation layers (tree: > 7 m; high shrub: 2–7 m; low shrub: 0.3–2  
35  
36 192 m; herb: < 0.3 m) was recorded. Species names were standardized following the  
37  
38 193 nomenclature of *Flora Europaea* (Tutin *et al.*, 2001). A total of 877 taxa were  
39  
40 194 recorded in all vegetation layers, including 751 occurring in the herb layer  
41  
42 195 (hereafter understory plants). Here, we restricted our analyses to understory plants,  
43  
44 196 including tree seedlings, because they represent a relevant component of forest  
45  
46 197 biodiversity which is not directly (but indirectly via the upper canopy layers)  
47  
48 198 impacted by forest management practices, and are expected to be the most  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 199 responsive to climate warming. Only taxa identified at the species or subspecies  
4  
5 200 levels were kept for analyses. Identifications at the genus level were considered as  
6  
7 201 incomplete and excluded from our analyses (i.e. 80 taxa excluded; see the list of  
8  
9 202 species provided in *Supporting Information App. Table 2*). Our dataset encompassed  
10  
11 203 a total of 782 vegetation records, including 347 records from the yearly dataset (492  
12  
13 204 species) and 435 records from the five-year dataset (593 species). Seasonal surveys  
14  
15 205 (spring and summer/autumn) were aggregated by year with the maximum cover-  
16  
17 206 abundance coefficient retained whenever a species was recorded at both dates. This  
18  
19 207 yielded to a total of 186 and 220 vegetation records for the yearly and five-year  
20  
21 208 dataset, respectively.

### 209 *Time series of temperature data*

210 To compare the velocity of mean annual temperature (MAT) changes with the  
211 velocity of temperature changes inferred from understory plant communities, we  
212 first extracted time series of MAT changes. We obtained macroclimate temperature  
213 time series from the global dataset TerraClimate (Abatzoglou *et al.*, 2018), which  
214 provides monthly climate data for every year from 1958 up to 2018 at a spatial  
215 resolution of 2.5 arc-minute ( $\sim 3.3 \text{ km} \times 4.64 \text{ km}$  at  $45^\circ$  of latitude). We also extracted  
216 climatic time series from the CHELSA database (Karger *et al.*, 2017) but found very  
217 similar trends between TerraClimate and CHELSA (*Supporting Information App.*  
218 *Fig. 2*). Thus, we decided to use only TerraClimate in our subsequent analyses as it  
219 better covers the end of our study period (i.e. 2017). **As TerraClimate provides only  
220 maximum and minimum monthly temperatures, we first calculated the monthly**

221 mean for each of the 44 studied plots (by averaging maximum and minimum  
 222 temperature values for each month), and then computed mean annual temperature  
 223 (hereafter MAT) for each year between 1958 and 2017. Between 1958 and 2017,  
 224 MAT increased across all 44 studied plots, leading to a shift from negative to only  
 225 positive anomalies towards the end of the reference period (1958–2017) (*Supporting*  
 226 *Information App. Fig. 3*). From the very beginning (1992) of the RENECOFOR  
 227 monitoring program until 2017, anomalies were mostly positive (values ranging  
 228 from -0.56 to 1.54 over the period 1992-2017), except for three years (1993, 1996 and  
 229 2010), for which anomalies were negative.

230 To infer temperature conditions from local species assemblages, we used the  
 231 community temperature index (CTI) (Devictor *et al.*, 2008, 2012) (see Equation 1).  
 232 The CTI was computed based on occurrence data by averaging the indicator values  
 233 (IVs) for temperature across all species  $i$  ( $Sp_i$ ) occurring in a given relevé  $j$   
 234 (containing  $S$  species, with  $S \geq 2$ ). Here, we used presence/absence data instead of  
 235 weighting IVs by each species' abundance/dominance in the relevés. We did that to  
 236 give the same weight to common and rare species. As each RENECOFOR plot  
 237 includes 8 subplots, we calculated the CTI values at the subplot level and kept this  
 238 information at the subplot level without aggregating it at the plot level.

$$(1) \quad CTI_j = \frac{\sum_{i=1}^S IV \text{ Temperature } Sp_i}{S}$$

240 Species' IVs for temperature were extracted from both the CATMINAT (Julve, 1998)  
 241 and EcoPlant (Gégout *et al.*, 2005) databases, and we used them in combination to  
 242 take advantage of their respective strengths (*Supporting Information App. Fig. 4*).

1  
2  
3 243 On the one hand, the CATMINAT database is based on expert knowledge similarly  
4  
5 244 to Ellenberg's IVs (Ellenberg et al., 1992), with temperature preferences of plant  
6  
7  
8 245 species coded along an ordinal scale from cold (1) to warm (9) conditions. It contains  
9  
10 246 almost all of species recorded in our dataset (858 out of the 877 species recorded in  
11  
12 247 all vegetation layers), but CTI values are unitless and direct comparisons with  
13  
14  
15 248 macroclimatic temperatures are not possible. On the other hand, EcoPlant IVs for  
16  
17 249 temperature are directly expressed in degrees Celsius (Gégout et al., 2005) allowing  
18  
19 250 the direct computation of CTI values in degrees Celsius, but much less species from  
20  
21 251 our dataset are recorded (396 out of 877 species). Based on these two databases, we  
22  
23  
24 252 calibrated a transfer function to produce CTI values directly expressed in degrees  
25  
26 253 Celsius and transform the unitless CTI values from the CATMINAT database into a  
27  
28 254 CTI value expressed in degrees Celsius (*Supporting Information App. Fig. 4 & 5*  
29  
30  
31 255 and see Lenoir *et al.*, 2013 for a similar approach).

32  
33  
34  
35 256 Based on the time series of MAT values and the time series of CTI values,  
36  
37 257 both expressed in degrees Celsius, we computed the lag as the difference between  
38  
39 258 the two (MAT – CTI), thus getting a new time series of the difference: the climatic  
40  
41 259 debt (Bertrand *et al.*, 2016).

### 42 43 44 45 260 *The main determinants of the climatic debt*

46  
47  
48  
49 261 Although we were primarily interested in assessing the temporal dynamic of the  
50  
51 262 difference between MAT and CTI values, thereby testing the effect of time itself on  
52  
53 263 the direction and magnitude of this difference – as we assumed the climatic debt to

1  
2  
3 264 increase over time – we also assessed the relative contributions of several abiotic  
4  
5 265 and biotic determinants known to potentially modulate the climatic debt. For  
6  
7  
8 266 instance, it has already been demonstrated that MAT during or prior to the baseline  
9  
10 267 survey was a strong and positive determinant of the climatic debt in understory  
11  
12 268 plant communities, suggesting greater lags in initially warmer areas (Bertrand *et*  
13  
14  
15 269 *al.*, 2016). Hence, to account for that potential confounding effect, we computed  
16  
17 270 baseline temperature conditions from TerraClimate as the average of MAT values  
18  
19  
20 271 during the 1961-1990 reference period (i.e. 30-yr average typically used to capture  
21  
22 272 long-term climatic conditions) and added it as a covariate to explain the climatic  
23  
24  
25 273 debt.

26  
27  
28 274 In addition to baseline temperature conditions, we gathered information on  
29  
30 275 anthropogenic and natural disturbances, and stand characteristics, all available  
31  
32 276 from the RENECOFOR monitoring program. Because local disturbances may have  
33  
34  
35 277 significant effects on understory plant communities and are known to accelerate  
36  
37 278 their thermophilization rates and reduce the climatic debt (Bengtsson *et al.*, 2000;  
38  
39  
40 279 Stevens *et al.*, 2015; Brice *et al.*, 2019), we built several variables capturing these  
41  
42 280 local disturbances (*Supporting Information App. Fig. 6 & 7*). First, we distinguished  
43  
44  
45 281 between natural (e.g. windstorm) and anthropogenic disturbances (i.e. local  
46  
47 282 disturbances associated to forest management practices such as thinning, cleaning  
48  
49  
50 283 and tree extraction). For each of these two types of disturbances, we computed the  
51  
52 284 cumulative sum of disturbance events occurring within a given forest plot prior to  
53  
54  
55 285 each survey year. Here, we assumed that repeated disturbance events are likely to



1  
2  
3 286 accelerate the thermophilization rate and thus reduce the climatic debt of  
4  
5 287 understory plant communities. Then, irrespective of the type of disturbance, we also  
6  
7  
8 288 computed the anteriority, i.e. the number of years, between the last disturbance  
9  
10 289 event and the year of the survey, assuming that a very short anteriority (less than a  
11  
12 290 year) will impact less the difference between MAT and CTI values. Indeed, plant  
13  
14  
15 291 community composition is unlikely to change during the vegetation season that is  
16  
17 292 just following a stand thinning event. However, significant changes in plant  
18  
19  
20 293 community composition may lag and happen during the next couple of years with  
21  
22 294 light demanding and warm-adapted species germinating from the soil seed bank and  
23  
24  
25 295 increasing CTI values, thus paying off part of the climatic debt. We also retrieved  
26  
27 296 information about stand characteristics at the plot level, including: stand basal area  
28  
29  
30 297 during the monitoring year; stand age in 1995; and the type of dominant tree species  
31  
32 298 (i.e. deciduous *vs.* coniferous). We hypothesized that plots with higher stand basal  
33  
34 299 area and located in older forest stands are likely to be more buffered against the  
35  
36  
37 300 adverse effect of macroclimate warming (Zellweger *et al.*, 2019), thus further  
38  
39 301 inflating the climatic debt of understory plant communities (De Frenne *et al.*, 2013).  
40  
41  
42 302 The enclosure effect (i.e. fenced area excluding ungulates and large herbivores  
43  
44 303 *vs.* control plots outside the fenced area) was added as a covariate in all our analyses  
45  
46  
47 304 on the temporal trends in CTI values and the climatic debt. According to the  
48  
49 305 scientific literature, climate change impacts on vegetation dynamics tend to be  
50  
51 306 inhibited by large herbivores (Post & Pedersen, 2008; Kaarlejärvi *et al.*, 2013), hence  
52  
53  
54 307 we hypothesized a greater climatic debt outside than inside the fenced area.  
55  
56  
57  
58  
59  
60

### 308 *Statistical analyses*

309 We used linear mixed-effects models (LMMs) to assess the velocity of MAT changes  
310 and thermophilization of understory plant communities. For CTI values, we used  
311 the continuous variable “year” of the monitoring as the main fixed effect while  
312 accounting for potentially diverging temporal dynamics between the outside and  
313 inside of the fenced area, by adding the interaction effect: year  $\times$  enclosure. The  
314 RENEFOFOR plot ID (44 or 8 levels depending on the dataset used) and the  
315 vegetation subplot ID (8 levels), nested within the plot ID, were used as a random  
316 term to account for the hierarchical structure of the sampling design and the  
317 repeated surveys over time. For MAT values, we only used the “year” of the  
318 monitoring as a fixed effect and the RENEFOFOR plot ID as a random term.  
319 Indeed, for MAT, the subplot level was not used in the random component of LMMs  
320 since all 8 subplots within a given plot have similar MAT values. To allow between-  
321 site variation in the temporal dynamics of MAT and CTI values, we set the random  
322 term (e.g. plot ID) of our LMMs so that it modifies the slope coefficient (i.e. random  
323 slope term) of the variable “year” for each level of the random term variable. The  
324 coefficient estimate of the “year” variable in our LMMs was used as a proxy for  
325 estimating the velocities of temperature changes, both for MAT and CTI values.  
326 Assumptions of normality and homoscedasticity of the residuals of our models were  
327 visually checked and no departure from normality and homoscedasticity were  
328 detected. Significance of the predictors in LMMs were assessed using Wald  $\chi^2$  tests,  
329 after fitting models with the restricted maximum likelihood estimation.

1  
2  
3 330 To analyze the temporal dynamic of the difference between MAT and CTI  
4  
5  
6 331 values over time (i.e. the climatic debt), we used LMMs with the value of the  
7  
8 332 difference (MAT – CTI) as the response variable and the continuous variable “year”  
9  
10 333 of monitoring as the main fixed effect variable. In addition to “year”, we added  
11  
12 334 several abiotic and biotic covariates as fixed effect variables to assess their relative  
13  
14  
15 335 contribution in modulating the climatic debt. These covariates were: baseline  
16  
17 336 temperature conditions; sum of natural disturbances; sum of anthropogenic  
18  
19  
20 337 disturbances; disturbance anteriority; stand basal area; stand age; **type of dominant**  
21  
22 338 **tree species in forest stand (deciduous vs. coniferous)**; and the exclosure effect. From  
23  
24  
25 339 the full model including all covariates together with time as fixed effect terms, we  
26  
27 340 ran all possible combinations of models and ranked all candidate models using the  
28  
29  
30 341 corrected Akaike information criteria (AICc). Based on the best candidate model  
31  
32 342 with the lowest AICc value, we then computed the difference in AICc values ( $\Delta\text{AICc}$ )  
33  
34  
35 343 for all models relative to the best candidate model. Finally, from the set of selected  
36  
37 344 candidate models with  $\Delta\text{AICc}$  values  $< 2$ , we used a model averaging approach to  
38  
39 345 calculate the conditional mean value of each coefficient and its associated 95%  
40  
41  
42 346 confidence interval (Burnham & Anderson, 2002). All continuous predictor variables  
43  
44 347 were standardized ( $z$ -score) before analyses to improve their interpretability on a  
45  
46  
47 348 comparable scale (Schielzeth, 2010). None of the fixed-effect variables included in  
48  
49 349 our models showed multicollinearity issues ( $\text{VIF} < 3$ ; Zuur *et al.*, 2007). Note that  
50  
51 350 correlation values among all our explanatory variables did not exceed a Spearman’s  
52  
53  
54 351  $r$  coefficient of 0.6 (*Supporting Information App. Fig. 8*). Again, we used the  
55  
56 352 RENECOFOR plot ID (44 or 8 levels depending on the dataset used) and the  
57  
58  
59  
60

1  
2  
3 353 vegetation subplot ID (8 levels), nested within the plot ID, as a random slope term  
4  
5 354 modifying the effect of “year” of monitoring. The explained variance of LMMs was  
6  
7  
8 355 determined by calculating the marginal ( $R^2_m$ , associated with fixed effects only) and  
9  
10 356 conditional ( $R^2_c$ , associated with both fixed and random effects)  $R^2$  values  
11  
12  
13 357 (Nakagawa & Schielzeth, 2013) using the `r.squared.GLMM` function from the MuMIn  
14  
15 358 package.

18 359 All statistical analyses were performed with the R software version 3.6.0 (R  
19  
20 360 Core Team, 2019). Models were performed with the nlme package v.3.1-140 (Pinheiro  
21  
22 361 *et al.*, 2019), and the model selection procedure was assessed with the MuMIn package  
23  
24 362 v.1.43.6 (Bartoń, 2019), using the dredge function. Other packages were used for  
25  
26 363 data handling extraction and graphical visualization: dplyr v.0.8.3 (Wickham *et al.*,  
27  
28 364 2019); raster v.2.9-23 (Hijmans, 2019); ggplot2 v.3.2.0 (Wickham, 2016); cowplot  
29  
30 365 v.1.0.0 (Wilke, 2019); and sf v.0.7-7 (Pebesma, 2018).

## 36 366 Results

### 37 367 *Temporal trends in macroclimate and community temperatures*

38  
39  
40 368 Using the 5-yr interval dataset, mean annual temperature (MAT) during 1995–2015  
41  
42 369 increased by  $0.23 \pm 0.02$  °C/decade (95 % Confidence Intervals (CI): 0.18-0.28,  $t =$   
43  
44 370 9.05,  $P < 0.001$ ; **Fig. 2a**). In the yearly dataset, MAT increased by  $0.31 \pm 0.05$   
45  
46 371 °C/decade (95 % CI: 0.20-0.42,  $t = 5.51$ ,  $P < 0.001$ ; **Fig. 2b**) during 1995–2017. Over  
47  
48 372 the same time periods, the velocities of CTI changes were weaker, but similar trends  
49  
50  
51 373 and magnitudes were observed in both the five-year ( $0.10 \pm 0.03$  °C/decade, 95 %

1  
2  
3 374 CI: 0.04-0.15,  $t = 3.62$ ,  $P < 0.001$ ; **Fig. 2a**) and yearly ( $0.08 \pm 0.03$  °C/decade, 95 %  
4  
5 375 CI: 0.01-0.14,  $t = 2.15$ ,  $P = 0.031$ ; **Fig. 2b**) datasets. No significant differences were  
6  
7 376 observed in CTI trends between inside and outside of the fenced area. Further, the  
8  
9 377 difference between MAT and CTI values showed a tendency to increase steadily over  
10  
11 378 time (**Fig. 2**): the lag in the response of herbaceous plant communities to climate  
12  
13 379 warming increased linearly over time. Although we found a significant effect of time,  
14  
15 380 it contributed weakly to the overall variation in MAT and CTI values:  $R^2_m \leq 0.01$   
16  
17 381 while  $R^2_c$  ranged between 0.92 to 0.96.  
18  
19  
20  
21  
22

### 23 382 *Factors contributing to the climatic debt*

24  
25  
26 383 For both the yearly and five-year datasets, the model averaging procedure to explain  
27  
28 384 the climatic debt (MAT – CTI) selected 4 and 3 candidate models ( $\Delta AIC_c < 2$ ),  
29  
30 385 respectively. The best candidate models explained 55% and 70% of the total  
31  
32 386 variation in the difference between MAT and CTI values for the five-year ( $R^2_m =$   
33  
34 387  $0.33$ ;  $R^2_c = 0.55$ ) and yearly ( $R^2_m = 0.46$ ;  $R^2_c = 0.70$ ) datasets, respectively. The best  
35  
36 388 candidate model for the five-year dataset contained 6 variables including the year of  
37  
38 389 sampling, stand age, the anteriority of the last disturbance, the cumulative number  
39  
40 390 of anthropogenic disturbances, stand basal area, and baseline temperature  
41  
42 391 conditions. For the yearly dataset, the best candidate model included the same 6  
43  
44 392 variables together with **the type of dominant tree species** (*Supporting Information*  
45  
46 393 *App. Tables 3 & 4 and App. Fig. 9 & 10*).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 394 Irrespective of the dataset considered in our analyses, we found very  
4  
5 395 consistent and similar trends in the factors contributing to the climatic debt (**Fig. 3**).  
6  
7  
8 396 For instance, the year of sampling had a positive effect on the magnitude of the  
9  
10 397 difference, confirming an increasing climatic debt of understory plant communities  
11  
12 398 over time (five-year dataset =  $0.06 \pm 0.03$  SE, *Fig. 3a*; yearly dataset =  $0.31 \pm 0.04$   
13  
14 399 SE, *Fig. 3b* and *Fig. 4a*).

15  
16  
17  
18 400 Baseline temperature conditions had the strongest positive influence on the  
19  
20 401 climatic debt (five-year dataset =  $0.77 \pm 0.04$  SE, *Fig. 3a*; yearly dataset =  $0.72 \pm$   
21  
22 402  $0.10$  SE, *Fig. 3b*), indicating greater lags in plots with warmer baseline temperature  
23  
24 403 conditions during the period 1961–1990. We also found a positive effect of stand  
25  
26 404 characteristics, through the variables stand age in 1995 and basal area during the  
27  
28 405 year of monitoring, on the magnitude of the lag. More specifically, the lag was larger  
29  
30 406 in denser (five-year dataset =  $0.22 \pm 0.04$  SE, *Fig. 3a*; yearly dataset =  $0.35 \pm 0.05$   
31  
32 407 SE, *Fig. 3b* and *Fig. 4b*) and older (five-year dataset =  $0.15 \pm 0.04$  SE, *Fig. 3a*;  
33  
34 408 yearly dataset =  $0.46 \pm 0.07$  SE, *Fig. 3b* and *Fig. 4c*) forests. We found no clear  
35  
36 409 differences in the climatic debt between coniferous and deciduous forests as well as  
37  
38 410 between inside and outside of the fenced area.

39  
40  
41  
42  
43  
44  
45 411 In contrast, forest disturbance had a negative effect on the climatic debt (*Fig.*  
46  
47 412 *3*). Notably, the magnitude of the lag was negatively correlated with the anteriority  
48  
49 413 of the last disturbance event (five-year dataset =  $-0.12 \pm 0.02$  SE, *Fig. 3a*; yearly  
50  
51 414 dataset =  $-0.10 \pm 0.03$  SE, *Fig. 3b*), revealing that lags decrease in plots recently  
52  
53 415 disturbed. Moreover, the cumulative number of anthropogenic disturbances were  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 416 negatively correlated to the climatic debt (five-year dataset =  $-0.17 \pm 0.04$  SE, *Fig.*  
4  
5 417 *3a*; yearly dataset =  $-0.19 \pm 0.07$  SE, *Fig. 3b*), indicating that lags decrease in the  
6  
7 418 most frequently disturbed plots, following logging events. The cumulative number of  
8  
9 419 natural disturbances was only retained in the models based on the five-year dataset,  
10  
11 420 albeit its tendency towards a negative effect did not differ from zero ( $-0.04 \pm 0.04$   
12  
13 421 SE, *Fig. 3a*).  
14  
15  
16  
17  
18

## 19 422 Discussion

### 20 21 22 423 *Velocities of thermophilization*

23  
24  
25 424 Irrespective of the temporal resolution in the data (i.e. the yearly vs. five-year  
26  
27 425 dataset), we show a very similar increase in temperature conditions inferred from  
28  
29 426 understory plant communities over the period 1995–2017, consistent with a  
30  
31 427 thermophilization process. This finding adds to a growing body of evidence  
32  
33 428 describing thermophilization of plant communities across many forest ecosystems  
34  
35 429 around the globe (e.g. in temperate, boreal and tropical ecosystems; Lenoir *et al.*,  
36  
37 430 2010; Bertrand *et al.*, 2011; De Frenne *et al.*, 2013; Savage & Vellend, 2015;  
38  
39 431 Fadrique *et al.*, 2018; Zellweger *et al.*, 2020), as well as across other ecosystems and  
40  
41 432 taxonomic groups (e.g. in mountain plant communities, Gottfried *et al.*, 2012;  
42  
43 433 marine fishes and invertebrates, Cheung *et al.*, 2013; birds and butterflies, Devictor  
44  
45 434 *et al.*, 2012; bumblebee communities, Fourcade *et al.*, 2018). The velocities of  
46  
47 435 thermophilization we report here (ranging from 0.08 to 0.10 °C/decade) are broadly  
48  
49 436 in the range of those reported in the scientific literature (from 0.03 to 0.38  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 437 °C/decade; *Supporting Information App. Table. 5*), but also are among the highest  
4  
5 438 rates recorded for plant communities (maximum value of 0.27 °C/decade; Duque *et*  
6  
7  
8 439 *al.*, 2015). Specifically, we show velocities that are twice as fast as the average  
9  
10 440 thermophilization rate estimated for understory plant communities across  
11  
12 441 temperate deciduous forests in Europe (i.e. 0.041 °C/decade; De Frenne *et al.*, 2013).  
13  
14  
15 442 Nevertheless, these rates are still much lower than those found in marine  
16  
17 443 ecosystems, where environmental constraints may weakly limit the dispersal of  
18  
19  
20 444 species in response to rising temperatures (Lenoir *et al.*, 2020).  
21  
22

23 445 Our findings consist in the most recent updates supporting and consolidating  
24  
25 446 previous reports of plant community thermophilization which happened across the  
26  
27  
28 447 same study area (France) (Bertrand *et al.*, 2011; Martin *et al.*, 2019). Focusing on a  
29  
30 448 completely independent dataset from the period 1965–2008, Bertrand *et al.* (2011)  
31  
32  
33 449 were the first to highlight a thermophilization of understory plant communities in  
34  
35 450 the French forests. More recently, Martin *et al.* (2019) have described a similar  
36  
37  
38 451 thermophilization trend extending to other plant communities at national and  
39  
40 452 regional scales in France over the period 2009–2017. Unlike these previous studies  
41  
42 453 relying either on massive collection of opportunistic surveys over time (Bertrand *et*  
43  
44  
45 454 *al.*, 2011) or citizen science data (Martin *et al.*, 2019), our study is the first that used  
46  
47 455 a long-term monitoring program based on strictly permanent plots intensively  
48  
49  
50 456 monitored over time (up to every year), providing a very detailed time series of  
51  
52 457 changes in understory plant community composition. Hence, we argue that the  
53  
54  
55 458 velocities we report here do not suffer from biases due to space-for-time  
56  
57  
58  
59  
60



1  
2  
3 459 substitutions and are thus likely to be more accurate than those previously reported  
4  
5 460 (Elmendorf *et al.*, 2015), which matters for informing our society on future scenarios  
6  
7 461 of biodiversity changes (IPBES, 2019). Furthermore, our yearly resurveys over a  
8  
9 462 period of 25 years confirm that the thermophilization rate of understory plant  
10  
11 463 communities in temperate forests is following a linear trajectory over time. This is  
12  
13 464 an important and novel finding that helps us to understand the temporal dynamic of  
14  
15 465 the biotic responses to anthropogenic climate change. Yet, time alone had a very  
16  
17 466 minor contribution to explain the variation in temperature conditions inferred from  
18  
19 467 understory plant communities. This suggests that other abiotic and biotic  
20  
21 468 determinants, such as the macroclimatic context as well as local stand  
22  
23 469 characteristics influenced by anthropogenic disturbances may potentially explain  
24  
25 470 local variations in temperature conditions.  
26  
27  
28  
29  
30  
31  
32

### 33 471 *The climatic debt and its potential determinants*

34  
35

36 472 One of our most striking results is that the velocity at which atmospheric air  
37  
38 473 temperatures are rising is twice faster, on average, than the velocity at which  
39  
40 474 understory plant communities are responding, generating a climatic debt that is  
41  
42 475 steadily growing over time, even after accounting for the relative contribution of  
43  
44 476 several abiotic and biotic determinants modulating it. Although the idea of an  
45  
46 477 increasing climatic debt over time has already been suggested (Bertrand *et al.*,  
47  
48 478 2011), we provide the first evidence that the climatic debt of understory plant  
49  
50 479 communities in response to the ongoing climate change is steadily increasing over  
51  
52 480 time. This contrasts with a previous report showing no clear temporal signal in the  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 481 climatic debt of herbaceous plant communities in French forests (Bertrand *et al.*,  
4  
5 482 2016). Considering the velocity of the predicted future climate change (Loarie *et al.*,  
6  
7 483 2009), this is an important result because it raises the question whether the lag will  
8  
9 484 continue to steadily accumulate and approach a potential critical breakpoint for  
10  
11 485 understory biodiversity and forest ecosystem functioning. Interestingly, the linear  
12  
13 486 increase of the climatic debt over time is clearer and more pronounced (higher effect  
14  
15 487 size) in the yearly dataset than in the five-year dataset. This is probably because the  
16  
17 488 long-term increasing trend in the climatic debt was better separated from inter-  
18  
19 489 annual variations in the yearly dataset than in the five-year dataset. Inter-annual  
20  
21 490 variations tend to play a bigger role in blurring the long-term response of understory  
22  
23 491 plant communities when the frequency between resurveys is lower.  
24  
25  
26  
27  
28  
29

30 492 Importantly, we demonstrate that forest stand characteristics and  
31  
32 493 anthropogenic disturbances significantly matter in modulating the climatic debt, by  
33  
34 494 either increasing (positive effect) or mitigating (negative effect) the lag. For  
35  
36 495 instance, both the anteriority of the last disturbance event and the cumulative  
37  
38 496 number of anthropogenic disturbances have a negative effect on the climatic debt.  
39  
40  
41 497 Such findings are in accordance with recent works indicating that local disturbances  
42  
43 498 speed up the process of thermophilization in plant communities (Stevens *et al.*,  
44  
45 499 2015; Brice *et al.*, 2019; Zellweger *et al.*, 2020). However, it remains to be  
46  
47 500 investigated in more details whether the thermophilization of understory plant  
48  
49 501 communities following anthropogenic disturbances is due to warm-adapted species  
50  
51 502 that are: (i) replacing more mesophilous species (pure turnover); (ii) increasing the  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 503 local species pool (nestedness); or both simultaneously. For instance, Closset-Kopp  
4  
5 504 *et al.* (2018) have shown that understory plant communities in heavily managed  
6  
7  
8 505 forest stands had the tendency to be richer in warm-adapted species in 2015 than in  
9  
10 506 1970, mostly due to an increase in local species richness but no species turnover. In  
11  
12  
13 507 our study, both processes seem to occur. Indeed, while the total number of species  
14  
15 508 per plot increases over time, suggesting a process of nestedness, the set of warm-  
16  
17  
18 509 adapted species that are colonizing (i.e. species gain) the plots of our monitoring  
19  
20 510 network between 1995 and 2015 are also replacing relatively more mesic species  
21  
22 511 that are disappearing (i.e. species loss) (*Supporting Information, App. Fig. 11*),  
23  
24  
25 512 suggesting a process of turnover. Not only the richness of warm-adapted species  
26  
27 513 increases at the expense of cold-adapted species, but this effect is concomitant with  
28  
29  
30 514 a relative increase in the richness of light- and nitrogen-demanding species at the  
31  
32 515 expense of species tolerating shade conditions and nitrogen-poor soils (*Supporting*  
33  
34 516 *Information, App. Fig. 12*). **These trends further confirm the importance of local**  
35  
36 517 **disturbances as key factors modulating the lag between macroclimatic conditions**  
37  
38 518 **and plant community thermophilization.** Noteworthy, our results contrast with  
39  
40  
41 519 those from Bertrand *et al.* (2016), who reported non-significant effects of  
42  
43  
44 520 anthropogenic disturbances on the climatic debt of understory plant communities.  
45  
46 521 Looking at the effects of stand characteristics, our results underline the key role of  
47  
48  
49 522 stand age and basal area to modulate the climatic debt. Older and denser (high  
50  
51 523 basal area) forest stands provide particular microclimatic conditions (e.g. cooler  
52  
53  
54 524 ground-level temperatures due to increased shading and humidity conditions) that  
55  
56 525 are likely to buffer the adverse effects of macroclimatic warming on understory  
57  
58  
59  
60

1  
2  
3 526 plant communities thus providing thermal microrefugia for species persistence  
4  
5 527 (Lenoir *et al.*, 2017; De Frenne *et al.*, 2019; Zellweger *et al.*, 2020). In contrast,  
6  
7 528 repeated disturbances are likely to create canopy openings in forest stands,  
8  
9 529 modifying the microhabitat and associated microclimate that strongly exacerbate  
10  
11 530 the impact of macroclimate temperature warming for understory vegetation  
12  
13 531 (Vanderwel & Purves, 2014; Zellweger *et al.*, 2020). This can in turn lead to drastic  
14  
15 532 changes in the overall composition of the plant community (i.e. by increasing the  
16  
17 533 proportion of warm-adapted species through either species turnover or an increase  
18  
19 534 in local species richness) that will reduce the climatic debt. Paying off the climatic  
20  
21 535 debt of understory plant communities in temperate forests implies to break off this  
22  
23 536 microclimate stability, which can consequently reduce the microrefugia capacity of  
24  
25 537 forest habitats for the survival of forest-dwelling species under anthropogenic  
26  
27 538 climate change (Lenoir *et al.*, 2017). Accepting the idea that the climatic debt of  
28  
29 539 understory plant communities might be a positive sign of local adaptation has  
30  
31 540 significant implications on the concept of climatic debt itself, which might be  
32  
33 541 interpreted as a macroclimatic debt but not a microclimatic debt.  
34  
35  
36  
37  
38  
39  
40  
41

42 542 Contrary to our initial expectation that large herbivores should contribute to  
43  
44 543 the climatic debt by further increasing the lag in the response of understory plant  
45  
46 544 communities to macroclimate warming, we found no difference in the observed  
47  
48 545 climatic debt between outside and inside the fenced area. This suggests that the  
49  
50 546 climatic debt generated by the buffering effect of microclimate in the understory  
51  
52 547 may mostly stem from the canopy layer and independently from the shrub layer, as  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 548 large herbivores are mostly impacting woody species from the understory shrub  
4  
5 549 layer (Olofsson *et al.*, 2009; Boulanger *et al.*, 2018). The indirect effects of herbivores  
6  
7  
8 550 on vegetation responses to climate warming can be highly dependent on the  
9  
10 551 ecological context, as pointed out by Vuorinen *et al.*, (2020).

11  
12  
13 552 Together with previous studies, our results show that the climatic debt in  
14  
15 553 understory plant communities seem to be much more pronounced in situations  
16  
17  
18 554 where temperature conditions were warmer during the period preceding the  
19  
20  
21 555 baseline surveys (Bertrand *et al.*, 2016), such as in the lowlands (Bertrand *et al.*,  
22  
23 556 2011). However, the fact that warmer baseline conditions increase the climatic debt  
24  
25  
26 557 of understory plant communities could also stem from a bias towards an  
27  
28 558 underestimation of CTI values in communities where immigrating species can only  
29  
30  
31 559 come from even warmer regions likely outside the calibration range used to compute  
32  
33 560 the CTI values. This could therefore lead to a systematic overestimation of the  
34  
35 561 observed climatic debt in the warmest part of the study area. This hypothesis is  
36  
37  
38 562 supported by the extreme values observed in the distribution of indicator values  
39  
40 563 (IVs) from the EcoPlant database (*Supporting Information App. Fig. 5*), which may  
41  
42  
43 564 indicate that the model calibration range is too limited and should be extended to  
44  
45 565 other environmental conditions (i.e. here, the Mediterranean regions to capture the  
46  
47 566 pool of immigrating species coming from for warmer conditions). This potential bias  
48  
49  
50 567 supports general recommendations to take into account existing differences in  
51  
52 568 sensitivity between species pools when assessing temporal changes in CTI values  
53  
54 569 (Burrows *et al.*, 2019; Williams *et al.*, 2020). More data on species IVs, either expert-

1  
2  
3 570 based or biogeographically-derived IVs, are needed from warmer regions to improve  
4  
5 571 estimates of CTI values in the current context of global warming.  
6  
7  
8

## 9 572 Conclusion

10  
11  
12 573 This study shows that the thermophilization of understory plant communities in  
13  
14 574 French forests continued during the last two decades with rapid but insufficient  
15  
16 575 rates to track the velocity of current climate warming, leading to a steadily increase  
17  
18 576 in the climatic debt. Additionally, we report that forest stand structure and  
19  
20 577 anthropogenic disturbances can be modulated to increase or decrease the climatic  
21  
22 578 debt. For instance, the positive effect of denser and older canopy layers on forest  
23  
24 579 microclimate dynamics could inflate the climatic debt of understory plant  
25  
26 580 communities and provide more time for understory plant communities to locally  
27  
28 581 adapt. However, it remains uncertain how big the climatic debt can grow over time  
29  
30 582 before a tipping point is reached such that the climatic debt will suddenly be paid  
31  
32 583 off. For instance, repeated drought events leading to extensive tree dieback and  
33  
34 584 massive crown defoliation could exceed this tipping point by suddenly recoupling the  
35  
36 585 understory layer with macroclimatic conditions prevailing outside forests. Such a  
37  
38 586 quick recovery of the climatic debt will have drastic consequences on forest  
39  
40 587 biodiversity and ecosystem functioning.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 589 References

- 590 Abatzoglou, J.T., Dobrowski, S.Z., Parks, S.A. & Hegewisch, K.C. (2018)  
591 TerraClimate, a high-resolution global dataset of monthly climate and climatic  
592 water balance from 1958–2015. *Scientific Data*, **5**, 170191.
- 593 Archaux, F., Camaret, S., Dupouey, J.L., Ulrich, E., Corcket, E., Bourjot, L., ...  
594 Touffet, J. (2009) Can we reliably estimate species richness with large plots? an  
595 assessment through calibration training. *Plant Ecology*, **203**, 303–315.
- 596 Ash, J.D., Givnish, T.J. & Waller, D.M. (2017) Tracking lags in historical plant  
597 species' shifts in relation to regional climate change. *Global Change Biology*, **23**,  
598 1305–1315.
- 599 Bartoń, K. (2019) MuMIn: Multi-Model Inference.
- 600 Becker-Scarpitta, A., Vissault, S. & Vellend, M. (2019) Four decades of plant  
601 community change along a continental gradient of warming. *Global Change Biology*,  
602 **25**, 1629–1641.
- 603 Bengtsson, J., Nilsson, S.G., Franc, A. & Menozzi, P. (2000) Biodiversity,  
604 disturbances, ecosystem function and management of European forests. *Forest  
605 Ecology and Management*, **132**, 39–50.
- 606 Bertrand, R., Lenoir, J., Piedallu, C., Dillon, G.R., De Ruffray, P., Vidal, C., ...  
607 Gégout, J.C. (2011) Changes in plant community composition lag behind climate  
608 warming in lowland forests. *Nature*, **479**, 517–520.

- 1  
2  
3 609 Bertrand, R., Riofrío-Dillon, G., Lenoir, J., Drapier, J., De Ruffray, P., Gégout, J.C.  
4  
5  
6 610 & Loreau, M. (2016) Ecological constraints increase the climatic debt in forests.  
7  
8 611 *Nature Communications*, **7**.  
9  
10  
11 612 Blonder, B., Moulton, D.E., Blois, J., Enquist, B.J., Graae, B.J., Macias-Fauria, M.,  
12  
13 613 ... Svenning, J.-C. (2017) Predictability in community dynamics. *Ecology Letters*, **20**,  
14  
15 614 293–306.  
16  
17  
18  
19 615 Boulanger, V., Dupouey, J., Archaux, F., Badeau, V., Baltzinger, C., Chevalier, R., ...  
20  
21 616 Ulrich, E. (2018) Ungulates increase forest plant species richness to the benefit of  
22  
23 617 non-forest specialists. *Global Change Biology*, **24**, e485–e495.  
24  
25  
26  
27 618 Brice, M., Cazelles, K., Legendre, P. & Fortin, M. (2019) Disturbances amplify tree  
28  
29 619 community responses to climate change in the temperate–boreal ecotone. *Global*  
30  
31 620 *Ecology and Biogeography*, geb.12971.  
32  
33  
34  
35 621 Burnham, K.P. & Anderson, D.R. eds. (2002) *Model Selection and Multimodel*  
36  
37 622 *Inference: A Practical Information-Theoretic Approach*. pp. 49–97. Springer-Verlag  
38  
39 623 New York.  
40  
41  
42  
43 624 Burrows, M.T., Bates, A.E., Costello, M.J., Edwards, M., Edgar, G.J., Fox, C.J.,  
44  
45 625 Halpern, B.S., Hiddink, J.G., Pinsky, M.L., Batt, R.D., García Molinos, J., Payne,  
46  
47 626 B.L., Schoeman, D.S., Stuart-Smith, R.D. & Poloczanska, E.S. (2019) Ocean  
48  
49 627 community warming responses explained by thermal affinities and temperature  
50  
51 628 gradients. *Nature Climate Change*, **9**, 959–963.  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 629 Camaret, S. & Brêthes, A. (2004) *Suivi de la composition floristique des placettes du*  
4  
5 630 *réseau (1994/95-2000) et élaboration d'un programme d'assurance qualité intensif,*  
6  
7  
8 631 Office National des Forêts-Direction Technique.  
9  
10  
11 632 Cheung, W.W.L., Watson, R. & Pauly, D. (2013) Signature of ocean warming in  
12  
13 633 global fisheries catch. *Nature*, **497**, 365–368.  
14  
15  
16  
17 634 Closset-Kopp, D., Hattab, T. & Decocq, G. (2018) Do drivers of forestry vehicles also  
18  
19 635 drive herb layer changes (1970-2015) in a temperate forest with contrasting habitat  
20  
21 636 and management conditions? *Journal of Ecology*.  
22  
23  
24  
25 637 Davis, M.B. (1984) *Climatic Instability, Time, Lags, and Community*  
26  
27 638 *Disequilibrium. Community ecology* Community ecology. (ed. by J.M. Diamond) and  
28  
29 639 T.J. Case), pp. 269–284. Harper; Row.  
30  
31  
32  
33 640 Davis, M.B. (1989) Lags in vegetation response to greenhouse warming. *Climatic*  
34  
35 641 *Change*, **15**, 75–82.  
36  
37  
38  
39 642 De Frenne, P., Rodríguez-Sánchez, F., Coomes, D.A., Baeten, L., Verstraeten, G.,  
40  
41 643 Vellend, M., ... Verheyen, K. (2013) Microclimate moderates plant responses to  
42  
43 644 macroclimate warming. *Proceedings of the National Academy of Sciences of the*  
44  
45 645 *United States of America*, **110**, 18561–5.  
46  
47  
48  
49 646 De Frenne, P., Zellweger, F., Rodríguez-Sánchez, F., Scheffers, B.R., Hylander, K.,  
50  
51 647 Luoto, M., ... Lenoir, J. (2019) Global buffering of temperatures under forest  
52  
53 648 canopies. *Nature Ecology and Evolution*, **3**, 744–749.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 649 Devictor, V., Julliard, R., Couvet, D. & Jiguet, F. (2008) Birds are tracking climate  
4  
5 650 warming, but not fast enough. *Proceedings of the Royal Society B: Biological*  
6  
7 651 *Sciences*, **275**, 2743–2748.
- 8  
9  
10  
11 652 Devictor, V., Swaay, C. van, Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J.,  
12  
13 653 ... Jiguet, F. (2012) Differences in the climatic debts of birds and butterflies at a  
14  
15 654 continental scale. *Nature Climate Change*, **2**, 121–124.
- 16  
17  
18  
19 655 Dornelas, M., Antão, L.H., Moyes, F., Bates, A.E., Magurran, A.E., Adam, D., ...  
20  
21 656 Zettler, M.L. (2018) BioTIME: A database of biodiversity time series for the  
22  
23 657 Anthropocene. *Global Ecology and Biogeography*, **27**, 760–786.
- 24  
25  
26  
27 658 Dullinger, S., Gattringer, A., Thuiller, W., Moser, D., Zimmermann, N.E., Guisan,  
28  
29 659 A., ... Hülber, K. (2012) Extinction debt of high-mountain plants under twenty-first-  
30  
31 660 century climate change. *Nature Climate Change*, **2**, 619–622.
- 32  
33  
34  
35 661 Duque, A., Stevenson, P.R. & Feeley, K.J. (2015) Thermophilization of adult and  
36  
37 662 juvenile tree communities in the northern tropical Andes. *Proceedings of the*  
38  
39 663 *National Academy of Sciences of the United States of America*, **112**, 10744–10749.
- 40  
41  
42  
43 664 Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W. & Paulißen, D. (1992)  
44  
45 665 Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica*.
- 46  
47  
48  
49 666 Elmendorf, S.C., Henry, G.H.R., Hollister, R.D., Fosaa, A.M., Gould, W.A.,  
50  
51 667 Hermanutz, L., ... Walker, M. (2015) Experiment, monitoring, and gradient methods  
52  
53 668 used to infer climate change effects on plant communities yield consistent patterns.
- 54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 669 *Proceedings of the National Academy of Sciences of the United States of America*,  
4  
5  
6 670 **112**, 448–52.  
7  
8  
9 671 Essl, F., Dullinger, S., Rabitsch, W., Hulme, P.E., Pyšek, P., Wilson, J.R.U. &  
10  
11 672 Richardson, D.M. (2015) Delayed biodiversity change: no time to waste. *Trends in*  
12  
13 673 *Ecology & Evolution*, **30**, 375–378.  
14  
15  
16  
17 674 Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., ... Feeley, K.J.  
18  
19 675 (2018) Widespread but heterogeneous responses of Andean forests to climate  
20  
21 676 change. *Nature*.  
22  
23  
24  
25 677 Fourcade, Y., Åström, S. & Öckinger, E. (2018) Climate and land-cover change alter  
26  
27 678 bumblebee species richness and community composition in subalpine areas.  
28  
29 679 *Biodiversity and Conservation*.  
30  
31  
32  
33 680 Gaüzère, P., Princé, K. & Devictor, V. (2017) Where do they go? The effects of  
34  
35 681 topography and habitat diversity on reducing climatic debt in birds. *Global Change*  
36  
37 682 *Biology*, **23**, 2218–2229.  
38  
39  
40  
41 683 Gégout, J.C., Coudun, C., Bailly, G. & Jabiol, B. (2005) EcoPlant: A forest site  
42  
43 684 database linking floristic data with soil and climate variables. *Journal of Vegetation*  
44  
45 685 *Science*, **16**, 257–260.  
46  
47  
48  
49 686 Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Barančok, P., Benito Alonso,  
50  
51 687 J.L., ... Grabherr, G. (2012) Continent-wide response of mountain vegetation to  
52  
53 688 climate change. *Nature Climate Change*, **2**, 111–115.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 689 **Hijmans, R.J. (2019) raster: Geographic Data Analysis and Modeling.**  
4  
5  
6 690 Hillebrand, H., Soininen, J. & Snoeijs, P. (2010) Warming leads to higher species  
7  
8 691 turnover in a coastal ecosystem. *Global Change Biology*, **16**, 1181–1193.  
9  
10  
11  
12 692 Hooper, D.U., Adair, E.C., Cardinale, B.J., Byrnes, J.E.K., Hungate, B.A., Matulich,  
13  
14 693 K.L., ... Connor, M.I. (2012) A global synthesis reveals biodiversity loss as a major  
15  
16 694 driver of ecosystem change. *Nature*, **486**, 105–108.  
17  
18  
19  
20 695 IPBES (2019) Summary for policymakers of the global assessment report on  
21  
22 696 biodiversity and ecosystem services of the Intergovernmental science-policy platform  
23  
24 697 on biodiversity and ecosystem services.  
25  
26  
27  
28 698 Julve, P. (1998) Baseflor: Index botanique, écologique et chorologique de la flore de  
29  
30 699 France. Version 2009.  
31  
32  
33  
34 700 Jump, A.S. & Peñuelas, J. (2005) Running to stand still: adaptation and the  
35  
36 701 response of plants to rapid climate change. *Ecology Letters*, **8**, 1010–1020.  
37  
38  
39 702 Kaarlejärvi, E., Eskelinen, A. & Olofsson, J. (2013) Herbivory prevents positive  
40  
41 703 responses of lowland plants to warmer and more fertile conditions at high altitudes.  
42  
43 704 *Functional Ecology*, **27**, 1244–1253.  
44  
45  
46  
47 705 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., ...  
48  
49 706 Kessler, M. (2017) Climatologies at high resolution for the earth's land surface  
50  
51 707 areas. *Scientific Data*, **4**, 170122.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 708 Lenoir, J., Gégout, J.C., Dupouey, J.L., Bert, D. & Svenning, J.C. (2010) Forest plant  
4  
5 709 community changes during 1989-2007 in response to climate warming in the Jura  
6  
7 710 Mountains (France and Switzerland). *Journal of Vegetation Science*, **21**, 949–964.  
8  
9  
10  
11 711 Lenoir, J., Graae, B.J., Aarrestad, P.A., Alsos, I.G., Armbruster, W.S., Austrheim,  
12  
13 712 G., ... Svenning, J.C. (2013) Local temperatures inferred from plant communities  
14  
15 713 suggest strong spatial buffering of climate warming across Northern Europe. *Global*  
16  
17 714 *Change Biology*, **19**, 1470–1481.  
18  
19  
20  
21 715 Lenoir, J., Hattab, T. & Pierre, G. (2017) Climatic microrefugia under anthropogenic  
22  
23 716 climate change: implications for species redistribution. *Ecography*, **40**, 253–266.  
24  
25  
26  
27 717 Lenoir, J. & Svenning, J.-C. (2015) Climate-related range shifts - a global  
28  
29 718 multidimensional synthesis and new research directions. *Ecography*, **38**, 15–28.  
30  
31  
32  
33 719 Lenoir, J., Bertrand, R., Comte, L. *et al.* (2020) Species better track climate warming  
34  
35 720 in the oceans than on land. *Nature Ecology Evolution* **4**, 1044–1059.  
36  
37  
38  
39 721 Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B. & Ackerly, D.D.  
40  
41 722 (2009) The velocity of climate change. *Nature*, **462**, 1052–1055.  
42  
43  
44 723 Martin, G., Devictor, V., Motard, E., Machon, N. & Porcher, E. (2019) Short-term  
45  
46 724 climate-induced change in French plant communities. *Biology Letters*, **15**, 20190280.  
47  
48  
49  
50 725 Menéndez, R., Megías, A.G., Hill, J.K., Braschler, B., Willis, S.G., Collingham, Y., ...  
51  
52 726 Thomas, C.D. (2006) Species richness changes lag behind climate change.  
53  
54 727 *Proceedings of the Royal Society B: Biological Sciences*, **273**, 1465–1470.  
55  
56  
57  
58  
59  
60

- 1  
2  
3 728 Nakagawa, S. & Schielzeth, H. (2013) A general and simple method for obtaining R<sup>2</sup>  
4  
5 729 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, **4**,  
6  
7 730 133–142.  
8  
9  
10  
11 731 Nicolas, M., Jolivet, C. & Jonard, M. (2014) How monitoring networks contribute to  
12  
13 732 the understanding and to the management of soil and forest ecosystems? *Revue*  
14  
15 733 *forestière française*.  
16  
17  
18  
19 734 Olofsson, J., Oksanen, L., Callaghan, T., Hulme, P.E., Oksanen, T. & Suominen, O.  
20  
21 735 (2009) Herbivores inhibit climate-driven shrub expansion on the tundra. *Global*  
22  
23 736 *Change Biology*, **15**, 2681–2693.  
24  
25  
26  
27 737 Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.-C., ...  
28  
29 738 Williams, S.E. (2017) Biodiversity redistribution under climate change: Impacts on  
30  
31 739 ecosystems and human well-being. *Science (New York, N.Y.)*, **355**, eaai9214.  
32  
33  
34  
35 740 **Pebesma, E. (2018) Simple Features for R: Standardized Support for Spatial Vector**  
36  
37 741 **Data. *The R Journal*, **10**, 439–446.**  
38  
39  
40  
41 742 Perring, M.P., Bernhardt-Römermann, M., Baeten, L., Midolo, G., Blondeel, H.,  
42  
43 743 Depauw, L., ... Verheyen, K. (2018) Global environmental change effects on plant  
44  
45 744 community composition trajectories depend upon management legacies. *Global*  
46  
47 745 *Change Biology*, **24**, 1722–1740.  
48  
49  
50  
51 746 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team (2019) nlme: Linear  
52  
53 747 and Nonlinear Mixed Effects Models.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 748 Post, E. & Pedersen, C. (2008) Opposing plant community responses to warming  
4  
5 749 with and without herbivores. *Proceedings of the National Academy of Sciences of the*  
6  
7 750 *United States of America*, **105**, 12353–12358.  
8  
9  
10  
11 751 R Core Team (2019) R: A Language and Environment for Statistical Computing.  
12  
13  
14 752 Savage, J. & Vellend, M. (2015) Elevational shifts, biotic homogenization and time  
15  
16 753 lags in vegetation change during 40 years of climate warming. *Ecography*, **38**, 546–  
17  
18 754 555.  
19  
20  
21  
22 755 Schielzeth, H. (2010) Simple means to improve the interpretability of regression  
23  
24 756 coefficients. *Methods in Ecology and Evolution*, **1**, 103–113.  
25  
26  
27  
28 757 Stevens, J.T., Safford, H.D., Harrison, S. & Latimer, A.M. (2015) Forest disturbance  
29  
30 758 accelerates thermophilization of understory plant communities. *Journal of Ecology*,  
31  
32 759 **103**, 1253–1263.  
33  
34  
35  
36 760 Tutin, T.G., Heywood, V.H., Burges, N.A., Valentine, D.H., Walters, S.M. & Webb,  
37  
38 761 D.A. (2001) Flora Europaea 5 Volume Set and CD-ROM Pack.  
39  
40  
41  
42 762 Ulrich, E. (1995) Le réseau RENECOFOR: objectifs et réalisation. *Revue forestière*  
43  
44 763 *française*.  
45  
46  
47 764 Vanderwel, M.C. & Purves, D.W. (2014) How do disturbances and environmental  
48  
49 765 heterogeneity affect the pace of forest distribution shifts under climate change?  
50  
51 766 *Ecography*, **37**, 10–20.  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 767 Vuorinen, K.E.M., Kolstad, A.L., De Vriendt, L., Austrheim, G., Tremblay, J.P.,  
4  
5 768 Solberg, E.J. & Speed, J.D.M. (2020) Cool as a moose: How can browsing counteract  
6  
7  
8 769 climate warming effects across boreal forest ecosystems? *Ecology*, **101**, 1–10.  
9  
10  
11 770 Walther, G.-R. (2010) Community and ecosystem responses to recent climate  
12  
13 771 change. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**,  
14  
15 772 2019–2024.  
16  
17  
18  
19 773 Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag  
20  
21 774 New York.  
22  
23  
24  
25 775 Wickham, H., François, R., Henry, L. & Müller, K. (2019) *dplyr: A Grammar of Data*  
26  
27 776 *Manipulation*.  
28  
29  
30 777 Wilke, C.O. (2019) *cowplot: Streamlined Plot Theme and Plot Annotations for*  
31  
32 778 *“ggplot2.”*  
33  
34  
35  
36 779 Williams, J.J., Bates, A.E. & Newbold, T. (2020) Human-dominated land uses favour  
37  
38 780 species affiliated with more extreme climates, especially in the tropics. *Ecography*,  
39  
40  
41 781 **43**, 391–405.  
42  
43  
44 782 Zellweger, F., Coomes, D., Lenoir, J., Depauw, L., Maes, S.L., Wulf, M., ... De  
45  
46 783 Frenne, P. (2019) Seasonal drivers of understory temperature buffering in  
47  
48 784 temperate deciduous forests across Europe. *Global Ecology and Biogeography*,  
49  
50 785 geb.12991.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 786 Zellweger, F., De Frenne, P., Lenoir, J., Vangansbeke, P., Verheyen, K., Bernhardt-  
4  
5 787 Römermann, M., ... Coomes, D. (2020) Forest microclimate dynamics drive plant  
6  
7  
8 788 responses to warming. *Science*, **368**, 772 LP–775.

9  
10  
11 789 Zuur, A., Ieno, E.N. & Graham, S.M. (2007) *Analyzing Ecological Data*, Springer-  
12  
13 790 Verlag New York.

14  
15  
16  
17 791

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2  
3 **792 Data Accessibility Statement**  
4

5  
6  
7 793 Datasets on CTI values are provided in Supporting Information, and will be  
8  
9 794 archived in an appropriate public repository after the process of review and  
10  
11 795 acceptance of the manuscript.  
12

13  
14  
15 796  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## 797 **Figures**

### 798 *List and captions*

#### 799 **Figure 1.**

800 Workflow implemented in the study. The 102 plots of the RENEFOFOR network are  
801 mapped, and the two datasets used in our analyses are represented in color (in red  
802 and blue for the five-year and yearly datasets, respectively). Analyses of temporal  
803 trends in mean annual temperature (MAT) and the community temperature index  
804 (CTI) and factors contributing to the lag between MAT and CTI (MAT-CTI) are  
805 conducted on the two datasets separately. For more details, see the Materials and  
806 Methods section.

#### 807 **Figure 2.**

808 Temporal trends for mean annual temperature (MAT) and the community  
809 temperature index (CTI) in (a) the five-year dataset from 1995 to 2015 and (b) the  
810 yearly dataset from 1995 to 2017. Linear trends (dashed lines) with standard errors  
811 (ribbons) of estimated MAT or CTI values are shown. For CTI, the values for  
812 enclosure or enclosure are plotted separately. Observed values for temperatures  
813 (MAT or CTI) are represented with their associated standard errors and symbol  
814 sizes are proportional to sample size (number of plots).

#### 815 **Figure 3.**


816 Coefficient estimates and 95% confidence intervals extracted from linear mixed-  
817 effects models testing the relative contribution of several potential abiotic and biotic  
818 determinants on the magnitude and direction of the lag between mean annual

1  
2  
3 819 temperature (MAT) and the community temperature index (CTI) values in (a) the  
4  
5 820 five-year dataset and (b) yearly dataset. Points (with 95% confidence intervals)  
6  
7  
8 821 represent the standardized mean coefficients averaged from the selected models ( $\Delta$   
9  
10 822  $AICc < 2$ ) in the model-averaging procedure. Colors show the magnitude and  
11  
12 823 significance of effects (red: significant amplification of the lag; blue: significant  
13  
14 824 mitigation of the lag; light grey: non-significant).

15  
16  
17  
18 825 **Figure 4.**

19  
20 826 Plots of partial residuals illustrating the contribution of three factors in explaining  
21  
22 827 the magnitude and direction of the lag between mean annual temperature (MAT)  
23  
24 828 and the community temperature index (CTI) for the yearly dataset. These plots  
25  
26 829 indicate the effects of each of the three factors when all others factors in the model  
27  
28 830 are accounted for and set to their mean value for quantitative variables. Qualitative  
29  
30 831 variables were set to deciduous stands and to outside the fence area here. The  
31  
32 832 factors (a) 'Years', (b) 'Stand Basal Area' and (c) 'Stand Age in 1995' were extracted  
33  
34 833 from the best model (in the model-averaging procedure) explaining the lagged  
35  
36 834 responses of forest plant communities in the yearly dataset. Solid lines and ribbons  
37  
38  
39 835 represent the linear trend and associated standard errors, respectively.  
40  
41  
42  
43  
44

45 836  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



837

838

839

Figure 1.




Figure 2.





Figure 3.



846

847

Figure 4.






Figure 1. Workflow implemented in the study. The 102 plots of the RENECOFOR network are mapped, and the two datasets used in our analyses are represented in color (in red and blue for the five-year and yearly datasets, respectively). Analyses of temporal trends in mean annual temperature (MAT) and the community temperature index (CTI) and factors contributing to the lag between MAT and CTI (MAT-CTI) are conducted on the two datasets separately. For more details, see the Materials and Methods section.




Figure 2. Temporal trends for mean annual temperature (MAT) and the community temperature index (CTI) in (a) the five-year dataset from 1995 to 2015 and (b) the yearly dataset from 1995 to 2017. Linear trends (dashed lines) with standard errors (ribbons) of estimated MAT or CTI are shown. For CTI, the values for enclosure or enclosure are plotted separately. Observed values for temperatures (MAT or CTI) are represented with their associated standard errors and symbol sizes are proportional to sample size (number of plots).

926x1296mm (72 x 72 DPI)




Figure 3. Coefficient estimates and 95% confidence intervals extracted from linear mixed-effects models testing the relative contribution of several potential abiotic and biotic determinants on the magnitude and direction of the lag between mean annual temperature (MAT) and the community temperature index (CTI) values in (a) the five-year dataset and (b) yearly dataset. Points (with 95% confidence intervals) represent the standardized mean coefficients averaged from the selected models ( $\Delta AICc < 2$ ) in the model-averaging procedure. Colors show the magnitude and significance of effects (red: significant amplification of the lag; blue: significant mitigation of the lag; light grey: non-significant).

987x1234mm (72 x 72 DPI)




Figure 4. Plots of partial residuals illustrating the contribution of three factors in explaining the magnitude and direction of the lag between mean annual temperature (MAT) and the community temperature index (CTI) for the yearly dataset. These plots indicate the effects of each of the three factors when all others factors in the model are accounted for and set to their mean value for quantitative variables. Qualitative variables were set to deciduous stands and to outside the fence area here. The factors (a) 'Years', (b) 'Stand Basal Area' and (c) 'Stand Age in 1995' were extracted from the best model (in the model-averaging procedure) explaining the lagged responses of forest plant communities in the yearly dataset. Solid lines and ribbons represent the linear trend and associated standard errors, respectively.

740x1358mm (72 x 72 DPI)

## Supplementary Information

### Title:

1. **The climatic debt is growing in the understory of temperate forests: stand characteristics matter**
2. **Running title:** Explaining the climatic debt of temperate forests

### List of appendices (figures and tables):

1. **App. Fig. 1** | Location of the monitored plots throughout France and the time-sequence of resurveys for each plot.
2. **App. Fig. 2** | Comparison between different dataset sources for macroclimate temperature time series.
3. **App. Fig. 3** | Trends in mean annual temperatures anomalies from 1958 to 2017 across the 44 study plots.
4. **App. Fig. 4** | Construction of the Community Temperature Indices (CTIs) and the transfer function.
5. **App. Fig. 5** | Information used to construct the transfer function for calculating CTI values.

- 1  
2  
3 17 6. **App. Fig. 6** | Illustration of the indices created to account for disturbance and  
4  
5 18 forest management.
- 6  
7  
8  
9 19 7. **App. Fig. 7** | Plots of disturbance and forest management indices with data  
10  
11 20 from each studied plot.
- 12  
13  
14 21 8. **App. Fig. 8** | Correlation matrix plots for explanatory variables used in our  
15  
16 22 models.
- 17  
18  
19  
20 23 9. **App. Fig. 9** | Partial residuals plots for variables explaining lags in the best  
21  
22 24 model in the five-year dataset.
- 23  
24  
25 25 10. **App. Fig. 10** | Partial residuals plots for variables explaining lags in the best  
26  
27 26 model in the yearly dataset.
- 28  
29  
30  
31 27 11. **App. Fig. 11** | Average of species richness over time (a), number of species  
32  
33 28 gained or lost between survey years (b), and mean of IV for Temperature over  
34  
35 29 time in group of species gained or lost between survey years (c).
- 36  
37  
38  
39 30 12. **App. Fig. 12** | Trends over time of species richness within each class of IV for  
40  
41 31 Temperature, Light and Nitrogen.
- 42  
43  
44  
45 32 13. **App. Table 1** | Informations on sites characteristics.
- 46  
47  
48 33 14. **App. Table 2** | List of 751 forest species found in the herbaceous layer in our  
49  
50 34 study.

1  
2  
3 35 15. **App. Tables 3 & 4** | Results of the model-averaging procedures and selected  
4  
5 36 models.

6  
7  
8 37 16. **App. Table 5** | Review of some scientific publications studying the  
9  
10 38 thermophilization rates (expressed in °C/decade) in different ecosystems and  
11  
12 39 taxonomic groups.

13  
14  
15  
16 40

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

### 41 *Captions:*

42 **App. Figure 1.** Location of the monitored plots throughout France (a) and the time-  
43 sequence plot for the repeated surveys (b). The two datasets are distinguished by  
44 color (in black and light blue for the five-year dataset and yearly dataset,  
45 respectively). Seasonal relevés are represented by different symbols (◦, Spring; +,  
46 Summer; ×, Autumn) and grouped by year. Labels refer to the names of plots in the  
47 RENECOFOR network. The baseline TerraClimate temperature conditions over the  
48 1960–1990 period is used as background raster.

49 **App. Figure 2.** Comparisons between three sources of time-series for macroclimate  
50 temperatures (TerraClimate, CHELSA and local meteorological stations). For  
51 comparisons, only 13 sites from the RENECOFOR network were used because they  
52 had local meteorological stations nearby. a) Location of the 13 sites from the  
53 RENECOFOR network used for comparisons. b) Time-series of macroclimate  
54 temperatures provided by TerraClimate, CHELSA, and the meteorological stations.  
55 Shaded ribbon (in grey) and dotted line show the beginning of vegetation surveys in  
56 the RENECOFOR monitoring program.

57 **App. Figure 3.** Trends in mean annual temperatures anomalies from 1958 to 2017 in  
58 the 44 permanent plots studied. Temperature anomalies are computed for each  
59 permanent plot as the difference between yearly temperatures and the mean  
60 temperature over the period 1958–2017. The black line depicts a 5-year moving  
61 average of MAT. Mean of temperature anomalies and standard errors are



1  
2  
3 62 represented, red and blue colors correspond to positive and negative values,  
4  
5 63 respectively. Shaded ribbon in grey and arrow show the beginning of vegetation  
6  
7  
8 64 surveys in the RENECOFOR monitoring program.  
9

10  
11 65 **App. Figure 4.** Workflow conducted to calculate the Community Temperature  
12  
13 66 Indices (CTIs) through the transfer function.  
14  
15

16  
17 67 **App. Figure 5.** Supporting information for the construction of Community  
18  
19 68 Temperature Indices (CTIs) using the transfer function.  
20  
21

22 69 a) Venn diagram depicting the number of taxa recorded in the CATMINAT and  
23  
24 70 EcoPlant databases for species indicator value of temperature. Our dataset  
25  
26 71 includes a total of 877 taxa (all strata of vegetation combined).  
27  
28  
29

30 72 b) Relationship between the IVs of temperature of CATMINAT and EcoPlant at  
31  
32 73 the species level.  
33  
34  
35

36 74 c) Map of the sites where the NFI floristic surveys have been performed, and  
37  
38 75 which have been used to model the transfer function.  
39  
40

41 76 d) Hexagonal binning plot between the CTIs calculated with CATMINAT and  
42  
43 77 EcoPlant at the community level, and the transfer function fitted to obtain CTI  
44  
45 78 values from CATMINAT in degrees Celsius.  
46  
47  
48

49 79 *Text details for App. Figure 5 :* To infer temperature from plant species  
50  
51 80 assemblages, we used an approach based on Community Temperature Indices  
52  
53 81 (CTIs) with species Indicator Values (IVs), and a transfer function to model the  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 82 relationship of two databases : CATMINAT (Julve, 1998) and EcoPlant (Gégout et  
4  
5 83 al., 2005). These two databases were chosen, because CATMINAT contains almost  
6  
7  
8 84 all of species recorded in our dataset and EcoPlant, poorer in species, has the  
9  
10 85 advantage to allow the direct computation of CTI values in degrees Celsius.  
11  
12 86 CATMINAT (Julve, 1998) is based on expert knowledge similarly to Ellenberg IVs  
13  
14  
15 87 (Ellenberg et al., 1992), with temperature preferences of plant species coded along  
16  
17 88 an ordinal scale from cold (1) to warm (9) environments. While the CATMINAT  
18  
19  
20 89 database contains almost all of the species sampled in the RENECOFOR monitoring  
21  
22 90 program (858 out of the 877 species recorded within all strata of vegetation; **App.**  
23  
24 91 **Fig. 5a**), CTI values estimated with these species IVs are unitless and direct  
25  
26 92 comparisons with macroclimatic temperatures are not possible. On the other hand,  
27  
28  
29 93 EcoPlant allows the computation of CTI values in degrees Celsius. EcoPlant is a  
30  
31 94 phyto-ecological database which provides information about ecological optimum for  
32  
33 95 key variables of forest plant species based on their distribution over the French  
34  
35 96 territory. In EcoPlant, species IV for temperature are expressed in degrees Celsius  
36  
37 97 as they are estimated using logistic regression models based on climatic data **from**  
38  
39 98 **AURELHY dataset** for the historical period (1961–1990). Nevertheless, the EcoPlant  
40  
41  
42 99 database covers much less species from our dataset (396 out of 877 species).  
43  
44  
45  
46 100 Therefore, we calibrated a transfer function to transform the unitless CTI values  
47  
48 101 from the CATMINAT database into a CTI value expressed in degrees Celsius. By  
49  
50  
51 102 regressing CTI values from EcoPlant against CTI values from CATMINAT, we can  
52  
53  
54 103 obtain predicted CTI values from CATMINAT expressed in degrees Celsius (see  
55  
56  
57  
58  
59  
60

1  
2  
3 104 Lenoir et al., 2013 for a similar approach). To improve the quality of the fit, we used  
4  
5 105 the community level and not directly the species level **App. Fig. 5b-d**).

6  
7  
8 106 To construct the transfer function, we processed as follows:

- 9  
10  
11 107 1. We used an independent dataset of floristic surveys gathered from the French  
12  
13 108 National Forest Inventory (NFI) for the 2005–2016 period to calibrate our  
14  
15 109 transfer function. Homogenization procedures were performed to avoid  
16  
17 110 taxonomic and nomenclatural issues between our dataset, the NFI data and the  
18  
19 111 IV databases. We selected only the floristic surveys from NFI performed during  
20  
21 112 the growing season and surveys recording at least 2 species. This yielded a total  
22  
23 113 of 45 379 surveys covering the whole French territory (**App. Fig. 5c**).
- 24  
25  
26 114 2. Then, we computed the CTI values for each floristic survey both with  
27  
28 115 CATMINAT and EcoPlant, using species presence/absence to give the same  
29  
30 116 weight to common and rare species. The relationship between CTI values from  
31  
32 117 the two IV datasets was also analyzed with species abundance and results were  
33  
34 118 similar (Pearson' correlation between occurrence and abundance data = 0.88,  $t =$   
35  
36 119 386,  $df = 45\ 377$ ,  $P < 0.001$ ).
- 37  
38  
39 120 3. To model the relationship between CTI values from CATMINAT and CTI from  
40  
41 121 EcoPlant, we used a Generalized Additive Model (GAM) with a  
42  
43 122 calibration/training procedure to select the best smoothness parameter  $\lambda$ . The  
44  
45 123 dataset was split into a training set and a test set (i.e. proportion of 70/30,  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 124 respectively). The final GAM was adjusted with  $\lambda = 6$  (model with the lowest  
4  
5 125 RMSE) and it explained 72.9% of the total deviance (**App. Fig. 5d**).

6  
7  
8  
9 126 **App. Figure 6.** Illustrative examples of the indices created to account for disturbance  
10  
11 127 and forest management. Two cases are presented : a) A site with two anthropogenic  
12  
13 128 events that occurred during several vegetation surveys ; and b) A site with one  
14  
15  
16 129 anthropogenic event and two natural events that occurred during several vegetation  
17  
18 130 surveys. For survey year, we recorded informations about : (i) Cumulated disturb.,  
19  
20  
21 131 as the cumulative number of disturbance known in a forest stand since the plot  
22  
23 132 establishment, whatever its nature (anthropogenic or natural event); (ii) the Type of  
24  
25  
26 133 last event (anthropogenic or natural event); (iii) Anteriority, as the anteriority in  
27  
28 134 year of the last disturbance whatever its nature; (iv) the Cumulated disturb. of  
29  
30  
31 135 anthropogenic events : (v) the Cumulated disturb. of natural events. Only the  
32  
33 136 columns filled in grey were used in our analyses. Anthropogenic events include  
34  
35 137 mainly silvicultural operations such as thinning, logging, tree extraction, cleaning  
36  
37  
38 138 brush and forestry machine passage. Natural events include for example, windfalls  
39  
40 139 and tree uprooting after a storm event.

41  
42  
43 140 **App. Figure 7.** Cumulative number of disturbance events along time for each studied  
44  
45  
46 141 plot. The type of disturbance is distinguished by different symbols, and the  
47  
48 142 anteriority (in years) of the last disturbance is shown by a color scale.

49  
50  
51 143 **App. Figure 8.** Correlation matrix plots between explanatory variables used in our  
52  
53  
54 144 analyses, showing scatterplots, density curves and Pearson correlation coefficients  
55  
56 145 for the explanatory variables tested in the models. 'lags': difference MAT – CTI;

1  
2  
3 146 'Years': Year of survey; 'Baseline T°C': the baseline temperature conditions during  
4  
5 147 the 1961–1990 period; 'Age in 1995': Age of forest stand in 1995 (year of the first  
6  
7 148 field campaign); 'Disturb. anterior': Anteriority (in years) of the last disturbance  
8  
9  
10 149 which occurred in the plot ; 'Cum.anthr.disturb.': cumulative sum of the number of  
11  
12 150 annual anthropogenic disturbances; 'Cum.nat.disturb.': cumulative sum of the  
13  
14  
15 151 number of natural disturbances ; 'Basal area': forest stand basal area. See Materials  
16  
17 152 and methods for further details.

19  
20  
21 153 **App. Figure 9.** Partial residuals plots for factors contributing to lags in the best  
22  
23 154 model selected by the model averaging procedure in the five-year dataset. Partial  
24  
25 155 residuals are the residuals that remain after removing the effect of all other  
26  
27  
28 156 covariates in the model.

29  
30  
31 157 **App. Figure 10.** Partial residuals plots for factors contributing to lags in the best  
32  
33 158 model selected by the model averaging procedure in the yearly dataset. Partial  
34  
35 159 residuals are the residuals that remain after removing the effect of all other  
36  
37  
38 160 covariates in the model.

39  
40  
41 161 **App. Figure 11.** Species richness changes over time (a), Number of species gained or  
42  
43 162 lost between survey years (b), and Temporal changes over time for averages of IVs  
44  
45 163 for Temperature within the group of species gained or lost between survey years (c).  
46  
47  
48 164 Points and error bars correspond to the average of values between the 44 studied  
49  
50  
51 165 sites and the standard errors, respectively. For the temporal changes, we compared  
52  
53 166 the species lists between the first survey in 1995 and the later surveys of each plot  
54  
55  
56 167 to identify the pools of species gained and lost between each pair of surveys. In

1  
2  
3 168 particular, these two pools are used to calculate temporal  $\beta$ -diversity between two  
4  
5 169 surveys of a given plot (Legendre, 2019). Then, we computed the average of IVs for  
6  
7  
8 170 Temperature within the gained and lost species.  
9

10  
11 171 **App. Figure 12.** Temporal changes over time in species richness within each class of  
12  
13 172 IV for Temperature, Light and Nitrogen (from CATMINAT, Julve, 1998). Each class  
14  
15 173 of IV are detailed in the online version of CATMINAT  
16  
17 174 (<http://philippe.julve.pagesperso-orange.fr/catminat.htm>). Species richness per  
18  
19 175 classes are expressed in pourcentage of the total species richness in a given plot.  
20  
21 176 Points correspond to the average of species richness between the 44 studied sites,  
22  
23 177 and only significant linear regressions are displayed ( $P < 0.05$ ).  
24  
25  
26  
27

28  
29 178 **App. Table 1** Informations on sites characteristics detailing the name of  
30  
31 179 RENECOFOR plots, the dominant tree species in the forest stands, the altitude, the  
32  
33 180 geographical coordinates (latitude and longitude in WGS84), and the dataset(s) in  
34  
35 181 which they occurred (see Materials and Methods for details on the yearly or five-  
36  
37 182 year datasets).  
38  
39  
40

41 183 **App. Table 2** List of 751 forest species inventoried in the herbaceous layer. Species  
42  
43 184 retained for analyses are marked with a “1” in the éponymous column.  
44  
45  
46


47 185 **App. Tables 3 & 4** Results of the model-averaging procedures and selected models  
48  
49 186 for the study of the effects of different variables on lag (difference between MAT  
50  
51 187 and CTI). Only the top 20 models are presented. Coefficients of variables were  
52  
53 188 estimated using linear mixed-effects models (LMMs including ‘plot’ and ‘nested  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 189 subplots' as random effects) in a model-averaging procedure. All variables were  
4  
5 190 standardized ( $z$ -score) to interpret estimates as relative effect sizes. Coefficient  
6  
7  
8 191 estimates for predictor variables, marginal  $R^2_m$  and conditional  $R^2_c$  are shown.  $R^2_m$   
9  
10 192 and  $R^2_c$  describe the variation explained by fixed effects only and variation  
11  
12  
13 193 explained by the fixed and random effects, respectively. Only models with  $\Delta AIC_c$   
14  
15 194 values  $< 2$  were used to estimate coefficients.


16  
17  
18 195 **App. Table 5** Review of some scientific publications studying the thermophilization  
19  
20 196 rates (expressed in  $^{\circ}\text{C}/\text{decade}$ ) in different ecosystems and taxonomic groups.  
21  
22  
23  
24 197

198 *Appendices:*

a)



b)




199

200 App. Figure 1.


201



a)



b)




Source — TerraClimate — CHELSA — Meteorological Station


202

203 App. Figure 2.

204




App. Figure 3.



208

209 App. Figure 4.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




210  
211 **App. Figure 5.**


view

**Disturbance and Forest management indices**

**a) Illustrative Case for Anthropogenic events (2 events)**



**b) Illustrative Case for Anthropogenic and Natural events (1 and 2 events, respectively)**




213

214 App. Figure 6.

215


Review



App. Figure 7.

Review


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



219

220 App. Figure 8.

221




222

223 App. Figure 9.

224






225

226 App. Figure 10.


227



view

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

App. Figure 11.



231

232 App. Figure 12.

233

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

234 **App. Table 1** cf. Excel file

235 **App. Table 2** cf. Excel files

236 **App. Tables 3 & 4** cf. Excel files

237 **App. Tables 5** cf. Excel files

238

For Peer Review

## 239 References

- 240 Ellenberg, H., Weber, H. E., Düll, R., Wirth, V., Werner, W., & Paulißen, D. (1992).  
241 Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica*.
- 242 Gégout, J. C., Coudun, C., Bailly, G., & Jabiol, B. (2005). EcoPlant: A forest site  
243 database linking floristic data with soil and climate variables. *Journal of Vegetation*  
244 *Science*, 16(2), 257–260. <https://doi.org/10.1111/j.1654-1103.2005.tb02363.x>
- 245 Julve, P. (1998). *Baseflor: Index botanique, écologique et chorologique de la flore de*  
246 *France. Version 2009*. <http://philippe.julve.pagesperso-orange.fr/catminat.htm>
- 247 Legendre, P. (2019). A temporal beta-diversity index to identify sites that have  
248 changed in exceptional ways in space–time surveys. *Ecology and Evolution*, 9(6),  
249 3500–3514. <https://doi.org/10.1002/ece3.4984>
- 250 Lenoir, J., Graae, B. J., Aarrestad, P. A., Alsos, I. G., Armbruster, W. S., Austrheim,  
251 G., Bergendorff, C., Birks, H. J. B., Bråthen, K. A., Brunet, J., Bruun, H. H.,  
252 Dahlberg, C. J., Decocq, G., Diekmann, M., Dynesius, M., Ejrnæs, R., Grytnes, J. A.,  
253 Hylander, K., Klanderud, K., ... Svenning, J. C. (2013). Local temperatures inferred  
254 from plant communities suggest strong spatial buffering of climate warming across  
255 Northern Europe. *Global Change Biology*, 19(5), 1470–1481.  
256 <https://doi.org/10.1111/gcb.12129>

	Site name	Dominant tree species	Altitude	Latitude_WGS84	Longitude_WGS84	Dataset(s)
1						
2						
3	CHP 18	Pedunculate oak	175	46.825	2.573	five-year
4	CHP 59	Pedunculate oak	149	50.171	3.754	five-year
5	CHP 70	Pedunculate oak	240	47.87	6.211	five-year
6						
7	CHS 18	Sessile oak	176	47.254	2.125	five-year
8	CHS 35	Sessile oak	80	48.177	-1.536	five-year
9	CHS 41	Sessile oak	127	47.569	1.259	five-year
10	CHS 51	Sessile oak	180	49.032	4.959	five-year
11	CHS 57a	Sessile oak	315	48.871	6.482	five-year; yearly
12	CHS 57b	Sessile oak	320	49.016	7.461	five-year
13						
14	CHS 68	Sessile oak	256	47.693	7.466	five-year
15	CHS 72	Sessile oak	170	47.796	0.379	five-year
16	CHS 86	Sessile oak	116	46.626	0.494	five-year
17	CHS 88	Sessile oak	330	48.026	6.039	five-year; yearly
18						
19	CPS 67	Sessile or pedunculate oa	350	48.99	7.728	five-year
20	CPS 77	Sessile or pedunculate oa	80	48.454	2.716	five-year
21	EPC 08	Norway spruce	480	49.948	4.809	five-year
22	EPC 63	Norway spruce	950	45.754	2.965	five-year
23	EPC 73	Norway spruce	1700	45.586	6.789	five-year; yearly
24	EPC 74	Norway spruce	1200	46.228	6.348	five-year; yearly
25						
26	EPC 87	Norway spruce	650	45.8	1.816	five-year
27	HET 14	Beech	90	49.182	-0.858	five-year
28	HET 26	Beech	1320	44.917	5.294	five-year; yearly
29	HET 29	Beech	50	47.836	-3.545	five-year
30	HET 30	Beech	1400	44.114	3.542	five-year
31						
32	HET 55	Beech	250	49.17	5.004	five-year
33	HET 60	Beech	138	49.324	2.875	five-year
34	HET 64	Beech	400	43.149	-0.66	five-year
35	HET 65	Beech	850	43.026	0.435	five-year
36	HET 88	Beech	400	48.106	6.244	five-year; yearly
37						
38	MEL 05	European larch	1850	44.704	6.561	five-year
39	PM 17	Maritime pine	15	45.982	-1.273	five-year
40	PM 40a	Maritime pine	27	43.942	-1.352	five-year
41	PM 40c	Maritime pine	150	44.045	-0.004	five-year
42						
43	PM 72	Maritime pine	153	47.746	0.333	five-year
44	PS 35	Scots pine	80	48.201	-1.558	five-year
45	PS 67a	Scots pine	175	48.85	7.71	five-year
46	PS 78	Scots pine	170	48.695	1.732	five-year
47	PS 88	Scots pine	500	48.225	6.695	five-year
48						
49	SP 05	Silver fir	1360	44.489	6.458	five-year; yearly
50	SP 11	Silver fir	950	42.866	2.101	five-year
51	SP 26	Silver fir	1150	44.947	5.331	five-year; yearly
52	SP 38	Silver fir	1100	45.42	6.13	five-year
53	SP 57	Silver fir	400	48.609	7.134	five-year
54	SP 68	Silver fir	680	47.933	7.124	five-year
55						
56						
57						
58						
59						
60						

1	Species	Species retained for analyses
2		
3	<i>Abies alba</i>	1
4	<i>Abies grandis</i>	1
5	<i>Acer campestre</i>	1
6	<i>Acer platanoides</i>	1
7	<i>Acer pseudoplatanus</i>	1
8	<i>Acer sp.</i>	0
9		
10	<i>Aceras anthropophorum</i>	1
11	<i>Achillea macrophylla</i>	1
12	<i>Achillea millefolium</i>	1
13	<i>Aconitum lycoctonum</i> subsp. <i>vulparia</i>	1
14	<i>Aconitum sp.</i>	0
15	<i>Aconitum variegatum</i> subsp. <i>paniculatum</i>	1
16	<i>Actaea spicata</i>	1
17	<i>Adenostyles alliariae</i>	1
18	<i>Adenostyles alpina</i>	1
19	<i>Adoxa moschatellina</i>	1
20	<i>Aegopodium podagraria</i>	1
21	<i>Aetheorhiza bulbosa</i>	1
22	<i>Aethusa cynapium</i> subsp. <i>cynapioides</i>	1
23	<i>Agrimonia eupatoria</i>	1
24	<i>Agrimonia procera</i>	1
25	<i>Agrostis agrostiflora</i>	1
26	<i>Agrostis canina</i>	1
27	<i>Agrostis capillaris</i>	1
28	<i>Agrostis curtisii</i>	1
29	<i>Agrostis sp.</i>	0
30	<i>Agrostis stolonifera</i>	1
31	<i>Ajuga pyramidalis</i>	1
32	<i>Ajuga reptans</i>	1
33	<i>Ajuga sp.</i>	0
34	<i>Alchemilla alpina</i>	1
35	<i>Alchemilla coriacea</i>	1
36	<i>Alchemilla glaucescens</i>	1
37	<i>Alchemilla xanthochlora</i>	1
38	<i>Alliaria petiolata</i>	1
39	<i>Alnus glutinosa</i>	1
40	<i>Alnus viridis</i>	1
41	<i>Alopecurus pratensis</i>	1
42	<i>Anchusa officinalis</i>	1
43	<i>Anemone apennina</i>	1
44	<i>Anemone nemorosa</i>	1
45	<i>Angelica sylvestris</i>	1
46	<i>Antennaria dioica</i>	1
47	<i>Anthoxanthum odoratum</i>	1
48	<i>Anthoxanthum sp.</i>	0
49	<i>Anthriscus sylvestris</i>	1
50	<i>Anthyllis vulneraria</i>	1
51	<i>Aquilegia vulgaris</i>	1
52	<i>Arabis brassica</i>	1

1		
2	<i>Arabis hirsuta</i>	1
3	<i>Arbutus unedo</i>	1
4	<i>Arctium nemorosum</i>	1
5	<i>Arenaria montana</i>	1
6	<i>Arnica montana</i>	1
7		
8	<i>Arrhenatherum elatius</i>	1
9	<i>Arum italicum</i>	1
10	<i>Arum maculatum</i>	1
11	<i>Arum sp.</i>	0
12	<i>Aruncus dioicus</i>	1
13		
14	<i>Aruncus sp.</i>	0
15	<i>Asparagus officinalis</i>	1
16	<i>Asphodelus albus</i>	1
17	<i>Asplenium trichomanes</i>	1
18		
19	<i>Asplenium trichomanes-ramosum</i>	1
20	<i>Astrantia major</i>	1
21	<i>Athyrium distentifolium</i>	1
22	<i>Athyrium filix-femina</i>	1
23	<i>Atropa bella-donna</i>	1
24		
25	<i>Avena sativa</i>	1
26	<i>Avena sp.</i>	0
27	<i>Betula pendula</i>	1
28	<i>Betula pubescens</i>	1
29	<i>Betula sp.</i>	0
30		
31	<i>Blackstonia perfoliata</i>	1
32	<i>Blechnum spicant</i>	1
33	<i>Brachypodium pinnatum</i>	1
34	<i>Brachypodium sp.</i>	0
35	<i>Brachypodium sylvaticum</i>	1
36		
37	<i>Briza media</i>	1
38	<i>Bromus benekenii</i>	1
39	<i>Bromus ramosus</i>	1
40	<i>Bromus sp.</i>	0
41	<i>Buxus sempervirens</i>	1
42		
43	<i>Calamagrostis epigejos</i>	1
44	<i>Calamagrostis sp.</i>	0
45	<i>Calamagrostis varia</i>	1
46	<i>Calamagrostis villosa</i>	1
47	<i>Calamintha grandiflora</i>	1
48	<i>Calamintha sylvatica</i>	1
49		
50	<i>Callitriche sp.</i>	0
51	<i>Calluna vulgaris</i>	1
52	<i>Caltha palustris</i>	1
53	<i>Campanula barbata</i>	1
54	<i>Campanula cervicaria</i>	1
55	<i>Campanula latifolia</i>	1
56	<i>Campanula rapunculoides</i>	1
57	<i>Campanula rhomboidalis</i>	1
58	<i>Campanula rotundifolia</i>	1
59		
60	<i>Campanula trachelium</i>	1



1		
2	<i>Cardamine flexuosa</i>	1
3	<i>Cardamine heptaphylla</i>	1
4	<i>Cardamine hirsuta</i>	1
5	<i>Cardamine impatiens</i>	1
6	<i>Cardamine pentaphyllos</i>	1
7	<i>Cardamine pratensis</i>	1
8	<i>Cardamine sp.</i>	0
9		
10	<i>Carduus carlinifolius</i>	1
11	<i>Carduus defloratus</i>	1
12	<i>Carex acutiformis</i>	1
13	<i>Carex alba</i>	1
14	<i>Carex arenaria</i>	1
15	<i>Carex bohémica</i>	1
16	<i>Carex brizoides</i>	1
17	<i>Carex caryophyllea</i>	1
18	<i>Carex curta</i>	1
19	<i>Carex demissa</i>	1
20	<i>Carex digitata</i>	1
21	<i>Carex divulsa</i>	1
22	<i>Carex echinata</i>	1
23	<i>Carex ferruginea</i> subsp. <i>australpina</i>	1
24	<i>Carex flacca</i>	1
25	<i>Carex flava</i>	1
26	<i>Carex hallerana</i>	1
27	<i>Carex hirta</i>	1
28	<i>Carex laevigata</i>	1
29	<i>Carex montana</i>	1
30	<i>Carex muricata</i>	1
31	<i>Carex muricata</i> subsp. <i>lamprocarpa</i>	1
32	<i>Carex ornithopoda</i>	1
33	<i>Carex ovalis</i>	1
34	<i>Carex pallescens</i>	1
35	<i>Carex panicea</i>	1
36	<i>Carex pendula</i>	1
37	<i>Carex pilosa</i>	1
38	<i>Carex pilulifera</i>	1
39	<i>Carex remota</i>	1
40	<i>Carex sp.</i>	0
41	<i>Carex strigosa</i>	1
42	<i>Carex sylvatica</i>	1
43	<i>Carex umbrosa</i>	1
44	<i>Carlina acaulis</i>	1
45	<i>Carpinus betulus</i>	1
46	<i>Carum verticillatum</i>	1
47	<i>Castanea sativa</i>	1
48	<i>Centaurea montana</i>	1
49	<i>Centaurea nigra</i>	1
50	<i>Centaurea uniflora</i>	1
51	<i>Centaureum erythraea</i>	1
52	<i>Cephalanthera damasonium</i>	1
53		
54		
55		
56		
57		
58		
59		
60		

1		
2	<i>Cephalanthera longifolia</i>	1
3	<i>Cephalanthera rubra</i>	1
4	<i>Cephalanthera</i> sp.	0
5	<i>Cerastium arvense</i>	1
6	<i>Cerastium fontanum</i>	1
7	<i>Cerastium fontanum</i> subsp. <i>vulgare</i>	1
8	<i>Cerastium glomeratum</i>	1
9	<i>Cerastium</i> sp.	0
10	<i>Ceratocapnos claviculata</i>	1
11	<i>Chaerophyllum aureum</i>	1
12	<i>Chaerophyllum hirsutum</i>	1
13	<i>Chaerophyllum</i> sp.	0
14	<i>Chaerophyllum villarsii</i>	1
15	<i>Chenopodium album</i>	1
16	<i>Chenopodium bonus-henricus</i>	1
17	<i>Chrysosplenium alternifolium</i>	1
18	<i>Chrysosplenium oppositifolium</i>	1
19	<i>Cicerbita alpina</i>	1
20	<i>Cicerbita plumieri</i>	1
21	<i>Circaea alpina</i>	1
22	<i>Circaea lutetiana</i>	1
23	<i>Circaea x-intermedia</i>	1
24	<i>Cirsium arvense</i>	1
25	<i>Cirsium eriophorum</i>	1
26	<i>Cirsium oleraceum</i>	1
27	<i>Cirsium palustre</i>	1
28	<i>Cirsium</i> sp.	0
29	<i>Cirsium spinosissimum</i>	1
30	<i>Cirsium vulgare</i>	1
31	<i>Cistus salvifolius</i>	1
32	<i>Clematis alpina</i>	1
33	<i>Clematis vitalba</i>	1
34	<i>Clinopodium vulgare</i>	1
35	<i>Colchicum autumnale</i>	1
36	<i>Conopodium majus</i>	1
37	<i>Convallaria majalis</i>	1
38	<i>Conyza canadensis</i>	1
39	<i>Corallorhiza trifida</i>	1
40	<i>Cornus mas</i>	1
41	<i>Cornus sanguinea</i>	1
42	<i>Cornus</i> sp.	0
43	<i>Corylus avellana</i>	1
44	<i>Cotoneaster nebrodensis</i>	1
45	<i>Crataegus laevigata</i>	1
46	<i>Crataegus monogyna</i>	1
47	<i>Crataegus</i> sp.	0
48	<i>Crepis biennis</i>	1
49	<i>Crepis conyzifolia</i>	1
50	<i>Crepis paludosa</i>	1
51	<i>Cynoglossum germanicum</i>	1

1		
2	<i>Cytisus scoparius</i>	1
3	<i>Cytisus</i> sp.	0
4	<i>Dactylis glomerata</i>	1
5	<i>Dactylis</i> sp.	0
6	<i>Dactylorhiza maculata</i>	1
7	<i>Danthonia decumbens</i>	1
8	<i>Danthonia</i> sp.	0
9	<i>Daphne cneorum</i>	1
10	<i>Daphne gnidium</i>	1
11	<i>Daphne laureola</i>	1
12	<i>Daphne mezereum</i>	1
13	<i>Daucus carota</i>	1
14	<i>Deschampsia cespitosa</i>	1
15	<i>Deschampsia flexuosa</i>	1
16	<i>Digitalis grandiflora</i>	1
17	<i>Digitalis lutea</i>	1
18	<i>Digitalis purpurea</i>	1
19	<i>Draba</i> sp.	0
20	<i>Dryopteris affinis</i>	1
21	<i>Dryopteris affinis</i> subsp. <i>borreri</i>	1
22	<i>Dryopteris carthusiana</i>	1
23	<i>Dryopteris dilatata</i>	1
24	<i>Dryopteris filix-mas</i>	1
25	<i>Dryopteris remota</i>	1
26	<i>Dryopteris</i> sp.	0
27	<i>Elymus caninus</i>	1
28	<i>Elymus</i> sp.	0
29	<i>Epilobium angustifolium</i>	1
30	<i>Epilobium collinum</i>	1
31	<i>Epilobium hirsutum</i>	1
32	<i>Epilobium montanum</i>	1
33	<i>Epilobium obscurum</i>	1
34	<i>Epilobium</i> sp.	0
35	<i>Epilobium tetragonum</i>	1
36	<i>Epipactis atrorubens</i>	1
37	<i>Epipactis helleborine</i>	1
38	<i>Epipactis microphylla</i>	1
39	<i>Epipactis</i> sp.	0
40	<i>Epipogium aphyllum</i>	1
41	<i>Equisetum arvense</i>	1
42	<i>Equisetum palustre</i>	1
43	<i>Equisetum ramosissimum</i>	1
44	<i>Equisetum sylvaticum</i>	1
45	<i>Erica ciliaris</i>	1
46	<i>Erica cinerea</i>	1
47	<i>Erica scoparia</i>	1
48	<i>Erica tetralix</i>	1
49	<i>Erigeron alpinus</i>	1
50	<i>Erigeron annuus</i>	1
51	<i>Euonymus europaeus</i>	1

1		
2	<i>Euonymus latifolius</i>	1
3	<i>Eupatorium cannabinum</i>	1
4	<i>Euphorbia amygdaloides</i>	1
5	<i>Euphorbia cyparissias</i>	1
6	<i>Euphorbia dulcis</i>	1
7	<i>Euphorbia hyberna</i>	1
8	<i>Euphorbia portlandica</i>	1
9	<i>Euphorbia serrulata</i>	1
10	<i>Fagus sylvatica</i>	1
11	<i>Festuca altissima</i>	1
12	<i>Festuca curvula</i>	1
13	<i>Festuca flavescens</i>	1
14	<i>Festuca gigantea</i>	1
15	<i>Festuca glauca</i>	1
16	<i>Festuca heterophylla</i>	1
17	<i>Festuca ovina</i>	1
18	<i>Festuca pratensis</i>	1
19	<i>Festuca rubra</i>	1
20	<i>Festuca rubra</i> subsp. <i>pruinosa</i>	1
21	<i>Festuca</i> sp.	0
22	<i>Festuca tenuifolia</i>	1
23	<i>Festuca trichophylla</i>	1
24	<i>Festuca vasconensis</i>	1
25	<i>Filipendula ulmaria</i>	1
26	<i>Fragaria</i> sp.	0
27	<i>Fragaria vesca</i>	1
28	<i>Frangula alnus</i>	1
29	<i>Fraxinus excelsior</i>	1
30	<i>Fraxinus</i> sp.	0
31	<i>Galeopsis tetrahit</i>	1
32	<i>Galium album</i>	1
33	<i>Galium aparine</i>	1
34	<i>Galium arenarium</i>	1
35	<i>Galium aristatum</i>	1
36	<i>Galium boreale</i>	1
37	<i>Galium glaucum</i>	1
38	<i>Galium mollugo</i>	1
39	<i>Galium odoratum</i>	1
40	<i>Galium palustre</i>	1
41	<i>Galium pumilum</i>	1
42	<i>Galium rotundifolium</i>	1
43	<i>Galium saxatile</i>	1
44	<i>Galium</i> sp.	0
45	<i>Galium sylvaticum</i>	1
46	<i>Galium uliginosum</i>	1
47	<i>Galium verum</i>	1
48	<i>Genista anglica</i>	1
49	<i>Genista pilosa</i>	1
50	<i>Gentiana acaulis</i>	1
51	<i>Gentiana lutea</i>	1

1		
2	<i>Gentiana purpurea</i>	1
3	<i>Gentiana verna</i>	1
4	<i>Gentianella ciliata</i>	1
5	<i>Geranium columbinum</i>	1
6	<i>Geranium nodosum</i>	1
7	<i>Geranium robertianum</i>	1
8	<i>Geranium sylvaticum</i>	1
9	<i>Geum montanum</i>	1
10	<i>Geum rivale</i>	1
11	<i>Geum sp.</i>	0
12	<i>Geum urbanum</i>	1
13	<i>Glechoma hederacea</i>	1
14	<i>Glechoma hirsuta</i>	1
15	<i>Glyceria fluitans</i>	1
16	<i>Glyceria striata</i>	1
17	<i>Gnaphalium sp.</i>	0
18	<i>Goodyera repens</i>	1
19	<i>Gymnocarpium sp.</i>	0
20	<i>Halimium alyssoides</i>	1
21	<i>Hedera helix</i>	1
22	<i>Helictotrichon sedenense</i>	1
23	<i>Helleborus foetidus</i>	1
24	<i>Helleborus viridis</i>	1
25	<i>Helleborus viridis subsp. occidentalis</i>	1
26	<i>Hepatica nobilis</i>	1
27	<i>Heracleum sphondylium</i>	1
28	<i>Hieracium bifidum</i>	1
29	<i>Hieracium juranum</i>	1
30	<i>Hieracium laevigatum</i>	1
31	<i>Hieracium murorum</i>	1
32	<i>Hieracium pilosella</i>	1
33	<i>Hieracium prenanthoides</i>	1
34	<i>Hieracium rapunculoides</i>	1
35	<i>Hieracium sp.</i>	0
36	<i>Hieracium umbellatum</i>	1
37	<i>Hieracium vulgatum</i>	1
38	<i>Hippocrepis comosa</i>	1
39	<i>Holcus lanatus</i>	1
40	<i>Holcus mollis</i>	1
41	<i>Holcus sp.</i>	0
42	<i>Homogyne alpina</i>	1
43	<i>Hordelymus europaeus</i>	1
44	<i>Hordeum vulgare</i>	1
45	<i>Humulus lupulus</i>	1
46	<i>Hyacinthoides non-scripta</i>	1
47	<i>Hypericum androsaemum</i>	1
48	<i>Hypericum hirsutum</i>	1
49	<i>Hypericum humifusum</i>	1
50	<i>Hypericum maculatum</i>	1
51	<i>Hypericum montanum</i>	1

1		
2	<i>Hypericum perforatum</i>	1
3	<i>Hypericum pulchrum</i>	1
4	<i>Hypericum richeri</i>	1
5	<i>Hypericum sp.</i>	0
6	<i>Hypochoeris maculata</i>	1
7	<i>Hypochoeris radicata</i>	1
8	<i>Hypochoeris sp.</i>	0
9	<i>Ilex aquifolium</i>	1
10	<i>Impatiens noli-tangere</i>	1
11	<i>Impatiens sp.</i>	0
12	<i>Inula conyza</i>	1
13	<i>Iris foetidissima</i>	1
14	<i>Isopyrum thalictroides</i>	1
15	<i>Jasione montana</i>	1
16	<i>Juglans regia</i>	1
17	<i>Juncus acutiflorus</i>	1
18	<i>Juncus bufonius</i>	1
19	<i>Juncus bulbosus</i>	1
20	<i>Juncus conglomeratus</i>	1
21	<i>Juncus effusus</i>	1
22	<i>Juncus inflexus</i>	1
23	<i>Juncus sp.</i>	0
24	<i>Juncus tenuis</i>	1
25	<i>Juniperus communis subsp. alpina</i>	1
26	<i>Knautia arvensis</i>	1
27	<i>Knautia dipsacifolia</i>	1
28	<i>Knautia dipsacifolia subsp. gracilis</i>	1
29	<i>Lactuca serriola</i>	1
30	<i>Lactuca virosa</i>	1
31	<i>Lamiastrum galeobdolon</i>	1
32	<i>Lamiastrum sp.</i>	0
33	<i>Lamium album</i>	1
34	<i>Lapsana communis</i>	1
35	<i>Larix decidua</i>	1
36	<i>Laserpitium halleri</i>	1
37	<i>Laserpitium siler</i>	1
38	<i>Laserpitium sp.</i>	0
39	<i>Lathyrus laevigatus subsp. occidentalis</i>	1
40	<i>Lathyrus montanus</i>	1
41	<i>Lathyrus niger</i>	1
42	<i>Lathyrus pratensis</i>	1
43	<i>Lathyrus sp.</i>	0
44	<i>Lathyrus sphaericus</i>	1
45	<i>Lathyrus tuberosus</i>	1
46	<i>Lathyrus vernus</i>	1
47	<i>Leontodon hispidus</i>	1
48	<i>Leontodon pyrenaicus subsp. helveticus</i>	1
49	<i>Leontodon sp.</i>	0
50	<i>Leucanthemum vulgare</i>	1
51	<i>Ligustrum vulgare</i>	1

1		
2	<i>Lilium martagon</i>	1
3	<i>Linaria repens</i>	1
4	<i>Lithospermum officinale</i>	1
5	<i>Lobelia urens</i>	1
6	<i>Lolium perenne</i>	1
7	<i>Lonicera alpigena</i>	1
8	<i>Lonicera caerulea</i>	1
9	<i>Lonicera nigra</i>	1
10	<i>Lonicera periclymenum</i>	1
11	<i>Lonicera xylosteum</i>	1
12	<i>Lotus corniculatus</i>	1
13	<i>Lotus sp.</i>	0
14	<i>Lotus uliginosus</i>	1
15	<i>Lunaria rediviva</i>	1
16	<i>Luzula campestris</i>	1
17	<i>Luzula forsteri</i>	1
18	<i>Luzula luzulina</i>	1
19	<i>Luzula luzuloides</i>	1
20	<i>Luzula multiflora</i>	1
21	<i>Luzula multiflora subsp. congesta</i>	1
22	<i>Luzula nivea</i>	1
23	<i>Luzula pilosa</i>	1
24	<i>Luzula sieberi</i>	1
25	<i>Luzula sp.</i>	0
26	<i>Luzula sylvatica</i>	1
27	<i>Lycopus europaeus</i>	1
28	<i>Lysimachia nemorum</i>	1
29	<i>Lysimachia nummularia</i>	1
30	<i>Lythrum salicaria</i>	1
31	<i>Maianthemum bifolium</i>	1
32	<i>Malus sylvestris</i>	1
33	<i>Marrubium sp.</i>	0
34	<i>Medicago lupulina</i>	1
35	<i>Medicago sp.</i>	0
36	<i>Melampyrum nemorosum</i>	1
37	<i>Melampyrum pratense</i>	1
38	<i>Melampyrum sylvaticum</i>	1
39	<i>Melica nutans</i>	1
40	<i>Melica uniflora</i>	1
41	<i>Melittis melissophyllum</i>	1
42	<i>Mentha aquatica</i>	1
43	<i>Mentha arvensis</i>	1
44	<i>Mercurialis perennis</i>	1
45	<i>Mercurialis sp.</i>	0
46	<i>Mespilus germanica</i>	1
47	<i>Milium effusum</i>	1
48	<i>Minuartia laricifolia</i>	1
49	<i>Moehringia muscosa</i>	1
50	<i>Moehringia trinervia</i>	1
51	<i>Molinia caerulea</i>	1

1		
2	<i>Molinia caerulea</i> subsp. <i>arundinacea</i>	1
3	<i>Monotropa hypopitys</i>	1
4	<i>Mycelis muralis</i>	1
5	<i>Myosotis alpestris</i>	1
6	<i>Myosotis scorpioides</i>	1
7	<i>Myosotis sylvatica</i>	1
8	<i>Myosoton aquaticum</i>	1
9	<i>Narcissus bulbocodium</i>	1
10	<i>Nardus stricta</i>	1
11	<i>Neottia nidus-avis</i>	1
12	<i>Odontites verna</i>	1
13	<i>Orchis purpurea</i>	1
14	<i>Oreopteris limbosperma</i>	1
15	<i>Origanum vulgare</i>	1
16	<i>Ornithogalum pyrenaicum</i>	1
17	<i>Ornithopus perpusillus</i>	1
18	<i>Orobanche lutea</i>	1
19	<i>Orobanche rapum-genistae</i>	1
20	<i>Orobanche</i> sp.	0
21	<i>Orthilia secunda</i>	1
22	<i>Osmunda regalis</i>	1
23	<i>Osyris alba</i>	1
24	<i>Oxalis acetosella</i>	1
25	<i>Oxalis europaea</i>	1
26	<i>Paris quadrifolia</i>	1
27	<i>Pastinaca sativa</i>	1
28	<i>Petasites albus</i>	1
29	<i>Peucedanum gallicum</i>	1
30	<i>Peucedanum ostruthium</i>	1
31	<i>Phalaris arundinacea</i>	1
32	<i>Phegopteris connectilis</i>	1
33	<i>Phleum alpinum</i>	1
34	<i>Phyteuma betonicifolium</i>	1
35	<i>Phyteuma ovatum</i>	1
36	<i>Phyteuma</i> sp.	0
37	<i>Phyteuma spicatum</i>	1
38	<i>Phytolacca americana</i>	1
39	<i>Picea abies</i>	1
40	<i>Picris hieracioides</i>	1
41	<i>Picris</i> sp.	0
42	<i>Pimpinella major</i>	1
43	<i>Pimpinella saxifraga</i>	1
44	<i>Pinus cembra</i>	1
45	<i>Pinus pinaster</i>	1
46	<i>Pinus pinaster</i> subsp. <i>atlantica</i>	1
47	<i>Pinus</i> sp.	0
48	<i>Pinus sylvestris</i>	1
49	<i>Plantago alpina</i>	1
50	<i>Plantago lanceolata</i>	1
51	<i>Plantago major</i>	1



1		
2	<i>Plantago maritima</i>	1
3	<i>Plantago media</i>	1
4	<i>Plantago</i> sp.	0
5	<i>Poa alpina</i>	1
6	<i>Poa annua</i>	1
7	<i>Poa bulbosa</i>	1
8	<i>Poa chaixii</i>	1
9	<i>Poa compressa</i>	1
10	<i>Poa nemoralis</i>	1
11	<i>Poa pratensis</i>	1
12	<i>Poa</i> sp.	0
13	<i>Poa supina</i>	1
14	<i>Poa trivialis</i>	1
15	<i>Poa trivialis</i> subsp. <i>sylvicola</i>	1
16	<i>Polygala serpyllifolia</i>	1
17	<i>Polygonatum multiflorum</i>	1
18	<i>Polygonatum</i> sp.	0
19	<i>Polygonatum verticillatum</i>	1
20	<i>Polygonum bistorta</i>	1
21	<i>Polygonum mite</i>	1
22	<i>Polygonum persicaria</i>	1
23	<i>Polygonum</i> sp.	0
24	<i>Polypodium vulgare</i>	1
25	<i>Polystichum aculeatum</i>	1
26	<i>Polystichum lonchitis</i>	1
27	<i>Polystichum setiferum</i>	1
28	<i>Polystichum</i> sp.	0
29	<i>Populus</i> sp.	0
30	<i>Populus tremula</i>	1
31	<i>Potentilla erecta</i>	1
32	<i>Potentilla grandiflora</i>	1
33	<i>Potentilla reptans</i>	1
34	<i>Potentilla sterilis</i>	1
35	<i>Prenanthes purpurea</i>	1
36	<i>Primula elatior</i>	1
37	<i>Primula veris</i>	1
38	<i>Prunella grandiflora</i>	1
39	<i>Prunella vulgaris</i>	1
40	<i>Prunus avium</i>	1
41	<i>Prunus laurocerasus</i>	1
42	<i>Prunus padus</i>	1
43	<i>Prunus serotina</i>	1
44	<i>Prunus</i> sp.	0
45	<i>Prunus spinosa</i>	1
46	<i>Prunus virginiana</i>	1
47	<i>Pseudarrhenatherum longifolium</i>	1
48	<i>Pseudorchis albida</i>	1
49	<i>Pseudotsuga menziesii</i>	1
50	<i>Pteridium aquilinum</i>	1
51	<i>Pulmonaria affinis</i>	1

1		
2	<i>Pulmonaria longifolia</i>	1
3	<i>Pulmonaria montana</i>	1
4	<i>Pulmonaria obscura</i>	1
5	<i>Pulmonaria saccharata</i>	1
6	<i>Pulsatilla alpina</i> subsp. <i>apiifolia</i>	1
7		
8	<i>Pyrola media</i>	1
9	<i>Pyrola minor</i>	1
10	<i>Pyrus pyraeaster</i>	1
11	<i>Quercus ilex</i>	1
12	<i>Quercus petraea</i>	1
13	<i>Quercus pubescens</i>	1
14	<i>Quercus pyrenaica</i>	1
15	<i>Quercus pyrenaica</i>	1
16	<i>Quercus robur</i>	1
17	<i>Quercus rubra</i>	1
18	<i>Quercus sp.</i>	0
19		
20	<i>Quercus suber</i>	1
21	<i>Ranunculus aconitifolius</i>	1
22	<i>Ranunculus acris</i>	1
23	<i>Ranunculus aduncus</i>	1
24	<i>Ranunculus auricomus</i>	1
25	<i>Ranunculus ficaria</i>	1
26	<i>Ranunculus flammula</i>	1
27	<i>Ranunculus flammula</i>	1
28	<i>Ranunculus lanuginosus</i>	1
29	<i>Ranunculus montanus</i>	1
30	<i>Ranunculus platanifolius</i>	1
31	<i>Ranunculus platanifolius</i>	1
32	<i>Ranunculus pyrenaicus</i>	1
33	<i>Ranunculus repens</i>	1
34	<i>Ranunculus serpens</i> subsp. <i>nemorosus</i>	1
35	<i>Ranunculus sp.</i>	0
36		
37	<i>Rhamnus catharticus</i>	1
38	<i>Rhododendron ferrugineum</i>	1
39	<i>Ribes alpinum</i>	1
40	<i>Ribes rubrum</i>	1
41	<i>Ribes uva-crispa</i>	1
42	<i>Rosa arvensis</i>	1
43	<i>Rosa arvensis</i>	1
44	<i>Rosa canina</i>	1
45	<i>Rosa pendulina</i>	1
46	<i>Rosa pimpinellifolia</i>	1
47	<i>Rosa villosa</i>	1
48	<i>Rubia peregrina</i>	1
49	<i>Rubia peregrina</i>	1
50	<i>Rubus caesius</i>	1
51	<i>Rubus canescens</i>	1
52	<i>Rubus fruticosus</i>	1
53	<i>Rubus idaeus</i>	1
54	<i>Rubus idaeus</i>	1
55	<i>Rubus questieri</i>	1
56	<i>Rubus saxatilis</i>	1
57	<i>Rubus sp.</i>	0
58	<i>Rubus ulmifolius</i>	1
59	<i>Rubus vestitus</i>	1
60	<i>Rumex acetosa</i>	1

1		
2	<i>Rumex acetosella</i>	1
3	<i>Rumex alpestris</i>	1
4	<i>Rumex alpinus</i>	1
5	<i>Rumex obtusifolius</i>	1
6	<i>Rumex sanguineus</i>	1
7	<i>Rumex scutatus</i>	1
8	<i>Rumex sp.</i>	0
9	<i>Ruscus aculeatus</i>	1
10	<i>Salix atrocineria</i>	1
11	<i>Salix caprea</i>	1
12	<i>Salix cinerea</i>	1
13	<i>Salix elaeagnos</i>	1
14	<i>Salix repens</i>	1
15	<i>Salix sp.</i>	0
16	<i>Sambucus ebulus</i>	1
17	<i>Sambucus nigra</i>	1
18	<i>Sambucus racemosa</i>	1
19	<i>Sambucus sp.</i>	0
20	<i>Sanguisorba minor</i>	1
21	<i>Sanicula europaea</i>	1
22	<i>Saxifraga cuneifolia</i> subsp. <i>robusta</i>	1
23	<i>Saxifraga granulata</i>	1
24	<i>Saxifraga rotundifolia</i>	1
25	<i>Scilla bifolia</i>	1
26	<i>Scirpus sylvaticus</i>	1
27	<i>Scorzonera humilis</i>	1
28	<i>Scrophularia alpestris</i>	1
29	<i>Scrophularia nodosa</i>	1
30	<i>Scutellaria minor</i>	1
31	<i>Scutellaria sp.</i>	0
32	<i>Sedum anacampseros</i>	1
33	<i>Sedum montanum</i>	1
34	<i>Sempervivum arachnoideum</i>	1
35	<i>Sempervivum tectorum</i>	1
36	<i>Senecio adonidifolius</i>	1
37	<i>Senecio jacobaea</i>	1
38	<i>Senecio nemorensis</i> subsp. <i>fuchsii</i>	1
39	<i>Senecio sp.</i>	0
40	<i>Senecio sylvaticus</i>	1
41	<i>Senecio viscosus</i>	1
42	<i>Senecio vulgaris</i>	1
43	<i>Serratula tinctoria</i>	1
44	<i>Seseli montanum</i>	1
45	<i>Sesleria albicans</i>	1
46	<i>Setaria sp.</i>	0
47	<i>Silene dioica</i>	1
48	<i>Silene nutans</i>	1
49	<i>Silene rupestris</i>	1
50	<i>Silene vulgaris</i>	1
51	<i>Simethis planifolia</i>	1
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2	<i>Sison amomum</i>	1
3	<i>Sisymbrium orientale</i>	1
4	<i>Solanum dulcamara</i>	1
5	<i>Solanum nigrum</i>	1
6	<i>Soldanella alpina</i>	1
7	<i>Solidago canadensis</i>	1
8	<i>Solidago gigantea</i>	1
9	<i>Solidago sp.</i>	0
10	<i>Solidago virgaurea</i>	1
11	<i>Sonchus arvensis</i>	1
12	<i>Sonchus asper</i>	1
13	<i>Sonchus oleraceus</i>	1
14	<i>Sonchus sp.</i>	0
15	<i>Sorbus aria</i>	1
16	<i>Sorbus aucuparia</i>	1
17	<i>Sorbus domestica</i>	1
18	<i>Sorbus sp.</i>	0
19	<i>Sorbus torminalis</i>	1
20	<i>Stachys alpina</i>	1
21	<i>Stachys officinalis</i>	1
22	<i>Stachys sylvatica</i>	1
23	<i>Stellaria graminea</i>	1
24	<i>Stellaria holostea</i>	1
25	<i>Stellaria media</i>	1
26	<i>Stellaria nemorum</i>	1
27	<i>Stellaria uliginosa</i>	1
28	<i>Succisa pratensis</i>	1
29	<i>Tamus communis</i>	1
30	<i>Taraxacum officinale</i>	1
31	<i>Taraxacum sp.</i>	0
32	<i>Taxus baccata</i>	1
33	<i>Teucrium scorodonia</i>	1
34	<i>Thalictrum aquilegifolium</i>	1
35	<i>Thalictrum sp.</i>	0
36	<i>Thesium alpinum</i>	1
37	<i>Thlaspi alpestre</i>	1
38	<i>Tilia cordata</i>	1
39	<i>Tilia platyphyllos</i>	1
40	<i>Tilia platyphyllos</i> subsp. <i>cordifolia</i>	1
41	<i>Torilis japonica</i>	1
42	<i>Tragopogon pratensis</i>	1
43	<i>Trifolium alpestre</i>	1
44	<i>Trifolium alpinum</i>	1
45	<i>Trifolium badium</i>	1
46	<i>Trifolium medium</i>	1
47	<i>Trifolium pratense</i>	1
48	<i>Trifolium repens</i>	1
49	<i>Trifolium sp.</i>	0
50	<i>Trisetum flavescens</i>	1
51	<i>Trochiscanthes nodiflora</i>	1
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2	<i>Trollius europaeus</i>	1
3	<i>Tussilago farfara</i>	1
4	<i>Ulex europaeus</i>	1
5	<i>Ulex minor</i>	1
6	<i>Ulmus glabra</i>	1
7	<i>Ulmus minor</i>	1
8	<i>Urtica dioica</i>	1
9	<i>Vaccinium myrtillus</i>	1
10	<i>Vaccinium vitis-idaea</i>	1
11	<i>Valeriana montana</i>	1
12	<i>Valeriana officinalis</i>	1
13	<i>Valeriana officinalis</i> subsp. <i>sambucifolia</i>	1
14	<i>Valeriana tripteris</i>	1
15	<i>Veratrum album</i>	1
16	<i>Verbascum lychnitis</i>	1
17	<i>Verbascum nigrum</i>	1
18	<i>Verbascum sinuatum</i>	1
19	<i>Verbascum thapsus</i>	1
20	<i>Veronica anagallis-aquatica</i>	1
21	<i>Veronica beccabunga</i>	1
22	<i>Veronica chamaedrys</i>	1
23	<i>Veronica hederifolia</i>	1
24	<i>Veronica hederifolia</i> subsp. <i>lucorum</i>	1
25	<i>Veronica montana</i>	1
26	<i>Veronica officinalis</i>	1
27	<i>Veronica serpyllifolia</i>	1
28	<i>Veronica serpyllifolia</i> subsp. <i>humifusa</i>	1
29	<i>Veronica</i> sp.	0
30	<i>Veronica urticifolia</i>	1
31	<i>Viburnum lantana</i>	1
32	<i>Viburnum opulus</i>	1
33	<i>Vicia cracca</i>	1
34	<i>Vicia disperma</i>	1
35	<i>Vicia hirsuta</i>	1
36	<i>Vicia pisiformis</i>	1
37	<i>Vicia sativa</i>	1
38	<i>Vicia sativa</i> subsp. <i>nigra</i>	1
39	<i>Vicia sepium</i>	1
40	<i>Vicia</i> sp.	0
41	<i>Vicia sylvatica</i>	1
42	<i>Vicia tetrasperma</i>	1
43	<i>Vinca minor</i>	1
44	<i>Vincetoxicum hirundinaria</i> subsp. <i>intermediur</i>	1
45	<i>Viola biflora</i>	1
46	<i>Viola canina</i>	1
47	<i>Viola hirta</i>	1
48	<i>Viola mirabilis</i>	1
49	<i>Viola odorata</i>	1
50	<i>Viola reichenbachiana</i>	1
51	<i>Viola riviniana</i>	1

1		
2	Viola sp.	0
3	Viola tricolor	1
4		
5		
6		
7		
8		
9		
10		
11		
12		
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

For Peer Review

	(Intercept)	Age_1995	Disturbanc	Years	Stand_Type	Exclosure	eCumulated	Cumulated	Stand basa
3	0.838	0.145	-0.115	0.058			-0.17		0.216
4	0.838	0.152	-0.113	0.059			-0.167	-0.042	0.207
5	0.811	0.16	-0.113	0.055	+		-0.165		0.208
6	0.83	0.145	-0.115	0.058		+	-0.17		0.216
8	0.805	0.17	-0.11	0.055	+		-0.161	-0.045	0.196
9	0.838	0.121	-0.087				-0.114		0.195
10	0.83	0.152	-0.113	0.059		+	-0.167	-0.042	0.207
11	0.838	0.127	-0.085				-0.111	-0.04	0.185
12	0.797	0.145	-0.087		+		-0.111		0.184
14	0.803	0.16	-0.113	0.055	+	+	-0.165		0.208
15	0.792	0.154	-0.084		+		-0.107	-0.044	0.172
16	0.797	0.17	-0.11	0.055	+	+	-0.161	-0.045	0.196
17	0.83	0.121	-0.087			+	-0.114		0.195
18	0.83	0.127	-0.085			+	-0.111	-0.04	0.185
20	0.789	0.145	-0.087		+	+	-0.111		0.184
21	0.784	0.154	-0.084		+	+	-0.107	-0.044	0.172
22	0.837		-0.084				-0.09		0.185
23	0.837		-0.099	0.033			-0.118		0.196
24	0.877		-0.104	0.041	+		-0.133		0.211
25	0.862		-0.085		+		-0.094		0.193

For Peer Review





References	Years	Journal	Methods to compute the CTI
Bertrand et al.	2011	Nature	CTI reconstructed using a combination of weighted averaging partial least squares and Breiman's random forest with transfer function
Brice et al.	2019	Global Ecology and Biogeography	CTI computed with the average of species temperature distributions (i.e. Wordclim) based on tree occurrence data (Forest inventory databases of eastern North America )
Cheung et al.	2013	Nature	CTI based on the average of temperature preferences of marine fishes and invertebrates in fisheries catch
De Frenne et al.	2013	PNAS	CTI computed with temperature preference distributions of species (during growing season April to September) in permanent or semipermanent plots
Devictor et al.	2008	PNAS	CTI calculated as the average of species thermal preferences
Devictor et al.	2012	Nature Climate Change	CTI calculated as the average of species thermal preferences
Devictor et al.	2012	Nature Climate Change	CTI calculated as the average of species thermal preferences
Fadrique et al.	2018	Nature	CTI calculated as the average of species thermal preferences
Flanagan et al.	2019	Ecography	CTI calculated as the average of species thermal preferences
Fourcade et al.	2018	Biodiversity and Conservation	CTI calculated as the average of species thermal preferences
Gotfried et al.	2012	Nature Climate Change	Temperature index with ecological indicator
Haase et al.	2019	Science of the Total Environment	CTI calculated as the average of species thermal preferences
Lindström et al.	2013	Ecography	CTI calculated as the average of species thermal preferences

1				
2	Savage et al.	2015	Ecography	CTI calculated as the average of
3				species thermal preferences
4	Elmendorf et al.	2015	PNAS	CTI calculated as the average of
5				species thermal preferences ;
6				Repeated sampling of historical
7				studies and warming experiments
8				
9				
10				
11	Duque et al.	2015	PNAS	CTI calculated as the average of
12				species thermal preferences
13				
14				
15	Zellweger et al.	2020	Science	CTI calculated as the average of
16				species thermal preferences
17				
18				
19				
20				
21				
22				
23				
24				
25				
26				
27				
28				
29				
30				
31				
32				
33				
34				
35				
36				
37				
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				

For Peer Review

**Methods to compute the velocity of changes**

	<b>Period</b>
Differences between CTI from between 1965-1986 and 1987-2008 periods	1965-1986 and 1987-2008
Comparison between historical and contemporary periods	1970 - 2016
Slope of linear regression	1970 - 2006
Comparison between historical and contemporary periods	depending on forest plots and regions; overall, from 1940 (minimum) to 2009 (maximum)
Slope of linear regression	1989 - 2006
Slope of linear regression	1990 - 2008
Slope of linear regression	1990 - 2008
Slope of linear regression	2000 - 2015
Slope of linear regression	1990 - 2014
Comparison between historical and contemporary periods	1960 - 2011
Comparison between historical and contemporary periods	2001 - 2008
Slope of linear regression	1990 - 2014
Slope of linear regression	1975 - 2009

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Comparison between historical and contemporary periods	1970 - 2012
Slope of linear regression	1970 - 2012

Rates calculated as the annualized change in the CTI values for each plot over the entire census period	2006-2014
---	-----------

Comparison between baseline surveys and resurveys	depending on forest plots and regions; overall, from 1934 (minimum) to 2017 (maximum) ; median time interval between two surveys of 38 years
---	--

For Peer Review

Taxonomic groups	Regions	Thermophilization rates
Forest understorey vegetation	France	$\Delta 0.02^{\circ}\text{C}$ and $\Delta 0.54^{\circ}\text{C}$ between 1965-1986 and 1987-2008 in lowland and highland, respectively.
Tree communities	Canada, Québec	$0.03^{\circ}\text{C}$ / decade
Marine fishes and invertebrates	Global	$0.19^{\circ}\text{C}$ / decade
Forest understorey vegetation	29 regions of Europe and North America	$0.041^{\circ}\text{C}$ / decade
Birds	France	$0.044^{\circ}\text{C}$ / decade
Birds	Europe	$0.044^{\circ}\text{C}$ / decade
Butterflies	Europe	$0.051^{\circ}\text{C}$ / decade
palms, tree ferns and lianas (Trees (including	The Andes (Peru, Colombia, Argentina, Ecuador)	$0.066^{\circ}\text{C}$ / decade
Marine fishes and invertebrates	Northeast U.S. continental shelf	in fall: $0.25^{\circ}\text{C}$ / decade; in spring: $0.38^{\circ}\text{C}$ / decade
Bumblebees	Norway	$0.14^{\circ}\text{C}$ / decade
Mountain vegetation	Europe	-
Stream invertebrate communities	Central Europe	$0.22^{\circ}\text{C}$ / decade (abundance data); $0.15^{\circ}\text{C}$ / decade (occurrence data)
Birds	Norway	$0.031^{\circ}\text{C}$ / decade (Old monitoring scheme); $0.096^{\circ}\text{C}$ / decade (New monitoring scheme)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Forest understorey vegetation	Canada, Québec	0.05° C / decade
Tundra vegetation	Canada, Québec	0.199° C / decade
Tree communities	Northern tropical Andes	adult trees: 0.11°C / decade; juvenile trees: 0.27°C / decade
Forest understorey vegetation	Europe	0.01°C / decade

For Peer Review