

Interaction of half-sandwich alkylmolybdenum(III) complexes with B(C6F5)3. The X-ray structure of $[CpMo(\eta 4-C4H6)(\mu-Cl)(\mu-CH2)(O)MoCp][CH3B(C6F5)3]$

Erwan Le Grognec, Rinaldo Poli, Philippe Richard, Rosa Llusar, Santiago

Uriel

▶ To cite this version:

Erwan Le Grognec, Rinaldo Poli, Philippe Richard, Rosa Llusar, Santiago Uriel. Interaction of half-sandwich alkylmolybdenum(III) complexes with B(C6F5)3. The X-ray structure of [CpMo(η 4-C4H6)(μ -Cl)(μ -CH2)(O)MoCp][CH3B(C6F5)3]. Journal of Organometallic Chemistry, 2001, 640 (1-2), pp.113-120. 10.1016/S0022-328X(01)01196-2. hal-03294431

HAL Id: hal-03294431 https://hal.science/hal-03294431

Submitted on 28 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Interaction of Half-Sandwich Alkylmolybdenum(III) Complexes with B(C₆F₅)₃. The X-ray Structure of [CpMo(η⁴-C₄H₆)(μ-Cl)(μ-CH₂)(O)MoCp][CH₃B(C₆F₅)₃]

Erwan Le Grognec,^a Rinaldo Poli^{*a}, Philippe Richard,^a Rosa Llusar,^b and Santiago Uriel^b

^aLaboratoire de Synthèse et d'Electrosynthèse Organométalliques, Faculté des Sciences "Gabriel", Université de Bourgogne, 6 Boulevard Gabriel, 21000 Dijon, France

^b Departament de Ciències Experimentals, Campus Riu Sec de la Universitat Jaume I, POB 224, 12080 Castelló, Spain

^{*} To whom correspondence should be addressed.

Tel: +33-3-80-39-68-81

Fax: +33-3-80-39-60-98

E-mail: Rinaldo.Poli@u-bourgogne.fr

Graphical Contents Entry



Summary

The reactions of the half-sandwich molybdenum(III) complexes $CpMo(\eta^4$ -

 $C_4H_4R_2)(CH_3)_2$, where $Cp = \eta^5 - C_5H_5$ and R = H or CH_3 , with equimolar amounts of $B(C_6F_5)_3$ have been investigated in toluene. EPR monitoring shows the formation of an addition product which does not readily react with Lewis bases such as ethylene, pyridine, or PMe₃. The analysis of the EPR properties and the X-ray structure of a decomposition product obtained from dichloromethane, $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp][CH_3B(C_6F_5)_3]$, indicate that the borane attack has occurred at the methyl position.

Introduction

Polymerization of α -olefins is one of the most important industrial processes. The combination of group 4 metallocenes and methylaluminoxane (MAO) has been actively investigated in view of its homogeneous process and the possibility to control the stereoregularity of the polymerization though the control of the ligand architecture.¹⁻⁵ A few studies have also been carried out with isoelectronic group 5 metal complexes such as $CpNb(\eta^4-diene)R_2^{6,7}$ and Tp'Nb(η^2 -alkyne)R₂ (Tp' = substituted hydrotris(pyrazolyl)borate).⁸ The prevalent concept of these works is the analogy (structure and electron count) to the architecture of the group 4 metallocenes, where the 14 electron metallocene alkyl cation $[Cp_2MR]^+$ is the proposed active species.^{1, 3, 4} By analogy with the work done on the group 5 metals, we considered the possibility that the half-sandwich 17-electrons molybdenum(III) complexes available in our laboratory99, 10 could also catalyze ethylene polymerization. Our initial experimental and theoretical studies of these complexes in ethylene polymerization catalysis have been recently reported.¹¹ In combination with methylaluminoxane, very low activities has been observed to yield polymers of high linearity and very high molecular weights. The theoretical studies have identified the low polarity of the Mo-CH₃ bonds, as compared with that of the Nb-CH₃ bonds in the analogous Nb(III) derivatives, as the cause of the low activity because the activation process is made less favorable. Other strong organo-Lewis acids such as $B(C_6F_5)_3$ have also been used to generate the active species in situ.¹² Under the assumption that the active specie is the $[CpMo(\eta^4-diene)R]^+$ complex, we were interested in probing the reactivity of the dimethyl complexes $CpMo(\eta^4$ diene)Me₂ and Cp^{*} analogues with the methyl abstracting agent $B(C_6F_5)_3$ in order to evaluate the

possibility to isolate the putative active species. The results of these investigations are reported in this contribution.

Results and Discussion

The addition of one equivalent of B(C₆F₅)₃ to solutions of CpMo(η^4 -diene)Me₂ (diene = 1,3-butadiene or 2,3-dimethyl-1,3-butadiene) in toluene at -20° C leads to a rapid color change from green to orange. Analogous color changes were also observed upon treating the same Mo complexes with MAO.¹¹ In each case, the product appears stable at room temperature in hydrocarbon solvents and under an inert atmosphere, at least for a few days. Unfortunately, all our crystallization attempts (see Experimental) have yielded oily products. Therefore, the only information available to us on the nature of the reaction products comes from solution spectroscopic studies. Monitoring the reactions by EPR at -20°C shows that, in each case, the characteristic pseudo-binomial 11-line pattern of the dimethyl Mo(III) complex,¹⁰ due to the apparent equivalence of 10 hydrogen atoms of the molecule, disappears and is replaced by a new, broader signal at g = 1.995. Both starting and final EPR spectra are essentially independent on the nature of the diene ligand, since the nature of the substitution of the diene ligand does not influence the g value nor the multiplicity of the spectrum. Figure 1 shows the example of the butadiene system. The new signal is slightly broader than that of the starting complex and is characterized by a different coupling pattern.

<Figure 1>

Several adducts of B(C₆F₅)₃ with transition metal complexes have been characterized spectroscopically and, in a few cases, also by X-ray crystallography.¹³⁻¹⁵ The borane reagent unfailingly attacks an alkyl group whenever at least one is present, to generate a M···R···B(C₆F₅)₃ moiety where the M-R bonding interaction is greatly weakened. In the presence of Lewis bases L (*i.e.*, THF, pyridine, phosphines), sometimes even very weak ones such as organic halides, the alkyl is typically abstracted resulting in charge separation, to generate $[M(L)]^+[R-B(C_6F_5)_3]^-$ species.¹⁵⁻¹⁸ However, boranes have also been shown to give rise to electrophilic additions to the terminal carbon atom of a coordinated diene to yield complexes containing a betaine allylic ligand, *e.g.* M[η³-C₃H₄-CH₂B(C₆F₅)₃] starting from a M(η⁴-C₄H₆) complex.¹⁹⁻²⁴ Therefore, the question of the site of attack (methyl or diene ligand) for complexes CpMo(η⁴-C₄H₆)Me₂ is not a trivial one *a priori*. A good spectroscopic handle on the nature of the interaction is provided by ¹¹B and ¹⁹F NMR, but this is unfortunately prevented in our case by the paramagnetism. The inability to observe a ¹⁹F NMR signal suggests, in fact, that the borane molecule is covalently linked to the paramagnetic Mo(III) complex.

The digital fitting of the EPR experimental spectrum provides useful information on the site of attack. The fitting was carried out only for the butadiene system, for which a slightly improved spectral resolution allowed a more accurate analysis. The fitting gives the best agreement for an $A_3M_3R_2X_2$ spin system (see Experimental section). This is interpreted as resulting from two sets of three equivalent H atoms from the two chemically inequivalent methyl groups, plus two sets of two pseudo-equivalent H atoms from the two chemically distinct diene terminal (*syn* and *anti*) positions. Previous EPR studies of paramagnetic diene complexes have systematically shown coupling to the terminal *syn* and *anti* H atoms and no observable coupling to the internal H atoms.^{9, 10, 25-27} Thus, the observed spectral modifications suggest that an interaction has occurred between the Lewis acid B(C₆F₅)₃ and one methyl group. The

nonequivalence of the two methyl groups would then arise from the fact that one is bonded only to the Mo atom whereas the other one is bonded to the Mo and B atoms. This asymmetry should make the two opposite termini of the diene ligand also nonequivalent, giving rise to a RSXY system rather than a R_2X_2 system for the diene H atoms. However, these atoms are far removed from the source of the asymmetry, yielding pseudo-equivalence, at least within the resolution limits of our spectrometer. Indeed, the quality of the fit significantly increases upon breaking the symmetry from 64 to 334 (the correlation increases from 0.990 to 0.992 and the error decreases from 1.743 10⁻² to 1.342 10⁻²). An almost insignificant improvement takes place upon further breaking the symmetry to 3322, because of the accidental equivalence of a_H to syn and anti diene protons. Previous studies have also highlighted the near-degeneracy of the hyperfine couplings to these two types of atoms.^{10, 26, 27} Further lowering the symmetry to a 331111 spin system does not further improve the quality of the fit. The alternative interpretation of a borane attack at one of the diene termini seems less likely for two reasons. First, the diene terminal H spin system would be expected to show greater asymmetry. Second, and more importantly, the resulting betaine allyl ligand would show coupling to the allylic terminal H atoms (thus 3 rather than 4 H atoms), according to the available precedents for similar allylmolybdenum(III) complexes.^{27, 28} Attempts to simulate the EPR spectrum on the basis of a 33111 model, however, systematically gave unacceptable fits. It is to be noted that the a_H values for the two inequivalent methyl groups (6.79 and 4.42 G) are not too different from each other, suggesting that the borane-bonded methyl group is still solidly attached to the Mo center. This is consistent with the observed reactivity (or lack thereof), vide infra. For comparison, the a_H value for the methyl H atoms in dimethylmolybdenym(III) complexes are in the 5.3-6.3 G range, depending on the nature of the diene and cyclopentadienyl ligands.^{10, 29} The increased broadness of the spectrum relative to that of the starting dimethyl complex may be related to the onset of a dynamic exchange of the two

methyl groups between the borane-bonded and nonbonded positions. A similar phenomenon has been shown to broaden the ¹H NMR resonances for diamagnetic compounds having the $M \cdots R \cdots B(C_6F_5)_3$ moiety.^{13, 30, 31} The EPR spectrum recorded at different temperatures (both lower and higher) has a poorer resolution and no useful information could be obtained from its analysis.

Additional indication for the occurrence of a methyl rather than diene attack comes from previous computational investigations, showing that the Mulliken charge on the methyl ligands is more negative than that on the diene terminal position.¹¹ This is true also for the corresponding Nb system, for which the formation of the $[CpNb(\eta^4-diene)Me]^+[Me-MAO]^-$ active site in ethylene polymerization catalysis has been proposed.^{7, 32} Finally, the borane attack at the methyl group for compound $CpMo(\eta^4-C_4H_6)Me_2$ is indicated by the characterization of the title compound, which results from a decomposition process (*vide infra*).

When the addition of CpMo(η^4 -C₄H₆)Me₂ to the B(C₆F₅)₃ solution was followed by the addition of a slight excess of PMe₃, no spectral modifications were observed by EPR. This surprising observation shows that the phosphine molecule is incapable of displacing the methyl group from either the boron atom [giving back the starting dimethyl Mo(III) complex and Me₃PB(C₆F₅)₃] or the molybdenum atom. The latter process would afford [CpMo(η^4 -C₄H₆)-(PMe₃)Me][MeB(C₆F₅)₃] by substitution of the labile [CH₃B(C₆F₅)₃]⁻ ligand with PMe₃ and should dramatically affect the shape of the EPR spectrum (notably a strong coupling to the P nucleus would be expected). A reaction of this type is undertaken, for instance, by the boratabenzene chromium complex (C₃H₅BMe)Cr(Me){MeB(C₆F₅)₃} upon addition of PMe₃.¹⁸ The resistance of the putative CpMo(η^4 -C₄H₆)Me{MeB(C₆F₅)₃} product against displacement of the borate anion is in agreement with the strength and low polarity of the Mo-CH₃ bonds in the

half sandwich Mo(III) precursors, as indicated theoretically by DFT calculations¹¹ and experimentally by protonation studies.¹⁰

Dissolution of the putative CpMo(η^4 -C₄H₆)Me {MeB(C₆F₅)₃} product in dichloromethane and diffusion of pentane, with participation of adventitious oxygen or water, afforded crystals of an oxygen and chlorine-containing decomposition product which were suitable for a full characterization, including an X-ray analysis. While the chlorine atom is obviously originating from the CH₂Cl₂ solvent, a likely source of the oxygen atom is residual water in the Celite used for the filtration. While oven-dried Celite was used and the product isolation was reproduced, omission of the filtration procedure only resulted in the formation of oily products. The compound corresponds to a salt of the dinuclear [CpMo(η^4 -C₄H₆)(μ -Cl)(μ -Cl₂)(O)MoCp]⁺ cation and the [B(C₆F₅)CH₃]⁻ anion. An ORTEP view of the two ions is shown in Figure 2 and selected metric parameters are given in Table 1.

<Figure 2 and Table 1>

Cation and anion are well separated from each other in the unit cell. The minimum contact is 2.368 Å Å between atoms H4 and H60b. Very weak interactions are observed between two symmetry-related cations (O1...H2, 2.478 Å; O1...H2-C2, 146.1°), which are therefore pairwise arranged in the unit cell as shown in Figure 3. The anion shows a regular tetrahedral coordination around the boron atom. The B-CH₃ distance of 1.658(6) Å is quite typical for this anion.^{33, 34} The cation is composed of the two fragments CpMo(η^4 -C₄H₆) and CpMo(O) bridged by a chlorine atom and a CH₂ group. The hydrogen atoms of the bridging methylene group were not directly located by the X-ray analysis, but their presence is unambiguously shown by the

NMR characterization (*vide infra*). The central $Mo_2(\mu-Cl)(\mu-CH_2)$ moiety is planar and the two Mo fragments are oriented in such a way as to place the two Cp rings on the opposite sides of the bridging plane (*anti*). The two metal atoms are sufficiently close to each other [2.8801(5) Å] to suggest the presence of a direct metal-metal interaction. The acute Mo-X-Mo angles at the bridging Cl and C(2A) atoms [72.00(4) and 86.62(3)°, respectively] reinforce this proposition.

<Figure 3>

Relative to the previously reported structure of $CpMo(\eta^4-C_4H_4Me_2-2,3)(CH_3)_2$, the Mo(1)-CNT(1) distance (Mo-Cp centroid) is slightly shorter [1.985(4) vs. 2.019(5) Å], while the Mo(1)-CNT(3) distance (Mo-diene centroid) is slightly longer [1.964(4) vs. 1.925(3) Å], and the Mo(1)-C(2A) distance is considerably shorter relative to the two Mo-CH₃ distances [2.090(10) Å vs. an average of 2.225(3) Å]. A shortening of all distances to the (formally) anionic ligands and a lengthening of those to the neutral ones are expected when an oxidation state increase occurs. However, we should not assign, from this observation alone, a formal oxidation state IV to the Mo(1) center (and consequently an oxidation state IV also to the Mo(2) center). The alternative assignment as a III,V compound would also be consistent with the observed trends if we suppose that the presence of the adjacent Mo(V) center increases, via the bridging ligands and the direct metal-metal bond, the effective positive charge on the Mo(III) center. The Mo-X (bridging atom) distances to Mo(1) and Mo(2), however, are rather similar [2.497(2) vs. 2.401(3) Å for Cl, 2.090(10) vs. 2.108(15) Å for C(2A)], in better agreement with a IV, IV formulation. While the two Mo-Cl distances are significantly different with that to Mo(2) being shorter, the two Mo-C(2A) distances are identical by the 3σ criterion. The Mo(2)-CNT(2) distance is longer than the Mo(1)-CNT(1) distance. Finally, the Mo(2)-O distance [1.673(3) Å] compares better with those of terminal Mo=O bonds in other cyclopentadienyl complexes of Mo(V), such as 1.684(4) Å for Cp*MoCl₂(O),³⁵ 1.668(7) Å for [CpMoOCl]₂(μ -O),³⁶ or 1.676(5) Å for [Cp*MoOCl]₂(μ -O),³⁷ than with analogous distances in cyclopentadienyl complexes of Mo(IV), such as 1.721(2) Å for (C₅H₄Me)₂Mo(O).³⁸

As mentioned above, the structure of this decomposition product is indirect evidence that the borane reagent has attacked the methyl group rather than the diene ligand in the dimethylmolybdenum(III) precursor, because one methyl group is found to bind the boron atom in the anion, while the putative residual $[CpMo(\eta^4-C_4H_6)(CH_3)]^+$ becomes incorporated, after deprotonation, into the cationic unit, the methyl ligand becoming the bridging methylene group. The $[CpMo(O)CI]^+$ fragment that completes the structure of the cation must derive from the decomposition of a second molecule of $[CpMo(\eta^4-C_4H_6)(CH_3)][CH_3B(C_6F_5)_3]$ (see Scheme 1). The high isolated yield (90%) excludes the fact that this complex is a by-product. The formation of the dinuclear cation can be ascribed to the instability of the putative 15-electron intermediate deriving from the methyl abstraction by $B(C_6F_5)_3$. An analogous instability has also been observed for related group 5 complexes, *e.g.* $[Cp^*Ta(\eta^4-C_4H_6)Me]^+$ and $[Tp'Nb(\eta^2 PhC=CMe)Me]^+$ (Tp' = hydridotris(pyrazolyl)borate),^{7, 8, 39} which, like our proposed Mo(III) intermediate, could not be isolated.

<Scheme 1>

Characterization of the title compound is completed by ¹H, ¹¹B and ¹⁹F NMR spectroscopy and by mass spectrometry. The ¹H NMR spectrum (CDCl₃) shows the presence of two different isomers in solution (in a 70:30 ratio by integration of the cyclopentadienyl signals). These two isomers are interpreted as having a different relative position of the CpMo(=O) and CpMo(η^4 -C4H₆) moieties across the central Mo₂(μ -Cl)(μ -CH₂) moiety, *i.e. syn* or *anti* (see Scheme 2) A similar behavior has been observed for complexes [CpMo(μ -X)(η^4 -C4H₆)]₂ (X=Cl, Br).²⁷ All protons show distinct signals for each isomer, including the six hydrogen atoms of the diene ligand. The two inequivalent methylene protons give rise to two distinct doublets (J_{HH} = 10.5 Hz) for the major isomer, while a singlet is observed for the minor one. The ¹¹B chemical shift and the difference of 2.5 ppm between the *meta*-F and *para*-F resonances in the ¹⁹F NMR are in good agreement with the formation of the borate ion and with a significant charge separation in the salt. ^{16, 40} The mass spectrum shows peaks in agreement with the given formulas for the cation and anion in the positive and negative chemical ionization modes, respectively.

<Scheme 2>

Conclusion

The borane reagent $B(C_6F_5)_3$ attacks the electron-rich dimethyl complexes $Cp*Mo(\eta^4-diene)Me_2$ at the methyl position, rather than at the diene ligand, to form relatively stable adducts that resist displacement of the $[CH_3B(C_6F_5)_3]^-$ anion by Lewis bases. The position of borane attack is suggested by the simulation of the EPR data and indirectly by the structural

characterization of the title product. The observed reactivity pattern is consistent with previously available Mulliken charges from DFT computations.

The present study has confirmed the great resistance of the Mo(III)-CH₃ bond toward heterolytic cleavage, previously shown by the fact that the bond can only be cleaved by acids stronger than H_3O^+ .¹⁰ Yet, compounds of type Cp*Mo(η^4 -diene)Me₂ has been shown capable to polymerize ethylene in combination with MAO, with a relatively rapid turnover for the active site. The activation step, however, is difficult.¹¹ The results shown here will guide us toward the development of modified systems with an improved catalytic activity.

Experimental section

General procedures. All reactions were carried out in a Jacomex glove box or by the use of standard Schlenk techniques under an argon atmosphere. The solvents were dried by conventional methods (THF, Et₂O, toluene and pentane from sodium benzophenone ketyl and CH₂Cl₂ from P₄O₁₀) and distilled under argon prior to use. EPR measurements were carried out at the X-band microwave frequency on a Bruker ESP300 spectrometer. The spectrometer frequency was calibrated with DPPH (g= 2.0037). NMR spectra were recorded using a Bruker AC-200 or DRX-500. ¹H NMR spectra were referenced to the residual solvent protons of the deuterated solvent. ¹⁹F (282.2 MHz) and ¹¹B (96.2 MHz) NMR spectra were referenced externally to CFCl₃ and BF₃:Et₂O, respectively. Cyclic voltammograms were recorded with an EG&G 362 potentiostat connected to a Macintosh computer through MacLab hardware/software. Electrospray mass spectra were recorded using a Micromass Quattro LC instrument. Isotope patterns were recorded for cation and anion and compared with theoretical patterns using the MassLynx 3.5 program. In both cases there was good agreement between the experimental and calculated isotopic mass distributions. The elemental analyses were carried out by the analytical service of the Laboratoire de Synthèse et d'Electrosynthèse Organométallique with a Fisons EA 1108 apparatus. Compound CpMo(diene)Cl₂ were prepared as described in the literature.¹⁰ $B(C_6F_5)_3$ was prepared as described⁴¹ and was sublimed in vacuo before use.

Reaction of CpMo(η^4 -C4H6)(CH3)2 with B(C6F5)3. A solution of CpMo(η^4 -C4H6)(CH3)2 (84 mg, 0,34 mmol) in toluene (10 mL) was added to a solution of B(C6F5)3 (175 mg, 0,34 mmol) in the same solvent (5 mL) at -20°C, resulting in an immediate color change from green to orange. The solution was stirred for 15 min at room temperature. An aliquot of the solution was examined by EPR, indicating the formation of a new compound (see Results and Figure 1): g = 1.995 (multiplet), $a_{M0} = 38.91$ G, $a_H = 6.79$ G (3H), $a_H = 4.42$ (3H), $a_H = 4.55$ G (2H), $a_H = 4.55$ G (2H). Cooling the filtered toluene solution to -20°C or -80°C gave either no precipitation or an oil, depending on concentration. The same occurred upon diffusion of pentane into a concentrated (and unfiltered) dichloromethane solution.

Reaction of CpMo(η^4 -C₄H₄Me₂-2,3)(CH₃)₂ with B(C₆F₅)₃. A procedure identical to that described in the previous section gave rise to the same observations. EPR (toluene): g = 1.995 (multiplet), $a_{Mo} = 34.5$ G. The values of the other hyperfine constants was not optimized by digital fitting. The appearance of this spectrum, however, was essentially identical with that of the butadiene analogue (*vide supra*), except for a slightly worse resolution.

Reaction of CpMo(η^4 -C4H₆)(CH₃)₂ with B(C₆F₅)₃ in the presence of PMe₃. Following the same procedure as described above, PMe₃ (1 mL, 1 M in toluene, 1 mmol) was added via a syringe at room temperature. No spectral modification were observed by ESR.

Formation of $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp][B(C_6F_5)CH_3]$. After carrying out a reaction between $CpMo(\eta^4-C_4H_6)(CH_3)_2$ and $B(C_6F_5)_3$ as indicated above, the toluene

solvent was completely evaporated. The residue was then extracted with 3 mL of CH_2Cl_2 , followed by filtration through Celite (0.5x1cm). A slow diffusion of pentane in this solution at room temperature afforded red crystals. Yield = 148 mg; 90 %. Anal. Calcd. for C₃₄H₁₉BClF₁₅Mo₂O: C, 42.16; H, 2.18. Found : C, 41.91; H, 2.09. Positive Electrospray-MS (Cone: 25V): m/z = 441 ([CpMo(η^4 -C₄H₆)(μ -Cl)(μ -CH₂)(O)MoCp]⁺).Negative Electrospray-MS (Cone: 25V) : m/z = 526 ([CH₃B(C₆F₅)₃]⁻). ¹H NMR (CDCl₃, 20°C, δ): [CH₃B(C₆F₅)₃]⁻ anion: 0.57 (br.). $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp]^+$ (major isomer): 6.1 (s, 5H, C₅H₅Mo(=O)); 5.6 (s, 5H, $C_5H_5Mo(\eta^4-C_4H_6)$); 8.44 (d, 1H, CH_2 , ${}^2J_{H-H} = 10.5$ Hz); 7.79 (d, 1H d, CH_2 , ${}^2J_{H-H} =$ 10.5 Hz); 4.80 (m, 1H, C₄H₆); 3.87 (m, 1H, C₄H₆); 3.39 (m, 1H, C₄H₆); 2.74 (m, 1H, C₄H₆); 2.03 (m, 1H, C₄ H_6); 1.67 (m, 1H, C₄ H_6). [CpMo(η^4 -C₄ H_6)(μ -Cl)(μ -Cl)(μ -Cl₂)(O)MoCp]⁺ (minor isomer): 6.05 (s, 5H, C₅H₅Mo(=O)); 5.32 (s, 5H, C₅H₅Mo(η^4 -C₄H₆)); 8.10 (s, 2H, CH₂); 4.54 (m, 1H, $C_{4}H_{6}$; 4.25 (m, 1H, $C_{4}H_{6}$); 3.33 (m, 1H, $C_{4}H_{6}$); 2.83 (m, 1H, $C_{4}H_{6}$); 2.33 (m, 1H, $C_{4}H_{6}$); 1.88 (m, 1H, C₄*H*₆). ¹¹B NMR (CDCl₃, 20°C, δ): -15,3 ppm. ¹⁹F NMR (CDCl₃, 20°C, δ): -132.8 (d, *o*-F, ${}^{3}J_{F-F} = 19.2$ Hz), -164.6 (t, *p*-F, ${}^{3}J_{F-F} = 20.8$ Hz), -167.2 (td, *m*-F, ${}^{3}J_{F-F} = 20.8$ Hz, ${}^{3}J_{F-F} = 19.2$ Hz).

X-ray crystallography for compound $[CpMo(\eta^4-C4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp]$ -[B(C₆F₅)₃CH₃]. The crystals are air stable and were mounted on the tip of a glass fiber. X-ray diffraction experiments were carried out on a Bruker SMART CCD diffractometer using Mo-K α radiation (λ 0.71073 A). The data were collected with a frame width of 0.3° in Ω and a counting time of 20 s per frame at a crystal to detector distance of 4 cm. The software SAINT⁴² was used for integration of intensity reflections and scaling and SADABS⁴³ for absorption correction. Final cell parameters were obtained by global refinement of reflections obtained from integration of all the frames data. The crystal parameters and basic information relating data collection and structure refinement for compound $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp][B(C_6F_5)_3CH_3]$ are summarized in Table 2. The structure was solved by direct methods and refined by the fullmatrix method based on F² using the SHELXTL software package.⁴⁴ The non-hydrogen atoms of structure was refined anisotropically; the positions of all hydrogen atoms were generated geometrically, assigned isotropic thermal parameters and allowed to ride on their respective parent carbon atoms. The chlorine and methylene bridge are disordered and were refined with a occupations factors of 0.62 for Cl1 and C2A and 0.38 for Cl2 and C1A.

<Table 2>

Acknowledgement. We are grateful to the CNRS (Programme Catalyse pour l'Industrie et l'Environnement) and COST D17 (Working group on "Environmentally friendly catalysts for olefin polymerization") for support of this work. E.L.G thanks the MENRT for a doctoral fellowship. Thanks are also extended to the Servei Central d'Instrumentació Científica (CSIC) of the university Jaume I for providing us with mass spectrometry and x-ray facilities.

Supporting information

Crystallographic data for the structural analysis for compound $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp][B(C_6F_5)_3CH_3]$ have been deposited with the Cambridge Crystallographic Data Centre 12 Union Road, Cambridge, CB2 1EZ UK, with the deposition number CCDC No. 164163 (11 pages).

References

- 1 R. F. Jordan, Adv. Organomet. Chem., 1991, **32**, 325-387.
- 2 P. C. Mohring and N. J. Coville, J. Organometal. Chem., 1994, 479, 1-29.
- H. H. Brintzinger, D. Fischer, R. Mülhaupt, B. Rieger and R. M. Waymouth, *Angew. Chem., Int. Ed. Engl.*, 1995, 34, 1143-1170.
- 4 M. Bochmann, J. Chem. Soc., Dalton Trans., 1996, 255-270.
- 5 W. Kaminsky, J. Chem. Soc., Dalton Trans., 1998, 1413-1420.
- 6 K. Mashima, S. Fujikawa, H. Urata, E. Tanaka and A. Nakamura, J. Chem. Soc., Chem. Commun., 1994, 1623-1624.
- K. Mashima, S. Fujikawa, Y. Tanaka, H. Urata, T. Oshiki, E. Tanaka and A. Nakamura, *Organometallics*, 1995, 14, 2633-2640.
- J. Jaffart, C. Nayral, R. Choukroun, R. Mathieu and M. Etienne, *Eur. J. Inorg. Chem.*, 1998, 425-428.
- 9 E. Le Grognec, R. Poli and L.-S. Wang, *Inorg. Chem. Commun.*, 1999, 2, 95-96.
- 10 E. Le Grognec, R. Poli and P. Richard, *Organometallics*, 2000, **19**, 3842-3853.
- 11 E. Le Grognec and R. Poli, *Chem. Eur. J.*, in press.
- 12 E. Y.-X. Chen and T. J. Marks, *Chem. Rev.*, 2000, **100**, 1391-1434.
- 13 X. Yang, C. L. Stern and T. J. Marks, J. Am. Chem. Soc., 1994, 116, 10015-10031.
- M. Bochmann, S. J. Lancaster, M. B. Hursthouse and K. M. A. Malik, *Organometallics*, 1994, 13, 2235-2243.
- G. C. Bazan, W. D. Cotter, Z. J. A. Komon, R. A. Lee and R. J. Lachicotte, *J. Am. Chem. Soc.*, 2000, **122**, 1371-1380.

- A. D. Horton, J. de With, A. J. van der Linden and H. van der Weg, *Organometallics*, 1996, 15, 2672-2674.
- F. Amor, A. Butt, K. E. du Plooy, T. P. Spaniol and J. Okuda, *Organometallics*, 1998, 17, 5836-5849.
- 18 J. S. Rogers, X. Bu and G. C. Bazan, J. Am. Chem. Soc., 2000, 730-731.
- B. Temme, G. Erker, J. Karl, H. Luftmann, R. Frohlich and S. Kotila, *Angew. Chem., Int. Ed. Eng.*, 1995, 34, 1755-1757.
- 20 B. Temme, J. Karl and G. Erker, *Chem. Eur. J.*, 1996, **2**, 919-924.
- J. Karl, M. Dahlmann, G. Erker and K. Bergander, J. Am. Chem. Soc., 1998, 120, 5643-5652.
- G. Jimenéz Pindado, S. J. Lancaster, M. Thornton-Pett and M. Bochmann, J. Am. Chem.
 Soc., 1998, 120, 6816-6817.
- A. H. Cowley, G. S. Hair, B. G. McBurnett and R. A. Jones, *Chem. Commun.*, 1999, 437-438.
- M. Dahlmann, G. Erker, R. Fröhlich and O. Meyer, *Organometallics*, 2000, 19, 2956-2967.
- J. L. Davidson, K. Davidson and W. E. Lindsell, J. Chem. Soc., Chem. Commun., 1983, 452-453.
- 26 J. L. Davidson, K. Davidson, W. E. Lindsell, N. W. Murrall and A. J. Welch, J. Chem. Soc., Dalton Trans., 1986, 1677-1688.
- 27 L.-S. Wang, J. C. Fettinger and R. Poli, J. Am. Chem. Soc., 1997, 119, 4453-4464.
- 28 R. Poli and L.-S. Wang, *Polyhedron*, 1998, 17, 3689-3700.
- 29 E. Le Grognec, J. Claverie and R. Poli, J. Am. Chem. Soc., in press.
- 30 X. Yang, C. L. Stern and T. J. Marks, J. Am. Chem. Soc., 1991, 113, 3623-3625.

- 31 C. L. Beswick and T. J. Marks, *Organometallics*, 1999, **18**, 2410-2412.
- 32 A. J. Sillanpää and K. E. Laasonen, J. Am. Chem. Soc., 2001, 20, 1334-1344.
- 33 H. C. Strauch, G. Erker, R. Frohlich and M. Nissinen, Organometallics, 1998, 17, 5746.
- H. C. Strauch, G. Erker, R. Fröhlich and M. Nissinen, *Eur. J. Inorg. Chem.*, 1999, 1453-1459.
- J. M. B. Blanchard, Q. Chen, Y.-D. Chang and M. B. Sponsler, *Inorg. Chim. Acta*, 1994,
 221, 147-150.
- 36 F. Bottomley, E. C. Ferris and P. S. White, *Organometallics*, 1990, 9, 1166-1171.
- 37 K. Umakoshi and K. Isobe, J. Organometal. Chem., 1990, **395**, 47-53.
- 38 N. D. Silavwe, M. Y. Chiang and D. R. Tyler, *Inorg. Chem.*, 1985, 24, 4219-4221.
- 39 K. Mashima, S. Fujikawa and A. Nakamura, J. Am. Chem. Soc., 1993, 115, 10990-10991.
- 40 A. D. Horton and J. de With, *Organometallics*, 1997, **16**, 5424-5436.
- 41 A. G. Massey and A. J. Park, J. Organomet. Chem., 1964, 2, 245-250.
- 42 Bruker Analytical X-ray System, 'SAINT, version 5.0', Madison WI, 1998.
- G. M. Sheldrick, 'SADABS empirical absorption program', University of Göttingen, 1996.
- 44 G. M. Sheldrick, 'SHELXTL, version 5.1', Bruker AXS, Inc., Madison WI, 1997.

Distances (Å)		Angles (°)	
(a) Cation			
Mo(1)-CNT(1)	1.985(4)	CNT(1)-Mo(1)-CNT(3)	130.99(5)
Mo(1)-CNT(3)	1.964(4)	CNT(1)-Mo(1)-C1	108.22(4)
Mo(1)-Cl	2.497(2)	CNT(1)-Mo(1)-C(2A)	111.80(4)
Mo(1)-C(2A)	2.090(10)	CNT(2)-Mo(2)-Cl	113.25(5)
Mo(1)-C(11)	2.288(5)	CNT(2)-Mo(2)-C(2A)	111.41(4)
Mo(1)-C(12)	2.353(4)	CNT(2)-Mo(2)-O	120.66(7)
Mo(1)-C(13)	2.349(4)	CNT(3)-Mo(1)-C(2A)	99.66(7)
Mo(1)-C(14)	2.298(4)	CNT(3)-Mo(1)-Cl	102.53(5)
C(11)-C(12)	1.409(7)	O-Mo(2)-Cl	102.21(7)
C(12)-C(13)	1.393(7)	O-Mo(2)-C(2A)	105.68(5)
C(13)-C(14)	1.393(6)	Mo(1)-Cl-Mo(2)	72.00(4)
Mo(2)-CNT(2)	2.067(4)	Mo(1)-C(2A)-Mo(2)	86.62(3)
Mo(2)-O	1.673(3)	Mo(1)-Mo(2)-O	107.18(7)
Mo(2)-Cl	2.401(3)	Mo(1)-Mo(2)-CNT(2)	131.99(5)
Mo(2)-C(2A)	2.108(15)	Mo(2)-Mo(1)-CNT(1)	116.74(5)
Mo(1)-Mo(2)	2.8801(5)	Mo(2)-Mo(1)-CNT(3)	112.23(4)
(b) Anion			
B(1)-C(20)	1.659(6)	C(20)-B(1)-C(30)	109.43(3)
B(1)-C(30)	1.662(6)	C(20)-B(1)-C(40)	110.26(3)
B(1)-C(40)	1.656(6)	C(30)-B(1)-C(40)	110.91(3)
B(1)-C(60)	1.658(6)	C(60)-B(1)-C(20)	109.77(3)
		C(60)-B(1)-C(30)	109.01(3)
		C(60)-B(1)-C(40)	107.43(3)

Table 1. Selected Bond Distances (Å) and Angles (°) for compound $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-Cl)(\mu-CH_2)(O)MoCp][B(C_6F_5)CH_3]$.^a

^a CNT(n) (n = 1, 2) are the center of gravity of the cyclopentadienyl ring [n = 1, atoms C(1)-C(5); n = 2, atoms C(6)-C(10)]; CNT(3) is the center of gravity of the butadiene ligand [atoms C(11)-C(14)]

Table 2. Crystal data and structure refinement for $[CpMo(\eta^4-C_4H_6)(\mu-Cl)(\mu-CH_2)(O)MoCp]-$ [B(C₆F₅)CH₃]

Empirical formula	CarHigBClErcMonO	
Empirical formula	$C_{34}\Pi_{19}DC_{1}\Gamma_{15}W_{102}O$	
Formula weight	908.03	
Temperature	173(2) K	
Crystal system	Monoclinic	
Space group	P21/c	
a (Å).	14.2045(12)	
b (Å)	18.5778(14)	
c (Å)	13.1036(10)	
β (°).	107.380(2)	
Volume (Å ³)	3300.0(5)	
Z	4	
Absorption coefficient (mm ⁻¹)	0.957	
Crystal size (mm ³)	0.113 x 0.044 x 0.016	
Theta range for data collection (°).	1.50 to 30.61	
Index ranges	18<=h<=19, -26<=k<=16, -18<=l<=18	
Reflections collected	26693	
Independent reflections	9775 [R(int) = 0.0792]	
Observed independent reflections [I>2sigma(I)]	4995	
Completeness to theta = 30.61°	96.1 %	
Goodness-of-fit on F ²	0.914	
Final R indices [I>2sigma(I)]	$R1^{a} = 0.0485, wR2^{b} = 0.0870$	
R indices (all data)	R1 = 0.1340, wR2 = 0.1066	
Largest diff. peak and hole (e.Å ⁻³)	0.718 and -0.552	

^a $R1 = \sum ||F_0| - |F_c|| / \sum F_0$ ^b $wR2 = \left[\sum \left[w (F_0^2 - F_c^2)^2\right] / \sum \left[w (F_0^2)^2\right]\right]^{1/2}$

Captions for Figures

- Figure 1. EPR monitoring of the reaction between $CpMo(\eta^4-C_4H_6)Me_2$ and $B(C_6F_5)_3$ in toluene at -20°C. (a) Before the borane addition. (b) After the borane addition.
- Figure 2. An ORTEP view of cation (above) and anion (below) of compound [CpMo(η⁴-C₄H₆)(μ-Cl)(μ-CH₂)(O)MoCp][B(C₆F₅)CH₃] with thermal ellipsoids drawn at the 30% probability level
- Figure 3. A view of the pairwise arrangement of the dinuclear molybdenum cations.

Figure 1.







Figure 3



Scheme 1



Scheme 2





