

# **Multi-Normex Distributions for the Sum of Random Vectors. Rates of Convergence**

Marie Kratz, Evgeny Prokopenko

## **To cite this version:**

Marie Kratz, Evgeny Prokopenko. Multi-Normex Distributions for the Sum of Random Vectors. Rates of Convergence.  $2021.$  hal- $03294714$ 

# **HAL Id: hal-03294714 <https://hal.science/hal-03294714>**

Preprint submitted on 21 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enlighten. Lead. Change.



# Multi-Normex Distributions for the Sum of Random Vectors. Rates of Convergence

Marie Kratz and Evgeny Prokopenko

**ESSEC RESEARCH CENTER** 

WORKING PAPER 2102

JULY 19, 2021



# Multi-Normex Distributions for the Sum of Random Vectors. Rates of Convergence

Marie Kratz† and Evgeny Prokopenko† ‡

† ESSEC Business School Paris, CREAR, France ‡ Sobolev Institute of Mathematics, Novosibirsk, Russia

July 19, 2021

#### Abstract

We build a sharp approximation of the whole distribution of the sum of iid heavy-tailed random vectors, combining mean and extreme behaviors. It extends the so-called 'normex' approach from a univariate to a multivariate framework. We propose two possible multinormex distributions, named d-Normex and MRV-Normex. Both rely on the Gaussian distribution for describing the mean behavior, via the CLT, while the difference between the two versions comes from using the exact distribution or the EV theorem for the maximum. The main theorems provide the rate of convergence for each version of the multi-normex distributions towards the distribution of the sum, assuming second order regular variation property for the norm of the parent random vector when considering the MRV-normex case. Numerical illustrations and comparisons are proposed with various dependence structures on the parent random vector, using QQ-plots based on geometrical quantiles.

2 010 AMS classification: 60E05; 60F05; 60G70; 60F15; 62G30; 41A25

Keywords: aggregation; central limit theorem; dependence; extreme value theorem; geometrical quantiles; multivariate regular variation; (multivariate) Pareto distribution; ordered statistics; QQ-plots; rate of convergence; second order regular variation; sum of random vectors

# Contents





# 1 Introduction

Motivation. Looking for the most accurate possible evaluation of the distribution of the sum of random variables, or vectors, or processes, with unknown distributions, has always been a classical problem in the probabilistic and statistical literature, with various answers depending on the given framework and on the specific application in view. On one hand, (uni- or multivariate) Central Limit Theorems (CLT) or Functional ones prove, under finite variance for the sum components or/and additional conditions, the asymptotic Gaussian behavior of the sum with some rate of convergence, focusing on the 'body' of the distribution. When considering heavy-tailed marginal distributions, Generalized CLT with the convergence to stable distributions, handle the case of infinite variance (see e.g. Samorodnitsky and Taqqu (1994) and Petrov (1995), and references therein), while, in the case of finite variance, an alternative way is to consider trimmed sums, removing extremes from the sample, to improve the rate of convergence; see e.g. Mori (1984) and Hahn (1991) and references therein.

When interested in tail distributions, CLTs may give poor results, especially when considering heavy tails. That is why different approaches have been developed, among which large deviation theorems (see e.g. Petrov (1975) and Borovkov (2020) for light tails and Mikosch and Nagaev (1998), Foss et al. (2013), and Lehtomaa (2017) for heavy tails, and references therein), extreme value theorems (EVT) focusing on the tail only (see e.g. Embrechts et al. (1997), de Haan and Ferreira (2006), and Resnick (2007)), and hybrid distributions combining (asymptotic) distributions for both the main and extreme behaviors when considering independent random variables (see e.g. Csörgö et al. (1988), Zaliapin et al. (2005), Kratz (2014), and Müller (2019); we use the name given in Kratz (2014) for this type of hybrid distribution/method/approach, namely Normex distribution/method/approach. Recall briefly the idea of Normex (for 'Norm(al)-Ex(tremes)') method. It consists of rewriting the sum of random variables as the sum of their ordered statistics, and splitting it into two main parts, a trimmed sum removing the extremes, and the extremes. Using that the trimmed sum of the first  $n - k - 1$  ordered statistics is conditionnally independent of the k largest order statistics, given the  $(n - k)$ -th order statistics, we can express the distribution of the sum, integrating w.r.t. to the  $(n-k)$ -th order statistics and using a CLT for the conditional trimmed sum, and an EVT one for the k largest order statistics. Note that a benefit of Normex approach is that it does not require any condition on the existence of moments, as the CLT applies on truncated random variables.

The following example (as developed in Kratz (2014)), simulating identically distributed and independent (iid) Pareto( $\alpha$ ) (such that  $\overline{F}(x) = x^{-\alpha}$  for  $x > 1$ ) random variables (rv), with  $\alpha = 2.3$  (finite variance, but no third moment), illustrates perfectly the adding value of combining main and extreme behaviors.

Indeed, on the QQ-plots given in Figure 1, we can compare the fit of the distribution of the empirical sample with the following three distributions: the Gaussian one (CLT approach), the Fréchet one (EVT approach), the hybrid one *(i.e.* Normex, combining the CLT and the distribution of the maximum). For the hybrid Normex distribution, we may consider either the exact distribution of the maximum, or its asymptotic approximation, the Fréchet distribution.



Figure 1: QQ-plots for the empirical distribution (sample size  $=10^7)$  of the sum of 52 iid Pareto $(\alpha=2.3)$ rv and three different approximations of the sum distribution using: CLT, EVT and Normex, respectively. The red circles and numbers denote the extreme quantiles and their levels.

Since both provide the same plot, we display it in the third plot on the right. Note that we choose a rather small number of components in the sum,  $n = 52$ , also to illustrate the speed of convergence when using asymptotic theorems. We observe that the CLT approach does not provide a sharp evaluation, even in the body of the distribution, due to this choice of  $n$ , which cannot compensate yet the fact that the distribution of the rv is asymmetric and skewed; increasing  $n$  will of course improve the fit in the body of the distribution. Given the fact that the Pareto distribution belongs to the Fréchet maximum domain of attraction, using the Fréchet distribution for the distribution of the sum of Pareto rv's gives a very sharp approximation in the tail (from the 93% quantile), but not for the average behavior, as expected. Finally, a perfect match between empirical quantiles and Normex ones is observed for the whole distribution in the right plot, even for a small number of summands.

GOAL OF THE STUDY. It is natural to extend the normex approach to a multivariate framework. With this goal of proposing a multi-normex method and distribution, we consider iid random vectors  $X_1, \ldots, X_n$ , with parent random vector X having a heavy-tailed d-dimensional distribution  $F_{\mathbf{X}}$  and density  $f_{\mathbf{X}}$  (when existing). Note that there are different ways to define multivariate extremes (see e.g. chap.8 in Beirlant et al. (2004)). The chosen way in this paper is w.r.t. the norm  $\|\cdot\|$  in  $\mathbb{R}^d$ , meaning that the ordered (w.r.t. the norm) vector of  $(\mathbf{X}_1, \ldots, \mathbf{X}_n)$ , denoted by  $(\mathbf{X}_{(1)}, \ldots, \mathbf{X}_{(n)})$ , satisfies

$$
\|\mathbf{X}_{(1)}\| \le \|\mathbf{X}_{(2)}\| \le \ldots \le \|\mathbf{X}_{(n)}\|.\tag{1.1}
$$

We propose two versions of multi-normex. The first one, named  $d$ -Normex, is a natural extension to any dimension d of the univariate  $(d = 1)$  normex method as developed in Kratz (2014): We approximate the distribution of the trimmed sum via the CLT and consider the distribution of the maximum  $\mathbf{X}_{(n)}$ . This latter distribution is approximated via the Extreme Value (EV) theorem in the second multi-normex version, named MRV-Normex.

Aiming at proving the benefit of using a multi-normex distribution for a better fit of the whole (unknown) distribution F, assuming F heavy-tailed in the sense of  $||\mathbf{X}|| \in \mathcal{RV}_{-\alpha}$ , *i.e.* regularly varying rv with  $\alpha > 0$  (which definition is recalled in Appendix A.1), we focus analytically on the case  $\alpha \in (2,3]$  (when ||X|| has a finite second moment, but no third moment), to compare the rates of convergence when using the CLT and the multi-normex approach, respectively. Note our focus on heavy tailed distributions (i.e. distributions belonging to the max domain of attraction of Fréchet), where the impact of using Normex distribution will be much stronger than in the light tail case (because of the one big jump principle), in particular for risk analysis and management. We prove that the normex approach leads, as expected, to a better speed of convergence for evaluating the distribution of the sum than the CLT does, for such type of heavy-tailed distributions. When varying the fatness of the tail measured by  $\alpha > 0$ , we draw this comparison numerically, using geometrical multivariate quantiles (see e.g. Chaudhuri (1996) and Dhar et al. (2014) or Appendix A.5).

STRUCTURE OF THE PAPER. In Section 2, besides general notations, we recall the normex approach and the generalized Berry-Esseen inequality. Then we give two specific results on conditional distributions of order statistics, which will be needed for the construction of multinormex distributions. The two next sections develop the two multi-normex versions, d-Normex in Section 3 and MRV-Normex in Section 4. These sections have the same structure: we first define the multi-normex distribution, then we study analytically its rate of convergence, before ending with some examples. In Section 5, we consider those examples to study numerically the two versions of multi-normex distribution, comparing them with the empirical distribution of the sum (obtained via simulation) as well as, if relevant, with the Gaussian approximation when applying the CLT. Geometrical multivariate quantiles are computed to this aim and represented on QQ-plots. Section 6 concludes. The appendix is divided in 5 parts. Section A.1 recalls the various definitions of regular variation and its related concepts, as well as two properties of second order regular variation that are needed for the proofs of the analytical results. The proofs of all analytical results are developed in Sections A.2 to A.4. An overview of geometrical multivariate quantiles is given in Section A.5.

## 2 Framework

### 2.1 Context and notations

NORMEX IDEA - The Normex method clearly adapts to a multivariate framework. Using this approach, we split the maximum from the rest of the sum:

$$
\mathbf{S}_n = \sum_{k=1}^{n-1} \mathbf{X}_{(k)} + \mathbf{X}_{(n)},
$$
\n(2.2)

taking into account the 'principle of one big jump', namely that the asymptotic tail behavior of the sum of heavy-tailed random vectors is led by that of the maximum. Indeed, this principle extends to dimension  $d > 1$  by borrowing the multivariate subexponential distribution definition of Samorodnitsky and Sun (2016) (see Section 4 therein):

$$
\mathbb{P}\left(\,\mathbf{S}_n>t\,\mathbf{A}\,\right)\;\underset{t\rightarrow\infty}{\sim}\;\mathbb{P}\left(\,\mathbf{X}_{(n)}>t\,\mathbf{A}\,\right),\,
$$

where  $\mathbf{A} \subset \mathbb{R}^d$  is open, increasing, such that  $\mathbf{A}^c$  convex and  $\mathbf{0} \notin \bar{\mathbf{A}}$ . Note also earlier works on that notion by Cline and Resnick (1992) and Omey (2006).

Combining the CLT for the trimmed sum given the maximum, and the distribution of the maximum or its asymptotic distribution, leads to a multivariate version of Normex distribution. We name this multi-normex distribution as d-Normex, when using the distribution of the maximum, and as MRV-Normex when considering its asymptotic distribution (when rescaled).

General notations - Before defining explicitly both versions of multi-normex distribution, let us introduce some general notations.

Let  $f_{(i)}$  denote the d-dimensional density, when existing, of the ordered vectors  $\mathbf{X}_{(i)}$ , for  $i =$  $1, \cdots, n$ .

The cumulative distribution function (cdf) of the norm  $\|\mathbf{X}\|$  is denoted as  $F_{\|\mathbf{X}\|}(\cdot)$  and its probability density function (pdf), when existing, as  $f_{\parallel {\bf X} \parallel}(\cdot)$ .

As we will work with truncated multidimensional distributions or vectors, let us introduce the following notions.

For any  $y > 0$ , we define the truncated (via the norm) multidimensional distribution  $F_{\mathbf{X} \mid \|\mathbf{X}\|}(\cdot | y)$ of **X** on  $\mathbb{R}^d$  as

$$
F_{\mathbf{X} \mid \|\mathbf{X}\|}(\mathbf{B} \mid y) := \mathbb{P}\left(\mathbf{X} \in \mathbf{B} \mid \|\mathbf{X}\| \leqslant y\right) \tag{2.3}
$$

for any event **B** of the Borel sigma-field  $\mathcal{B}(\mathbb{R}^d)$ . We denote by  $f_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot)$  its pdf, when existing:

$$
f_{\mathbf{X} \mid \|\mathbf{X}\|}(\mathbf{x} \mid y) = \frac{f(\mathbf{x}) \, \mathbf{1}_{\left(\|\mathbf{x}\| \leq y\right)}}{F_{\|\mathbf{X}\|}(y)}.\tag{2.4}
$$

Let  $\mathbf{X}_y \in \mathbb{R}^d$  denote the random vector with distribution  $F_{\mathbf{X}_y}$  on the  $(d-1)$ -sphere  $\mathcal{S}_y = \left\{ \mathbf{x} \in \mathbb{R}^d : ||\mathbf{x}|| = y \right\}$  for  $y > 0$ , defined by

$$
F_{\hat{\mathbf{X}}_y}(\mathbf{B}_y) = \mathbb{P}\left(\overset{\circ}{\mathbf{X}}_y \in \mathbf{B}_y\right) := \mathbb{P}\left(\mathbf{X} \in \mathbf{B}_y \,|\, \|\mathbf{X}\| = y\right) \text{ for } \mathbf{B}_y \subseteq \mathcal{S}_y. \tag{2.5}
$$

With this definition, for any  $\mathbf{A} \subseteq \mathbb{R}^d$ , we can write

$$
\mathbb{P}\left(\mathbf{X} \in \mathbf{A}\right) = \mathbb{E}\left[\mathbb{E}[\mathbb{I}_{\mathbf{X} \in \mathbf{A}} \mid \|\mathbf{X}\|]\right] = \int_0^\infty \mathbb{E}[\mathbb{I}_{\mathbf{X} \in \mathbf{A}} \mid \|\mathbf{X}\| = y] f_{\|\mathbf{X}\|}(y) dy,
$$

thus

$$
\mathbb{P}\left(\mathbf{X} \in \mathbf{A}\right) = \int_{\mathbb{R}_{+}} \mathbb{P}\left(\mathbf{X}_{y} \in \mathbf{A}_{y}\right) f_{\|\mathbf{X}\|}(y) \mathrm{d}y, \quad \text{where} \quad \mathbf{A}_{y} := \mathbf{A} \cap \mathcal{S}_{y}.
$$
 (2.6)

GENERALIZED BERRY-ESSEEN INEQUALITY - As we are going to compare the rates of convergence when using, respectively, the CLT and the Normex method, let us recall the rate of convergence in the CLT provided by the Generalized Berry-Esseen inequality (see e.g. Corollary 18.3 of Bhattacharya and Rao (2010)), when assuming a 'moderate' heavy tail.

Proposition 2.1 (Generalized Berry-Esseen inequality).

Let  $X_1, \ldots, X_n$  be i.i.d. centered random vectors with parent random vector X with values in  $\mathbb{R}^d$ , with positive-definite covariance matrix  $\Sigma$ . If

$$
\mathbb{E} \left\| \mathbf{X} \right\|^{\alpha} < \infty, \quad \text{for some } \alpha \in [2, 3],
$$

then

$$
\sup_{\mathbf{B}\in\mathcal{C}}|\mathbb{P}\left(\mathbf{S}_n\in\mathbf{B}\right)-\Phi_{\mathbf{0},\Sigma}(\mathbf{B})|\leqslant c\,\mathbb{E}\left(\left\|\Sigma^{-1/2}\mathbf{X}\right\|^{\alpha}\right)\,n^{-(\alpha-2)/2},\tag{2.7}
$$

where C is the class of all Borel-measurable convex subsets of  $\mathbb{R}^d$ , c is a positive universal constant, and  $\Phi_{0,\Sigma}$  is the cdf of the centered normal multivariate distribution with covariance matrix  $\Sigma$ .

Note that the (non-generalized) Berry-Esseen inequality, which holds for any  $\alpha \geqslant 3$ , corresponds to (2.7) when taking  $\alpha = 3$ .

#### 2.2 Preliminary results on order statistics

First, we have a quite elegant result:

**Lemma 2.1.** The distribution of the first  $n-1$  order statistics  $X_{(1)}, \ldots, X_{(n-1)}$ , conditionally on the event  $\|\mathbf{X}_{(n)}\| = y$ , is the distribution of the n - 1 ordered statistics from the truncated distribution  $F_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot | y)$ :

$$
\mathcal{L}\left(\mathbf{X}_{(1)},\ldots,\mathbf{X}_{(n-1)}\,\middle|\, \|\mathbf{X}_{(n)}\| = y\right) = \mathcal{L}\left(\mathbf{Y}_{(1)},\ldots,\mathbf{Y}_{(n-1)}\right),
$$

where  $Y_1, \ldots, Y_{n-1}$  are i.i.d. random vectors with multidimensional distribution  $F_{\mathbf{X} \mid \mathbf{X} \mathbf{X}}(\cdot | y)$ defined in (2.3).

Similarly, we have the following result.

**Lemma 2.2.** The distribution of the order statistics  $X_{(1)}, \ldots, X_{(n-1)}, X_{(n)}$ , conditionally on the event  $\|\mathbf{X}_{(n)}\| = y$ , is the distribution of the n-1 ordered statistics from the truncated distribution  $F_{\mathbf{X}||\mathbf{X}||}(\cdot | y)$ , and of an independent random vector  $\hat{\mathbf{X}}_y$  defined on the  $(d-1)$ -sphere  $\mathcal{S}_y$ , for  $y > 0$ :

$$
\mathcal{L}\left(\mathbf{X}_{(1)},\ldots,\mathbf{X}_{(n-1)},\mathbf{X}_{(n)}\,\big|\,\|\mathbf{X}_{(n)}\|=y\right)=\mathcal{L}\left(\mathbf{Y}_{(1)},\ldots,\mathbf{Y}_{(n-1)}\right)\times\mathcal{L}\left(\overset{\circ}{\mathbf{X}}_{y}\right),
$$

where  $Y_1, \ldots, Y_{n-1}$  are i.i.d. random vectors with multidimensional truncated distribution  $F_{\mathbf{X}||\mathbf{X}\|}(\cdot | y)$  defined in (2.3), and the random vector  $\mathbf{X}_y$  has the distribution  $F_{\mathbf{X}_y}(\cdot)$  defined in  $(2.5).$ 

These lemmas are proved in Appendix A.2.

## 3 A first multi-normex version: d-Normex

We start building a first multi-normex version, using the Normex approach  $(2.2)$ , then approximating the distribution of the trimmed sum via the CLT and keeping the distribution of the maximum  $\mathbf{X}_{(n)}$ . It is a natural extension to any dimension d of the univariate  $(d = 1)$  Normex distribution as developed in Kratz (2014). Note that, when turning to data, the distribution of  $\mathbf{X}_{(n)}$  may be approximated e.g. via simulations or, as will be done in the MRV-Normex, using another asymptotic theorem, the EV one.

### 3.1 Definition

**Definition 3.1.** The so-called **d-Normex** distribution function is defined, for  $B \subset \mathbb{R}^d$ , as:

$$
G_n(\mathbf{B}) = \mathbb{P}(\mathbf{Z} + \mathbf{X}_{(n)} \in \mathbf{B})
$$

where **Z** is, conditionally on the event  $(\|\mathbf{X}_{(n)}\| = y)$ , a Gaussian random vector with mean  $(n-1)\mu(y)$  and covariance matrix  $(n-1)\Sigma(y)$ , the functions  $\mu(\cdot)$  and  $\Sigma(\cdot)$  are, respectively, the mean vector and covariance matrix of the truncated distribution  $F_{\mathbf{X}||\mathbf{X}||}$  defined in (2.3).

Another way to formulate Definition 3.1 is the following:

$$
G_n(\mathbf{B}) = \int_{\mathbb{R}^d} f_{(n)}(\mathbf{x}) \ \Phi_{(n-1)\mu(\|\mathbf{x}\|), (n-1)\Sigma(\|\mathbf{x}\|)}(\mathbf{B} - \mathbf{x}) \ \mathrm{d}\mathbf{x},\tag{3.8}
$$

where  $\Phi_{m,\Gamma}$  denotes the cdf of the Gaussian vector with mean m and covariance matrix Γ.

#### 3.2 Rate of convergence

Let us turn to the evaluation of the Normex distribution for approximating the distribution of the sum of iid random vectors, studying its rate of convergence. We do it analytically. Although the result holds for any  $\alpha \geq 2$ , we state it when assuming  $\alpha \in (2,3]$ , so that we can compare it with the result obtained via the generalized Berry-Esseen inequality. Then we show numerically the general good fit of d-Normex in an example (see Section 3.3).

The analytical result given in Theorem 3.1 shows that applying Normex method rather than the multivariate CLT improves, as expected, the accuracy of the evaluation of the (tail) distribution of the sum of heavy tailed vectors, with a better rate of convergence than the one of the CLT whenever the shape parameter  $\alpha \in (2,3)$ .

**Theorem 3.1.** Let  $X_1, \ldots, X_n$  be i.i.d. random vectors with parent random vector X with values in  $\mathbb{R}^d$  such that:

- (C1) For all  $y > 0$ , the truncated (w.r.t. the norm) distribution  $F_{\mathbf{X}||\mathbf{X}||}(\cdot | y)$  defined in (2.3) is nondegenerate.
- (C2) The distribution of the rv  $\|\mathbf{X}\|$  is absolutely continuous and regularly varying at infinity:  $\|\mathbf{X}\| \in \mathcal{RV}_{-\alpha}$ , with  $\alpha > 0$ .

Then, for any  $\alpha \in (2,3]$ , there exists a slowly varying function  $L(\cdot)$  such that

$$
\sup_{\mathbf{B}\in\mathcal{C}}\left|\mathbb{P}\left(\mathbf{S}_{\mathbf{n}}\in\mathbf{B}\right)-G_{n}(\mathbf{B})\right|\leqslant L(n)\,n^{-\frac{1}{2}+\frac{3-\alpha}{\alpha}},
$$

where C is the class of all Borel-measurable convex subsets of  $\mathbb{R}^d$ .

Let us briefly indicate how to reach the upper bound of this main result; for more details, see the proof developed in Appendix A.3. First, we use the law of total probability conditioning by  $X_{(n)}$ . Second, we apply the (non-generalized) Berry-Esseen inequality for the truncated r.v.  ${\bf \{Y}_i\}_{i\leq n-1}$  given the event  $(\|{\bf X}_{(n)}\|=y)$ . The right-hand side of the inequality is of the order of  $\frac{1}{\sqrt{2}}$  $\frac{1}{n}\mathbb{E} \|\mathbf{Y}\|^3$ , which is equivalent to  $\frac{1}{\sqrt{3}}$  $\frac{1}{n}\mathbb{E} \|\mathbf{X}_{(n)}\|^{3-\alpha}$  whenever  $\alpha \leqslant 3$ . Finally, to derive the upper bound of the main result, we use that  $\|\mathbf{X}_{(n)}\|$  is of the order  $n^{1/\alpha}$  under  $(C2)$ .

#### Remark 3.1.

- (i) Note that we consider the case  $\alpha \in (2,3]$  since it is the condition under which the Generalized Berry-Esseen holds. For  $\alpha > 3$ , the bound given in Theorem 3.1 is the same as that of Berry-Esseen inequality, making the analytical comparison useless. Indeed, in such a case, the bound  $\frac{1}{\sqrt{2}}$  $\frac{1}{n}\mathbb{E}\|\mathbf{Y}\|^3$  reduces simply to the order  $\frac{1}{\sqrt{3}}$  $\frac{1}{n}$  (see (1.40) in the proof), giving back the same rate as for the CLT. It means to look for an alternative way if we want to study analytically the Normex rate of convergence. We might use Edgeworth expansions, but it evolves too heavy computations (as we could experience for rv (the case  $d = 1$ ), conditioning on  $X_{(n)}$ ). This is why we show numerically the benefit of using Normex distribution, as illustrated in Section 5.
- (ii) The rate of convergence given in Theorem 3.1 is better than the one provided in the generalized Berry-Esseen inequality (Proposition 2.1), whenever  $\alpha \in (2,3)$  (and whatever  $n$ , as

$$
\frac{\alpha-2}{2} < \frac{1}{2} - \frac{3-\alpha}{\alpha}.
$$

Note also that in the case  $\alpha = 3$  and  $\mathbb{E} \|\mathbf{X}\|^3 = \infty$ , the inequality in Theorem 3.1 is slightly sharper than the inequality that can be obtained by the Berry-Esseen theorem (replacing  $n^{-1/2+\varepsilon}, \varepsilon > 0, \text{ by } L(n)n^{-1/2}.$ 

(iii) One can apply the Normex method with any norm on  $\mathbb{R}^d$ , for instance the  $l_1$  norm defined,  $for \mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d, by \|\mathbf{x}\|_1 := \sum_{i=1}^d x_i.$ 

(iv) Considering the  $l_1$  norm, for positive random variables, Condition (C2) translates into the assumption

$$
(C2^*) \qquad S_d := \sum_{i=1}^d X^{(i)} \in \mathcal{RV}_{-\alpha},
$$

where  $X^{(i)}$ , for  $i = 1, \dots, d$ , denote the components of **X**.

We may want to relate this  $\mathcal{RV}$  property on the sum, with conditions on the random vector itself. This topic has already been investigated in the literature; see e.g. Basrak et al. (2002), Barbe et al. (2006), Mainik and Embrechts (2013), and Cuberos et al. (2015). For instance, assuming X multivariate regularly varying,  $X \in \mathcal{MRV}_{-\alpha}(b, \nu)$  (as defined in Definition A.5, Appendix A.1), implies that the sum  $S_d \in \mathcal{RV}_{-\alpha}(b)$ . We will come back on the MRV notion in Section 4.

### 3.3 Example of the Multivariate Pareto-Lomax distribution

Let us consider a d-dimensional random vector  $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})$  having a multivariate Pareto-Lomax( $\alpha$ ) distribution, with  $\alpha > 0$ , *i.e.* with survival distribution function defined, for any non-negative real numbers  $x_1, \ldots, x_d$ , by

$$
\overline{F}_{\mathbf{X}}(x_1,\ldots,x_d):=\mathbb{P}\left(X^{(i)}>x_i,\,i=1,\cdots,d\right)=\left(1+\sum_{i=1}^dx_i\right)^{-\alpha}.
$$

First we compute the statistical characteristics, as (truncated) moments, needed for applying d-Normex. We consider the case  $\alpha \in (2,3]$ , as in Theorem 3.1 (even if the method works for any  $\alpha > 0$ ).

It is straightforward to compute the pdf, marginal distributions, expectation and covariance matrix of a multivariate Pareto-Lomax vector, namely:

$$
f_{\mathbf{X}}(x_1,\ldots,x_d) = \frac{\alpha(\alpha+1)\ldots(\alpha+d-1)}{(1+\sum_{i=1}^d x_i)^{\alpha+d}}, \quad x_1,\ldots,x_d \geq 0; \tag{3.9}
$$

$$
\mathbb{P}(X_j > x) = (1+x)^{-\alpha}, \quad x \ge 0, \ j \in \{1, \dots, d\};\tag{3.10}
$$

$$
\mathbb{E}[X_j] = \frac{1}{\alpha - 1}, \quad \text{Var}(X_j) = \frac{\alpha}{(\alpha - 1)^2 (\alpha - 2)}, \quad j \in \{1, ..., d\} ;
$$

$$
\mathbb{E}[X_j X_i] = \frac{1}{(\alpha - 1)(\alpha - 2)}, \quad \text{Cov}(X_i, X_j) = \frac{1}{(\alpha - 1)^2 (\alpha - 2)}, \quad j \neq i \in \{1, ..., d\} .
$$

Given the expression of the survival distribution of  $X$ , and given that we can apply d-Normex for any norm on  $\mathbb{R}^d$ , we choose the example of the  $L_1$  norm  $\|\mathbf{x}\|_1 := \sum$ d  $i=1$  $x_i$  to simplify the computations. We also take the example of  $d = 3$  for illustration.

We can express the cdf of the rv  $||\mathbf{X}||$ , for  $y > 0$ , as

$$
F_{\|\mathbf{X}\|}(y) = \mathbb{P}\left(\|\mathbf{X}\| \leqslant y\right) = 1 - (1+y)^{-\alpha} - \alpha y (1+y)^{-(\alpha+1)} - \frac{\alpha(\alpha+1)}{2} y^2 (1+y)^{-(\alpha+2)}.
$$

Note that those expressions may also be obtained when applying the more general results (1.61) and (1.62) developed for any norm on  $\mathbb{R}^d$  and for any d; see Appendix A.4.3.

Now, let us compute the moments for the d-dimensional truncated Pareto-Lomax random vector, denoted by Y, having cdf  $F_{\mathbf{X}||\mathbf{X}||}(\cdot | y)$  (see Definition 2.3), expectation  $\mu(y)$  and covariance matrix  $\Sigma(y)$ . We have, for any  $j \in \{1, 2\}$ , if  $\alpha \neq 1$  (which is our case),

$$
\mu_j(y) := \mathbb{E}[Y_j] = \mathbb{E}\left[X_j \mid \|X\| \leq y\right]
$$
  
= 
$$
\frac{1}{F_{\|X\|}(y)} \frac{(y+1)^{-\alpha-2} \left( (y+1)^{\alpha+2} - (\alpha+2) y (1+y (\alpha+1) (\alpha y+3) /6) - 1 \right)}{(\alpha-1)},
$$

and, if  $\alpha \neq 1, 2$  (also our case),

$$
\mathbb{E}\left[Y_{1}Y_{2}\right] = \mathbb{E}\left[X_{1}X_{2} \mid \|\mathbf{X}\| \leqslant y\right] = \frac{1}{F_{\|\mathbf{X}\|}(y)} \frac{1}{(\alpha - 2)(\alpha - 1)} \Big(1 - (y + 1)^{-\alpha - 2} \times
$$
\n
$$
\left((\alpha + 2) y \left((\alpha + 1) y \left(\alpha y \left((\alpha - 1) y/24 + 1/6\right) + 1/2\right) + 1\right) - 1\right)\right),
$$
\n
$$
\mathbb{E}\left[Y_{1}^{2}\right] = \mathbb{E}\left[X_{1}^{2} \mid \|\mathbf{X}\| \leqslant y\right] = \frac{1}{F_{\|\mathbf{X}\|}(y)} \frac{2}{(\alpha - 2)(\alpha - 1)} \Big(1 - (y + 1)^{-\alpha - 2} \times
$$
\n
$$
\Big[ (\alpha + 2) y \Big( y \left(\alpha + 1\right) \left(\alpha y \left(y \left(\alpha - 1\right)/24 + 1/6\right) + 1/2\right) + 1 \Big) \Big),
$$

from which can be deduced the covariance matrix  $\Sigma(y) = (\Sigma_{ij}(y))_{i,j}$ .

Therefore, the Gaussian cdf  $\Phi_{(n-1)\mu(\|\mathbf{x}\|), (n-1)\Sigma(\|\mathbf{x}\|)}$  introduced in Definition 3.1 is explicitly determined, and so is the d-Normex distribution  $G_n$  defined in (3.8).

To illustrate the benefit of the Normex method (see Section 5), whatever the value  $\alpha > 0$ , we draw QQ-plots (see Figure 2 to Figure 4). For that, we simulate a sample from the multivariate Pareto-Lomax( $\alpha$ ) distribution (3.9). We proceed by induction: the first coordinate  $X^{(1)}$  has (1-dimensional) Pareto-Lomax distribution (3.10) with parameter  $\alpha > 0$ ; for the rest of the components, we can easily derive their conditional distribution:

$$
\mathcal{L}\left(X^{(k+1)}\,\big|\,\left(X^{(1)},\cdots,X^{(k)}\right)=(x^{(1)},\cdots,x^{(k)})\right)=(1+x^{(1)}+\cdots+x^{(k)})\,\widehat{X},\quad k\geqslant 1,
$$

where  $\hat{X}$  has the Pareto-Lomax distribution (3.10) with parameter  $\alpha + k$  and is independent of  $X^{(1)}, \cdots, X^{(k)}$ .

# 4 MRV-Normex

Here we investigate a more general version of multi-normex, named MRV-Normex, using an asymptotic theorem for the maximum, namely the Extreme Value (EV) one. Given our focus on the sum of iid heavy-tailed (w.r.t. the norm) random vectors, we consider the standard extreme value theory (EVT) framework of multivariate regularly varying (MRV), a natural extension of the regular variation in a multivariate framework. In fact, to obtain the rate of convergence of this multi-normex approximation, we assume slightly stronger assumption than MRV, asking for a uniform asymptotic independence of the polar coordinates of the random vector, as made explicit in Condition  $(M_{\Theta})$  of Theorem 4.1 below.

In order to obtain the rate of convergence for the MRV-Normex approximation of the sum, we first need to discuss the rate of convergence in EVT to control the difference between the norm of the maximum  $\|\mathbf{X}_{(n)}\|$  and the limit Fréchet distribution. This is the object of the next subsection.

### 4.1 Rate of convergence in the EV theorem. Discussion of its assumptions

After recalling the Extreme Value Theorem, we discuss its rate of convergence, depending on the assumptions.

Extreme Value (EV) Theorem - Let  $\{X_n, n \geq 1\}$  be i.i.d. random variables with c.d.f.  $F_X$ . Denote the maximum of this sequence as  $\max_{1 \leqslant i \leqslant n} X_i$ . Assume that  $F_X$  is in the maximum domain of attraction (MDA) of an extreme-value distribution  $G_{\gamma}$ , denoted by  $F_X \in MDA(G_{\gamma})$ with  $\gamma \in \mathbb{R}$ , i.e. there exist normalizing constants  $a_n > 0$  and  $b_n \in \mathbb{R}$  such that

$$
\mathbb{P}\left(\max_{1\leq i\leq n} X_i \leq a_n x + b_n\right) = F_X^n(a_n x + b_n) \underset{n\to\infty}{\longrightarrow} G_\gamma(x). \tag{4.11}
$$

Let us introduce the real function g defined on  $\mathbb{R}^+$  by :

$$
g := \left(\frac{1}{-\log F_X}\right)^{\leftarrow} \tag{4.12}
$$

 $($   $\leftarrow$  denoting the left-continuous inverse function).

It is straightforward to show that the convergence (4.11) is equivalent to

$$
\lim_{t \to \infty} \frac{g(tx) - g(t)}{a(t)} = \frac{x^{\gamma} - 1}{\gamma}, \quad \forall x > 0,
$$
\n(4.13)

for some  $\gamma \in \mathbb{R}$  and auxiliary positive function a defined on  $\mathbb{R}^+$ , *i.e.* g is of extended regular variation (see Appendix A.1), denoted by  $g \in ERV_{\gamma}(a)$ :

$$
F_X \in MDA(G_\gamma) \quad \Leftrightarrow \quad g \in E\mathcal{RV}_\gamma(a).
$$

Note that this equivalence holds true when replacing g by  $U =$  $\begin{pmatrix} 1 \end{pmatrix}$  $1-F_X$  $\setminus$ , for some auxiliary positive function a (see e.g. de Haan and Ferreira  $(2006)$ , Theorem 1.1.6).

If  $F_X$  is differentiable, then the auxiliary function a in (4.13) can be chosen as  $a(t) := tg'(t)$ , so that

$$
\lim_{t \to \infty} \frac{g(tx) - g(t)}{tg'(t)} = \frac{x^{\gamma} - 1}{\gamma}, \quad \forall x > 0, \quad \text{with } \gamma \in \mathbb{R}.
$$
\n(4.14)

On the assumptions for the rate of convergence in the EV theorem - To describe the rate of convergence in the EV theorem, we first refer to two studies developed under slightly different assumptions, namely Falk and Marohn (1993) with a direct condition on the derivative of the distribution  $F_X$  (see (4.17)), and de Haan and Resnick (1996) assuming a second-order von Mises condition on  $g$  defined in  $(4.12)$ . Note that the results obtained in both studies hold for  $F_X$  belonging to any MDA. Then, focusing on the case of  $F_X \in MDA(Fref)$ , we look for a condition on F involving RV properties to replace the second-order von Mises condition on g and to retrieve the exact rate of convergence described in de Haan and Resnick (1996). This is presented in Proposition 4.2.

 $\triangleright$  DE HAAN AND RESNICK'S RESULT. In de Haan and Resnick (1996), the authors assume that the function g defined in  $(4.12)$  satisfies the second-order von Mises condition, which we recall here.

**Definition 4.1.** (second-order von Mises condition) A twice differentiable function  $g : (0, \infty) \rightarrow$ R satisfies the second-order von Mises condition (shortly,  $g \in 2$ -von Mises( $\gamma, -\rho$ )) for  $\gamma \in \mathbb{R}$ and  $\rho > 0$ , if g' is eventually positive and the function

$$
A(t) := \frac{t g''(t)}{g'(t)} - \gamma + 1
$$

has constant sign near infinity and is such that

$$
A(t) \to 0, \text{ as } t \to \infty, \text{ and } |A| \in RV_{-\rho}.
$$

An equivalent definition (see de Haan and Resnick  $(1996)$ , Theorem 2.1) is that g' has the representation:

$$
g'(t) = k t^{\gamma - 1} \exp\left\{ \int_1^t \frac{A(u)}{u} du \right\}, \quad \text{with } k \neq 0.
$$
 (4.15)

In particular, it implies that (see de Haan and Resnick (1996), Theorem 2.1)

$$
g' \in 2\mathcal{RV}_{\gamma-1,-\rho},
$$

where the  $2\mathcal{RV}$ -class can be defined as (4.18) below (see also Lemma A.1 in Appendix A.1). De Haan and Resnick use the 2-von Mises condition representation (4.15) to derive Potter bounds for  $g'$  and obtain the rate of convergence for  $(4.14)$ :

$$
\frac{[g(tx) - g(t)]/(tg'(t)) - (x^{\gamma} - 1)/\gamma}{A(t)} \to \int_1^x u^{\gamma - 1} \left(\frac{u^{\rho} - 1}{\rho}\right) du, \text{ as } t \to \infty,
$$

and the following rate of convergence in (4.11) for the total variation metric, given in Proposition 4.1.

#### Proposition 4.1 (Theorem 4.1 in de Haan and Resnick (1996)).

Let g be defined in (4.12) such that  $g \in 2$ -von Mises( $\gamma$ , - $\rho$ ) with  $\gamma \in \mathbb{R}$  and  $\rho > 0$ . Then there exists a constant  $C > 0$  (that is defined explicitly) such that

$$
\lim_{n \to \infty} \frac{\sup_{A \in B(\mathbb{R})} \left| \mathbb{P}\left(a_n^{-1}\left(\max_{1 \le i \le n} X_i - b_n\right) \in A\right) - G_\gamma(A) \right|}{|A(n)|} = C,\tag{4.16}
$$

where  $a_n = ng'(n)$  and  $b_n = g(n)$ .

**Remark 4.1.** From the definitions of  $2$ −von Mises and  $RV_{-0}$ , we have

$$
A(t) = L(t)t^{-\rho},
$$

where  $L(\cdot)$  is a slowly-varying function at infinity. From Proposition 4.1, it follows that

$$
\sup_{A\in B(R)} \left| \mathbb{P}\left(a_n^{-1}\left(\max_{1\leqslant i\leqslant n} X_i - b_n\right) \in A\right) - G_{\gamma}(A)\right| \underset{n\to\infty}{\sim} L(n)n^{-\rho}.
$$

 $\blacktriangleright$  FALK AND MAROHN'S RESULT. When assuming that the distribution F belongs to the Fréchet maximum domain of attraction (taking  $\gamma = -\alpha < 0$ ), which is our general assumption, Falk and Marohn propose von-Mises type conditions to obtain an asymptotic bound for the total variation distance in the EV theorem (see Theorem 3.1 in Falk and Marohn (1993)). They show (see Theorems  $2.2 \& 2.4$  in Falk and Marohn (1993)) that these conditions are equivalent to

$$
f_X(t) = ct^{-\alpha - 1}(1 + h(t)), \quad \text{with} \quad h(t) = O(t^{-\rho}), \, \rho > 0,\tag{4.17}
$$

where  $f_X(t)$  denotes the density function of the cdf  $F_X$ . Note that it implies that  $f_X \in RV_{-\alpha-1}$ taking the slowly varying function as a constant.

Looking at the converse statement, the authors propose conditions on the remainder terms of the von-Mises type conditions (see Theorem 3.2 in Falk and Marohn (1993)) to ensure the slowly varying function to be a constant, so that, under those conditions, the total variation distance in the EV theorem implies (4.17).

Note that Condition (4.17) is very close to the characterization of the  $2RV$  class given in (4.18) (see Lemma A.1 in Appendix A.1); nevertheless these conditions do not imply each other.

#### ► OUR PROPOSITION. Assuming  $F \in MDA(Frefchet)$ , we can state the following:

**Proposition 4.2.** Suppose  $\bar{F}_X \in RV_{-\alpha}$ , with  $\alpha > 0$ . The rate of convergence for the EV theorem given in Proposition 4.1 holds when replacing the condition  $g \in 2$ –von Mises( $-\alpha, -\rho$ ) (q being defined in (4.12)), where  $\rho > 0$ , by the condition

$$
f_X(t) = ct^{-\alpha - 1}(1 + h(t)), \text{ with } h \in \mathcal{RV}_{-\beta}, \ \beta > 0. \tag{4.18}
$$

In this case, the function  $A(\cdot)$  in (4.16) belongs to the  $\mathcal{RV}_{-\rho}$  class with  $\rho := -\min\left\{1, \frac{\beta}{\alpha}\right\}$  $\frac{\beta}{\alpha}$ . Indeed, since  $F_X \in MDA(Fref)$ , the Potter bounds (1.29) can be directly established from a  $2RV$  condition on g' via Proposition 4 in Hua and Joe (2011) (see Lemma A.2 in Appendix A.1), which in turn follows from a  $2RV$  condition on  $f_X$  by Lemma 4.1 below (which proof is provided in Appendix A.4.1). This latter condition is equivalent to our assumption (4.18), as stated in Hua and Joe (2011), Lemma 3 (recalled as Lemma A.1 in Appendix A.1). Once we have the Potter bounds, we can replicate exactly the proof of Proposition 4.1 as given in de Haan and Resnick (1996), obtaining the same rate of convergence.

**Lemma 4.1.** If  $f_X \in 2\mathcal{RV}_{-\alpha-1,-\beta}$ , with  $\alpha > 0$  and  $\beta > 0$ , then the derivative g' of g defined in (4.12), satisfies  $g' \in 2\mathcal{RV}_{\frac{1}{\alpha}-1,\rho}$ , where  $\rho := -\min\left\{1, \frac{\beta}{\alpha}\right\}$  $\frac{\beta}{\alpha}\big\}$  (< 0).

Now, let us comment in Remark 4.2 the choice of the normalizing sequences  $(a_n)$  and  $(b_n)$ in  $(4.11)$  of the EV theorem. First, notice that the limit in  $(4.11)$  remains unchanged when replacing  $(a_n)$  and  $(b_n)$  with  $(\tilde{a}_n)$  and  $(\tilde{b}_n)$ , as long as

$$
\lim_{n \to \infty} \frac{\tilde{a}_n}{a_n} = 1 \quad \text{and} \quad \lim_{n \to \infty} \frac{b_n - \tilde{b}_n}{a_n} = 0. \tag{4.19}
$$

#### Remark 4.2.

(i) Remark p.117 in de Haan and Resnick (1996). When  $F_X$  has a pdf, de Haan and Resnick provide universal norming sequences  $(a_n)$  and  $(b_n)$  in (4.11) for the EV theorem, namely

$$
a_n = n g'(n) \quad and \quad b_n = g(n).
$$

They point out that this choice is optimal in the following sense: If one changes  $(a_n)$  and  $(b_n)$  into  $(\tilde{a}_n)$  and  $(\tilde{b}_n)$  such that for some  $c_a, c_b \in \mathbb{R}$ , we have

$$
\lim_{n \to \infty} \frac{1}{A(n)} \left( \frac{\tilde{a}_n}{a_n} - 1 \right) = c_a \quad \text{and} \quad \lim_{n \to \infty} \frac{1}{A(n)} \left( \frac{b_n - \tilde{b}_n}{a_n} \right) = c_b,
$$

then the rate of convergence in (4.16) remains the same up to a constant. But if any of the above limits is infinite, then the supremum in  $(4.16)$  is no longer  $O(A(n))$ .

(ii) If the distribution  $F_X$  of a rv X belongs to the  $2RV_{-\alpha,-\beta}$  class, with  $\alpha > 0$  and  $\beta > 0$ , then we can choose the normalizing sequences  $(a_n)$  and  $(b_n)$  in (4.11) for the EV theorem as

$$
a_n = c^{1/\alpha} n^{1/\alpha}
$$
 with  $c := \lim_{x \to \infty} x^{\alpha} \overline{F}_X(x)$ , and  $b_n = 0$ .

(iii) In fact, the sequences  $(a_n)$  and  $(b_n)$  suggested in (i) and (ii) do not provide an optimal rate of convergence, as can be observed in the following simple example.

**Example 4.1.** Let  $X_1, \dots, X_n$  be a sample from a Pareto-Lomax distribution with parameter  $\alpha > 0$  and survival function

$$
\bar{F}_X(x) = (1+x)^{-\alpha}, \ x > 0.
$$

The Pareto distribution belonging to the maximum domain of attraction of the Fréchet distribution, there exist  $a_n > 0, b_n \in \mathbb{R}$ , such that

$$
F_X^n(a_n x + b_n) \underset{n \to \infty}{\to} e^{-x^{-\alpha}}, \ x > 0. \tag{4.20}
$$

If we choose  $a_n = n^{1/\alpha}$  and  $b_n = 0$ , then we have

$$
F_X^n(a_n x + b_n) = (1 - (1 + n^{1/\alpha} x)^{-\alpha})^n
$$

and the difference in (4.20) is  $O(n^{-1/\alpha})$ . Now, if we take  $b_n = -1$  instead of 0, then

$$
F_X^n(a_n x + b_n) = \left(1 - \frac{x^{-\alpha}}{n}\right)^n
$$

and the difference in (4.20) is  $O(1/n)$ , which is much smaller for  $\alpha > 1$ . So playing with the sequence  $(b_n)$  in this simple example provides a strong improvement in the rate of convergence in (4.20).

### 4.2 Rate of convergence for MRV-Normex

Definitions - While the standard MRV definition is recalled in Appendix A.1 (see Definition A.5), let us give an equivalent MRV definition using the pseudo-polar representation, to better understand why the introduction of this multi-normex version, namely MRV-Normex.

**Definition 4.2.** The random vector  $X \in \mathcal{MRV}_{-\alpha}$ , with  $\alpha > 0$ , if there exists a d-dimensional random vector  $\Theta$  with values in the unit sphere  $S_1$  in  $\mathbb{R}^d$  w.r.t. the norm  $\|\cdot\|$ , such that,  $\forall t > 0$ ,

$$
\frac{\mathbb{P}(\|\mathbf{X}\| > t u, \mathbf{X}/\|\mathbf{X}\| \in \cdot)}{\mathbb{P}(\|\mathbf{X}\| > u)} \xrightarrow{v} t^{-\alpha} \mathbb{P}(\Theta \in \cdot) \quad \text{as} \quad u \to \infty. \tag{4.21}
$$

Using this latter definition of MRV, in particular the random vector  $\Theta$ , and Remark 4.2, we can define the MRV-Normex distribution as follows.

**Definition 4.3.** The so-called MRV-Normex distribution function is defined, for  $B \subset \mathbb{R}^d$ , by:

$$
GM_n(\mathbf{B}) := \mathbb{P}\left(H_{\alpha,n}\,\mathbf{\Theta} + Z \in \mathbf{B}\right). \tag{4.22}
$$

with  $H_{\alpha,n} := a_n H_\alpha + b_n$  where the random variable  $H_\alpha$  (with  $\alpha > 0$ ) is Fréchet distributed (i.e.  $\mathbb{P}(H_{\alpha} \leqslant x) = e^{-x^{-\alpha}}$ , for  $x > 0$ ) and independent of the random vector  $\Theta$  introduced in (4.21), the normalizing sequences satisfy the standard conditions of EV theorem, namely  $a_n = c n^{1/\alpha}$  with  $c^{\alpha} := \lim_{y \to \infty} y^{\alpha} \overline{F}_{\|X\|}(y)$ , and  $b_n = 0$ . The d-dimensional random vector  $Z$ , also assumed to be independent of  $\Theta$ , is, conditionally to the event  $(H_{\alpha,n} = y)$ , with  $y > 0$ , normally  $\mathcal{N}_{(n-1)\mu(y),(n-1)\Sigma(y)}$ -distributed, the mean vector and covariance matrix being those of the truncated distribution  $F_{\mathbf{X}||\mathbf{X}\|}(\cdot|y)$  defined in (2.3).

The MRV-Normex cdf can be rewritten as

$$
GM_n(\mathbf{B}) := \int\limits_0^\infty f_{H_{\alpha,n}}(y) \, \mathbb{P}\left(\, y \, \mathbf{\Theta} + Z_y \in \mathbf{B}\,\right) \mathrm{d}y,\tag{4.23}
$$

where  $Z_y$  is  $\mathcal{N}_{(n-1)\mu(y),(n-1)\Sigma(y)}$ -distributed and independent of  $\Theta$ .

Note that we may choose other normalizing sequences that satisfy (4.19) in the definition of MRV-Normex distribution. Moreover, this distribution may be defined mathematically for any MDA, using the Generalized Extreme Value distribution and the appropriate normalizing sequences  $(a_n)$  and  $(b_n)$ , as the EV theorem (see (4.11)) holds for any MDA. Nevertheless, we focus on the MDA(Fréchet) for which a naive use of a light tail distribution for the sum would lead to a large error.

Rate of convergence for the MRV-Normex approximation - We have the following result, which proof can be found in Appendix A.4.2.

**Theorem 4.1.** Let  $X_1, \ldots, X_n$  be i.i.d. random vectors with parent random vector X with values in  $\mathbb{R}^d$ . Assume the following conditions:

- (C1) given in Theorem 3.1 (namely,  $F_{\mathbf{X} \mid \|\mathbf{X}\| \cdot \| y}$  non degenerate  $\forall y > 0$ );
- $(M_{\|\cdot\|})$  The distribution of the rv  $\|\mathbf{X}\|$  is absolutely continuous and its pdf  $f_{\|\mathbf{X}\|} \in 2\mathcal{RV}_{-\alpha-1,-\beta}$ with  $\alpha > 0, \ \beta > 0;$
- (M<sub>Θ</sub>) There exists a function A such that  $A(t) \to 0$ ,  $|A(t)| \in RV_{-\rho}$  with  $\rho > 0$ , and

$$
\sup_{\mathbf{B}\in\mathcal{S}_1}\left|\mathbb{P}\left(\frac{\boldsymbol{X}}{\|\boldsymbol{X}\|}\in\mathbf{B}\,\bigg|\,\|\boldsymbol{X}\|>t\,\right)-\mathbb{P}\left(\,\boldsymbol{\Theta}\in\mathbf{B}\,\right)\right|\,\underset{t\to\infty}{\sim}A(t).
$$

Then, for any  $\alpha \in (2,3], \beta > 0$  and  $\rho > 0$ , there exists a slowly varying function  $L(\cdot)$  such that

$$
\sup_{\mathbf{B}\in\mathcal{C}}|\mathbb{P}\left(\mathbf{S}_{\mathbf{n}}\in\mathbf{B}\right)-GM_{n}(\mathbf{B})|\leqslant\left(n^{-\frac{1}{2}+\frac{3-\alpha}{\alpha}}+n^{-\frac{\beta}{\alpha}}+n^{-\min(1,\frac{\beta}{\alpha})}\right)L(n),\tag{4.24}
$$

where C is the class of all Borel-measurable convex subsets of  $\mathbb{R}^d$  and  $GM_n$  is defined in (4.23).

#### Remark 4.3.

1. If  $\rho > \alpha$  and  $\beta > \alpha$ , then (4.24) rewrites as

$$
\sup_{\mathbf{B}\in\mathcal{C}}\left|\mathbb{P}\left(\,\mathbf{S}_{\mathbf{n}}\in\mathbf{B}\,\right)-GM_{n}(\mathbf{B})\right|\,\leqslant\,n^{-\frac{1}{2}+\frac{3-\alpha}{\alpha}}L(n),
$$

providing the same rate of convergence given in Theorem 3.1 for d-Normex.

- 2. If the supremum considered in  $(M_{\Theta})$  has a faster speed of convergence than that of  $RV$ , then we can set  $\rho = \infty$  and exclude the term  $n^{-\frac{\rho}{\alpha}}$  from inequality (4.24).
- 3. Discussion on Condition  $(M_{\Theta})$ :
	- (a) First note that assuming  $\|\mathbf{X}\| \in RV$  and Condition  $(M_{\Theta})$ , which requires uniform convergence, is closely related to the MRV definition (4.21). Nevertheless, replacing this technical condition  $(M_{\Theta})$  with (4.21) might be investigated further.
	- (b) Independent polar coordinates If the norm  $\|\mathbf{X}\|$  and the direction  $\mathbf{X}/\|\mathbf{X}\|$  of X are independent, then  $(M_{\Theta})$  is satisfied. Moreover, in the absolutely continuous case,  $\|\mathbf{X}\|$  and  $\mathbf{X}/\|\mathbf{X}\|$  are independent if and only if there exist non negative functions  $h_1$  and  $h_2$  such that the pdf of **X** can be factorized as:

$$
f_{\mathbf{X}}(\mathbf{x}) = h_1(||\mathbf{x}||) h_2(\mathbf{x}/||\mathbf{x}||), \ \mathbf{x} \in \mathbb{R}^d. \tag{4.25}
$$

(c) In some cases, the distribution of the random vector  $\Theta$  can be made explicit, as follows. For any  $t > 0$  and  $B \in \mathcal{S}_1$ ,

$$
\mathbb{P}\left(\left.\frac{\mathbf{X}}{\|\mathbf{X}\|}\in \mathbf{B}\,\middle\|\, \|\mathbf{X}\|\geqslant t\,\right)=\frac{1}{\overline{F}_{\|\mathbf{X}\|}(t)}\int\limits_{\frac{\mathbf{u}}{\|\mathbf{u}\|}\in \mathbf{B},\|\mathbf{u}\|\geqslant 1}\frac{f(t\mathbf{u})\,t^d}{\overline{F}_{\|\mathbf{X}\|}(t)}\mathrm{d}\,\mathbf{u}.
$$

If the latter integral has a limit, as  $t \to \infty$ , of the form  $\frac{\mathbf{u}}{\|\mathbf{u}\|} \in B, \|\mathbf{u}\| \geqslant 1$  $f_{\Theta}(\mathbf{u}) \, \mathrm{d}\mathbf{u},$ 

then this limit defines the distribution of  $\Theta$ :

$$
\mathbb{P}(\Theta \in \mathbf{B}) = \int_{\frac{\mathbf{u}}{\|\mathbf{u}\| \in \mathbf{B}, \|\mathbf{u}\| \geqslant 1}} f_{\Theta}(\mathbf{u}) \, \mathrm{d}\mathbf{u},
$$

from which we deduce that

$$
\sup_{\mathbf{B}}\left|\mathbb{P}\left(\left.\frac{\mathbf{X}}{\|\mathbf{X}\|}\in\mathbf{B}\,\middle\|\,\|\mathbf{X}\|\geqslant t\,\right)-\mathbb{P}\left(\,\Theta\in\mathbf{B}\,\right)\right|=\frac{1}{2}\int\limits_{\|\mathbf{u}\|\geqslant 1}\left|\frac{f(t\mathbf{u})\,t^d}{\overline{F}_{\|\mathbf{X}\|}(t)}-f_\Theta(\mathbf{u})\right|\mathrm{d}\mathbf{u}.
$$

Further details are given in Appendix A.4.3.

### 4.3 Examples

Let us develop two examples such that the marginal distributions are Pareto-Lomax, as in Example 3.3, so that it allows comparison with d-Normex. We consider two cases for the parent random vector  $X$ , when assuming its components to be, on one hand, independent, on the other hand, related with a survival Clayton copula. These are standard examples considered

in the actuarial and risk literature (see e.g. Das and Kratz (2020)), in particular in reinsurance context for the Clayton copula is (see e.g. Dacorogna et al. (2018) and references therein). This second example includes itself two cases, when the polar coordinates of the considered vector X are dependent (but asymptotically independent), and when they are independent. This latter case corresponds to Example 3.3.

We check that the conditions of Theorem 4.1 are satisfied whenever  $\alpha \in (2,3]$  (recall that this constraint on  $\alpha$  appears only for the analytical comparison with the generalized Berry-Esseen inequality), but apply the MRV-Normex distribution for any positive  $\alpha$ , as the construction via asymptotic theorems remains valid whatever the value of this parameter.

#### 4.3.1 Independent Pareto-Lomax marginals

Assume the components of the random vector **X** to be iid with Pareto-Lomax $(\alpha)$  distribution (3.10). Then its (non truncated) moments remain the same as in Example 3.3 and its covariance matrix is diagonal. As we developed Example 3.3 with the  $L_1$ -norm, let us switch here to the L<sup>∞</sup>-norm  $\|\cdot\|_{\infty}$ , more convenient in terms of computations in this framework.

The cdf of the norm of the vector **X** being, for  $\alpha > 0$ ,

$$
F_{\|\mathbf{X}\|_{\infty}}(y) = \mathbb{P}\left(\max\left\{X^{(1)}, \cdots, X^{(d)}\right\} \leqslant y\right) = (1 - (1 + y)^{-\alpha})^d, \text{ for } y > 0,
$$

straightforward calculations give the following expressions for the truncated moments:

$$
\mu^{(1)}(y) = \mathbb{E}\left(X^{(1)} \mid \|\mathbf{X}\|_{\infty} \leqslant y\right) = \mathbb{E}\left(X^{(1)} \mid X^{(1)} \leqslant y\right)
$$
  
= 
$$
\frac{1}{1 - (1 + y)^{-\alpha}} \left(\frac{1}{\alpha - 1} \left(1 - (1 + y)^{-\alpha + 1}\right) - y(1 + y)^{-\alpha}\right),
$$
  

$$
\mathbb{E}\left(\left(X^{(1)}\right)^{2} \mid \|\mathbf{X}\|_{\infty} \leqslant y\right) = \frac{1}{1 - (1 + y)^{-\alpha}} \frac{2 - (1 + y)^{-\alpha} \left(\alpha y \left((\alpha - 1) y + 2\right) + 2\right)}{(\alpha - 2) (\alpha - 1)},
$$

and zero for the truncated covariances.

When looking for the distribution of  $\Theta$ , notice that, for any  $i \neq j$ , for any  $\varepsilon_i > 0$  and  $\varepsilon_j > 0$ , we have

$$
\lim_{n\to\infty}\mathbb{P}\left(X^{(i)}>\varepsilon_i t, X^{(j)}>\varepsilon_j t \mid ||\mathbf{X}||_{\infty}>t\right)=0.
$$

Therefore, the distribution of the random vector  $\Theta$  is discrete on the unit sphere  $S_1$ , with values given by the basis vectors  $e_i = (0, \dots, 1, \dots, 0)$  (where 1 is for the *i*-th component). It is straightforward to verify that  $F_{\|\mathbf{X}\|} \in 2\mathcal{RV}_{-\alpha,-\alpha}$ , so that  $(M_{\|\cdot\|})$  is satisfied, and that Condition  $(M_{\Theta})$  holds with auxiliary function  $A(\cdot) \in \mathcal{RV}_{-\alpha}$ .

Finally, one may chose the normalizing sequences as  $a_n = (dn)^{1/\alpha}$  (according to Remark 4.2(i) and  $b_n = -1$  as in Example 4.1 of Remark 4.2(iii).

The numerical implementation of this MRV-Normex approximation is then developed in Section 5, along with that of the d-Normex one, for any positive  $\alpha$ , both multi-normex methods being compared to the Gaussian approximation whenever  $\alpha \geq 2$ . The QQ-plots are drawn in Figure 5.

#### 4.3.2 Pareto-Lomax marginal distribution with survival Clayton copula

We introduce in this example some dependence among the components of  $X$ , choosing a survival Clayton copula (so, with upper tail dependence). To lighten the expressions of the computed moments, we choose  $d = 2$ .

We consider  $\mathbf{X} = (X_1, X_2)$  with identical Pareto-Lomax  $(\alpha, 1)$  marginal distributions,  $\alpha > 1$ , i.e.

$$
\bar{F}_1(x) = \bar{F}_2(x) = (1+x)^{-\alpha}, \quad \forall x > 0,
$$

and survival Clayton copula on  $[0, 1]^2$ , with parameter  $\theta > 0$ , defined by

$$
\mathbb{P}\left(X_1 > x_1, X_2 > x_2\right) = \left[ (1+x_1)^{\alpha\theta} + (1+x_2)^{\alpha\theta} - 1 \right]^{-1/\theta},
$$

with pdf  $f(x_1, x_2) = \alpha^2 (1 + \theta) (1 + x_1)^{\alpha \theta - 1} (1 + x_2)^{\alpha \theta - 1} ((1 + x_1)^{\alpha \theta} + (1 + x_2)^{\alpha \theta} - 1)^{-\frac{1}{\theta} - 2}.$ 

Considering the  $L^{\infty}$ -norm,  $\|\cdot\|_{\infty}$ , the survival cdf of the norm of the vector is:

$$
\overline{F}_{\|\mathbf{X}\|_{\infty}}(t) = \mathbb{P}\left(\max\left\{X_1, X_2\right\} > t\right) = 2(1+t)^{-\alpha} - (2(1+t)^{\alpha\theta} - 1)^{-1/\theta}.
$$

We computed the truncated moments (of order 1 and 2), using an integral calculator (based on Maxima, a computer algebra system developed by W. Schelter, MIT), providing explicit but long expressions, that is why we do not display them here (and give them in Appendix A.7). For positive  $u_1$  and  $u_2$  such that max  $\{u_1, u_2\} \geq 1$ , we can write

$$
\frac{f(t\mathbf{u})t^d}{\overline{F}_{\|\mathbf{X}\|_{\infty}}(t)} = \frac{\alpha^2(1+\theta)\left(u_1 + \frac{1}{t}\right)^{\alpha\theta-1}\left(u_2 + \frac{1}{t}\right)^{\alpha\theta-1}}{(2-2^{-1/\theta})\left(1+O(\frac{1}{t})\right)\left(\left(u_1 + \frac{1}{t}\right)^{\alpha\theta} + \left(u_2 + \frac{1}{t}\right)^{\alpha\theta}\right)^{1/\theta+2}}
$$

from which we deduce the limit as  $t \to \infty$ , namely

$$
\lim_{t \to \infty} \frac{f(t\mathbf{u}) t^d}{\overline{F}_{\|\mathbf{X}\|_{\infty}}(t)} = \frac{\alpha^2 (1+\theta) u_1^{\alpha \theta - 1} u_2^{\alpha \theta - 1}}{(2 - 2^{-1/\theta}) \left(u_1^{\alpha \theta} + u_2^{\alpha \theta}\right)^{1/\theta + 2}} =: f_{\Theta}(\mathbf{u}).
$$

Note that, for  $\|\cdot\| = \|\cdot\|_{\infty}$ , the function  $\frac{f(t\mathbf{u})t^d}{\overline{F}_{\text{max}}(t)}$  $\frac{f(t\mathbf{u})t^{\alpha}}{F_{\|\mathbf{x}\|}(t)}$  depends on t, therefore the random variable  $\|\mathbf{X}\|$  and random vector  $\mathbf{X}/\|\mathbf{X}\|$  are not independent.

They will be independent when replacing the  $L^{\infty}$ -norm with the  $L^{1}$ -norm  $(\Vert \cdot \Vert = \Vert \cdot \Vert_{1})$  and choosing  $\alpha\theta = 1$ ; this corresponds to the Pareto-Lomax Example 3.3.

Let us check the conditions of Theorem 4.1. It is straightforward to check that  $F_{\|\mathbf{X}\|} \in$  $2\mathcal{RV}_{-\alpha,-\min(\alpha\theta,1)}$ , so that  $(M_{\|\cdot\|})$  is satisfied. Some computations are required for Condition  $(M_{\Theta})$ . We keep the maximum norm, *i.e.*  $\|\cdot\| = \|\cdot\|_{\infty}$ , so that we exhibit an example with dependence between the polar coordinates (but with asymptotic independence), but consider

the case  $\alpha\theta = 1$  to simplify the computations. We obtain:

$$
\sup_{B} \left| \mathbb{P} \left( \frac{\mathbf{X}}{\|\mathbf{X}\|} \in B \,|\, \|\mathbf{X}\| \geq t \right) - \mathbb{P} \left( \Theta \in B \right) \right| = \frac{1}{2} \int_{\|\mathbf{u}\| \geq 1} \left| \frac{f(t\mathbf{u})t^{d}}{\overline{F}_{\|\mathbf{X}\|}(t)} - f_{\Theta}(\mathbf{u}) \right| d\mathbf{u}
$$
\n
$$
= \frac{\alpha(\alpha + 1)}{2t} \int_{\|\mathbf{u}\| \geq 1} t \left| \frac{t^{2}}{\overline{F}(t)(1 + t|\mathbf{u}|)^{\alpha + 2}} - \frac{1}{c_{\alpha}|\mathbf{u}|^{\alpha + 2}} \right| d\mathbf{u}
$$
\n
$$
= \frac{\alpha(\alpha + 1)}{2t} \int_{\|\mathbf{u}\| \geq 1} t \left| \frac{c_{\alpha}|\mathbf{u}|^{\alpha + 2} - \overline{F}(t)t^{\alpha}(|\mathbf{u}| + \frac{1}{t})^{\alpha + 2}}{\overline{F}(t)t^{\alpha}(|\mathbf{u}| + \frac{1}{t})^{\alpha + 2}} c_{\alpha}|\mathbf{u}|^{\alpha + 2}} \right| d\mathbf{u}, \tag{4.26}
$$

where  $c_{\alpha} = (2 - 2^{-\alpha})$  and  $|u| = u_1 + u_2$ .

We can easily find a majorant of type  $c/|\mathbf{u}|^{\alpha+2}$  for the integrand (4.26), and, noticing that this integrand converges, as  $t \to \infty$ , to  $\frac{|\hat{c}_{\alpha}|\mathbf{u}| + c_{\alpha}(\alpha+2)|}{2 + |\alpha|^2}$  $\hat{c}_{\alpha}^{2}|\mathbf{u}|^{\alpha+3}$ , with  $\hat{c}_{\alpha} := 2^{-\alpha-1} - 2$ , we can conclude, via the dominated convergence theorem, that

$$
\int_{\|\mathbf{u}\| \geqslant 1} t \left| \frac{c_{\alpha} |\mathbf{u}|^{\alpha+2} - \overline{F}(t) t^{\alpha} (|\mathbf{u}| + \frac{1}{t})^{\alpha+2}}{\overline{F}(t) t^{\alpha} ((|\mathbf{u}| + \frac{1}{t})^{\alpha+2}) c_{\alpha} |\mathbf{u}|^{\alpha+2}} \right| d\mathbf{u} \underset{t \to \infty}{\to} \int_{\|\mathbf{u}\| \geqslant 1} \frac{|\hat{c}_{\alpha} |\mathbf{u}| + c_{\alpha} (\alpha+2)|}{c_{\alpha}^2 |\mathbf{u}|^{\alpha+3}} d\mathbf{u} < \infty. \tag{4.27}
$$

Combining (4.26) and (4.27) provides that Condition  $(M_{\Theta})$  holds with  $A(t) = \frac{C_{\alpha}}{t}$  for some constant  $C_{\alpha} \in (0,\infty)$ .

As in the previous example (case of independent components), one may chose the normalizing sequences  $a_n = (c_\alpha n)^{1/\alpha}$  (see Remark 4.2, (i)) and  $b_n = -1$  (see Remark 4.2, (iii)).

We refer to the next section for the numerical implementation of this example; see Figures 6a & 6b for the QQ-plots. As we are in dimension 2, we also provide (ranked) scatter plots to compare the various approximations of the sum distribution, varying the parameters, in particular the dependence parameter  $\theta$ ; see Figures 7 & 8.

# 5 QQ-plots of the various examples, illustrating both versions of the multi-normex method

### 5.1 Construction of the QQ-plots

Considering the examples given so far, we illustrate the benefit of the multi-normex method on d−dimensional QQ-plots based on geometrical quantiles. We propose a brief overview of this latter notion in Appendix A.5, but refer mainly to Dhar et al. (2014) for definitions and detailed explanations. Here are a few first key ideas of these objects, to help interpret the plots displayed in this section.

The geometrical quantile, as given in Definition A.7 (see Appendix A.5), is a generalization to

higher dimension of the 1-dimensional quantile that can be defined as the solution of some optimization problem (see (1.63) in Appendix A.5). So, one can formulate the same optimization problem in the d−dimensional case, for which the solution will be named geometrical quantile. In this way, the geometrical quantile is a point of  $\mathbb{R}^d$ . While 1-dimensional quantiles are parameterized by the interval (0, 1), multidimensional ones are parameterized by a d−dimensional unit ball from  $\mathbb{R}^d$ . In this context, we refer to vectors of the unit ball as *levels*. Although geometrical quantiles reflect the structure of a d−dimensional distribution, they are abstract objects and do not have a nice interpretation as 1-dimensional quantiles do. Only the median has a geometrical sense: given a random vector, the median is a point of  $\mathbb{R}^d$  such that the overall sum of the distances from this point to all values of the random vector is minimum (note that distances are multiplied by the 'probabilities' that the vector takes the considered values, respectively). Moreover, if the vector has a finite second moment, then its extreme quantiles will share the same speed of convergence towards infinity as any vector having the same covariance matrix (see Girard and Stupfler (2015) and Girard and Stupfler (2017)). Nevertheless, we can construct QQ-plots with these geometrical quantiles, with the aim at comparing d−dimensional distributions by comparing the plots between each other.

The construction of QQ-plots is similar as in the 1-dimensional case. Considering two distributions on  $\mathbb{R}^d$ , we can solve the optimization problem for a fixed number of levels, say N, and obtain two sets of N geometrical quantiles:  $\{q_1, \ldots, q_N\}$  for the first distribution, and  $\{q'_1, \ldots, q'_N\}$  for the second. Then we draw a 2-dimensional QQ-plot for each component of the  $\mathbb{R}^d$ -geometrical quantiles to get a visualization, obtaining d QQ-plots:  $\big\{\big(q_{i,j}, q'_{i,j}\big)$  ;  $j = 1, \cdots, N\big\}$ , for  $i = 1, \dots, d$ . In Dhar et al. (2014), the authors proved that all the points (pairs of quantiles) lie close to a straight line with slope 1 and intercept 0 if and only if the two distributions are equal.

Turning to our previous examples, we consider multidimensional distributions with Pareto-Lomax( $\alpha$ ) marginals defined in (3.10), varying their heaviness through the parameter  $\alpha \in$  $\{1.5, 2.3, 3.5\}$ , with the different structures of dependence given in Examples 3.3 & 4.3. We choose  $n = 52$  and  $d \in \{2, 3\}$ . For each case, we evaluate both multi-normex distributions and the normal distribution (except for  $\alpha = 1.5$ ) obtained by the CLT; those distributions are evaluated empirically, via simulations, each sample being of size  $10<sup>7</sup>$ . Then we proceed to their comparisons via the QQ-plots.

We construct the QQ-plots in 4 main steps:

- 1. Simulate all the distributions to be compared: the distribution of the sum  $S_n$ , the normal distribution from CLT, the d-Normex distribution  $G_n$ , and the MRV-Normex distribution  $MG_n$ . Namely,
	- (a) To obtain a simulated sample (of size  $10^7$ ) for the sum  $S_n$ , we simulate  $n \times 10^7$ random vectors X from the considered distribution.
	- (b) To obtain a simulated sample from the d-Normex distribution defined in (3.8): First, we build n samples (of size  $10^7$ ) from the d-dimensional multivariate Pareto distribution, from which we deduce a sample (of size  $10<sup>7</sup>$ ) for the (d-dimensional) maximum  $\mathbf{X}_{(n)}$  (see (1.1)). Second, for each element of the latter sample, we calculate its norm

 $y = ||\mathbf{X}_{(n)}||$  and simulate a normal vector with mean  $(n-1)\mu(y)$  and covariance matrix  $(n-1)\Sigma(y)$  (described in Definition 3.1), collecting then 10<sup>7</sup> Gaussian vectors. Finally, we sum maximum and normal vectors to produce a sample (of size  $= 10^7$ ) from the d-Normex distribution.

- (c) In order to simulate the MRV-Normex distribution defined in (4.22), we start simulating a sample (of size  $10^7$ ) for the vector  $\Theta$  (representing the direction) and an independent sample for the Fréchet distributed rv  $H_{\alpha,n}$  introduced in (4.23). Next, we collect the 10<sup>7</sup> normal vectors with mean  $(n - 1)\mu(y)$  and covariance matrix  $(n-1)\Sigma(y)$ , where, now,  $y = H_{\alpha,n}$ . Finally, we aggregate all the constructed samples according to (4.22).
- 2. Fix the set of levels  $\mathcal{L} \subset \{ \mathbf{v} \in \mathbb{R}^d, \|\mathbf{v}\| < 1 \}$  with different lengths and directions. We choose 10 lengths,  $||\mathbf{v}|| \in \{0, 0.2, 0.4, 0.6, 0.8, 0.9, 0.9225, 0.945, 0.9675, 0.99\}$  (half for the body of the distribution and half for its tail), and all the directions with factor  $\frac{\pi}{4}$ . It represents a total of 235 vectors for  $d = 3$  and of 145 for  $d = 2$ .
- 3. Calculate the geometrical quantiles for the simulated samples of all considered distributions. It means to solve the numerically optimization problem (1.65) for each empirical distribution and for all levels **v** from the set  $\mathcal{L}$ .
- 4. Draw a QQ-plot for the three pairs (or two, when the CLT cannot be applied): (sum, CLT), (sum, d-Normex) and (sum, MRV-Normex).

Note that the numerical implementation has been performed with Python (SciPy library). The scipy.optimize.minimize function based on the quasi-Newton method of Broyden, Fletcher, Goldfarb, and Shanno (see p.136 in Nocedal and Wright (2006)) has helped solve the numerical optimization problem (1.65). The gradient for the function in (1.65) has been calculated analytically. The computation time of one geometrical quantile was, on average, 53 seconds (on the computer i7 2GHz, 16 GB RAM).

### 5.2 Multivariate Pareto-Lomax distribution - Example 3.3

We first show the QQ-plots for the multivariate Pareto distribution, developed in Section 3.3, considering the dimension  $d = 3$ , the number of summands  $n = 52$ , and varying  $\alpha$ . We choose a small number of summands to highlight the performance of multi-normex methods, even in this case. The QQ-plots are given on a same row for each of the three components, and for each given approximation method (CLT, d-Normex and MRV-Normex). As expected, the plots are similar componentwise. A zoom of the center of the graph is given in the upper left corner of each plot, since it is the only region where the quantiles are so concentrated, making it more difficult to judge if they form a straight line (Note that an html version of all plots, with the possibility to zoom anywhere, is available on the website [crear.essec.edu\)](http://crear.essec.edu/members). It should be noted that the center corresponds not only to levels with length less than or equal to 0.9 (marked in blue), but also to extreme levels (in red) with length greater than 0.9.

CASE  $\alpha \in (2,3]$ . Figure 2 exhibits the QQ-plots in the case  $\alpha = 2.3$ , i.e. in the framework of the multi-normex theorems, when using the Gaussian approximation (via the CLT) and both multinormex approximations. We observe that all the points (pairs of quantiles) for d-Normex and MRV-Normex lie much closer to the line with slope 1 and intercept 0 than for the Gaussian distribution. Thus, we can see numerically that both multi-normex approximations better describe the distribution of the vectorial sum  $S_n$ , as proved analytically in Theorems 3.1 and 4.1. While the QQ-plots look quite similar for the two versions of multi-normex approximations, when zooming, the fit is slightly better when using the d-Normex distribution than the MRV-Normex one that uses the Fréchet distribution for the distribution of the rescaled maximum.

ARBITRARY  $\alpha > 0$ . Here we consider other examples of heavy tails than the case  $\alpha \in (2,3]$ , to illustrate the benefit of using Normex distributions rather than applying the generalized CLT with a Gaussian distribution (for finite variance) or a stable one (when the variance is infinite). We do it numerically as the upper bound of the Berry-Esseen inequality does not allow us to compare analytically the rates of convergence in terms of the number of summands n. Nevertheless, the constant given in terms of  $\alpha$  and the number k of largest order statistics when trimming the sum, may make a difference, as noticed in the 1-dimensional case (see Kratz (2014)). We give two examples, when  $\alpha = 1.5$  (Figure 3), case where the summands have no variance, and when  $\alpha = 3.5$  (Figure 4), *i.e.* beyond the frame of the generalized Berry-Esseen inequality. In Figure 3, we only have two rows as the CLT does not apply and we did not build the QQ-plot for the stable distribution. The fit looks very good for both multinormex methods, with barely no difference of fit between the two, as shown in the zoomed part. Whenever  $\alpha = 3.5$ , we again clearly observe in Figure 4 an overall fit that gives the advantage to the multi-normex distributions, but with more difference, when zooming, between  $d$ -normex and MRV-normex than in the previous cases  $\alpha = 1.5$  and  $\alpha = 2.3$ .



Figure 2: 3-dimensional QQ-plots for the empirical distribution (sample size  $= 10^7$ ) of the sum of 52 iid trivariate Pareto ( $\alpha = 2.3$ ) random vectors, with three different approximations of the sum distribution: CLT (first row), d-Normex (second row) and MRV-Normex (third row). Each column corresponds to a component (from the 1st to the 3rd). The red points on the plots correspond to extreme geometric quantiles (when the norm of the parameterized vectors is greater than 0.9)



Figure 3: 3-dimensional QQ-plots for the empirical distribution (sample size  $= 10^7$ ) of the sum of 52 iid trivariate Pareto ( $\alpha = 1.5$ ) random vectors, with two different approximations of the sum distribution: d-Normex (first row) and MRV-Normex (second row). Each column corresponds to a component (from the 1st to the 3rd). The red points on the plots correspond to extreme geometric quantiles (when the norm of the parameterized vectors is greater than 0.9)



Figure 4: 3-dimensional QQ-plots for the empirical distribution (sample size  $= 10^7$ ) of the sum of 52 iid trivariate Pareto ( $\alpha = 3.5$ ) random vectors, with three different approximations of the sum distribution: CLT (first row), d-Normex (second row) and MRV-Normex (third row). Each column corresponds to a component (from the 1st to the 3rd). The red points on the plots correspond to extreme geometric quantiles (when the norm of the parameterized vectors is greater than 0.9).

### 5.3 Pareto-Lomax marginal distributions with various dependence structures - Examples 4.3

Here, we consider the examples developed in Section 4.3, with Pareto-Lomax $(\alpha)$  marginal distributions, various dependence structures and the norm  $\|\cdot\|_{\infty}$  instead of  $\|\cdot\|_{1}$  as in the previous subsection. Figure 5 displays QQ-plots for independent components of the vector X, taking  $\alpha = 2.3$  to be in the half-closed interval  $(2,3]$  considered in our theorems. Here also, the good fit of the multi-normex distributions appears clearly, in particular when comparing with the Gaussian approximation. Nevertheless, the difference between the two multi-normex distributions is more pronounced in the center, as can be observed in the zoomed part.

When the dependence structure is given via a survival Clayton copula with parameter  $\theta$ , we provide the QQ-plots in Figure 6a assuming  $\alpha\theta = 0.5$ . The previous observations hold too, with an increasing difference in the center between the two multi-normex distributions. The case  $\alpha\theta = 1$  is illustrated in Figure 6b. Note that this choice of  $\theta = 1/\alpha$  is a standard example in the literature as it makes analytical computations much more tractable.

This case  $\alpha\theta = 1$  would also correspond to Example 3.3 if choosing the  $L_1$ -norm  $\|\cdot\|_1$ , allowing then a comparison of the results obtained respectively with the two norms. To easy such a numerical comparison based on norms, we reduce the dimension of the vectors to  $d = 2$  and provide (ranked) scatter plots, simulating  $52 \times 10^4$  observations for **X** (hence a sample of size  $10<sup>4</sup>$  for the bivariate sum  $\mathbf{S}_n$ , with  $n = 52$ ). We display the (ranked) scatter plots on two rows, the first one for the case  $\|\cdot\|_{\infty}$  and the second for  $\|\cdot\|_{1}$ . Fixing  $\alpha\theta = 1$ , we propose two choices for  $\alpha$  to also observe the impact of the upper tail dependence in the survival Clayton copula: (i)  $\alpha = 2.3$ , as previously, implying that  $\theta \approx 0.43$  (weak upper tail dependence) and (ii)  $\alpha = 1.01$ to increase the value of  $\theta$  (close to 1), hence the upper tail dependence, while keeping  $\alpha > 1$ (to have a finite expectation). Note that the CLT does not hold in the case (ii). We exhibit the scatter plots for the case (i) in Figure 7: It is obvious that the Gaussian approximation given by the CLT does not reflect the distribution of  $S_n$ , while both multi-normex scatter plots look more or less similar as that of  $S_n$ , whatever the norm. It corresponds to the analytical results (Theorems 3.1 and 4.1), which are independent of the norm. Turning to (ii), as we have a stronger upper tail dependence, we display ranked scatter plots (rather than scatter plots). We observe in Figure 8 that the dependence structure looks alike among all the plots. Also for this range of  $\alpha$ , the norm does not seem to have any impact.



Figure 5: 3-dimensional QQ-plots for the empirical distribution (sample size  $= 10^7$ ) of the sum of 52 iid random vectors with independent Pareto-Lomax components of the vectors ( $\alpha = 2.3$ ), with three different approximations of the sum distribution: CLT (first row), d-Normex (second row) and MRV-Normex (third row). Each column corresponds to a component (from the 1st to the 3rd). The red points on the plots correspond to extreme geometric quantiles (when the norm of the parameterized vectors is greater than 0.9)



#### Pareto-Lomax with various Clayton copulas ( $\alpha = 2.3$ )

Figure 6: 2-dimensional QQ-plots for the empirical distribution (sample size  $= 10^7$ ) of the sum of 52 iid random vectors with Pareto-Lomax marginal distributions ( $\alpha = 2.3$ ) joint by a Clayton copula with parameter  $\theta$  such that  $\alpha\theta = 0.5$  for the case (a), and  $\alpha\theta = 1$  for the case (b). The three rows correspond to the approximations of the sum distribution: CLT (first row), d-Normex (second row) and MRV-Normex (third row). For each case (a) and (b), the two columns correspond to the 1st and the 2nd components. The red points on the plots correspond to extreme geometric quantiles (when the norm of the parameterized vectors is greater than 0.9).



Figure 7: Scatter plots for the Pareto-Lomax( $\alpha = 2.3$ ) marginal distributions with survival Clayton copula with parameter  $\theta$  such that  $\alpha\theta = 1$ . The first column corresponds to the simulated sample (sample size  $= 10<sup>4</sup>$ ) of the sum of 52 iid random vectors while the three next columns consider the different approximations of the sum distribution: CLT (2nd column), d-Normex (3rd column) and MRV-Normex (4th column). For the multi-normex approximations, two norms have been chosen, the  $\|\cdot\|_{\infty}$ -norm given in the 1st row, and  $\|\cdot\|_1$ -norm in the second one.



Figure 8: Ranked scatter plots for the Pareto-Lomax $(\alpha = 1.01)$  marginal distributions with survival Clayton copula with parameter  $\theta$  such that  $\alpha\theta = 1$ . The first column corresponds to the simulated sample (sample size  $=10^7$ ) of the sum of 52 iid random vectors while the two next columns consider the multi-normex approximations: d-Normex (2nd column) and MRV-Normex (3rd column). Two norms have been chosen, the  $\|\cdot\|_{\infty}$ -norm given in the 1st row, and  $\|\cdot\|_{1}$ -norm in the second one.

# 6 Conclusion

The purpose of this study was to build a sharp approximation of the whole distribution of the sum of iid random vectors under the presence of heavy tails. It has been done by extending the normex approach from a univariate to a multivariate framework, combining mean and extreme behaviors. We proposed two possible multi-normex distributions, named d-Normex and MRV-Normex. Both rely on the Gaussian distribution for describing the mean behavior, via the CLT, while the difference between the two versions comes from using the EV theorem or the exact distribution for the maximum. The main theorems establish the rate of convergence of each version of the multi-normex distributions towards the distribution of the sum. This is done analytically whenever the shape parameter  $\alpha$  of the tail of the marginal distribution belongs to the interval (2, 3], making the comparison with the generalized Berry-Esseen inequality relevant. For the MRV-Normex, second order regular variation conditions are needed to obtain the main theorem. These conditions are discussed w.r.t. the Extreme Value Theory literature. Numerical comparisons are developed for any value of  $\alpha$ , for both multi-normex distributions, considering examples with different structure of dependence for the random vectors. Illustrations are made through multidimensional QQ-plots based on geometrical quantiles.

We focused on the case of heavy tailed random vectors, as it is of most interest in the risk literature. Nevertheless, this method could be extended to light tails vectors (whenever  $1/\alpha =$ 0) as the rate of convergence for the EV theorem is also known in such a case. It would then require to introduce a specific metric to evaluate the error for the whole distribution taking into account the impact of extremes.

Moreover, the MRV-Normex approach has been developed conditioning on the norm of the maximum. It could be done conditioning on the maximum itself (a vector). To widen the applicability of the multi-normex methods, simple approximations of truncated moments could also be suggested (as numerical ones or evaluating them using e.g. a Pareto approximation). Finally, generalization of multi-normex distributions will be studied when introducing dependence between random vectors, then considering random processes. We intend to explore such topics in the near future.

# Acknowledgments

Evgeny Prokopenko acknowledges the support received from the National Research Agency of the French government through the program "Investment for the future" (ANR-16-IDEX-0008 CY Initiative) during his postdoctoral fellowship at ESSEC CREAR.

# References

- Barbe, P., A. Fougères, and C. Genest (2006). "On the Tail Behavior of Sums of Dependent Risks". In:  $ASTIN$  Bulletin 36.2, pp. 361–373. DOI: [10.1017/S0515036100014550](http://dx.doi.org/10.1017/S0515036100014550).
- Basrak, B., R. Davis, and T. Mikosch (2002). "A characterization of multivariate regular varia-tion". In: Annals of Applied Probability 12.3, pp. 908–920. DOI: [10.1214/aoap/1031863174](http://dx.doi.org/10.1214/aoap/1031863174).
- Beirlant, J., Y. Goegebeur, J. Segers, and J.L. Teugels (2004). Statistics of extremes: theory and applications. Wiley Series in Probability and Statistics. John Wiley and Sons, Ltd. DOI: [10.1002/0470012382](http://dx.doi.org/10.1002/0470012382).
- Bhattacharya, R. N. and R. R. Rao (2010). Normal Approximation and Asymptotic Expansions (Classics in Applied Mathematics). Classics in applied mathematics. SIAM,  $xxii+316$  p. doi: [10.1137/1.9780898719895](http://dx.doi.org/10.1137/1.9780898719895).
- Billingsley, P. (1968). Convergence of Probability Measures. John Wiley and Sons.
- Bingham, N. H., C. M. Goldie, and J. L. Teugels (1987). Regular variation. Vol. 27. Encyclopedia of Mathematics and its Applications. Cambridge University Press, pp.  $xx+494$ . DOI: [10.](http://dx.doi.org/10.1017/CBO9780511721434) [1017/CBO9780511721434](http://dx.doi.org/10.1017/CBO9780511721434).
- Borovkov, A. A. (2020). Asymptotic Analysis of Random Walks: Light-Tailed Distributions. Encyclopedia of Mathematics and its Applications. Cambridge University Press, pp. v–viii. doi: [10.1017/9781139871303](http://dx.doi.org/10.1017/9781139871303).
- Chakraborty, B. (2001). "On Affine Equivariant Multivariate Quantiles". In: Annals of the *Institute of Statistical Mathematics* 53.2, pp. 380–403. DOI: [10.1023/A:1012478908041](http://dx.doi.org/10.1023/A:1012478908041).
- Chaudhuri, P. (1996). "On a Geometric Notion of Quantiles for Multivariate Data". In: Journal of the American Statistical Association 91.434, pp. 862–872. doi: [10.1080/01621459.1996.](http://dx.doi.org/10.1080/01621459.1996.10476954) [10476954](http://dx.doi.org/10.1080/01621459.1996.10476954).
- Cline, D.B.H. and S. I. Resnick (1992). "Multivariate subexponential distributions". In: Stochastic Processes and their Applications 42.1, pp. 49–72. DOI: [https://doi.org/10.1016/](http://dx.doi.org/https://doi.org/10.1016/0304-4149(92)90026-M) [0304-4149\(92\)90026-M](http://dx.doi.org/https://doi.org/10.1016/0304-4149(92)90026-M).
- Csörgö, S., E. Haeusler, and D. M. Mason (1988). "A probabilistic approach to the asymptotic distribution of sums of independent, identically distributed random variables". In: Advances in Applied Mathematics 9.3, pp. 259–333. DOI:  $10.1016/0196-8858(88)90016-4$ .
- Cuberos, A., E. Masiello, and V. Maume-Deschamps (2015). "High level quantile estimations of sums of risks". In: *Dependence Modeling* 3, pp. 141–158. DOI: [10.1515/demo-2015-0010](http://dx.doi.org/10.1515/demo-2015-0010).
- Dacorogna, M., L. Elbahtouri, and M. Kratz (2018). "Model Validation for Aggregated Risks". In: Annals of Actuarial Science 12.2, pp. 433–454. DOI: [10.1017/S1748499517000227](http://dx.doi.org/10.1017/S1748499517000227).
- Das, B. and M. Kratz (2020). "Risk concentration under second order regular variation". In: *Extremes* 23, pp. 381–410. DOI: [10.1007/s10687-020-00382-3](http://dx.doi.org/10.1007/s10687-020-00382-3).
- David, H. A. and H. N. Nagaraja (2004). Order Statistics. Wiley Series in Probability and Statistics. Wiley. DOI: [10.1002/0471667196.ess6023](http://dx.doi.org/10.1002/0471667196.ess6023).
- Dhar, S., B. Chakraborty, and P. Chaudhuri (2014). "Comparison of multivariate distributions using quantile-quantile plots and related tests". In:  $Bernoulli$  20.3, pp. 1484–1506. DOI: [10.3150/13-BEJ530](http://dx.doi.org/10.3150/13-BEJ530).
- Dobrushin, R. L. (1970). "Prescribing a System of Random Variables by Conditional Distri-butions". In: Theory of Probability and Its Applications 15.3, pp. 458–486. DOI: [10.1137/](http://dx.doi.org/10.1137/1115049) [1115049](http://dx.doi.org/10.1137/1115049).
- Embrechts, P., C. Klüppelberg, and T. Mikosch (1997). Modelling Extreme Events for Insurance and Finance. Stochastic Modelling and Applied Probability. Springer-Verlag Berlin Heidelberg, pp. XV, 648. doi: [10.1007/978-3-642-33483-2](http://dx.doi.org/10.1007/978-3-642-33483-2).
- Falk, M. and F. Marohn (1993). "Von Mises Conditions Revisited". In: Annals of Probability 21.3, pp. 1310 -1328. DOI: [10.1214/aop/1176989120](http://dx.doi.org/10.1214/aop/1176989120).
- Foss, S., D. Korshunov, and S. Zachary (2013). An Introduction to Heavy-Tailed and Subexponential Distributions. Springer Series in Operations Research and Financial Engineering. Springer-Verlag New York, pp. XI, 157. DOI: [10.1007/978-1-4614-7101-1](http://dx.doi.org/10.1007/978-1-4614-7101-1).
- Geluk, J. L. and L. de Haan (1987). Regular Variation, Extensions and Tauberian Theorems. Vol. 40. CWI Tract. Amsterdam: Centrum voor Wiskunde en Informatica, pp. iv+132. isbn: 90-6196-324-9.
- Girard, S. and G. Stupfler (2015). "Extreme geometric quantiles in a multivariate regular variation framework". In: *Extremes* 18.4, pp. 629–663. DOI: [10.1007/s10687-015-0226-0](http://dx.doi.org/10.1007/s10687-015-0226-0).
- (2017). "Intriguing properties of extreme geometric quantiles". In: Revstat Statistical Journal 15.1, pp. 107–139.
- de Haan, L. (1970). On regular variation and its applications to the weak convergence of sample extremes. Amsterdam: Mathematical Centre Tracts Vol. 32.
- de Haan, L. and A. Ferreira (2006). Extreme Value Theory: An Introduction. Springer Series in Operations Research and Financial Engineering. New York: Springer, p. XVIII 418. doi: [10.1007/0-387-34471-3](http://dx.doi.org/10.1007/0-387-34471-3).
- de Haan, L. and S. I. Resnick (1993). "Estimating the limit distribution of multivariate extremes". In: *Communications in Statistics. Stochastic Models* 9.2, pp. 275–309. DOI: [10.](http://dx.doi.org/10.1080/15326349308807267) [1080/15326349308807267](http://dx.doi.org/10.1080/15326349308807267).
- (1996). "Second-Order Regular Variation and Rates of Convergence in Extreme-Value The-ory". In: Annals of Probability 24.1, pp. 97–124. DOI: [10.1214/aop/1042644709](http://dx.doi.org/10.1214/aop/1042644709).
- de Haan, L. and U. Stadtm $\widehat{A}$ ijller (1996). "Generalized regular variation of second order". In: Journal of the Australian Mathematical Society. Series A. Pure Mathematics and Statistics 61.3, pp. 381–395. doi: [10.1017/S144678870000046X](http://dx.doi.org/10.1017/S144678870000046X).
- Hahn, M. G. (1991). Sums, Trimmed Sums and Extremes. Ed. by M. G. Hahn, D. M. Mason, and D. C. Weiner. Vol. 23. Progress in Probability. Birkhäuser Boston. doi: [10.1007/978-](http://dx.doi.org/10.1007/978-1-4684-6793-2) [1-4684-6793-2](http://dx.doi.org/10.1007/978-1-4684-6793-2).
- Hua, L. and H. Joe (2011). "Second order regular variation and conditional tail expectation of multiple risks". In: *Insurance Mathematics and Economics* 49.3, pp. 537–546. DOI: [10.](http://dx.doi.org/10.1016/j.insmatheco.2011.08.013) [1016/j.insmatheco.2011.08.013](http://dx.doi.org/10.1016/j.insmatheco.2011.08.013).
- Karamata, J. (1933). "Sur un mode de croissance régulière. Théorèmes fondamentaux". In: Bulletin de la Société Mathématique de France 61, pp.  $55-62$ . DOI: [10.24033/bsmf.1196](http://dx.doi.org/10.24033/bsmf.1196).
- Koltchinskii, V. I. (1997). "M-estimation, convexity and quantiles". In: Annals of Statistics 25.2, pp. 435 –477. doi: [10.1214/aos/1031833659](http://dx.doi.org/10.1214/aos/1031833659).
- Kratz, M. (2014). "Normex, a new method for evaluating the distribution of aggregated heavy tailed risks. Application to risk measures". In: *Extremes* 17.4, pp. 661–691. DOI: [10.1007/](http://dx.doi.org/10.1007/s10687-014-0197-6) [s10687-014-0197-6](http://dx.doi.org/10.1007/s10687-014-0197-6).
- Lehtomaa, J. (2017). "Large deviations of means of heavy-tailed random variables with finite moments of all orders". In: *Journal of Applied Probability* 54.1, pp.  $66-81$ . DOI: [10.1017/](http://dx.doi.org/10.1017/jpr.2016.87) [jpr.2016.87](http://dx.doi.org/10.1017/jpr.2016.87).
- Lv, W., T. Mao, and T. Hu (2012). "Properties of second-order regular variation and expansions for risk concentration". In: *Probability in the Engineering and Informational Sciences* 26.4, pp. 535–559. doi: [10.1017/S0269964812000174](http://dx.doi.org/10.1017/S0269964812000174).
- Mainik, G. and P. Embrechts (2013). "Diversification in heavy-tailed portfolios: Properties and Pitfalls". In: *Annals of Actuarial Science* 7.1, pp. 26–45. DOI: [10.1017/S1748499512000280](http://dx.doi.org/10.1017/S1748499512000280).
- Mikosch, T. and A. Nagaev (1998). "Large Deviations of Heavy-Tailed Sums with Applications in Insurance". In: *Extremes* 1, pp. 81–110. DOI:  $10.1023/A:1009913901219$ .
- Mori, T. (1984). "On the limit distributions of lightly trimmed sums". In: *Mathematical Proceed*ings of the Cambridge Philosophical Society  $96.3$ , pp. 507–516. DOI: [10.1017/S0305004100062447](http://dx.doi.org/10.1017/S0305004100062447).
- Müller, U. K. (2019). "Refining the central limit theorem approximation via extreme value theory". In: *Statistics and Probability Letters* 155, pp. 1–7. DOI: [10.1016/j.spl.2019.](http://dx.doi.org/10.1016/j.spl.2019.108564) [108564](http://dx.doi.org/10.1016/j.spl.2019.108564).
- Nocedal, J. and S. Wright (2006). Numerical optimization. Springer Series in Operations Research and Financial Engineering. Springer, pp. XXII, 664. DOI: 10.1007/978-0-387-[40065-5](http://dx.doi.org/10.1007/978-0-387-40065-5).
- Omey, E. (2006). "Subexponential Distribution Functions in  $\mathbb{R}^{d}$ ". In: *Journal of Mathematical*  $Sciences$  138.1, pp. 5434-5449. DOI: [10.1007/s10958-006-0310-8](http://dx.doi.org/10.1007/s10958-006-0310-8).
- Petrov, V. V. (1975). Sums of Independent Random Variables. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, pp. X, 348. doi: [10.1007/978-3-642-65809-9](http://dx.doi.org/10.1007/978-3-642-65809-9).
- Petrov, V. V. (1995). Limit Theorem of Probability Theory: Sequences of Independent Random Variables. Vol. 4. Oxford Studies in Probability. Oxford Sciences Publications. ISBN: 0-19-853499-X.
- Resnick, S. I. (1987). Extreme Values, Regular Variation and Point Processes. Springer Series in Operations Research and Financial Engineering. New York: Springer, pp.  $xii+320$ . DOI: [10.1007/978-0-387-75953-1](http://dx.doi.org/10.1007/978-0-387-75953-1).
- (2002). "Hidden Regular Variation, Second Order Regular Variation and Asymptotic Independence". In: *Extremes* 5.4, pp. 303–336. DOI: [10.1023/A:1025148622954](http://dx.doi.org/10.1023/A:1025148622954).
- (2007). Heavy Tail Phenomena: Probabilistic and Statistical Modeling. Springer Series in Operations Research and Financial Engineering. New York: Springer, pp. xx+404. doi: [10.1007/978-0-387-45024-7](http://dx.doi.org/10.1007/978-0-387-45024-7).
- Samorodnitsky, G. and J. Sun (2016). "Multivariate subexponential distributions and their applications". In: *Extremes* 19, pp. 171–196. DOI: [10.1007/s10687-016-0242-8](http://dx.doi.org/10.1007/s10687-016-0242-8).
- Samorodnitsky, G. and M. S. Taqqu (1994). Stable non-Gaussian Random Processes: Stochastic Models with Infinite Variance. Chapman and Hall. DOI: [10.1201/9780203738818](http://dx.doi.org/10.1201/9780203738818).
- Seneta, E. (1976). Regularly Varying Functions. Vol. 508. Lecture Notes Math. Springer-Verlag Berlin Heidelberg. DOI: [10.1007/BFb0079658](http://dx.doi.org/10.1007/BFb0079658).
- Zaliapin, I. V., Y. Y. Kagan, and F. P. Schoenberg (2005). "Approximating the Distribution of Pareto Sums". In: *Pure and Applied Geophysics* 162, pp. 1187–1228. DOI: [10.1007/s00024-](http://dx.doi.org/10.1007/s00024-004-2666-3) [004-2666-3](http://dx.doi.org/10.1007/s00024-004-2666-3).

# A Appendices

### A.1 Regular Variation and related notions

Let us recall the definitions and properties of regular variation in both univariate and multivariate cases (see Bingham et al. (1987), Geluk and de Haan (1987), de Haan (1970), de Haan and Ferreira (2006), Resnick (2002), Resnick (1987), and Seneta (1976)), and point out some useful relations between each other.

#### Univariate distributions -

**Definition A.1** (Regular variation in one dimension; see Bingham et al. (1987)). A rv X with cdf F has a regularly varying (right) tail with index  $\alpha > 0$  if  $\overline{F} = 1 - F \in RV_{-\alpha}$  (or, by abuse of notation,  $X \in \mathcal{RV}_{-\alpha}$ , i.e.

$$
\lim_{t \to \infty} \frac{F(tx)}{\overline{F}(t)} = x^{-\alpha}, \quad x > 0.
$$

Alternatively, we say that X has a regularly varying tail if there exists a function  $b : \mathbb{R}_+ \to \mathbb{R}_+$ with  $b(t) \uparrow \infty$  as  $t \to \infty$  such that

$$
\lim_{t \to \infty} t \, \mathbb{P} \left( \, X > b(t)x \, \right) = x^{-\alpha}.
$$

We write  $\overline{F} \in \mathcal{RV}_{-\alpha}(b)$  or, by abuse of notation,  $X \in \mathcal{RV}_{-\alpha}(b)$ .

As a consequence,  $b(\cdot) \in \mathcal{RV}_{1/\alpha}$  and a natural choice is  $b(t) = (1/\overline{F})^{\leftarrow}(t)$ .

Definition A.2 (Extended Regular variation; see e.g. de Haan and Ferreira (2006)). A measurable function f is of extended regular variation, denoted by  $f \in ERV(\gamma)$ , if for some  $\gamma \in \mathbb{R}$ and positive function a,

$$
\lim_{t \to \infty} \frac{f(tx) - f(t)}{a(t)} = \frac{x^{\gamma} - 1}{\gamma}
$$

for all  $x > 0$ .

When looking at the speed of convergence to the limits given in the definition of  $(E)RV$ , comes the notion of "second-order (extended) regular variation".

**Definition A.3** (Second order regular variation; see e.g. de Haan and Resnick (1993)). A rv X with cdf F such that  $F \in RV_{-\alpha}(b)$  with  $\alpha > 0$ , possesses second order regular variation with parameter  $\rho \leq 0$ , denoted by  $X \in 2\mathcal{RV}(-\alpha,\rho)$ , if there exists a function  $A(t) \rightarrow 0$  that is ultimately of constant sign,  $|A(t)| \in \mathcal{RV}_\rho$  with  $\rho \leq 0$  and  $c \neq 0$  such that

$$
\frac{t\bar{F}(b(t)x) - x^{-\alpha}}{A(b(t))} \xrightarrow{t \to \infty} cx^{-\alpha} \frac{x^{\rho} - 1}{\rho}, \quad x > 0
$$
\n(1.28)

The right hand side of Eq. (1.28) is interpreted as the function  $cx^{-\alpha} \log(x)$  when  $\rho = 0$ .

Definition A.4 (Second order extended regular variation; see e.g. de Haan and StadtmÄijller (1996)). A measurable function f is said to be of second-order extended regular variation (with second-order parameter  $\rho \leq 0$ , denoted by  $f \in 2ERV(\gamma, \rho)$ , if there exist positive functions a and A with  $\lim_{t\to\infty} A(t) = 0$  such that

$$
\frac{\frac{f(tx)-f(t)}{a(t)} - \frac{x^{\gamma}-1}{\gamma}}{A(t)} \underset{t \to \infty}{\longrightarrow} H(x)
$$

for all  $x > 0$ , where the limit function H is not a multiple of  $(x^{\gamma} - 1)/\gamma$ .

An equivalent representation of 2RV functions and "potter bounds"

It appears that a  $2\mathcal{RV}$  function has a specific form.

**Lemma A.1** (Lemma 3 in Hua and Joe (2011)). Suppose  $g \in 2\mathcal{RV}_{\alpha,\rho}$ ,  $\alpha \in \mathbb{R}, \rho < 0$ , then we can write  $g = kt^{\alpha}(1 + l(t)), t > 0$  with some constant  $k \neq 0$  and regularly varying function  $l(t) \in \text{RV}_{\rho}$ .

In other words, Lemma A.1 says that a slowly varying function  $L(\cdot)$  can be a  $2\mathcal{RV}_{0,\rho}$  function with parameters  $0, \rho < 0$ , only in the case  $\lim_{t\to\infty} L(t) \neq 0, \pm\infty$ . In the case  $\rho = 0$ , the function L belongs to so called Π-class and has the representation (see theorem B.2.12 in de Haan and Ferreira (2006))

$$
L(t) = c_1 + c_2 a_1(t) + \int_1^t \frac{a_2(s)}{s} ds,
$$

where  $c_1, c_2 \in \mathbb{R}$  are constants and  $a_1(\cdot), a_2(\cdot)$  are slowly varying functions such that  $a_1(t) \underset{t\to\infty}{\sim} a_2(t)$ . The following lemma will be useful to prove Proposition 4.2.

**Lemma A.2** (Proposition 4 in Hua and Joe (2011)). Suppose  $g \in 2RV_{\alpha,\rho}$  with  $\alpha \in \mathbb{R}$  and  $\rho < 0$ , then for any  $\epsilon, \delta > 0$ , there exists  $t_0 = t_0(\epsilon, \delta)$  such that for all  $t, tx \geq t_0$  and  $x > 0$ 

$$
\left| \frac{g(tx)/g(t) - x^{\alpha}}{a(t)} - x^{\alpha} \frac{x^{\rho} - 1}{\rho} \right| \leq \epsilon \max \left( x^{\alpha + \rho + \delta}, x^{\alpha + \rho - \delta} \right), \tag{1.29}
$$

where  $a(t) = -\rho[\ell(t)]/(1 + \ell(t))$  with  $g(t) = kt^{\alpha}(1 + \ell(t)), 0 < k < +\infty$  and  $l(\cdot) \in \text{RV}_{\rho}$ .

In fact, Hua and Joe use Lemmas A.1 and A.2 for  $g(t) = \overline{F}(t)$ , and, for this reason, proved the lemmas in the case  $\alpha < 0$ . But their proof can be repeated line by line for an arbitrary  $\alpha \in \mathbb{R}$ . We believe that Lemma A.2 should also hold true when  $\rho = 0$ , but we will not develop further on this question.

Finally, let us end this paragraph by representing the relations between the various notions of (related) regular variation in Figure 9.



Figure 9: Relations between RV related notions, where F is the cdf of a real rv X and the function g is defined by  $g(x) := (-1/\log F(x))$ <sup>←</sup>

Multivariate Distributions - Multivariate regular variation appears as a natural extension of Definition A.1, when studying jointly heavy-tailed random variables. The notion of vague convergence of measures is used for convergence of measures on the non-negative Euclidean orthant  $\mathbb{R}^d_+$  and its subsets; see Resnick (2007) for further details.

**Definition A.5** (Multivariate regular variation, Resnick (2007)). Suppose  $\mathbf{X} = (X_1, \ldots, X_d)$ is a random vector in  $[0,\infty)^d$ . Then **X** is multivariate regularly varying if there exist  $b(t) \uparrow \infty$ with  $b(\cdot) \in \mathcal{RV}_{1/\alpha}$ ,  $\alpha > 0$ , and a Radon measure  $\nu \neq 0$  such that, as  $t \to \infty$ ,

$$
t\,\mathbb{P}\left(\frac{\boldsymbol{X}}{b(t)}\in\ \cdot\ \right)\ \stackrel{v}{\rightarrow}\ \nu(\cdot)\quad on\quad \mathbb{M}_+(E),
$$

where  $\stackrel{v}{\rightarrow}$  denotes vague convergence of measures on  $\mathbb{M}_+(E)$ , the class of Radon measures on Borel subsets of  $E := [0, \infty]^d \setminus \{0\}$ . We write  $\mathbf{X} \in \mathcal{MRV}_{-\alpha}(b, \nu)$ .

The measure  $\nu(\cdot)$  has a scaling property for relatively compact  $A \subset E$ , given by

$$
\nu(kA) = k^{-\alpha} \nu(A), \quad k > 0.
$$

As already mentioned in the paper, an equivalent definition is through the pseudo-polar representation:  $\mathbf{X} \in \mathcal{MRV}_{-\alpha}$  if there exists a random vector  $\Theta$  with values in the unit sphere  $\mathcal{S}_{d-1}$ in  $\mathbb{R}^d$  w.r.t. the norm  $\|\cdot\|$ , such that, for all  $t > 0$ ,

$$
\frac{\mathbb{P}(\|\mathbf{X}\| > t u, \mathbf{X}/\|\mathbf{X}\| \in \cdot)}{\mathbb{P}(\|\mathbf{X}\| > u)} \xrightarrow{v} t^{-\alpha} \mathbb{P}(\Theta \in \cdot) \quad \text{as } u \to \infty.
$$

Note that a characterization of MRV for X is that any linear combination  $\langle t, X \rangle, t \in \mathbb{R}^d$ , is regularly varying,  $\langle \cdot, \cdot \rangle$  denoting the usual inner product in  $\mathbb{R}^d$ , namely (see Basrak et al.

### $(2002)$ : There exist  $\alpha > 0$  and a slowly varying function L such that, for all x,

$$
\lim_{u\to\infty}\frac{\mathbb{P}\left(\langle \boldsymbol{x}, \boldsymbol{X}\rangle\geqslant u\right)}{u^{-\alpha}L(u)}=w(\boldsymbol{x})\text{ exists and }\exists \boldsymbol{x}_0\neq \boldsymbol{0} \text{ s.t. } w(\boldsymbol{x}_0)>0.
$$

As for the dimension 1, we can define the notion of second order multivariate regular variation, namely:

Definition A.6 (Second order multivariate regular variation; see Resnick (2002)). Suppose  $\mathbf{X} \in \mathcal{MRV}_{-\alpha}(b, \nu)$  and there exists  $A(t) \to 0$  that is ultimately of constant sign with  $|A(\cdot)| \in$  $\mathcal{RV}_\rho, \rho \leqslant 0$ , such that

$$
\frac{t \mathbb{P}\left(\frac{\mathbf{X}}{b(t)} \in [\mathbf{0}, \mathbf{x}]^c\right) - \nu([\mathbf{0}, \mathbf{x}]^c)}{A(b(t))} \underset{t \to \infty}{\to} H(\mathbf{x})
$$
\n(1.30)

locally uniformly in  $x \in (0,\infty]^d \setminus \{\infty\}$ , where H is a function that is non-zero and finite. Then X is second order regularly varying with parameters  $\alpha > 0$  and  $\rho \leq 0$ . We write  $X \in$  $2MRV_{-\alpha,\rho}(b, A, \nu, H).$ 

### A.2 Proofs of Lemmas of Section 2

We can prove Lemma 2.1 and Lemma 2.2 using pdf, when existing, or characteristic functions. We proceed for instance via pdf for the proof of the first lemma, as it is a very standard way (see e.g. David and Nagaraja (2004)), and via characteristic function for the second one to show that it can be done in full generality.

Starting with Lemma 2.1, let us prove the following result.

Let  $f_{\mathbf{X}_{(1)},\cdots,\mathbf{X}_{(n-1)}\|\|\mathbf{X}_{(n)}\|}(\mathbf{x}_1,\cdots,\mathbf{x}_{n-1}\|y)$  be the conditional pdf of  $(\mathbf{X}_{(1)},\ldots,\mathbf{X}_{(n-1)})$  given the event  $(\|\mathbf{X}_{(n)}\| = y)$ . For any  $\mathbf{x}_1, \ldots, \mathbf{x}_{n-1} \in \mathbb{R}^d$ ,  $y \geqslant 0$ , such that  $\|\mathbf{x}_1\| \leqslant \ldots \leqslant \|\mathbf{x}_{n-1}\| \leqslant y$ , we have

$$
f_{\mathbf{X}_{(1)},\cdots,\mathbf{X}_{(n-1)}\|\|\mathbf{X}_{(n)}\|}(\mathbf{x}_1,\ldots,\mathbf{x}_{n-1}\|y)=(n-1)!\prod_{i=1}^{n-1}f_{\mathbf{X}\|\|\mathbf{X}\|}(\mathbf{x}_i\|y),
$$
\n(1.31)

where  $f_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot | y)$  is defined in (2.4).

*Proof*. Let  $[y, y + \delta y]$  be a 'small' interval from  $\mathbb{R}^+$  and, for  $i = 1, ..., n - 1$ , let  $[\mathbf{x}_i, \mathbf{x}_i + \delta \mathbf{x}_i]$ denote a 'small' cube in  $\mathbb{R}^d$  with initial vertex  $\mathbf{x}_i$  and measure  $\delta \mathbf{x}_i$ . As  $\mathbf{X}_{(1)}, \ldots, \mathbf{X}_{(n)}$  are the ordered (by norm) statistics (see (1.1)), there are n! permutations of the vector  $(\mathbf{X}_1, \ldots, \mathbf{X}_n)$ to be  $(X_{(1)},...,X_{(n)})$ , hence we can write

$$
\mathbb{P}\left(\mathbf{X}_{(1)} \in [\mathbf{x}_1, \mathbf{x}_1 + \delta \mathbf{x}_1), \cdots, \mathbf{X}_{(n-1)} \in [\mathbf{x}_{n-1}, \mathbf{x}_{n-1} + \delta \mathbf{x}_{n-1}), ||\mathbf{X}_{(n)}|| \in [y, y + \delta y)\right)
$$
  
= n! 
$$
\prod_{i=1}^{n-1} \left(f(\mathbf{x}_i)\delta \mathbf{x}_i\right) f_{\|\mathbf{X}\|}(y)\delta y + O((\delta y)^2).
$$

Also, as  $\|\mathbf{X}_{(n)}\|$  is the maximum of the iid rv  $\|\mathbf{X}_{1}\|, \ldots, \|\mathbf{X}_{n}\|$ , we have

$$
f_{\|\mathbf{X}_{(n)}\|}(y) = n F_{\|\mathbf{X}\|}^{n-1}(y) f_{\|\mathbf{X}\|}(y)
$$
\n(1.32)

and

$$
\mathbb{P}\left(\left\|\mathbf{X}_{(n)}\right\| \in [y, y + \delta y)\right) = n F_{\|\mathbf{X}\|}^{n-1}(y) f_{\|\mathbf{X}\|}(y) \delta y + O\left((\delta y)^2\right).
$$

Then

$$
f_{\mathbf{X}_{(1)},\cdots,\mathbf{X}_{(n-1)}\|\|\mathbf{X}_{(n)}\|}(\mathbf{x}_1,\ldots,\mathbf{x}_{n-1}\|y)=(n-1)!\prod_{i=1}^{n-1}\frac{f(\mathbf{x}_i)}{F_{\|\mathbf{X}\|}(y)},
$$

hence the result  $(1.31)$ .

Turning to Lemma 2.2, it can be formalized as follows, using characteristic functions: For any vectors  $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^d$  and any real  $y > 0$ , we have

$$
\mathbb{E}\left[e^{\sum_{k=1}^{n}i\left\langle\mathbf{x}_{k},\mathbf{X}_{(k)}\right\rangle}\middle|\left\|\mathbf{X}_{(n)}\right\|=y\right]=\mathbb{E}\left[e^{\sum_{k=1}^{n-1}i\left\langle\mathbf{x}_{k},\mathbf{Y}_{(k)}\right\rangle}\right]\mathbb{E}\left[e^{i\left\langle\mathbf{x}_{n},\mathbf{X}_{n}\right\rangle}\middle|\left\|\mathbf{X}_{n}\right\|=y\right].
$$
 (1.33)

*Proof.* To simplify the notations, we denote by  $R_1, \dots, R_n$  the norms  $\|\mathbf{X}_1\|, \dots, \|\mathbf{X}_n\|$ . Since  $(R_1, \dots, R_n)$  are iid, for all  $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{R}^d, y > 0$ , we have

$$
\mathbb{E}\left[e^{\sum_{k=1}^{n}i\left\langle\mathbf{x}_{k},\mathbf{X}_{(k)}\right\rangle}\middle|R_{(n)}=y\right]=n!\,\mathbb{E}\left[e^{\sum_{k=1}^{n}i\left\langle\mathbf{x}_{k},\mathbf{X}_{k}\right\rangle}\mathbb{1}_{\{R_{1}\leqslant\cdots\leqslant R_{n-1}\leqslant R_{n}\}}\middle|R_{(n)}=y\right]\right]
$$
\n
$$
=(n-1)!\,\mathbb{E}\left[\left(e^{\sum_{k=1}^{n-1}i\left\langle\mathbf{x}_{k},\mathbf{X}_{k}\right\rangle}\mathbb{1}_{\{R_{1}\leqslant\cdots\leqslant R_{n-1}\leqslant y\}}\right)\left(ne^{i\left\langle\mathbf{x}_{n},\mathbf{X}_{n}\right\rangle}\mathbb{1}_{\{R_{n}=y\}}\right)\middle|R_{(n)}=y\right].
$$

The random variables in the two big brackets being independent, since one depends on  $X_1, \dots, X_{n-1}$ and the other on  $\mathbf{X}_n$ , the last expression equals

$$
\mathbb{E}\left[(n-1)!\,e^{\sum_{k=1}^{n-1}i\langle\mathbf{x}_k,\mathbf{X}_k\rangle}\mathbb{1}_{\{R_1\leqslant\cdots\leqslant R_{n-1}\leqslant y\}}\Big|R_{(n)}=y\right]\mathbb{E}\left[n e^{i\langle\mathbf{x}_n,\mathbf{X}_n\rangle}\mathbb{1}_{\{R_n=y\}}\Big|R_{(n)}=y\right].\tag{1.34}
$$

Note that, by Lemma 2.1, the first expectation in (1.34) equals to  $\mathbb{E}\left[e^{\sum_{k=1}^{n-1} i\langle\mathbf{x}_k,\mathbf{Y}_{(k)}\rangle}\right]$ . Let us calculate the second expectation in (1.34). Considering the  $\sigma$ -algebra  $\sigma(R)$  generated by the iid rv  $(R_1, \dots, R_n)$  (with parent rv R), we can write, for any rv  $\xi$ ,  $\mathbb{E}\left[\xi \mid R_{(n)}\right] = \mathbb{E}\left[\mathbb{E}\left[\xi \mid \sigma(R)\right]\right]$ <br>  $\mathbb{E}\left[\mathbb{E}\left[\tau \mid R_{(n)}\right]\right]$  $R_{(n)}$  and  $\mathbb{E}\left[e^{i\langle \mathbf{x}_n,\mathbf{X}_n\rangle}\right]$  $\sigma(R)$ ] =  $\mathbb{E}\left[e^{i\langle \mathbf{x}_n, \mathbf{X}_n\rangle}\right]$  $R_n$  = :  $h_{\mathbf{x}_n}(R_n)$ . We deduce that, for all  $y > 0$ ,

$$
\mathbb{E}\left[n\mathbb{1}_{\{R_n=y\}}e^{i\langle\mathbf{x}_n,\mathbf{X}_n\rangle}\middle|R_{(n)}=y\right] = \mathbb{E}\left[n\mathbb{1}_{\{R_n=y\}}h_{\mathbf{x}_n}(R_n)\middle|R_{(n)}=y\right]
$$

$$
= h_{\mathbf{x}_n}(y)\mathbb{E}\left[n\mathbb{1}_{\{R_n=R_{(n)}\}}\middle|R_{(n)}=y\right] = h_{\mathbf{x}_n}(y).
$$

 $\Box$ 

Combining the results obtained for the two expectations in (1.34) provides (1.33).

### A.3 Proof of Theorem 3.1

PRELIMINARY RESULT - To prove Theorem 3.1, we need to show that the inverse matrix of the covariance matrix  $\Sigma(y)$  of the truncated random vector **Y** is bounded, namely:

**Lemma A.3.** Under the settings of Theorem 3.1, for some  $C > 0$ , there exists  $\delta > 0$  such that  $\mathbb{P}(\|\mathbf{X}\| \leqslant \delta) < 1$  and

$$
\|\Sigma(y)^{-1/2}\| \leqslant C \quad \text{for all } y \geqslant \delta.
$$

*Proof.* From the definition of the truncated distribution  $F_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot |y)$  in (2.3) and continuity of  $F_{\parallel {\bf X} \parallel}(\cdot)$ , one can show that, for any sequence  $(y_n)$  converging to  $y_0 \in \mathbb{R}^+$ , with  $\mathbb{P}(\parallel {\bf X} \parallel \leq y_0)$ 0, we have

$$
\Sigma(y_n) \to \Sigma(y_0)
$$
 as  $n \to \infty$ ,

where, for convenience, we suppose  $\Sigma(\infty) := \Sigma$ , covariance matrix of the initial random vector **X** with c.d.f. F. Condition (C2) in Theorem 3.1 and the definition of  $y_0$  imply that  $\Sigma(y_0)$  is positive definite. Consequently, the square root of its inverse matrix exists, with finite norm:

$$
\|\Sigma(y_0)^{-1/2}\| < \infty. \tag{1.35}
$$

Now, assume that the statement of Lemma A.3 is false. Then, for all  $C > 0$  and  $\delta > 0$  such that  $\mathbb{P}(\|\mathbf{X}\| \leq \delta) < 1$ , there exists  $y \geq \delta$  such that  $\|\Sigma(y)^{-1/2}\| > C$ . Therefore, there exists a sequence  $(y_n)_n \geq \delta$  such that

$$
\|\Sigma(y_n)^{-1/2}\| \to \infty \text{ as } n \to \infty.
$$
\n(1.36)

But we can choose a subsequence  $(y_{n_k})_{k\geqslant 1}$  such that, as  $k\to\infty$ , either (a)  $y_{n_k}\to\infty$ , or (b)  $y_{n_k} \to y_0 \geq \delta$  for some point  $y_0 \geq \delta$ . In both cases, (1.36) contradicts (1.35). We conclude that the statement of Lemma A.3 is true. □

PROOF OF THEOREM 3.1 - Let C denote a positive constant that may vary from line to line, all along the proof. Let  $Y_1, \ldots, Y_{n-1}$  be i.i.d. random vectors, with parent random vector **Y**, having the truncated distribution  $F_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot |y)$  defined in (2.3). We have, using conditional probabilities and applying Lemma 2.1,

$$
\mathbb{P}\left(\mathbf{S}_{n} \in \mathbf{B}\right) = \mathbb{P}\left(\mathbf{X}_{(1)} + \cdots + \mathbf{X}_{(n)} \in \mathbf{B}\right)
$$
  
=  $\int_{\mathbb{R}^{d}} f_{(n)}(\mathbf{x}) \mathbb{P}\left(\mathbf{X}_{(1)} + \cdots + \mathbf{X}_{(n-1)} \in \mathbf{B} - \mathbf{x} \mid \mathbf{X}_{(n)} = \mathbf{x}\right) d\mathbf{x}$   
=  $\int_{\mathbb{R}^{d}} f_{(n)}(\mathbf{x}) \mathbb{P}\left(\mathbf{T}_{n-1} \in \mathbf{B} - \mathbf{x}\right) d\mathbf{x}$ , where  $T_{n-1} := \sum_{k=1}^{n-1} \mathbf{Y}_{k}$ .

Using Definition 3.1 of the d-Normex distribution, we can write, for any  $\mathbf{B} \in \mathcal{C}$ ,

$$
|\mathbb{P}\left(\mathbf{S}_n \in \mathbf{B}\right) - G_n(\mathbf{B})| = \int_{\mathbb{R}^d} f_{(n)}(\mathbf{x}) |\mathbb{P}\left(\mathbf{T}_{n-1} \in \mathbf{B} - \mathbf{x}\right) - \Phi_{(n-1)\mu(\|\mathbf{x}\|), (n-1)\Sigma(\|\mathbf{x}\|)}(\mathbf{B} - \mathbf{x}) |\,\mathrm{d}\mathbf{x}.\tag{1.37}
$$

Further, under (C2), the pdf of the rv  $\|\mathbf{X}\|$  is regularly varying,  $f_{\|X\|}(\cdot) \in \mathcal{RV}_{-\alpha-1}$ , *i.e.* (see e.g., Bingham et al. Bingham et al. (1987)) there exists a slowly varying function  $L(\cdot)$  such that

$$
f_{\|X\|}(y) = L(y)y^{-\alpha - 1}.
$$
\n(1.38)

Let us choose  $\delta > 0$  that satisfies Lemma A.3 and such that the function of y defined by sup  $L(t)$  is slowly varying (see e.g. ex.4 p.58 in Bingham et al. (1987)).  $t\in[\delta,y]$ 

Splitting the integration domain of the integral of (1.37) into two disjoint sets  $\{x : ||x|| < \delta\}$ and  $\{x : ||x|| \ge \delta\}$ , we have, on one hand,

$$
\int_{\|\mathbf{x}\| < \delta} f_{(n)}(\mathbf{x}) \left| \mathbb{P}\left(\mathbf{T}_{n-1} \in \mathbf{B} - \mathbf{x}\right) - \Phi_{(n-1)\mu(\|\mathbf{x}\|), (n-1)\Sigma(\|\mathbf{x}\|)} (\mathbf{B} - \mathbf{x}) \right| d\mathbf{x} \n\leqslant \int_{\|\mathbf{x}\| < \delta} f_{(n)}(\mathbf{x}) d\mathbf{x} = \mathbb{P}\left(\|\mathbf{X}_{(n)}\| < \delta\right) = \left(\mathbb{P}\left(\|\mathbf{X}\| < \delta\right)\right)^n.
$$
\n(1.39)

On the other hand, since  $Y_1, \ldots, Y_{n-1}$  are i.i.d bounded random vectors with a non degenerate distribution (via  $(C2)$ ), we can use the non-generalized Berry-Esseen inequality recalled in Proposition 2.1, and obtain

$$
\int_{\|\mathbf{x}\| \geqslant \delta} f_{(n)}(\mathbf{x}) \left| \mathbb{P} \left( \mathbf{T}_{n-1} \in \mathbf{B} - \mathbf{x} \right) - \Phi_{(n-1)\mu(\|\mathbf{x}\|), (n-1)\Sigma(\|\mathbf{x}\|)} (\mathbf{B} - \mathbf{x}) \right| d\mathbf{x} \n\leqslant \frac{C}{\sqrt{n}} \int_{\|\mathbf{x}\| \geqslant \delta} f_{(n)}(\mathbf{x}) \mathbb{E} \left( \left\| (\Sigma(\|\mathbf{x}\|))^{-1/2} \left( \mathbf{Y} - \mu(\|\mathbf{x}\|) \right) \right\|^3 \right) d\mathbf{x},
$$
\n(1.40)

Y denoting the parent random vector of  $(Y_1, \ldots, Y_{n-1})$ . It is straightforward to see that the last integral term in (1.40) can be written only in terms of  $\|\mathbf{x}\|$ , which we denote by y, and of the pdf of the rv  $\|\mathbf{X}_{(n)}\|$ :

$$
\int_{\|\mathbf{x}\| \geqslant \delta} f_{(n)}(\mathbf{x}) \, \mathbb{E}\left(\left\| \left(\Sigma(\|\mathbf{x}\|)\right)^{-1/2} (\mathbf{Y} - \mu(\|\mathbf{x}\|)) \right\|^3 \right) \mathrm{d}\mathbf{x} = \int_{y \geqslant \delta} f_{\|\mathbf{X}_{(n)}\|}(y) \, \mathbb{E}\left(\left\| \left(\Sigma(y)\right)^{-1/2} (\mathbf{Y} - \mu(y)) \right\|^3 \right) \mathrm{d}y. \tag{1.41}
$$

For  $y \geq \delta$ , using Lemma A.3, it is straightforward to see that

$$
\mathbb{E}\left(\left\|\Sigma^{-1/2}(y)\left(\mathbf{Y}-\mu(y)\right)\right\|^3\right) \leqslant C \,\mathbb{E}\left\|\mathbf{Y}-\mu(y)\right\|^3 \leqslant C \left(\mathbb{E}\left\|\mathbf{Y}\right\|^3 + \|\mu(y)\|^3\right) \leqslant C \,\mathbb{E}\left\|\mathbf{Y}\right\|^3.
$$

Splitting the latter expectation into 2 parts according to ( $\|\mathbf{X}\| \leq \delta$ ) or ( $\|\mathbf{X}\| \in [\delta, y]$ ) (and recalling that Y has the truncated cdf  $F_{\mathbf{X} \parallel \mathbf{X} \parallel}(\cdot |y)$ , we can write

$$
\mathbb{E}\left(\left\|\Sigma^{-1/2}(y)\left(\mathbf{Y}-\mu(y)\right)\right\|^3\right) \leqslant C\left(\delta^3 + \mathbb{E}\left(\left\|\mathbf{X}\right\|^3 \mathbb{1}_{\{\left\|\mathbf{X}\right\| \in [\delta,y]\}}\right)\right) \leqslant C\left(\delta^3 + \int_{\delta}^y t^3 L(t)t^{-\alpha-1}dt\right) \leqslant CL(y)y^{3-\alpha},
$$
\n(1.42)

where the last inequality comes from the Karamata's integral theorem (Karamata (1933); see e.g. Proposition 1.5.8 p.26 in Bingham et al. (1987)). So, from  $(1.37)$ – $(1.42)$  we obtain

$$
|\mathbb{P}\left(\mathbf{S}_n \in \mathbf{B}\right) - G_n(\mathbf{B})| \leqslant \frac{C}{\sqrt{n}} \int_{y \geqslant \delta} f_{\|\mathbf{X}_{(n)}\|}(y) L(y) y^{3-\alpha} dy = \frac{C}{\sqrt{n}} \mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|) \|\mathbf{X}_{(n)}\|^{3-\alpha} \mathbb{1}_{\{\|\mathbf{X}_{(n)}\| \geqslant \delta\}}\right).
$$
\n(1.43)

For *n* large enough such that  $n^{1/\alpha} > \delta$ , using (1.32) and (1.38) gives

$$
\mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|)\|\mathbf{X}_{(n)}\|^{3-\alpha}1_{\{|X_{(n)}\|>n^{1/\alpha}\}}\right) = \int_{n^{1/\alpha}}^{\infty} L(y)y^{3-\alpha}nF_{\|\mathbf{X}\|}^{n-1}(y)f_{\|\mathbf{X}\|}(y) \,dy
$$

$$
\leqslant n \int_{n^{1/\alpha}}^{\infty} L(y)y^{2-2\alpha} \,dy \underset{n\to\infty}{\sim} L(n^{1/\alpha})n^{\frac{3-\alpha}{\alpha}}.
$$

Consequently, we have, L denoting a slowly varying function at infinity that may vary from line to line,

$$
\mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|)\|\mathbf{X}_{(n)}\|^{3-\alpha}\mathbb{1}_{\{\|\mathbf{X}_{(n)}\|\geqslant\delta\}}\right) \n\leqslant \mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|)\|\mathbf{X}_{(n)}\|^{3-\alpha}\mathbb{1}_{\{\|\mathbf{X}_{(n)}\|\in[\delta,n^{1/\alpha}]\}}\right) + \mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|)\|\mathbf{X}_{(n)}\|^{3-\alpha}\mathbb{1}_{\{\|\mathbf{X}_{(n)}\|>n^{1/\alpha}\}}\right) \n\leqslant \sup_{y\in[\delta,n^{1/\alpha}]} L(y)n^{\frac{3-\alpha}{\alpha}} + L(n^{1/\alpha})n^{\frac{3-\alpha}{\alpha}} = L(n)n^{\frac{3-\alpha}{\alpha}}.
$$
\n(1.44)

Combining inequalities  $(1.43)$  and  $(1.44)$  provides the result.  $\Box$ 

### A.4 Proof of the results given in Section 4

#### A.4.1 Proof of Lemma 4.1

First let us recall Lemma 4.1 for convenience:

If  $f_X \in 2\mathcal{RV}_{-\alpha-1,-\beta}$ , with  $\alpha > 0$  and  $\beta > 0$ , then the derivative g' of g defined in (4.12), satisfies  $g' \in 2\mathcal{RV}_{\frac{1}{\alpha}-1, \rho}$  , where  $\rho := -\min\left\{1, \frac{\beta}{\alpha}\right\}$  $\frac{\beta}{\alpha}$ .

*Proof.* First we prove that g belongs to  $2\mathcal{RV}_{\gamma-1,\rho}$ , with  $\gamma = 1/\alpha$ . It is straightforward to check that:

$$
g(t) = F_X^{\leftarrow}(e^{-\frac{1}{t}}) = \bar{F}_X^{\leftarrow}(1 - e^{-\frac{1}{t}}).
$$
\n(1.45)

Notice that the function  $h(t) := 1 - e^{-\frac{1}{t}}$  belongs to  $2\mathcal{RV}_{-1,-1}$ . Now, from the  $2RV$  condition on  $f_X$ , we obtain that  $\bar{F}_X \in 2\mathcal{RV}_{-\alpha,-\beta}$  with parameters  $\alpha > 0, \beta > 0$ ; see e.g. Proposition 6 in Hua and Joe (2011). Then, applying Proposition 2.6 in Lv et al. (2012) and denoting by  $2\mathcal{RV}_*(0+)$  the class of regularly varying functions at 0+, we have

$$
\bar{F}_X^{\leftarrow} \in 2\mathcal{RV}_{-\frac{1}{\alpha},-\frac{\beta}{\alpha}}(0+),
$$

from which we deduce, using Proposition 2.8 in Lv et al. (2012) (for a composition of functions) and (1.45),

$$
g\in 2\mathcal{RV}_{\frac{1}{\alpha},-\min\left\{1,\frac{\beta}{\alpha}\right\}}.
$$

Now let us look on the derivative g'. From (1.45), we have  $g'(t) = \frac{e^{-\frac{1}{t}}t^{-2}}{e^{-\frac{1}{t}}(t)}$  $\frac{\partial}{\partial f_X(g(t))}$ .

Again, by Proposition 2.8 and Proposition 2.5 in Lv et al. (2012), we have

$$
\frac{1}{f_X\left(g(t)\right)} \in 2\mathcal{RV}_{\frac{1}{\alpha}+1,-\min\left\{1,\frac{\beta}{\alpha}\right\}} \quad \text{and} \quad e^{-\frac{1}{t}}t^{-2} \in 2\mathcal{RV}_{-2,-1}.
$$

 $\Box$ 

Hence, multiplying the  $2\mathcal{RV}$  functions, we obtain  $g' \in 2\mathcal{RV}_{\frac{1}{\alpha}-1,-\min\left\{1,\frac{\beta}{\alpha}\right\}}$ .

#### A.4.2 Proof of Theorem 4.1

Recall that  $\mathbf{Y}_1, \ldots, \mathbf{Y}_{n-1}$  denote i.i.d. random vectors having the truncated distribution  $F_\mathbf{X}(\cdot \mid y)$ defined in (2.3), while  $\mathbf{X}_y$  is a family of random vectors with distribution (2.5), independent of  ${Y_k}$ . Using conditional probabilities and applying Lemma 2.2, we have

$$
\mathbb{P}\left(\mathbf{S}_{n} \in \mathbf{B}\right) = \int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y)\mathbb{P}\left(\mathbf{X}_{(1)} + \cdots + \mathbf{X}_{(n-1)} + \mathbf{X}_{(n)} \in \mathbf{B}\,|\, \|\mathbf{X}_{(n)}\| = y\right) \mathrm{d}y
$$
\n
$$
= \int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y)\mathbb{P}\left(\mathbf{T}_{n-1} + \mathbf{X}_{y} \in \mathbf{B}\right) \mathrm{d}y, \text{ where } \mathbf{T}_{n-1} := \sum_{k=1}^{n-1} \mathbf{Y}_{k}.
$$

Using the definition (4.23) of the MRV-Normex cdf, then the triangle inequality, we have, for any  $\mathbf{B} \in \mathcal{C}$ ,

$$
|\mathbb{P}(S_n \in \mathbf{B}) - MG_n(\mathbf{B})| = \left| \int_0^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \mathbb{P}\left(\mathbf{\hat{X}}_y + \mathbf{T}_{n-1} \in \mathbf{B}\right) dy - \int_0^{\infty} f_{H_{\alpha,n}}(y) \mathbb{P}(y \cdot \mathbf{\Theta} + Z_y \in \mathbf{B}) dy \right|
$$
  
\n
$$
\leq \int_0^{\infty} \left| f_{\|\mathbf{X}_{(n)}\|}(y) - f_{H_{\alpha,n}}(y) \right| dy
$$
  
\n
$$
+ \int_0^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{\hat{X}}_y + \mathbf{T}_{n-1} \in \mathbf{B}\right) - \mathbb{P}\left(\mathbf{\hat{X}}_y + Z_y \in \mathbf{B}\right) \right| dy
$$
  
\n
$$
+ \int_0^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{\hat{X}}_y + Z_y \in \mathbf{B}\right) - \mathbb{P}(y \cdot \mathbf{\Theta} + Z_y) \right| dy.
$$
\n(1.46)

Recall that, via the Scheffé theorem (see e.g. Billingsley (1968), p. 224), if the cdf F and G are absolutely continuous, then sup  $A \in B(R)$  $|F(A) - G(A)| = \frac{1}{2}$ 2  $\int^{\infty}$ −∞  $|F'(x) - G'(x)| dx.$ 

Therefore, for the first integral in the right-hand side of the inequality (1.46), using Proposition 4.2 and this recall, there exists a slowly-varying function at infinity L such that, for all  $n \geqslant 1$ ,  $\sim$ 

$$
\int_{0}^{\infty} \left| f_{\|\mathbf{X}_{(n)}\|}(y) - f_{H_{\alpha,n}}(y) \right| \mathrm{d}y \leqslant L(n)n^{-\min\left\{1,\frac{\beta}{\alpha}\right\}}.
$$
\n(1.47)

For the second integral appearing in (1.46), following the same approach as for the proof of Theorem 3.1, we can prove that there exists a slowly varying function at infinity, L, such that, for all  $n \geqslant 1$ ,

$$
\int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y)\|\mathbb{P}\left(\mathbf{X}_{y} + \mathbf{T}_{n-1} \in \mathbf{B}\right) - \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in B\right) \,|\mathrm{d}y \leqslant L(n)n^{-\frac{1}{2} + \frac{3-\alpha}{\alpha}}.\tag{1.48}
$$

Indeed, choosing the constant  $\delta > 0$  as in the proof of Theorem 3.1 and splitting the integration domain of the integral in (1.48) into two disjoint sets  $\{y : y < \delta\}$  and  $\{y : y \geq \delta\}$ , we can write

$$
\int_{0}^{\delta} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{T}_{n-1} + \mathbf{X}_{y} \in \mathbf{B}\right) - \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in B\right) \right| \mathrm{d}y \leqslant \int_{0}^{\delta} f_{\|\mathbf{X}_{(n)}\|}(y) |\mathrm{d}y = (F_{\|\mathbf{X}\|}(\delta))^{n}.
$$
\n(1.49)

Let  $C$  denote a positive constant and  $L$  a slowly varying function at infinity, which may vary from line to line. Since  $Y_1, \ldots, Y_{n-1}$  are i.i.d bounded random vectors (Y denoting their parent random vector) with non degenerate distribution (condition  $(C2)$ ), we can use once again the non-generalized Berry-Esseen inequality, and obtain

$$
\int_{\delta}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{T}_{n-1} + \mathbf{X}_y \in \mathbf{B}\right) - \mathbb{P}\left(\mathbf{X}_y + Z_y \in \mathbf{B}\right) \right| dy
$$
\n
$$
\leq \int_{\delta}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \sup_{\mathbf{B} \in \mathcal{C}} |\mathbb{P}(\mathbf{T}_{n-1} \in \mathbf{B}) - \mathbb{P}(Z_y \in \mathbf{B})| dy
$$
\n
$$
\leq \frac{C}{\sqrt{n}} \int_{\delta}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \mathbb{E}\left(\left\| (\Sigma(y))^{-1/2} (\mathbf{Y} - \mu(y))\right\|_{2}^{3}\right) dy. \tag{1.50}
$$

.

Replicating the arguments in the proof of Theorem 3.1, we obtain  $(1.42)$ , *i.e.* 

$$
\mathbb{E}\left(\left\|\Sigma^{-1/2}(y)\left(\mathbf{Y} - \mu(y)\right)\right\|_{2}^{3}\right) \leqslant CL(y) y^{3-\alpha}
$$

We deduce that

$$
\int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{T}_{n-1} + \mathbf{X}_{y} \in \mathbf{B}\right) - \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in B\right) \right| dy
$$
\n
$$
\leq \frac{C}{\sqrt{n}} \int_{y \geq \delta} f_{\|\mathbf{X}_{(n)}\|}(y) L(y) y^{3-\alpha} dy = \frac{C}{\sqrt{n}} \mathbb{E}\left(L(\|\mathbf{X}_{(n)}\|) \|\mathbf{X}_{(n)}\|^{3-\alpha} \mathbb{1}_{\{\|\mathbf{X}_{(n)}\| \geq \delta\}}\right).
$$
\n(1.51)

We are back to the upper bound given in  $(1.43)$ . Combining it with  $(1.44)$  provides the claimed result (1.48).

The last integral in (1.46) is taken care of, in the following technical lemma.

**Lemma A.4.** There exists a slowly varying function at infinity, L, such that, for all  $n \geq 1$ , for any  $\mathbf{B} \in \mathcal{C}$ ,

$$
\int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y)\bigg|\mathbb{P}\left(\mathring{\mathbf{X}}_y + Z_y \in \mathbf{B}\right) - \mathbb{P}\left(y \cdot \mathbf{\Theta} + Z_y \in \mathbf{B}\right)\bigg|{\rm d}y \leqslant L(n)n^{-\frac{\rho}{\alpha}}.
$$

Combining inequalities (1.46), (1.47), (1.48) and Lemma A.4 provides the statement of Theorem 4.1.  $\Box$ 

Let us turn to the proof of Lemma A.4.

*Proof of Lemma A.4.* Define the sequence  $(a_n)_{n\geqslant1}$  with  $a_n := F_{\|\mathbf{X}\|}^{\leftarrow} \left( n^{-\frac{\ln n}{n}} \right) \underset{n\to\infty}{\to} \infty$ . We have

$$
\mathbb{P}\left(\left\|\mathbf{X}_{(n)}\right\| \leqslant a_n\right) = (F_{\left\|\mathbf{X}\right\|}(a_n))^n = n^{-\ln n}.\tag{1.52}
$$

Note that  $a_n = \bar{F}_{\|\mathbf{X}\|}^{\leftarrow} \left(1 - n^{-\frac{\ln n}{n}}\right)$ . Since the function  $u(n) := 1 - n^{-\frac{\ln n}{n}} \in RV_{-1}$  (at infinity) and  $||\mathbf{X}|| \in \mathcal{RV}_{-\alpha}$ , so  $\overline{F}_{||\mathbf{X}||} \in \mathcal{RV}_{-1/\alpha}$  at  $0^+$ , therefore

$$
a_n = \bar{F}_{\|\mathbf{X}\|}^{\leftarrow} \circ u(n) \in \mathcal{RV}_{1/\alpha} \quad \text{(at infinity)} \tag{1.53}
$$

(for a reminder on properties of  $RV$ , see e.g. Lv et al.  $(2012)$ ).

As  $\bar{F}_{\|\mathbf{X}\|}(a_n) = n\left(1 - n^{-\frac{\ln n}{n}}\right)$ , there exists also a slowly varying function at infinity, L, such that

$$
n\bar{F}_{\|\mathbf{X}\|}(a_n) = L(n). \tag{1.54}
$$

Splitting the integral of Lemma A.4 into two sets  $\{y \leq a_n\}$  and  $\{y > a_n\}$  and using the pdf for

the maximum (1.32), we can write

$$
\int_{0}^{\infty} f_{\|\mathbf{X}_{(n)}\|}(y) \left| \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in \mathbf{B}\right) - \mathbb{P}\left(y \cdot \mathbf{\Theta} + Z_{y} \in \mathbf{B}\right) \right| dy
$$
\n
$$
\leq \mathbb{P}\left(\|\mathbf{X}_{(n)}\| \leq a_{n}\right) + \int_{a_{n}}^{\infty} n f_{\|\mathbf{X}\|}(y) F_{\|\mathbf{X}\|}^{n-1}(y) \left| \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in \mathbf{B}\right) - \mathbb{P}\left(y \cdot \mathbf{\Theta} + Z_{y} \in \mathbf{B}\right) \right| dy
$$
\n
$$
\leq n^{-\ln n} + \int_{a_{n}}^{\infty} n f_{\|\mathbf{X}\|}(y) \left| \mathbb{P}\left(\mathbf{X}_{y} + Z_{y} \in \mathbf{B}\right) - \mathbb{P}\left(y \cdot \mathbf{\Theta} + Z_{y} \in \mathbf{B}\right) \right| dy, \quad \text{using (1.52).}
$$
\n(1.55)

To estimate the integral given in (1.55), we use a coupling technique, namely: For any random vector  $(\xi, \eta)$  in some complete separable metric space, and for any measurable set B, we have

$$
|\mathbb{P}(\xi \in B) - \mathbb{P}(\eta \in B)| \leq \mathbb{P}(\xi \neq \eta).
$$

So, for any joint distribution of  $\mathbf{\hat{X}}_y$  and  $\Theta$  (or, more generally, for any joint distribution of X and  $\Theta$ ), we have

$$
\left|\mathbb{P}\left(\mathbf{X}_y + Z_y \in \mathbf{B}\right) - \mathbb{P}\left(y \cdot \mathbf{\Theta} + Z_y \in \mathbf{B}\right)\right| \leqslant \mathbb{P}\left(\mathbf{X}_y \neq y \cdot \mathbf{\Theta}\right),
$$

which gives the following upper bound for the integral in  $(1.55)$ , using  $(2.5)$ :

$$
\int_{a_n}^{\infty} n f_{\|{\bf X}\|}(y) \mathbb{P}\left(\frac{\hat{\bf X}_y}{y} \neq {\bf \Theta}\right) dy = n \mathbb{P}\left(\frac{{\bf X}}{\|{\bf X}\|} \neq {\bf \Theta}, \|{\bf X}\| > a_n\right).
$$

Therefore, we have

$$
(1.55) \leqslant n^{-\ln n} + n \mathbb{P}(\|\mathbf{X}\| > a_n) \mathbb{P}\left(\frac{\mathbf{X}}{\|\mathbf{X}\|} \neq \mathbf{\Theta} \Big| \|\mathbf{X}\| > a_n\right). \tag{1.56}
$$

Now, at given n, we define the joint distribution of **X** and  $\Theta$  such that:

(i)  $\Theta$  is independent of the event  $\{\Vert \mathbf{X} \Vert > a_n\},\$ 

(ii)  $(X / ||X||, \Theta)$  has a joint distribution defined by Dobrushin's theorem (see Dobrushin  $(1970)$ .

Hence we can write, using Condition  $(M_{\Theta})$  to get the asymptotic behavior,

$$
\mathbb{P}\left(\left.\frac{\mathbf{X}}{\|\mathbf{X}\|}\neq\mathbf{\Theta}\,\right|\|\mathbf{X}\|>a_n\right)=\sup_{\mathbf{B}\subseteq\mathcal{S}_1}\left|\mathbb{P}\left(\left.\frac{\mathbf{X}}{\|\mathbf{X}\|}\in\mathbf{B}\,\right|\|\mathbf{X}\|>a_n\right)-\mathbb{P}\left(\mathbf{\Theta}\in\mathbf{B}\right)\right|\right|\underset{n\to\infty}{\sim}A(a_n),
$$

where  $A(a_n) \in \mathcal{RV}_{-\rho/\alpha}$  by combining  $(M_\Theta)$  and  $(1.53)$ . Reporting this last result in  $(1.56)$  and using  $(1.54)$ , we obtain

$$
(1.55) \leqslant n^{-\ln n} + n\bar{F}_{\|\mathbf{X}\|}(a_n) A(a_n) = n^{-\ln n} + L(n) A(a_n),
$$

from which the result of Lemma A.4 follows.

 $\Box$ 

#### A.4.3 Discussion on Condition  $(M_{\Theta})$

Proof of 3(b) given in Remark 4.3 - Let us prove that Condition (4.25) given on the form of the pdf of a random vector is a necessary and sufficient condition to have the independence between the norm and direction of this vector.

*Proof.* Notice that the unit ball  $\{x : ||x|| \leq 1\}$  is a convex set, which boundary  $S_1$  is a differentiable manifold. Further we want to integrate on  $S_1$ , and assume w.l.o.g. that the atlas for  $S_1$  consists of one chart only. (Indeed, to provide the integration in the case of many charts, one would just need to use a partition and sum the integrals over all charts.)

If a map  $\varphi : \mathbb{R}^{d-1} \to \mathcal{S}_1$  parameterizes a set B on the unit sphere  $\mathcal{S}_1$ , one can introduce the map

$$
\widehat{\varphi} : \mathbb{R}^{d-1} \to \mathcal{S}_y
$$

$$
\mathbf{u} \to \widehat{\varphi}(\mathbf{u}) := y \, \varphi(\mathbf{u})
$$

that parameterizes a set  $B_y := \{ \mathbf{x} : \mathbf{x}/\|\mathbf{x}\| \in B, \|\mathbf{x}\| = y \}$  on the sphere  $\mathcal{S}_y := \{ \mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| = y \}.$ Note that both functions  $\varphi$ ,  $\widehat{\varphi}$  are differentiable a.e.

Now, assuming that the pdf of  $X$  satisfies (4.25), the joint distribution of the random variable  $\|\mathbf{X}\|$  and the random vector  $\mathbf{X}/\|\mathbf{X}\|$  can be expressed as:

$$
\mathbb{P}\left(\left.\|\mathbf{X}\|\leqslant t,\, \frac{\mathbf{X}}{\|\mathbf{X}\|}\in B\right.\right)=\int\limits_{\left\|\mathbf{x}\right\|\leqslant t,\, \frac{\mathbf{x}}{\|\mathbf{x}\|}\in B}h(\|\mathbf{x}\|)g(\mathbf{x}/\|\mathbf{x}\|) \mathrm{d}\mathbf{x}=\int\limits_{y\in[0,t],\, \mathbf{u}\in\varphi^{-1}(B)}\hspace{-0.5cm}h(y)g(\varphi(\mathbf{u}))J(y,\mathbf{u}) \mathrm{d}y \mathrm{d}\mathbf{u}
$$

where  $J(y, u)$  denotes the Jacobian for the change of variables

$$
\mathbf{x} = y\varphi(\mathbf{u}).\tag{1.57}
$$

 $\bigg\}$  $\begin{array}{c} \end{array}$  $\overline{\phantom{a}}$  $\overline{\phantom{a}}$  $\overline{\phantom{a}}$  $\overline{\phantom{a}}$  $\overline{\phantom{a}}$  $\bigg\}$  $\begin{array}{c} \end{array}$ 

Let us compute  $J(y, u)$ , which is the determinant of the  $d \times d$ -matrix  $\left(\frac{\partial \mathbf{x}}{\partial x}\right)$  $\partial(y,\mathbf{u})$  $\setminus$ .

$$
J(y, \mathbf{u}) := \begin{vmatrix} \frac{\partial x_1}{\partial y} & \frac{\partial x_2}{\partial y} & \cdots & \frac{\partial x_d}{\partial y} \\ \frac{\partial x_1}{\partial u_1} & \frac{\partial x_2}{\partial u_1} & \cdots & \frac{\partial x_d}{\partial u_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_1}{\partial u_{d-1}} & \frac{\partial x_2}{\partial u_{d-1}} & \cdots & \frac{\partial x_d}{\partial u_{d-1}} \end{vmatrix} = \begin{vmatrix} \varphi_1 & \varphi_2 & \cdots & \varphi_d \\ y \frac{\varphi_1}{\partial u_1} & y \frac{\varphi_2}{\partial u_1} & \cdots & y \frac{\varphi_d}{\partial u_1} \\ \vdots & \vdots & \ddots & \vdots \\ y \frac{\varphi_1}{\partial u_{d-1}} & y \frac{\varphi_2}{\partial u_{d-1}} & \cdots & y \frac{\varphi_d}{\partial u_{d-1}} \end{vmatrix}
$$
  
=  $y^{d-1} \times \begin{vmatrix} \varphi_1 & \varphi_2 & \cdots & \varphi_d \\ \frac{\varphi_1}{\partial u_1} & \frac{\varphi_2}{\partial u_1} & \cdots & \frac{\varphi_d}{\partial u_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\varphi_1}{\partial u_{d-1}} & \frac{\varphi_2}{\partial u_{d-1}} & \cdots & \frac{\varphi_d}{\partial u_{d-1}} \end{vmatrix} = y^{d-1} \times J(1, \mathbf{u}).$ 

Therefore, we can write

$$
\mathbb{P}\left(\|\mathbf{X}\| \leqslant t, \mathbf{X}/\|\mathbf{X}\| \in B\right) = \int h(y)y^{d-1}g(\varphi(\mathbf{u}))J(1, \mathbf{u}) \, \mathrm{d}y \, \mathrm{d}\mathbf{u}
$$
\n
$$
y \in [0, t], \mathbf{u} \in \varphi^{-1}(B)
$$
\n
$$
= \int \int \int \int \int \int \int g(\varphi(\mathbf{u}))J(1, \mathbf{u}) \, \mathrm{d}\mathbf{u} = \mathbb{P}\left(\|\mathbf{X}\| \leqslant t\right) \mathbb{P}\left(\frac{\mathbf{X}}{\|\mathbf{X}\|} \in B\right)
$$
\n
$$
\text{with } c_d := \int \int \int \int \int \int g(\varphi(\mathbf{u}))J(1, \mathbf{u}) \, \mathrm{d}\mathbf{u} = \left(\int \int \int \int h(y)y^{d-1} \, \mathrm{d}y\right)^{-1}.
$$
\n
$$
\text{with } c_d := \int \int \int \int \int \int g(\varphi(\mathbf{u}))J(1, \mathbf{u}) \, \mathrm{d}\mathbf{u} = \left(\int \int \int \int h(y)y^{d-1} \, \mathrm{d}y\right)^{-1}.
$$
\n
$$
(1.58)
$$

We deduce that  $\|\mathbf{X}\|$  and  $\frac{\mathbf{X}}{\|\mathbf{X}\|}$  $\|\mathbf{X}\|$ are independent.

Let us prove the converse, assuming that  $\|\mathbf{X}\|$  and  $\mathbf{X}/\|\mathbf{X}\|$  are independent.

Note that, using the same change of variable (1.57), we can write

$$
\int_{\|\mathbf{x}\| \leqslant t, \mathbf{x}/\|\mathbf{x}\| \in B} f_{\mathbf{X}}(\mathbf{x}) \mathrm{d}\mathbf{x} = \int_{y \in [0,t], \mathbf{u} \in \varphi^{-1}(B)} f_{\mathbf{X}}(y\varphi(\mathbf{u})) y^{d-1} J(1, \mathbf{u}) \mathrm{d}y \mathrm{d}\mathbf{u}.
$$
 (1.59)

Moreover, we have

$$
\mathbb{P}\left(\|\mathbf{X}\| \leqslant t, \mathbf{X}/\|\mathbf{X}\| \in B\right) = \mathbb{P}\left(\|\mathbf{X}\| \leqslant t\right) \mathbb{P}\left(\mathbf{X}/\|\mathbf{X}\| \in B\right)
$$
\n
$$
= \int_{\|\mathbf{x}\| \leqslant t} f_{\mathbf{X}}(\mathbf{x}) \, \mathrm{d}\mathbf{x} \int_{\|\mathbf{x}\| \in B} f_{\mathbf{X}}(\mathbf{x}) \, \mathrm{d}\mathbf{x}
$$
\n
$$
= \int_{y \in [0,t]; \, \mathbf{u} \in \mathbb{R}^{d-1}} f_{\mathbf{X}}(y\varphi(\mathbf{u})) y^{d-1} J(1, \mathbf{u}) \, \mathrm{d}y \, \mathrm{d}\mathbf{u} \int_{y \in \mathbb{R}^{+}; \, \mathbf{u} \in \varphi^{-1}(B)} f_{\mathbf{X}}(y\varphi(\mathbf{u})) y^{d-1} J(1, \mathbf{u}) \, \mathrm{d}y \, \mathrm{d}\mathbf{u}.
$$
\n(1.60)

Since  $\mathbb{P}(\|\mathbf{X}\| \leq t, \mathbf{X}/\|\mathbf{X}\| \in B) = \int f_{\mathbf{X}}(\mathbf{x})d\mathbf{x}$ , then the right hand side of (1.59) equals that  $\Vert \mathbf{x} \Vert \leqslant t, \mathbf{x}/\Vert \mathbf{x} \Vert \in B$ of (1.60). Consequently, for  $t_0 \in (0,\infty)$  and  $\varphi(\mathbf{u}_0) \in \mathcal{S}_1$ , we can write

$$
f_{\mathbf{X}}(t_0 \varphi(\mathbf{u}_0)) t_0^{d-1} J(1, \mathbf{u}_0) = \int_{\varphi(\mathbf{u}) \in \mathcal{S}_1} f_{\mathbf{X}}(t_0 \varphi(\mathbf{u})) t_0^{d-1} J(1, \mathbf{u}) d\mathbf{u} \times \int_0^{\infty} f_{\mathbf{X}}(y \varphi(\mathbf{u}_0)) y^{d-1} J(1, \mathbf{u}_0) dy \text{ a.e. },
$$

thus, simplifying, we obtain

$$
f_{\mathbf{X}}(t_0\varphi(\mathbf{u}_0)) = \int_{\varphi(\mathbf{u}) \in \mathcal{S}_1} f_{\mathbf{X}}(t_0\varphi(\mathbf{u})) J(1, \mathbf{u}) \mathrm{d}\mathbf{u} \times \int_{0}^{\infty} f_{\mathbf{X}}(y\varphi(\mathbf{u}_0)) y^{d-1} \mathrm{d}y =: h_1(t_0) \times h_2(\varphi(\mathbf{u}_0)),
$$

 $\Box$ 

proving the existence of the function  $h_1$  and  $h_2$  satisfying (4.25).

Let us end with a remark. It appears that assuming that the function related to the direction  $\mathbf{X}/\|\mathbf{X}\|$  (here,  $h_2$ ) is a constant, does not guarantee that the distribution of  $\mathbf{X}/\|\mathbf{X}\|$  is uniform on the unit sphere  $S_1$ . Nevertheless, for  $L^1$ ,  $L^2$  and  $L^{\infty}$  norms (or their weighted versions), this distribution is uniform on the unit sphere  $S_1$ :

$$
\mathbb{P}(\mathbf{X}/\|\mathbf{X}\| \in B) = \frac{\mu(B)}{\mu(S_1)}
$$

where  $\mu$  is a measure on a sphere. But for  $L^p$ -norms with  $p \neq 1, 2, \infty$ , it is not uniform, as a measure on the unit sphere is not proportional to a measure on the unit ball.

Using integration on sphere: an illustration with the norm's cdf - Assume the random vector **X** has a pdf defined, for any  $\mathbf{x} \in \mathbb{R}^d$ , by

$$
f(\mathbf{x}) = h_1(||\mathbf{x}||) h_2(\mathbf{x}/||\mathbf{x}||) \quad \text{with} \quad h_1(||\mathbf{x}||) := C_d (1 + ||\mathbf{x}||)^{-(\alpha + d)}, \tag{1.61}
$$

,

 $C_d > 0$  being the normalizing constant (making f a pdf).

Note that, if  $h_2$  is a constant, then we get back the multivariate Pareto pdf considered in the example developed in Section 3.3.

Changing to polar coordinates (1.57), setting  $r = ||\mathbf{x}||$  and  $\theta = \frac{\mathbf{x}}{||\mathbf{x}||}$  $\frac{\mathbf{x}}{\|\mathbf{x}\|}$ , we can write

$$
\frac{1}{C_d} = \int_{\mathbb{R}^d} \frac{h_2(\mathbf{x}/\|\mathbf{x}\|)}{(1+\|\mathbf{x}\|)^{\alpha+d}} \,\mathrm{d}\mathbf{x} = c_d \int_0^\infty \frac{r^{d-1}}{(1+r)^{\alpha+d}} \,\mathrm{d}r = c_d \frac{\Gamma(d)\Gamma(\alpha)}{\Gamma(\alpha+d)},
$$

where  $c_d$  is defined after (1.58). Then the cdf of  $\|\mathbf{X}\|$  can be expressed as

$$
F_{\|\mathbf{X}\|}(t) = \mathbb{P}\left(\|\mathbf{X}\| \leq t\right) = C_d c_d \int_0^t r^{d-1} \frac{1}{(1+r)^{\alpha+d}} \, dr = 1 - \sum_{k=0}^{d-1} \frac{\Gamma(\alpha+k)}{\Gamma(k+1)\Gamma(\alpha)} \frac{t^k}{(1+t)^{\alpha+k}},\tag{1.62}
$$

proceeding successively by an integration by parts and using the fact that, for any  $m \in \mathbb{N}$ ,  $\alpha > 0$  such that  $\alpha > m + 1$ ,

$$
\int_0^\infty \frac{x^m}{(1+x)^\alpha} dx = \frac{\Gamma(m+1)\Gamma(\alpha - (m+1))}{\Gamma(\alpha)},
$$

to obtain the last equality in (1.62).

Note that this result is independent of the norm choice and it will be the same when applying Normex method to any  $X$  having its pdf satisfying  $(1.61)$ .

The result (1.62) will be useful for applying both versions of multi-normex, d-Normex and MRV-Normex, in the various examples considered in the paper.

### A.5 Geometric notion of multivariate quantiles

In this section, we give an overview of the notion of multivariate quantiles (see e.g. Chaudhuri (1996) and Dhar et al. (2014)) and QQ-plots.

#### One dimensional case

In the one dimensional case, the quantile  $q_\alpha(X)$  of level  $\alpha \in (0,1)$  for a random variable  $X \in \mathbb{R}$ can be defined as follows

$$
q_{\alpha}(X) := \arg\min_{q \in \mathbb{R}} \left\{ \mathbb{E} \left( |X - q| - |X| \right) - q(2\alpha - 1) \right\}. \tag{1.63}
$$

The proof of (1.63) is easy to obtain by the following representation for the expectation:

$$
\mathbb{E}X = \int_0^\infty \bar{F}(t)dt - \int_{-\infty}^0 F(t)dt.
$$

If the first moment of X exists, the term  $|X|$  can be dropped out in Definition 1.63, hence we have

$$
q_{\alpha}(X) := \arg\min_{q \in \mathbb{R}} \left\{ \mathbb{E}|X - q| - q(2\alpha - 1) \right\}, \quad \alpha \in (0, 1).
$$

Sometimes, it is convenient to parameterize the levels  $\alpha \in (0,1)$  by the set  $(-1,1)$  with the linear transformation  $u = 2\alpha - 1$ , so that the median  $(\alpha = 1/2)$  corresponds to  $u = 0$ . In such a case, we have

$$
q_u(X) := \arg\min_{q \in \mathbb{R}} \{ \mathbb{E}|X - q| - uq \}, \quad u \in (-1, 1).
$$

#### Multidimensional case

Let F denote the multivariate cdf of the random vector  $\mathbf{X} \in \mathbb{R}^d$ , with  $d \geq 1, \langle \cdot, \cdot \rangle$  be the Euclidean inner product and  $\|\cdot\|$  the Euclidean norm induced by the inner product.

Definition A.7 (Geometrical quantile, see Chaudhuri (1996)). For a random vector X with a probability distribution  $F$  on  $\mathbb{R}^d$ , the d-dimensional spatial quantile or geometrical quantile  $(GQ) Q_F(\mathbf{u}) = (Q_{F,1}(\mathbf{u}), \ldots, Q_{F,d}(\mathbf{u}))$  is defined as

$$
Q_F(\mathbf{u}) = \arg\min_{\mathbf{Q} \in \mathbb{R}^d} \mathbb{E}\{\|\mathbf{X} - \mathbf{Q}\| - \|\mathbf{X}\| - \langle \mathbf{u}, \mathbf{Q} \rangle\},\tag{1.64}
$$

with  $\mathbf{u} \in B^d = \{ \mathbf{v} \in \mathbb{R}^d, ||\mathbf{v}|| < 1 \}$ .

If  $u = 0$ , then  $Q_F(0)$  is a spatial median - a point from  $\mathbb{R}^d$  that is equidistant from all values of a random vector X.

GQs satisfy the equivariant property under rotation and shifting: For deterministic vector  $\mathbf{a} \in \mathbb{R}^d$  and orthogonal matrix A we can find a GQ of  $A\mathbf{X} + \mathbf{a}$  as

$$
\mathbf{Q}_{A\mathbf{X} + \mathbf{a}}(\mathbf{u}) = A\mathbf{Q}_{\mathbf{X}}(A^{\mathrm{T}}\mathbf{u}) + \mathbf{a},
$$

where  $\mathbf{Q}_{\mathbf{X}}(\mathbf{u})$  is a GQ of **X**.

Unfortunately, GQs are not equivariant under arbitrary affine transformations. In Chakraborty (2001), the author used a transformation-retransformation approach to construct affine equivariant estimates of multivariate quantiles, but the object became quite complicated and untractable.

Nevertheless, GQs characterize the distribution, namely, if two random vectors  $X$  and  $Y$  yield the same quantile function  $q$ , then they have the same distribution (see Koltchinskii (1997)). For extreme multivariate quantiles, this characterization does not hold true whenever  $\mathbb{E} \|\mathbf{X}\|^2 < \infty$ : The GQs converge to  $\infty$  as the length  $\lambda = ||\mathbf{u}|| \to 1$  with the rate  $(1 - \lambda)^{-1/2}$ , more precisely

$$
\|\mathbf{Q}(\lambda\mathbf{u})\|^2(1-\lambda) \to \frac{1}{2} \left( \operatorname{tr} \Sigma - u^{\mathrm{T}} \Sigma u \right) \text{ as } \lambda \uparrow 1,
$$

where  $\Sigma$  denotes the covariance matrix of **X** (see Girard and Stupfler (2015)). Thus the notion of GQ is not relevant for extreme multivariate quantiles of  $L^p$ -vectors with  $p \geq 2$ , since for different distributions having the same covariance matrix, GQs equivalently converge to  $\infty$ . But, in the case  $\mathbb{E} \|\mathbf{X}\|^2 = \infty$ , the same authors prove in Girard and Stupfler (2015) that, for MRV distributions with parameter  $\alpha \in (0, 2]$ , the asymptotics of GQs depend on the regular function introduced in the definition of MRV (A.5), making then the GQs more relevant.

If all first moments of the random vector  $\bf{X}$  exist, then Equation (1.64) can be written as

$$
Q_F(\mathbf{u}) = \arg\min_{\mathbf{Q} \in \mathbb{R}^d} \mathbb{E} \left[ \|\mathbf{X} - \mathbf{Q}\| - \langle \mathbf{u}, \mathbf{Q} \rangle \right].
$$

Considering a sample  $\mathbf{X}_1, \ldots, \mathbf{X}_n$  of iid random vectors of  $\mathbb{R}^d$  with distribution F,  $Q_F(\mathbf{u})$  can be estimated by the empirical quantile

$$
\hat{Q}_F(\mathbf{u}) = \arg\min_{\mathbf{Q} \in \mathbb{R}^d} \frac{1}{n} \sum_{i=1}^n \left( \|\mathbf{X}_i - \mathbf{Q}\| - \langle \mathbf{u}, \mathbf{Q} \rangle \right). \tag{1.65}
$$

In this case, the notion of **spatial rank** may be introduced for each element  $X_j$  of the sample, namely

$$
\mathbf{u}_{j} := n^{-1} \sum_{i \neq j} \left\| \mathbf{X}_{j} - \mathbf{X}_{i} \right\|^{-1} \left( \mathbf{X}_{j} - \mathbf{X}_{i} \right).
$$

It is straightforward to verify that the GQ  $\hat{Q}_F(\mathbf{u}_j)$  evaluated at the spatial rank  $\mathbf{u}_j$  is equal to the appropriate element  $X_j$  of the sample:

$$
\hat{Q}_F(\mathbf{u}_j) = \mathbf{X}_j. \tag{1.66}
$$

This is a key property in the construction of QQ-plots of multivariate quantiles for small samples.

To compute the GQs, an algorithm has been proposed in Chaudhuri (1996), based on the optimization problem (1.65), while another algorithm to construct QQ-plots has been developed in Dhar et al. (2014).

#### Multivariate QQ-plot

To construct QQ-plots, we follow the algorithm from Dhar et al. (2014) but with a small modification, given the fact that we use  $10<sup>7</sup>$  simulations to evaluate the considered distribution. Instead of considering a number of points (in the plot) equal to the general volume of compared samples, as suggested in Dhar et al. (2014), which in our case would represent a huge amount of points, we decide to fix a set of 'levels'  $\mathbf{u} \in B^d$  and compare multi-quantiles from different distributions for these levels.

More precisely, we consider two independent d-dimensional data sets, namely,  $\mathcal{X} = {\mathbf{x}_1, \dots, \mathbf{x}_n}$ and  $\mathcal{Y} = {\mathbf{y}_1, \dots, \mathbf{y}_m}$ , where  $\mathbf{x}_i = (x_{i,1}, \dots, x_{i,d})$  are the realizations of a random sample with underlying distribution F, and  $\mathbf{y}_j = (y_{j,1}, \ldots, y_{j,d})$  with distribution G. Let  $\mathcal{U} = {\mathbf{u}_1, \ldots, \mathbf{u}_n}$ be a set of vectors from  $B^d$ . We compute  $Q_{\mathcal{X}}(\mathbf{u}_k)$  and  $Q_{\mathcal{Y}}(\mathbf{u}_k)$  for  $k = 1, \ldots, n$ , using the algorithm from Chaudhuri (1996). Then, we can match the two sets of quantiles by setting the correspondence between  $Q_{\mathcal{X}}(\mathbf{u}_k)$  and  $Q_{\mathcal{Y}}(\mathbf{u}_k)$  for  $k = 1, \ldots, n$ . Finally, we construct the QQ-plots as a collection of d 2-dimensional plots, where each plot corresponds to a component of the spatial quantile. In Dhar et al. (2014), Theorem 2.2, the authors proved that for all  $i = 1, \dots, d$ , the points in the *i*-th 2-dimensional plot will lie close to a straight line with slope 1 and intercept 0 if and only if  $F = G$ .



#### A.6 Extra scatter plots

Figure 10: Pareto-Lomax marginal distribution with survival Clayton copula ( $\alpha\theta = 1$ ) with parameter  $\alpha = 2.3$ . First row –  $\|\cdot\|_{\infty}$  norm, second row –  $\|\cdot\|_{1}$  norm. Sample size = 10<sup>4</sup>.



Figure 11: Pareto-Lomax marginal distribution with survival Clayton copula ( $\alpha\theta = 1$ ) with parameter  $\alpha = 1.01$ . First row –  $\|\cdot\|_{\infty}$  norm, second row –  $\|\cdot\|_{1}$  norm. Sample size = 10<sup>4</sup>.

# A.7 Truncated moments for certain distributions to apply multinormex

### A.7.1 Example 4.3.2 with Clayton copula

In this section we provide truncated moments for the example 4.3.2 in the case  $d = 2, \alpha\theta = 1$ and  $\|\cdot\|_{\infty}$  norm in programming language to be able to apply multi-normex methods.  $\mathbb{E}(X_1; \|\mathbf{X}\|_{\infty} \leqslant y) =$ 

```
((((-2*a-2)*sqrt(y+1)+(2*a^2+a+1)*y+2*a^2+a+2)*exp(-2*a*log(2*sqrt(y+1)-1)))+(exp(-a*log(y+1))*(((2*sqrt(y+1)-1)^(2*a)*((2*a-1)*(y+1)^a+2*a^2-3*a+1)+(-
2*a^2+3*a-1)*(y+1)^a)*exp(a*log(y+1))+(y+1)^a*(2*sqrt(y+1)-1)^(2*a)*(2*2*a))*sqrt(y+1)+(-2*a^2+a-1)*y-2*a^2+a-1)))/((y+1)^a*(2*sqrt(y+1)-1)^(2*a)))/(2
*a^2-3*a+1))
```

```
\mathbb{E}(X_1^2; \|\mathbf{X}\|_{\infty} \leq y) =
```

```
(((2*a-3)*(2*a-1)*((2*sqrt(y+1)-1)^(2*a)*(2*(y+1)^a-(a-2)*(a-1))+(a-2)*(a-1))*(y+1)^a) * exp(a * log(y+1)) - (y+1)^a(a+1)*(2 * sqrt(y+1)-1)^(2 *a)*(12*(a-2)*sqrt(y+1))+1)+(4*a^4-12*a^3+11*a^2-3*a+6)*y-4*a^4+20*a^3-19*a^2-19*a+30))*exp(2*a*log(
2*sqrt(y+1)-1))-(y+1)^a*(-y*((4*a^4-4*a^3+11*a^2+7*a+6)*y+2*(8*a^3+10*a^2+23
*a+18))+4*sqrt(y+1)*((a+1)*(2*a^2+a+6)*y+8*a^2+11*a+6)+4*a^4-20*a^3-5*a^2-53
```
 $*a-24)*(2*sqrt(y+1)-1)^{2*a)*exp(a*log(y+1)))*exp(-a*(2*log(2*sqrt(y+1)-1)+l)$  $o(g(y+1)))/((a-2)*(a-1)*(2*a-3)*(2*a-1)*(y+1)^a*(2*sqrt(y+1)-1)^(2*a))$ 

 $\mathbb{E}(X_1X_2; \|\mathbf{X}\|_{\infty} \leq y) =$ 

 $(((((2*sqrt(y+1)-1)^{2}(2*a)*((4*a^3-10*a^2+5*a)*(y+1)^a+(-4*a^4+16*a^3-17*a^2)))$ -a+6)\*sqrt(y+1)+(4\*a^5-16\*a^4+17\*a^3+a^2-6\*a)\*y+4\*a^4-16\*a^3+17\*a^2+a-6)+(y +1)^a\*((4\*a^4-16\*a^3+17\*a^2+a-6)\*sqrt(y+1)+(-4\*a^5+16\*a^4-17\*a^3-a^2+6\*a)\*y  $-4*a^4+16*a^3-17*a^2-a+6)$ )\*exp(a\*log(y+1))+(y+1)^a\*((-4\*a^3+10\*a^2-6\*a)\*y^2 +sqrt(y+1)\*((-8\*a^4+28\*a^3-28\*a^2+8\*a)\*y+4\*a^4-24\*a^3+41\*a^2-15\*a-6)+(-4\*a^ 5+16\*a^4-21\*a^3+5\*a^2+6\*a)\*y-4\*a^4+16\*a^3-21\*a^2+5\*a+6)\*(2\*sqrt(y+1)-1)^(2\* a))\*exp(2\*a\*log(2\*sqrt(y+1)-1))+(y+1)^a\*((4\*a^5-4\*a^4-5\*a^3-a^2+6\*a)\*y^2+sq rt(y+1)\*((-8\*a^4+4\*a^3+12\*a^2-8\*a)\*y-4\*a^4-13\*a^2+15\*a+6)+(4\*a^5+a^3-3\*a^2- 6\*a)\*y+4\*a^4+4\*a^3+3\*a^2-10\*a-6)\*(2\*sqrt(y+1)-1)^(2\*a)\*exp(a\*log(y+1)))\*exp  $(-2*a*log(2*sqrt(y+1)-1)-a*log(y+1)))/(a*(4*a^2-20*a^3+35*a^2-25*a+6)*(y+1)$  $\hat{a}*(2*sqrt(y+1)-1)^{2}(2*a)$ 

### A.7.2 Multivariate Pareto-Lomax with  $L^{\infty}$ -norm

In this section we provide an example of Multivariate Pareto-Lomax distribution, defined by  $L^{\infty}$ -norm, and calculations of truncated (by  $L^{\infty}$ -norm as well) moments for an arbitrary dimension  $d \geq 1$ . One can use this calculations to apply multi-normex methods. Let the density function of the initial random vector is

$$
f_{\mathbf{X}}(\mathbf{x}) := \frac{C_d}{(1 + \|\mathbf{x}\|_{\infty})^{\alpha + d}}, x_i > 0, i = 1, \cdots, d.
$$

As usual, parameter  $\alpha > 0$ , and for the constant  $c_d$  we have

$$
\frac{1}{C_d} = \iint_{\mathbb{R}_+^d} f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x} = d \int_0^\infty \frac{x^{d-1}}{(1+x)^{\alpha+d}} dx = \frac{\Gamma(d+1)\Gamma(\alpha)}{\Gamma(\alpha+d)}.
$$

$$
\mathbb{E}X_{1} = C_{d} \iint_{\mathbb{R}_{d}^{+}} \frac{x_{1}}{(1 + \max_{i} x_{i})^{\alpha+d}} d\mathbf{x}
$$
  
\n
$$
= C_{d} \left[ \int_{x_{1} - \max_{i} x_{i}} \frac{x_{1}}{(1 + x_{1})^{\alpha+d}} d\mathbf{x} + (d - 1) \iint_{x_{2} = \max_{i} x_{i}} \frac{x_{1}}{(1 + x_{2})^{\alpha+d}} d\mathbf{x} \right]
$$
  
\n
$$
= C_{d} \left[ \int_{0}^{\infty} \frac{x_{1}^{d}}{(1 + x_{1})^{\alpha+d}} d\mathbf{x}_{1} + (d - 1) \int_{0}^{\infty} \frac{1}{2} \frac{x_{2}^{d}}{(1 + x_{2})^{\alpha+d}} d\mathbf{x}_{2} \right]
$$
  
\n
$$
= C_{d} \cdot \left( 1 + \frac{(d - 1)}{2} \right) \int_{0}^{\infty} \frac{x_{1}^{d}}{(1 + x_{1})^{\alpha+d}} d\mathbf{x}_{1} = \frac{d + 1}{2(\alpha - 1)}.
$$

Using the same arguments

$$
\mathbb{E}\left(X_1; \|\mathbf{X}\|_{\infty} \leq t\right) = C_d \cdot \left(1 + \frac{(d-1)}{2}\right) \int_0^t \frac{x_1^d}{\left(1 + x_1\right)^{\alpha + d}} dx_1
$$

$$
= \left(\frac{d+1}{2}\right) \left(\frac{1}{\alpha - 1} - \sum_{k=0}^d \frac{\Gamma(\alpha + k - 1)}{\Gamma(k + 1)\Gamma(\alpha)} \frac{t^k}{(1 + t)^{\alpha + k - 1}}\right).
$$

$$
\mathbb{E}X_1^2 = C_d \iint_{\mathbb{R}_d^+} \frac{x_1^2}{(1 + \max_i x_i)^{\alpha + d}} d\mathbf{x}
$$
  
=  $C_d \cdot \left(1 + \frac{(d-1)}{3}\right) \int_0^\infty \frac{x_1^{d+1}}{(1 + x_1)^{\alpha + d}} dx_1 = \frac{(d+1)(d+2)}{3(\alpha - 1)(\alpha - 2)}$ .  

$$
\mathbb{E}\left(X_1^2; \|\mathbf{X}\|_{\infty} \leq t\right) = C_d \cdot \left(1 + \frac{(d-1)}{3}\right) \int_0^t \frac{x_1^{d+1}}{(1 + x_1)^{\alpha + d}} dx_1
$$

$$
=\frac{(d+1)(d+2)}{3}\left(\frac{1}{(\alpha-1)(\alpha-2)}-\sum_{k=0}^{d+1}\frac{\Gamma(\alpha+k-2)}{\Gamma(k+1)\Gamma(\alpha)}\frac{t^k}{(1+t)^{\alpha+k-2}}\right).
$$

$$
\mathbb{E}X_{1}X_{2} = C_{d} \iint_{\mathbb{R}_{d}^{+}} \frac{x_{1}x_{2}}{(1 + \max_{i} x_{i})^{\alpha + d}} d\mathbf{x}
$$
\n
$$
= C_{d} \left[ 2 \int_{x_{1} = \max_{i} x_{i}} \frac{x_{1}x_{2}}{(1 + x_{1})^{\alpha + d}} d\mathbf{x} + (d - 2) \iint_{x_{3} = \max_{i} x_{i}} \frac{x_{1}x_{2}}{(1 + x_{3})^{\alpha + d}} d\mathbf{x} \right]
$$
\n
$$
= C_{d} \left[ \int_{0}^{\infty} \frac{x_{1}^{d+1}}{(1 + x_{1})^{\alpha + d}} d\mathbf{x}_{1} + (d - 2) \int_{0}^{\infty} \frac{1}{4} \frac{x_{3}^{d+1}}{(1 + x_{3})^{\alpha + d}} d\mathbf{x}_{3} \right]
$$
\n
$$
= C_{d} \cdot \left( 1 + \frac{(d - 2)}{4} \right) \int_{0}^{\infty} \frac{x^{d+1}}{(1 + x)^{\alpha + d}} d\mathbf{x} = \frac{(d + 1)(d + 2)}{4(\alpha - 1)(\alpha - 2)}.
$$
\n
$$
\mathbb{E} (X_{1}X_{2}; \|\mathbf{X}\|_{\infty} \leq t) = C_{d} \cdot \left( 1 + \frac{(d - 2)}{4} \right) \int_{0}^{t} \frac{x_{1}^{d+1}}{(1 + x_{1})^{\alpha + d}} d\mathbf{x}_{1}
$$
\n
$$
= \frac{(d + 1)(d + 2)}{4} \left( \frac{1}{(\alpha - 1)(\alpha - 2)} - \sum_{i=0}^{d+1} \frac{\Gamma(\alpha + k - 2)}{\Gamma(k + 1)\Gamma(\alpha)} \frac{t^{k}}{(1 + t)^{\alpha + k - 2}} \right)
$$

=

4 
$$
(\alpha - 1)(\alpha - 2) \sum_{k=0}^{\infty} \Gamma(k+1)\Gamma(\alpha)
$$

 $(1+t)^{\alpha+k-2}$ 

.



### **ESSEC Business School**

3 avenue Bernard-Hirsch **CS 50105 Cergy** 95021 Cergy-Pontoise Cedex France Tél. +33 (0)1 34 43 30 00 www.essec.edu

**ESSEC Executive Education** CNIT BP 230 92053 Paris-La Défense France Tél. +33 (0)1 46 92 49 00 www.executive-education.essec.edu

**ESSEC Asia-Pacific** 5 Nepal Park Singapore 139408 Tél. +65 6884 9780 www.essec.edu/asia

ESSEC | CPE Registration number 200511927D<br>Period of registration: 30 June 2017 - 29 June 2023 Committee of Private Education (CPE) is part of SkillsFuture Singapore (SSG)

**ESSEC Afrique** Plage des Nations - Golf City Route de Kénitra - Sidi Bouknadel (Rabat-Salé) Morocco Tél. +212 (0)5 37 82 40 00 www.essec.edu

## CONTACT

#### Prénom NOMNOMNOM

Fonction fonctionr Tel. +33 (0)1 34 43 00 00 nnomnom@essec.edu







