



HAL
open science

FURTHER ENUMERATION RESULTS CONCERNING A RECENT EQUIVALENCE OF RESTRICTED INVERSION SEQUENCES

Toufik Mansour, Mark Shattuck

► **To cite this version:**

Toufik Mansour, Mark Shattuck. FURTHER ENUMERATION RESULTS CONCERNING A RECENT EQUIVALENCE OF RESTRICTED INVERSION SEQUENCES. 2021. hal-03295362v1

HAL Id: hal-03295362

<https://hal.science/hal-03295362v1>

Preprint submitted on 22 Jul 2021 (v1), last revised 31 Jan 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FURTHER ENUMERATION RESULTS CONCERNING A RECENT EQUIVALENCE OF RESTRICTED INVERSION SEQUENCES

TOUFIK MANSOUR AND MARK SHATTUCK

ABSTRACT. Let asc and desc denote respectively the statistics recording the number of ascents or descents in a sequence having non-negative integer entries. In a recent paper by Andrews and Chern, it was shown that the distribution of asc on the inversion sequence avoidance class $I_n(\geq, \neq, >)$ is the same as that of $n - 1 - \text{asc}$ on the class $I_n(>, \neq, \geq)$. In this paper, we consider some further enumerative aspects related to this equivalence. In particular, we find recurrence relations for the joint distribution on $I_n(\geq, \neq, >)$ of asc and desc along with two other parameters, and do the same for $n - 1 - \text{asc}$ and desc on $I_n(>, \neq, \geq)$. By employing a functional equation approach together with the kernel method, we are able to compute explicitly the generating function for both of the aforementioned joint distributions, which extends (and provides a new proof of) the recent result $|I_n(\geq, \neq, >)| = |I_n(>, \neq, \geq)|$. In both cases, an algorithm is formulated for computing the generating function of the asc distribution on members of each respective class having a fixed number of descents.

1. INTRODUCTION

An *inversion* within $\pi = \pi_1 \cdots \pi_n \in \mathcal{S}_n$ is an ordered pair (a, b) where $a, b \in [n] = \{1, \dots, n\}$ with $a < b$ and $\pi_a > \pi_b$. The *inversion sequence* of π is defined by $x = x_1 \cdots x_n$, where x_i for each $i \in [n]$ records the number of inversions for which $\pi_a = i$, that is, inversions caused by i when its position relative to the elements in $[i - 1]$ is decided. Note that such sequences x are characterized by the property $0 \leq x \leq i - 1$ for all i . For example, if $\pi = 451632 \in \mathcal{S}_6$, then $x = 001332$. Let I_n denote the set of all inversion sequences of length n . The systematic study of patterns in members of I_n is a topic that has only recently been initiated in [3, 11], starting with the avoidance of a single pattern of length three.

Martinez and Savage [13] extended the notion of pattern avoidance by considering a fixed triple of relations $(\rho_1, \rho_2, \rho_3) \in \{<, >, \leq, \geq, =, \neq, -\}$, where $-$ denotes the universal relation (i.e., $x - y$ for all $x, y \in [n]$). They studied the set $I_n(\rho_1, \rho_2, \rho_3)$ consisting of those $e = e_1 \cdots e_n \in I_n$ for which there exist no indices $1 \leq i < j < k \leq n$ such that $e_i \rho_1 e_j$, $e_j \rho_2 e_k$ and $e_i \rho_3 e_k$. Note that sequences in $I_n(\rho_1, \rho_2, \rho_3)$ are synonymous with members of $I_n(\{\tau_1, \dots, \tau_r\})$ for some patterns τ_1, \dots, τ_r of length three, where $I_n(\{\tau_1, \dots, \tau_r\})$ denotes the subset of I_n whose members avoid each τ_i for $i \in [r]$ in the classical sense. For example, we have $I_n(>, <, \geq) = I_n(\{101, 201\})$ and $I_n(<, -, <) = I_n(\{011, 012, 021\})$.

Since their introduction in [13], the problem of enumerating members of an avoidance class $I_n(\rho_1, \rho_2, \rho_3)$ is one that has been investigated extensively, with many connections having been made to sequences in the OEIS [14]. We refer the reader to [9] and references contained therein. Moreover, several equivalences among the 343 possible sets $I_n(\rho_1, \rho_2, \rho_3)$ were conjectured in [13] and later proven in [2, 4, 6, 8, 9, 16]. Paralleling the study of pattern avoidance on permutations represented in the one-line notation (see, e.g., [7]), analogous problems, such as avoidance of vincular [10] or multiple [15] patterns, have been considered on inversion sequences.

A further direction in the study of inversion sequences avoiding a pattern of relation triples is obtained by considering the distribution of a statistic on members of an avoidance class; see, e.g., [2, 6, 9, 12, 13]. In this paper, we consider the distribution of the number of ascents, descents and levels over certain avoidance classes of I_n involving a pattern of relation triples. Recall that

Date: July 22, 2021.

2010 Mathematics Subject Classification. 05A15, 05A05, 65Q20, 65Q30.

Key words and phrases. pattern avoidance, combinatorial statistic, kernel method, inversion sequence.

given a sequence $w = w_1 \cdots w_n$, an *ascent* is an index $i \in [n - 1]$ such that $w_i < w_{i+1}$, a *descent* is one with $w_i > w_{i+1}$ and a *level* one with $w_i = w_{i+1}$. Let $\text{asc}(w)$, $\text{desc}(w)$ and $\text{lev}(w)$ denote respectively the number of ascents, descents and levels in the sequence w . Further, recall that within a descent $w_j > w_{j+1}$ for some $j \in [n - 1]$, the entries w_j and w_{j+1} are referred to as a *descent top* and a *descent bottom*, respectively. Note that $\text{desc}(w) + \text{lev}(w) = n - 1 - \text{asc}(w)$ for all w of length n .

Lin [9] conjectured the following equivalence involving the ascents statistic on the avoidance classes $I_n(\geq, \neq, >)$ and $I_n(>, \neq, \geq)$:

$$(1) \quad \sum_{e \in I_n(\geq, \neq, >)} q^{\text{asc}(e)} = \sum_{e \in I_n(>, \neq, \geq)} q^{n-1-\text{asc}(e)}, \quad n \geq 1.$$

This equivalence was shown by Andrews and Chern [1] using a functional equation approach. Here, we consider some further combinatorial aspects of (1). In particular, we consider a refinement of both sides of (1) by introducing a variable p which marks the number of descents in members of each class. We compute an explicit formula for the generating function of the joint distribution of desc and asc on $I_n(\geq, \neq, >)$, and also of desc and $n - 1 - \text{asc}$ on $I_n(>, \neq, \geq)$, using the kernel method [5].

Comparing the $p = 1$ cases of our main results below (see Theorems 4 and 13), in addition to providing a new proof of (1), yields a formula for the generating function of both sides of (1). Such a formula was not given explicitly in [1]. Further, taking $p = q = 1$ yields a new proof of the fact first shown by Lin [9] that

$$(2) \quad |I_n(\geq, \neq, >)| = |I_n(>, \neq, \geq)| = |\mathcal{S}_n(4231, 42513)|, \quad n \geq 1,$$

which confirmed a conjecture made originally in [13]. Note that the sequence of cardinalities $|\mathcal{S}_n(4231, 42513)|$ for $n \geq 1$ corresponding to the $q = 1$ case of (1) occurs as entry A098746 in [14]. To show (2), note that Lin did not enumerate either $I_n(\geq, \neq, >)$ or $I_n(>, \neq, \geq)$ directly, but rather defined a bijection between the two and enumerated instead the set $I_n(>, -, >)$, which had been shown in [13] to be equinumerous with $I_n(\geq, \neq, >)$ via a bijection. Thus, the method employed here allows for a unified proof of (1) and (2), which were shown previously by seemingly unrelated approaches, and leads to a refinement of these relations in terms of the descents statistic (and other parameters).

The organization of this paper is as follows. In the next section, we consider the joint distribution of desc and asc on $I_n(\geq, \neq, >)$ and find its generating function. To do so, a recurrence is found for a refinement of this distribution involving two additional parameters which is then converted to a system of functional equations involving the corresponding generating functions. Further, an algorithm is devised for finding the generating function for the distribution of asc on members of $I_n(\geq, \neq, >)$ having a prescribed number of descents. In the third section, we consider the distribution of desc and $n - 1 - \text{asc}$ on $I_n(>, \neq, \geq)$ and compute the generating function of this distribution. Comparable formulas are found and an algorithm is given for determining explicitly the coefficient of p^m for a fixed m in this generating function.

We remark further that in order to obtain recurrences for the joint distribution of $(\text{desc}, \text{asc})$ on $I_n(\geq, \neq, >)$ and of $(\text{desc}, n - 1 - \text{asc})$ on $I_n(>, \neq, \geq)$, we must refine these sets according to a pair of new parameters on each. In particular, to study members $e \in I_n(\geq, \neq, >)$, it is useful to consider a parameter which we call the *height* of e and is defined as the maximum letter that starts either a descent or a level of e . On the other hand, for $e \in I_n(>, \neq, \geq)$, it is convenient to consider the statistic which tracks the smallest letter serving as a descent top for the largest descent bottom in e . These parameters (and variants thereof) may very well prove interesting to study in their own right on other types of discrete structures that are often represented as sequences, such as finite set partitions or functions between two finite sets of prescribed size.

2. THE DESCENTS AND ASCENTS STATISTICS ON $I_n(\geq, \neq, >)$.

We first define two new concepts related to the relative sizes of the non-ascent entries within an inversion sequence. Let the *height* of $e = e_1e_2 \cdots e_n \in I_n$ be given by

$$\text{hgt}(e) = \max\{e_i : 1 \leq i \leq n-1 \text{ and } e_i \geq e_{i+1}\}.$$

If $a = \text{hgt}(e)$ with $j \in [n-1]$ minimal such that $e_j = a$, then let the *depth* of e be defined as $\text{dep}(e) = e_{j+1}$. Here, and in the subsequent section, we find it more convenient notationally to represent members of I_n using positive instead of non-negative integers, which is achieved by adding one to each entry of the standard representation.

Let $\mathcal{A}_n = I_n(\geq, \neq, >)$ and suppose $e \in \mathcal{A}_n$ has height and depth values of a and b , respectively. If $a > b$, then there exists a single descent ab and at most two runs of the letter a , the first of which has length one. On the other hand, if $a = b$ within e , there can exist only a single run of a . Within a (maximal) subsequence of the form $ab \cdots b$, any letter beyond the second will be referred to as a *redundant bottom*, regardless of whether or not a and b are distinct. For example, if $e = 123116333669 \in \mathcal{A}_{12}$, then $\text{hgt}(e) = 6$ and $\text{dep}(e) = 3$, with the last two 3's redundant bottoms. If $e = 1132267779 \in \mathcal{A}_{10}$, then $\text{hgt}(e) = \text{dep}(e) = 7$, with only the third 7 a redundant bottom.

We now decompose \mathcal{A}_n into disjoint subsets as follows. Given $1 \leq i \leq n-1$ and $1 \leq j \leq n$, let $\mathcal{B}_n(i, j)$ denote the subset of \mathcal{A}_n whose members have height i and last letter j , where the last letter is not a redundant bottom. Let $\mathcal{C}_n(i, j)$ be defined the same as $\mathcal{B}_n(i, j)$, but where the last letter is a redundant bottom. Note that $\mathcal{C}_n(i, j)$ can be nonempty only when $n \geq 3$ and $1 \leq j \leq i \leq n-2$. Define the distribution polynomial $b_n(i, j) = b_n(i, j; p, q)$ by

$$b_n(i, j) = \sum_{\pi \in \mathcal{B}_n(i, j)} p^{\text{desc}(\pi)} q^{\text{asc}(\pi)},$$

and likewise for $c_n(i, j) = c_n(i, j; p, q)$. For example, we have

$$\mathcal{B}_5(3, 4) = \{11314, 11324, 11334, 12314, 12324, 12334\}$$

and $\mathcal{C}_5(3, 2) = \{11322, 12322\}$, which implies $b_5(3, 4) = q^2 + q^3 + 2pq^2 + 2pq^3$ and $c_5(3, 2) = pq + pq^2$. Assume $b_n(i, j)$ or $c_n(i, j)$ to be zero if the subset of \mathcal{A}_n corresponding to i and j is empty.

Let

$$b_n = \sum_{i=1}^{n-1} \sum_{j=1}^n b_n(i, j), \quad n \geq 2,$$

and

$$c_n = \sum_{i=1}^{n-2} \sum_{j=1}^i c_n(i, j), \quad n \geq 3,$$

and put $b_1 = 0$ and $c_1 = c_2 = 0$. Note that b_n and c_n are polynomials in p and q . Then we seek a formula for $a_n = a_n(p, q)$ defined as

$$a_n = b_n + c_n + q^{n-1}, \quad n \geq 1.$$

Note that a_n gives the joint distribution of desc and asc on \mathcal{A}_n , where the q^{n-1} term accounts for the sequence $12 \cdots n$ which belongs to no subset $\mathcal{B}_n(i, j)$ or $\mathcal{C}_n(i, j)$.

The arrays $b_n(i, j)$ and $c_n(i, j)$ satisfy the following system of recurrences.

Lemma 1. *We have*

$$\begin{aligned} b_n(i, j) &= \delta_{i, n-2} \cdot q^{n-2} + qc_{n-1}(i, i) + q \sum_{\ell=i+1}^{j-1} b_{n-1}(i, \ell) + q \sum_{k=1}^{i-1} b_{n-2}(k, i) + q \sum_{\ell=1}^{i-1} c_n(i, \ell) \\ (3) \quad &+ q^2 \sum_{\ell=1}^{i-1} \sum_{s=1}^{n-i-2} c_{n-s}(i, \ell), \quad 1 \leq i \leq n-2 \text{ and } i < j \leq n, \end{aligned}$$

with $b_n(n-1, n) = 0$ for $n \geq 2$,

$$(4) \quad b_n(i, i) = \delta_{i, n-1} \cdot q^{n-2} + \sum_{k=1}^{i-1} b_{n-1}(k, i) + q \sum_{\ell=1}^{i-1} \sum_{s=1}^{n-i-1} c_{n-s+1}(i, \ell), \quad 1 \leq i \leq n-1,$$

$$(5) \quad b_n(i, j) = \delta_{i, n-1} \cdot pq^{n-2} + p \sum_{k=1}^j b_{n-1}(k, i) + p \sum_{k=j+1}^{i-1} \sum_{s=0}^{i-k-1} \binom{i-k-1}{s} q^{s+1} c_{n-s-1}(k, j),$$

for $1 \leq j < i \leq n-1$,

$$(6) \quad c_n(i, j) = b_{n-1}(i, j) + c_{n-1}(i, j), \quad 1 \leq j < i \leq n-2,$$

$$(7) \quad c_n(i, i) = \delta_{i, n-2} \cdot q^{n-3} + c_{n-1}(i, i) + \sum_{k=1}^{i-1} b_{n-2}(k, i), \quad 1 \leq i \leq n-2.$$

Furthermore, we have the following alternative recurrences to (3) and (5) above when $i < j$ and $i > j$, respectively:

$$(8) \quad b_n(i, j) = \sum_{\ell=1}^i \sum_{t=0}^{j-i-1} \binom{j-i-1}{t} q^{t+1} c_{n-t}(i, \ell) + \sum_{\ell=1}^{i-1} \sum_{t=0}^{j-i-1} \sum_{s=1}^{n-i-t-2} \binom{j-i-1}{t} q^{t+2} c_{n-s-t}(i, \ell)$$

and

$$(9) \quad \begin{aligned} \frac{1}{pq} (b_n(i, j) - q \sum_{k=j+1}^{i-1} b_{n-1}(k, j)) &= \delta_{j, n-2} \cdot q^{n-3} + \sum_{k=j+1}^{i-1} c_{n-1}(k, j) + \sum_{k=1}^j \sum_{\ell=1}^k c_{n-1}(k, \ell) \\ &+ \sum_{\ell=2}^j \sum_{k=1}^{\ell-1} b_{n-2}(k, \ell) + q \sum_{k=2}^j \sum_{\ell=1}^{k-1} \sum_{r=1}^{n-k-2} c_{n-r-1}(k, \ell). \end{aligned}$$

Proof. To show (3), let $\pi \in \mathcal{B}_n(i, j)$ where $j > i$. Note that $1 \leq i \leq n-2$ is required since the height cannot be achieved for the first time by the penultimate letter if $j > i$. Suppose that π has depth ℓ for some $\ell \in [i]$. Then it may be verified that π can be decomposed uniquely as

$$\pi = \pi' i \ell^r i^s \alpha,$$

where $s \geq 0$, α is a (strictly) increasing sequence in $[i+1, j]$ ending in j and π' contains only letters in $[i-1]$, with $r \geq 1$ if $\ell < i$ and $r = 1$ if $\ell = i$. To see this, note that π' cannot contain any letters greater than i , for otherwise i would fail to be the height of π . Further, if $\ell < i$, then π' cannot contain i either, for then $(\geq, \neq, >)$ would be realized by $i i \ell$. On the other hand, if $\ell = i$, one need only consider the leftmost occurrence of i to obtain the stated decomposition.

If $|\alpha| > 1$, then the weight of all possible such π is given by $q \sum_{\ell=i+1}^{j-1} b_{n-1}(i, \ell)$, upon deleting j and considering the penultimate letter which belongs to $[i+1, j-1]$ in this case. So assume α consists of only the terminal j , i.e., $\pi = \pi' i \ell^r i^s j$, where ℓ, r and s are as before. We consider then the following cases on s and ℓ : (i) $s = 0, \ell = i$; (ii) $s \geq 1, \ell = i$; (iii) $s = 0, \ell < i$; (iv) $s \geq 1, \ell < i$. One may verify that the respective contributions towards $b_n(i, j)$ in the four cases are given by (i) $\delta_{i, n-2} \cdot q^{n-2} + q \sum_{k=1}^{i-1} b_{n-2}(k, i)$, (ii) $q c_{n-1}(i, i)$, (iii) $q \sum_{\ell=1}^{i-1} (b_{n-1}(i, \ell) + c_{n-1}(i, \ell))$ and (iv) $q^2 \sum_{\ell=1}^{i-1} \sum_{s=1}^{n-i-2} (b_{n-s-1}(i, \ell) + c_{n-s-1}(i, \ell))$. Note that in (iii), deletion of the final j results in an arbitrary member of $\mathcal{B}_{n-1}(i, \ell) \cup \mathcal{C}_{n-1}(i, \ell)$ for some $\ell \in [i-1]$, upon considering whether $r = 1$ or $r > 1$ in the decomposition of π above, where the factor of q accounts for the ascent arising due to j . In (iv), on the other hand, there are two extra ascents that arise since $j > i > \ell$ with $s \geq 1$ so that deletion of i^s along with the terminal j from π results in a sequence enumerated by $b_{n-s-1}(i, \ell) + c_{n-s-1}(i, \ell)$. Now observe that (6) follows from the definitions, upon removing the final letter which must be equal to its predecessor in this case. Formula (3) then follows from (6) and combining all the previous cases.

To show (4), let $\pi \in \mathcal{B}_n(i, i)$ where $1 \leq i \leq n-1$. First suppose that π ends in i, i with no other i 's occurring in π . If $i < n-1$, then deleting the final letter results in a member of $\mathcal{B}_{n-1}(k, i)$ for some $k \in [i-1]$, whereas if $i = n-1$, then in addition one can have $\pi = 12 \cdots (n-1)(n-1)$,

which is accounted for by the extra term $\delta_{i,n-1} \cdot q^{n-2}$. So assume it is not the case that π ends in i, i with no others i 's occurring. Note that it is not possible for π to have a single run of i 's which occurs at the end and has length three or more, for that would imply that the terminal i is a redundant bottom with such π not being enumerated by $b_n(i, i)$. Then we must have $\pi = \pi' i \ell^r i^s$, where π' has letters only in $[i-1]$, $r, s \geq 1$ and $\ell \in [i-1]$. Note that the first i occurring in the j -th position for some $j \geq i$, together with $r \geq 1$, implies $s \leq n-i-1$. Then deletion of i^s from π results in members of $\mathcal{B}_n(i, i)$ whose weight is given by the second sum in (4), upon applying (6). Combining the previous cases now yields (4).

To show (5), let $\pi \in \mathcal{B}_n(i, j)$ where $2 \leq i \leq n-1$ and $j \in [i-1]$. Then j not a redundant bottom implies we must have $\pi = \pi' i j$ where $\max(\pi') < i$. If $i = n-1$, then $\pi' = 12 \cdots (n-2)$ is possible, which is accounted for by $\delta_{i,n-1} \cdot pq^{n-2}$, so assume π' has height k for some $k \in [n-2]$. Note that $k \geq i$ is impossible, for otherwise kij would correspond to an occurrence of $(\geq, \neq, >)$. If $k \in [j]$, then the terminal j may be deleted yielding a $p \sum_{k=1}^j b_{n-1}(k, i)$ possibilities, where the factor of p accounts for the descent between i and j . So assume $k \in [j+1, i-1]$. Then π must have the form $\pi = \alpha k j^r \beta j$, where $\max(\alpha) < k$, $r \geq 1$ and β is increasing on $[k+1, i]$ with last letter i . If $|\beta| = s+1$, then there are $\binom{i-k-1}{s}$ ways in which to choose the members of β . Deleting β from π (keeping the terminal j) yields a member of $\mathcal{C}_{n-s-1}(k, j)$, as the resulting sequence would end in a redundant (descent) bottom in this case. Considering all possible k and s then gives the weight of the remaining members of $\mathcal{B}_n(i, j)$ and implies (5), where the pq^{s+1} factor in the final sum accounts for the ascents caused by the members of β and the descent between the last two letters.

To show (7), first note that $\pi \in \mathcal{C}_n(i, i)$ must end in a run of i 's of length at least three with all other letters outside of this run strictly less than i . Upon deleting the final i , there are $c_{n-1}(i, i)$ possibilities if π ends in four or more i 's. Otherwise, $\pi = \pi' i i$, where $\pi' \in \mathcal{B}_{n-2}(k, i)$ for some $k < i$ or $\pi' = 12 \cdots (n-2)$, the latter applying only if $i = n-2$, which implies (7). To show (8), consider the same four cases (i)–(iv) used in the proof of (1) above, but where the single terminal j is replaced with an increasing sequence β in $[i+1, j]$ whose last letter is j . Then the first sum on the right side of (8) gives the contribution towards the overall weight of those members of $\mathcal{B}_n(i, j)$ in which cases (i)–(iii) apply. This is seen upon deleting all letters in β where $|\beta| = t+1$ and adding back an extra copy of the letter ℓ to the end, which results in a member of $\mathcal{C}_{n-t}(i, \ell)$. To find the weight of members of $\mathcal{B}_n(i, j)$ in case (iv), first delete from π both β and i^s (which directly precedes β) and then add back a letter ℓ to the end. Note that the resulting sequence belongs to $\mathcal{C}_{n-s-t}(i, \ell)$ for some $\ell < i$, with $i \leq n-s-t-2$ implying $s \in [n-i-t-2]$. Considering all possible ℓ, t and s then gives the second sum on the right side of (8) and completes the proof.

To show (9), suppose $\pi \in \mathcal{B}_n(i, j)$ where $i > j$. Then we may write $\pi = \rho i j$, where $\max(\rho) < i$, and consider cases based on the last letter t of ρ . If $t > j$, then deleting i from π results in a member of $\mathcal{B}_{n-1}(t, j)$ and considering all possible t yields a contribution of $q \sum_{t=j+1}^{i-1} b_{n-1}(t, j)$ towards the weight in this case. Now suppose $t = j$. If $\rho = 12 \cdots (n-2)$, then $i = n-1, j = n-2$ and π has weight pq^{n-2} . So assume $\rho \neq 12 \cdots (n-2)$ and let $\text{hgt}(\rho) = k$. If $k < j$, then deleting i and the terminal j implies a contribution of $pq \sum_{k=1}^{j-1} b_{n-2}(k, j)$ towards the weight. If $k = j$, then either $\rho = \rho' j^{r+1}$ or $\rho = \rho' j \ell^s j^r$, where $r, s \geq 1, \ell \in [j-1]$ and $\max(\rho') < j$. Removing i from π in the first case and removing the terminal section $j^r i j$ from π and adding back a copy of ℓ in the second (so that the resulting sequence is $\rho' j \ell^{s+1}$) yields respective contributions of $pq c_{n-1}(j, j)$ and $pq^2 \sum_{\ell=1}^{j-1} \sum_{r=1}^{n-j-2} c_{n-r-1}(j, \ell)$, upon considering all possible ℓ and r in the latter case. Note that $r, s \geq 1$ and $\ell < j$ implies two ascents (and a descent) are lost with the removal of $j^r i j$ from π . The remaining possibility when $t = j$ is for $j+1 \leq k \leq i-1$, in which case $\rho = \rho' k j^r$, where $\max(\rho') < k$ and $r \geq 1$. Deleting i and considering all k yields $pq \sum_{k=j+1}^{i-1} c_{n-1}(k, j)$, as the resulting sequence ends in $k j^{r+1}$ where $k > j$.

Now assume $t < j$, and again let $k = \text{hgt}(\rho)$. Note that $k > j$ is not possible in this case, for otherwise ktj is an occurrence of $(\geq, \neq, >)$. If $k = j$, then $\rho = \rho' j \ell^r$, where $\ell \in [j-1]$ and $r \geq 1$, and thus there are $pq \sum_{\ell=1}^{j-1} c_{n-1}(j, \ell)$ possibilities, upon deleting i and the terminal j and putting back an ℓ . So assume $k < j$. Considering whether $t \leq k$ or $k < t < j$ implies ρ must have one of the following four forms: (a) $\rho = \rho' k \ell^s k^r$, (b) $\rho = \rho' k^r + 2$, (c) $\rho = \rho' k \ell^s k^r \alpha$ or (d) $\rho = \rho' k^{r+2} \alpha$,

where $r \geq 0$, $s \geq 1$, $\ell \in [k-1]$, $\max(\rho') < k$ and α is a nonempty (strictly) increasing sequence in $[k+1, j-1]$. Deleting i and j from π in (b) or in the $r = 0$ case of (a), and adding a copy of the current final letter, yields a contribution of $pq \sum_{k=1}^{j-1} \sum_{\ell=1}^k c_{n-1}(k, \ell)$, upon considering all possible k and ℓ . If $r \geq 1$ in (a), then removal of the final $r+2$ letters of π , followed by adding an ℓ , yields $pq^2 \sum_{k=2}^{j-1} \sum_{\ell=1}^{k-1} \sum_{r=1}^{n-k-2} c_{n-r-1}(k, \ell)$, as the resulting sequence is of the form $\rho' k \ell^{s+1}$ where $s \geq 1$. Further, it is seen that cases (c) and (d) combine to yield $pq \sum_{k=1}^{j-2} \sum_{\ell=k+1}^{j-1} b_{n-2}(k, \ell)$, where t denotes the last letter of α .

Finally, combining the ten contributions towards the weight coming from all the cases above, and observing the simplifications

$$c_{n-1}(j, j) + \sum_{\ell=1}^{j-1} c_{n-1}(j, \ell) + \sum_{k=1}^{j-1} \sum_{\ell=1}^k c_{n-1}(k, \ell) = \sum_{k=1}^j \sum_{\ell=1}^k c_{n-1}(k, \ell),$$

$$\sum_{k=1}^{j-1} b_{n-2}(k, j) + \sum_{k=1}^{j-2} \sum_{\ell=k+1}^{j-1} b_{n-2}(k, \ell) = \sum_{k=1}^{j-1} b_{n-2}(k, j) + \sum_{\ell=2}^{j-1} \sum_{k=1}^{\ell-1} b_{n-2}(k, \ell) = \sum_{\ell=2}^j \sum_{k=1}^{\ell-1} b_{n-2}(k, \ell)$$

and

$$\sum_{\ell=1}^{j-1} \sum_{r=1}^{n-j-2} c_{n-r-1}(j, \ell) + \sum_{k=2}^{j-1} \sum_{\ell=1}^{k-1} \sum_{r=1}^{n-k-2} c_{n-r-1}(k, \ell) = \sum_{k=2}^j \sum_{\ell=1}^{k-1} \sum_{r=1}^{n-k-2} c_{n-r-1}(k, \ell),$$

yields (9) and completes the proof. \square

From the recurrences in the prior lemma, we have that the nonzero values of $b_n(i, j)$ and $c_n(i, j)$ are given for $n = 2$ by $b_2(1, 1) = 1$, for $n = 3$ by $b_3(1, 2) = b_3(1, 3) = b_3(2, 2) = q$, $b_3(2, 1) = pq$, $c_3(1, 1) = 1$ and for $n = 4$ by

$$\begin{array}{llll} b_4(1, 2) = q & b_4(1, 3) = q + q^2 & b_4(1, 4) = q + 2q^2 & b_4(2, 1) = pq \\ b_4(2, 2) = q + pq^2 & b_4(2, 3) = q^2 + pq^2 & b_4(2, 4) = q^2 + pq^2 & b_4(3, 1) = pq + pq^2 \\ b_4(3, 2) = pq + pq^2 & b_4(3, 3) = q + q^2 & c_4(1, 1) = 1 & c_4(2, 1) = pq \\ c_4(2, 2) = q, & & & \end{array}$$

which may be verified directly using the definitions.

Define now the following generating functions: $A(x) = \sum_{n \geq 1} a_n x^n$, $B(x) = \sum_{n \geq 2} b_n x^n$ and $C(x) = \sum_{n \geq 3} c_n x^n$. Then clearly,

$$A(x) = B(x) + C(x) + \frac{x}{1-xq}.$$

Note that $A(x)$ is the (ordinary) generating function for the joint distribution of desc and asc on $I_n(\geq, \neq, >)$ for $n \geq 1$. In order to study $B(x)$ and $C(x)$, we refine them as follows. Define

$$B_0(x, v) = \sum_{n \geq 2} \sum_{i=1}^{n-1} b_n(i, i) v^i x^n,$$

$$B^+(x, v, w) = \sum_{n \geq 2} \sum_{i=1}^{n-1} \sum_{j=i+1}^n b_n(i, j) v^i w^j x^n,$$

$$B^-(x, v, w) = \sum_{n \geq 3} \sum_{i=2}^{n-1} \sum_{j=1}^{i-1} b_n(i, j) v^i w^j x^n,$$

with $C_0(x, v) = \sum_{n \geq 3} \sum_{i=1}^{n-2} c_n(i, i) v^i x^n$ and $C^-(x, v, w) = \sum_{n \geq 4} \sum_{i=2}^{n-2} \sum_{j=1}^{i-1} c_n(i, j) v^i w^j x^n$.

Translating (3)–(9) above in terms of these generating functions yields the following system of functional equations.

Lemma 2. We have $B(x) = B_0(x, 1) + B^+(x, 1, 1) + B^-(x, 1, 1)$ and $C(x) = C_0(x, 1) + C^-(x, 1, 1)$, where

$$\begin{aligned}
 B^+(x, v, w) &= \frac{qvw^2x^3(w+1)}{1-qvwx} + \frac{qwx}{1-w}(C_0(x, vw) - wC_0(wx, v)) \\
 &+ \frac{qwx}{1-w}(B^+(x, v, w) - wB^+(wx, v, 1)) + \frac{qwx^2}{1-w}(B^+(x, 1, vw) - w^2B^+(wx, 1, v)) \\
 &+ \frac{qw}{1-w}(C^-(x, vw, 1) - C^-(wx, v, 1)) \\
 &+ \frac{q^2w}{1-w} \left(\frac{x}{1-x}C^-(x, vw, 1) - \frac{wx}{1-wx}C^-(wx, v, 1) \right), \\
 B_0(x, v) &= \frac{vx^2}{1-qvx} + xB^+(x, 1, v) + \frac{q}{1-x}C^-(x, v, 1), \\
 B^-(x, v, w) &= \frac{pqv^2wx^3}{(1-qvx)(1-qvwx)} + \frac{px}{1-w}(B^+(x, w, v) - B^+(x, 1, vw)) \\
 &+ \frac{pqvx}{1-v-qvx} \left(C^-(x, v, w) - C^-\left(\frac{vx}{1-qvx}, 1-qvx, w\right) \right), \\
 C^-(x, v, w) &= \frac{x}{1-x}B^-(x, v, w), \\
 C_0(x, v) &= \frac{vx^3}{(1-x)(1-qvx)} + \frac{x^2}{1-x}B^+(x, 1, v), \\
 B^+(x, v, w) &= \frac{qw}{1-w-qwx} \left(C_0(x, vw) - C_0\left(\frac{wx}{1-qwx}, v(1-qwx)\right) \right) \\
 &+ \frac{qw}{1-w-qwx} \left(C^-(x, vw, 1) - C^-\left(\frac{wx}{1-qwx}, v(1-qwx), 1\right) \right) \\
 &+ \frac{q^2wx}{1-w-qwx} \left(\frac{1}{1-x}C^-(x, vw, 1) - \frac{w}{1-wx-qwx}C^-\left(\frac{wx}{1-qwx}, v(1-qwx), 1\right) \right)
 \end{aligned}$$

and

$$\begin{aligned}
 &\frac{1}{pq}B^-(x, v, w) \\
 &= \frac{vx}{p(1-v)}(B^-(x, v, w) - B^-(vx, 1, w)) + \frac{v^2wx^3}{1-qvwx} + \frac{vx}{1-v}(C^-(x, v, w) - C^-(vx, 1, w)) \\
 &+ \frac{x}{1-w} \left(\frac{v}{1-v}(C^-(x, vw, 1) + C_0(x, vw) - C^-(vx, w, 1) - C_0(vx, w)) \right. \\
 &\left. - \frac{vw}{1-vw}(C^-(x, vw, 1) + C_0(x, vw) - C^-(vwx, 1, 1) - C_0(vwx, 1)) \right) \\
 &+ \frac{x^2}{1-w} \left(\frac{v}{1-v}(B^+(x, 1, vw) - vB^+(vx, 1, w)) - \frac{vw}{1-vw}(B^+(x, 1, vw) - vwB^+(vwx, 1, 1)) \right) \\
 &+ qx^2 \left(\frac{v}{(1-v)(1-vw)(1-x)}C^-(x, vw, 1) - \frac{v^2w}{(1-v)(1-vw)(1-vwx)}C^-(vwx, 1, 1) \right. \\
 &\left. - \frac{v^2}{(1-v)(1-w)(1-vx)}C^-(vx, w, 1) + \frac{v^2w}{(1-v)(1-w)(1-vwx)}C^-(vwx, 1, 1) \right).
 \end{aligned}$$

By use of the first, fourth and fifth equations in Lemma 2 with $v = 1$, we find a formula for $B^+(x, 1, w)$:

$$\begin{aligned}
 B^+(x, 1, w) &= \frac{qwx(qx - x + 1)}{(qwx - wx + w + x - 1)(1-x)}B^-(x, w, 1) \\
 &+ \frac{qw^2x(1-x)(qwx - wx + 1)}{(qwx - wx + w + x - 1)(1-wx)^2}B^-(wx, 1, 1)
 \end{aligned}$$

$$\begin{aligned}
& + \frac{qw^2x(1-x)}{(qwx-wx+w+x-1)(1-wx)}B^+(wx, 1, 1) \\
& + \frac{qw^2x^3(1-w)(wx-w-1)}{(qwx-wx+w+x-1)(1-qwx)(1-wx)}.
\end{aligned}$$

Using this last expression twice (both for $B^+(x, 1, vw)$ and $B^+(vx, 1, w)$), substituting into the final equation in Lemma 2 and replacing x with x/v , we obtain the following result.

Lemma 3. *The generating function $A(x)$ is given by*

$$A(x) = \frac{1}{1-x}B^+(x, 1, 1) + \frac{1-x+qx}{(1-x)^2}B^-(x, 1, 1) + \frac{x}{(1-x)(1-qx)},$$

where

$$K(x, v)B^-(x/v, 1, v) = A_1(x, v)B^-(x, 1, 1) + A_2(x, v)B^+(x, 1, 1) + A_3(x, v),$$

with

$$\begin{aligned}
K(x, v) &= (x-v)(qx+v-1)(qvx+v^2-vx-v+x) - qx^2(qvx-qx+v^2-vx-2v+2x)p, \\
A_1(x, v) &= (x-v)(qvx+v^2-vx-v+x)xq \\
&+ \frac{(v-x)x^2((qv-q-v+2)x^2 + (v^2-v-2)x - v^2 + 2v)qp}{(1-x)^2}, \\
A_2(x, v) &= \frac{(x-v)(1-v)(qx^2+vx-x^2-v+x)xp}{1-x}, \\
A_3(x, v) &= \frac{(x-v)(qx^2-qx+vx-x^2-v+x+1)(1-v)pqx^3}{(1-x)(1-qx)}.
\end{aligned}$$

Note that the kernel equation $K(x, v) = 0$ (see prior lemma) has two power series solutions $v_1(x)$ and $v_2(x)$, where

$$\begin{aligned}
v_1(x) &= 1 + (\sqrt{pq} - q)x - \frac{q((2p+1)\sqrt{pq} - 2pq - p)}{2\sqrt{pq}}x^2 \\
&- \frac{q(4(5pq + 3p + q + 1)\sqrt{pq} - 12p^2q - 8pq^2 - 16pq - 3p - q)}{8\sqrt{pq}}x^3 + \dots, \\
v_2(x) &= 1 - (\sqrt{pq} + q)x - \frac{q((2p+1)\sqrt{pq} + 2pq + p)}{2\sqrt{pq}}x^2 \\
&- \frac{q(4(5pq + 3p + q + 1)\sqrt{pq} + 12p^2q + 8pq^2 + 16pq + 3p + q)}{8\sqrt{pq}}x^3 + \dots.
\end{aligned}$$

Substituting $v = v_1(x)$ and $v = v_2(x)$ into

$$K(x, v)B^-(x/v, 1, v) = A_1(x, v)B^-(x, 1, 1) + A_2(x, v)B^+(x, 1, 1) + A_3(x, v),$$

and solving for $B^-(x, 1, 1)$ and $B^+(x, 1, 1)$, we obtain

$$\begin{aligned}
B^-(x, 1, 1) &= \frac{px(v_2(x) - 1)(v_1(x) - 1)}{((q-1)x^2 - v_1(x)v_2(x) + xv_1(x) + xv_2(x))(p-q)}, \\
B^+(x, 1, 1) &= \frac{qx(qx-1)(qx-x+1)(v_2(x)-1)(v_1(x)-1)}{(1-x)(1-qx)((q-1)x^2 - v_1(x)v_2(x) + xv_1(x) + xv_2(x))(p-q)} \\
&+ \frac{qx^2((q-1)x(v_1(x)v_2(x) + x^2) + (x^2 - x + 1 - qx)(v_1(x) + v_2(x)) + (1-2q)x^2 + (1+2q)x - 2)}{(1-x)(1-qx)((q-1)x^2 - v_1(x)v_2(x) + xv_1(x) + xv_2(x))}.
\end{aligned}$$

Hence, by Lemma 3, we have the following result.

Theorem 4. *The generating function $A(x)$ is given by*

$$A(x) = (1 - v_1(x) - v_2(x) - (q-1)x)x(q-1)x^2 - v_1(x)v_2(x) + x(v_1(x) + v_2(x)),$$

where $v_1(x)$ and $v_2(x)$ are the power series solutions to $K(x, v) = 0$ as defined above.

2.1. **Case $p = q = 1$.** Letting $p = q = 1$ in Lemma 3 gives

$$K'(x, v)B^-(x/v, 1, v) = A'(x, v)B^-(x, 1, 1) + (1-x)A'(x, v)B^+(x, 1, 1) + A''(x, v),$$

where

$$\begin{aligned} K'(x, v) &= v^3 - v^2 + 2vx - x^2, \\ A'(x, v) &= \frac{x(v-x)(vx-v+x)}{(1-x)^2}, \\ A''(x, v) &= \frac{x^3(v-x)(vx-v+1)}{(1-x)^3}. \end{aligned}$$

Thus, by taking $v = 1$ or

$$v = v_3(x) = 1 + \frac{16}{9} \sin^4\left(\frac{1}{3} \arcsin\left(\frac{3\sqrt{3x}}{2}\right)\right) - \frac{8}{3} \sin^2\left(\frac{1}{3} \arcsin\left(\frac{3\sqrt{3x}}{2}\right)\right) = 1 - 2x - 3x^2 - 10x^3 + \dots,$$

the latter being a root of $K'(x, v) = 0$, we obtain

$$\begin{aligned} (2x-1)(x-1)B^+(x, 1, 1) - (x^2-x+1)B^-(x, 1, 1) - x^3 &= 0, \\ \frac{xv_3(x) - v_3(x) + x}{(1-x)^2}B^-(x, 1, 1) + \frac{xv_3(x) - v_3(x) + x}{1-x}B^+(x, 1, 1) + \frac{x^2(xv_3(x) - v_3(x) + 1)}{(1-x)^3} &= 0. \end{aligned}$$

Theorem 5. Let $y = \cos(\frac{1}{3} \arcsin(\frac{3\sqrt{3x}}{2}))$. Then the generating function $\sum_{n \geq 1} a_n(1, 1)x^n$ is given by

$$-\frac{(4y^2-1)^2-9x}{2A} \left(B + (64y^6 - 96y^4 + 36y^2 + 27x - 4)(64y^6 - 12y^2 - 27x + 2)\sqrt{1-4x} \right),$$

where $A = (256y^8 - 256y^6 - 48y^4 + 56y^2 + 81x - 8)(x(4y^2 - 1)^4 + (x-1)((4y^2 - 1)^2 - 9x)^2)$ and $B = (4y^2 - 1)^6 - 9(32y^4x + 16y^4 - 16y^2x - 8y^2 - 7x + 1)((4y^2 - 1)^2 - 9x)$.

2.2. **Coefficient of p^m .** Note that Lemma 2 implies the following formula.

Lemma 6. The generating function $A(x)$ is given by

$$A(x) = \frac{1}{1-x}B^+(x, 1, 1) + \frac{1-x+qx}{(1-x)^2}B^-(x, 1, 1) + \frac{x}{(1-x)(1-qx)},$$

where

$$\begin{aligned} \left(1 - \frac{pqvx^2}{(1-x)(1-v-qvx)}\right)B^-(x, v, w) &= \frac{pqv^2wx^3}{(1-qvwx)(1-qvx)} + \frac{px(B^+(x, w, v) - B^+(x, 1, vw))}{1-w} \\ - \frac{pqv^2x^2}{(1-v-qvx)(1-vx-qvx)}B^-\left(\frac{vx}{1-qvx}, 1-qvx, w\right) & \end{aligned}$$

and

$$\begin{aligned} (1-w-qwx)B^+(x, v, w) &= \frac{qvw^2x^3(1+w-wx)(1-w)}{(1-x)(1-wx)(1-qvwx)} + \frac{qwx^2}{1-x}B^+(x, 1, vw) \\ - \frac{qw^3x^2}{1-wx}B^+(wx, 1, v) - qw^2xB^+(wx, v, 1) + \frac{qwx(1-x+qx)}{(1-x)^2}B^-(x, vw, 1) & \\ - \frac{qw^2x(1-wx+qwx)}{(1-wx)^2}B^-(wx, v, 1), & \end{aligned}$$

with

$$\begin{aligned} B^+(x, v, 1) &= \frac{x}{(1-qx)(1-x-qx)}B^+\left(\frac{x}{1-qx}, 1, v(1-qx)\right) - \frac{x}{1-x}B^+(x, 1, v) \\ &\quad - \frac{1-x+qx}{(1-x)^2}B^-(x, v, 1) + \frac{1-x}{(1-x-qx)^2}B^-\left(\frac{x}{1-qx}, v(1-qx), 1\right) \\ &\quad - \frac{(qx+x-2)qvx^3}{(1-x-qx)(1-qx)(1-x)(1-qvx)}. \end{aligned}$$

Given a generating function $f(x) = f(x, p)$, we denote its coefficient of p^m by $f(x|m)$. From Lemma 6, one obtains the following result.

Proposition 7. *We have*

$$A(x|m) = \frac{1}{1-x}B^+(x, 1, 1|m) + \frac{1-x+qx}{(1-x)^2}B^-(x, 1, 1|m) + \frac{x}{(1-x)(1-qx)}\delta_{m,0},$$

where $B^+(x, v, w|m)$ and $B^-(x, v, w|m)$ satisfy

$$\begin{aligned} B^-(x, v, w|m) &= \frac{qv^2x^2}{(1-x)(1-v-qvx)}B^-(x, v, w|m-1) + \frac{qv^2wx^3}{(1-qvwx)(1-qvx)}\delta_{m,1} \\ &+ \frac{x(B^+(x, w, v|m-1) - B^+(x, 1, vw|m-1))}{1-w} \\ (10) \quad &- \frac{qv^2x^2}{(1-v-qvx)(1-vx-qvx)}B^-\left(\frac{vx}{1-qvx}, 1-qvx, w|m-1\right), \\ (1-w-qwx)B^+(x, v, w|m) &= \frac{qv^2x^3(1+w-wx)(1-w)}{(1-x)(1-wx)(1-qvwx)}\delta_{m,0} + \frac{qwx^2}{1-x}B^+(x, 1, vw|m) \\ &- \frac{qw^3x^2}{1-wx}B^+(wx, 1, v|m) - qw^2xB^+(wx, v, 1|m) + \frac{qwx(1-x+qx)}{(1-x)^2}B^-(x, vw, 1|m) \\ (11) \quad &- \frac{qw^2x(1-wx+qwx)}{(1-wx)^2}B^-(wx, v, 1|m) \end{aligned}$$

and

$$\begin{aligned} B^+(x, v, 1|m) &= \frac{x}{(1-qx)(1-x-qx)}B^+\left(\frac{x}{1-qx}, 1, v(1-qx)|m\right) - \frac{x}{1-x}B^+(x, 1, v|m) \\ &- \frac{1-x+qx}{(1-x)^2}B^-(x, v, 1|m) + \frac{1-x}{(1-x-qx)^2}B^-\left(\frac{x}{1-qx}, v(1-qx), 1|m\right) \\ (12) \quad &- \frac{(qx+x-2)qv^3x^3}{(1-x-qx)(1-qx)(1-x)(1-qvx)}\delta_{m,0}, \end{aligned}$$

with $B^-(x, v, w|m) = 0$ for $m < 0$.

2.2.1. *Case $m = 0$.* By Proposition 7, we see that $B^-(x, v, w|0) = 0$ and thus

$$\left(1-w-\frac{qwx}{w-x}\right)B^+(x/w, 1, w|0) = \frac{qx^3(1+w-x)(1-w)}{(w-x)(1-x)(1-qx)} - \frac{qwx}{1-x}B^+(x, 1, 1|0).$$

Let $w = w_0(x) = \frac{1}{2}(1+x-qx + \sqrt{(1+x-qx)^2 - 4x})$. Then

$$B^+(x, 1, 1|0) = \frac{(w_0(x)+1-x)(w_0(x)-1)x^2}{w_0(x)(qx-1)(w_0(x)-x)}$$

and

$$B^+(x, 1, w|0) = \frac{(w_0(wx)-w)(w+w_0(wx)-wx)qw^2x^3}{w_0(wx)(w_0(wx)-wx)(qwx-1)(qwx+w-wx-1+x)}.$$

Thus, by Lemma 2, we have $C^-(x, v, w|0) = B^-(x, v, w|0) = 0$, and then by Proposition 7, we get

$$\begin{aligned} B^+(x, v, w|0) &= \frac{qv^2x^2(vwx-vw-v+1)\sqrt{q^2v^2w^2x^2-2qv^2w^2x^2+v^2w^2x^2-2qvwx-2vwx+1}}{2(qv^2x^2-qvwx+qwx-vwx+wx+v-1)(qvwx-vwx+vw+x-1)} \\ &+ \frac{qv^2x^2(2qwx^2-qwx+2wx^2-2wx-2x+1)}{2(qvwx-vwx+vw+x-1)(qwx-1)(qv^2x^2-qvwx+qwx-vwx+wx+v-1)} \\ &+ \frac{qv^2w^2x^2(2x-1-w(2qx^2+2x^2-4x+1)+xw^2(q^2x+2qx^2-2qx+q-2x+2))}{2(qvwx-vwx+vw+x-1)(qwx-1)(qv^2x^2-qvwx+qwx-vwx+wx+v-1)} \\ &+ \frac{qv^3w^3x^3(wx-w-1)(q-1)(qwx-1)}{2(qvwx-vwx+vw+x-1)(qwx-1)(qv^2x^2-qvwx+qwx-vwx+wx+v-1)}. \end{aligned}$$

Hence, Proposition 7 implies the following result.

Theorem 8. *We have*

$$A(x|0) = \frac{2x}{1-x-qx + \sqrt{(1-x-qx)^2 - 4qx^2}}.$$

2.2.2. *Case $m = 1$.* By the $m = 0$ case and Proposition 7, we have

$$\begin{aligned} B^-(x, v, w|1) &= \frac{qv^2wx^3}{(1-qvwx)(1-qvx)} + \frac{x(B^+(x, w, v|0) - B^+(x, 1, vw|0))}{1-w} \\ &= \frac{vwx(qvx - vx^2 + vx + x - 1)\sqrt{(1 - (1+q)vwx)^2 - 4qv^2w^2x^2}}{2(qv^2wx^2 - qvwx + qvx - vwx + vx + w - 1)(qvwx - vwx + vw + x - 1)} \\ &\quad + \frac{vwx((1-x)(1-vx)(1-vwx) + qvx(vwx^2 - wx - w + 2x - 1) + q^2v^2wx^2)}{2(qv^2wx^2 - qvwx + qvx - vwx + vx + w - 1)(qvwx - vwx + vw + x - 1)}. \end{aligned}$$

So the generating function $B^+(x/w, 1, w|1)$ satisfies

$$\begin{aligned} &(1-w - \frac{qwx}{w-x})B^+(x/w, 1, w|1) \\ &= -\frac{qwx}{1-x}B^+(x, 1, 1|1) + \frac{qwx(w-x+qx)}{(w-x)^2}B^-(x/w, w, 1|1) - \frac{qwx(1-x+qx)}{(1-x)^2}B^-(x, 1, 1|1). \end{aligned}$$

By substituting $w = w_0(x)$, we obtain

$$B^+(x, 1, 1|1) = \frac{(w_0(x) - x + qx)(1-x)}{(w_0(x) - x)^2}B^-(x/w_0(x), w_0(x), 1|1) - \frac{1-x+qx}{1-x}B^-(x, 1, 1|1),$$

which implies that the expression of $B^+(x, 1, w|1)$ can be found from

$$\begin{aligned} &(1-w - \frac{qwx}{1-x})B^+(x, 1, w|1) \\ &= -\frac{qw^2x}{1-wx}B^+(wx, 1, 1|1) + \frac{qwx(1-x+qx)}{(1-x)^2}B^-(x, w, 1|1) - \frac{qw^2x(1-wx+qwx)}{(1-wx)^2}B^-(wx, 1, 1|1). \end{aligned}$$

Hence, by Proposition 7, one gets an explicit formula for $B^+(x, v, 1|1)$, and subsequently for $B^+(x, v, w|1)$, which implies the following result.

Theorem 9. *We have*

$$\begin{aligned} A(x|1) &= \frac{q^3(q-1)x^4 - (4q^3 + q^2 + q + 1)x^3 + (6q^2 + 5q + 3)x^2 - (4q + 3)x + 1}{2q^2x^4\sqrt{(1-x-qx)^2 - 4qx^2}} \\ &\quad + \frac{q^3x^3 - (3q^2 + 2q + 1)x^2 + (3q + 2)x - 1}{2q^2x^4}. \end{aligned}$$

2.2.3. *General Case.* Note that our calculation may be extended to any $m > 1$. More precisely, to find the generating function $A(x|m)$, we have the following algorithm:

- Let $m > 1$. Suppose we have found the generating functions $B^+(x, v, w|m-1)$ and $B^-(x, v, w|m-1)$.
- By (10), we have an explicit formula for the generating function $B^-(x, v, w|m)$.
- By (11), we have

$$\begin{aligned} &(1-w - \frac{qwx}{w-x})B^+(x/w, 1, w|m) \\ &= -\frac{qwx}{1-x}B^+(x, 1, 1|m) \\ &\quad + \frac{qwx(w-x+qx)}{(w-x)^2}B^-(x/w, w, 1|m) - \frac{qw^2x(1-wx+qwx)}{(1-wx)^2}B^-(wx, 1, 1|m). \end{aligned}$$

Thus, by taking $w = w_0(x)$ and using the formula of $B^-(x/w, w, 1|m)$, we obtain an explicit formula for the generating function $B^+(x, 1, 1|m)$.

- By (11) with $v = 1$, we obtain an explicit formula for $B^+(x, 1, w|m)$.
- By (12), we derive a formula for $B^+(x, v, 1|m)$.

- Thus, by (11) and the results of the previous steps, we obtain a formula for $B^+(x, v, w|m)$.
- Hence, by Proposition 7, namely,

$$A(x|m) = \frac{1}{1-x}B^+(x, 1, 1|m) + \frac{1-x+qx}{(1-x)^2}B^-(x, 1, 1|m),$$

we derive a formula for the generating function $A(x|m)$.

Applying this algorithm for $m = 2, 3$ yields

$$A(x|m) = \frac{a_m}{2q^{m+1}x^{3m+1}(qx+x-1)^{2m-3}\sqrt{(1-x-qx)^2-4qx^2}^{2m-1}} + \frac{b_m}{2q^{m+1}x^{3m+1}(qx+x-1)^{2m-3}},$$

where

$$\begin{aligned} a_2 &= (x+1)(x-1)^7 + x(2x^3 - 5x^2 - 3x - 9)(x-1)^5q - 2x^2(x^4 + 6x^3 - 13x^2 + 9x - 18)(x-1)^3q^2 \\ &\quad - 2x^3(x-1)(3x^4 - 30x^3 + 53x^2 - 63x + 42)q^3 - 2x^4(3x^4 - 36x^3 + 80x^2 - 105x + 63)q^4 \\ &\quad + x^5(2x^5 + x^4 - 9x^3 + 64x^2 - 126x + 126)q^5 - 2x^6(3x^4 + 5x^3 + 10x^2 - 21x + 42)q^6 \\ &\quad + 2x^7(3x^3 + 4x^2 - 3x + 18)q^7 - x^8(2x^2 + 9)q^8 + x^9q^9, \end{aligned}$$

$$\begin{aligned} b_2 &= (x+1)(x-1)^4 + x(2x^3 - 2x^2 + 3x - 6)(x-1)^2q + x^2(x-1)(4x^3 - 11x^2 + 15x - 15)q^2 \\ &\quad + 2x^3(3x^3 - 10x^2 + 15x - 10)q^3 + x^4(8x^2 - 15x + 15)q^4 - x^5(2x^2 - 3x + 6)q^5 + x^6q^6, \end{aligned}$$

$$\begin{aligned} a_3 &= -(2x^2 + 2x + 1)(x-1)^{12} - x(4x^4 - 6x^3 - 25x^2 - 18x - 15)(x-1)^{10}q \\ &\quad - x^2(3x^6 - 18x^5 - 15x^4 - 4x^3 + 147x^2 + 34x + 105)(x-1)^8q^2 + x^3(4x^7 + 29x^6 - 144x^5 \\ &\quad + 74x^4 - 404x^3 + 692x^2 - 286x + 455)(x-1)^6q^3 + x^4(6x^8 - 40x^7 - 195x^6 + 714x^5 - 1325x^4 \\ &\quad + 2944x^3 - 3211x^2 + 2262x - 1365)(x-1)^4q^4 - x^5(12x^8 + 15x^7 + 89x^6 - 1131x^5 + 2715x^4 \\ &\quad - 5925x^3 + 7381x^2 - 5291x + 3003)(x-1)^3q^5 + x^6(5x^8 + 10x^7 - 65x^6 + 1320x^5 - 4210x^4 \\ &\quad + 9650x^3 - 12837x^2 + 9724x - 5005)(x-1)^2q^6 - x^7(x-1)(5x^9 - 11x^8 - 15x^7 + 73x^6 - 1422x^5 \\ &\quad + 5370x^4 - 13143x^3 + 17853x^2 - 14157x + 6435)q^7 + x^8(22x^9 - 25x^8 - 28x^7 - 112x^6 + 1448x^5 \\ &\quad - 6076x^4 + 15192x^3 - 20526x^2 + 16302x - 6435)q^8 - x^9(35x^8 + 26x^6 + 214x^5 - 1130x^4 + 5170x^3 \\ &\quad - 9878x^2 + 9724x - 5005)q^9 + x^{10}(20x^7 - 2x^6 + 20x^5 - 62x^4 + 1132x^3 - 3619x^2 + 4290x \\ &\quad - 3003)q^{10} + x^{11}(5x^6 + 20x^5 - 4x^4 - 114x^3 + 1045x^2 - 1352x + 1365)q^{11} - x^{12}(10x^5 + 9x^4 \\ &\quad + 12x^3 + 245x^2 - 286x + 455)q^{12} + x^{13}(3x^4 + 4x^3 + 43x^2 - 36x + 105)q^{13} - x^{14}(4x^2 - 2x + 15)q^{14} \\ &\quad + x^{15}q^{15}, \end{aligned}$$

$$\begin{aligned} b_3 &= -(2x^2 + 2x + 1)(x-1)^7 - x(2x^2 + x + 2)(2x^2 + x - 5)(x-1)^5q - x^2(3x^6 + 2x^5 + 5x^4 - 49x^3 \\ &\quad + 57x^2 - 61x + 45)(x-1)^3q^2 - x^3(11x^6 - 33x^5 + 86x^4 - 218x^3 + 276x^2 - 236x + 120)(x-1)^2q^3 \\ &\quad - x^4(x-1)(19x^6 - 97x^5 + 287x^4 - 588x^3 + 707x^2 - 532x + 210)q^4 - x^5(27x^6 - 166x^5 + 532x^4 \\ &\quad - 1007x^3 + 1134x^2 - 770x + 252)q^5 - x^6(35x^5 - 181x^4 + 474x^3 - 665x^2 + 532x - 210)q^6 + x^7(5x^5 \\ &\quad - 34x^4 + 128x^3 - 240x^2 + 236x - 120)q^7 + x^8(3x^4 - 16x^3 + 48x^2 - 61x + 45)q^8 - x^9(4x^2 \\ &\quad - 7x + 10)q^9 + x^{10}q^{10}. \end{aligned}$$

3. THE DESC AND $n-1$ -ASC STATISTICS ON $I_n(>, \neq, \geq)$.

We start by decomposing the set $\mathcal{T}_n = I_n(>, \neq, \geq)$ as follows. To do so, it is useful to consider the largest descent bottom x of $e = e_1e_2 \cdots e_n \in \mathcal{T}_n$ where e contains at least one descent; that is, $x = \max\{e_{i+1} : 1 \leq i \leq n-1 \text{ and } e_i > e_{i+1}\}$. If w is the leftmost letter of e forming a descent with x , then w will be referred to as the *leftmost top* of e . Let $\mathcal{U}_n(i, j)$ for $i < j$ denote the subset of \mathcal{T}_n consisting of those e having leftmost top i and largest letter j , where it is assumed further that e ends in a j . Let $\mathcal{V}_n(i, j)$ for $i < j$ be the same as $\mathcal{U}_n(i, j)$ except that j is not the last letter of e . Let $\mathcal{W}_n(i, j)$ for $i > j$ be the subset of \mathcal{T}_n whose members e have leftmost top i and are

expressible as $e = e'ij^s$ for some $s \geq 1$ where $\max(e') \leq i$. Finally, let $\mathcal{Z}_n(i)$ be the subset of \mathcal{T}_n whose members end in i and are weakly increasing (i.e., contain no descents).

A few remarks are in order concerning these definitions. Note that the leftmost letter i forming a descent with x within any member of \mathcal{T}_n is also the smallest such letter, for otherwise a $(>, \neq, \geq)$ would be present with ixz , where z is the smallest. Further, no element greater than i can occur anywhere to the left of the first i due to avoidance of $(>, \neq, \geq)$, with all i 's occurring as a single run (directly preceding the first x). Similarly, x is the only element less than i occurring somewhere to the right of the run of i 's. Thus, members of $\mathcal{V}_n(i, j)$ must end in x , for otherwise there would be an occurrence of $(>, \neq, \geq)$ involving the last letter or a descent bottom that exceeds x contradicting its maximality. Further, members of $\mathcal{W}_n(i, j)$ are seen to have largest descent bottom j . Finally, a common interpretation for the parameter j which applies to all three of the subsets $\mathcal{U}_n(i, j)$, $\mathcal{V}_n(i, j)$ and $\mathcal{W}_n(i, j)$ is that it represents the largest element occurring anywhere to the right of the run of the letter i .

Define the distribution polynomial $u_n(i, j) = u_n(i, j; p, q)$ by

$$u_n(i, j) = \sum_{\pi \in \mathcal{U}_n(i, j)} p^{\text{desc}(\pi)} q^{n-1-\text{asc}(\pi)},$$

and likewise for $v_n(i, j)$, $w_n(i, j)$ and $z_n(i)$. For example, we have $\mathcal{V}_5(2, 4) = \{12141\}$,

$$\mathcal{U}_5(2, 4) = \{11214, 12114, 12134, 12144, 12214\},$$

$$\mathcal{W}_5(3, 2) = \{11132, 11232, 11322, 11332, 12232, 12322, 12332\}$$

and

$$\mathcal{Z}_4(4) = \{1114, 1124, 1134, 1224, 1234\},$$

which implies $v_5(2, 4) = p^2q^2$, $u_5(2, 4) = pq + 4pq^2$, $w_5(3, 2) = 4pq^2 + 3pq^3$ and $z_4(4) = 1 + 3q + q^2$. Put zero for any of these polynomial arrays if the subset of \mathcal{T}_n corresponding to the parameters in question is empty. Thus, these arrays may be nonzero only for the following parameter values: (i) $u_n(i, j)$: $n \geq 4$, $2 \leq i \leq n-2$ and $i < j \leq n$, (ii) $v_n(i, j)$: $n \geq 5$, $2 \leq i \leq n-3$ and $i < j \leq n-1$, (iii) $w_n(i, j)$: $n \geq 3$ and $1 \leq j < i \leq n-1$ and (iv) $z_n(i)$: $n \geq 1$ and $1 \leq i \leq n$.

Define $u_n = \sum_{i=2}^{n-2} \sum_{j=i+1}^n u_n(i, j)$ for $n \geq 4$, and likewise v_n for $n \geq 5$, w_n for $n \geq 3$ and z_n for $n \geq 1$. Then we seek a formula for $t_n = t_n(p, q)$ given by

$$t_n = u_n + v_n + w_n + z_n, \quad n \geq 1,$$

where we put zero for u_n , v_n or w_n if the corresponding subset of \mathcal{T}_n is empty. Note that t_n gives the joint distribution of the desc and $n-1-\text{asc}$ statistics on all of \mathcal{T}_n . Recall that $n-1-\text{asc}(\pi) = \text{desc}(\pi) + \text{lev}(\pi)$ for all $\pi \in I_n$.

The arrays $u_n(i, j)$, $v_n(i, j)$, $w_n(i, j)$ and $z_n(i)$ satisfy the following system of intertwined recurrences.

Lemma 10. *We have*

$$(13) \quad u_n(i, j) = qu_{n-1}(i, j) + \sum_{\ell=1}^{i-1} w_{n-1}(i, \ell) + \sum_{\ell=i+1}^{j-1} (u_{n-1}(i, \ell) + v_{n-1}(i, \ell)),$$

for $2 \leq i \leq n-2$ and $i < j \leq n$,

$$(14) \quad v_n(i, j) = q(pu_{n-1}(i, j) + v_{n-1}(i, j)),$$

for $2 \leq i \leq n-3$ and $i < j \leq n-1$,

$$(15) \quad \begin{aligned} w_n(i, j) &= pqz_{n-1}(i) + q(2w_{n-1}(i, j) - qw_{n-2}(i, j) - pqz_{n-2}(i)) + pq \sum_{k=2}^{j-1} u_{n-2}(k, j) \\ &+ pq \sum_{k=2}^{j-1} (u_{n-1}(k, j) - qu_{n-2}(k, j)) + \sum_{k=j+1}^{i-1} (w_{n-1}(k, j) - qw_{n-2}(k, j) - pqz_{n-2}(k)), \end{aligned}$$

for $1 \leq j < i \leq n-1$,

$$(16) \quad z_n(i) = qz_{n-1}(i) + \sum_{j=1}^{i-1} z_{n-1}(j), \quad 1 \leq i \leq n-1,$$

$$(17) \quad z_n(n) = \sum_{j=1}^{n-1} z_{n-1}(j), \quad n \geq 2,$$

with the initial condition $z_1(1) = 1$.

Proof. To show (13), first suppose $\pi \in \mathcal{U}_n(i, j)$ where i and j are as given. If the penultimate letter of π is also j , then there are clearly $qu_{n-1}(i, j)$ possibilities, so assume that this is not the case. Let ℓ be the second largest letter of π occurring to the right of the last i , x be the maximum descent bottom and y be the penultimate letter. Then $y < x$ is not possible, for otherwise ixy would be an occurrence of $(>, \neq, \geq)$. Further, $x < y < \ell$ is also not possible. To see this, suppose to the contrary that $x < y < \ell$ and let w denote the rightmost letter in π that is distinct from both y and j . If $w > y$, then y would be a descent bottom, contradicting that x is the maximum descent bottom. If $w < y$, then an occurrence of $(>, \neq, \geq)$ would be witnessed by lwy . Thus, we must have $y = \ell$ or $y = x$. If $y = \ell$ and $\ell \in [i+1, j-1]$, then there are $u_{n-1}(i, \ell)$ possibilities by definition, upon deleting j . If $y = x$ and $\ell \in [i+1, j-1]$, then π is expressible as $\pi = \pi'lx^rj$ for some $r \geq 1$ where $\max(\pi') \leq \ell$. Note that the letter directly preceding the run x^r must be ℓ , for otherwise membership in \mathcal{T}_n would be violated. Removing j from π then gives $v_{n-1}(i, \ell)$ for the contribution towards the weight in this case. Thus, the total weight of the members of $\mathcal{U}_n(i, j)$ in question for which $i < \ell < j$ is given by $\sum_{\ell=i+1}^{j-1} (u_{n-1}(i, \ell) + v_{n-1}(i, \ell))$. On the other hand, only $\ell = x$ is possible if it is assumed that $\ell \leq i$. In this case, all letters to the right of the last i (except j) are x . Thus, removing j and considering all possible values of x yields the remaining sum on the right side of (13), which finishes the proof of (13).

To show (15), suppose $\pi \in \mathcal{W}_n(i, j)$ where $1 \leq j < i \leq n-1$. If π ends in two or more j 's or if it is of the form $\pi = \pi'ij$, where π' has no descents, then there are contributions towards the weight of $qw_{n-1}(i, j)$ and $pqz_{n-1}(i)$, respectively. Note that the extra q factor in the former case accounts for the level between the last two letters of π which is lost when the final letter is deleted, whereas pq in the latter case accounts for the descent between i and j which is lost. So assume henceforth $\pi = \pi'ij$, where π' contains at least one descent; note that $\max(\pi') \leq i$. Let k denote the final letter of π' . If $k \in [j+1, i-1]$, then there are $w_{n-1}(k, j) - qw_{n-2}(k, j) - pqz_{n-2}(k)$ possibilities for each k by subtraction, upon deleting i , and summing over all k gives the last summation expression on the right side of (15). If $k = i$, then one gets $q(w_{n-1}(i, j) - qw_{n-2}(i, j) - pqz_{n-2}(i))$, where the q factor accounts for the extra level occurring between the last two letters i .

Suppose now $k = j$ and let z and ℓ denote the largest descent bottom and leftmost top of π' , respectively. Then $\ell = i$ is not possible, for otherwise $(>, \neq, \geq)$ is realized with iji , so we must have $\ell < i$. If $j < \ell < i$ and $z \neq j$, then $\ell z j$ would not be allowed, whereas if $z = j$, then the resulting inversion sequence would be of a form not even enumerated by $w_n(i, j)$ since ℓ , and not i , would be the leftmost top of π . If $\ell = j$, then jzj would occur with $z < j$, which is again not allowed. Thus, we must have $2 \leq \ell \leq j-1$ and deleting ij from π yields a member of $\mathcal{U}_{n-2}(\ell, j)$. Considering all possible ℓ in this case then gives a contribution towards the weight of $pq \sum_{\ell=2}^{j-1} u_{n-2}(\ell, j)$. Finally, if $k < j$, then $\ell \geq j$ would imply $\ell k j$ is not permissible, whence $2 \leq \ell \leq j-1$. Then deleting i from π in this case results in a member of $\mathcal{U}_{n-1}(\ell, j)$ in which the next-to-last letter is not j . By subtraction, there is a contribution of $pq(u_{n-1}(\ell, j) - qu_{n-2}(\ell, j))$ for each ℓ , which we then sum over ℓ . Combining now all of the previous cases implies (15).

The remaining formulas can be shown quickly. To realize (14), consider cases based on whether the terminal run of the maximum descent bottom of $\pi \in \mathcal{V}_n(i, j)$ is of length one or greater than one. Note that in the former case, the next-to-last letter must be j resulting in a descent involving the final two letters. To show (16) and (17), we delete the final i from $\pi \in \mathcal{Z}_n(i)$ and consider cases on whether the next-to-last letter equals i or is less than i , noting that the first option is not possible in the case $i = n$ since $\pi \in I_n$. \square

From the recurrences, we have that the nonzero values of the arrays above are given for $n = 2$ by $z_2(1) = q$, $z_2(2) = 1$, for $n = 3$ by $w_3(2, 1) = pq$, $z_3(1) = q^2$, $z_3(2) = 2q$, $z_3(3) = 1 + q$ and for $n = 4$ by

$$\begin{array}{lll} u_4(2, 3) = pq & u_4(2, 4) = pq & w_4(2, 1) = 3pq^2 \\ w_4(3, 1) = pq + pq^2 & w_4(3, 2) = pq + pq^2 & z_4(1) = q^3 \\ z_4(2) = 3q^2 & z_4(3) = 3q + 2q^2 & z_4(4) = 1 + 3q + q^2, \end{array}$$

which may be verified directly.

Now define

$$\begin{aligned} U(x, v, w) &= \sum_{n \geq 4} \sum_{i=2}^{n-2} \sum_{j=i+1}^n u_n(i, j) v^i w^j x^n, \\ V(x, v, w) &= \sum_{n \geq 5} \sum_{i=2}^{n-3} \sum_{j=i+1}^{n-1} v_n(i, j) v^i w^j x^n, \\ W(x, v, w) &= \sum_{n \geq 3} \sum_{i=2}^{n-1} \sum_{j=1}^{i-1} w_n(i, j) v^i w^j x^n, \end{aligned}$$

and $Z(x, v) = \sum_{n \geq 1} \sum_{i=1}^n z_n(i) v^i x^n$. By translating (13)–(17) above in terms of these generating functions, we obtain the following system of functional equations.

Lemma 11. *We have*

$$\begin{aligned} U(x, v, w) &= qxU(x, v, w) + \frac{wx}{1-w}(W(x, vw, 1) - wW(wx, v, 1)) \\ &\quad + \frac{wx}{1-w}(U(x, v, w) + V(x, v, w) - wU(wx, v, 1) - wV(wx, v, 1)), \\ V(x, v, w) &= pqxU(x, v, w) + qxV(x, v, w), \\ W(x, v, w) &= \frac{pqx}{1-w}(wZ(x, v) - Z(x, vw)) + 2qxW(x, v, w) - q^2x^2W(x, v, w) \\ &\quad - \frac{pq^2x^2}{1-w}(wZ(x, v) - Z(x, vw)) + \frac{pqvx^2}{1-v}(U(x, 1, vw) - vU(vx, 1, w)) \\ &\quad + \frac{pqvx}{1-v}(U(x, 1, vw) - U(vx, 1, w) - qxU(x, 1, vw) + qvxU(vx, 1, w)) \\ &\quad + \frac{v}{1-v}(xW(x, v, w) - qx^2W(x, v, w)) - \frac{pqvx^2}{(1-v)(1-w)}(wZ(x, v) - Z(x, vw)) \\ &\quad - \frac{vx}{1-v}(W(vx, 1, w) - qvxW(vx, 1, w)) + \frac{pqv^2x^2}{(1-v)(1-w)}(wZ(vx, 1) - Z(vx, w)), \\ Z(x, v) &= vxZ(vx, 1) + vx + qxZ(x, v) + \frac{vx}{1-v}(Z(x, v) - Z(vx, 1)). \end{aligned}$$

The last equation in Lemma 11 implies

$$\left(1 - \frac{qx}{v} - \frac{x}{1-v}\right) Z(x/v, v) = x + xZ(x, 1) - \frac{x}{1-v}Z(x, 1).$$

By taking $v = v_0(x) = \frac{1}{2}(1 - x + qx + \sqrt{(1 - x + qx)^2 - 4qx})$, we obtain

$$Z(x, 1) = \frac{1 - v_0(x)}{v_0(x)},$$

and hence

$$(18) \quad Z(x, v) = \frac{vx(v_0(vx) - v)}{v_0(vx)((1-v)(1-qx) - vx)}.$$

Further, Lemma 11 gives

$$V(x, v, w) = \frac{qpx}{1-qx}U(x, v, w).$$

Thus Lemma 11 implies that the generating functions $U(x, 1, v)$ and $W(x, v, 1)$ satisfy

$$\begin{aligned} U(x, 1, v) &= qxU(x, 1, v) + \frac{xv}{1-v}(W(x, v, 1) - vW(vx, 1, 1)) \\ &\quad + \frac{vx}{1-v}(U(x, 1, v) + \frac{pqx}{1-qx}U(x, 1, v) - vU(vx, 1, 1) - \frac{pqvx}{1-qx}U(vx, 1, 1)), \\ W(x, v, 1) &= 2qxW(x, v, 1) - q^2x^2W(x, v, 1) + \frac{pqvx^2}{1-v}(U(x, 1, v) - vU(vx, 1, 1)) \\ &\quad + \frac{pqvx}{1-v}(U(x, 1, v) - U(vx, 1, 1) - qxU(x, 1, v) + qvxU(vx, 1, 1)) \\ &\quad + \frac{v}{1-v}(xW(x, v, 1) - qx^2W(x, v, 1)) - \frac{vx}{1-v}(W(vx, 1, 1) - qvxW(vx, 1, 1)) \\ &\quad + F(x, v), \end{aligned}$$

where

$$\begin{aligned} F(x, v) &= \lim_{w \rightarrow 1} \left(\frac{pqx(1-qx)}{1-w}(wZ(x, v) - Z(x, vw)) - \frac{pqvx^2}{(1-v)(1-w)}(wZ(x, v) - Z(x, vw)) \right. \\ &\quad \left. + \frac{pqv^2x^2}{(1-v)(1-w)}(wZ(vx, 1) - Z(vx, w)) \right). \end{aligned}$$

Hence, by finding $U(x, 1, v)$ in the first equation and substituting it into the second with use of (18), one obtains the following result.

Lemma 12. *We have*

$$K(x, v)W(x, v, 1) = A(x, v)W(vx, 1, 1) + B(x, v)U(vx, 1, 1) + F(x, v),$$

where

$$\begin{aligned} K(x, v) &= \frac{(qx-1)^2((qx-1)^2 - (pqx^2 + 2q^2x^2 - 2qx^2 - 4qx + 2x + 2)v + (qx-x-1)^2v^2)}{(1-v)((qx-1)^2 - (pqx^2 + q^2x^2 - qx^2 - 2qx + x + 1)v)}, \\ A(x, v) &= -\frac{xv((qx-1)^2 + (q^3x^3 + q(p-q-1)x^2 - (q-1)x + 1)v + qx(q(pq+q-1)x^2 + (1-2q)(p+1)x + p+1)v^2)}{(1-v)((qx-1)^2 - (pqx^2 + q^2x^2 - qx^2 - 2qx + x + 1)v)}, \\ B(x, v) &= -\frac{pqvx((qx-1)^2 - qx(q(2q-1)x^2 + (p-4q+1)x + 2)v + qx^2(q^2(q-1)x^2 - (1-2q)(p-q)x - p+q)v^2)}{(1-qvx)((qx-1)^2 - (pqx^2 + q^2x^2 - qx^2 - 2qx + x + 1)v)}, \\ F(x, v) &= \frac{p(qv(q-1)x^2 - q(v+1)x + 1)}{2((q-1)vx - qx - v + 1)} \sqrt{(1-vx+qvx)^2 - 4qvx} \\ &\quad + \frac{(qv^2(q-1)^2x^3 - qv(qv-2q+v)x^2 + (2qv+q+v)x - 1)}{2(qvx - qx - vx - v + 1)}. \end{aligned}$$

For the kernel equation $K(x/v, v) = 0$, namely,

$$pqvx^2 - ((qx-v)(v-1) - vx)^2 = 0,$$

we have two different power series roots $v(x) = f_1(x), f_2(x)$ such that $v(0) = 1$. Here,

$$\begin{aligned} f_1(x) &= 1 - (1 - \sqrt{pq})x - \frac{q}{2\sqrt{pq}}(\sqrt{pq}(p+2) - p(2q+1))x^2 \\ &\quad - \frac{q}{8\sqrt{pq}}(8\sqrt{pq}(2pq+q+p+1) - p(5pq+8q^2+24q+3))x^3 + \dots, \\ f_2(x) &= 1 - (1 + \sqrt{pq})x - \frac{q}{2\sqrt{pq}}(\sqrt{pq}(p+2) + p(2q+1))x^2 \\ &\quad - \frac{q}{8}(8\sqrt{pq}(2pq+p+q+1) + p(5pq+8q^2+24q+3))x^3 + \dots. \end{aligned}$$

Replacing x by x/v in Lemma 12, we have

$$K(x/v, v)W(x/v, v, 1) = A(x/v, v)W(x, 1, 1) + B(x/v, v)U(x, 1, 1) + F(x/v, v).$$

By taking $v = f_1(x)$ and $v = f_2(x)$ in this last equation, we obtain the system

$$A(x/f_1(x), f_1(x))W(x, 1, 1) + B(x/f_1(x), f_1(x))U(x, 1, 1) + F(x/f_1(x), f_1(x)) = 0,$$

$$A(x/f_2(x), f_2(x))W(x, 1, 1) + B(x/f_2(x), f_2(x))U(x, 1, 1) + F(x/f_2(x), f_2(x)) = 0,$$

which leads to

$$(19) \quad U(x, 1, 1) = -\frac{A(x/f_1(x), f_1(x))F(x/f_2(x), f_2(x)) - A(x/f_2(x), f_2(x))F(x/f_1(x), f_1(x))}{A(x/f_1(x), f_1(x))B(x/f_2(x), f_2(x)) - A(x/f_2(x), f_2(x))B(x/f_1(x), f_1(x))},$$

$$(20) \quad W(x, 1, 1) = -\frac{F(x/f_1(x), f_1(x))B(x/f_2(x), f_2(x)) - F(x/f_2(x), f_2(x))B(x/f_1(x), f_1(x))}{A(x/f_1(x), f_1(x))B(x/f_2(x), f_2(x)) - A(x/f_2(x), f_2(x))B(x/f_1(x), f_1(x))}.$$

Moreover, we have $V(x, 1, 1) = \frac{qpx}{1-qx}U(x, 1, 1)$. Combining the results above yields the following generating function formula.

Theorem 13. *The generating function $\sum_{n \geq 1} t_n(p, q)x^n$ is given by*

$$\frac{1 - qx + qpx}{1 - qx}U(x, 1, 1) + W(x, 1, 1) + Z(x, 1),$$

where $Z(x, 1)$, $U(x, 1, 1)$ and $W(x, 1, 1)$ are given by (18), (19) and (20), respectively.

3.1. **Case $p = q = 1$.** Then the kernel equation $K(x/v, v) = 0$ in this case has for two of its roots

$$v = 1 \text{ and } v = 1 + \frac{16}{9} \sin^4\left(\frac{1}{3} \arcsin\left(\frac{3\sqrt{3x}}{2}\right)\right) - \frac{8}{3} \sin^2\left(\frac{1}{3} \arcsin\left(\frac{3\sqrt{3x}}{2}\right)\right).$$

Proceeding as in the proof of Theorem 13, we obtain the following result.

Theorem 14. *Let $y = \cos\left(\frac{1}{3} \arcsin\left(\frac{3\sqrt{3x}}{2}\right)\right)$. Then the generating function $\sum_{n \geq 1} t_n(1, 1)x^n$ is given by*

$$-\frac{(4y^2 - 1)^2 - 9x}{2A} \left(B + (64y^6 - 96y^4 + 36y^2 + 27x - 4)(64y^6 - 12y^2 - 27x + 2)\sqrt{1 - 4x} \right),$$

where $A = (256y^8 - 256y^6 - 48y^4 + 56y^2 + 81x - 8)(x(4y^2 - 1)^4 + (x - 1)((4y^2 - 1)^2 - 9x)^2)$ and $B = (4y^2 - 1)^6 - 9(32y^4x + 16y^4 - 16y^2x - 8y^2 - 7x + 1)((4y^2 - 1)^2 - 9x)$.

Remark: Comparing Theorems 5 and 14 implies (2), which was originally shown in [9] by a different method.

3.2. **Case $p = 1$.** Comparing the formulas from Theorems 4 and 13 when $p = 1$, one can show with the aid of mathematical programming the following result which is equivalent to (1).

Corollary 15. *We have $a_n(1, q) = t_n(1, q)$ for all $n \geq 1$.*

Remark: The joint distributions $a_n(p, q)$ and $t_n(p, q)$ are seen however to differ for general p , where p marks the number of descents (upon comparing the $n = 5$ terms of the corresponding generating functions).

3.3. **Coefficient of p^m .** Recall that we denote the coefficient of p^m in a generating function $f(x)$ by $f(x|m)$. Note that by (18), we get

$$Z(x, v|0) = \frac{vx(v_0(vx) - v)}{v_0(vx)((1 - v)(1 - qx) - vx)},$$

with $Z(x, v|m) = 0$ for all $m \geq 1$.

By Lemma 12, we get

$$K'(x, v)W(x, v, 1) = A'(x, v)W(vx, 1, 1) + B'(x, v)U(vx, 1, 1) + F'(x, v),$$

where

$$K'(x, v) = \frac{(qx - 1)^2((q - 1)vx - qx - v + 1)^2}{1 - v} - \frac{(qx - 1)^2vx^2qp}{1 - v},$$

$$A'(x, v) = \frac{vx(qx - 1)(1 - qvx)((q - 1)vx - qx - v + 1)}{1 - v} - \frac{v^2x^2q(q^2vx^2 + (1 - 2q)vx + v - x)}{1 - v}p,$$

$$B'(x, v) = (qx - 1)(q(q - 1)vx^2 - qvx - qx + 1)qvxvp + \frac{q^2x^3v^2((1 - 2q)vx + v + 1)}{1 - qvx}p^2,$$

$$F'(x, v) = (q(q-1)vx^2 - qvx - qx + 1) \left(\frac{1}{2}(1-qx)p - \frac{vx^2q}{2((q-1)vx - qx - v + 1)}p^2 \right) \sqrt{(1-vx+qvx)^2 - 4qvx} \\ + (vx(q(1-2q)vx^2 + qvx + 1) - (1-qx)(1-qvx)^2) \left(\frac{1}{2}(1-qx)p - \frac{vx^2q}{2((q-1)vx - qx - v + 1)}p^2 \right).$$

Thus,

$$(21) \quad \begin{aligned} & K'(x/v, v|0)W(x/v, v, 1|m) + K'(x/v, v|1)W(x/v, v, 1|m-1) \\ &= A'(x/v, v|0)W(x, 1, 1|m) + A'(x/v, v|1)W(x, 1, 1|m-1) \\ &+ B'(x/v, v|1)U(x, 1, 1|m-1) + B'(x/v, v|2)U(x, 1, 1|m-2) \\ &+ F'(x/v, v|1)\delta_{m,1} + F'(x/v, v|2)\delta_{m,2}. \end{aligned}$$

This equation performs a procedure which computes the coefficient of p^m for all $m \geq 0$. Clearly,

$$V(x, v, w|m) = U(x, v, w|m) = W(x, v, w|m) = 0, \quad m \leq 0.$$

Hence, by Theorem 13, we have that the coefficient of p^0 in the generating function $\sum_{n \geq 1} t_n x^n$ is given by $Z(x, 1) = \frac{1-v_0(x)}{v_0(x)}$.

3.3.1. *Case $m = 1$.* Suppose $m = 1$. Clearly, $Z(x, v|1) = 0$. By Lemma 11, we have $V(x, v, w|1) = 0$. Then by (21), we get

$$\begin{aligned} & \frac{((q-1)vx - qx - v^2 + v)^2(qx - v)^2}{v^2(1-v)} W(x/v, v, 1|1) \\ &= - \frac{((q-1)vx - qx - v^2 + v)x(qx - 1)(qx - v)}{1-v} W(x, 1, 1) \\ &- \frac{1}{2}(qx - v)(q(q-1)x^2 - qvx - qx + v)\sqrt{(q-1)^2x^2 - 2(q+1)x + 1} \\ &+ \frac{1}{2}(qx - v)(q(q-1)^2x^3 - q(q-1)vx^2 - 2q^2x^2 + (2q+1)vx + qx - v). \end{aligned}$$

Differentiating with respect to v and then substituting $v = v_0(x)$, we obtain

$$W(x, 1, 1|1) = \frac{q(q-1)^2x^3 - q(3q-1)x^2 + (3q+1)x - 1}{2x(qx-1)\sqrt{(q-1)^2x^2 - 2(q+1)x + 1}} + \frac{(qx-1)^2 - qx^2}{2x(qx-1)},$$

which leads to an explicit formula for $W(x/v, v, 1)$ and hence for $W(x, v, 1)$.

To find $U(x, 1, 1|1)$, we use Lemma 11 to get

$$(1 - \frac{qx}{w} - \frac{x}{1-w})U(x/w, 1, w|1) = \frac{x}{1-w}(W(x/w, w, 1|1) - wW(x, 1, 1|1)) - \frac{xw}{1-w}U(x, 1, 1|1).$$

By taking $w = v_0(x)$ and using the expressions for $W(x/w, w, 1|1)$ and $W(x, 1, 1|1)$ (from the previous step), we obtain explicit formulas for $U(x, 1, 1|1)$ and $U(x/w, 1, w|1)$. In particular,

$$U(x, 1, 1|1) = \frac{q(q-1)^4x^5 - q(q-1)(5q^2 - 3q - 1)x^4 + (10q^3 + 1)x^3 - 2(5q^2 + 4q + 2)x^2 + (5q + 4)x - 1}{2x(1-qx)\sqrt{((q-1)^2x^2 - (2q+1)x + 1)^3}} \\ + \frac{(q(q-1)^3x^4 - q^2(4q-3)x^3 + (6q^2 + 3q + 1)x^2 - (4q+3)x + 1)}{2x(1-qx)((q-1)^2x^2 - (2q+1)x + 1)}.$$

By Theorem 13, we have that the coefficient of p^1 in the generating function $\sum_{n \geq 1} t_n x^n$ is given by

$$U(x, 1, 1|1) + W(x, 1, 1|1).$$

3.3.2. *Case $m \geq 2$.* Let $m \geq 2$ and suppose we have determined the generating functions $W(x, v, 1|m-1)$ and $U(x, 1, w|m-1)$. Now let us describe an algorithm for finding $W(x, v, 1|m)$ and $U(x, 1, w|m)$.

- By Lemma 11, we have $Z(x, v|m) = 0$ and $V(x, 1, w|m) = \frac{qx}{1-qx}U(x, 1, w|m-1)$. Hence, we know $V(x, 1, w|m)$.

- By Lemma 11, and noting $Z(x, v|m-1) = 0$, we have

$$\begin{aligned} & (1 - qx/v)(1 - \frac{qx}{v} - \frac{x}{1-v})W(x/v, v, 1|m) \\ &= \frac{qx^2}{v(1-v)}(U(x/v, 1, v|m-1) - vU(x, 1, 1|m-1)) \\ &+ \frac{qx}{1-v}((1 - qx/v)U(x/v, 1, v|m-1) - (1 - qx)U(x, 1, 1|m-1)) - \frac{x(1 - qx)}{1-v}W(x, 1, 1|m). \end{aligned}$$

Note that $v = v_0(x)$ is a solution to $1 - \frac{qx}{v} - \frac{x}{1-v} = 0$. Hence, by taking $v = v_0(x)$ and using the prior established results for $W(x, v, 1|m-1)$ and $U(x, 1, w|m-1)$, we obtain an explicit formula for $W(x, 1, 1|m)$, which leads to a formula for $W(x/v, v, 1|m)$ and thus $W(x, v, 1|m)$.

- By Lemma 11, we have

$$\begin{aligned} (1 - \frac{qx}{w} - \frac{x}{1-w})U(x/w, 1, w|m) &= \frac{x}{1-w}(W(x/w, w, 1|m) - wW(x, 1, 1|m)) \\ &+ \frac{x}{1-w}(V(x/w, 1, w|m) - wU(x, 1, 1|m) - wV(x, 1, 1|m)). \end{aligned}$$

By taking $w = v_0(x)$ and using expressions of $W(x, v, 1|m)$ (see second step of the algorithm) and $V(x, 1, w|m)$ (see first step), we obtain $U(x, 1, 1|m)$, which leads to an explicit formula for $U(x, 1, w|m)$.

Hence, by applying the algorithm above, one can find the generating functions $W(x, v, 1|m)$ and $U(x, 1, w|m)$ from $W(x, v, 1|m-1)$ and $U(x, 1, w|m-1)$. Then the coefficient of p^m in the generating function $\sum_{n \geq 1} t_n x^n$ is given by

$$U(x, 1, 1|m) + V(x, 1, 1|m) + W(x, 1, 1|m), \quad m \geq 2.$$

REFERENCES

- [1] G. E. Andrews and S. Chern, A proof of Lin’s conjecture on inversion sequences avoiding patterns of relation triples, *J. Combin. Theory Ser. A* **179** (2021), Art. 105388.
- [2] W. Cao, E. Y. Jin and Z. Lin, Enumeration of inversion sequences avoiding triples of relations, *Discrete Appl. Math.* **260** (2019), 86–97.
- [3] S. Corteel, M. Martinez, C. D. Savage and M. Weselcouch, Patterns in inversion sequences I, *Discrete Math. Theor. Comput. Sci.* **18** (2016), #2.
- [4] V. Guerrini, On enumeration sequences generalising Catalan and Baxter numbers, Ph.D. Thesis, Università di Siena, 2017.
- [5] Q. Hou and T. Mansour, Kernel method and linear recurrence system, *J. Comput. Appl. Math.* **216**(1) (2008), 227–242.
- [6] D. Kim and Z. Lin, Refined restricted inversion sequences, *Sémin. Lothar. Comb.* **78B** (2017), 52.
- [7] S. Kitaev, *Patterns in Permutations and Words*, Springer-Verlag, Berlin, 2011.
- [8] Z. Lin, Restricted inversion sequences and enhanced 3-noncrossing partitions, *European J. Combin.* **70** (2018), 202–211.
- [9] Z. Lin, Patterns of relation triples in inversion and ascent sequences, *Theoret. Comput. Sci.* **804** (2020), 115–125.
- [10] Z. Lin and S. H. F. Yan, Vincular patterns in inversion sequences, *Appl. Math. Comput.* **364** (2020), Art. 124672.
- [11] T. Mansour and M. Shattuck, Pattern avoidance in inversion sequences, *Pure Math. Appl. (P.U.M.A.)* **25** (2015), 157–176.
- [12] T. Mansour and M. Shattuck, On a conjecture of Lin and Kim concerning a refinement of Schröder numbers, pre-print (2021).
- [13] M. A. Martinez and C. D. Savage, Patterns in inversion sequences II: Inversion sequences avoiding triples of relations, *J. Integer Seq.* **21** (2018), Art. 18.2.2.
- [14] N. J. Sloane, The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>, 2010.
- [15] C. Yan and Z. Lin, Inversion sequences avoiding pairs of patterns, *Discrete Math. Theor. Comput. Sci.* **22**(1) (2020), Art. 23.
- [16] S. H. F. Yan, Bijections for inversion sequences, ascent sequences and 3-nonnesting set partitions, *Appl. Math. Comput.* **325** (2018), 24–30.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, 3498838 HAIFA, ISRAEL
Email address: `tmansour@univ.haifa.ac.il`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE, 37996 KNOXVILLE, TN
Email address: `shattuck@math.utk.edu`