



The S-measure for algebraic integers having all their conjugates in a sector

Valérie Flammang

► To cite this version:

Valérie Flammang. The S-measure for algebraic integers having all their conjugates in a sector. Rocky Mountain Journal of Mathematics, 2020, 50 (4), pp.1313-1321. 10.1216/rmj.2020.50.1313 . hal-03295906

HAL Id: hal-03295906

<https://hal.science/hal-03295906>

Submitted on 22 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE S-MEASURE FOR ALGEBRAIC INTEGERS HAVING ALL THEIR CONJUGATES IN A SECTOR.

V. FLAMMANG

Abstract

Let α be a nonzero algebraic integer of degree d whose all conjugates $\alpha_1 = \alpha, \alpha_2, \dots, \alpha_d$ lie in a sector $|\arg z| \leq \theta$, $0 < \theta \leq 90^\circ$. We define the *S-measure* of α by $S(\alpha) = \sum_{i=1}^d |\alpha_i|$ and the *absolute S-measure* of α by $s(\alpha) = S(\alpha)/d$. We compute the greatest lower bound $c(\theta)$ of $s(\alpha)$ for α belonging to twelve subintervals of $(0, \theta)$. Among these subintervals, three are complete. These computations use the principle of explicit auxiliary functions and our recursive algorithm.

1 Introduction

Let α be a nonzero algebraic integer of degree d . We define the *S-measure* of α by:

$$S(\alpha) = \sum_{i=1}^d |\alpha_i|.$$

The *absolute S-measure* of α is the quantity $s(\alpha) = S(\alpha)/d$.

If P is the minimal polynomial of α , then $S(P) = S(\alpha)$ and $s(P) = s(\alpha)$.

The arithmetic-geometric inequality shows that $s(\alpha) \geq 1$. Since α is a nonzero algebraic integer the quantity $|\alpha_1 \dots \alpha_d|$ is a positive integer, and therefore $|\alpha_1 \dots \alpha_d| \geq 1$. The equality holds if and only if α is a root of unity. Indeed, if $s(\alpha) = 1$, then $\frac{\sum_{i=1}^d |\alpha_i|}{d} = \prod_{i=1}^d |\alpha_i|^{1/d}$. Thus, for $i = 1, \dots, d$, all the $|\alpha_i|$ are equal to each other and then equal to 1. The theorem of Kronecker [K] implies that α is a root of unity.

In this paper, we consider an algebraic integer α whose all conjugates lie in a sector $|\arg z| \leq \theta$, $0 < \theta \leq 90^\circ$.

Note that the case θ equals to 0 is well known. α is then a *totally positive* algebraic integer i.e., all its conjugates are positive real numbers. Thus, the S-measure of α is then the usual trace of α and was already studied by a lot of authors. For more details, see for example [S], [ABP], [AP1], [AP2], [LW], [DW], [F1].

Hence, we focus here our attention on $\theta > 0$. We follow a work of M. Langevin [La] on the absolute Mahler measure $\Omega(\alpha) = (\prod_{i=1}^d \max(1, |\alpha_i|))^{1/d}$ of algebraic integers α having all their conjugates in a sector. He proved that there exists a function $c(\theta)$ on $[0, 180^\circ]$, always > 1 , such that if $\alpha \neq 0$ is not a root of unity, whose conjugates all lie in $S_\theta = \{z \mid |\arg z| \leq \theta, 0 \leq \theta \leq 180^\circ\}$, then $\Omega(\alpha) \geq c(\theta)$. But his result was not numerical. G. Rhin and C. Smyth [RS] were the first to succeed in finding the exact value of $c(\theta)$ for θ in nine subintervals of $[0, 120^\circ]$ and conjectured that $c(\theta)$ is a “staircase” function of θ , which is constant except for finitely many left discontinuities in any closed subinterval of $[0, 180^\circ]$. Their method used explicit auxiliary polynomials found heuristically. In 2004, thanks to Wu’s algorithm [Wu], G. Rhin and Q. Wu [RW] gave the exact value of $c(\theta)$ for four new subintervals of $[0, 140^\circ]$ and extended four existing subintervals. In 2013, the author and G. Rhin [FR] found for the first time a complete subinterval and a fourteenth subinterval. A complete subinterval is an interval on which the function $c(\theta)$ is constant, with jump discontinuities at each end. These improvements were due to our recursive algorithm. From there, we studied most of the usual measures for α having all its conjugates in a sector, in particular the usual trace [F2].

Here, we prove the following result

Theorem 1. *There exist a left discontinuous, strictly positive, staircase function g on $[0, 90^\circ)$ and a positive, continuous, monotonically decreasing function f on $[0, 90^\circ)$ such that:*

for $\theta \in [0, 90^\circ)$, if α is a nonzero algebraic integer of degree ≥ 2 whose conjugates all lie in $S_\theta = \{z \mid |\arg z| \leq \theta, 0 \leq \theta \leq 90^\circ\}$ then

$$s(\alpha) \geq \min(f(\theta), g(\theta))$$

Moreover, the exact value of $c(\theta) = \inf_{|\arg \alpha| \leq \theta} s(\alpha)$ is known on twelve subintervals of $[0, 90^\circ)$.

These intervals are given in Table 1.

The function $g(\theta)$ is a decreasing staircase function having left discontinuities. The function $g(\theta)$ is the smallest value of $s(\alpha)$ that could be found for α having all its conjugates in $|\arg z| \leq \theta$.

The function $f(\theta)$ is given by $f(\theta) = \max_{1 \leq i \leq 12} (f_i(\theta))$, and the functions $f_i(\theta)$ are of the type:

$$f_i(\theta) = \min_{z \in S_\theta} \left(|z| - \sum_{1 \leq j \leq J} c_{ij} \log |Q_{ij}(z)| \right)$$

where the polynomials Q_{ij} with integer coefficients and the positive numbers c_{ij} are read off from Table 2.

Table 1 Intervals $[\theta_i, \theta'_i]$ where $c(\theta)$ is known exactly. Here $c(\theta) = c(\theta_i) = s(P)$ for $\theta \in [\theta_i, \theta'_i]$.

P denotes the polynomial whose all conjugates lie in S_{θ_i} . The angles are given in degrees.

i	$c(\theta)$	θ_i	θ'_i	P
1	1.5	0	20.1	$x^2 - 3x + 1$
2	1.428241	21.640384	26.408740	$x^3 - 4x^2 + 5x - 1$
3	1.395643	26.408740	29.86	$x^4 - 5x^3 + 9x^2 - 5x + 1$
4	1.365660	32.569940	33.14	$x^6 - 7x^5 + 21x^4 - 32x^3 + 25x^2 - 8x + 1$
5	1.280776	38.668282	40.65	$x^4 - 4x^3 + 7x^2 - 4x + 1$
6	1.221499	47.941432	50.882970	$x^6 - 5x^5 + 13x^4 - 17x^3 + 13x^2 - 5x + 1$
7	1.214392	50.882970	53.32	$x^6 - 5x^5 + 13x^4 - 18x^3 + 15x^2 - 6x + 1$
8	1.196310	56.004692	57.66	$x^8 - 6x^7 + 20x^6 - 40x^5 + 53x^4 - 45x^3 + 24x^2 - 7x + 1$
9	1.190591	59.066144	63.91	$x^6 - 4x^5 + 10x^4 - 13x^3 + 11x^2 - 4x + 1$
10	1.144122	64.086354	72.86	$x^4 - 2x^3 + 4x^3 - 2x + 1$
11	1.094450	76.795630	80.656154	$x^4 - x^3 + 3x^2 - x + 1$
12	1.073092	80.656154	87.5	$x^3 - x^2 + 2x - 1$

2 The explicit auxiliary functions

Let α be an algebraic integer of degree d all of which conjugates $\alpha_1 = \alpha, \dots, \alpha_d$ lie in $S_\theta = \{z \mid |\arg z| \leq \theta\}$, with minimal polynomial P . The auxiliary functions f_i , $1 \leq i \leq 12$, are of the following type:

$$\forall z \in S_\theta, f(z) = |z| - \sum_{1 \leq j \leq J} c_j \log |Q_j(z)|, \quad (2.1)$$

where the coefficients c_j are positive real numbers and the polynomials Q_j are nonzero in $\mathbb{Z}[z]$, not necessarily irreducible, but not divisible by P . The main point is to choose the numbers c_j and the polynomials Q_j in order to maximize the minimum m of f on S_θ .

If we have $\sum_{i=1}^d f(\alpha_i) \geq md$ then

$$S(\alpha) = \sum_{i=1}^d |\alpha_i| \geq md + \sum_{j=1}^J c_j \log |\prod_{i=1}^d Q_j(\alpha_i)|.$$

Since P does not divide any Q_j , $\prod_{i=1}^d Q_j(\alpha_i)$ is a nonzero integer because it is the resultant of P and Q_j . Therefore, we have

$$s(\alpha) \geq m.$$

The main difficulty in this procedure is to find a good list of polynomials Q_j which gives a value of m as large as possible. For this purpose, we link the auxiliary function to the integer transfinite diameter in order to find our polynomials by the recursive algorithm.

3 Auxiliary functions and integer transfinite diameter

In this section, we will need the following definition:

Let K be a compact subset of \mathbb{C} . We define the *integer transfinite diameter of K* by

$$\begin{aligned} t_{\mathbb{Z}}(K) &= \liminf_{\substack{n \geq 1 \\ n \rightarrow \infty}} \inf_{\substack{P \in \mathbb{Z}[z] \\ \deg(P) = n}} |P|_{\infty, K}^{\frac{1}{n}}, \end{aligned}$$

where $|P|_{\infty, K} = \sup_{z \in K} |P(z)|$. If φ is a positive function defined on K , the φ -*integer transfinite diameter of K* is defined as

$$\begin{aligned} t_{\mathbb{Z}, \varphi}(K) &= \liminf_{\substack{n \geq 1 \\ n \rightarrow \infty}} \inf_{\substack{P \in \mathbb{Z}[z] \\ \deg(P) = n}} \sup_{z \in K} \left(|P(z)|^{\frac{1}{n}} \varphi(z) \right). \end{aligned}$$

This weighted version of the integer transfinite diameter was introduced by F. Amoroso [A]. It is an important tool in the study of rational approximations of logarithms of rational numbers. Inside the auxiliary function (2.1), we replace the numbers c_j by rational numbers a_j/q where q is a common denominator of the c_j for $1 \leq j \leq J$. Then we can write:

$$\text{for } z \in S_\theta, \quad f(z) = |z| - \frac{t}{r} \log |Q(z)| \geq m \tag{3.1}$$

where $Q = \prod_{j=1}^J Q_j^{a_j} \in \mathbb{Z}[z]$ is of degree $r = \sum_{i=1}^J a_j \deg Q_j$ and $t = \sum_{j=1}^J c_j \deg Q_j$. We want to get a function whose minimum m in the sector S_θ is as large as possible. Thus we search for a polynomial $Q \in \mathbb{Z}[z]$ such that

$$\sup_{z \in S_\theta} |Q(z)|^{t/r} (e^{|z|})^{-1} \leq e^{-m}.$$

If we suppose that t is fixed, it is clear that we need an effective upper bound for the quantity

$$\begin{aligned} t_{\mathbb{Z}, \varphi}(S_\theta) &= \liminf_{\substack{r \geq 1 \\ r \rightarrow +\infty}} \inf_{\substack{P \in \mathbb{Z}[z] \\ \deg(P) = r}} \sup_{z \in S_\theta} \left(|P(z)|^{\frac{t}{r}} \varphi(z) \right) . \end{aligned}$$

where we use the weight $\varphi(z) = (e^{|z|})^{-1}$.

4 Construction of the auxiliary functions

The polynomials involved in the auxiliary functions are found by our recursive algorithm developed in [F3] from Wu's algorithm [Wu]. It replaces a heuristic search for suitable polynomials by a systematic inductive search. Suppose that we have already found a list Q_1, Q_2, \dots, Q_J of suitable polynomials. Then we use semi-infinite linear programming (introduced into number theory by C.J. Smyth [S]) to optimize f for this set of polynomials (i.e., to get the greatest possible m). We obtain the real positive numbers c_1, c_2, \dots, c_J and then f in the form (3.1) as above. The function f is invariant under complex conjugation so we can limit ourselves to the sector $S'_\theta = \{z \in \mathbb{C} \text{ such that } 0 \leq \arg z \leq \theta\}$. Since the function f is harmonic in S'_θ outside the union of arbitrarily small disks around the roots of the polynomials Q_j , the minimum is taken on the boundary of S'_θ . Thus, it is sufficient to consider the minimum of f on the half line $R_\theta = \{z = se^{i\theta}, s \geq 0\}$.

The auxiliary function on the half line is:

$$f(z) = s - \sum_{1 \leq j \leq J} c_j \log |Q_j(z)|.$$

Thus, by our algorithm, we seek a polynomial $R(z) = \sum_{l=0}^k a_l z^l \in \mathbb{Z}[z]$, where k is varying from 4 to 15 successively, such that

$$\sup_{s \geq 0} |Q(se^{i\theta})R(se^{i\theta})|^{\frac{t}{r+k}} (e^s)^{-1} \leq e^{-m},$$

i.e., such that

$$\sup_{s \geq 0} |Q(se^{i\theta})R(se^{i\theta})| (e^s)^{-\frac{r+k}{t}}$$

is as small as possible.

But, here, $R(se^{i\theta})$ is not a linear forms in the unknown coefficients a_i of R . We replace them by their real parts and their imaginary parts. Then, we get the following real linear forms

$$|Q(s_n e^{i\theta})| \operatorname{Re}(R(s_n e^{i\theta})) (e^{s_n})^{-\frac{r+k}{t}} \quad \text{and} \quad |Q(s_n e^{i\theta})| \operatorname{Im}(R(s_n e^{i\theta})) (e^{s_n})^{-\frac{r+k}{t}}.$$

The s_n are suitable points in $(0, \infty)$, including the points where f has its least local minima. All these linear forms define a real lattice on \mathbb{Z}^{k+1} . We use algorithm LLL to obtain a small vector in this lattice.

Then, we get a polynomial R whose factors R_j are good candidates to enlarge the set of polynomials (Q_1, Q_2, \dots, Q_J) . We only keep the polynomials R_j which have a nonzero coefficient c_j in the newly optimized auxiliary function f . After optimization, some previous polynomials Q_j may have a zero coefficient c_j and so are removed.

Table 2: The auxiliary functions f_i , $1 \leq i \leq 12$.

f_1	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.54074137	0	x
	1.0000000	0.64261192	0	$x - 1$
	1.5000000	0.13252114	0	$x^2 - 3x + 1$
	1.5303349	0.00694451	19.533923	$x^{11} - 16x^{10} + 112x^9 - 447x^8 + 1115x^7 - 1798x^6 + 1883x^5 - 1264x^4 + 531x^3 - 133x^2 + 18x - 1$
	1.4282414	0.08169266	21.640384	$x^3 - 4x^2 + 5x - 1$
	1.4806329	0.02908250	32.317239	$x^{10} - 14x^9 + 85x^8 - 291x^7 + 613x^6 - 815x^5 + 677x^4 - 339x^3 + 98x^2 - 15x + 1$
f_2	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.56975402	0	x
	1.0000000	0.77184525	0	$x - 1$
	1.4282414	0.07029371	21.640384	$x^3 - 4x^2 + 5x - 1$
	1.4782611	0.00634725	25.535459	$x^5 - 7x^4 + 17x^3 - 18x^2 + 7x - 1$
	1.3956439	0.03715149	26.408740	$x^4 - 5x^3 + 9x^2 - 5x + 1$
	1.4666102	0.00582468	27.821143	$x^6 - 8x^5 + 26x^4 - 41x^3 + 31x^2 - 9x + 1$
	1.5087080	0.00143308	28.849859	$x^8 - 11x^7 + 52x^6 - 134x^5 + 201x^4 - 174x^3 + 82x^2 - 20x + 2$
	1.4320220	0.00119483	32.281080	$x^7 - 9x^6 + 34x^5 - 67x^4 + 72x^3 - 40x^2 + 11x - 1$
	1.4331455	0.01240370	32.422861	$x^7 - 9x^6 + 34x^5 - 67x^4 + 72x^3 - 39x^2 + 10x - 1$
	1.4410315	0.00448025	32.724266	$x^7 - 9x^6 + 34x^5 - 66x^4 + 69x^3 - 37x^2 + 10x - 1$
	1.4312869	0.00316917	37.481532	$x^7 - 9x^6 + 34x^5 - 66x^4 + 68x^3 - 35x^2 + 9x - 1$
f_3	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.58114845	0	x
	1.0000000	0.76150091	0	$x - 1$
	1.3956439	0.03170582	26.408740	$x^4 - 5x^3 + 9x^2 - 5x + 1$
	1.5062565	0.00371221	29.619471	$x^6 - 8x^5 + 27x^4 - 46x^3 + 40x^2 - 15x + 2$
	1.7320508	0.01765706	30.000000	$x^2 - 3x + 3$
	1.4798946	0.00283543	30.333904	$x^{10} - 13x^9 + 76x^8 - 257x^7 + 546x^6 - 743x^5 + 636x^4 - 326x^3 + 96x^2 - 15x + 1$
	1.4856233	0.00365455	31.605909	$x^7 - 9x^6 + 35x^5 - 71x^4 + 77x^3 - 40x^2 + 10x - 1$
	1.5478075	0.01357778	32.185004	$x^3 - 4x^2 + 6x - 2$
	1.4177256	0.00425036	32.536719	$x^{12} - 15x^{11} + 103x^{10} - 422x^9 + 1135x^8 - 2084x^7 + 2641x^6 - 2294x^5 + 1341x^4 - 515x^3 + 124x^2 - 17x + 1$
	1.3656607	0.02118463	32.569940	$x^6 - 7x^5 + 21x^4 - 32x^3 + 25x^2 - 8x + 1$
	1.4725753	0.00603112	32.870830	$x^7 - 9x^6 + 35x^5 - 72x^4 + 81x^3 - 45x^2 + 11x - 1$
	1.4650511	0.00585525	34.381807	$x^{11} - 14x^{10} + 89x^9 - 332x^8 + 794x^7 - 1254x^6 + 1308x^5 - 885x^4 + 382x^3 - 101x^2 + 15x - 1$
	1.3521869	0.00668035	34.985334	$x^5 - 6x^4 + 14x^3 - 15x^2 + 6x - 1$
f_4	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.56838131	0	x
	1.0000000	0.77021312	0	$x - 1$
	1.5478075	0.00171754	32.185004	$x^3 - 4x^2 + 6x - 2$
	1.3656607	0.02265560	32.569940	$x^6 - 7x^5 + 21x^4 - 32x^3 + 25x^2 - 8x + 1$
	1.4910451	0.00213815	34.193790	$x^4 - 5x^3 + 10x^2 - 7x + 2$
	1.3521869	0.01440308	34.985334	$x^5 - 6x^4 + 14x^3 - 15x^2 + 6x - 1$
	1.3973223	0.00151006	35.031799	$x^{11} - 13x^{10} + 78x^9 - 278x^8 + 643x^7 - 996x^6 + 1037x^5 - 715x^4 + 320x^3 - 89x^2 + 14x - 1$
	1.4102499	0.00407700	35.274805	$x^{10} - 12x^9 + 66x^8 - 214x^7 + 446x^6 - 614x^5 + 556x^4 - 322x^3 + 115x^2 - 23x + 2$
	1.4226863	0.00818534	35.357349	$x^{10} - 12x^9 + 66x^8 - 213x^7 + 438x^6 - 585x^5 + 500x^4 - 262x^3 + 82x^2 - 14x + 1$
	1.4206206	0.00058659	36.322451	$x^{15} - 18x^{14} + 153x^{13} - 805x^{12} + 2906x^{11} - 7561x^{10} + 14526x^9 - 20798x^8 + 22166x^7 - 17417x^6 + 9923x^5 - 4004x^4 + 1108x^3 - 199x^2 + 21x - 1$
	1.4665625	0.00084278	37.323334	$x^{13} - 16x^{12} + 120x^{11} - 550x^{10} + 1700x^9 - 3702x^8 + 5778x^7 - 6461x^6 + 5105x^5 - 2780x^4 + 1013x^3 - 236x^2 + 32x - 2$
	1.3808230	0.00322863	37.650150	$x^{10} - 12x^9 + 65x^8 - 205x^7 + 410x^6 - 533x^5 + 447x^4 - 235x^3 + 75x^2 - 13x + 1$
	1.3737014	0.00116992	38.363201	$x^{11} - 13x^{10} + 77x^9 - 270x^8 + 615x^7 - 944x^6 + 984x^5 - 688x^4 + 312x^3 - 88x^2 + 14x - 1$
	1.2807764	0.02023734	38.668282	$x^4 - 4x^3 + 7x^2 - 4x + 1$
	1.3500797	0.00237483	39.090356	$x^9 - 10x^8 + 46x^7 - 123x^6 + 207x^5 - 222x^4 + 148x^3 - 57x^2 + 12x - 1$
f_5	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.56549176	0	x
	1.0000000	0.82799146	0	$x - 1$
	1.2807764	0.01663368	38.668282	$x^4 - 4x^3 + 7x^2 - 4x + 1$
	1.2887120	0.04592145	40.894844	$x^3 - 3x^2 + 4x - 1$
	1.3025959	0.01073176	41.920317	$x^6 - 6x^5 + 17x^4 - 25x^3 + 20x^2 - 7x + 1$
	1.2993422	0.00373700	43.441992	$x^7 - 7x^6 + 23x^5 - 41x^4 + 43x^3 - 25x^2 + 8x - 1$
	1.2882975	0.00079616	44.070606	$x^8 - 8x^7 + 30x^6 - 64x^5 + 83x^4 - 65x^3 + 31x^2 - 8x + 1$
	1.2703626	0.00312681	45.173320	$x^6 - 6x^5 + 16x^4 - 22x^3 + 17x^2 - 6x + 1$
	1.3256092	0.00502304	46.440516	$x^{10} - 10x^9 + 48x^8 - 138x^7 + 259x^6 - 324x^5 + 270x^4 - 145x^3 + 50x^2 - 10x + 1$
	1.2971977	0.00828626	51.015623	$x^7 - 7x^6 + 23x^5 - 41x^4 + 42x^3 - 23x^2 + 7x - 1$
f_6	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.62533984	0	x
	1.0000000	0.13129538	60.000000	$x^2 - x + 1$
	1.4142136	0.01666709	45.000000	$x^2 - 2x + 2$
	1.2214993	0.00012139	47.941432	$x^6 - 5x^5 + 13x^4 - 17x^3 + 13x^2 - 5x + 1$
	1.1513878	0.08048976	49.353681	$x^4 - 3x^3 + 5x^2 - 3x + 1$
	1.2143924	0.03396658	50.882970	$x^6 - 5x^5 + 13x^4 - 18x^3 + 15x^2 - 6x + 1$
	1.2257003	0.01375343	53.350320	$x^8 - 6x^7 + 20x^6 - 38x^5 + 48x^4 - 38x^3 + 20x^2 - 6x + 1$
	1.2237982	0.02183003	54.861223	$x^8 - 6x^7 + 20x^6 - 39x^5 + 50x^4 - 40x^3 + 21x^2 - 6x + 1$

f_7	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.64156873	0	x
	1.4142136	0.02560466	45.000000	$x^2 - 2x + 2$
	1.1513878	0.08849685	49.353681	$x^4 - 3x^3 + 5x^2 - 3x + 1$
	1.2143924	0.02688682	50.882970	$x^6 - 5x^5 + 13x^4 - 18x^3 + 15x^2 - 6x + 1$
	1.2257003	0.01486371	53.350320	$x^8 - 6x^7 + 20x^6 - 38x^5 + 48x^4 - 38x^3 + 20x^2 - 6x + 1$
	1.2237982	0.01372826	54.861223	$x^8 - 6x^7 + 20x^6 - 39x^5 + 50x^4 - 40x^3 + 21x^2 - 6x + 1$
	1.0000000	0.12411259	60.000000	$x^2 - x + 1$
f_8	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.66258051	0	x
	1.1513878	0.08777864	49.353681	$x^4 - 3x^3 + 5x^2 - 3x + 1$
	1.1963101	0.00140181	56.004692	$x^8 - 6x^7 + 20x^6 - 40x^5 + 53x^4 - 45x^3 + 24x^2 - 7x + 1$
	1.1598550	0.01093296	59.015770	$x^3 - 2x^2 + 3x - 1$
	1.1905913	0.02037738	59.066144	$x^6 - 4x^5 + 10x^4 - 13x^3 + 11x^2 - 4x + 1$
	1.0000000	0.19173198	60.000000	$x^2 - x + 1$
f_9	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.69106953	0	x
	1.1905913	0.00047744	59.066144	$x^6 - 4x^5 + 10x^4 - 13x^3 + 11x^2 - 4x + 1$
	1.0000000	0.20669082	60.000000	$x^2 - x + 1$
	1.2329260	0.00070272	63.571914	$x^7 - 4x^6 + 12x^5 - 19x^4 + 23x^3 - 15x^2 + 7x - 1$
	1.1441228	0.06530631	64.086354	$x^4 - 2x^3 + 4x^2 - 2x + 1$
	1.2104994	0.02478898	64.985950	$x^7 - 4x^6 + 12x^5 - 19x^4 + 22x^3 - 14x^2 + 5x - 1$
f_{10}	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.65189622	0	x
	1.0000000	0.20329094	60.000000	$x^2 - x + 1$
	1.1441228	0.05925279	64.086354	$x^4 - 2x^3 + 4x^2 - 2x + 1$
	1.4142136	0.01542631	69.295189	$x^2 - x + 2$
	1.1660867	0.01043488	73.001452	$x^8 - 3x^7 + 10x^6 - 14x^5 + 20x^4 - 14x^3 + 10x^2 - 3x + 1$
	1.1710692	0.02295000	74.639018	$x^8 - 3x^7 + 10x^6 - 15x^5 + 22x^4 - 16x^3 + 11x^2 - 3x + 1$
	1.1435156	0.00095069	77.579788	$x^6 - 2x^5 + 6x^4 - 5x^3 + 6x^2 - 2x + 1$
f_{11}	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.67127297	0	x
	1.0944505	0.029235734	76.795630	$x^4 - x^3 + 3x^2 - x + 1$
	1.1307279	0.01817266	84.841604	$x^6 - x^5 + 5x^4 - 3x^3 + 5x^2 - x + 1$
	1.1476672	0.02486566	86.221414	$x^6 - x^5 + 5x^4 - 3x^3 + 6x^2 - x + 1$
	1.0000000	0.22968256	90.000000	$x^2 + 1$
f_{12}	$s(Q_j)$	c_j	$\theta(Q_j)$	Q_j
	0	0.74200412	0	x
	1.0944505	0.08810863	76.795630	$x^4 - x^3 + 3x^2 - x + 1$
	1.0730921	0.002505677	80.656154	$x^3 - x^2 + 2x - 1$
	1.0000000	0.14047981	90.000000	$x^2 + 1$

References

- [A] F. Amoroso. *f-transfinite diameter and number-theoretic applications*, Ann. Inst. Fourier, Grenoble, **43** (1993), 1179–1198.
- [ABP] J. Aguirre, M. Bilbao, J. C. Peral. *The trace of totally positive algebraic integers*, Math. Comp. **75** (2006), no 253, 385–393.
- [AP1] J. Aguirre and J.C. Peral. *The integer Chebyshev constant of Farey intervals*, Publ. Mat. 2007, Proceedings of the Primeras Jornadas de Teoría de Números, 11–27.
- [AP2] J. Aguirre and J.C. Peral. *The trace problem for totally positive algebraic integers. With an appendix by Jean-Pierre Serre*, London Math. Soc. Lecture Note Ser., **352**, Number theory and polynomials, 1–19, Cambridge Univ. Press, Cambridge, 2008.
- [DW] X. Dong and Q. Wu. *The absolute trace of totally positive reciprocal algebraic integers*, J. Number Theory 170 (2017), 66–74.
- [F1] V. Flammang. *Une nouvelle minoration pour la trace absolue des entiers algébriques totalement positifs* available on hal-01346165v1, (2016)
- [F2] V. Flammang. *On the absolute trace of polynomials having all zeros in a sector*, Experiment. Math. 17 (2008), no. 4, 443–450.
- [F3] V. Flammang. *Trace of totally positive algebraic integers and integer transfinite diameter*, Math. Comp. **78** (2009), no. 266, 1119–1125.
- [FR] V. Flammang, G. Rhin. *On the absolute Mahler measure of polynomials having all zeros in a sector. III*, Math. Comp. **84** (2015), no. 296, 2927–2938.
- [K] L. Kronecker. *Zwei Sätze über Gleichungen mit ganzzahligen Koeffizienten*, J. reine angew. Math. **53** (1857), 173–175.
- [La] M. Langevin. *Minorations de la maison et de la mesure de Mahler de certains entiers algébriques*, C. R. Acad. Sci. Paris, **303** (1986), 523–526.

[LW] Y. Liang and Q. Wu. *The trace problem for totally positive algebraic integers*, J. Aust. Math. Soc. **90** (2011), no. 3, 341–354.

[RS] G. Rhin, C.J. Smyth. *On the absolute Mahler measure of polynomials having all zeros in a sector*, Math. Comp. **64** (209) (1995), 295–304.

[RW] G. Rhin, Q. Wu. *On the absolute Mahler measure of polynomials having all zeros in a sector II*, Math. Comp. **74** (249)(2004), 383-388.

[S] C.J. Smyth. *The mean value of totally real algebraic numbers*, Math. Comp. **42** (1984), 663–681.

[Wu] Q. Wu. *On the linear independence measure of logarithms of rational numbers*, Math. Comp. **72** (2003), 901–911.

UMR CNRS 7502. IECL, Université de Lorraine, site de Metz,
Département de Mathématiques, UFR MIM,
3 rue Augustin Fresnel BP 45112 57073 Metz cedex 3 E-mail address : valerie.flammang@univ-lorraine.fr