



HAL
open science

Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen

Adélaïde Lerebours, Svetlana Murzina, You Song, Knut Erik Tollefsen, Maura Benedetti, Francesco Regoli, Jeanette M Rotchell, Jasmine Nahrgang

► To cite this version:

Adélaïde Lerebours, Svetlana Murzina, You Song, Knut Erik Tollefsen, Maura Benedetti, et al.. Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen. *Marine Environmental Research*, 2021, 170, pp.105434. 10.1016/j.marenvres.2021.105434 . hal-03298908

HAL Id: hal-03298908

<https://hal.science/hal-03298908v1>

Submitted on 25 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Susceptibility of polar cod (*Boreogadus saida*) to a model carcinogen**

2

3 Adélaïde Lerebours^{1,2,*}, ³Svetlana Murzina, ⁴You Song, ⁴Knut Erik Tollefsen, ⁵Maura
4 Benedetti, ⁵Francesco Regoli, ⁶Jeanette M. Rotchell and ⁷Jasmine Nahrgang

5

6 ¹School of Biological Sciences, University of Portsmouth, Portsmouth, PO1 2DY, United
7 Kingdom

8 ²UMR CNRS LIENSs, Littoral Environnement et Sociétés, Université de La Rochelle, La
9 Rochelle, 17 000, France

10 ³Institute of Biology of the Karelian Research Centre of the Russian Academy of Sciences
11 (IB KarRC RAS) 185910 Petrozavodsk, Russia

12 ⁴Norwegian Institute for Water Research (NIVA), Section of Ecotoxicology and Risk
13 Assessment, Gaustadalléen 21, N-0349 Oslo, Norway

14 ⁵ Department of Life and Environmental Sciences, Polytechnic University of Marche,
15 Ancona, 60131, Italy

16 ⁶Department of Biological and Marine Sciences, University of Hull, Cottingham Road, Hull,
17 HU6 7RX, United Kingdom

18 ⁷Department of Arctic and Marine Biology, Faculty of Biosciences, Fisheries and Economics
19 UiT The Arctic University of Norway, 9037 Tromsø, Norway

20

21 **Keywords:** Carcinogenesis, polar cod, benzo(a)pyrene, cancer genes, DNA repair genes

22

23 * Corresponding author: Adélaïde Lerebours, phone number: +33 (0) 5 46 50 76 23, email:
24 adelaidelerebours@univ-lr.fr

25

26 **Abstract**

27 Studies that aim to characterise the susceptibility of the ecologically relevant and non-model
28 fish polar cod (*Boreogadus saida*) to model carcinogens are required. Polar cod were exposed
29 under laboratory conditions for six months to control, 0.03 µg BaP/ g fish/ week and 0.3 µg
30 BaP/ g fish/ week dietary benzo(a)pyrene (BaP), a reference carcinogen. The concentrations
31 of the 3-OH-BaP bile metabolite and transcriptional responses of genes involved in DNA
32 adduct recognition (*xpc*), helicase activity (*xpd*), DNA repair (*xpf*, *rad51*) and tumour
33 suppression (*tp53*) were assessed after 0, 1, 3 and 6 months of exposure, alongside body
34 condition indexes (gonadosomatic index, hepatosomatic index and condition factor).
35 Micronuclei and nuclear abnormalities in blood and spleen, and liver histopathological
36 endpoints were assessed at the end of the experiment.

37 Fish grew steadily over the whole experiment and no mortality was recorded. The
38 concentrations of 3-OH-BaP increased significantly after 1 month of exposure to the highest
39 BaP concentration and after 6 months of exposure to all BaP concentrations showing the
40 biotransformation of the mother compound. Nevertheless, no significant induction of gene
41 transcripts involved in DNA damage repair or tumour suppression were observed at the
42 selected sampling times. These results together with the absence of chromosomal damage in
43 blood and spleen cells, the subtle increase in nuclear abnormalities observed in spleen cells
44 and the low occurrence of foci of cellular alteration suggested that the exposure was below
45 the threshold of observable effects. Taken together, the results showed that polar cod was not
46 susceptible to carcinogenesis using the BaP exposure regime employed herein.

47

48

49

50

51 **Introduction**

52 Polycyclic aromatic hydrocarbons (PAHs) are a very large group of ubiquitous organic
53 compounds that can originate from petrogenic, pyrogenic, biogenic and diagenic sources. A
54 number of PAHs are identified as having carcinogenic properties and have been associated to
55 an increased incidence of liver tumours in flatfish in highly polluted environments (Malins *et al.*
56 *al.*, 1985; Myers *et al.*, 1991; Harshbarger and Clark, 1990; Vogelbein *et al.*, 1990; Baumann
57 and Harshbarger, 1998). This pathology has been used to monitor the effects of exposure to
58 PAHs and the health of marine ecosystem since the 1980s (Malins *et al.*, 1985; Veethaak and
59 Ap Rheinallt, 1992) and its assessment recommended by the International Council for
60 Exploration of the Sea (ICES) and the Oslo and Paris Convention (OSPAR) Joint
61 Assessments and Monitoring Programme (JAMP) (Lyons *et al.*, 2010).

62 Benzo(a)pyrene (BaP) is a well-known pyrogenic carcinogen in a plethora of animals
63 such as marine mammals (Acevedo-Whitehouse *et al.*, 2018; Poirier *et al.*, 2019), fish (Wang
64 *et al.*, 2010; Wills *et al.*, 2010) and mice (Kasala *et al.*, 2015; Chen *et al.*, 2019). The
65 reference oral dose below which no effect is expected is 3.10^{-4} $\mu\text{g BaP /g per day}$, based on
66 animal and human studies (reviewed in EPA/635/R-17/003). The metabolites generated by
67 endogenous metabolism (biotransformation) are highly genotoxic. Phase I biotransformation
68 of BaP is mediated by cytochrome P450 (CYP) enzymes and produces highly reactive
69 metabolic intermediates such as diol-epoxide, dihydrodiol and 3-hydroxybenzo(a)pyrene (3-
70 OH-BaP) (Karle *et al.*, 2004; Zhu *et al.*, 2008; Rey-Salgueiro *et al.*, 2011). Those metabolites
71 form DNA adducts that interfere with DNA repair and replication (Phillips and Arlt, 2007).
72 This represents a critical event in the initiation of tumorigenesis, potentially leading to
73 mutations within specific regions of DNA, such as proto-oncogenes and tumour suppressor
74 genes (Rotchell *et al.*, 2001, Du Corbier *et al.*, 2005, Lerebours *et al.*, 2014, 2016). The
75 carcinogenicity of BaP has been well studied in several temperate fish species where specific

76 DNA adducts are used as markers for exposure and potential genotoxic effects. Exposure to
77 BaP specifically caused DNA adducts in fish such as pale chub (*Zacco platypus*) (Lee *et al.*,
78 2014) and killifish species (*Fundulus grandis* and *F. similis*) (Willett *et al.*, 1995; Rose *et al.*,
79 2000, 2001). Moreover, BaP exposure was associated with neoplastic lesions in brown
80 bullhead (*Ameiurus nebulosus*) (Ploch *et al.*, 1998), English sole (*Parophrys vetulus*)
81 (Reichert *et al.*, 1998) and rainbow trout (Hendricks *et al.*, 1985). PAH-induced lesions have
82 also recently been suggested in marine mammals such as harbour porpoises (*Phocoena*
83 *phocoena*) (Acevedo-Whitehouse *et al.*, 2018) and beluga whales (Poirier *et al.*, 2019).
84 Pollution induced cancer affects many aquatic species and represents a growing concern for
85 aquatic wildlife (for a review see Baines *et al.*, 2021).

86 Tumourigenesis is a progressive process characterised by different stages for which
87 the underlying molecular steps and the role of environmental exposure are not always well-
88 known. Nonetheless, liver tumourigenesis has been well studied in flatfish (Stentiford *et al.*,
89 2010; Lerebours *et al.*, 2013; 2014; 2017) and in the model fish Japanese medaka (*Oryzias*
90 *latipes*) (Rotchell *et al.*, 2001) and zebrafish (*Danio rerio*) (Li *et al.*, 2017; 2019). DNA
91 repair mechanisms have been associated with tumour formation perturbing several steps of
92 the nucleotide excision repair (NER) pathway, which recognises and repairs DNA adducts
93 induced by numerous environmental mutagens, including PAHs (Gillet and Schärer, 2006;
94 Rastogi *et al.*, 2010). While such mechanisms involved in the development of tumours in
95 certain fish species are well characterised, a substantial knowledge gap exists for non-model
96 and ecologically important species inhabiting remote regions in particular. The Arctic is
97 currently experiencing a rapid decline in sea ice (Kumar *et al.*, 2021) that may lead to a
98 significant increase in marine shipping (Ho *et al.*, 2010), oil and gas exploration and
99 operation (Elias, 2018), and tourism (Meier *et al.*, 2014) and associated release of potential
100 carcinogenic contaminants (Elias, 2018). The polar cod (*Boreogadus saida*) is a keystone fish

101 species in the arctic marine ecosystem due to its abundance, distribution and central role in
102 the food web (Welch *et al.*, 1992). Polar cod has been considered a model fish for arctic
103 ecotoxicology studies (Jonsson *et al.*, 2010; Nahrgang *et al.*, 2009, 2010a,b,c). The toxicity of
104 petroleum compounds on the physiology of polar cod has been well studied (Geraudie *et al.*,
105 2014; Bender *et al.*, 2016; Nahrgang *et al.*, 2016; Vieweg *et al.*, 2018; Nahrgang *et al.*, 2019)
106 but the tumourigenic potential of a potent carcinogen remains unknown in that species. A few
107 studies however have reported a potential susceptibility to carcinogenic contaminants
108 including BaP. The hepatic metabolism of BaP is particularly efficient in polar cod and a
109 significant increase of covalently bound reactive intermediates of BaP in the bile of fish has
110 been found after dietary exposure to BaP (Ingebrigtsen *et al.*, 2000; Bakke *et al.*, 2016).
111 These reactive intermediates were found to induce the formation of DNA adducts in the liver
112 of that species (Aas *et al.*, 2003). This genotoxic effect can in turn result in cellular
113 abnormalities and cancer initiation. Finally, a recent study showed that expression of genes
114 involved in DNA repair and cell cycle regulation processes was modified in liver of polar cod
115 dietary exposed to BaP (Song *et al.*, 2019).

116 In order to evaluate the susceptibility of polar cod to a carcinogenic compound, adult
117 specimens were exposed under laboratory conditions for six months to control, 0.03 µg BaP/
118 g fish/ week and 0.3 µg BaP/ g fish/ week dietary BaP. Selected body condition indexes, bile
119 metabolite concentrations and transcriptional responses of genes involved in DNA adduct
120 recognition (*xpc*), helicase activity (*xpd*), DNA repair (*xpf*, *rad51*) and tumour suppression
121 (*tp53*) were assessed after 0, 1, 3 and 6 months of exposure. Blood and spleen micronuclei,
122 nuclear abnormalities and liver histopathological endpoints were assessed at the end of the
123 experiment.

124 The sampling times were selected because carcinogenesis is a long-term process.
125 They were comparable to the exposure durations used in several studies interested in

126 carcinogenesis in European eel (Nogueira *et al.*, 2006), brown bullhead and channel catfish
127 (Ploch *et al.*, 1985) rainbow trout (Hendricks *et al.*, 1985, Black *et al.*, 1985) and coho
128 salmon (Black *et al.*, 1985) exposed to BaP. The BaP doses selected were lower than the
129 concentrations frequently used in previous studies. They were 10 and 100 times lower than
130 the concentration of 3 µg BaP/g of fish /week (Colli-Dula *et al.*, 2018) that induced a
131 decrease of body indexes in Nile tilapia after one month of exposure. In addition, our highest
132 concentration was 4 times lower than the lowest concentration used in the study of Song *et*
133 *al.*, (2019) that found gene expression changes in polar cod after two weeks of exposure, a
134 twelve times shorter exposure duration.

135

136 **Methods**

137 *Fish collection and exposure*

138 Adult polar cod (4 years old) were collected along the west coast of the Svalbard archipelago
139 (Norway) onboard RV Helmer Hanssen in January 2014 using a Campelen bottom trawl (at
140 200m depth) and a fish-lift (Holst and McDonald, 2000). At the Tromsø aquaculture research
141 station (Havbrukstasjon i Tromsø), fish were kept in 3000 L acclimation tank under a natural
142 light and temperature (1.5 - 3 °C) regime of 79°N (based on mooring data in Wallace *et al.*,
143 2010). During this period, fish were fed until satiation with thawed *Calanus sp.* copepods
144 (*Calanus AS*, Tromsø). Ninety fish were selected based on similar length (15 ± 1 cm) and
145 weight (25 ± 7 g) for the experiment (June 2014).

146 Polar cod were dietarily exposed to 0, 0.03 and 0.3 µg BaP per gram fish per week,
147 for 6 months (2nd of July 2014 to 31st January 2015). The experiment was conducted in
148 compliance with the policies of the Norwegian animal welfare authorities (application ID
149 6571). Briefly, a BaP (Sigma Aldrich, St. Louis, USA) solution in acetone was mixed with
150 *Calanus spp* (*Calanus AS*) to yield 0.5 or 5 µg BaP per g feed or acetone alone (acetone

151 control). The acetone was volatilized by constant stirring on a magnetic stirrer for 2.5 hours
152 at 30 °C. Small pellets were then created with the addition of 0.5 mL gelatin per g feed. Fish
153 were fed pellets corresponding to 4% of their body wet weight (bw) 5 days a week. On the
154 first, third and fifth day of a week, fish were exposed to dietary BaP or a solvent control by
155 receiving the 2% bw exposed feed (or solvent control) and 2% bw of unexposed feed (no
156 BaP, no acetone). Feeding was done by distributing the pellets to the surface of the tank.
157 Thus, feeding hierarchies may have occurred resulting in some intra-tank individual exposure
158 variations. On the remaining 2 days of a week, all fish were fed 4% bw of unexposed feed.
159 The amount of food given to each tank was adjusted at each sampling point to account for
160 both growth and sampling of specimens. With this feeding regime, the fish nominally
161 received an average of 0, 0.03 and 0.3 µg BaP per gram of fish per week. After 1 (2nd
162 August), 3 (3rd October) and 6 (31th of January) months, 10 fish per condition were
163 anaesthetized and killed by a sharp blow to the head. Total body weight (g) and fork length
164 (cm) were measured and the presence of parasites recorded. Liver and gonads were removed
165 and weighed. Bile was snap frozen in liquid nitrogen and stored at -80°C until 3-OH BaP
166 metabolite determination. A liver section was snap frozen in liquid nitrogen and stored at -
167 80°C for molecular analyses. During the final sampling (6 months of exposure), a
168 standardized liver cross-section was fixed for 24 hrs in neutral buffered formaldehyde (4%)
169 before being transferred to 70% ethanol for subsequent histological assessment. Blood and
170 spleen samples were preserved in Carnoy solution (3 methanol: 1 acetic acid) and stored at
171 +4°C for subsequent identification of nuclear abnormalities and micronuclei. Finally, somatic
172 weight (g) was determined as weight of eviscerated fish. Gonadosomatic index (GSI) and
173 hepatosomatic index (HSI) were calculated as follows:

174
$$\text{GSI} = (\text{gonad weight} / \text{somatic weight}) \times 100$$

175
$$\text{HSI} = (\text{liver weight} / \text{somatic weight}) \times 100.$$

176 *3-OH-benzo[a]pyrene measurement*

177 Biliary 3-OH-benzo[a]pyrene metabolite concentration was determined after 1, 3 and 6
178 months of exposure following the procedure detailed in Song *et al.*, (2018). Preparation of
179 hydrolysed bile samples was performed as described in Krahn *et al.*, (1992). Briefly, bile (1-
180 20 μ L) was mixed with an internal standard (triphenylamine) and diluted with demineralised
181 water (10-50 μ L) and hydrolysed with β -glucuronidasearylsulphatase (20 μ L, 1 h at 37
182 $^{\circ}$ C). Methanol (75-200 μ L) was added and the sample was mixed thoroughly before
183 centrifugation. The supernatant was then transferred to vials and analysed. High pressure
184 liquid chromatography (Waters 2695 Separations Module) was used to separate 3-OH-BaP in
185 a Waters PAH C18 column (4.6 \times 250 mm, 5 μ m particle size). The mobile phase consisted
186 of a gradient from 40:60 acetonitrile:ammonium acetate (0.05 M, pH 4.1) to 100%
187 acetonitrile at a flow rate of 1 mL/min, and the column was heated to 35 $^{\circ}$ C. A 2475
188 fluorescence detector measured fluorescence at the optimum for each analyte
189 (excitation/emissions: 380/430). A total of 25 μ L extract was injected for each analysis. The
190 results were calculated by use of the internal standard method (Grung *et al.*, 2009). The
191 calibration standards utilized were obtained from Chiron AS, Trondheim, Norway, and were
192 in the range 0.2-200 ng/g. Values below the limit of detection were considered as equal to 0
193 ng/g in the analyses.

194 *Histopathological analyses*

195 Livers were processed in a vacuum infiltration processor (Shandon Citadel 1000) using
196 standard histological protocols (Feist *et al.*, 2004). The tissues were embedded in paraffin
197 using an STP-120 spin tissue processor (Thermo Fisher Scientific, USA). Sections of 4 μ m
198 thickness were cut using a microtome HM 450 (Thermo Fisher Scientific, USA) and
199 subsequently stained with haematoxylin and eosin (H&E). The liver sections were examined

200 for microscopic pre-tumour and tumour lesions according to BEQUALM and ICES criteria
201 (Feist *et al.*, 2004). The pre-tumour lesions sought were the vacuolated, basophilic and
202 eosinophilic foci of cellular alteration (FCA). Tumour lesions were the benign hepatocellular
203 adenoma and the malignant hepatocellular carcinoma (HCC). Lesions associated to nuclear
204 and cellular polymorphism, cell death, inflammation and regeneration were also examined. A
205 total of 5, 6 and 4 fish were assessed from control, low and high exposure condition,
206 respectively.

207 *Micronucleus test and nuclear abnormalities*

208 The micronuclei and nuclear abnormalities frequencies were measured in blood and spleen of
209 polar cod tissues fixed in Carnoy's solution; subsequently separated cells were dispersed on
210 glass slides, and stained with the fluorescent dye 4',6-diamidino- 2-phenylindole at 100 ng/
211 mL. For each experimental condition, a range of 6 to 8 fish were investigated, and for each
212 specimen 2000 cells with preserved cytoplasm were scored to assess the presence of
213 micronuclei and nuclear abnormalities. Micronuclei are defined as round structures, smaller
214 than 1/3 of the main nucleus diameter, on the same optical plan and clearly separated from
215 nucleus; Nuclear abnormalities include (i) binucleated: cell with two nuclei, (ii) notch nuclei:
216 looks like nucleus but do not have nuclear materials, (iii) nuclear bud: evagination of bud-like
217 structure from the nucleus, and (iv) blebbed nuclei: small euchromatin evagination of the
218 nuclear membrane (Gorbi *et al.*, 2009; Islam *et al.*, 2021).

219 *Gene expression analyses*

220 Following 1, 3 and 6 months of exposure, a cross section of each liver, next to the one
221 dedicated to histological analyses at 6 months was used for gene transcriptional response
222 analyses. Total RNAs were extracted using the High Pure RNA Tissue kit (Roche
223 Diagnostics Ltd, West Sussex, U.K.) according to the supplier's instructions which included
224 a DNase treatment. RNA quality (integrity of 18S and 28S ribosomal bands) was evaluated

225 by electrophoresis on a 1% agarose-formaldehyde gel. RNA purity was assessed by
226 measuring the ratios of absorbance: A_{260}/A_{280} and A_{260}/A_{230} using a spectrophotometer
227 (NanoDrop, ThermoFisher). All samples were of high purity (ratios' values > 2.1).

228 First strand cDNAs were synthesized from 1 μ g of total RNA using the AffinityScript
229 Multiple Temperature cDNA Synthesis Kit (Agilent Technologies, Stockport, U. K.) using
230 random hexamer primers and according to the supplier's instructions. Putative coding
231 sequences (Figure S1) were identified by nucleotide and protein BLAST searches on the
232 NCBI database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and sequence homologies across fish
233 species on the EMBL-EBI platform (<https://www.ebi.ac.uk/Tools/msa/clustalo/>). The contigs
234 produced in the study of Song *et al.*, (2019) were also used. Primer pairs and FAMTM-
235 TAMRATM dye probes used to amplify the target sequences were designed using the Prime
236 Express software (Applied Biosystem) (Table 1). Ten ng of the reverse transcribed product
237 measured by a qubit fluorometer (Thermo Fisher Scientific) was used as a template for
238 subsequent polymerase chain reaction (PCR) in a 20 μ L final volume using 1x of TaqMan[®]
239 Fast Advanced Master Mix (Life technologies, Paisley, U.K.), 900 nM primers and 250 nM
240 probe (final concentrations) according to the supplier's protocol. PCR reactions were
241 performed in the Applied BiosystemsTM ViiATM 7 Real-Time PCR System using the
242 following programme: one cycle at 95°C for 20 s and 40 amplification cycles at 95°C for 3 s
243 and 60°C for 30 s. Primer efficiencies were determined by 10 times dilution series of the
244 cDNA template and were about 100%. The optimal normalization gene was selected by
245 testing the expressions of 3 reference genes (*β tubulin*, *hprt1* and *28S*) on all the samples
246 using the NormFinder algorithm. The expression of the *β tubulin* gene displayed the highest
247 stability. The melting curves were carefully checked after each qPCR run. The gene
248 expression was calculated according to the delta delta Ct method.

249

250 *Statistical analyses*

251 Statistical analyses were performed using R (version 3.1.2). The effect of the BaP exposure
252 concentration and time of exposure were assessed on all the biological parameters measured
253 using 1-way ANOVA. When the normality of the residuals was not verified by the Shapiro-
254 Wilk test, the non-parametric Kruskal-Wallis tests were used. Post-hoc comparisons were
255 performed using the least-square mean test for parametric test and the Wilcoxon rank test for
256 non-parametric test. The α error was adjusted using the Bonferroni correction for each post-
257 hoc test.

258 **Results**

259 Over the course of the six-month exposure, all specimens grew significantly in weight and
260 underwent gonadal maturation with mean GSI ranging from 1.5 ± 0.7 to 20.8 ± 3.9 % (Table
261 2). Endoparasites were commonly found across all treatments and sampling times.
262 Nematodes on the liver surface were the most common parasites with a frequency of
263 occurrence of 26%. Parasites of the phylum Platyhelminthes were less common (7%). No
264 mortality was observed.

265 The dietary BaP exposure of polar cod led to a dose-dependent production of biliary
266 3-OH BaP metabolites for the low ($0.03 \mu\text{g BaP/g}$ of fish/week) and high ($0.3 \mu\text{g BaP/g}$ of
267 fish/week) BaP exposure conditions after 1 and 6 months of exposure (Figure 1). The
268 concentrations of bile 3-OH BaP ranged from 20 to 40 ng/g of bile for the low exposure
269 condition and were approximately 10 times higher, from 132 to 390 ng/g of bile, for the
270 highest exposure condition.

271 The transcriptional responses related to DNA adduct recognition (*xpc*), helicase
272 activity (*xpd*), DNA repair (*xpf*, *rad51*) and tumour suppression (*tp53*) were not significantly
273 changed by any BaP dietary exposures as compared to controls ($p > 0.05$) (Figure 2).

274 The number of micronuclei recorded in the blood and spleen of polar cod ($p > 0.05$)
275 exposed to BaP did not significantly vary as compared to control (Figure 3 A, B). Nuclear
276 abnormalities in polar cod spleen were significantly increased in the high dose group ($p =$
277 0.03), while close to significant in the low dose group ($p = 0.057$) (Figure 3 C, D). No
278 significant nuclear abnormalities were observed after BaP exposure in blood cells ($p > 0.05$).

279 Histopathological analyses revealed one basophilic focus of cellular alteration in liver
280 of two individuals exposed to the low exposure condition after six months of exposure
281 (Figure S2). No tumour-related lesions were observed in livers of control and highly exposed
282 individuals.

283 **Discussion**

284 The present study showed that polar cod grew steadily through the entire experiments for all
285 exposure conditions (0, 0.03 and 0.3 μg BaP/g of fish/ week). These results were expected
286 under chronic low dose exposure scenarios and are consistent with similar results obtained in
287 polar cod exposed to higher dietary BaP concentrations (1.2 and 60.9 μg BaP/ g of fish/
288 week) but for a shorter period of two weeks (Song *et al.*, 2019). In Nile tilapia exposed to
289 intraperitoneal injections of 3 μg BaP/g of fish/ week for 4 weeks, K, GSI and GSI were
290 slightly decreased (Colli-Dula *et al.*, 2018). This suggests that there is a dose and time
291 dependent threshold above which exposure to BaP induce significant body condition indexes
292 changes. The fish species and the mode of BaP administration (injected intraperitoneally
293 versus dietary) may also be important factors to consider.

294 The bile concentration of 3-OH-BaP has been used as an indicator of BaP exposure
295 and biotransformation in many fish species including polar cod (Baake *et al.*, 2016; Baali *et*
296 *al.*, 2016; Kammann *et al.*, 2017; Song *et al.*, 2019). Indeed, previous studies led on polar cod
297 exposed to either PAHs or crude oil have shown a very high correlation between bile
298 metabolites of PAHs and both cyp1a mRNA expression and EROD activity (Bakke *et al.*,

299 2016, Bender *et al.*, 2016, Vieweg *et al.*, 2018, Nahrgang *et al.*, 2019, Song *et al.*, 2019). The
300 increase in biliary 3-OH-BaP metabolite concentrations after 1 and 6 months exposure
301 supported that of a similar, albeit shorter, exposure study (Song *et al.*, 2019). When exposed
302 to a four times higher exposure dose (1.2 µg BaP/ g of fish/ week) than the highest dose used
303 in the present study, a 3-OH-BaP concentration of 800 ng/ g of bile was found, which was
304 two to six times higher than the metabolite concentration range identified in our study. The
305 reactive BaP intermediates have been found to accumulate and covalently bind DNA in the
306 biliary system of polar cod one month after exposure to a single dietary concentration (Baake
307 *et al.*, 2016) equivalent to the cumulative dose received in the high BaP exposure condition
308 during the first month of our study. Those reactive BaP metabolites covalently bind to
309 biological molecules such nucleic acids and form DNA adducts that can lead to tumour
310 formation. For instance, higher levels of BaP-7,8-diol metabolites and DNA binding activity
311 were found in bile of English sole (*Parophrys vetulus*) a fish species more sensitive to
312 carcinogenesis than the more resistant starry flounder (*Platichthys stellatus*) (Varanasi *et al.*,
313 1986). In polar cod dietarily exposed to higher BaP concentrations (from 5 µg BaP/ g of fish
314 in a single injection) DNA adducts were found (Aas *et al.*, 2003), revealing an increased risk
315 of liver tumour formation later on. Indeed, 50% of rainbow trout displayed pre-tumour
316 (basophilic FCA) and tumour (HCC) liver lesions after six months of exposure to a similar
317 dose injected intraperitoneally (Hendricks *et al.*, 1985). In their study, 25 % of the trouts
318 displayed similar liver lesions after twelve months of dietary exposure to a high dose of BaP
319 (estimated to 1-2 mg BaP/ g fish/ week) (Hendricks *et al.*, 1985). In the present study, the
320 potential genotoxic damage generated by the BaP metabolites produced did not cause
321 significant tumour lesions. This could be the result of several factors potentially in
322 combination, including low dose, low exposure duration and effective DNA repair
323 mechanisms.

324 The DNA repair system and cell cycle regulators can prevent DNA adducts and the
325 onset of tumorous events. In the present study, the transcriptional response of genes involved
326 in the nucleotide excision repair (NER) process (*xpc*, *xpd*, *xpf*), DNA double strand breaks
327 repair (*rad51*) and cell division regulation (*tp53*) did not vary significantly after 1, 3, and 6
328 months of exposure to both BaP dietary concentrations. The exposure levels of BaP may have
329 been too low to cause significant accumulation of cell damage and trigger a significant gene
330 transcriptional response. Interestingly, a dose-specific transcriptional response of some genes
331 has been observed in liver of polar cod dietary exposed to BaP (Song *et al.*, 2019). For
332 instance, some genes involved in apoptosis (*bax* and *casp9*), a process that eliminates
333 damaged cells and prevent the proliferation of abnormal cells in tumour formation, were
334 upregulated in polar cod exposed to the high exposure level (60.9 µg BaP/ g of fish/ week).
335 The expression levels of those genes were not modified in fish exposed to the low exposure
336 dose (1.2 µg BaP/ g of fish/week) suggesting a threshold above which gene transcription is
337 modified (Song *et al.*, 2019). The basal gene expression level may also be sufficient to repair
338 DNA and/or delay the cell cycle to maintain the genetic integrity. Moreover, the DNA repair
339 gene measured in our study, *rad51*, may not be involved in the repair of specific DNA
340 damage induced. Similarly to the results herein, this gene was not differentially expressed in
341 liver of polar cod dietary exposed to BaP (Song *et al.*, 2019). *Rad51* is involved in the repair
342 of DNA double strand breaks, which belong to a different pathway than the NER. The
343 mechanism of DNA damage induced by BaP exposure is more likely to involve DNA
344 adducts than double strand breaks. Other genes involved in DNA repair processes and control
345 of cell cycle have been found induced at higher exposure regimes. For example, the gene
346 encoding for the growth arrest and DNA damage inducible beta gene (*gadd45b*) was induced
347 in the liver of the tropical fish, Nile Tilapia (*Oreochromis niloticus*) after one month of
348 exposure to 3 µg BaP/ g of fish/week (Colli-Dula *et al.*, 2018). In polar cod exposed to 1.2

349 and 60.9 µg BaP/g of fish/ week, genes involved in the excision DNA repair process (such as
350 *hmg2b* and *rad23a*) were differentially expressed (Song *et al.*, 2019). Shorter timepoints
351 may have also been necessary to observe a gene expression modulation as an early response
352 to stressors. In the liver of polar cod, transcriptional responses of genes involved in DNA
353 damage repair were changed after two weeks of dietary exposure to BaP (Song *et al.*, 2019).
354 Some studies using a reference genotoxic compound showed that DNA damage was rapidly
355 repaired with increased transcription of DNA repair genes such as *rad51* in zebrafish larvae,
356 as early as 6 hours (Reinardy *et al.* 2013). The addition of early sampling times seems
357 relevant to include in future studies.

358 Other biological processes such as detoxification mechanisms could have prevented
359 polar cod from the genotoxic effects of BaP exposure. Activation of detoxification events
360 could explain the resistance of polar cod to BaP exposure and the absence of liver tumours in
361 the present study. Variation in the expression of genes and proteins belonging to the
362 cytochrome P450 family involved in phase I of BaP detoxification process has been well
363 described in liver of fish (Nahgang *et al.*, 2009; Lee *et al.*, 2014; Colli-Dula *et al.*, 2018).
364 Interestingly, *cyp1a1* and *cyp1b1* genes were upregulated in liver of polar cod following
365 dietary exposure to 60.9 µg BaP/g of fish/ week but were not differentially expressed after
366 exposure to a lower dose of BaP (1.2 µg BaP/g of fish/ week) (Song *et al.*, 2019). This
367 suggests a dose threshold for activating the detoxification mechanisms during a chronic
368 exposure. Activation of genes and proteins involved in phase II detoxification process has
369 been also described in liver of fish exposed to BaP (Nahgang *et al.*, 2009). For instance,
370 *gstA1* gene expression was modified in the liver of Nile tilapia exposed to 3 µg BaP/ g of
371 fish/ week (Colli-Dula *et al.*, 2018). Interestingly, GST activity was higher in starry flounder,
372 a tumour resistant species, than in English sole, a tumour sensitive species, after exposure to
373 a BaP dose that induced carcinogenesis (Varanasi *et al.*, 1987). Finally, phase III

374 detoxification process based on active efflux of chemicals by ATP-binding cassette (ABC)
375 transporters could be involved in BaP elimination. For example, a rainbow trout ABCG2
376 transporter was found to interact with BaP (Zaja *et al.*, 2016).

377 In accordance with relatively low biliary BaP metabolite concentrations observed
378 throughout the exposure and limited responses of genes involved in DNA damage
379 identification and repair, no effect of dietary BaP on the micronuclei and nuclear
380 abnormalities was recorded in the present study except in spleen where nuclear abnormalities
381 increased in the high dose group. Micronuclei are formed during the anaphase stage of the
382 cell division. They are considered as a reliable index of chromosomal breakage,
383 chromosomal loss and cellular spindle malfunction (Bolognesi and Hayashi, 2011).
384 Additionally, micronuclei constitute an irreversible form of genotoxic damage compared to
385 DNA strand breaks and their induction are regulated by a large number of experimental
386 carcinogens, including chlorinated hydrocarbons, benzidine, aflatoxins, methylcholanthrene,
387 and common carcinogenic pollutants, such as PAHs, heavy metals, and pesticides (Bolognesi
388 and Hayashi, 2011). Many research studies reported the increased in micronuclei frequency
389 in erythrocytes of different fish species exposed to PAHs (Shirmohammadi *et al.*, 2018).
390 Contrary to micronuclei, nuclear abnormalities origin has not been clearly explained; some
391 suggest that nuclear abnormalities can be a primary response, prior to the micronuclei
392 formation, highlighting their relevance in the evaluation of genotoxic damage (Bolognesi and
393 Hayashi, 2011; Seriani *et al.*, 2011). An increase of erythrocytic nuclear abnormalities and
394 strand breaks was observed in eels (*Anguilla anguilla* L.) and juvenile sea bass
395 (*Dicentrarchus labrax*) exposed to a range of 0.3 to 2.7 μM of BaP and naphthalene (Maria *et*
396 *al.*, 2002; Teles *et al.*, 2003; Gravato and Santos, 2002), while on the contrary, lower
397 concentrations of BaP (0.1 μM) did not affect DNA integrity (Nogueira *et al.*, 2006). The
398 induction of micronuclei and other nuclear abnormalities were also caused by crude oil

399 exposure in turbot (*Scophthalmus maximus*) and Atlantic cod (*Gadus morua*) (Baršienė *et al.*,
400 2004; 2006). The exposure duration and levels are extremely important in determining
401 micronuclei and nuclear abnormalities formation; long-term chemical exposures can cause
402 genetic changes and consequently physiological alterations or pathologies including cancer
403 development (Depledge and Hopkin, 1995). In a study led on the European flounder,
404 *Platichthys flesus*, Köhler and Ellesat, (2008), first suggested that nuclear anomalies inside
405 liver lesions of hepatocellular cancers were correlated with micronuclei frequencies in fish
406 blood and that the histopathological grading of cancers from preneoplastic, benign to
407 malignant types was clearly associated with micronuclei increase.

408 The present study showed that polar cod were consistently exposed to dietary BaP
409 through the entire experiment and biotransformed the mother compound to intermediate
410 metabolites. However, this exposure did not lead to significant changes in the transcription of
411 selected genes, nor in chromosomal alterations and significant tissue lesions. Some early
412 responses to stress may have occurred prior to the first sampling time point at one month of
413 exposure, and basal expression of genes or potentially activated compensatory mechanisms
414 may have been sufficient to control the damage caused by the reactive metabolites.
415 Moreover, protective mechanisms such as detoxification and apoptosis could have prevented
416 the cells from the accumulation of cell damage caused by the reactive metabolites. Therefore,
417 we deduce that the BaP exposure concentrations were below the threshold of observable
418 effects. As a whole, our results showed that polar cod exposed to 0.03 and 0.3 µg BaP/ g fish/
419 week was not sensitive to the model carcinogen and liver carcinogenesis. The present work
420 encourages the addition of earlier sampling points and indicators of detoxification
421 mechanisms in future studies.

422

423

424

425

426

427

428

429

430

431 **References**

432

433 Aas E, Liewenborg B, Grøsvik BE, Camus L, Jonsson G, Børseth JF, Balk L. 2003. DNA
434 adduct levels in fish from pristine areas are not detectable or low when analysed using the
435 nuclease P1 version of the ³²P-postlabelling technique. *Biomarkers*, 8: 445-460.

436

437 Acevedo-Whitehouse K, Cole KJ, Philips DH, Jepson PD, Deaville R, Arlt VM. 2018.
438 Hepatic DNA Damage in Harbour Porpoises (*Phocoena phocoena*) Stranded along the
439 English and Welsh Coastlines. *Environmental and Molecular Mutagenesis*, 59: 613-624.

440

441 Baali A, Kammann U, Hanel R, El Qoraychy I, Yahyaoui A. 2016. Bile metabolites of
442 polycyclic aromatic hydrocarbons (PAHs) in three species of fish from Morocco.
443 *Environmental Sciences Europe*. 28, 25.

444

445 Baines C, Lerebours A, Thomas F, Fort J, Kreitsberg R, Gentes S, Meitern R, Saks L, Ujvari
446 B, Giraudeau M, Stepp T. 2021. Linking pollution and cancer in aquatic environments: A
447 review. *Environment International* 149: 106391.

448

449 Bakke MJ, Nahrgang J, Ingebrigtsen K. 2016. Comparative absorption and tissue distribution
450 of ¹⁴C-benzo(a)pyrene and ¹⁴C-phenanthrene in the polar cod (*Boreogadus saida*) following
451 oral administration. *Polar biology*, 39: 165–1173.

452

453 Baršienė J, Dedonytė V, Rybakovas A, Andreikėnaitė L, Andersen OK. 2006. Investigation
454 of micronuclei and other nuclear abnormalities in peripheral blood and kidney of marine fish
455 treated with crude oil. *Aquatic Toxicology*, 78: S99-S104

456

457 Baršienė J, Lazutka J, Šyvokienė J, Dedonytė V, Rybakovas A, Bagdonas E, Bjornstad A,
458 Andersen OK. 2004. Analysis of micronuclei in blue mussels and fish from the Baltic and
459 North Seas. *Environmental Toxicology*, 19: 365-371.

460

461 Baumann, PC, Harshbarger JC. 1998. Long term trends in liver neoplasm epizootics of brown
462 bullhead in the Black River, Ohio. *Environmental Monitoring and Assessment* 53: 213-223.

463

464 Bender ML, Frantzen M, Vieweg I, Falk-Petersen IB, Johnsen HK, Rudolfson G, Tollefsen
465 KE, Dubourg P, Nahrgang J. 2016. Effects of chronic dietary petroleum exposure on
466 reproductive development in polar cod (*Boreogadus saida*). *Aquatic Toxicology*, 180: 196–
467 208.

468

469 Black JJ, Maccubbin AE and Martin M. 1985. A Reliable, Efficient, Microinjection
470 Apparatus and Methodology for the In Vivo Exposure of Rainbow Trout and Salmon
471 Embryos to Chemical Carcinogen. *Journal of the National Cancer Institute*, 75: 1123-8.

472

473 Bolognesi C, Hayashi M. 2011. Micronucleus assay in aquatic animals. *Mutagenesis*, 26:
474 205–213.
475

476 Chen W, Zhang H and Liu Y. 2019. Anti-Inflammatory and Apoptotic Signaling Effect of
477 Fucoxanthin on Benzo(A)Pyrene-Induced Lung Cancer in Mice. *Journal of Environmental*
478 *Pathology, Toxicology and Oncology*, 38: 239-251.
479

480 Colli-Dula RC, Fang X, Moraga-Amador D, Albornoz-Abud N, Zamora-Bustillos R, Conesa
481 A, Zapata-Perez O, Moreno D, Hernandez-Nuñez E. 2018. Transcriptome analysis reveals
482 novel insights into the response of low-dose benzo(a)pyrene exposure in male tilapia. *Aquatic*
483 *Toxicology*, 162-173.
484

485 Depledge M.H., Hopkin S.P. 1995. Methods to assess effects on brackish, estuarine, and
486 near-coastal water organisms. Wiley (Ed.), *SCOPE 53-Methods to Assess the Effects of*
487 *Chemicals on Ecosystems, the Scientific Committee On Problems of the Environment*
488 *(SCOPE)*.
489

490 Du Corbier, FA, Stentiford GD, Lyons BP, Rotchell JM. 2005. Isolation of the
491 retinoblastoma cDNA from the marine flatfish dab (*Limanda limanda*) and evidence of
492 mutational alterations in liver tumors. *Environmental Science and Technology*, 39: 9785-
493 9790.
494

495 Elias, 2018. Conservation issues: Tundra ecosystems. *Encyclopedia of the Anthropocene*,
496 2, 2018,165-175
497

498 EPA/635/R-17/003Fa. Toxicological Review of Benzo[a]pyrene. 2017. U.S. Environmental
499 Protection Agency Washington, DC.

500

501 Feist SW. 2004. Biological effects of contaminants: use of liver pathology of the European
502 flatfish dab (*Limanda limanda L.*) and flounder (*Platichthys flesus L.*) for monitoring
503 (International Council for the Exploration of the Sea, Copenhagen).

504

505 Geraudie P, Nahrgang J, Forget-Leray J, Minier C, Camus L. 2014. In vivo effects of en
506 vironmental concentrations of produced water on the reproductive function of polar cod
507 (*Boreogadus saida*). Journal of Toxicology and Environmental Health A, 77: 557–573.

508

509 Gillet LCJ and Scharer OD. 2006. Molecular mechanisms of mammalian global genome
510 nucleotide excision repair. Chemical Reviews, 106: 253–276.

511 Gorbi S., Benedetti M., Virno Lamberti C., Pisanelli B., Moltedo G., Regoli F. 2009.
512 Biological effects of diethylene glycol (DEG) and produced waters (PWs) released from
513 offshore activities: A multi-biomarker approach with the sea bass *Dicentrarchus labrax*.
514 Environmental Pollution, 157: 3166-3173.

515

516 Gravato C, Santos MA. 2002. Juvenile sea bass liver P450, EROD induction, and
517 erythrocytic genotoxic responses to PAH and PAH-like compounds. Ecotoxicology and
518 Environmental Safety, 51: 115-127

519

520 Harshbarger JC, Clark JB. 1990. Epizootiology of neoplasms in bony fish of North America.
521 Science of the Total Environment, 94: 1-32.

522

523 Hendricks JD, Meyers TR, Shelton DW, Casteel JL, Bailey GS. 1985. Hepatocarcinogenicity
524 of benzo[a]pyrene to rainbow trout by dietary exposure and intraperitoneal injection. Journal
525 of national cancer institute, 74 : 839-851.

526

527 Ho, J. 2010. The implications of Arctic sea ice decline on shipping. Marine Policy, 34 : 713–
528 715.

529

530 Holst, JC, McDonald, A. 2000. Fish-lift: a device for sampling live fish with trawls. Fish.
531 Research, 48 : 87–91.

532

533 Ingebrigtsen K, Christiansen JS, Lindhe Ö, Brandt I. 2000. Disposition and cellular
534 binding of ³H-benzo(a)pyrene at subzero temperatures: studies in an aglomerular arctic
535 teleost fish - the polar cod (*Boreogadus saida*). Polar Biology, 23: 503-509.

536

537 Islam MJ, Slater MJ, Thiele R, Kunzmann A. 2021. Influence of extreme ambient cold stress
538 on growth, hematological, antioxidants, and immune responses in European seabass,
539 *Dicentrarchus labrax* acclimatized at different salinities. Ecological Indicators, 122: 107280.

540

541 Jonsson H, Sundt RC, Aas E, Sanni S. 2010. The Arctic is no longer put on ice: evalua- tion
542 of polar cod (*Boreogadus saida*) as a monitoring species of oil pollution in cold waters.
543 Marine Pollution Bulletin, 60: 390–395.

544

545 Kammann U, Akcha F, Budzinski H, Burgeot T, Gubbins MJ, Lang T, Le Menach K,
546 Vethaak AD, Hylland K. 2017. PAH metabolites in fish bile: from the seine estuary to
547 Iceland. Marine Environmental Research, 124: 41–45.

548

549 Karle IL, Yagi H, Sayer JM, and Donald MJ. 2004. Crystal and molecular structure of a
550 benzo[a]pyrene 7,8-diol 9,10-epoxide N2-deoxyguanosine adduct: Absolute configuration
551 and conformation. Proceedings of the National Academy of Sciences of the United States of
552 America, 6: 1433–1438.

553

554 Kasala ER, Bodduluru LN, Barua CC, Sriram CS, Gogoi R. 2015. Benzo(a)pyrene induced
555 lung cancer: Role of dietary phytochemicals in chemoprevention. Pharmacological Reports.
556 67: 996-1009.

557

558 Köhler A, Ellesat K. 2008. Nuclear changes in blood, early liver anomalies and
559 hepatocellular cancers in flounder (*Platichthys flesus* L.) as prognostic indicator for a higher
560 cancer risk? Marine Environmental Research, 66 : 149-150

561

562 Krahn, MM, Burrows DG, Ylitalo GM, Brown DW, Wigren CA, Collier TK, Chan SL,
563 Varanasi U. 1992. Mass-spectrometric analysis for aromatic-compounds in bile of fish
564 sampled after the Exxon-Valdez oil-spill. Environmental Science and Technology, 26: 116–
565 126.

566

567 Kumar A, Yadav J, Mohan R. 2021. Spatio-temporal change and variability of Barents-Kara
568 sea ice, in the Arctic: Ocean and atmospheric implications. Science of the Total Environment,
569 753: 142046.

570

571 Lee, JW, Kim, YH, Yoon, S, Lee SK. 2014. Cytochrome P450 System Expression and DNA
572 Adduct Formation in the Liver of *Zacco platypus* Following Waterborne Benzo(a)pyrene

573 Exposure: Implications for Biomarker Determination. *Environmental Toxicology*, 29: 1032-
574 1042.

575

576 Lerebours A, Bignell JP, Stentiford GD, Feist SW, Lyons BP, Rotchell JM. 2013. Advanced
577 diagnostics applied to fish liver tumours: relating pathology to underlying molecular
578 aetiology. *Marine Pollution Bulletin*, 72 : 94-98.

579

580 Lerebours A, Stentiford GD, Lyons BP, Bignell JP, Derocles SA, Rotchell JM. 2014. Genetic
581 alterations and cancer formation in a European flatfish at sites of different contaminant
582 burdens. *Environmental Science and Technology*, 48 :10448-10455.

583

584 Lerebours A, Chapman E, Lyons BP, Bignell JP, Stentiford GD, Rotchell JM. 2017.
585 Hepatocellular adenoma in a European flatfish (*Limanda limanda*): genetic alterations in
586 laser-capture micro-dissected tissue and global transcriptomic approach. *Marine Pollution*
587 *Bulletin*, 119: 120-127.

588

589 Li Y, Li H, Spitsbergen JM, Gong Z. 2017. Males develop faster and more severe
590 hepatocellular carcinoma than females in *krasV12* transgenic zebrafish. *Scientific Reports*, 7:
591 41280.

592

593 Li H, Li Y, Lu JW, Huo X, Gong Z. 2019. Liver-specific androgen receptor knockout
594 attenuates early liver tumor development in zebrafish. *Scientific Reports*, 9: 10645.

595

596 Lyons, BP, Thain, JE, Stentiford, GD, Hylland, K, Davies, IM, Vethaak, AD. 2010. Using
597 biological effects tools to define Good Environmental Status under the European Union
598 Marine Strategy Framework Directive. *Marine Pollution Bulletin*, 60 : 1647-1651.
599

600 Malins, DC, Krahn MM, Brown DW, Rhodes LD, Myers MS, McCain BB, Chan SL. 1985.
601 Toxic-Chemicals in Marine Sediment and Biota from Mukilteo, Washington - Relationships
602 with Hepatic Neoplasms and Other Hepatic-Lesions in English Sole (*Parophrys Vetulus*).
603 *Journal of the National Cancer Institute*, 74: 487-494.
604

605 Maria VL, Correia AC, Santos MA. 2002. *Anguilla anguilla* L. biochemical and genotoxic
606 responses to benzo[a]pyrene. *Ecotoxicology and Environmental Safety*, 53: 86-92.
607

608 Meier WN, Hovelsrud GK, van Oort BEH, Key JR, Kovacs KM, Michel C, Haas C, Mats A
609 G, Gerland S, Donald KP, Makshtas A, Reist JD. 2014. Arctic sea ice in transformation: A
610 review of recent observed changes and impacts on biology and human activity. *Reviews of*
611 *Geophysics*, 51: 185–217
612

613 Myers, MS, Landahl JT, Krahn MM, McCain BB. 1991. Relationships between hepatic
614 neoplasms and related lesions and exposure to toxic chemicals in marine fish from the U.S.
615 west coast. *Environmental Health Perspectives*, 90: 7-15.

616 Nahrgang J, Camus L, Gonzalez P, Goksøyr A, Christiansen J, Hop H, 2009. PAH biomarker
617 responses in polar cod (*Boreogadus saida*) exposed to benzo(a)pyrene. *Aquatic Toxicology*.
618 14, 309-319.

619 Nahrgang J, Camus L, Broms F, Christiansen JS, Hop H. 2010a. Seasonal baseline levels of
620 physiological and biochemical parameters in polar cod (*Boreogadus saida*): implications for
621 environmental monitoring. *Marine Pollution Bulletin*, 60: 1336–1345.

622 Nahrgang J, Camus L, Carls MG, Gonzalez P, Jonsson M, Taban IC, Bechmann RK,
623 Christiansen JS, Hop H. 2010b. Biomarker responses in polar cod (*Boreogadus saida*)
624 exposed to the water soluble fraction of crude oil. *Aquatic Toxicology*, 97: 234–242.

625 Nahrgang J, Camus L, Gonzalez P, Jonsson M, Christiansen JS, Hop H. 2010c. Biomarker
626 responses in polar cod (*Boreogadus saida*) exposed to dietary crude oil. *Aquatic Toxicology*,
627 96: 77–83.

628 Nahrgang J, Dubourg P, Frantzen M, Storch D, Dahlke F, Meador JP. 2016. Early life stages
629 of an arctic keystone species (*Boreogadus saida*) show high sensitivity to a water-soluble
630 fraction of crude oil. *Environmental Pollution*, 218: 605–614.

631 Nahrgang J, Bender ML, Meier S, Nechev J, Berge J, Frantzen M. 2019. Growth and
632 metabolism of adult polar cod (*Boreogadus saida*) in response to dietary crude oil.
633 *Ecotoxicology and Environmental Safety*, 180: 53–62.

634 Nogueira PR, Lourenço J, Mendo S, Rotchell LM. 2006. Mutation analysis of ras gene in the
635 liver of European eel (*Anguilla anguilla* L.) exposed to benzo[a]pyrene. *Marine Pollution*
636 *Bulletin*, 52: 1611-1616.

637

638 Phillips and Artlt. 2007. The ³²P-postlabeling assay for DNA adducts. *Nature Protocols*, 2:
639 2772–2781.

640

641 Ploch SA, King LC, Kohan MJ, Di Giulio RT.1998. Comparative *in Vitro* and *in Vivo*

642 Benzo[*a*]pyrene–DNA Adduct Formation and Its Relationship to CYP1A Activity in Two
643 Species of Ictalurid Catfish. *Toxicology and applied pharmacology* 149: 90–98.

644 Poirier et al 2019. Intestinal Polycyclic Aromatic Hydrocarbon-DNA Adducts in a Population
645 of Beluga Whales with High Levels of Gastrointestinal Cancers *Environ Mol Mutagen*, 60:
646 29–41.

647

648 Rastogi A, Richa, Kumar A, Tyagi MB, Sinha RP. 2010. Molecular mechanisms of
649 ultraviolet radiation-induced DNA damage and repair. *Journal of Nucleic Acids*, 2010:
650 592980.

651

652

653 Reinardy KC, Dharamshi J, Jha AN, Henry TB. 2013. Changes in expression profiles of
654 genes associated with DNA repair following induction of DNA damage in larval zebrafish
655 *Danio rerio*. *Mutagenesis*, 28: 601-608.

656

657 Rei-Salgueiro L, Costa J, Ferreira M Reil-Henriques MA. 2011. Evaluation of 3-hydroxy-
658 benzo[*a*]pyrene levels in Nile tilapia (*Oreochromis niloticus*) after waterborne exposure to
659 Benzo[*a*]pyrene. *Ecotoxicology*, 2040-2054.

660

661 Reichert WL, Myers MS, Peck-Miller K, French B, Anulacion BF, Collier TK, Stein JE,
662 Varanasi U. 1998. Molecular epizootiology of genotoxic events in marine fish: Linking
663 contaminant exposure, DNA damage, and tissue-level alterations. *Mutation Research*, 411:
664 215-225

665

666 Rose, WL, French, BL, Reichert, WL, Faisal, M. 2001. Persistence of benzo[a]pyrene–DNA
667 adducts in hematopoietic tissues and blood of the mummichog, *Fundulus heteroclitus*.
668 *Aquatic Toxicology*, 52: 319–328.

669

670 Rose, WL, French, BL, Reichert, WL, Faisal, M., 2000. DNA adducts in hematopoietic
671 tissues and blood of the mummichog (*Fundulus heteroclitus*) from a creosote-contaminated
672 site in the Elizabeth River, Virginia. *Marine Environmental Research*, 50: 581–589.

673

674 Rotchell, JM, Blair, JB, Shim, JK, Hawkins, WE, Ostrander, GK. 2001. Cloning of the
675 Retinoblastoma cDNA from the Japanese medaka (*Oryzias latipes*) and preliminary evidence
676 of mutational alterations in chemically-induced retinoblastomas. *Gene*, 263: 231-237.

677

678 Seriani R, Tavares Ranzani-Paiva M J., Silva- Souza A. T., Silva Roseli N. 2011.
679 Hematology, micronuclei and nuclear abnormalities in fishes from São Francisco River,
680 Minas Gerais state, Brazil. *Acta Scientiarum Biological Sciences*. 33: 107–112.

681

682 Shirmohammadi M., Mohammad N. S., Ronagh T., Movahedinia A., Hamidian G. 2018.
683 Using cell apoptosis, micronuclei and immune alternations as biomarkers of phenanthrene
684 exposure in yellowfin seabream (*Acanthopagrus latus*). *Fish and Shellfish Immunology*, 72:
685 37-47.

686

687 Song Y, Nahrgang Y, Tollefden KE. 2019. Transcriptomic analysis reveals dose-dependent
688 modes of action of benzo(a)pyrene in polar cod (*Boreogadus saida*). *Science of the total*
689 *environment*, 653: 176-189.

690

691 Stentiford, GD, Bignell, JP, Lyons, BP, Thain, JE, Feist, SW. 2010. Effect of age on liver
692 pathology and other diseases in flatfish: Implications for assessment of marine ecological
693 health status. *Marine Ecology Progress Series*, 411: 215-230.

694

695 Teles M, Pacheco M, Santos MA. 2003. *Anguilla anguilla* L. liver ethoxyresorufin O-
696 deethylation, Glutathione S-transferase, erythrocytic nuclear abnormalities, and endocrine
697 responses to naphthalene and β -naphthoflavone. *Ecotoxicology and Environmental Safety*,
698 55: 98-107

699

700 Varasini U, Nishimoto M, Reichert WL, Le Eberhart BT. 1986. Comparative Metabolism of
701 Benzo(a)pyrene and Covalent Binding to Hepatic DNA in English Sole, Starry Flounder, and
702 Rat. *Cancer Research*, 46: 3817-3824.

703

704 Varasini U, Stein JE, Nishimoto M, Reichert WL, Collier TK. 1987. Chemical
705 Carcinogenesis in Feral Fish: Uptake, Activation, and Detoxication of Organic Xenobiotics.
706 *Environmental Health Perspectives*, 71: 155-170.

707

708 Vethaak, AD, Rheinallt Tap. 1992. Fish disease as a monitor for marine pollution: the case of
709 the North Sea. *Reviews in Fish Biology and Fisheries*, 2: 1-32.

710

711 Vieweg I, Bilbao E, Meador JP, Cancio I, Bender ML, Cajaraville MP, Nahrgang J. 2018.
712 Effects of dietary crude oil exposure on molecular and physiological parameters related to
713 lipid homeostasis in polar cod (*Boreogadus saida*). *Comparative Biochemistry and*
714 *Physiology, Part C: Toxicology and Pharmacology*. 206-207, 54–64.

715

716 Vogelbein, WK, Fournie JW, Van Veld PA, Huggett RJ. 1990. Hepatic neoplasms in the
717 mummichog *Fundulus heteroclitus* from a creosote-contaminated site. *Cancer Research* 50:
718 5978-5986.

719

720 Wallace MI, Cottier FR, Berge J, Tarling GA, Griffiths C, et al. 2010. Comparison of
721 zooplankton vertical migration in an ice-free and a seasonally ice-covered Arctic fjord: An
722 insight into the influence of sea ice cover on zooplankton behavior. *Limnology and*
723 *Oceanography* 55: 831–845.

724

725 Wang L, Camus AC, Donga W, Thorntona C, Willetta KL. 2010. Expression of CYP1C1
726 and CYP1A in *Fundulus heteroclitus* during PAH-induced carcinogenesis. *Aquatic*
727 *Toxicology*, 99: 439–447.

728

729 Welch HE, Bergmann MA, Siferd TD, Martin KA, Curtis MF, Crawford RE Conover RJ,
730 Hop H. 1992. Energy flow through the marine ecosystem of the Lancaster Sound region,
731 Arctic Canada. *Arctic* 45, 343-375.

732

733 Willett, K, Steinberg M, Thomsen J, Narasimhan TR, Safe S, McDonald S, Beatty K,
734 Kennicutt, MC. 1995. Exposure of killifish to benzo[a]pyrene: comparative metabolism,
735 DNA adduct formation and aryl hydrocarbon (Ah) receptor agonist activities. *Comparative*
736 *Biochemistry and Physiology. Part B: Biochemistry and Molecular Biology*, 112: 93–103.

737

738 Wills, LP, Jung D, Koehn K, Zhu S, Willett KL, Hinton DE, di Giulio RT. 2010.
739 Comparative chronic liver toxicity of benzo[a]pyrene in two populations of the atlantic

740 killifish (*Fundulus heteroclitus*) with different exposure histories. Environmental Health
741 Perspectives, 118: 1376-1381.

742 Zaja R, Popovic M, Loncar J, Smital T. 2016. Functional characterization of rainbow trout
743 (*Oncorhynchus mykiss*) Abcg2a (Bcrp) transporter. Comparative Biochemistry and
744 Physiology, Part C, 190: 15–2.

745 Zhu S, Li L, Thornton C, Carvalho P, Avery BA, Willett KL. 2008. Simultaneous
746 determination of benzo[*a*]pyrene and eight of its metabolites in *Fundulus heteroclitus* bile
747 using ultra-performance liquid chromatography with mass spectrometry. Journal of
748 Chromatography B, 863: 141-149.

749

750

751 **Figure and Table Legend.**

752

753 **Figure 1.** Concentrations of biliary 3-OH-benzo[*a*]pyrene (ng/g bile, $n = 4$ to 9) in fish
754 exposed to three treatments of BaP (acetone control, low and high) during 0, 1, 3 and 6
755 months. Plots represent the median (line), 25–75% percentiles (box), non-outlier range
756 (whisker), outliers (circle) and extreme values (coloured triangle). The effect of the dose and
757 time on the metabolite concentrations were assessed using the Kruskal-Wallis rank test.
758 When significant, a Wilcoxon test and a Bonferroni correction were applied. Asterisks (*)
759 show significant difference from the control treatment ($p < 0.05$). Numbers above boxes
760 represent the n .

761 **Figure 2.** Relative expression of genes (mean \pm SD, arbitrary units) in liver of polar cods ($n =$
762 10 per treatment and time) exposed to acetone control, low and high BaP treatments after 1, 3
763 and 6 months. Plots represent the median (line), 25–75% percentiles (box), non-outlier range

764 (whisker), outliers (circle) and extreme values (coloured triangle). The effect of the dose on
765 the gene expression levels was assessed using the Kruskal-Wallis rank test. No significant
766 differences ($p > 0.05$) among treatments were found. Numbers above boxes represent the n .


767 **Figure 3.** DNA damage in the form of micronuclei per thousand in the blood (A) and spleen
768 (B) and nuclear abnormalities in the blood (C) and spleen (D) of polar cod sampled after 6
769 month of exposure. Plots represent the median (line), 25–75% percentiles (box), non-outlier
770 range (whisker), outliers (circle) and extreme values (coloured triangle). The effect of the
771 dose on the number of micronuclei and nuclear abnormalities was assessed using the
772 Kruskal-Wallis rank test. When significant, a Wilcoxon test and a Bonferroni correction were
773 applied. Asterisks (*) show significant difference from the control treatment ($p < 0.05$).
774 Numbers above boxes represent the n .

775 **Table 1.** Sequences of primer pairs and FAM/TAMRA probes used in RT-qPCR reactions
776 for each of the target genes studied. *β tubulin* was used as the reference gene.

777 **Table 2.** Fulton condition (K), hepatosomatic index (HSI), gonadosomatic index (GSI), liver,
778 gonad and body weight (g), and fork length (cm) (mean \pm SD, $n = 10$) and sex ratio
779 determined after 0, 1, 3, and 6 months of exposure to different BaP treatments (acetone
780 control, low and high exposures).

781

782 Figure 1



783

784

785

786

787

788

789


790

791

792

793

794 Figure 2



795

796


797

798

799

800

801 Figure 3



802

803

804

805

Gene name	Primers and Probe Sequences 5'-3'
<i>β tubulin</i>	F: GCCCGGCACCATGGA R: TGGCCGAAAACGAAGTTGTC P: TCCGGTGCTTTCGGTCAGATCTCA
<i>XPC</i>	F: GCTTCGACTTCCATGGAGGAT R: CTTCGTGCTCCTCACACACAA P: CGCATGCTGTGACCGACGGCTAC
<i>XPD</i>	F: TCATGTTCCGGAGTCCCTTATGTT R: GGAAGTGGTCCCGGAGGTA P: ACACACAGAGCCGCATTCTGAAGGC
<i>XPF</i>	F: ATCTGGACCTGGCGAGGAA R: TCCTGCTTTGCGGGTGTT P: CTGGAGCCCGCCAACGCTACC
<i>Rad51</i>	F: AAGAAGCCGATTGGAGGAAAC R: CGCCCCTTCCTCAGGTACA P: TCATGGCCCACGCCTCCACC
<i>tp53</i>	F: CCTCTGAGGGGCATGTTCTC R: GGGGCTCTTTCTTTTTTTTGG P: TCCTGGGCGCGACCGCA

807

808

809

810

811

812

813

814

815

816

817

818

819 Table 2

	0 month	1 month			3 months			6 months		
		Control	Low	High	Control	Low	High	Control	Low	High
Fork l. (cm)	15.9 ± 1.6	16.4 ± 1	16.7 ± 1.6	16.9 ± 1.7	17.8 ± 1.6	16.8 ± 1.9	17.3 ± 1.5	16.4 ± 1.5	17.2 ± 0.7	17.3 ± 1.2
Total w. (g)	25.6 ± 7.1	29.3 ± 5.6	32.4 ± 8.8	33.2 ± 9.1	38.3 ± 11.6	32.1 ± 8.5	35.6 ± 10.4	32.7 ± 10	37.3 ± 4.5	36.7 ± 8
K	5.2 ± 0.5	5.2 ± 0.4	5.5 ± 0.5	5.4 ± 0.3	5.3 ± 0.7	5.4 ± 0.5	5.3 ± 0.4	5.2 ± 0.7	5.3 ± 0.4	5 ± 0.4
Liver w. (g)	2 ± 0.9	2.2 ± 0.7	2.5 ± 0.9	2.6 ± 0.7	2.7 ± 1.2	2.3 ± 0.6	2.8 ± 1	2.7 ± 0.9	2.6 ± 0.5	3 ± 0.9
HSI	9.6 ± 3.3	9.4 ± 2.1	9.4 ± 2.1	10.4 ± 2.8	8.6 ± 2.6	9.3 ± 2.3	9.8 ± 2.1	12 ± 3.9	9.5 ± 2	11.3 ± 2.3
Gonad w. (g)	0.3 ± 0.2	0.5 ± 0.2	0.5 ± 0.3	0.5 ± 0.3	1.4 ± 0.5	1.1 ± 0.6	1.6 ± 0.9	4.2 ± 2.3	5.6 ± 1.3	5 ± 1.8
GSI	1.5 ± 0.7	2 ± 0.5	2.2 ± 1	1.8 ± 0.6	4.6 ± 1.2	4.1 ± 1.8	5.6 ± 2.3	16.9 ± 6.8	20.8 ± 3.9	19.3 ± 6.4
Sex ratio	40	60	30	50	50	40	30	50	22	50

820

821

Supplementary material

Figure S1. Putative coding sequences used to design the primers and probe for the 6 genes studied. The pink marks indicate the location of the introns in the DNA sequence. Primers were designed to overlap the introns whenever possible to check the specificity of the qPCR reactions.

>*β tubulin*

```
TCTACTACAATGAGGCCTCAGGTAATTAAGCAATGTCAAATGGATGACGATACTTCTCTCGTCTGTTTCAGCTCTAACTGTT
TCTAATCTATTAGGTGGCAAATACGTCCCCCGCGCTGTTCTGGTTCGATCTTGAGCCCCGGCACCATGGACTCTGTGAGGTCCGG
TGCTTTCGGTCAGATCTTCAGGCCAGACAACCTTCGTTTTCCGCCAGAGTGGTGCTGGCAACAACCTGGGCCAAGGGTCACTACA
CGGAAGTGCCGAGCTGGTGACTCTGTGCTCGACGTGGTGAGGAAAGAGGCAGAGAGCTGTGACTGCCTGCAGGGCTTCCAG
CTC
```

>*XPC*

```
ACGGGCTTCGACTTCCATGGAGGATACTCGCATGCTGTGACCGACGGCTACATTGTGTGTGAGGAGCACGAAGAGATTCTCAG
AGCAGCTGGGAGGAAGATCAAGCGCTCCAGAAACAGAGAGAGATTGAGAAGCGAGAGAAGCGGGCCACCACCAACTGGAAGC
TACTGGTGAAGGGCTTCTGATCAGGGAGAGGCTCCAGCTACGATACGCCAA
```

>*XPD*

```
CGGGGTGGTCCCTGACGGCATCGTGGCGTTCCTTACCAGCTACATGTACATGGAGAACATCGTGGCGTCCGGTATGAACAAG
GAATCTGGAGAACATCCAGAGGAACAAGCTGATCTTCATTGAGACGCCAGATGCTGCAGAGACCAGCATGGCTCTGGAGAAA
TACCAGGAGGCATGTGAGAACGGGAGAGAGCCATCCTTCTGTCTGTGGCCGAGGAAAAGTGTGGAAGGAATCGATTTTCT
GCACCACTTTCGGTCGGGCAGTGATCATGTTCCGAGTCCCTTATGTTTACACACAGAGCCGCATTTCTGAAAGCGCGTCTGGAGT
ACCTCCGGGACCAGTTCAGATCCGGGAGAACGACTTCCTGACGTTCCGACCCATGCGCCATGCGGCCAGTGCCTGGGCCGG
GTTCATCAGGGGCAAGACGACTACGGACTCATGATCTTCGCTGACAAACGCTACGCCGGGCGGACAAGCGGGGAAGCTGCC
CCGCTGGATCCAGGAGCACATCAGCGACGGCAGCCTGAACCTCACGGTGGACGAGACGGTGCAGCTCTCCAAGCACTTCCTGA
GGCAGATGGCCAGCCCTTCAACAGGAGGACCAGCTGGGTCTGTCACTGCTGACGATAGAACAGCTGGAGTCAGAGGAGATG
CTGAAGAAGATCAGCCAAATGGCTCACCAGGCCTGACCACAT
```

>*XPF*

```
GATCTACAAGGCCAACCGCCCCGGGAAGACGCTGCGGGTGTATTTTCTGATCTATGGAGGATCCACAGAGGAACAGAAGTATC
TCACCGCGCTCTCCAAGGAGAAGAAAGCCTTCGAACACCTCATCAGGGAGAAGGCGACCATGGTTGTGCCGAGGAGCGAGAG
GGTCGAGAAGACACCAATCTGGACCTGGCGAGGAATCTGGAGCCCGCCAACGCTACCACCAACACCCGCAAGCAGGAGGCCA
GGACCAGCCAGGGAGCCCTCCCGGTTCATCGTGGACATGCGGGAGTTCGCGAGCGAGCTGCCCTCCCTGCTGCACCGCCGCG
GGCTGGATCATCGAGCCGTCACCTGGAGGTGGGCGACTACATCCTGACGGCGGACACCTGCGTGGAGCGCAAGAGCGTGAGC
GACCTGATCGGCTCGCTGCAGAGCGGCCCGCTTACACGCAGTGCCTGTCCATGACCGCTACTACAAGCGCGCCGTGCTGCT
CATCGAGTTCGACCCGGCCAAGCCCTTCTCGTGGTGGCGCGCTCCGAGTTCGCCACGAGCTGTGCGCCAACGACGTCACGT
CCAAGCTGACGCTGCTACCCCTGCATTCGCGCGCTGGCCCTCTCTGGTGCCCTCGCCCCACGCCAGCGCCGAGCTCTTC
GAGGAGATGAAGCGGGGCGCGGCGAGCCGAGCGCCGCGCCCGCGCAGGCCATCGCGGCCGATCGGACGCCAGGACGACGGC
GGAGCTGTACAACCCGGCGCCGTACGACTTCCTGCTGAAGATGCCGGGGTCAACGCCAAGAACGTCGGGGCGCTGGTGAGCA
AGGCGGACAGCCTGGCCGCGCTGGCCGAGTTCAGCCAGGAGAGGCTGGCGCAGGCTCCTGGGGCACACCGGCCAACGCCAAGATG
CTCTACGAGTTCCTGCACAACGTGGCCGACGTGCCCGCCAGCTGCCAAGGGCAGACGGACGTGAAGGGAAGACT
```

>*rad51*

```
ATGGCTATGAGGAGTGAAGTGCCTTTGGAGGAGGAGGTAGAGGTGGAGGAGAACTTCGGCCCTCAGCCCATCAGTTCGACTGGA
GCAAAGCGGTGTGAGCAGCAGTGACCTGAAGAAGCTGGAGGAGGCGGGCTTCCACACCATCGAGGCCGTGGCTACACCCCCA
AGAAAGAGCTGCTCCACATCAAGGGCATCAGCGAGGCCAAGGCCGACAAGATCCTGCGGAGGCAGCCAAGCTGGTGCCCATG
GGCTTACCACGGCAACGGAGTTCACCAGCGGGCGGGCGGAGATCATCCAGATCTCCACCGCTCCAAGGAGCTGGACAAATTT
GTTAATGGGGGAATGGAGACGGGTTCATCACGGAGATGTTTGGAGAGTTCGGACGGGGAAGACACAGCTGTGCCACACTC
TGGCTGTCACTGCCAGTGCCTGCCCCATCGACCAGGGCGGGGAGAGGGCAAGGCCATGTACATCGACACCGAGGGAACCTTCCGG
CCGGAGCGCTGCTGGCCGTGGCCGAGAGGTACGGGCTGGTGGGCGAGCAGCTCCTGGACACGTTGGCGTACGCCCGGCCCTT
CAACACCGACCACCAGACCCAGCTGCTGTACCAGGCCCTGCGCCATGATGGCAGAGTACCTATGCCCTGCTCATCGTGGACA
GCGCCACCGCTCTGTACCGGACGGACTACTCCGGTCCGGGGGAGCTGTCCGGCCGGCAGGGCCACCTCGGCCGCTTCCCTCCGC
ATGCTGCTGCGGCTGGCCGACGAGTTTGGCGTTCGCGTGGTGTATCACAACAGGTTGGTGGCCAGGTTGGACGGGGCGGCCAT
GTTCTCGGCCGACCCCAAGAAGCCGATTGGAGGAAACATCATGGCCACGCTCCACCACGCGCTGTACCTGAGGAAGGGGC
GGGGAGAGACCCGATCTGTAAGATCTACGACTCCCCCTGCCTCCCGGAGTCCGGAGGCCATGTTCCGCCATCAACGCCGACGGC
GTGGGCGACGCCAAGGACTGA
```

>p53

```
CGACGCTCCTTCTGAACTACATGTGCAACAGCTCCTGCATGGGAGGGATGAACCGGAGAGCCATCCTGACCATCCTGACCCTG
GAGTCCTCTGACGGGCATGTTCTCGGGCGGCGTTGCTTCGAGGTGCGCGTCTGTGCCTGTCTGGGCGCGACCGCAAGACGGA
GGAGGGCAACGTGGAGAAAAAGACGGAGGGATCCAAGCCCACCAAAAAAAGAAAAGAGCCCCCACTCGGCCCCACGGCT
CCGCCAAGAGGGTCTGTCCGCTCCAGCGCTGAAGAGGAGGATAAGGAGGTGTTTGTGCTACAGGTCGTTGGCCGGAAGAG
ATTTCGAGATCCTGAGGCAGATAAACGATGCACTCGCGCTGCAGGAGAGGATGACAGTCAAGCAGGAGGTCCAAGGAGGGCCGT
CGCGGGAAAGAGACGGCTGGGGGACCGGACAGACGAGGGGACCGACTGAGCGACCGTCCAACCGACCGTCCAACATAACACT
GCCACAGCGAACCCCGTTATTTTCTACACTTTTCTTTTGTCAATTCCTATTTTATTTTTTTTCATAGCTTTTTTTTCAGCATA
TAGTTTTATATACAATGTATTTTTATTTTTCATGTATTTTTTCTTTCTTGATACTAATTCTTTTTTATTTTCGTTTTTATAAG
AGGCCATG
```

Figure S2. Basophilic foci of cellular alteration diagnosed in liver of a polar cod exposed to the low BaP concentration (magnification x40).

