



HAL
open science

Fundamental limits for learning hidden Markov model parameters

Kweku Abraham, Elisabeth Gassiat, Zacharie Naulet

► **To cite this version:**

Kweku Abraham, Elisabeth Gassiat, Zacharie Naulet. Fundamental limits for learning hidden Markov model parameters. 2021. hal-03301592

HAL Id: hal-03301592

<https://hal.science/hal-03301592>

Preprint submitted on 27 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fundamental limits for learning hidden Markov model parameters

Kweku Abraham

Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay,
91405 Orsay, France
kweku.abraham@universite-paris-saclay.fr

Elisabeth Gassiat

Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay,
91405 Orsay, France
elisabeth.gassiat@universite-paris-saclay.fr

Zacharie Naulet

Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay,
91405 Orsay, France
zacharie.naulet@universite-paris-saclay.fr

Abstract

We study the frontier between learnable and unlearnable hidden Markov models (HMMs). HMMs are flexible tools for clustering dependent data coming from unknown populations. The model parameters are known to be identifiable as soon as the clusters are distinct and the hidden chain is ergodic with a full rank transition matrix. In the limit as any one of these conditions fails, it becomes impossible to identify parameters. For a chain with two hidden states we prove nonasymptotic minimax upper and lower bounds, matching up to constants, which exhibit thresholds at which the parameters become learnable.

1 Introduction

We consider the learnability of hidden Markov model parameters. Hidden Markov models offer an attractive approach to clustering problems. Such problems are ubiquitous in machine learning. A natural probabilistic approach to modelling the clusters views data as independent draws from a mixture of distributions and seeks to identify the labels. Without strong assumptions about the form of the distributions, this approach fails: an unconstrained mixture model is non-identifiable. This difficulty can be overcome by assuming that the clusters are well separated.

An alternative approach is to remove the assumption that the data points are independent. Perhaps the simplest, but nevertheless a powerful, alternative structure is to model the cluster labels as a time-homogeneous Markov chain; then the data itself is described as following a hidden Markov model (HMM). In sharp contrast to the independent setting, with hidden Markov structure one can recover the distribution of data for each cluster absent virtually any constraint on these distributions: indeed, once all submodels in which the data is in fact independent are ruled out, no further constraint is necessary when there are two hidden states. [With more than two hidden states, current state-of-the-art results have the mild further requirement that the Markov transition matrix be full-rank.] Note that, given consistent estimators of the model parameters, one can construct an empirical Bayes classifier which mimics the optimal clustering performance of the Bayes classifier.

The present work initiates an exploration of the limits of learnability of the hidden Markov parameters as the independent subcase is approached. We focus on the setting of two hidden states and multinomial data, and exhibit principles which should generalise to much wider settings. Our main result, [Theorem 1](#), gives upper and lower bounds showing the minimax estimation rate for the model parameters, exhibiting that these parameters can be learned if and only if the sample size n is large enough compared to a suitable measure of the closeness of the data to the independent subcase.

2 Related work

Clustering is one of the most classical tasks in modern machine learning. Given data (Y_1, \dots, Y_n) , one seeks a partition where Y_i and Y_j are in the same block if they share similarities: typically a proximity condition to a certain centroid. Algorithms for clustering can be divided into two categories according to whether they rely on geometric arguments or probabilistic modelling. Geometric approaches usually do not assume any generating model for (Y_1, \dots, Y_n) , but rather that the clusters can be identified from the data via a suitable separation condition. Undoubtedly the most famous algorithm of this type is the celebrated *k-means* algorithm [29] whose ability to recover well-separated clusters has been investigated in [22].

The other main class of clustering algorithms posit a probabilistic generating model for (Y_1, \dots, Y_n) and seeks to recover hidden cluster labels associated with the generating model. One of the most popular models assumes the Y_i 's are drawn independently from a mixture of normal distributions, where the cluster labels identify from which component a data point is drawn. The model parameters are traditionally learned using the famous EM algorithm [12], and points are assigned to the cluster with the closest centroid. Let us again emphasise that a general independent mixture model is non-identifiable, so that the assumption of gaussianity plays an essential role in allowing the recovery of the clusters. A solution to allow less rigid modelling of the data distribution for each cluster is to drop the independence condition and to use a dependent probabilistic approach, such as the HMM approach considered here.

HMMs have long been used as a flexible tool for clustering questions: see for instance [9, 25, 23, 35, 38, 37]. In this context the distributions of data for the clusters are known as the emission distributions. The proof that parameters can be identified with minimal assumptions is comparatively recent, given for HMMs taking values in a finite set in [4, 21, 5] and extended to allow for emission distributions modelled nonparametrically (but still with the underlying Markov chain having finite state space) in [15, 3]. Results in [11] control the propagation of errors from parameter estimation to the posterior probabilities when calculating the latter via plug-in, implying that good control on the risk of the estimators will ensure the performance of the empirical Bayes classifier is close to that of the true Bayes classifier (whose optimality for clustering is a standard result in decision theory [13]).

A topic closely related to binary classification/clustering is multiple testing, in which one aims to identify within some large data set a collection of data points which come from a ‘discovery’ hypothesis, rather than from the conservative null hypothesis. In this setting control of the false discovery rate has been obtained recently for a knockoffs-based method in [34] and for an empirical Bayes method in [1]; in each case estimation of the HMM parameters is an essential first step. Modelling the proportion of non-null signals as vanishingly small, as our results permit, would allow for further links to the setting of *sparse* multiple testing, considered for example (with independent data) in [2, 8].

Theoretical justification of a range of learning methods for HMMs with emission distributions modelled parametrically or nonparametrically have been developed in recent years: moment and tensor methods in [5, 11], and model selection using penalized least squares estimation in [10, 27], using penalized likelihood methods in [28], or using other techniques in [26]. These works all give both asymptotic and nonasymptotic upper bounds controlling the distance between estimators and the unknown parameters. All require the data to truly be dependent, but none quantify explicitly how their sample complexity results depend on the ‘distance’ to independence. Indeed, quantifying this dependence requires a sharp understanding of how the distances between distributions evolve with respect to the distances between parameters, as done for particular parametric finite mixture models in [18, 20, 14].

To the best of our knowledge no prior theoretical result exists addressing the learning of parameters of a HMM when approaching the independent case. A phase transition between learnable and

unlearnable HMMs as exhibited in our current work was previously exhibited experimentally in [33] when the multinomial emission distributions approach each other.

Finally, let us mention that departure from the independence assumption has been noted to allow much better possible learning also in HMM settings free from the assumption that the Markov chain has a finite state space [17, 6] (at the expense of stricter assumptions on the emission distributions), and also in other problems including dynamic networks [30, 7], image denoising [31], and deconvolution [16].

3 Setting

Consider a two-state HMM with multinomial emissions, in which we observe the first n entries of a sequence $\mathbf{Y} = (Y_1, Y_2, \dots) \in \{1, \dots, K\}^{\mathbb{N}}$ which satisfies

$$\begin{aligned} \mathbb{P}(Y_n = k \mid \mathbf{X}) &= f_{X_n}(k), \\ \mathbf{X} = (X_n)_{n \in \mathbb{N}} &\sim \text{Markov}(\pi, Q), \end{aligned} \quad (1)$$

with the Y_j , $j \in \mathbb{N}$ conditionally independent given \mathbf{X} . The vector \mathbf{X} of ‘hidden states’ takes values in $\{0, 1\}^{\mathbb{N}}$ and, for a parameter $\theta = (p, q, f_0, f_1)$, the transition matrix of the chain is given by

$$Q := \begin{pmatrix} 1-p & p \\ q & 1-q \end{pmatrix}, \quad (2)$$

with the convention that for $j \geq 1$, $\mathbb{P}_\theta(X_{j+1} = 0 \mid X_j = 0) = 1 - p < 1$ and $\mathbb{P}_\theta(X_{j+1} = 0 \mid X_j = 1) = q > 0$. The densities f_0, f_1 are the ‘emission densities’ with respect to counting measure on $\{1, \dots, K\}$. Grant also that X_1 is drawn from the stationary distribution of the chain, i.e. $\mathbb{P}_\theta(X_1 = 1) = p/(p+q)$.

In the limit where the sequence \mathbf{Y} becomes independent and identically distributed (i.i.d.), learning the parameters becomes impossible due to standard identifiability issues for mixture models. This i.i.d. limit can be approached in three ways:

1. $p \approx 0$ or $q \approx 0$, and thus the chain \mathbf{X} passes long periods of time in one of the two states;
2. the transition matrix Q is nearly singular, so that \mathbf{X} itself is almost i.i.d; this is the case if $|1 - p - q| \approx 0$;
3. the emission distributions are close to each other: $\|f_0 - f_1\| \approx 0$, where $\|\cdot\|$ denotes the usual Euclidean norm, $\|f\|^2 = \sum |f(k)|^2$.

We adopt a minimax point of view and encapsulate all the above scenarios within the class of parameters defined, for some $\delta, \epsilon \in (0, 1)$ and some $\zeta > 0$, by

$$\Theta = \Theta(\delta, \epsilon, \zeta) = \{\theta : p, q \geq \delta, |1 - p - q| \geq \epsilon, \|f_0 - f_1\| \geq \zeta\}.$$

Introduce also the subset

$$\Theta_L = \Theta_L(\delta, \epsilon, \zeta) = \Theta \cap \{1 - |1 - p - q| \geq L\}.$$

Remark 1. Note that $1 - |1 - p - q|$ is the absolute spectral gap of the chain \mathbf{X} , and hence the mixing time of the chain can be upper bounded uniformly in Θ_L . Here L may be arbitrarily small but we think of it as fixed, in contrast to δ, ϵ and ζ which are allowed to depend on n . With the introduction of this lower bound we still allow one of p, q to be vanishingly small (or arbitrarily close – even equal – to 1), but not both.

Remark 2. If ζ is too large compared to $1/K$, $\Theta(\delta, \epsilon, \zeta)$ may be empty, or too small to be an interesting parameter space. To avoid this we assume a mild compatibility condition: that

$$\zeta \leq \frac{\sqrt{2\lfloor K/2 \rfloor}}{4K}. \quad (3)$$

4 Main results

To avoid a label-switching issue discussed in the next section we assume that $f_0 - f_1$ lies in some specified half-plane. Our main result is the following.

Theorem 1. *There exist an estimator $\hat{\theta} = (\hat{p}, \hat{q}, \hat{f}_0, \hat{f}_1)$ and a constant $C = C(K, L) > 0$ such that for all $1 \leq x^2 \leq n\delta^2\epsilon^4\zeta^6$*

$$\sup_{\theta \in \Theta_L} \mathbb{P}_\theta \left(\max(|\hat{p} - p|, |\hat{q} - q|) > \frac{Cx}{\sqrt{n\delta^2\epsilon^4\zeta^6}} \max(\delta, \epsilon\zeta) \right) \leq e^{-x^2},$$

$$\sup_{\theta \in \Theta_L} \mathbb{P}_\theta \left(\max(\|\hat{f}_0 - f_0\|, \|\hat{f}_1 - f_1\|) > \frac{Cx}{\sqrt{n\delta^2\epsilon^4\zeta^4}} \right) \leq e^{-x^2}.$$

Grant condition (3), assume $\delta \leq 1/6$, $\epsilon \leq 1/3$, $L \leq 1/3$ and $n\delta^2\epsilon^4\zeta^6 \geq 1$. Then for some $c = c(K) > 0$,

$$\inf_{\check{\theta}} \sup_{\theta \in \Theta_L} \mathbb{P}_\theta \left(\max(|\check{p} - p|, |\check{q} - q|) > \frac{c}{\sqrt{n\delta^2\epsilon^4\zeta^6}} \max(\delta, \epsilon\zeta) \right) \geq 1/4,$$

$$\inf_{\check{\theta}} \sup_{\theta \in \Theta_L} \mathbb{P}_\theta \left(\max(\|\check{f}_0 - f_0\|, \|\check{f}_1 - f_1\|) > \frac{c}{\sqrt{n\delta^2\epsilon^4\zeta^4}} \right) \geq 1/4,$$

where the infima are over all estimators $\check{\theta} = (\check{p}, \check{q}, \check{f}_0, \check{f}_1)$.

The estimator $\hat{\theta}$ is built via plug-in from those constructed later in [Theorem 2](#). Note that the maxima are genuinely required in the lower bounds: in the extreme case where p is close to zero and q is close to 1, one has many samples with $X_i = 0$ and few with $X_i = 1$, so that p and f_0 are easier to estimate accurately than q and f_1 .

We deduce immediately the sample complexity for learning the parameters.

Corollary 1. *Fix a target error magnitude $E > 0$ and a probability level $\alpha > 0$. For the same estimators as in [Theorem 1](#), there exists a constant $C = C(K, L)$ such that for any $\theta \in \Theta_L$ we have*

$$n \geq \frac{\log(1/\alpha)}{\delta^2\epsilon^4\zeta^6} \max\left(\frac{C\delta^2}{E^2}, \frac{C\epsilon^2\zeta^2}{E^2}, 1\right) \implies \mathbb{P}_\theta(\max(|\hat{p} - p|, |\hat{q} - q|) > E) \leq \alpha,$$

$$n \geq \frac{\log(1/\alpha)}{\delta^2\epsilon^4\zeta^4} \max\left(\frac{C}{E^2}, \frac{1}{\zeta^2}\right) \implies \mathbb{P}_\theta(\max(\|\hat{f}_0 - f_0\|, \|\hat{f}_1 - f_1\|) > E) \leq \alpha.$$

Conversely there exists a constant $c(K) > 0$ such that for all $0 < E \leq c(K)$ and for any estimator $\check{\theta} = (\check{p}, \check{q}, \check{f}_0, \check{f}_1)$ there exists $\theta \in \Theta_L$ such that

$$n \leq \frac{\max(\delta^2, \epsilon^2\zeta^2)}{\delta^2\epsilon^4\zeta^6} \implies \mathbb{P}_\theta(\max(|\check{p} - p|, |\check{q} - q|) > E) \geq 1/4,$$

$$n \leq \frac{1}{\delta^2\epsilon^4\zeta^4} \implies \mathbb{P}_\theta(\max(\|\check{f}_0 - f_0\|, \|\check{f}_1 - f_1\|) > E) \geq 1/4.$$

Note that to apply [Theorem 1](#) for the lower bounds we would initially also need $n \geq (\delta^2\epsilon^4\zeta^6)^{-1}$ but by monotonicity — i.e. the fact that any measurable function of (Y_1, \dots, Y_n) is also a measurable function of (Y_1, \dots, Y_N) for $N \geq n$ — the restriction can be removed.

5 Proof outline

We sketch out the main ideas behind the proof of [Theorem 1](#). The full proof is deferred to the supplementary material, along with all other proofs for this article.

The minimax lower bounds are obtained by an argument *à la* Le Cam. In particular, it is a famous result of Le Cam [[24](#), [36](#)] that the minimax rate (under quadratic loss) of estimating a functional $g : \Theta \rightarrow \mathbb{R}$ is always greater than the maximum value that $|g(\theta) - g(\tilde{\theta})|^2$ can take for $\theta, \tilde{\theta} \in \Theta$ under the constraint that $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)}) \leq c$, where $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)})$ denotes the *Kullback-Leibler* (KL) divergence between the laws of (Y_1, \dots, Y_n) under parameters θ and $\tilde{\theta}$, and $0 < c < 1$ is a small positive constant (see [Lemma 3](#) for the precise formulation we use). Understanding bounds on $|g(\theta) - g(\tilde{\theta})|$ in terms of bounds on $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)})$ is also sufficient for obtaining an upper bound on the minimax estimation rate. Since the observations are not i.i.d, it is not the case that $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)}) = nK(p_\theta^{(1)}; p_{\tilde{\theta}}^{(1)})$, and,

while it is true that $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)}) \leq nK(p_\theta^{(1)}; p_{\tilde{\theta}}^{(1)})$, this bound is not useful: if it were, it could equally be applied to give bounds in a general mixture model, in which parameter estimation is known to be impossible. Building on the work of [5, 15] we know that θ is identifiable from $p_\theta^{(3)}$ and we record in Lemma 2 the more useful bound $K(p_\theta^{(n)}; p_{\tilde{\theta}}^{(n)}) \leq \lceil n/3 \rceil K(p_\theta^{(3)}; p_{\tilde{\theta}}^{(3)})$, the right side of which we further bound by a constant multiple of $n\|p_\theta^{(3)} - p_{\tilde{\theta}}^{(3)}\|^2$ since the KL divergence is equivalent to the squared Euclidean distance under mild assumptions. The main difficulty of the proof is to relate $\|p_\theta^{(3)} - p_{\tilde{\theta}}^{(3)}\|$ to a suitable notion of distance between θ and $\tilde{\theta}$. We propose a reparametrisation of the model problem to simplify the analysis. Indeed, motivated by a desire to simplify the expression for $p^{(3)}$ (see equations (9) and (10) in Section 6), we introduce new parameters ϕ, ψ . A key result is Proposition 1, which establishes that $\|p_{\theta(\phi, \psi)}^{(3)} - p_{\theta(\tilde{\phi}, \tilde{\psi})}^{(3)}\|$ is equivalent to $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi})$, where ρ is defined in the proposition and can be seen as the “natural” statistical distance of the problem. The parameterization (ϕ, ψ) turns out to be of special interest: the component of ϕ determine how “close” the sequence \mathbf{Y} is to being i.i.d in an interpretable way (see Section 6), and the parameter ψ is related to the stationary distribution of the sequence \mathbf{Y} . For this reason, we also establish minimax lower bounds for the estimation of ϕ and ψ themselves in Section 8, Theorem 3. Summarizing all the paragraph above, the lower bounds for ϕ (respectively ψ) in Theorem 3 are obtained by lower bounding the value of the optimization problems $\max |\phi_j - \tilde{\phi}_j|^2$ (respectively $\max |\psi_j - \tilde{\psi}_j|^2$) subject to $n\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq c$ and $\theta(\phi, \psi), \theta(\tilde{\phi}, \tilde{\psi}) \in \Theta$ for a small enough constant $c > 0$. Finally the lower bounds for the original parameters in the Theorem 1 are essentially deduced from the bounds for (ϕ, ψ) and inversion of the parameterization.

The minimax upper bounds are obtained by producing an estimator that attains the bounds. To do so, we leverage that $p_\theta^{(3)}$ can be estimated in Euclidean distance at the parametric rate $n^{-1/2}$ by the empirical estimator $\hat{p}_n^{(3)}$ defined in Section 7, Lemma 1. Then, we produce an estimator $(\hat{\phi}, \hat{\psi})$ for (ϕ, ψ) by solving for $(\hat{\phi}, \hat{\psi}) \in \arg \min_{\phi, \psi} \|p_{\theta(\phi, \psi)}^{(3)} - \hat{p}_n^{(3)}\|$. By standard calculations and using the equivalence between $\|p_{\theta(\phi, \psi)}^{(3)} - p_{\theta(\tilde{\phi}, \tilde{\psi})}^{(3)}\|$ and $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi})$ derived in the Proposition 1, we show in Theorem 2 that any such $(\hat{\phi}, \hat{\psi})$ attains the minimax lower bounds for estimating (ϕ, ψ) . Finally, the upper bounds for the original parameters in the Theorem 1 are obtained by taking $\hat{\theta} = \theta(\hat{\phi}, \hat{\psi})$.

6 Change of parameterisation

We reparametrise the model in such a way that the i.i.d. limiting cases are highlighted, by changing variables to $\phi = (\phi_1, \phi_2, \phi_3)$ and $\psi = (\psi_1, \psi_2)$ defined as

$$\phi(\theta) = \left(\frac{q-p}{p+q} \quad 1-p-q \quad \|f_0 - f_1\| \right), \quad \psi(\theta) = \left(\frac{qf_0 + pf_1}{p+q} \quad \frac{f_0 - f_1}{\|f_0 - f_1\|} \right).$$

Here we have separated the scalar parameters ϕ from the vector parameters ψ . Defining

$$r(\phi) = \frac{1}{4}(1 - \phi_1^2)\phi_2\phi_3^2, \quad (4)$$

it follows from the discussion in Section 3 that the data \mathbf{Y} is close to i.i.d. exactly when $r(\phi) \approx 0$. [This is of course true also of other combinations of the components of ϕ , but as equation (10) will show, $r(\phi)$ is the appropriate combination measuring the “distance” to the i.i.d. case.]

Define

$$\begin{aligned} \Phi &= \Phi(\delta, \epsilon, \zeta) = \{(\phi(\theta), \psi(\theta)) : \theta \in \Theta(\delta, \epsilon, \zeta)\}, \\ \Phi_L &= \Phi_L(\delta, \epsilon, \zeta) = \{(\phi(\theta), \psi(\theta)) : \theta \in \Theta_L(\delta, \epsilon, \zeta)\}, \end{aligned}$$

and note that for $(\phi, \psi) \in \Phi$, we have

$$-\frac{1-\delta}{1+\delta} \leq \phi_1 \leq \frac{1-\delta}{1+\delta}, \quad \epsilon \leq |\phi_2| \leq 1-2\delta, \quad \phi_3 \geq \zeta, \quad |r(\phi)| \geq \delta\epsilon\zeta^2/4, \quad (5)$$

while for $(\phi, \psi) \in \Phi_L$ we additionally have

$$|\phi_2| \leq 1-L. \quad (6)$$

Remark 3. When $K = 2$, in view of identifiability issues discussed in the next subsection, ψ_2 is not needed in the parametrisation, since we may universally make the choice

$$\psi_2 = \left(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}} \right).$$

Remark 4. The parametrisation $\theta \mapsto (\phi, \psi)$ is invertible: we calculate

$$\begin{aligned} p &= \frac{1}{2}(1 - \phi_2)(1 - \phi_1), \\ q &= \frac{1}{2}(1 - \phi_2)(1 + \phi_1), \\ f_0 &= \psi_1 - \frac{1}{2}\phi_1\phi_3\psi_2 + \frac{1}{2}\phi_3\psi_2, \\ f_1 &= \psi_1 - \frac{1}{2}\phi_1\phi_3\psi_2 - \frac{1}{2}\phi_3\psi_2. \end{aligned}$$

Remark 5. Suppose ψ_1 is a probability density function with respect to counting measure on $\{1, \dots, K\}$, ψ_2 is a function satisfying $\|\psi_2\| = 1$, $\sum_k \psi_2(k) = 0$, and ϕ satisfies $|\phi_1| \leq 1$, $|\phi_2| \leq 1$ and $\phi_3 \geq 0$. Then (ϕ, ψ) lies in $\Phi(\delta, \epsilon, \zeta)$ if and only if

$$\frac{1}{2}(1 - \phi_2)(1 - |\phi_1|) \geq \delta, \quad \frac{1}{2}(1 - \phi_2)(1 + |\phi_1|) \leq 1, \quad |\phi_2| \geq \epsilon, \quad \phi_3 \geq \zeta, \quad (7)$$

$$\psi_1(k) - \frac{1}{2}\phi_1\phi_3\psi_2(k) - \frac{1}{2}\phi_3|\psi_2(k)| \geq 0, \quad \forall k \leq K. \quad (8)$$

6.1 Identifiability of the model

The model (1) is identifiable for the parameter set Θ only up to ‘label-switching’, since \mathbf{Y} has the same distribution under the parameters (p, q, f_0, f_1) and (q, p, f_1, f_0) ; in the parametrisation (ϕ, ψ) , the distribution of \mathbf{Y} is the same under $(\phi_1, \phi_2, \phi_3, \psi_1, \psi_2)$ and under $(-\phi_1, \phi_2, \phi_3, \psi_1, -\psi_2)$. However, it was proved in [5] that aside from this label-switching, the model parameters can be identified from the law of just three consecutive observations. To that end, for any integer m denoting by $P_\theta^{(m)}$ the law of m consecutive observations with parameter $\theta \in \Theta$, and by $p_\theta^{(m)}$ the corresponding density with respect to counting measure on $\{1, \dots, K\}^m$, we calculate

$$p_\theta^{(3)} = \left(\frac{q}{p+q} \right) g \otimes f_0 \otimes g + \left(\frac{p}{p+q} \right) h \otimes f_1 \otimes h, \quad (9)$$

where $g = (1-p)f_0 + pf_1$ and $h = qf_0 + (1-q)f_1$, and where \otimes denotes the tensor product so that $(f \otimes g \otimes h)(a, b, c) = f(a)g(b)h(c)$, $(a, b, c) \in \{1, \dots, K\}^3$.

In the (ϕ, ψ) parametrisation, writing, in a slight abuse of notation, just $p_{\phi, \psi}^{(3)}$ for $p_{\theta(\phi, \psi)}^{(3)}$, we have

$$\begin{aligned} p_{\phi, \psi}^{(3)} &= \psi_1 \otimes \psi_1 \otimes \psi_1 + r(\phi)(\psi_2 \otimes \psi_2 \otimes \psi_1 + \psi_1 \otimes \psi_2 \otimes \psi_2) \\ &\quad + \phi_2 r(\phi) \psi_2 \otimes \psi_1 \otimes \psi_2 - \phi_1 \phi_2 \phi_3 r(\phi) \psi_2 \otimes \psi_2 \otimes \psi_2, \end{aligned} \quad (10)$$

where we recall the notation $r(\phi) = \frac{1}{4}(1 - \phi_1^2)\phi_2^2\phi_3^2$,

Optimal estimation rates can be obtained if we adequately understand the Kullback–Leibler divergence between distributions with different parameters. The Kullback–Leibler divergence between $P_{\theta(\phi, \psi)}^{(n)}$ and $P_{\theta(\tilde{\phi}, \tilde{\psi})}^{(n)}$ can be related to the Euclidean distance between the densities $p_{\phi, \psi}^{(3)}$ and $p_{\tilde{\phi}, \tilde{\psi}}^{(3)}$ (see Lemma 2), and we define a statistical distance ρ directly on the parameter space Φ which is equivalent to this Euclidean distance. The function ρ is not a true metric because it may not satisfy the triangle inequality and because, due to the identifiability issues reflected by the appearance of factors of $\text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle)$ in its definition, we may have $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) = 0$ with $(\phi, \psi) \neq (\tilde{\phi}, \tilde{\psi})$. Here $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product on \mathbb{R}^K , $\langle f, g \rangle = \sum_{i=1}^K f(i)g(i)$.

Proposition 1. For r as in equation (4) define m by

$$m(\phi) := (r(\phi), \phi_2 r(\phi), \phi_1 \phi_2 \phi_3 r(\phi)), \quad (11)$$

and define

$$\begin{aligned} \rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) &= \max\{|m_1(\phi) - m_1(\tilde{\phi})|, |m_2(\phi) - m_2(\tilde{\phi})|, |m_3(\phi) - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot m_3(\tilde{\phi})|, \\ &\quad \|\psi_1 - \tilde{\psi}_1\|, \max\{|m_1(\phi)|, |m_1(\tilde{\phi})|\} \cdot \|\psi_2 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\|\}. \end{aligned}$$

There exist constants $c_1, c_2 > 0$ (which depend on K) such that for all $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \bigcup_{\delta, \epsilon, \zeta} \Phi(\delta, \epsilon, \zeta)$ we have

$$c_1 \rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\| \leq c_2 \rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}).$$

7 Upper bounds

We obtain the following upper bounds for estimating ϕ and ψ . Since we are studying limits as the quantities of interest become small, the relative risk may be of as much interest as the absolute risk, and we provide bounds for both quantities. The bounds demonstrate that learning model parameters is possible in the regime where n is large enough in relation to δ , ϵ and ρ . Observe firstly that estimation of $p^{(3)}$ is possible at a parametric rate.

Lemma 1. Define the empirical estimator $\hat{p}_n^{(3)} : \{1, \dots, K\}^n \rightarrow [0, 1]$ by

$$\hat{p}_n^{(3)}(a, b, c) = \frac{1}{n} \sum_{i=1}^{n-2} \mathbb{1}\{Y_i = a, Y_{i+1} = b, Y_{i+2} = c\}.$$

Then for some constant $C = C(K, L)$ and any $x \geq 1$

$$\sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{(\phi, \psi)}(\|\hat{p}^{(3)} - p^{(3)}\| \geq Cx/\sqrt{n}) \leq e^{-x^2}.$$

Theorem 2. Assume Φ_L is non-empty and let $\hat{\phi}, \hat{\psi}$ be any measurable functions satisfying

$$\|p_{\hat{\phi}, \hat{\psi}}^{(3)} - \hat{p}_n^{(3)}\| \leq 2 \inf_{(\tilde{\phi}, \tilde{\psi}) \in \Phi_L} \|p_{\tilde{\phi}, \tilde{\psi}}^{(3)} - \hat{p}_n^{(3)}\|.$$

There exists a constant $C = C(K, L) > 0$ such that the following hold.

1. Assume $1 \leq x^2 \leq n\delta^2\epsilon^4\zeta^6$. Then

$$\begin{aligned} & \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{1 - \hat{\phi}_1^2}{1 - \phi_1^2} - 1 \right|^2 \geq \frac{2Cx^2}{n\delta^2\epsilon^4\zeta^6} \right) \\ & \leq \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\min(|\hat{\phi}_1 - \phi_1|, |\hat{\phi}_1 + \phi_1|)^2 \geq \frac{Cx^2}{n\epsilon^4\zeta^6} \right) \leq e^{-x^2}. \end{aligned}$$

2. Assume $1 \leq x^2 \leq n\delta^2\epsilon^2\zeta^4$. Then

$$\begin{aligned} & \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{\hat{\phi}_2}{\phi_2} - 1 \right|^2 \geq C \frac{x^2}{n\delta^2\epsilon^4\zeta^4} \right) \\ & \leq \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(|\hat{\phi}_2 - \phi_2|^2 \geq C \frac{x^2}{n\delta^2\epsilon^2\zeta^4} \right) \leq e^{-x^2}. \end{aligned}$$

3. Assume $1 \leq x^2 \leq n\delta^2\epsilon^4\zeta^6$. Then

$$\begin{aligned} & \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{\hat{\phi}_3}{\phi_3} - 1 \right|^2 \geq C \frac{x^2}{n\delta^2\epsilon^4\zeta^6} \right) \\ & \leq \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(|\hat{\phi}_3 - \phi_3|^2 \geq C \frac{x^2}{n\delta^2\epsilon^4\zeta^4} \right) \leq e^{-x^2}. \end{aligned}$$

4. Assume $1 \leq x^2 \leq n$. Then

$$\sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\|\hat{\psi}_1 - \psi_1\|^2 \geq \frac{Cx^2}{n} \right) \leq e^{-x^2}.$$

5. Assume $1 \leq x^2 \leq n\delta^2\epsilon^2\zeta^4$ and $K > 2$. Then

$$\sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\min(\|\hat{\psi}_2 - \psi_2\|^2, \|\hat{\psi}_2 + \psi_2\|^2) \geq \frac{Cx^2}{n\delta^2\epsilon^2\zeta^4} \right) \leq e^{-x^2}.$$

Recall that estimating ψ_2 is unnecessary when $K = 2$ (see [Remark 3](#)). Note that the absolute loss in each case is bounded, and one can deduce that the bounds for ϕ_2 and for ψ hold without an upper bound on x , with e^{-x^2} on the right replaced by zero (for C large enough).

8 Lower bounds

We prove lower bounds, matching the previous upper bounds in a suitable regime and demonstrating the impossibility of learning model parameters when n is not large enough in relation to δ , ϵ and ρ . The particular value $1/4$ on the right sides in the following is not essential: what is important is that the probabilities are bounded away from zero. The lower bounds over Φ_L remain true over the larger set $\tilde{\Phi}$.

Theorem 3. *Grant the compatibility condition (3) and assume $\epsilon \leq 1/3$, $\delta \leq 1/6$ and $L \leq 1/3$. There exists a constant $c = c(K)$ such that the following hold. The infima are over all estimators, i.e. all measurable functions of the data (Y_1, \dots, Y_n) .*

1. Assume $n\delta^2\epsilon^4\zeta^6 \geq 1$. Then

$$\begin{aligned} \inf_{\hat{\phi}_1} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\min(|\hat{\phi}_1 - \phi_1|^2, |\hat{\phi}_1 + \phi_1|^2) \geq \frac{c}{n\epsilon^4\zeta^6} \right) \\ \geq \inf_{\hat{\phi}_1} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{1 - \tilde{\phi}_1^2}{1 - \phi_1^2} - 1 \right|^2 \geq \frac{2c}{n\delta^2\epsilon^4\zeta^6} \right) \geq 1/4. \end{aligned}$$

2. Assume $n\delta^2\epsilon^4\zeta^4 \geq 1$. Then

$$\begin{aligned} \inf_{\hat{\phi}_2} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(|\hat{\phi}_2 - \phi_2|^2 \geq \frac{c}{n\delta^2\epsilon^2\zeta^4} \right) \\ \geq \inf_{\hat{\phi}_2} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{\hat{\phi}_2}{\phi_2} - 1 \right|^2 \geq \frac{c}{n\delta^2\epsilon^4\zeta^4} \right) \geq 1/4. \end{aligned}$$

3. Assume $n\delta^2\epsilon^4\zeta^6 \geq 1$. Then

$$\begin{aligned} \inf_{\hat{\phi}_3} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(|\hat{\phi}_3 - \phi_3|^2 \geq \frac{c}{n\delta^2\epsilon^4\zeta^4} \right) \\ \geq \inf_{\hat{\phi}_3} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\left| \frac{\hat{\phi}_3}{\phi_3} - 1 \right|^2 \geq \frac{c}{n\delta^2\epsilon^4\zeta^6} \right) \geq 1/4. \end{aligned}$$

4. For any n , δ , ϵ and ζ ,

$$\inf_{\hat{\psi}_1} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\|\hat{\psi}_1 - \psi_1\|^2 \geq \frac{c}{n} \right) \geq 1/4.$$

5. Assume $n\delta^2\epsilon^2\zeta^4 \geq 1$ and $K > 2$. Then

$$\inf_{\hat{\psi}_1} \sup_{(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)} \mathbb{P}_{\phi, \psi} \left(\min(\|\hat{\psi}_2 - \psi_2\|, \|\hat{\psi}_2 + \psi_2\|)^2 \geq \frac{c}{n\delta^2\epsilon^2\zeta^4} \right) \geq 1/4.$$

9 Conclusions, limitations and future directions

In this work we have quantified the impact on learnability of approaching the i.i.d. boundary within the set of parameters of a hidden Markov model. The limiting cases occur when one hidden state is absorbing, when the underlying Markov chain becomes a sequence of independent variables, or when the emission distributions are equal. We have proved both upper and lower bounds for the estimation rates of the parameters in a hidden Markov models with two hidden states and finitely many possible outcomes. Our results show that a phase transition occurs near the boundary and quantify how large the sample has to be in order to get estimators with prescribed error with high probability.

Some tricky regions of the parameter space are not fully captured in the upper and lower bounds. Specifically, the condition on n in the lower bound for estimating ϕ_2 differs by a factor of ϵ^2 from the corresponding condition in the upper bound. In the upper bound for ϕ_1 , in the region $n\delta^2\epsilon^4\zeta^6 < x^2 \leq n\epsilon^4\zeta^6$ we do not obtain the correct rate of decay in the probability as x increases.

In this range we can obtain something by applying the bound with $y^2 = \min(x^2, n\delta^2\epsilon^4\zeta^6)$ but we cannot expect that this gives the correct dependence on y . [When $x^2 \geq n\epsilon^4\zeta^6$ we may replace e^{-x^2} with zero, as noted after the theorem for ϕ_2 and ψ .] A similar gap exists for estimating ϕ_3 . Our results already work for a wide range of parameters, and extending to the few remaining cases is an interesting issue for future research.

Our results are limited to the context of latent variables with two hidden states and multinomial emission distributions. We believe similar results hold with more than two hidden states and with arbitrary nonparametric emission distributions. Investigation of phase transitions for more general HMMs and misspecified modelling will be the object of further work. Developments of our findings for clustering, multiple testing and sparse settings will also be the object of further work, and all will depend fundamentally on the results obtained here.

On the practical side, usual estimation algorithms can be expected to exhibit bad computational behaviour when the unknown true parameters lie near the learning frontier. We have not tackled this issue here and we believe it merits substantive investigation, both in building robust algorithms and in detecting the poor performance in the problematic region. This last question is interesting both from a practical and a theoretical point of view.

Acknowledgments and Disclosure of Funding

This work was supported by a public grant as part of the Investissement d’avenir project, reference ANR-11-LABX-0056-LMH, LabEx LMH, and by Institut Universitaire de France.

A Proofs

The following lemma exhibits the fact that $p^{(3)}$ contains qualitatively the same information as $p^{(m)}$ for $m \geq 3$. The information here is quantified by the Kullback–Leibler divergence, defined for distributions with densities p, q with respect to some common dominating measure by

$$K(p; q) = \mathbb{E}_p \left(\log \frac{p}{q} \right).$$

Lemma 2. *For any $n \in \mathbb{N}$ and any $\theta, \tilde{\theta} \in \Theta$,*

$$K(p_{\theta}^{(n)}; p_{\tilde{\theta}}^{(n)}) \leq \lceil n/3 \rceil K(p_{\theta}^{(3)}; p_{\tilde{\theta}}^{(3)}).$$

If further for all k we have $\min(\tilde{f}_0(k), \tilde{f}_1(k)) \geq c$ for some constant $c > 0$, then

$$K(p_{\theta}^{(n)}; p_{\tilde{\theta}}^{(n)}) \leq \lceil n/3 \rceil c^{-3} \|p_{\theta}^{(3)} - p_{\tilde{\theta}}^{(3)}\|_2^2.$$

Proof. For any $k, k' \geq 1$ satisfying $k' \leq 3k$, we have

$$K(p_{\theta}^{(k')} ; p_{\tilde{\theta}}^{(k')}) \leq K(p_{\theta}^{(3k)} ; p_{\tilde{\theta}}^{(3k)}) \leq k K(p_{\theta}^{(3)} ; p_{\tilde{\theta}}^{(3)}),$$

which yields the first claim.

For the second claim, recall that $K(p; q)$ is upper bounded by the chi-square distance $\chi^2(p, q) = \mathbb{E}_q[(p/q - 1)^2]$ (e.g. [36, Lemma 2.7]). The result then follows from the fact that if \tilde{f}_0, \tilde{f}_1 are bounded below by c then $p_{\tilde{\theta}}^{(3)}$ is lower bounded by c^3 . \square

A.1 Proof of Proposition 1

Recall the definition (11) of m as

$$m(\phi) := (r(\phi), \phi_2 r(\phi), \phi_1 \phi_2 \phi_3 r(\phi)), \quad r(\phi) = \frac{1}{4}(1 - \phi_1^2)\phi_2\phi_3^2.$$

We write $\tilde{m} = m(\tilde{\phi})$, and we write ψ_{ijk} for $\psi_i \otimes \psi_j \otimes \psi_k$ and $\tilde{\psi}_{ijk}$ for $\tilde{\psi}_i \otimes \tilde{\psi}_j \otimes \tilde{\psi}_k$. Then from equation (10) we have

$$\begin{aligned} p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)} &= (\psi_{111} - \tilde{\psi}_{111}) + \{m_1(\psi_{221} + \psi_{122}) - \tilde{m}_1(\tilde{\psi}_{221} + \tilde{\psi}_{122})\} \\ &\quad + \{m_2\psi_{212} - \tilde{m}_2\tilde{\psi}_{212}\} - \{m_3\psi_{222} - \tilde{m}_3\tilde{\psi}_{222}\}. \end{aligned} \tag{12}$$

Recalling that $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product on \mathbb{R}^K , we have $\langle \psi_1, 1 \rangle = 1$, $\langle \psi_2, 1 \rangle = 0$, $\|\psi_2\| = 1$ and $\|1\| = K^{1/2}$. Let $\langle \cdot, \cdot \rangle$ also denote the Euclidean inner product on $\mathbb{R}^{3 \times 3 \times 3}$, wherein for functions $f_i, \tilde{f}_i : \{1, \dots, K\} \rightarrow \mathbb{R}$, $i \leq 3$ we have

$$\langle f_1 \otimes f_2 \otimes f_3, \tilde{f}_1 \otimes \tilde{f}_2 \otimes \tilde{f}_3 \rangle = \langle f_1, \tilde{f}_1 \rangle \langle f_2, \tilde{f}_2 \rangle \langle f_3, \tilde{f}_3 \rangle.$$

Lower bounding $\|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\|$ For any function $f : \{1, \dots, K\} \rightarrow \mathbb{R}$, we have

$$\langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, f \otimes 1 \otimes 1 \rangle = \langle \psi_{111} - \tilde{\psi}_{111}, f \otimes 1 \otimes 1 \rangle = \langle \psi_1 - \tilde{\psi}_1, f \rangle.$$

Then

$$\begin{aligned} \|\psi_1 - \tilde{\psi}_1\| &= \sup_{\|f\|=1} |\langle \psi_1 - \tilde{\psi}_1, f \rangle| \\ &= \sup_{\|f\|=1} |\langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, f \otimes 1 \otimes 1 \rangle| \\ &\leq \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\| \sup_{\|f\|=1} \|f \otimes 1 \otimes 1\| = K \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\|, \end{aligned} \quad (13)$$

and similarly,

$$\begin{aligned} \langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, 1 \otimes f \otimes f \rangle &= \langle \psi_{111} - \tilde{\psi}_{111}, 1 \otimes f \otimes f \rangle + \langle m_1 \psi_{122} - \tilde{m}_1 \tilde{\psi}_{122}, 1 \otimes f \otimes f \rangle \\ &= \langle \psi_1 - \tilde{\psi}_1, f \rangle^2 + m_1 \langle \psi_2, f \rangle^2 - \tilde{m}_1 \langle \tilde{\psi}_2, f \rangle^2. \end{aligned} \quad (14)$$

Choosing $f = \psi_2 + \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2$ (with the convention that $\text{sgn}(0) = +1$), we observe that

$$\langle \psi_2, f \rangle = 1 + |\langle \psi_2, \tilde{\psi}_2 \rangle| = \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \langle \tilde{\psi}_2, f \rangle.$$

In particular we note that $\langle \psi_2, f \rangle^2 = \langle \tilde{\psi}_2, f \rangle^2 = (1 + |\langle \psi_2, \tilde{\psi}_2 \rangle|)^2 \geq 1$. Since also $\|f\|^2 = 2 + 2|\langle \psi_2, \tilde{\psi}_2 \rangle| \leq 4$, returning to (14) we observe that

$$\begin{aligned} |m_1 - \tilde{m}_1| &\leq \|f\|^2 \|\psi_1 - \tilde{\psi}_1\|^2 + \|1 \otimes f \otimes f\| \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\| \\ &\leq 4 \|\psi_1 - \tilde{\psi}_1\|^2 + 4K^{1/2} \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\| \\ &\leq 4(K^{7/2} + K^{1/2}) \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\|, \end{aligned} \quad (15)$$

where for the last line we have used [equation \(13\)](#) and the fact that $\|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\|^2 \leq K^3$. We continue by considering the expression $f \otimes 1 \otimes f$, for which we have

$$\begin{aligned} \langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, f \otimes 1 \otimes f \rangle &= \langle \psi_{111} - \tilde{\psi}_{111}, f \otimes 1 \otimes f \rangle + \langle m_2 \psi_{212} - \tilde{m}_2 \tilde{\psi}_{212}, f \otimes 1 \otimes f \rangle \\ &= \langle \psi_1 - \tilde{\psi}_1, f \rangle^2 + m_2 \langle \psi_2, f \rangle^2 - \tilde{m}_2 \langle \tilde{\psi}_2, f \rangle^2 \end{aligned}$$

Recognising symmetry with [equation \(14\)](#), we again choose $f = \psi_2 + \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2$ to obtain

$$|m_2 - \tilde{m}_2| \leq 4(K^{7/2} + K^{1/2}) \|p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}\|. \quad (16)$$

Finally, considering the expression $f \otimes f \otimes f$, we observe that

$$\begin{aligned} \langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, f \otimes f \otimes f \rangle &= \langle \psi_{111} - \tilde{\psi}_{111}, f \otimes f \otimes f \rangle + \langle m_1(\psi_{221} + \psi_{122}) - \tilde{m}_1(\tilde{\psi}_{221} + \tilde{\psi}_{122}), f \otimes f \otimes f \rangle \\ &\quad + \langle m_2 \psi_{212} - \tilde{m}_2 \tilde{\psi}_{212}, f \otimes f \otimes f \rangle - \langle m_3 \psi_{222} - \tilde{m}_3 \tilde{\psi}_{222}, f \otimes f \otimes f \rangle. \end{aligned}$$

In other words,

$$\begin{aligned} \langle p_{\phi, \psi}^{(3)} - p_{\tilde{\phi}, \tilde{\psi}}^{(3)}, f \otimes f \otimes f \rangle &= \langle \psi_1 - \tilde{\psi}_1, f \rangle^3 + 2(m_1 \langle \psi_2, f \rangle^2 \langle \psi_1, f \rangle - \tilde{m}_1 \langle \tilde{\psi}_2, f \rangle^2 \langle \tilde{\psi}_1, f \rangle) \\ &\quad + (m_2 \langle \psi_2, f \rangle^2 \langle \psi_1, f \rangle - \tilde{m}_2 \langle \tilde{\psi}_2, f \rangle^2 \langle \tilde{\psi}_1, f \rangle) - (m_3 \langle \psi_2, f \rangle^3 - \tilde{m}_3 \langle \tilde{\psi}_2, f \rangle^3). \end{aligned}$$

Once more choosing $f = \psi_2 + \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2$, we obtain (recall that by construction $\|f\| \leq 2$, $1 \leq \langle \psi_2, f \rangle^2 = \langle \tilde{\psi}_2, f \rangle^2 \leq 4$, and also $\text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \langle \psi_2, f \rangle = \langle \tilde{\psi}_2, f \rangle$)

$$\begin{aligned} |m_3 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{m}_3| &\leq 8\|\psi_1 - \tilde{\psi}_1\|^3 + 8\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\| \\ &\quad + 8|m_1 \langle \psi_1, f \rangle - \tilde{m}_1 \langle \tilde{\psi}_1, f \rangle| + 4|m_2 \langle \psi_1, f \rangle - \tilde{m}_2 \langle \tilde{\psi}_1, f \rangle|. \end{aligned}$$

For some constant $C = C(K)$ we have

$$\begin{aligned} |m_1 \langle \psi_1, f \rangle - \tilde{m}_1 \langle \tilde{\psi}_1, f \rangle| &\leq |\langle \psi_1, f \rangle| |m_1 - \tilde{m}_1| + |\tilde{m}_1| |\langle \psi_1 - \tilde{\psi}_1, f \rangle| \\ &\leq 2\|\psi_1\| |m_1 - \tilde{m}_1| + 2|\tilde{m}_1| \|\psi_1 - \tilde{\psi}_1\| \\ &\leq C\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|, \end{aligned}$$

where for the last line we have used [equations \(13\) and \(15\)](#) and that $\|\psi_1\| \leq K^{1/2}$ and $|\tilde{m}_1| \leq \tilde{\phi}_3^2/4 \leq \|\tilde{f}_0 - \tilde{f}_1\|^2/4 \leq K/4$.

Similarly, using [equation \(16\)](#) and the fact that $|\tilde{m}_2|$ is suitably bounded, we have for some $C = C(K)$

$$|m_2 \langle \psi_1, f \rangle - \tilde{m}_2 \langle \tilde{\psi}_1, f \rangle| \leq C\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|.$$

We deduce for some different constant $C = C(K)$ that

$$|m_3 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{m}_3| \leq C\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|. \quad (17)$$

Finally, for ψ_2 we show that for some C we have

$$\max(|m_1|, |\tilde{m}_1|) \|\psi_2 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\| \leq C\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|. \quad (18)$$

If $\psi_2 = \tilde{\psi}_2$ there is nothing to prove, so we assume without loss of generality that $\psi_2 \neq \tilde{\psi}_2$. Also assume that $|m_1| \geq |\tilde{m}_1|$, the final bound then following by symmetry. Returning to [equation \(14\)](#) with f to be chosen, we see that

$$m_1(\langle \psi_2, f \rangle^2 - \langle \tilde{\psi}_2, f \rangle^2) = \langle \psi_1 - \tilde{\psi}_1, f \rangle^2 - \langle p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}, 1 \otimes f \otimes f \rangle - \langle \tilde{\psi}_2, f \rangle^2 (m_1 - \tilde{m}_1).$$

Since $\langle \psi_2, f \rangle^2 - \langle \tilde{\psi}_2, f \rangle^2 = \langle \psi_2 - \tilde{\psi}_2, f \rangle \langle \psi_2 + \tilde{\psi}_2, f \rangle$ we obtain

$$|m_1 \langle \psi_2 - \tilde{\psi}_2, f \rangle \langle \psi_2 + \tilde{\psi}_2, f \rangle| \leq \|f\|^2 \|\psi_1 - \tilde{\psi}_1\|^2 + K\|f\|^2 \|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\| + |m_1 - \tilde{m}_1| \langle \tilde{\psi}_2, f \rangle^2. \quad (19)$$

Observe that $\psi_2 + \tilde{\psi}_2$ is orthogonal to $\psi_2 - \tilde{\psi}_2$ (this arises from the fact that ψ_2 and $\tilde{\psi}_2$ have unit norms) and choose

$$f = \frac{\psi_2 + \tilde{\psi}_2}{\|\psi_2 + \tilde{\psi}_2\|} + \frac{\psi_2 - \tilde{\psi}_2}{\|\psi_2 - \tilde{\psi}_2\|};$$

note that

$$\langle \psi_2 - \tilde{\psi}_2, f \rangle \langle \psi_2 + \tilde{\psi}_2, f \rangle = \|\psi_2 - \tilde{\psi}_2\| \|\psi_2 + \tilde{\psi}_2\|.$$

Since also $\|f\| \leq 2$ and $|\langle \tilde{\psi}_2, f \rangle| \leq 2$, continuing from [equation \(19\)](#) and using [equations \(13\) and \(15\)](#) we see that for a constant $C = C(K)$

$$\begin{aligned} |m_1| \|\psi_2 - \tilde{\psi}_2\| \|\psi_2 + \tilde{\psi}_2\| &\leq 4\|\psi_1 - \tilde{\psi}_1\|^2 + 4K\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\| + 4|m_1 - \tilde{m}_1| \\ &\leq 2C\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|. \end{aligned}$$

Observing that

$$\begin{aligned} \|\psi_2 - \tilde{\psi}_2\|^2 \|\psi_2 + \tilde{\psi}_2\|^2 &= \|\psi_2 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\|^2 \|\psi_2 + \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\|^2 \\ &= \|\psi_2 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\|^2 (2 + 2|\langle \psi_2, \tilde{\psi}_2 \rangle|) \\ &\geq 2\|\psi_2 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\psi}_2\|^2, \end{aligned}$$

and recalling we assumed that $|m_1| \geq |\tilde{m}_1|$, [equation \(18\)](#) follows.

The proof that $\|p_{\phi, \psi}^{(3)} - p_{\phi, \tilde{\psi}}^{(3)}\|$ is lower bounded up to a constant by $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi})$ follows by combining [equations \(13\) and \(15\)–\(18\)](#)

Upper bounding $\|p_{\phi,\psi}^{(3)} - p_{\tilde{\phi},\tilde{\psi}}^{(3)}\|$ From [equation \(12\)](#),

$$\begin{aligned} \|p_{\phi,\psi}^{(3)} - p_{\tilde{\phi},\tilde{\psi}}^{(3)}\| &\leq \|\psi_{111} - \tilde{\psi}_{111}\| + |m_1 - \tilde{m}_1| \|\psi_{221} + \psi_{122}\| + |\tilde{m}_1| \|\psi_{221} - \tilde{\psi}_{221}\| \\ &\quad + |\tilde{m}_1| \|\psi_{122} - \tilde{\psi}_{122}\| + |m_2 - \tilde{m}_2| \|\psi_{212}\| + |\tilde{m}_2| \|\psi_{212} - \tilde{\psi}_{212}\| \\ &\quad + |m_3 - \tilde{m}_3| \|\psi_{222}\| + |\tilde{m}_3| \|\psi_{222} - \tilde{\psi}_{222}\|. \end{aligned} \quad (20)$$

Note that the bound remains valid if we replace the final two terms by

$$|m_3 + \tilde{m}_3| \|\psi_{222}\| + |\tilde{m}_3| \|\psi_{222} + \tilde{\psi}_{222}\|;$$

we focus on the case where $\text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) = +1$ for which the original decomposition yields suitable bounds, but the proof in the other case is similar using the alternative decomposition.

As used already in proving the lower bound on $\|p_{\phi,\psi}^{(3)} - p_{\tilde{\phi},\tilde{\psi}}^{(3)}\|$, we note that

$$\max(\|\psi_{221}\|, \|\psi_{122}\|, |\tilde{m}_1|, \|\psi_{212}\|, |\tilde{m}_2|, \|\psi_{222}\|, |\tilde{m}_3|) \leq C,$$

for some $C = C(K)$. To conclude the proof it thus suffices to bound the tensor product terms $\|\psi_{ijk} - \tilde{\psi}_{ijk}\|$ in terms of the differences $\|\psi_1 - \tilde{\psi}_1\|, \|\psi_2 - \tilde{\psi}_2\|$. First we decompose

$$\begin{aligned} \|\psi_1 \otimes \psi_1 \otimes \psi_1 - \tilde{\psi}_1 \otimes \tilde{\psi}_1 \otimes \tilde{\psi}_1\| &\leq \|\psi_1 \otimes \psi_1 \otimes \psi_1 - \tilde{\psi}_1 \otimes \psi_1 \otimes \psi_1\| \\ &\quad + \|\tilde{\psi}_1 \otimes \psi_1 \otimes \psi_1 - \tilde{\psi}_1 \otimes \tilde{\psi}_1 \otimes \psi_1\| + \|\tilde{\psi}_1 \otimes \tilde{\psi}_1 \otimes \psi_1 - \tilde{\psi}_1 \otimes \tilde{\psi}_1 \otimes \tilde{\psi}_1\|, \end{aligned}$$

so that

$$\|\psi_{111} - \tilde{\psi}_{111}\| \leq \|\psi_1 - \tilde{\psi}_1\| (\|\psi_1\|^2 + \|\psi_1\| \|\tilde{\psi}_1\| + \|\tilde{\psi}_1\|^2) \leq 3K \|\psi_1 - \tilde{\psi}_1\|. \quad (21)$$

We also note, recalling that ψ_2 and $\tilde{\psi}_2$ have unit norms, that

$$\begin{aligned} \|\psi_{221} - \tilde{\psi}_{221}\|^2 &= \|\psi_{221}\|^2 + \|\tilde{\psi}_{221}\|^2 - 2\langle \psi_{221}, \tilde{\psi}_{221} \rangle \\ &= \|\psi_1\|^2 + \|\tilde{\psi}_1\|^2 - 2\langle \psi_2, \tilde{\psi}_2 \rangle^2 \langle \psi_1, \tilde{\psi}_1 \rangle \\ &= \|\psi_1 - \tilde{\psi}_1\|^2 + 2\langle \psi_1, \tilde{\psi}_1 \rangle (1 - \langle \psi_2, \tilde{\psi}_2 \rangle^2) \\ &\leq \|\psi_1 - \tilde{\psi}_1\|^2 + 2\|\psi_1\| \|\tilde{\psi}_1\| |1 - \langle \psi_2, \tilde{\psi}_2 \rangle|. \end{aligned}$$

Observe that

$$\|\psi_2 - \tilde{\psi}_2\|^2 = 2(1 - \langle \psi_2, \tilde{\psi}_2 \rangle), \quad (22)$$

and hence

$$|1 - \langle \psi_2, \tilde{\psi}_2 \rangle| = |1 + \langle \psi_2, \tilde{\psi}_2 \rangle| |1 - \langle \psi_2, \tilde{\psi}_2 \rangle| \leq 2|1 - \langle \psi_2, \tilde{\psi}_2 \rangle| = \|\psi_2 - \tilde{\psi}_2\|^2.$$

We deduce that

$$\|\psi_{221} - \tilde{\psi}_{221}\|^2 \leq \|\psi_1 - \tilde{\psi}_1\|^2 + 2\|\psi_1\| \|\tilde{\psi}_1\| \|\psi_2 - \tilde{\psi}_2\|^2 \leq \|\psi_1 - \tilde{\psi}_1\|^2 + 2K \|\psi_2 - \tilde{\psi}_2\|^2. \quad (23)$$

By symmetry, the same bound holds for $\|\psi_{122} - \tilde{\psi}_{122}\|$ and for $\|\psi_{212} - \tilde{\psi}_{212}\|$. Furthermore $\|\psi_{222}\| = 1$, and using [\(22\)](#),

$$\begin{aligned} \|\psi_{222} - \tilde{\psi}_{222}\|^2 &= \|\psi_{222}\|^2 + \|\tilde{\psi}_{222}\|^2 - 2\langle \psi_{222}, \tilde{\psi}_{222} \rangle \\ &= 2 - 2\langle \psi_2, \tilde{\psi}_2 \rangle^3 \\ &= 2(1 - \langle \psi_2, \tilde{\psi}_2 \rangle) (1 + \langle \psi_2, \tilde{\psi}_2 \rangle + \langle \psi_2, \tilde{\psi}_2 \rangle^2) \\ &\leq 3\|\psi_2 - \tilde{\psi}_2\|^2. \end{aligned} \quad (24)$$

The claim follows from inserting [equations \(21\)](#), [\(23\)](#) and [\(24\)](#) into [equation \(20\)](#).

A.2 Proof of [Theorem 2](#)

We start with the proof of [Lemma 1](#), that $p^{(3)}$ can be estimated at a parametric rate.

Proof of Lemma 1. We use a Markov chain concentration result from [32]. Theorem 3.4 therein (but note there is an updated version of the paper on arXiv) tells us that for any stationary Markov chain $\mathbf{Z} = (Z^{(1)}, Z^{(2)}, \dots)$ of pseudo-spectral gap γ_{ps} (defined as in [32]) and any function h satisfying $\mathbb{E}[h(Z^{(1)})^2] \leq \sigma^2$ and $\|h\|_\infty \leq b$,

$$\mathbb{P}\left(\left|\sum_{i=1}^n h(Z^{(i)}) - \mathbb{E}h(Z^{(1)})\right| \geq x\right) \leq 2 \exp\left(-\frac{x^2 \gamma_{\text{ps}}}{8(n+1/\gamma_{\text{ps}})\sigma^2 + 20bx}\right). \quad (25)$$

We apply to the chain \mathbf{Z} defined by $Z^{(n)} = (X_n, X_{n+1}, X_{n+2}, Y_n, Y_{n+1}, Y_{n+2})$; we begin by showing the pseudo-spectral gap of this chain is bounded from below. Proposition 3.4 of the same reference shows that the reciprocal of the pseudo-spectral gap of any chain is bounded above by twice the mixing time $t_{\text{mix}}^{\mathbf{Z}}$ of the chain, defined as the first time that the law of \mathbf{Z} , regardless of the starting distribution, is within 1/4 of its invariant distribution in total variation distance. We note that $t_{\text{mix}}^{\mathbf{Z}}$ is equal to the mixing time $t_{\text{mix}}^{\mathbf{X}^{(3)}}$ of the chain $((X_n, X_{n+1}, X_{n+2})_{n \geq 0})$. This latter quantity is equal to $t_{\text{mix}}^{\mathbf{X}} + 2$ where $t_{\text{mix}}^{\mathbf{X}}$ denotes the mixing time of the chain \mathbf{X} itself. Finally, the matrix Q has eigenvalues 1 and ϕ_2 , and an explicit computation yields that $\max_{i,j} |Q_{ij}^n - \pi_j| = \max_i (\pi_i) |\phi_2|^n$ so that the mixing time of \mathbf{X} is at most

$$\left\lceil \frac{\log 4}{\log(1/|\phi_2|)} \right\rceil \leq \left\lceil \frac{\log 4}{\log(1/(1-L))} \right\rceil \leq \left\lceil \frac{\log 4}{L} \right\rceil,$$

which is a constant since L is fixed. The pseudo-spectral gap of the chain \mathbf{Z} is thus lower bounded by some constant $\gamma = \gamma(L)$.

Applying equation (25) with $h(Z) = \mathbb{1}\{Z_4 = a, Z_5 = b, Z_6 = c\}$, which satisfies $\mathbb{E}h^2 \leq 1$ and $\|h\|_\infty \leq 1$, we see that

$$\mathbb{P}_{\phi, \psi}\left(n|\hat{p}^{(3)}(a, b, c) - p_{\phi, \psi}^{(3)}(a, b, c)| \geq x\right) \leq 2 \exp\left(-\frac{\gamma x^2}{8n + 8/\gamma + 20x}\right),$$

hence for some constant $c' > 0$

$$\mathbb{P}_{\phi, \psi}\left(|\hat{p}^{(3)}(a, b, c) - p_{\phi, \psi}^{(3)}(a, b, c)| \geq x/\sqrt{n}\right) \leq 2 \exp\left(-c' \min\left(x^2, x^2 n, x\sqrt{n}\right)\right).$$

Using that $\|\hat{p}^{(3)} - p^{(3)}\| \leq K^3 \max_{a,b,c} |\hat{p}^{(3)}(a, b, c) - p^{(3)}(a, b, c)|$ and a union bound, we deduce for some $C = C(K, L)$ and for $x \leq \sqrt{n}$ that

$$\mathbb{P}_{\phi, \psi}\left(\|\hat{p}^{(3)} - p^{(3)}\| \geq K^3 x/\sqrt{n}\right) \leq 2K^3 \exp(-Cx^2).$$

For $x \geq 1$ we may absorb the factor $2K^3$ into the exponential by changing the constant C , and by replacing x with $C'x$ we can remove this constant, yielding the result in the case where $C'x \leq \sqrt{n}$. In the other case, since $\|\hat{p}^{(3)} - p^{(3)}\|$ is bounded (by $K^{3/2}$), by increasing the constant C' if necessary we have $C'x/\sqrt{n} \geq K^{3/2}$ so that the probability in question is equal to $0 \leq e^{-x^2}$. \square

To prove Theorem 2, observe that by Lemma 1 there exist events \mathcal{A}_n of probability at least e^{-x^2} on which

$$\|\hat{p}_n^{(3)} - p_{\phi, \psi}^{(3)}\| \leq Cx/\sqrt{n}.$$

The true parameter (ϕ, ψ) lies in Φ_L so that any estimators constructed in Theorem 2 satisfy

$$\|p_{\hat{\phi}, \hat{\psi}}^{(3)} - \hat{p}_n^{(3)}\| \leq 2\|p_{\phi, \psi}^{(3)} - \hat{p}_n^{(3)}\|,$$

and hence on the event \mathcal{A}_n further satisfy

$$\|p_{\hat{\phi}, \hat{\psi}}^{(3)} - p_{\phi, \psi}^{(3)}\| \leq \|p_{\hat{\phi}, \hat{\psi}}^{(3)} - \hat{p}_n^{(3)}\| + \|p_{\phi, \psi}^{(3)} - \hat{p}_n^{(3)}\| \leq 3\|p_{\phi, \psi}^{(3)} - \hat{p}_n^{(3)}\| \leq 3Cx/\sqrt{n},$$

By Proposition 1 we deduce for a constant C' that $\rho(\hat{\phi}, \hat{\psi}; \phi, \psi) \leq C'x/\sqrt{n}$ on \mathcal{A}_n . For estimating ψ , observe that $\|\hat{\psi}_1 - \psi_1\| \leq \rho(\hat{\phi}, \hat{\psi}; \phi, \psi)$ and $|r(\phi)| \min(\|\hat{\psi}_2 - \psi_2\|, \|\hat{\psi}_2 + \psi_2\|) \leq \rho(\hat{\phi}, \hat{\psi}; \phi, \psi)$. The upper bound for estimating ψ_1 is immediate and, recalling from equation (5) that $|r(\phi)| \geq \delta\epsilon\zeta^2/4$, we also deduce the bound for ψ_2 .

For the bounds on ϕ , observe firstly that it suffices to prove the upper bounds on the absolute risk since, taking ϕ_2 as an example, for $(\phi, \psi) \in \Phi_L(\delta, \epsilon, \zeta)$ we have

$$\begin{aligned} \mathbb{P}_{\phi, \psi} \left(\left| \hat{\phi}_2 / \phi_2 - 1 \right|^2 \geq \frac{C}{n\delta^2\epsilon^4\zeta^4} \right) &= \mathbb{P}_{\phi, \psi} \left(\left| \hat{\phi}_2 - \phi_2 \right|^2 \geq \frac{C\phi_2^2}{n\delta^2\epsilon^4\zeta^4} \right) \\ &\leq \mathbb{P}_{\phi, \psi} \left(\left| \hat{\phi}_2 - \phi_2 \right| \geq \frac{C\epsilon^2}{n\delta^2\epsilon^4\zeta^4} \right). \end{aligned} \quad (26)$$

(See also after [equation \(27\)](#) for a similar argument with ϕ_1 .) Define

$$\omega_1(\phi, \psi; \eta) := \sup \left\{ \left| \phi_1 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\phi}_1 \right| : \rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq \eta \right\},$$

and

$$\omega_j(\phi, \psi; \eta) := \sup \left\{ \left| \phi_j - \tilde{\phi}_j \right| : \rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq \eta \right\}, \quad j = 2, 3.$$

We have the following.

Proposition 2. *Let $\eta \in [0, 1]$. There exist constants c, C for which the following hold.*

$$\begin{aligned} \eta < c(1 - \phi_1^2)\phi_2^2\phi_3^3 &\implies \omega_1(\phi, \psi; \eta) \leq \frac{C\eta}{\phi_2^2\phi_3^3}, \\ \eta < c(1 - \phi_1^2)|\phi_2|\phi_3^2 &\implies \omega_2(\phi, \psi; \eta) \leq \frac{C\eta}{(1 - \phi_1^2)|\phi_2|\phi_3^2}, \\ \eta < c(1 - \phi_1^2)\phi_2^2\phi_3^3 &\implies \omega_3(\phi, \psi; \eta) \leq \frac{C\eta}{(1 - \phi_1^2)\phi_2^2\phi_3^3}. \end{aligned}$$

The conditions of [Theorem 2](#) ensure that on the event \mathcal{A}_n we may apply [Proposition 2](#) with $\eta = C'x/\sqrt{n}$. We deduce the upper bounds for estimating the components of ϕ immediately upon replacing ϕ_1, ϕ_2 and ϕ_3 on the right sides in [Proposition 2](#) by their lower bounds [for ϕ_1 we note that $\min(|\hat{\phi}_1 - \phi_1|, |\hat{\phi}_1 + \phi_1|) \leq |\phi_1 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\phi}_1|$].

Proof of [Proposition 2](#). Recall that $m(\phi) = (r(\phi), \phi_2 r(\phi), \phi_1 \phi_2 \phi_3 r(\phi))$ with $r(\phi) = \frac{1}{4}(1 - \phi_1^2)\phi_2\phi_3^2$. If $r(\phi) = 0$ then for each item no $\eta \in [0, 1]$ satisfies the conditions and so there is nothing to prove. Otherwise, note that m is invertible when restricted to $\{\phi : r(\phi) \neq 0\} \supset \Phi(\delta, \epsilon, \zeta)$ and its inverse is given by $\phi(m)$ defined by

$$\begin{aligned} \phi_1(m) &= m_3 / (4m_1^2 m_2 + m_3^2)^{1/2} \\ \phi_2(m) &= m_2 / m_1, \\ \phi_3(m) &= (4m_1^2 m_2 + m_3^2)^{1/2} / m_2. \end{aligned}$$

For arbitrary $(\phi, \psi; \tilde{\phi}, \tilde{\psi})$ satisfying $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq \eta$, we define

$$\Delta_1 := m_1(\tilde{\phi}) - m_1(\phi), \quad \Delta_2 := m_2(\tilde{\phi}) - m_2(\phi), \quad \Delta_3 := \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot m_3(\tilde{\phi}) - m_3(\phi).$$

Define also

$$\begin{aligned} g(\phi) &:= 4m_1(\phi)^2 m_2(\phi) + m_3(\phi)^2 \\ &= \{m_2(\phi)\phi_3\}^2 \\ &= \left\{ \frac{1}{4}(1 - \phi_1^2)\phi_2^2\phi_3^3 \right\}^2, \end{aligned}$$

and, for $\Delta = (\Delta_1, \Delta_2, \Delta_3)$,

$$\begin{aligned} h_\phi(\Delta) &:= g(\tilde{\phi}) - g(\phi) \\ &= 4(m_1(\phi) + \Delta_1)^2 (m_2(\phi) + \Delta_2) + (m_3(\phi) + \Delta_3)^2 - \{4m_1(\phi)^2 m_2(\phi) + m_3(\phi)^2\}. \end{aligned}$$

Observe that

$$\begin{aligned} h_\phi(\Delta) &= 8m_1(\phi)m_2(\phi)\Delta_1 + 8m_1(\phi)\Delta_1\Delta_2 + 4m_2(\phi)\Delta_1^2 \\ &\quad + 4\Delta_1^2\Delta_2 + 4m_1(\phi)^2\Delta_2 + 2m_3(\phi)\Delta_3 + \Delta_3^2. \end{aligned}$$

Bounding ω_1 We decompose,

$$\begin{aligned}
\phi_1 - \operatorname{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\phi}_1 &= \frac{m_3(\phi)}{\sqrt{4m_1(\phi)^2 m_2(\phi) + m_3(\phi)^2}} - \frac{\operatorname{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot m_3(\tilde{\phi})}{\sqrt{4m_1(\tilde{\phi})^2 m_2(\tilde{\phi}) + m_3(\tilde{\phi})^2}} \\
&= \frac{m_3(\phi)}{\sqrt{g(\phi)}} - \frac{m_3(\phi) + \Delta_3}{\sqrt{g(\phi) + h_\phi(\Delta)}} \\
&= m_3(\phi) \left\{ \frac{1}{\sqrt{g(\phi)}} - \frac{1}{\sqrt{g(\phi) + h_\phi(\Delta)}} \right\} - \frac{\Delta_3}{\sqrt{g(\phi) + h_\phi(\Delta)}} \\
&= \frac{m_3(\phi)}{\sqrt{g(\phi)(g(\phi) + h_\phi(\Delta))}} \left\{ \sqrt{g(\phi) + h_\phi(\Delta)} - \sqrt{g(\phi)} \right\} - \frac{\Delta_3}{\sqrt{g(\phi) + h_\phi(\Delta)}} \\
&= \frac{m_3(\phi)}{\sqrt{g(\phi)(g(\phi) + h_\phi(\Delta))}} \frac{h_\phi(\Delta)}{\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)}} - \frac{\Delta_3}{\sqrt{g(\phi) + h_\phi(\Delta)}}.
\end{aligned}$$

Now we observe that $m_3(\phi)/\sqrt{g(\phi)}$ is equal to ϕ_1 , so indeed

$$\phi_1 - \operatorname{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\phi}_1 = \frac{\phi_1 h_\phi(\Delta) - \Delta_3(\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)})}{\sqrt{g(\phi) + h_\phi(\Delta)}(\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)})}.$$

Call the numerator of this last fraction N and call its denominator D . Writing $h_\phi(\Delta)$ as $h_\phi(\Delta) = \xi_\phi(\Delta) + \gamma_\phi(\Delta)$, where $\gamma_\phi(\Delta) := 2m_3(\phi)\Delta_3 + \Delta_3^2$, we see that

$$N = \phi_1 \xi_\phi(\Delta) + \phi_1 \gamma_\phi(\Delta) - \Delta_3 \{ (g(\phi) + h_\phi(\Delta))^{1/2} + g(\phi)^{1/2} \}.$$

In order to obtain the optimal upper bound, we need to do a fine analysis of this expression. To this end, we calculate

$$\begin{aligned}
A &:= \phi_1 \gamma_\phi(\Delta) - \Delta_3 \{ (g + h)^{1/2} + g^{1/2} \} \\
&= 2\Delta_3 \{ \phi_1 m_3(\phi) - g^{1/2} \} + \phi_1 \Delta_3^2 - \Delta_3 \{ (g + h)^{1/2} - g^{1/2} \} \\
&= -2\Delta_3 (1 - \phi_1^2) g^{1/2} + \phi_1 \Delta_3^2 - \frac{\Delta_3 h}{(g + h)^{1/2} + g^{1/2}} \\
&= -2\Delta_3 (1 - \phi_1^2) g^{1/2} - \Delta_3 \frac{\gamma_\phi(\Delta) - \phi_1 \Delta_3 ((g + h)^{1/2} + g^{1/2})}{(g + h)^{1/2} + g^{1/2}} - \frac{\Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}},
\end{aligned}$$

where the last line follows because $\phi_1 m_3(\phi) = \phi_1^2 g(\phi)^{1/2}$. We now focus on the middle term of this last display, which we will express as a function of A .

$$\begin{aligned}
B &:= \gamma_\phi(\Delta) - \phi_1 \Delta_3 ((g + h)^{1/2} + g^{1/2}) \\
&= 2\Delta_3 (m_3(\phi) - \phi_1 g^{1/2}) + \Delta_3^2 - \phi_1 \Delta_3 \{ (g + h)^{1/2} - g^{1/2} \} \\
&= \Delta_3^2 - \frac{\phi_1 \Delta_3 h}{(g + h)^{1/2} + g^{1/2}} \\
&= \Delta_3^2 - \frac{\phi_1 \Delta_3 \gamma_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}} - \frac{\phi_1 \Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}} \\
&= \Delta_3 \frac{\Delta_3 \{ (g + h)^{1/2} + g^{1/2} \} - \phi_1 \gamma_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}} - \frac{\phi_1 \Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}} \\
&= -\frac{\Delta_3 A}{(g + h)^{1/2} + g^{1/2}} - \frac{\phi_1 \Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}}.
\end{aligned}$$

Thus,

$$\begin{aligned}
A &= -2\Delta_3 (1 - \phi_1^2) g^{1/2} - \frac{\Delta_3 B}{(g + h)^{1/2} + g^{1/2}} - \frac{\Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}} \\
&= -2\Delta_3 (1 - \phi_1^2) g^{1/2} + \frac{\Delta_3^2 A}{\{ (g + h)^{1/2} + g^{1/2} \}^2} + \frac{\phi_1 \Delta_3^2 \xi_\phi(\Delta)}{\{ (g + h)^{1/2} + g^{1/2} \}^2} - \frac{\Delta_3 \xi_\phi(\Delta)}{(g + h)^{1/2} + g^{1/2}},
\end{aligned}$$

from which we deduce that

$$N = \phi_1 \xi_\phi(\Delta) + \frac{-2\Delta_3(1 - \phi_1^2)g^{1/2} + \frac{\phi_1 \Delta_3^2 \xi_\phi(\Delta)}{\{(g+h)^{1/2} + g^{1/2}\}^2} - \frac{\Delta_3 \xi_\phi(\Delta)}{(g+h)^{1/2} + g^{1/2}}}{1 - \Delta_3^2 / \{(g+h)^{1/2} + g^{1/2}\}^2}.$$

Since $m_2(\phi) \geq 0$, we see that $\xi_\phi(\Delta)$ has maximal amplitude when $\Delta_1 = \text{sgn}(m_1(\phi))\eta$ and when $\Delta_2 = \eta$, in which case we have

$$\begin{aligned} |\xi_\phi(\Delta)| &= 8|m_1(\phi)|m_2(\phi)\eta + 8|m_1(\phi)|\eta^2 + 4m_2(\phi)\eta^2 + 4\eta^3 + 4m_1(\phi)^2\eta \\ &\leq 12m_1(\phi)^2\eta + 12|m_1(\phi)|\eta^2 + 4\eta^3, \end{aligned}$$

where the last line follows since $m_2(\phi) \leq |m_1(\phi)|$. Now we observe that under the condition of the lemma, we have $\eta \lesssim |m_1(\phi)|$, and so we can find a constant $C > 0$ such that

$$|\xi_\phi(\Delta)| \leq C m_1(\phi)^2 \eta.$$

Also, we have that $|\gamma_\phi(\Delta)| \leq 2|m_3(\phi)|\eta + \eta^2$, and so

$$|h_\phi(\Delta)| \leq C m_1(\phi)^2 \eta + 2|m_3(\phi)|\eta + \eta^2,$$

Noting that $\phi_3 \leq \sqrt{K}$, for $c_0 = c_0(K)$ sufficiently small in the assumption of the proposition we have $|h_\phi(\Delta)| \leq g(\phi)/2$. Consequently, noting also that $|\Delta_3| \leq \eta$ and $\eta \leq 4c_0 g^{1/2}$, we find that

$$\begin{aligned} |N| &\lesssim |\phi_1| m_1(\phi)^2 \eta + \eta(1 - \phi_1^2)g(\phi)^{1/2} \\ &\lesssim \eta(1 - \phi_1^2)^2 \phi_2^2 \phi_3^3, \end{aligned}$$

and

$$|D| \gtrsim g(\phi) \gtrsim (1 - \phi_1^2)^2 \phi_2^4 \phi_3^6.$$

Hence we have

$$|\phi_1 - \text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) \cdot \tilde{\phi}_1| \lesssim \frac{\eta}{\phi_2^2 \phi_3^3}.$$

Bounding ω_2 We rewrite,

$$\begin{aligned} \phi_2 - \tilde{\phi}_2 &= \frac{m_2(\phi)}{m_1(\phi)} - \frac{m_2(\phi) + \Delta_2}{m_1(\phi) + \Delta_1} \\ &= \frac{m_2(\phi)(m_1(\phi) + \Delta_1) - (m_2(\phi) + \Delta_2)m_1(\phi)}{m_1(\phi)(m_1(\phi) + \Delta_1)}. \end{aligned}$$

Hence,

$$\phi_2 - \tilde{\phi}_2 = \frac{\Delta_1 m_2(\phi) - \Delta_2 m_1(\phi)}{m_1(\phi)(m_1(\phi) + \Delta_1)}.$$

Under the assumptions of the theorem, we have that $\eta \leq m_1(\phi)/2$, and thus

$$\begin{aligned} |\phi_2 - \tilde{\phi}_2| &\leq \frac{2\eta(m_2(\phi) + |m_1(\phi)|)}{m_1(\phi)^2} \\ &\leq \frac{4\eta}{|m_1(\phi)|} \\ &= \frac{16\eta}{(1 - \phi_1^2)|\phi_2| \phi_3^2}. \end{aligned}$$

Bounding ω_3 We rewrite,

$$\begin{aligned} \phi_3 - \tilde{\phi}_3 &= \frac{\sqrt{g(\phi)}}{m_2(\phi)} - \frac{\sqrt{g(\phi) + h_\phi(\Delta)}}{m_2(\phi) + \Delta_2} \\ &= \frac{m_2(\phi)(\sqrt{g(\phi)} - \sqrt{g(\phi) + h_\phi(\Delta)})}{m_2(\phi)(m_2(\phi) + \Delta_2)} + \frac{\Delta_2 \sqrt{g(\phi)}}{m_2(\phi)(m_2(\phi) + \Delta_2)} \\ &= \frac{-h_\phi(\Delta)}{(m_2(\phi) + \Delta_2)(\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)})} + \frac{\Delta_2 \phi_3}{m_2(\phi) + \Delta_2} \\ &= \frac{-h_\phi(\Delta) + \Delta_2 \phi_3(\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)})}{(m_2(\phi) + \Delta_2)(\sqrt{g(\phi) + h_\phi(\Delta)} + \sqrt{g(\phi)})} \end{aligned}$$

Let us call the numerator of the rhs of the last display N , and the denominator D . We further decompose $h_\phi(\Delta)$ as $h_\phi(\Delta) = \xi_\phi(\Delta) + \gamma_\phi(\Delta)$, where $\gamma_\phi(\Delta) := 4m_1(\phi)^2\Delta_2$. We see that

$$\begin{aligned} N &= -\xi_\phi(\phi) - 4m_1(\phi)^2\Delta_2 + \phi_3\Delta_2((g+h)^{1/2} + g^{1/2}) \\ &= -\xi_\phi(\phi) - 4m_1(\phi)^2\Delta_2 + \phi_3\Delta_2\{(g+h)^{1/2} + g^{1/2}\} \\ &= -\xi_\phi(\phi) - 4m_1(\phi)^2\Delta_2 + 2\phi_3\Delta_2g^{1/2} + \phi_3\Delta_2\{(g+h)^{1/2} - g^{1/2}\} \\ &= -\xi_\phi(\phi) + \Delta_2(1 + \phi_1^2)\phi_3g^{1/2} + \frac{\phi_3\Delta_2h}{(g+h)^{1/2} + g^{1/2}}, \end{aligned}$$

where the last line follows because $m_1(\phi)^2 = \frac{1}{4}(1 - \phi_1^2)\phi_3g^{1/2}$. Since $m_2(\phi) \geq 0$, we see that $\xi_\phi(\Delta)$ has maximal amplitude when $\Delta_1 = \text{sgn}(m_1(\phi))\eta$ and when $\Delta_2 = \eta$, in which case we have

$$\begin{aligned} |\xi_\phi(\Delta)| &= 8|m_1(\phi)|m_2(\phi)\eta + 8|m_1(\phi)|\eta^2 + 4m_2(\phi)\eta^2 + 4\eta^3 + 2|m_3(\phi)|\eta + \eta^2 \\ &\lesssim \{|m_1(\phi)|m_2(\phi) + 2|m_3(\phi)|\}\eta + \eta^2 \\ &\lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{(1 - \phi_1^2)|\phi_2|\phi_3, |\phi_1|\}\eta + \eta^2, \end{aligned}$$

where the second line follows because under the assumptions of the proposition we have that $m_2(\phi) \lesssim |m_1(\phi)|$ and $\eta \leq m_2(\phi)/2$ (note that $\phi_3 \leq K^{1/2}$). Since $h_\phi(\Delta) = \xi_\phi(\Delta) + 4m_1(\phi)^2\Delta_2$, we also have

$$|h_\phi(\Delta)| \lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{(1 - \phi_1^2)\phi_3, |\phi_1|\}\eta + \eta^2,$$

Hence,

$$\begin{aligned} |N| &\lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{(1 - \phi_1^2)|\phi_2|\phi_3, |\phi_1|\}\eta + \eta^2 \\ &\quad + \eta\phi_3g^{1/2} + \frac{\eta^2\phi_3(1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{(1 - \phi_1^2)\phi_3, |\phi_1|\} + \eta^3\phi_3}{g^{1/2}} \\ &\lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{(1 - \phi_1^2)|\phi_2|\phi_3, |\phi_1|\}\eta + \eta^2 \\ &\quad + \eta\phi_3g^{1/2} + \eta^2\phi_3 \max\{(1 - \phi_1^2)\phi_3, |\phi_1|\} + \frac{\eta^3}{(1 - \phi_1^2)\phi_2^2\phi_3^2} \end{aligned}$$

But by assumption $\eta \lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3$, and $4g^{1/2} = (1 - \phi_1^2)\phi_2^2\phi_3^2$, thus

$$|N| \lesssim (1 - \phi_1^2)\phi_2^2\phi_3^3 \max\{\phi_3, |\phi_1|\}\eta + \eta^2.$$

Note that $\max(\phi_3, |\phi_1|) \leq \sqrt{K}$. Moreover, under the assumptions of the proposition and using that $\phi_3 \leq \sqrt{K}$, it is the case that $|\Delta_2| \leq \eta \lesssim m_2(\phi)$. Therefore $|D| \gtrsim m_2(\phi)\sqrt{g(\phi)}$, and

$$|\phi_3 - \tilde{\phi}_3| \lesssim \frac{\eta}{(1 - \phi_1^2)\phi_2^2\phi_3^2} + \frac{\eta^2}{(1 - \phi_1^2)^2\phi_2^4\phi_3^5}.$$

Finally, since we have assumed that $\eta < \frac{(1 - \phi_1^2)\phi_2^2\phi_3^2}{8}$, we see that the second term is at most a constant times the first, so that it can be absorbed by increasing the constant C . \square

A.3 Proof of Theorem 3

We give a standard two-point testing lower bound, summarising ideas that can be found for example in Chapter 2 of [36].

Lemma 3. *Given data $X^{(n)} \sim p_u^{(n)}$ for parameter $u \in \mathcal{U}$, the following lower bounds hold for estimating u .*

Suppose $\mathcal{U} \subseteq \mathbb{R}$ and for some $r \leq 1/2$ assume that there exist parameters u_0, u_1 satisfying

i. $|u_1/u_0 - 1| \geq 4r$,

ii. $K(p_{u_1}^{(n)}; p_{u_0}^{(n)}) \leq 1/100$.

where we recall K denotes the Kullback–Leibler divergence. Then

$$\inf_{\hat{u}} \sup_{u \in \mathcal{U}} \mathbb{P}_u(|\hat{u}/u - 1| \geq r) \geq 1/4,$$

where the infimum is over all estimators \hat{u} based on the data $X^{(n)}$.

If instead (\mathcal{U}, d) is a pseudo-metric space and for some $r \geq 0$ there exist parameters u_0, u_1 satisfying

i. $d(u_0, u_1) \geq 2r$

ii. $K(p_{u_1}^{(n)}, p_{u_0}^{(n)}) \leq 1/100$,

then

$$\inf_{\hat{u}} \sup_{u \in \mathcal{U}} \mathbb{P}_u(d(\hat{u}, u) \geq r) \geq 1/4.$$

Proof. Given an estimator \hat{u} we may construct a test T of $u = u_0$ vs $u = u_1$,

$$T = \mathbb{1}\left\{\left|\frac{\hat{u}}{u_0} - 1\right| > \left|\frac{\hat{u}}{u_1} - 1\right|\right\}.$$

Observe that

$$\begin{aligned} \left|\frac{\hat{u}}{u_0} - 1\right| &= \left|\frac{u_1}{u_0} - 1 + \frac{\hat{u} - u_1}{u_1} \frac{u_1}{u_0}\right| \\ &\geq 4r - \left|\frac{\hat{u}}{u_1} - 1\right|(1 + 4r). \end{aligned}$$

Then

$$\begin{aligned} \mathbb{P}_{u_1}(T = 0) &= \mathbb{P}_{u_1}\left(\left|\frac{\hat{u}}{u_0} - 1\right| \leq \left|\frac{\hat{u}}{u_1} - 1\right|\right) \\ &\leq \mathbb{P}_{u_1}\left(4r - \left|\frac{\hat{u}}{u_1} - 1\right|(1 + 4r) \leq \left|\frac{\hat{u}}{u_1} - 1\right|\right) \\ &\leq \mathbb{P}_{u_1}\left(\left|\frac{\hat{u}}{u_1} - 1\right| \geq r\right), \end{aligned}$$

where for the last line we have used that $4r/(2 + 4r) \geq r$ for $r \leq 1/2$. Also note that on the event $\{T = 1\} \cap \{|\hat{u}/u_0 - 1| < r\}$ we have also $|\hat{u}/u_1 - 1| < r$ and hence

$$\begin{aligned} |u_1/u_0 - 1| &= |\hat{u}/u_0 - 1 - (\hat{u}/u_1 - 1) - (\hat{u}/u_1 - 1)(u_1/u_0 - 1)| \\ &< 2r + r|u_1/u_0 - 1|, \end{aligned}$$

so that $|u_1/u_0 - 1| < 2r/(1 - r)$ on this event. Having assumed $r \leq 1/2$ and $|u_1/u_0 - 1| \geq 4r$ we deduce that $\{T = 1\} \cap \{|\hat{u}/u_0 - 1| < r\} = \emptyset$ so that $\{T = 1\} \subseteq \{|\hat{u}/u_0 - 1| \geq r\}$, and hence we have shown

$$\inf_{\hat{u}} \sup_u \mathbb{P}_u\left(\left|\frac{\hat{u}}{u} - 1\right| \geq r\right) \geq \inf_{\hat{u}} \max_{i=0,1} \mathbb{P}_{u_i}\left(\left|\frac{\hat{u}}{u_i} - 1\right| \geq r\right) \geq \inf_T \max_{i=0,1} \mathbb{P}_{u_i}(T \neq i),$$

where the latter infimum is over all tests T . In the pseudo-metric case a reduction considering the test $T = \mathbb{1}\{d(\hat{u}, u_0) > d(\hat{u}, u_1)\}$ and directly using the triangle inequality likewise yields

$$\inf_{\hat{u}} \sup_u \mathbb{P}_u(d(\hat{u}, u) \geq r) \geq \inf_T \max_{i=0,1} \mathbb{P}_{u_i}(T \neq i).$$

It remains to lower bound the maximum probability of testing error by 1/4. Introducing the event

$$A = \left\{\frac{p_{u_0}^{(n)}}{p_{u_1}^{(n)}} \geq 1/2\right\},$$

we see

$$\mathbb{P}_{u_0}(T \neq 0) \geq \mathbb{E}_{u_1}\left[\frac{p_{u_0}^{(n)}}{p_{u_1}^{(n)}} \mathbb{1}_A T\right] \geq \frac{1}{2}[\mathbb{P}_{u_1}(T = 1) - \mathbb{P}_{u_1}(A^c)]$$

Thus, writing $p_1 = \mathbb{P}_{u_1}(T = 1)$, we see

$$\begin{aligned} \max(\mathbb{P}_{u_0}(T \neq 0), \mathbb{P}_{u_1}(T \neq 1)) &\geq \max\left(\frac{1}{2}(p_1 - \mathbb{P}_{u_1}(A^c)), 1 - p_1\right) \\ &\geq \inf_{p \in [0,1]} \max\left(\frac{1}{2}(p - \mathbb{P}_{u_1}(A^c)), 1 - p\right). \end{aligned}$$

The infimum is attained when $\frac{1}{2}(p - \mathbb{P}_{u_1}(A^c)) = 1 - p$ and takes the value $\frac{1}{3}\mathbb{P}_{u_1}(A)$ so that

$$\inf_T \max_{i=0,1} \mathbb{P}_{u_i}(T \neq i) \geq \frac{1}{3}\mathbb{P}_{u_1}(A).$$

Next observe

$$\begin{aligned} \mathbb{P}_{u_1}(A) &= \mathbb{P}_{u_1}\left[\frac{p_{u_1}^{(n)}}{p_{u_0}^{(n)}} \leq 2\right] = 1 - \mathbb{P}_{u_1}\left[\log\left(\frac{p_{u_1}^{(n)}}{p_{u_0}^{(n)}}\right) > \log 2\right] \geq 1 - \mathbb{P}_{\theta_1}\left[|\log\left(\frac{p_{u_1}^{(n)}}{p_{u_0}^{(n)}}\right)| > \log 2\right] \\ &\geq 1 - (\log 2)^{-1}\mathbb{E}_{u_1}\left|\log\left(\frac{p_{u_1}^{(n)}}{p_{u_0}^{(n)}}\right)\right|, \end{aligned}$$

where we have used Markov's inequality to attain the final expression. By the second Pinsker inequality (e.g. Proposition 6.1.7b in [19]), using the upper bound on the Kullback–Leibler divergence we can continue the chain of inequalities to see

$$\mathbb{P}_{u_1}(A) \geq 1 - (\log 2)^{-1}\left[K(p_{u_1}^{(n)}, p_{u_0}^{(n)}) + \sqrt{2K(p_{u_1}^{(n)}, p_{u_0}^{(n)})}\right] \geq 1 - (\log 2)^{-1}(\mu + \sqrt{2\mu}).$$

For any $c < 1/3$, we may choose $\mu = \mu(c)$ small enough that the testing error satisfies

$$\inf_T \max_{i=0,1} \mathbb{P}_{u_i}(T \neq i) \geq \frac{1}{3}\left(1 - \frac{\mu + \sqrt{2\mu}}{\log 2}\right) > c,$$

and in particular a numerical calculation shows that $\mu = 1 + \frac{1}{4}\log 2 - \sqrt{1 + \frac{1}{2}\log 2} > 1/100$ works for $c = 1/4$. \square

In view of [Lemma 2](#) and [Proposition 1](#), for any $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \Phi$ corresponding to strictly positive emission densities, we have

$$K(p_{\phi, \psi}^{(n)}, p_{\tilde{\phi}, \tilde{\psi}}^{(n)}) \leq Cn\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi})^2,$$

where $C > 0$ is a constant depending only on K and a lower bound for the emission densities. To prove [Item \(1\)](#) we apply [Lemma 3](#) to $u = 1 - \phi_1^2$ to see that there exists a constant c_1 for which the relative risk bound is proved if we can exhibit, for a constant c_2 which we may choose arbitrarily, parameters $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \Phi_L(\delta, \epsilon, \zeta)$ satisfying

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq c_1/\sqrt{n}, \quad \text{and} \quad \left|\frac{1 - \tilde{\phi}_1^2}{1 - \phi_1^2} - 1\right| \geq c_2/\sqrt{n\delta^2\epsilon^4\zeta^6}. \quad (27)$$

To prove the absolute risk bound we then note that for any $a \geq 0$, since $|\phi_1| \leq 1$ and $1 - \phi_1^2 \geq \delta$, so that we may assume the same of $\hat{\phi}_1$, we have

$$\begin{aligned} \mathbb{P}_{\phi, \psi}(\min(|\hat{\phi}_1 - \phi_1|, |\hat{\phi}_1 + \phi_1|) \geq a) &\geq \mathbb{P}_{\phi, \psi}(|(1 - \hat{\phi}_1^2) - (1 - \phi_1^2)| \geq 2a) \\ &\geq \mathbb{P}_{\phi, \psi}(|(1 - \hat{\phi}_1^2)/(1 - \phi_1^2) - 1| \geq 2a/\delta). \end{aligned}$$

(See also [equation \(26\)](#) for a similar calculation with ϕ_2 .)

Similar conditions to [\(27\)](#) suffice for proving the other parts of [Theorem 3](#) and we proceed now to verifying the existence of suitable parameters (ϕ, ψ) and $(\tilde{\phi}, \tilde{\psi})$, with the help of the following lemma.

Lemma 4. *For a given ϕ , assume conditions [\(6\)](#) and [\(7\)](#) and assume that $\phi_3 \leq \sqrt{2\lfloor K/2 \rfloor}/(2K)$. Then there exists ψ such that (ϕ, ψ) lies in Φ_L and the corresponding emission densities f_0, f_1 are bounded below by some constant $c = c(K) > 0$.*

In particular, for $|\phi_1| \leq 1 - 3\delta$, $\epsilon \leq \phi_2 \leq \min(1/3, 1 - L)$, $\zeta \leq \phi_3 \leq 2\zeta$, then such a ψ exists under the compatibility condition [\(3\)](#).

Proof. Take

$$\psi_1(k) = 1/K, \quad \psi_2(k) = (2\lfloor K/2 \rfloor)^{-1/2}(\mathbb{1}\{k \text{ odd}, k < K\} - \mathbb{1}\{k \text{ even}\}), \quad k \leq K.$$

Under the assumed condition on ϕ_3 and recalling that $|\phi_1| \leq 1$ by assumption, we observe from the expressions for f_0, f_1 given in [Remark 4](#) that these are lower bounded by $1/(2K)$. In the particular case, one simply notes that all the conditions hold for such ϕ . \square

Proof of Items (1) and (3) We prove the lower bounds for estimating ϕ_1 and ϕ_3 together. For some small constant $c > 0$, set $R = c\epsilon^{-2}\zeta^{-3}n^{-1/2}$ and, writing $S = (2 - 6\delta - R)R/(6\delta - 9\delta^2)$, set

$$\begin{aligned}\phi &= (1 - 3\delta, \epsilon, \zeta\sqrt{1 + S}), \\ \tilde{\phi} &= (1 - 3\delta - R, \epsilon, \zeta).\end{aligned}$$

Recalling the definition $r(\phi) = (1 - \phi_1^2)\phi_2\phi_3^2/4$, the choice of ϕ_3 ensures that $r(\phi) = r(\tilde{\phi})$, and we note that under the assumptions of the theorem we have $R \leq \delta \leq 1/6$ so that $S \leq R/\delta \leq 1$ and $\zeta \leq \phi_3 \leq 2\zeta$. By [Lemma 4](#) there exists ψ such that $(\phi, \psi), (\tilde{\phi}, \psi) \in \Phi_L$ and for this ψ we see that

$$\rho(\phi, \psi; \tilde{\phi}, \psi) = |\phi_1\phi_2\phi_3r(\phi) - \tilde{\phi}_1\tilde{\phi}_2\tilde{\phi}_3r(\tilde{\phi})| = \phi_2r(\phi)|\phi_1\phi_3 - \tilde{\phi}_1\tilde{\phi}_3|.$$

Using that $\sqrt{1+t} \leq 1+t$ for $t \geq 0$ we have

$$|\phi_1\phi_3 - \tilde{\phi}_1\tilde{\phi}_3| = (1 - 3\delta)\zeta(\sqrt{1 + S} - 1) + R\zeta \leq (S + R)\zeta \leq 2R\zeta/\delta,$$

hence since $r(\phi) = (6\delta - 9\delta^2)\epsilon\zeta^2(1 + S)/4 \leq 3\delta\epsilon\zeta^2$, we obtain

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq 6\epsilon^2\zeta^3R \leq 6cn^{-1/2}.$$

Recalling that $R \leq \delta \leq 1/6$ and that $r(\phi) = r(\tilde{\phi})$ one calculates

$$\frac{1 - \tilde{\phi}_1^2}{1 - \phi_1^2} - 1 = S \geq R/(12\delta).$$

For c small enough we see that the conditions in [equation \(27\)](#) are satisfied, yielding the claimed bound for estimating ϕ_1 .

To prove the lower bound for estimating ϕ_3 it suffices to lower bound $|\phi_3/\tilde{\phi}_3 - 1|$. Here we use the bound $\sqrt{1+x} - 1 \geq x/(2\sqrt{1+x}) \geq x/(2\sqrt{2})$ for $0 \leq x \leq 1$ to see for a constant $c' > 0$ that

$$|\phi_3/\tilde{\phi}_3 - 1| \geq c'R/\delta.$$

The bound for ϕ_3 follows from applying [Lemma 3](#).

Proof of Item (2) For a constant $c > 0$, define $R = c\delta^{-1}\epsilon^{-1}\zeta^{-2}n^{-1/2}$, define $\phi, \tilde{\phi}$ by

$$\begin{aligned}\phi &= (1 - 3\delta, \epsilon, \zeta(1 + R/\epsilon)^{1/2}) \\ \tilde{\phi} &= (1 - 3\delta, \epsilon + R, \zeta),\end{aligned}$$

and observe that by construction $r(\phi) = r(\tilde{\phi})$. Noting that $\phi_2 \leq 2\epsilon \leq 1 - L$ and $\phi_3 \leq 2\zeta$ because the assumptions of [Theorem 3](#) ensure that $R \leq \epsilon \leq 1/3$, we deduce using [Lemma 4](#) that there exists some $\psi = \tilde{\psi}$ such that $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \Phi_L(\delta, \epsilon, \zeta)$.

Next observe, using that $(1+x)^{1/2} \leq 1+x$,

$$\phi_1|\phi_2\phi_3 - \tilde{\phi}_2\tilde{\phi}_3| \leq |\phi_2||\phi_3 - \tilde{\phi}_3| + |\tilde{\phi}_3||\phi_2 - \tilde{\phi}_2| = \epsilon\zeta(\sqrt{1 + R/\epsilon} - 1) + \zeta R \leq 2\zeta R \leq R,$$

the last inequality following from the fact that under the compatibility condition (3) we have $\zeta \leq 1/2$. We deduce

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) = r(\phi) \max(|\phi_2 - \tilde{\phi}_2|, |\phi_1\phi_2\phi_3 - \tilde{\phi}_1\tilde{\phi}_2\tilde{\phi}_3|) = Rr(\phi).$$

Again using that $\phi_3 \leq 2\zeta$ and noting also that $(1 - \phi_1^2) = 6\delta - 9\delta^2 \leq 6\delta$, we see that for some $C' > 0$ we have

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq C'\delta\epsilon\zeta^2R \leq cC'n^{-1/2}.$$

As with [Items \(1\) and \(3\)](#), for c small enough in the definition of R we may apply [Lemma 3](#) to deduce the claimed lower bound since $|\tilde{\phi}_2/\phi_2 - 1| = R/\epsilon$.

Proof of Item (4) Set $\phi = \tilde{\phi} = (0, \epsilon, \zeta)$, set, as in [Lemma 4](#),

$$\psi_1(k) = 1/K, \quad \psi_2(k) = (2\lfloor K/2 \rfloor)^{-1/2}(\mathbb{1}\{k \text{ odd}, k < K\} - \mathbb{1}\{k \text{ even}\}), \quad k \leq K,$$

and define $\tilde{\psi}_1 = \psi_1 + cn^{-1/2}\psi_2$. Note that under the compatibility condition (3) we have $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \Phi_L$ for n larger than some $C = C(K, c)$, or for all $n \geq 1$ if c is small enough. Then

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) = cn^{-1/2}$$

and we apply [Lemma 3](#) to deduce the result.

Proof of Item (5) Set $\phi = \tilde{\phi} = (1 - 3\delta, \epsilon, \zeta)$, choose $\psi_1 = \tilde{\psi}_1$ to be the uniform density on $\{1, \dots, K\}$. As with the previous parts, an application of [Lemma 3](#) will yield the theorem if we can exhibit $\psi_2, \tilde{\psi}_2$ such that the induced emission densities are bounded below by some $c' = c'(K) > 0$, $\|\psi_2 - \tilde{\psi}_2\| = R := c(n\delta^2\epsilon^2\zeta^4)^{-1/2}$ for some $c > 0$, $\text{sgn}(\langle \psi_2, \tilde{\psi}_2 \rangle) = +1$, and $\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) \leq c_1 n^{-1/2}$ for a small constant c_1 . Such a choice is possible, for a small enough constant c , under the compatibility condition (3) and for $n\delta^2\epsilon^2\zeta^4 \geq 1$; for example, define ψ_2 as in [Lemma 4](#) by

$$\psi_2(k) = (2\lfloor K/2 \rfloor)^{-1/2}(\mathbb{1}\{k \text{ odd}, k < K\} - \mathbb{1}\{k \text{ even}\}), \quad k \leq K,$$

and, for h defined by $h(1) = 2^{-1/2}$, $h(3) = -2^{-1/2}$ and $h(k) = 0$ for all other k , define

$$\tilde{\psi}_2 = (\psi_2 + \alpha h)/(1 + \alpha), \quad \alpha = R/(2 - R).$$

This satisfies $\|\tilde{\psi}_2 - \psi_2\| = R$, $\|\tilde{\psi}_2\| = 1$, $\langle \tilde{\psi}_2, 1 \rangle = 0$ and $\langle \tilde{\psi}_2, \psi_2 \rangle \geq 0$. For $k \notin \{1, 3\}$ the condition (8) of [Remark 5](#) holds with $1/(2K)$ in place of 0 on the right, and for $k \in \{1, 3\}$ a direct calculation shows that the condition with $1/(4K)$ in on the right if R is upper bounded by some $c' = c'(K)$, which is the case for $c = c(K)$ sufficiently small. Then

$$\rho(\phi, \psi; \tilde{\phi}, \tilde{\psi}) = |r(\phi)|\|\psi_2 - \tilde{\psi}_2\| \leq \delta\epsilon\zeta^2 R \leq cn^{-1/2}.$$

A.4 Proofs for [Section 4](#)

Proof of [Theorem 1](#). We begin with the upper bounds. From the inversion formulae in [Remark 4](#) we have

$$\max(\|\hat{f}_0 - f_0\|, \|\hat{f}_1 - f_1\|) \leq \|\hat{\psi}_1 - \psi_1\| + \frac{1}{2}\|\hat{\phi}_1\hat{\phi}_3\hat{\psi}_2 - \phi_1\phi_3\psi_2\| + \frac{1}{2}\|\hat{\phi}_3\hat{\psi}_2 - \phi_3\psi_2\|$$

Recalling that $|\hat{\phi}_1| \leq 1$, that $0 \leq \phi_3 \leq K^{1/2}$ and that $\|\psi_2\| = \|\hat{\psi}_2\| = 1$, we decompose the second term on the right, with an implicit decomposition of the third term included:

$$\begin{aligned} \|\hat{\phi}_1\hat{\phi}_3\hat{\psi}_2 - \phi_1\phi_3\psi_2\| &\leq |\hat{\phi}_1|\|\hat{\phi}_3\hat{\psi}_2 - \phi_3\psi_2\| + |\phi_3|\|\hat{\phi}_1 - \phi_1\| \\ &\leq |\hat{\phi}_3 - \phi_3| + K^{1/2}\|\hat{\psi}_2 - \psi_2\| + \phi_3|\hat{\phi}_1 - \phi_1|. \end{aligned}$$

It follows that for some constant C we have

$$\max(\|\hat{f}_0 - f_0\|, \|\hat{f}_1 - f_1\|) \leq C \max(\|\hat{\psi}_1 - \psi_1\|, \|\hat{\psi}_2 - \psi_2\|, |\hat{\phi}_3 - \phi_3|, \phi_3|\hat{\phi}_1 - \phi_1|).$$

Applying [Proposition 2](#) as in the proof of [Theorem 2](#), one can show that for some $C > 0$

$$\mathbb{P}_{\phi, \psi} \left(\phi_3^2 |\hat{\phi}_1 - \phi_1|^2 \geq \frac{Cx^2}{n\epsilon^4\zeta^4} \right) \leq e^{-x^2}.$$

The upper bounds for estimating f_0 and f_1 then follow from [Theorem 2](#).

Similarly, [Remark 4](#) and the fact that $|\phi_2| \leq 1$ give

$$\max(|\hat{p} - p|, |\hat{q} - q|) \leq \frac{1}{2}(1 + |\hat{\phi}_1|)|\hat{\phi}_2 - \phi_2| + \frac{1}{2}|\hat{\phi}_1 - \phi_1||1 - \phi_2| \leq 2 \max(|\hat{\phi}_1 - \phi_1|, |\hat{\phi}_2 - \phi_2|).$$

The upper bounds then again follow from [Theorem 2](#).

For the lower bounds, writing $\theta(\phi, \psi) = (p, q, f_0, f_1)$ and $\theta(\tilde{\phi}, \tilde{\psi}) = (\tilde{p}, \tilde{q}, \tilde{f}_0, \tilde{f}_1)$, observe by [Lemma 3](#) that it suffices to lower bound $\max(|p - \tilde{p}|, |q - \tilde{q}|)$ and $\max(\|f_0 - \tilde{f}_0\|, \|f_1 - \tilde{f}_1\|)$ corresponding to choices of $(\phi, \psi), (\tilde{\phi}, \tilde{\psi})$ made in the proof of [Theorem 3](#).

From the inversion formulae in [Remark 4](#) we calculate, for any $\phi, \tilde{\phi}$,

$$\begin{aligned} &2 \max(|p - \tilde{p}|, |q - \tilde{q}|) \\ &\geq \max((1 + |\phi_1|)|\phi_2 - \tilde{\phi}_2| - |1 - \tilde{\phi}_2||\phi_1 - \tilde{\phi}_1|, |\phi_1 - \tilde{\phi}_1||1 - \tilde{\phi}_2| - (1 - |\phi_1|)|\phi_2 - \tilde{\phi}_2|). \end{aligned} \tag{28}$$

If $\delta > \epsilon\zeta$ set $\phi = (1 - 3\delta, \epsilon, \zeta(1 + S)^{1/2})$ and $\tilde{\phi} = (1 - 3\delta - R, \epsilon, \zeta)$, where $R = c(n\epsilon^4\zeta^6)^{-1/2}$ for some $c > 0$ and where $S \in [R/(12\delta), R/\delta]$ is, as in the proof of [Theorem 3 Item \(1\)](#), such that

$r(\phi) = r(\tilde{\phi})$. If $\delta \leq \epsilon\zeta$ instead set $\phi = (1 - 3\delta, \epsilon, \zeta(1 + R/\epsilon)^{1/2})$, $\tilde{\phi} = (1 - 3\delta, \epsilon + R, \zeta)$ with $R = c(n\epsilon^2\delta^2\zeta^4)^{-1/2}$. In either case the proof of [Theorem 3](#) demonstrates that for suitable $\psi = \tilde{\psi}$ we have $K(p_{\phi, \psi}^{(n)}, p_{\tilde{\phi}, \tilde{\psi}}^{(n)}) \leq 1/100$ for c small enough hence by [Lemma 3](#)

$$\inf_{\tilde{\theta}} \sup_{\theta \in \Theta} \mathbb{P}_{\theta} \left(\max(|\check{p} - p|, |\check{q} - q|) > c' \max(|p - \tilde{p}|, |q - \tilde{q}|) \right) \geq 1/4.$$

Inserting from [equation \(28\)](#) we conclude the bound in either case.

For (f_0, f_1) , again set $\phi = (1 - 3\delta, \epsilon, \zeta(1 + S)^{1/2})$, $\tilde{\phi} = (1 - 3\delta - R, \epsilon, \zeta)$ where $R = c\epsilon^{-2}\zeta^{-3}n^{-1/2}$, and choose $\psi = \tilde{\psi}$ by [Lemma 4](#). As with p and q we deduce that for some $c' > 0$ we have

$$\inf_{\tilde{\theta}} \sup_{\theta \in \Theta} \mathbb{P}_{\theta} \left(\max(\|\check{f}_0 - f_0\|, \|\check{f}_1 - f_1\|) > c' \max(\|f_0 - \tilde{f}_0\|, \|f_1 - \tilde{f}_1\|) \right) \geq 1/4.$$

Using the inversion formulae in [Remark 4](#) and the fact that $\psi = \tilde{\psi}$ and $\|\psi_2\| = 1$, one calculates

$$2 \max(\|f_0 - \tilde{f}_0\|, \|f_1 - \tilde{f}_1\|) = |\phi_1\phi_3 - \tilde{\phi}_1\tilde{\phi}_3| + |\phi_3 - \tilde{\phi}_3| \geq |\phi_3 - \tilde{\phi}_3|$$

For the current choice of $\phi, \tilde{\phi}$, calculating as in proving [Theorem 3 Item \(3\)](#), we have $|\phi_3 - \tilde{\phi}_3| \geq C\zeta R/\delta$ for some $C > 0$ and we deduce the lower bound. \square

Proof of [Corollary 1](#). It suffices to substitute $\alpha = e^{-x^2}$ into [Theorem 1](#) and solve for error equal to E , while ensuring that $x^2 = \log(1/\alpha)$ is suitably bounded. \square

References

- [1] K. Abraham, I. Castillo, and É. Gassiat. Multiple testing in nonparametric hidden markov models: An empirical Bayes approach. *arXiv:2101.03838*, 2021.
- [2] F. Abramovich, Y. Benjamini, D. L. Donoho, and I. M. Johnstone. Adapting to unknown sparsity by controlling the false discovery rate. *Ann. Statist.*, 34(2):584–653, 2006.
- [3] G. Alexandrovich, H. Holzmam, and A. Leister. Nonparametric identification and maximum likelihood estimation for hidden Markov models. *Biometrika*, 103(2):423–434, 2016.
- [4] E. S. Allman, C. Matias, and J. A. Rhodes. Identifiability of parameters in latent structure models with many observed variables. *Ann. Statist.*, 37(6A):3099–3132, 2009.
- [5] A. Anandumar, R. Ge, D. Hsu, S. M. Kakade, and M. Telgarsky. Tensor decompositions for learning latent variable models. *J. Mach. Learn. Res.*, 15:2773–2832, 2014.
- [6] P. Andreoletti, D. Loukianova, and C. Matias. Hidden Markov model for parameter estimation of a random walk in a Markov environment. *ESAIM Probab. Stat.*, 19:605–625, 2015.
- [7] A. Becker and H. Holzmam. Nonparametric identification in the dynamic stochastic block model. *IEEE Trans. Inform. Theory*, 65(7):4335–4344, 2019.
- [8] I. Castillo and É. Roquain. On spike and slab empirical Bayes multiple testing. *Ann. Statist.*, 48(5):2548–2574, 2020.
- [9] L. Couvreur and C. Couvreur. Wavelet based non-parametric HMMs: theory and methods. In *ICASSP '00 Proceedings*, pages 604–607, 2000.
- [10] Y. De Castro, É. Gassiat, and C. Lacour. Minimax adaptive estimation of nonparametric hidden Markov models. *J. Mach. Learn. Res.*, 17:Paper No. 111, 43, 2016.
- [11] Y. De Castro, É. Gassiat, and S. Le Corff. Consistent estimation of the filtering and marginal smoothing distributions in nonparametric hidden Markov models. *IEEE Trans. Inform. Theory*, 63(8):4758–4777, 2017.
- [12] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 39(1):1–22, 1977.
- [13] L. Devroye, L. Györfi, and G. Lugosi. *A probabilistic theory of pattern recognition*, volume 31. Springer Science & Business Media, 2013.

- [14] N. Doss, Y. Wu, P. Yang, and H. H. Zhou. Optimal estimation of high-dimensional gaussian mixtures. *arXiv:2002.05818*, 2020.
- [15] É. Gassiat, A. Cleynen, and S. Robin. Inference in finite state space non parametric hidden Markov models and applications. *Stat. Comput.*, 26(1-2):61–71, 2016.
- [16] É. Gassiat, S. Le Corff, and L. Lehéricy. Deconvolution with unknown noise distribution is possible for multivariate signals. *arXiv:2006.14226*, 2020.
- [17] É. Gassiat, S. Le Corff, and L. Lehéricy. Identifiability and consistent estimation of nonparametric translation hidden Markov models with general state space. *J. Mach. Learn. Res.*, 21:Paper No. 115, 40, 2020.
- [18] É. Gassiat and R. van Handel. The local geometry of finite mixtures. *Trans. Amer. Math. Soc.*, 366(2):1047–1072, 2014.
- [19] Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*. Cambridge Series in Statistical and Probabilistic Mathematics, [40]. Cambridge University Press, New York, 2016.
- [20] P. Heinrich and J. Kahn. Strong identifiability and optimal minimax rates for finite mixture estimation. *Ann. Statist.*, 46(6A):2844–2870, 2018.
- [21] D. Hsu, S. M. Kakade, and T. Zhang. A spectral algorithm for learning hidden Markov models. *J. Comput. System Sci.*, 78(5):1460–1480, 2012.
- [22] A. Kumar and R. Kannan. Clustering with spectral norm and the k-means algorithm. In *Proceedings of the 2010 IEEE 51st Annual Symposium on Foundations of Computer Science, FOCS '10*, page 299–308, USA, 2010. IEEE Computer Society.
- [23] M. F. Lambert, J. P. Whiting, and A. V. Metcalfe. A non-parametric hidden Markov model for climate state identification. *Hydrology and Earth System Sciences*, 7 (5):652–667, 2003.
- [24] L. Le Cam. *Asymptotic methods in statistical decision theory*. Springer-Verlag, New York, 1986.
- [25] F. Lefèvre. Non-parametric probability estimation for HMM-based automatic speech recognition. *Computer Speech and Language*, 17:113–136, 2003.
- [26] L. Lehéricy. State-by-state minimax adaptive estimation for nonparametric hidden Markov models. *J. Mach. Learn. Res.*, 19:Paper No. 39, 46, 2018.
- [27] L. Lehéricy. Consistent order estimation for nonparametric hidden Markov models. *Bernoulli*, 25(1):464–498, 2019.
- [28] L. Lehéricy. Nonasymptotic control of the MLE for misspecified nonparametric hidden Markov models. *arXiv:1807.03997*, 2018.
- [29] S. Lloyd. Least squares quantization in PCM. *IEEE transactions on information theory*, 28(2):129–137, 1982.
- [30] C. Matias and V. Miele. Statistical clustering of temporal networks through a dynamic stochastic block model. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 79(4):1119–1141, 2017.
- [31] J. Ollion, C. Ollion, É. Gassiat, L. Lehéricy, and S. Le Corff. Joint self-supervised blind denoising and noise estimation. *arXiv:2102.08023*, 2021.
- [32] D. Paulin. Concentration inequalities for Markov chains by Marton couplings and spectral methods. *Electron. J. Probab.*, 20:no. 79, 32, 2015.
- [33] N. Rau, J. Lucke, and A. K. Hartmann. Phase transition for parameter learning of Hidden Markov Models. *arXiv:2003.11680*, 2020.
- [34] M. Sesia, C. Sabatti, and E. J. Candès. Gene hunting with hidden Markov model knockoffs. *Biometrika*, 106(1):1–18, 2019.
- [35] L. Shang and K.P. Chan. Nonparametric discriminant HMM and application to facial expression recognition. In *2009 IEEE Computer Society Conference on Computer Vision and Pattern Recognition Workshops*, pages 2090–2096, 2009.
- [36] A. B. Tsybakov. *Introduction to nonparametric estimation*. Springer-Verlag, New York, 2009.
- [37] S. Volant, C. Bérard, M.-L. Martin-Magniette, and S. Robin. Hidden Markov Models with mixtures as emission distributions. *Statistics and Computing*, pages 1–12, 2013.

- [38] C. Yau, O. Papaspiliopoulos, G. O. Roberts, and C. Holmes. Bayesian non-parametric hidden Markov models with applications in genomics. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 73(1):37–57, 2011.