



HAL
open science

Bifunctional N-Heterocyclic Carbene-Catalyzed Highly Enantioselective Trans-Cyclopentannulation of Enals and Enones via Homoenolate

Zhiwei Jiang, Martial Toffano, Giang Vo-thanh, Chloée Bournaud

► **To cite this version:**

Zhiwei Jiang, Martial Toffano, Giang Vo-thanh, Chloée Bournaud. Bifunctional N-Heterocyclic Carbene-Catalyzed Highly Enantioselective Trans-Cyclopentannulation of Enals and Enones via Homoenolate. *ChemCatChem*, 2021, 13 (2), pp.712-717. 10.1002/cctc.202001513 . hal-03303929

HAL Id: hal-03303929

<https://hal.science/hal-03303929>

Submitted on 28 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bifunctional N-Heterocyclic Carbene-Catalyzed Highly Enantioselective *Trans*-Cyclopentannulation of Enals and Enones via Homo-enolate

Zhiwei Jiang, Martial Toffano, Giang Vo-Thanh,* and Chloée Bournaud*[a]

[a] Z. Jiang, Dr. M. Toffano, Prof. G. Vo-Thanh, Dr. C. Bournaud
Institut de Chimie Moléculaire et des Matériaux d'Orsay, CNRS UMR 8182, Université Paris-Saclay
Rue du doyen Georges Poitou, 91405 Orsay Cedex, France
E-mail: chloe.bournaud@universite-paris-saclay.fr; giang.vo-thanh@universite-paris-saclay.fr

Supporting information for this article is given via a link at the end of the document.

Abstract: An efficient and flexible synthesis of a new class of chiral bifunctional NHC catalyst has been reported. These new imidazolylidene NHCs, bearing a (thio)urea function as a hydrogen bond donor promoted efficiently highly diastereoselective *trans*-cyclopentannulation of enals and enones in good yield (up to 69 % yield) along with excellent enantioselectivity (up to 96 % ee). This methodology could be applied to a large variety of substrates (30 examples).

Introduction

Nowadays, developing more sustainable chemistry seems to be essential. Various strategies have therefore been adopted in order to reduce the environmental impact of chemistry. Chemists were able to develop catalyzed transformations improving selectivity while decreasing reaction time and chemical waste.^[1] Among various catalytic systems reported, organocatalysis, using a small organic molecule under mild reaction conditions, appears as a very attractive area in modern organic synthesis and is now considered the third pillar of catalysis, alongside metal catalysis and biocatalysis. The main advantages of organocatalysis lie in their tolerance towards many functional groups and therefore many types of organocatalyst have been described. For example, *N*-heterocyclic carbenes (NHCs) are a class of organocatalyst that have been studied in depth due to their efficiency in forming C-C and C-X bonds.^[2] They are able to trigger non-conventional reactions thanks to the formation of reactive intermediates, such as Breslow intermediate, homo-enolate intermediate, azolium enolate, acyl azolium, α,β -unsaturated acyl azolium or azolium dienolate, generated by nucleophilic addition of the lone pair of the carbene to an aldehyde or its equivalent.^[3] The first racemic reaction promoted by a thiazolium salt (precursor to NHC) is the benzoin condensation and dated since 1943 with Ukai^[4] while the first umpolung reaction was described in 1832 by Wöhler and Liebig.^[5] Since then, research on NHCs has intensified, and enantioselective reactions catalyzed by NHCs are now extremely efficient, thanks to the discovery of Breslow's mechanism,^[6] the isolation of the first stable carbene by Bertrand^[7] and Arduengo,^[8] and the first example of enantioselective transformation with triazolium salt precursor.^[9] Although there are manifold chiral NHC catalysts such as thiazolylidene, imidazolylidene, imidazolinylidene or triazolylidene, the latter is by far the most used in asymmetric organocatalysis. Moreover, achiral imidazolylidene derived from imidazolium salt promotes effectively racemic umpolung reactions involving homo-enolate intermediate but it should be noted that only a few chiral imidazolylidene organocatalyst have been reported so far delivering poor level of enantioselectivity in most cases.^[10] So, the design and synthesis of chiral imidazolium, precursor to imidazolylidene NHCs, could extend the range of reactions catalyzed by chiral NHCs. Moreover, a few triazolylidene, thiazolylidene and imidazolinylidene (saturated analogue imidazolylidene) NHC organocatalysts bearing an additional hydrogen bond donor have been reported in the literature.^[11,12] For example, combination of an hydroxyl group with triazolylidene is common.^[11] Amide was combined with triazolylidene^[12c] or thiazolylidene.^[12e-f] Furthermore, thiourea was associated with triazolylidene^[12a-b] or imidazolinylidene^[12d] but the results have been disappointing or still need to be enhanced to have a highly efficient chiral NHC catalyst.

To the best of our knowledge, the combination of chiral imidazolium salt precursor to imidazolylidene NHC and a thiourea function has never been reported so far. In a continuation of our research on the design and synthesis of chiral bifunctional organocatalyst^[13] starting from inexpensive and commercially available precursor from the natural chiral pool, we report herein the synthesis of a new family of chiral bifunctional imidazolylidene NHC-thioureas (Figure 1) and their application in organocatalyzed asymmetric *trans*-cyclopentannulation of enals and chalcones.





Figure 1. Design of new chiral bifunctional imidazolylidene-NHC organocatalyst.


Results and Discussion

Imidazolium scaffolds of the NHC catalyst were prepared in one step by mixing commercially available

aminoalcohols, mesityl amines, formaldehyde, glyoxal and acetic acid.^[14] Hydroxy-imidazolium **2a-d** were isolated in moderate yields (Scheme 1). Then amino-imidazolium **3a-d** were obtained in three steps including an activation of the hydroxyl group by a mesylate followed by a nucleophilic substitution with sodium azide and a reduction by hydrogen over Pd/C. Amino-imidazolium **3a-d** were engaged in the next step without further purification. Finally, reaction with various iso(thio)cyanates afforded expected bifunctional NHC pre-catalyst **4** in good yields.




Scheme 1. Synthesis of bifunctional imidazolylidene-NHC organocatalysts.




i) MesNH₂, glyoxal, formaldehyde, AcOH, 60 °C, 10 min, then KPF₆, rt, 1 h ;
 ii) MsCl, Et₃N, DCM, rt, 2 h ; iii) NaN₃, DMF, 80 °C, 48 h ; iv) Pd/C, H₂, MeOH, rt, 2h; v) Ar-N=C=X, CH₃CN, rt, overnight.

transformation. For example, the use of additives (such as Ti(OiPr)₄)^[16] gave the *cis* products in good yields and ee. In 2013, Coquerel and co-workers^[12d] described the use of NHC functionalized hydrogen-bond donors to promote the formation of *trans*-cyclopentene in low yield (18%) and encouraging 78% ee compared to the previously reported enantioselectivity.^[15a] Moreover, saturated carboxylic acid derivatives were efficiently used instead of enals.^[17] It should be noted that an excess of DBU is needed to generate the homoenolate intermediate due to the release of nitrophenol during the reaction. Finally, various Michael acceptors such as oxindole, benzofuranones and tetrahydroquinolinone were also investigated to afford cyclopentene fused heterocycles adducts.^[18] Despite these results, formation of *trans*-cyclopentene adducts in high yields and enantioselectivity starting from enals and chalcones via homoenolate intermediate was still unsolved. In light of this, we envisioned that *trans*-cyclopentannulation could be efficiently and stereoselectively promoted by our chiral bifunctional imidazolylidene-NHC organocatalysts (scheme 2, Eq. 2).

1) Enantioselective NHC-Catalyzed *cis*-cycloannulation of enals and oxoenoates (previous work, Eq. 1)




2) Enantioselective bifunctional NHC-Catalyzed *trans*-cycloannulation of enals and enones (this work, Eq.2)



function gave slightly better yield and enantiomeric excess than urea one (entries 2 and 4). Moreover 3,5-bis(trifluoromethyl)phenyl derived thiourea gave the best ee (entries 1-3). Finally, the influence of chiral scaffold of the pre-catalyst (R group) was evaluated (entries 2, 7-9), the best yield and enantioselectivity were provided by the pre-catalyst incorporating an *iso*-butyl substituent (entry 7).

Scheme 2. Synthesis of 1,3,4-trisubstituted cyclopentenes catalyzed by NHC via homoenolate intermediate.


We began our study with the cyclopentannulation of 4-methoxy-cinnamaldehyde **5a** and chalcone **6a** in THF at 30 °C using **4af** as pre-catalyst in the presence of potassium carbonate. The cyclopentene adduct was isolated in low yield and moderate ee (Table 1, entry1). It is noteworthy that only the *trans* diastereoisomer **7a** was observed (*dr*>25:1). Next, catalyst screening was studied. All the previously synthesized pre-catalysts were then evaluated in the same reaction conditions (Table 1). This screening revealed that thiourea

Table 1. Catalyst screening.^[a]

Entry	Catalyst	Yield (%) ^[b]	ee (%) ^[c]
1	4af	28	51
2	4ae	32	76
3	4ag	36	25
4	4ah	28	73
5	4ai	21	68
6	4aj	45	58
7	4be	36	82
8	4ce	20	50
9	4de	34	81

[a] Reaction conditions: **5a** (0.28 mmol), **6a** (0.2 mmol), cat. (0.04 mmol), K₂CO₃ (0.08 mmol), THF (2 mL). [b] Yield of isolated product. [c] Determined by chiral HPLC analysis.

With the best pre-catalyst in hand, we explored reaction conditions to increase the efficiency of this transformation. To obtain the *trans*-cyclopentene adduct in high ee, the use of K₂CO₃ as a base was crucial (Table 2, entries 1-4). Solvent was also screened. Among the solvents tested, DIPE was found to be the best to afford the expected product in 43 % yield and 89% ee. Next, different phenol derivatives were used as additives in this transformation.^[19] The number of hydroxyl group on the benzene ring was crucial (Table 2, entries 8-10, see Supplementary Part, Table S3), the catechol amount could be reduced to 20 mol % (entry 11) and better yield was obtained at 40 °C (entry 12). The increase in yield by using this additive might be explained by the fact that catechol could assist a proton transfer to generate homoenolate intermediate^[19c] or by the activation of the enone via a formation of hydrogen bonding.^[19d] In summary, using 20 mol% of catalyst **4be**, 40 mol% K₂CO₃ and 20 mol% catechol in DIPE at 40 °C afforded only the *trans*-diastereoisomer (> 25:1 dr) in 50% yield with 93% ee. Optimization of the reaction conditions allowed us to increase the yield from 36% to 50%. Generally speaking, although yields remained moderate in almost all cases, starting materials could be recovered and reused without difficulty. Moreover, it should be emphasized that this cyclopentannulation is an atom economy transformation since only carbon dioxide is released. Besides, the absolute configuration of the product was assigned as (*S,S*) in comparison with data reported in the literature.^[17a]

Table 2. Optimization of the Reaction Conditions.




Entry	Base	Solvent	Additive (1 equiv)	Yield (%) ^[c]	ee (%) ^[d]
1	K ₂ CO ₃	THF	-	36	82
2	DBU	THF	-	58	36

3	KHMDS	THF	-	40	51
4	t-BuOK	THF	-	50	46
5	K ₂ CO ₃	Et ₂ O	-	48	80
6	K ₂ CO ₃	toluene	-	28	81
7	K ₂ CO ₃	DIPE	-	43	89
8	K ₂ CO ₃	DIPE	catechol	24	95
9	K ₂ CO ₃	DIPE	pyrogallol	5	94
10	K ₂ CO ₃	DIPE	phenol	45	89
11	K ₂ CO ₃	DIPE	catechol ^[a]	42	94
12	K ₂ CO ₃	DIPE	Catecho ^[a,b]	50	93


[a] 20 mol% of catechol was used. [b] The reaction was performed at 40 °C. [c] Yield of isolated product. [d] Determined by chiral HPLC analysis.

With the optimized reaction conditions, we next evaluated the scope of cycloannulation by varying the enal substrates **5a-l**. As outlined in Scheme 3, (*Z*) or (*E*)-cinnamaldehyde afforded the same *trans*-cycloadduct **7c** with comparable yield and enantioselectivity likely due to an isomerization during the formation of the homoenolate intermediate.^[20] Enals with substituents of different electronic nature on phenyl ring could be tolerated affording the cycloadduct **7a-h** in the same range of enantioselectivity (89-96%) in 41-60% yield. The best result with 60% yield and 93% ee was observed using *p*-fluorocinnamaldehyde **5f** as substrate. When the phenyl ring was changed to the naphthyl or furane ring, the reaction also worked well and the products **7i** and **7j** were isolated in 43-48% yield and 90-92% ee. The reaction was also compatible with enals bearing an alkyl chain, producing **7k** with 93% ee, albeit in lower yield (30 %). Furthermore, a styryl-substituted enal underwent the cycloannulation to yield **7l** in 54% yield and with 90% ee.


Scheme 3. Enal scope. [a] **5** (0.28 mmol), **6a** (0.2 mmol), cat. (0.04 mmol), K₂CO₃ (0.08 mmol), catechol (0.04 mmol), DIPE (2 mL). [b] Yield of isolated product. [c] ee determined by chiral HPLC analysis.




7i 43% yield, 92% ee



7j 48% yield, 90% ee



7k 30% yield, 93% ee



7l 54% yield, 90% ee

To broaden the scope of the *trans*-cycloannulation, enal **5f** was chosen to react with various unsaturated ketones or esters **6a-s** (Scheme 4). First, the scope of chalcone derivatives was examined. Electron-donating and electron-withdrawing aromatic substituents at either the β -position (R²) or the carbonyl (R³) of the chalcone are well accommodated, *trans*-cyclopentene were isolated in 41-69% yield and 86-96% ee with the exception of **7fc** (28% yield and 83% ee). This result might be explained by the presence of a nitro substituent, which could coordinate to the catalyst, thereby reducing its efficiency. Furthermore, heteroaromatic or vinylphenyl on the β -position or the carbonyl of the chalcone was also well tolerated, giving the products **7fo-fq** in moderate yield and high enantioselectivity. Finally, the reaction was conducted with 4-oxoenone and 4-oxoenone affording respectively *trans*-cyclopentene **7fs** and **7fr** as the major diastereoisomer but in low yield and enantioselectivity along with lower diastereoselectivity. The formation of the major *trans* diastereoisomer is notable considering the result previously reported by Bode.^[15]

DIPE and the only side product of this reaction is CO₂. Moreover, the efficiency of this methodology was proved by the large scope of this transformation and its scalability. Further investigations to expand the utilization of this new class of bifunctional NHC catalysts are currently on-going in our laboratory.

Experimental Section

General procedure for the synthesis of 7. To an oven-dried 10 mL Schlenk tube equipped with a magnetic stirring bar was charged with enone **6** (0.2 mmol, 1.0 equiv.), enal **5** (0.28 mmol, 1.4 equiv.), K₂CO₃ (11.1 mg, 0.08 mmol, 0.4 equiv.), catechol (4.5 mg, 0.04 mmol, 0.2 equiv.) and NHC precatalyst **4be** (28.1 mg, 0.04 mmol, 0.2 equiv.). This tube was closed with a septum, evacuated, and back-filled with argon. To this mixture, at room temperature, was added freshly distilled diisopropyl ether (2 mL). The reaction mixture was warmed to 40 °C and stirred for 72 h under argon atmosphere. After removing solvent, the residue was directly subjected to silica gel column chromatography (petroleum ether/ethyl acetate as eluent, typically 100/1 to 100/1.5) to give desired single *trans*-product **7** (unless otherwise noted). All the corresponding racemic samples for the standard of chiral HPLC spectra were obtained according to the known procedure by using achiral NHC precatalyst 1,3-dimesityl imidazolium chloride.^[10]

For further experimental details including experimental procedures for preparation of organocatalyst, copies of NMR spectra and HPLC data, see Supporting Information.

Acknowledgements

We are grateful to the 'Chinese Government' (China Scholarship Council) for a doctoral fellowship to Z. Jiang, the CNRS (UMR 8182) and the University Paris-Sud/ University Paris Saclay for financial supports. We also thank Emilie Kolodziej for technical assistance with the HPLC analysis. This work has benefited from the facilities and expertise of the Small Molecule Mass Spectrometry platform of ICSN (Centre de Recherche de Gif - www.icsn.cnrs-gif.fr)

Keywords: Asymmetric catalysis • cyclopentannulation • *N*-heterocyclic carbenes • homoenolate • organocatalysis

- [1] For recent selected reviews, see: a) J. G. Hernandez, E. Juaristi, *Chem. Commun.* **2012**, *48*, 5396-5409; b) E. Negishi, *J. Synth. Org. Chem., Jpn.* **2011**, *69*, 1201-1201; c) M. Tavakolian, S. Vahdati-Khajeh, S. Asgari, *ChemCatChem* **2019**, *11*, 2943-2977; d) C. Jamieson, A. J. B. Watson, in *Catalysis for Sustainability: Goals, Challenges, and Impacts*, Chap. 3 (Ed.: T. A. Umile), Taylor and Francis, Abingdon, **2016**, pp. 47-82; e) G. Guillena, D. Alonso, A. Baeza, R. Chinchilla, J. Flores-Ferrandiz, M. Gomez-Martinez, P. Trillo, *Current Organocatalysis* **2015**, *2*, 102-123; f) P. T. Anastas, J. B. Zimmerman, *Green Chem.* **2019**, *21*, 6545-6566; g) A. Collado, A. Gómez-Suárez, S. P. Nolan in *Sustainable Catalysis : With Non-endangered Metals, Part 2, Chap. 16* (Ed.: M. North), The Royal Society of Chemistry, London, **2016**, pp. 41-90; h) N. V. Tzouras, I. K. Stamatopoulos, A. T. Papastavrou, A. A. Liori, G. C. Vougioukalakis, *Coord. Chem. Rev.* **2017**, *343*, 25-138; i) R. A. Sheldon, D. Brady, *ChemSusChem* **2019**, *12*, 2859-2881; j) P. Chauhan, S. S. Chimni. *Beilstein J. Org. Chem.* **2012**, *8*, 2132-2141.
- [2] For recent selected reviews, see: a) S. Mondal, S. R. Yetra, S. Mukherjee, A. T. Biju, *Acc. Chem. Res.* **2019**, *52*, 425-436; b) S. Mukherjee, A. T. Biju, *Chem. Asian J.* **2018**, *13*, 2333-2349; c) X.-Y. Chen, S. Li, F. Vetica, M. Kumar, D. Enders, *iScience* **2018**, *2*, 1-26; d) M. Zhao, Y.-T. Zhang, J. Chen, L. Zhou, *Asian J. Org. Chem.* **2018**, *7*, 54-69; e) E. Reyes, U. Uria, L. Carrillo, J. L. Vicario, *Synthesis* **2017**, *49*, 451-471; f) A. Levens, D. W. Lupton, *Synlett* **2017**, *28*, 415-424; g) Q. Ren, M. Li, L. Yuan, J. Wang, *Org. Biomol. Chem.* **2017**, *15*, 4731-4749; h) M. H. Wang, K. A. Scheidt, *Angew. Chem.* **2016**, *128*, 15134-15145; *Angew. Chem. Int. Ed.* **2016**, *55*, 14912-14922; i) D. M. Flanagan, F. Romanov-Michailidis, N. A. White, T. Rovis, *Chem. Rev.* **2015**, *115*, 9307-9387; j) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, *Nature* **2014**, *510*, 485-496; k) S. J. Ryan, L. Candish, D. W. Lupton, *Chem. Soc. Rev.* **2013**, *42*, 4906-4917; l) J. W. Bode, *Nat. Chem.* **2013**, *5*, 813-815; m) A. Grossmann, D. Enders, *Angew. Chem.* **2012**, *124*, 320-332; *Angew. Chem. Int. Ed.* **2012**, *51*, 314-325; n) D. T. Cohen, K. A. Scheidt, *Chem. Sci.* **2012**, *3*, 53-57; o) J. Izquierdo, G. E. Hutson, D. T. Cohen, K. A. Scheidt, *Angew. Chem.* **2012**, *124*, 11854-11866; *Angew. Chem. Int. Ed.* **2012**, *51*, 11686-11698; p) X. Bugaut, F. Glorius, *Chem. Soc. Rev.* **2012**, *41*, 3511-3522; q) A. T. Biju, N. Kuhl, F. Glorius, *Acc. Chem. Res.* **2011**, *44*, 1182-1195; r) D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.* **2007**, *107*, 5606-5655.
- [3] For NHC-catalyzed reactions via the Breslow intermediates, see: a) R. S. Menon, A. T. Biju, V. Nair, *Beilstein J. Org. Chem.* **2016**, *12*, 444-461; b) S. R. Yetra, A. Patra, A. T. Biju, *Synthesis* **2015**, *47*, 1357-1378; c) J. Read de Alaniz, T. Rovis, *Synlett* **2009**, 1189-1207; d) D. Enders, T. Balensiefer, *Acc. Chem. Res.* **2004**, *37*, 534-541. For NHC-catalyzed reactions via homoenolate equivalents, see: e) R. S. Menon, A. T. Biju, V. Nair, *Chem. Soc. Rev.* **2015**, *44*, 5040-5052; f) V. Nair, R. S. Menon, A. T. Biju, C. R. Sinu, R. R. Paul, A. Jose, V. Sreekumar, *Chem. Soc. Rev.* **2011**, *40*, 5336-5346; g) V. Nair, S. Vellalath, B. P. Babu, *Chem. Soc. Rev.* **2008**, *37*, 2691-2698; h) K. J. R. Murauski, A. A. Jaworski, K. A. Scheidt, *Chem. Soc. Rev.* **2018**, *47*, 1773-1782. For NHC-catalyzed reactions via azolium enolates, see: i) X.-Y. Chen, S. Ye, *Synlett* **2013**, *24*, 1614-1622; j) H. U. Vora, P. Wheeler, T. Rovis, *Adv. Synth. Catal.* **2012**, *354*, 1617-1639; k) J. Douglas, G. Churchill, A. D. Smith, *Synthesis* **2012**, *44*, 2295-2309. For NHC-catalyzed reactions via acyl azoliums, see: l) J. Mahatthananchai. *J. W. Bode, Acc. Chem. Res.* **2014**, *47*, 696-707; m) C. E. I. Knappke, A. Imami, A. Jacobi von Wangelin, *ChemCatChem* **2012**, *4*, 937-941; n) H. U. Vora, T. Rovis, *Aldrichimica Acta* **2011**, *44*, 3-11. For NHC-catalyzed reactions via α,β -unsaturated acyl azolium intermediates, see: o) C. Zhang, J. F. Hooper, D. W. Lupton, *ACS Catal.* **2017**, *7*, 2583-2596; p) S. De Sarkar, A. Biswas, R. C. Samanta, A. Studer, *Chem. Eur. J.* **2013**, *19*, 4664-4678. For NHC-catalyzed reactions via azolium dienolates, see: q) X.-Y. Chen, Q. Liu, P. Chauhan, D. Enders, *Angew. Chem.* **2018**, *130*, 3924-3935; *Angew. Chem. Int. Ed.* **2018**, *57*, 3862-3873.
- [4] T. Ukai, R. Tanaka, T. Dokawa, *J. Pharm. Soc. Jpn.* **1943**, *63*, 296-300.
- [5] F. Wöhler, J. Liebig, *Ann. Der. Pharm.* **1832**, *3*, 249-282.
- [6] R. Breslow, *J. Am. Chem. Soc.* **1958**, *80*, 3719-3726.
- [7] A. Igau, H. Grutzmacher, A. Baccaredo, G. Bertrand, *J. Am. Chem. Soc.* **1988**, *110*, 6463-6466.
- [8] A. J. Arduengo III, R. L. Harlow, M. A. Kline, *J. Am. Chem. Soc.* **1991**, *113*, 361-363.

- [9] J. C. Sheehan, D. H. Hunneman, *J. Am. Chem. Soc.* **1966**, *88*, 3666-3667.
- [10] For selected references, see a) For a review see ref 2i and 3e-h. b) S. S. Sohn, E. L. Rosen, J. W. Bode, *J. Am. Chem. Soc.* **2004**, *126*, 14370-14371; c) C. Burstein, F. Glorius, *Angew. Chem.* **2004**, *116*, 6331-6334; *Angew. Chem. Int. Ed.* **2004**, *43*, 6205-6208; d) V. Nair, M. Poonoth, S. Vellalath, E. Suresh, R. Thirumalai, *J. Org. Chem.* **2006**, *71*, 8964-8965; e) M. He, J. W. Bode, *Org. Lett.* **2005**, *7*, 3131-3134; f) J. Seayad, P. K. Patra, Y. Zhang, J. Y. Ying, *Org. Lett.* **2008**, *10*, 953-956; g) J. R. Struble, J. Kaeobamrung, J. W. Bode, *Org. Lett.* **2008**, *10*, 957-960; h) J. Kaeobamrung, J. W. Bode, *Org. Lett.* **2009**, *11*, 677-680; i) V. Nair, S. Vellalath, M. Poonoth, E. Suresh, *J. Am. Chem. Soc.* **2006**, *128*, 8736-8737; j) A. Chan, K. A. Scheidt, *J. Am. Chem. Soc.* **2007**, *129*, 5334-5335; k) V. Nair, S. Vellalath, M. Poonoth, R. Mohan, E. Suresh, *Org. Lett.* **2006**, *8*, 507-509; l) V. Nair, B. P. Babu, S. Vellalath, E. Suresh, *Chem. Commun.* **2008**, 747-749; m) L. Yang, B. Tan, F. Wang, G. Zhong, *J. Org. Chem.* **2009**, *74*, 1744-1746; n) I. R. Siddiqui, A. Srivastava, S. Shamim, A. Srivastava, M. A. Waseem, R. K. P. Singh, *Synlett* **2013**, *24*, 2586-2590; o) S. Urban, M. Tursky, R. Fröhlich, F. Glorius, *Dalton Trans.* **2009**, 6934-6940; p) R. Kyan, K. Sato, N. Mase, N. Watanabe, T. Narumi, *Org. Lett.* **2017**, *19*, 2750-2753. r) C. T. Check, K. P. Jang, C. B. Schwamb, A. S. Wong, M. H. Wang, K. A. Scheidt, *Angew. Chem.* **2015**, *127*, 4338-4342; *Angew. Chem. Int. Ed.* **2015**, *54*, 4264-4268.
- [11] a) L. He, Y.-R. Zhang, X.-L. Huang, S. Ye, *Synthesis* **2008**, 2825-2829; b) X.-L. Huang, X.-Y. Chen, S. Ye, *J. Org. Chem.* **2009**, *74*, 7585-7587; c) T. Wang, X.-L. Huang, S. Ye, *Org. Biomol. Chem.* **2010**, *8*, 5007-5011; d) X.-N. Wang, Y.-Y. Zhang, S. Ye, *Adv. Synth. Catal.* **2010**, *352*, 1892-1895; e) X.-N. Wang, L.-T. Shen, S. Ye, *Org. Lett.* **2011**, *13*, 6382-6385; f) P.-L. Shao, X.-Y. Chen, L.-H. Sun, S. Ye, *Tetrahedron Lett.* **2010**, *51*, 2316-2318; g) T.-Y. Jian, P.-L. Shao, S. Ye, *Chem. Commun.* **2011**, 47, 2381-2383; h) C. Zheng, W. Yao, Y. Zhang, C. Ma, *Org. Lett.* **2014**, *16*, 5028-5031; i) X.-Y. Chen, Z.-H. Gao, S. Ye, *Acc. Chem. Res.* **2020**, *53*, 690-702.
- [12] a) J. P. Brand, J. I. O. Siles, J. Waser, *Synlett* **2010**, 6, 881-884; b) P. Zheng, C. A. Gondo, J. W. Bode, *Chem. Asian J.* **2011**, *6*, 614-620; c) L. Baragwanath, C. A. Rose, K. Zeitler, S. J. Connon, *J. Org. Chem.* **2009**, *74*, 9214-9217; d) F. Nawaz, M. Zaghouni, D. Bonne, O. Chuzel, J. Rodriguez, Y. Coquerel, *Eur. J. Org. Chem.* **2013**, 8253-8264; e) S. M. Mennen, J. T. Blank, M. B. Tran-Dubé, J. E. Imbriglio, S. J. Miller, *Chem. Commun.* **2005**, 195-197; f) S. M. Mennen, J. D. Gipson, Y. R. Kim, S. J. Miller, *J. Am. Chem. Soc.* **2005**, *127*, 1654-1655.
- [13] a) T.-T.-D. Ngo, T.-H. Nguyen, C. Bournaud, R. Guillot, M. Toffano, G. Vo-Thanh, *Asian J. Org. Chem.* **2016**, *5*, 895-899; b) T.-H. Nguyen, M. Toffano, C. Bournaud, G. Vo-Thanh, *Tetrahedron Lett.* **2014**, *55*, 6377-6380; c) A. Thomasset, L. Boucardy, C. Bournaud, R. Guillot, M. Toffano, G. Vo-Thanh, *Synthesis* **2014**, *46*, 242-250; d) A. Aupoix, C. Bournaud, G. Vo-Thanh, *Eur. J. Org. Chem.* **2011**, *15*, 2772-2776.
- [14] a) C. Jahier-Diallo, M. S. T. Morin, P. Queval, M. Rouen, I. Artur, P. Querard, L. Toupet, C. Crévisy, O. Baslé, M. Mauduit, *Chem. Eur. J.* **2015**, *21*, 993-997; b) R. Tarrieu, A. Dumas, J. Thongpaen, T. Vives, T. Roisnel, V. Dorcet, C. Crévisy, O. Baslé, M. Mauduit, *J. Org. Chem.* **2017**, *82*, 1880-1887; c) P. Queval, C. Jahier, M. Rouen, I. Artur, J.-C. Legeay, L. Falivene, L. Toupet, C. Crévisy, L. Cavallo, O. Baslé, M. Mauduit, *Angew. Chem.* **2013**, *125*, 14353-14357; *Angew. Chem. Int. Ed.* **2013**, *52*, 14103-14107.
- [15] a) P.-C. Chiang, J. Kaeobamrung, J. W. Bode, *J. Am. Chem. Soc.* **2007**, *129*, 3520-3521; b) P.-C. Chiang, M. Rommel, J. W. Bode, *J. Am. Chem. Soc.* **2009**, *131*, 8714-8718.
- [16] B. Cardinal-David, D. E. A. Raup, K. A. Scheidt, *J. Am. Chem. Soc.* **2010**, *132*, 5345-5347.
- [17] a) Z. Fu, J. Xu, T. Zhu, W. W. Y. Leong, Y. R. Chi, *Nature Chemistry* **2013**, *5*, 835-839; b) Y. Xie, C. Yu, T. Li, S. Tu, C. Yao, *Chem. Eur. J.* **2015**, *21*, 5355-5359; c) Z. Fu, K. Jiang, T. Zhu, J. Torres, Y. R. Chi, *Angew. Chem.* **2014**, *126*, 6624-6628; *Angew. Chem. Int. Ed.* **2014**, *53*, 6506-6510; d) Z. Jin, S. Chen, Y. Wang, P. Zheng, S. Yang, Y. R. Chi, *Angew. Chem.* **2014**, *126*, 13724-13727; *Angew. Chem. Int. Ed.* **2014**, *53*, 13506-13509; e) Y. Wang, D. Wei, Y. Wang, W. Zhang, M. Tang, *ACS Catal.* **2016**, *6*, 279-289.
- [18] a) Z.-F. Zhang, K.-Q. Chen, C.-L. Zhang, S. Ye, *Chem. Commun.* **2017**, 53, 4327-4330; b) K. C. Seetha Lakshmi, J. Krishnan, C. R. Sinu, S. Varughese, V. Nair, *Org. Lett.* **2014**, *16*, 6374-6377; c) L. Zhao, S. Li, L. Wang, S. Yu, G. Raabe, D. Enders, *Synthesis* **2018**, *50*, 2523-2532.
- [19] a) Y. Reddi, R. B. Sunoj, *ACS Catal.* **2015**, *5*, 1596-1603; b) D. A. DiRocco, T. Rovis, *J. Am. Chem. Soc.* **2011**, *133*, 10402-10405; c) C. M. Filloux, S. P. Lathrop, T. Rovis, *PNAS* **2010**, *107*, 20666-20671; d) T. J. Peelen, Y. Chi, S.H. Gellman, *J. Am. Chem. Soc.* **2005**, *127*, 11598-11599.
- [20] X. Chen, X. Fang, Y. R. Chi, *Chem. Sci.* **2013**, *4*, 2613-2618.
- [21] a) see ref 10i; b) P. Verma, P. A. Patni, R. B. Sunoj, *J. Org. Chem.* **2011**, *76*, 5606-8630; c) L. R. Domingo, R. J. Zaragoza, M. Arnó, *Org. Biomol. Chem.* **2010**, *8*, 4884-4891; d) Y. Wang, D. Wei, W. Zhang, *ChemCatChem* **2018**, *10*, 338-360.