



An introduction to the topological derivative

Samuel Amstutz

► To cite this version:

| Samuel Amstutz. An introduction to the topological derivative. 2021. hal-03311517v1

HAL Id: hal-03311517

<https://hal.science/hal-03311517v1>

Preprint submitted on 1 Aug 2021 (v1), last revised 9 Sep 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An introduction to the topological derivative

S. Amstutz

CMAP - Ecole polytechnique
Route de Saclay, 91120 Palaiseau, France

Abstract

This paper provides a self-contained introduction to the mathematical aspects of the topological derivative. Full justifications are given on simple model problems following a modern approach. Methodological aspects and extensions are discussed, in relation with the literature on the field.

1 Introduction

Let D be a bounded, open subset of \mathbb{R}^N , the "hold-all" domain. With applications to shape optimization in mind, it is mostly assumed that $N = 2$ or $N = 3$, but the case $N = 1$ may have its own interest, and the cases $N > 3$ may be considered for mathematical generality. To fix ideas let $\mathcal{A} = \{\Omega \subset D, \Omega \text{ open}\}$, and consider a function

$$\Omega \in \mathcal{A} \mapsto \mathcal{J}(\Omega) \in \mathbb{R}$$

indifferently called shape functional, or cost function, or criterion. The concept of topological derivative was formally introduced in [48, 26] as the "bubble method" for the optimal design of structures, and the first mathematical justifications appeared in [49, 28]. It can be defined as follows.

Definition 1.1 *Let ω be a bounded, open subset of \mathbb{R}^N . We say that \mathcal{J} admits a topological derivative at $\Omega_0 \in \mathcal{A}$ and at the point $z \in \Omega_0$ with respect to ω if there exists a function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\lim_{\varepsilon \searrow 0} f(\varepsilon) = 0$ such that the following limit exists:*

$$d_T \mathcal{J}(\Omega_0, \omega, z) = \lim_{\varepsilon \searrow 0} \frac{\mathcal{J}(\Omega_0 \setminus (\overline{z + \varepsilon \omega})) - \mathcal{J}(\Omega_0)}{f(\varepsilon)}. \quad (1)$$

Of course, (1) is equivalent to the "topological asymptotic expansion":

$$\mathcal{J}(\Omega_0 \setminus (\overline{z + \varepsilon \omega})) - \mathcal{J}(\Omega_0) = f(\varepsilon) d_T \mathcal{J}(\Omega_0, \omega, z) + o(f(\varepsilon)).$$

The set $\omega_\varepsilon := z + \varepsilon \omega$ is the domain perturbation, see Figure 1. It is often called hole, or cavity, or more specifically inclusion when it is filled with a material phase different from the background one. The set $\Omega_\varepsilon := \Omega_0 \setminus \overline{\omega_\varepsilon}$ is the perturbed or perforated domain. The set ω is the perturbation at scale 1. It is typically chosen as the unit ball, but other cases, like cracks, can be of interest. Clearly, for the volume functional $\mathcal{J}(\Omega) = |\Omega|$, we have $d_T \mathcal{J}(\Omega, \omega, z) = -1$ for all $z \in \Omega$ with $f(\varepsilon) = |\omega_\varepsilon| = \varepsilon^N |\omega|$. The perimeter functional $\mathcal{J}(\Omega) = \int_{\partial \Omega} ds$ yields $d_T \mathcal{J}(\Omega, \omega, z) = 1$ for all $z \in \Omega$ with $f(\varepsilon) = \varepsilon^{N-1} \int_{\partial \omega} ds$.

In most practical cases the shape functional involves the solution of a boundary value problem, the so-called (direct) state. Then the type of boundary condition around the hole has a drastic effect on

its asymptotic behavior. One case turns out to be much easier to analyze than the others: it is when the perturbation is on coefficients of non-principal parts of the differential operator. This situation is studied first, while the main content of this presentation concerns perturbations of the principal part in the form of transmission, Neumann and Dirichlet conditions. In this introductory paper we devote to provide a rigorous mathematical analysis of the aforementioned issues on simple model problems. In particular we restrict ourselves to Laplace-type elliptic boundary value problems and simple shape functionals. To ease the reading, the use of boundary integral equations is bypassed, and non-standard function spaces are described in some details as soon as they are needed. Extensions and refinements are discussed in comments, in connection with the related literature.

From the technical standpoint, a crucial aspect of the topological sensitivity analysis is the incorporation of the adjoint state while we are typically not working in a differentiable framework. This is in contrast with smooth shape variations and the more classical notion of shape derivative [38, 1, 33, 50]. Several approaches have been developed to overcome this issue, including the domain truncation method [36, 28, 29, 42, 41], the generalized Lagrangian method [4, 5, 8, 6], and the topological shape sensitivity method [39, 40]. Other authors only incorporate the adjoint state in the end of the derivations [31, 21]. Here we will follow the averaged adjoint method recently introduced in [51, 27] as a modification of the generalized Lagrangian method. This formulation, initially mainly developed for nonlinear purposes, turns out to be somehow natural. It is also elegant in the linear context as it expresses the variation of the cost function exactly as the variation of the Lagrangian.

The paper is organized as follows. Section 2 is dedicated to the perturbation of non-principal parts, where differential calculus and the standard adjoint state method can be used. The averaged adjoint method is presented section 3 in an abstract linear framework. Section 4 deals with the cases of transmission and Neumann conditions in dimension $N \geq 1$. Sections 5 and 6 address the Dirichlet case in 2D and in 3D, respectively, as here these two space dimensions need to be distinguished and the 1D case is meaningless.




Figure 1: Domain perturbation.

2 Perturbation of non-principal parts

In this section we specifically assume that $N \in \{1, 2, 3\}$, although higher dimensions could be addressed upon minor adaptations. A key element of the analysis will be the Sobolev embedding $H_0^1(D) \hookrightarrow L^p(D)$ with p recalled in Table 1, see e.g. [13, 22, 24]. For any measurable set $\Omega \subset D$ we consider the boundary value problem

$$\begin{cases} -\Delta u_\Omega + h_\Omega u_\Omega = f_\Omega & \text{in } D \\ u_\Omega = 0 & \text{on } \partial D, \end{cases}$$

where the functions h_Ω and f_Ω are defined using the characteristic function χ_Ω of Ω by

$$h_\Omega = \chi_\Omega h_1 + (1 - \chi_\Omega)h_0, \quad f_\Omega = \chi_\Omega f_1 + (1 - \chi_\Omega)f_0.$$

The above functions $h_0, h_1 \in L^q(D)$, $f_0, f_1 \in L^r(D)$ are supposed to be given, with $h_0, h_1 \geq 0$ and q, r as in Table 1. Finally we consider a shape functional of the form $\mathcal{J}(\Omega) = J(u_\Omega)$ where $J : H_0^1(D) \rightarrow \mathbb{R}$ is of class \mathcal{C}^2 .

N	p	q	r
1	$\leq +\infty$	> 2	> 2
2	$< +\infty$	> 2	> 2
3	≤ 6	> 6	> 3

Table 1: Lebesgue exponents

The following result extends ideas from [32], see also [40].

Theorem 2.1 *The above shape functional admits at a.e. $z \in \Omega$ the topological derivative*

$$d_T \mathcal{J}(\Omega, \omega, z) = [(h_0 - h_1)u_\Omega v_\Omega - (f_0 - f_1)v_\Omega](z)$$

with $f(z) = \varepsilon^N |\omega|$, where the adjoint state v_Ω is the solution of

$$\int_D (\nabla v_\Omega \cdot \nabla \varphi + h_\Omega v_\Omega \varphi) dx = -dJ(u_\Omega) \varphi dx \quad \forall \varphi \in H_0^1(D).$$

PROOF. Given $p > 2$ as in Table 1 we define, for all $(u, v, h, f) \in H_0^1(D) \times H_0^1(D) \times L^{p/(p-2)}(D) \times L^{p/(p-1)}(D)$, the quantities

$$a(h, u, v) = \int_D (\nabla u \cdot \nabla v + huv) dx, \quad l(f, v) = \int_D f v dx.$$

These expressions are well defined since $uv \in L^{p/2}(D)$ by the Cauchy-Schwarz inequality. For any $(h, f) \in L^{p/(p-2)}(D) \times L^{p/(p-1)}(D)$ we define $A(h) \in \mathcal{L}(H_0^1(D), H^{-1}(D))$ and $L(f) \in H^{-1}(D)$ by

$$\langle A(h)u, v \rangle_{H^{-1}(D), H_0^1(D)} = a(h, u, v), \quad \langle L(f), v \rangle_{H^{-1}(D), H_0^1(D)} = l(f, v).$$

Set

$$\mathcal{U} = \left\{ (h, f) \in L^{p/(p-2)}(D) \times L^{p/(p-1)}(D), h \geq 0 \text{ a.e.} \right\}.$$

By the Lax-Milgram theorem and the Poincaré inequality we have $A(h) \in \text{isom}(H_0^1(D), H^{-1}(D))$ for every (h, f) in a neighborhood of any $(f_0, h_0) \in \mathcal{U}$. In addition, due to its affine structure, the map $(h, f) \in L^{p/(p-2)}(D) \times L^{p/(p-1)}(D) \mapsto (A(h), L(f))$ is of class \mathcal{C}^∞ . It follows that the map $(h, f) \mapsto u(h, f) := A(h)^{-1}L(f)$ is of class \mathcal{C}^∞ in a neighborhood of any element of \mathcal{U} . In order to find a convenient expression of the derivative we use the classical adjoint method, based on the Lagrangian

$$\mathcal{L}(h, f, u, v) = J(u) + a(h, u, v) - l(f, v).$$

We differentiate $j(h, f) := J(u(h, f)) = \mathcal{L}(h, f, u(h, f), v)$ at (h_Ω, f_Ω) for an arbitrary $v \in H_0^1(D)$:

$$dj(h_\Omega, f_\Omega)(\hat{u}, \hat{f}) = d_h \mathcal{L}(h_\Omega, f_\Omega, u_\Omega, v) \hat{h} + d_f \mathcal{L}(h_\Omega, f_\Omega, u_\Omega, v) \hat{f} + d_u \mathcal{L}(h_\Omega, f_\Omega, u_\Omega, v) du(h_\Omega, f_\Omega)(\hat{h}, \hat{f}).$$

Choosing v as the adjoint state v_Ω cancels the last term by construction. We arrive at

$$dj(h_\Omega, f_\Omega)(\hat{u}, \hat{f}) = \int_D \hat{h} u_\Omega v_\Omega dx - \int_D \hat{f} v_\Omega dx.$$

By Taylor-Lagrange expansion this implies

$$j(h_\Omega + \hat{h}, f_\Omega + \hat{f}) - j(h_\Omega, f_\Omega) = \int_D \hat{h} u_\Omega v_\Omega dx - \int_D \hat{f} v_\Omega dx + O(\|\hat{h}\|_{L^{p/(p-2)}(D)}^2 + \|\hat{f}\|_{L^{p/(p-1)}(D)}^2).$$

Choose $z \in \Omega$ and ε small enough so that

$$\hat{h} := h_{\Omega \setminus (\overline{z + \varepsilon \omega})} - h_\Omega = \chi_{z + \varepsilon \omega}(h_0 - h_1), \quad \hat{f} := f_{\Omega \setminus (\overline{z + \varepsilon \omega})} - f_\Omega = \chi_{z + \varepsilon \omega}(f_0 - f_1).$$

It is straightforward from Hölder's inequality that

$$\begin{aligned} \|\hat{h}\|_{L^{p/(p-2)}(D)} &\leq (\varepsilon^N |\omega|)^{1 - \frac{2}{p} - \frac{1}{q}} \|h_0 - h_1\|_{L^q(D)} \\ \|\hat{f}\|_{L^{p/(p-1)}(D)} &\leq (\varepsilon^N |\omega|)^{1 - \frac{1}{p} - \frac{1}{r}} \|f_0 - f_1\|_{L^r(D)}. \end{aligned}$$

With the assumptions made on q and r we can adjust p in accordance with Table 1 in order to have

$$\|\hat{h}\|_{L^{p/(p-2)}(D)}^2 + \|\hat{f}\|_{L^{p/(p-1)}(D)}^2 = o(\varepsilon^N).$$

So far we have shown that

$$\mathcal{J}(\Omega \setminus (\overline{z + \varepsilon \omega})) - \mathcal{J}(\Omega) = \int_{z + \varepsilon \omega} ((h_0 - h_1)u_\Omega v_\Omega - (f_0 - f_1)v_\Omega) dx + o(\varepsilon^N).$$

This can be rewritten as

$$\begin{aligned} \mathcal{J}(\Omega \setminus (\overline{z + \varepsilon \omega})) - \mathcal{J}(\Omega) &= |\omega| \varepsilon^N [(h_0 - h_1)u_\Omega v_\Omega - (f_0 - f_1)v_\Omega](z) \\ &+ \int_{z + \varepsilon \omega} (((h_0 - h_1)u_\Omega v_\Omega - (f_0 - f_1)v_\Omega)(x) - ((h_0 - h_1)u_\Omega v_\Omega - (f_0 - f_1)v_\Omega)(z)) dx + o(\varepsilon^N). \end{aligned}$$

By Lebesgue's differentiation theorem this latter integral is a $o(\varepsilon^N)$ for a.e. $z \in \Omega$. \square

We observe here that the topological derivative does not depend on the shape of ω . We will see that this property is usually lost when principal parts are perturbed.

3 A generalized adjoint method

In the previous section we have been able to use Fréchet's differential calculus and the classical Lagrangian method thanks to Sobolev embeddings and the fact that the characteristic function of the hole was small in an appropriate L^p norm. When we perturb the principal part of the differential operator, differential calculus applies in L^∞ , but the L^∞ norm of the characteristic function of a set of positive Lebesgue measure does not go to zero with the diameter of the set. As explained in the introduction we are going to use a generalization of the Lagrangian framework based on the averaged adjoint state idea from [51, 27].

Proposition 3.1 *Let H be a Hilbert space and $\varepsilon_0 > 0$. For every $\varepsilon \in [0, \varepsilon_0]$ we consider :*

- a bilinear form $a_\varepsilon(\cdot, \cdot)$ on H ,

- a linear form $l_\varepsilon(\cdot)$ on H ,
- a direct state $u_\varepsilon \in H$ solution of

$$a_\varepsilon(u_\varepsilon, \varphi) = l_\varepsilon(\varphi) \quad \forall \varphi \in H,$$

- a cost function $J_\varepsilon(\cdot)$ continuously Fréchet differentiable on H ,
- an adjoint state $v_\varepsilon \in H$ solution of

$$a_\varepsilon(\varphi, v_\varepsilon) = - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0)\varphi dt \quad \forall \varphi \in H.$$

Then we have for all $\varepsilon \in [0, \varepsilon_0]$

$$J_\varepsilon(u_\varepsilon) - J_0(u_0) = (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon),$$

with the Lagrangian

$$\mathcal{L}_\varepsilon(u, v) = J_\varepsilon(u) + a_\varepsilon(u, v) - l_\varepsilon(v) \quad \forall (\varepsilon, u, v) \in [0, \varepsilon_0] \times H \times H.$$

PROOF. We have the easy equalities:

$$\begin{aligned} J_\varepsilon(u_\varepsilon) - J_0(u_0) &= \mathcal{L}_\varepsilon(u_\varepsilon, v_\varepsilon) - \mathcal{L}_0(u_0, v_\varepsilon) \\ &= \mathcal{L}_\varepsilon(u_\varepsilon, v_\varepsilon) - \mathcal{L}_\varepsilon(u_0, v_\varepsilon) + \mathcal{L}_\varepsilon(u_0, v_\varepsilon) - \mathcal{L}_0(u_0, v_\varepsilon) \\ &= J_\varepsilon(u_\varepsilon) + a_\varepsilon(u_\varepsilon, v_\varepsilon) - J_\varepsilon(u_0) - a_\varepsilon(u_0, v_\varepsilon) + \mathcal{L}_\varepsilon(u_0, v_\varepsilon) - \mathcal{L}_0(u_0, v_\varepsilon) \\ &= J_\varepsilon(u_\varepsilon) - J_\varepsilon(u_0) + a_\varepsilon(u_\varepsilon - u_0, v_\varepsilon) + \mathcal{L}_\varepsilon(u_0, v_\varepsilon) - \mathcal{L}_0(u_0, v_\varepsilon) \\ &= J_\varepsilon(u_\varepsilon) - J_\varepsilon(u_0) - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0)(u_\varepsilon - u_0)dt + \mathcal{L}_\varepsilon(u_0, v_\varepsilon) - \mathcal{L}_0(u_0, v_\varepsilon). \end{aligned}$$

The first three terms cancel out, leading to the claim. \square

We stress that the variation of the Lagrangian needs to be evaluated at the variable adjoint state v_ε . We will see that in case of topology perturbations, approximating v_ε by v_0 usually yields an error of first order. Note that v_0 is the standard unperturbed adjoint state, solution of

$$a_0(\varphi, v_0) = -dJ(u_0)\varphi \quad \forall \varphi \in H.$$

4 Inclusion and Neumann cases

4.1 Problem formulation

Let Ω be an open and bounded subset of \mathbb{R}^N , $N \in \mathbb{N}^*$, and ω be a bounded, smooth open subset of \mathbb{R}^N . We consider a point $z \in \Omega$ and, for $\varepsilon \geq 0$ small enough, the set

$$\omega_\varepsilon = z + \varepsilon\omega \subset \Omega.$$

We recall the notation $\Omega_\varepsilon = \Omega \setminus \overline{\omega}_\varepsilon$, see Fig. 1. We focus our attention on the problem

$$\text{find } u_\varepsilon \in H_0^1(\Omega) \quad \text{s.t.} \quad \int_\Omega \sigma_\varepsilon \nabla u_\varepsilon \cdot \nabla \varphi dx = \int_\Omega f \varphi dx \quad \forall \varphi \in H_0^1(\Omega), \quad (2)$$

with $f \in L^2(\Omega)$ and the piecewise constant conductivity

$$\sigma_\varepsilon = \chi_{\Omega_\varepsilon} \alpha + \chi_{\omega_\varepsilon} \beta, \quad \alpha > 0, \beta \geq 0.$$

We will distinguish the two cases:

- $\beta > 0$, called the inclusion case,
- $\beta = 0$, called the Neumann case.

For simplicity, and to make the same setting applicable to the two above cases, we assume that $f = 0$ in a neighborhood z , and that ε is small enough so that $f = 0$ in ω_ε .

The inclusion case obviously admits a unique solution, and it corresponds to the strong form

$$\begin{cases} -\operatorname{div}(\sigma_\varepsilon \nabla u_\varepsilon) = f & \text{in } \Omega \\ u_\varepsilon = 0 & \text{on } \partial\Omega. \end{cases} \quad (3)$$

The Neumann case can be equivalently reformulated as

$$\text{find } u_\varepsilon \in H_0^1(\Omega) \quad \text{s.t.} \quad \int_{\Omega_\varepsilon} \alpha \nabla u_\varepsilon \cdot \nabla \varphi dx = \int_{\Omega_\varepsilon} f \varphi dx \quad \forall \varphi \in H_0^1(\Omega). \quad (4)$$

Since every function in $H^1(\Omega_\varepsilon)$ can be extended to a function in $H^1(\Omega)$, we recognize that $u_\varepsilon|_{\Omega_\varepsilon}$ is the weak solution of

$$\begin{cases} -\alpha \Delta u_\varepsilon = f & \text{in } \Omega_\varepsilon \\ \frac{\partial u_\varepsilon}{\partial n} = 0 & \text{on } \partial\omega_\varepsilon \\ u_\varepsilon = 0 & \text{on } \partial\Omega. \end{cases} \quad (5)$$

In this case u_ε is undefined inside ω_ε . The "Neumann" terminology, of course, refers to the boundary condition on the hole. The boundary condition on $\partial\Omega$ plays no significant role in the present study.

In the two cases, (2) corresponds for $\varepsilon = 0$ to the unperturbed problem

$$\begin{cases} -\alpha \Delta u_0 = f & \text{in } \Omega \\ u_0 = 0 & \text{on } \partial\Omega. \end{cases} \quad (6)$$

For consistence with the notation of Proposition 3.1 we set

$$a_\varepsilon(u, v) = \int_{\Omega} \sigma_\varepsilon \nabla u \cdot \nabla \varphi dx, \quad l_\varepsilon(v) = \int_{\Omega} f v dx \quad \forall u, v \in H_0^1(\Omega). \quad (7)$$

For the sake of clarity, we will at first concentrate on the inclusion case, and we will afterwards discuss the adaptations to the Neumann case.

For simplicity we consider a cost function of the form

$$J_\varepsilon(u) = \hat{J}(u|_{\hat{\Omega}}), \quad (8)$$

where $\hat{\Omega}$ is an open subset of Ω excluding a neighborhood of z and $\hat{J} : H^1(\hat{\Omega}) \rightarrow \mathbb{R}$ is Fréchet differentiable. We further assume that $d\hat{J}$ is Lipschitz continuous. Comments will be made regarding more general cost functions in Remark 4.10.

4.2 Variation of the direct state

In order to apply Proposition 3.1, it is needed that the variation of the Lagrangian be evaluated at the averaged adjoint state v_ε . Since the construction of this adjoint state involves the direct state u_ε , our first step is to analyze the behavior of u_ε . We adapt the approach of [27].

Set $\tilde{u}_\varepsilon = u_\varepsilon - u_0$. Subtracting the variational formulations for u_ε and u_0 results in

$$\int_{\Omega} \sigma_\varepsilon \nabla \tilde{u}_\varepsilon \cdot \nabla \varphi dx = (\alpha - \beta) \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla \varphi dx \quad \forall \varphi \in H_0^1(\Omega).$$

We now define the rescaled function

$$U_\varepsilon(y) = \frac{1}{\varepsilon} \tilde{u}_\varepsilon(z + \varepsilon y), \quad y \in \varepsilon^{-1}(\Omega - z),$$

so that

$$\tilde{u}_\varepsilon(x) = \varepsilon U_\varepsilon\left(\frac{x - z}{\varepsilon}\right), \quad \nabla \tilde{u}_\varepsilon(x) = \nabla U_\varepsilon\left(\frac{x - z}{\varepsilon}\right) \quad \forall x \in \Omega.$$

A straightforward change of variables leads to

$$\int_{\varepsilon^{-1}(\Omega - z)} \sigma_\varepsilon(z + \varepsilon y) \nabla U_\varepsilon(y) \cdot \nabla \varphi(z + \varepsilon y) dy = (\alpha - \beta) \int_\omega \nabla u_0(z + \varepsilon y) \cdot \nabla \varphi(z + \varepsilon y) dy \quad \forall \varphi \in H_0^1(\Omega).$$

Changing test functions yields

$$\int_{\varepsilon^{-1}(\Omega - z)} \sigma_\varepsilon(z + \varepsilon y) \nabla U_\varepsilon(y) \cdot \nabla \Phi(y) dy = (\alpha - \beta) \int_\omega \nabla u_0(z + \varepsilon y) \cdot \nabla \Phi(y) dy \quad \forall \Phi \in H_0^1(\varepsilon^{-1}(\Omega - z)). \quad (9)$$

Let us define the reference conductivity field

$$\sigma(y) = \begin{cases} \beta & \text{if } y \in \omega \\ \alpha & \text{if } y \in \mathbb{R}^N \setminus \omega. \end{cases}$$

With the aim of passing to the limit in (9), we will work with the space

$$X = \{u \in L_{\text{loc}}^2(\mathbb{R}^N) : \nabla u \in L^2(\mathbb{R}^N)\},$$

from the family of Beppo-Levi spaces [25], and the associated quotient space X/\mathbb{R} by the equivalence relation

$$u \sim v \Rightarrow \exists c \in \mathbb{R} \text{ s.t. } u - v = c.$$

When it is equipped with the inner product

$$\langle u, v \rangle_{X/\mathbb{R}} = \int_{\mathbb{R}^N} \nabla u \cdot \nabla v dx,$$

the space X/\mathbb{R} is a Hilbert space [44]. By construction the function U_ε belongs to $H_0^1(\varepsilon^{-1}(\Omega - z))$. We implicitly consider an extension by 0 over \mathbb{R}^N . By the Lax-Milgram theorem there exists a unique $U \in X/\mathbb{R}$ solution of

$$\int_{\mathbb{R}^N} \sigma(y) \nabla U(y) \cdot \nabla \Phi(y) dy = (\alpha - \beta) \int_\omega \nabla u_0(z) \cdot \nabla \Phi(y) dy \quad \forall \Phi \in X/\mathbb{R}. \quad (10)$$

The key point is the following convergence result, again adapted from [27]. Alternative estimates can be derived with the help of the fundamental solution of the principal part of the differential operator, see e.g. [5], or by comparison principles [7]. The advantage of the presented approach is that it does not require any knowledge of the behavior of U at infinity. In particular it extends relatively easily to some nonlinear problems [27].

Proposition 4.1 *We have the strong convergence $\nabla U_\varepsilon \rightarrow \nabla U$ in $L^2(\mathbb{R}^N)$ when $\varepsilon \searrow 0$, provided that ∇u_0 be continuous at point z .*

PROOF. Step 1. By the extension convention, $H_0^1(\varepsilon^{-1}(\Omega - z))/\mathbb{R}$ is a closed linear subspace of X/\mathbb{R} . We denote by P_ε the projection of U onto $H_0^1(\varepsilon^{-1}(\Omega - z))/\mathbb{R}$. By a small abuse of notation we will assume that P_ε stands for the representative in $H_0^1(\varepsilon^{-1}(\Omega - z))$. By definition we have

$$P_\varepsilon = \operatorname{argmin}_{\Phi \in H_0^1(\varepsilon^{-1}(\Omega - z))} \|\nabla \Phi - \nabla U\|_{L^2(\mathbb{R}^N)}.$$

Standard properties of the projection onto a linear subspace ensure that $\|\nabla P_\varepsilon\|_{L^2(\varepsilon^{-1}(\Omega - z))} \leq \|\nabla U\|_{L^2(\mathbb{R}^N)}$ and

$$\int_{\mathbb{R}^N} (\nabla P_\varepsilon - \nabla U) \cdot \nabla \Phi dx = 0 \quad \forall \Phi \in H_0^1(\varepsilon^{-1}(\Omega - z)).$$

The first assertion yields that there exists $Q \in X/\mathbb{R}$ such that $\nabla P_\varepsilon \rightharpoonup \nabla Q$ weakly in $L^2(\mathbb{R}^N)$, up to a subsequence. The second assertion implies that

$$\int_{\mathbb{R}^N} (\nabla Q - \nabla U) \cdot \nabla \Phi dx = 0 \quad \forall \Phi \in H_0^1(B(0, R)), \forall R > 0.$$

Let $\zeta : \mathbb{R}^N \rightarrow [0, 1]$ be a smooth function such that $\zeta = 1$ in $B(0, 1)$ and $\zeta = 0$ outside $B(0, 2)$. Set $\zeta_n(x) = \zeta(x/n)$ and $\Phi_n(x) = (Q - U + \lambda_n)\zeta_n \in H_0^1(B(0, 2n))$, with $\lambda_n \in \mathbb{R}$ at the moment arbitrary. This yields

$$\int_{\mathbb{R}^N} |\nabla Q - \nabla U|^2 \zeta_n dx + \int_{\mathbb{R}^N} (Q - U + \lambda_n)(\nabla Q - \nabla U) \cdot \nabla \zeta_n dx = 0. \quad (11)$$

A change of variables entails

$$\|(Q - U + \lambda_n)\nabla \zeta_n\|_{L^2(\mathbb{R}^N)}^2 = n^{N-2} \int_{\mathbb{R}^N} |(Q - U + \lambda_n)(ny)\nabla \zeta(y)|^2 dy \leq cn^{N-2} \int_{R(0,1,2)} |(Q - U + \lambda_n)(ny)|^2 dy,$$

with $R(0, 1, 2)$ the ring centered at 0 with radii 1 and 2. We now fix λ_n such that

$$\int_{R(0,1,2)} (Q - U + \lambda_n)(ny) dy = 0.$$

By the Poincaré-Wirtinger inequality we infer

$$\|(Q - U + \lambda_n)\nabla \zeta_n\|_{L^2(\mathbb{R}^N)}^2 \leq cn^N \int_{R(0,1,2)} |\nabla(Q - U)(ny)|^2 dy \leq c \int_{R(0,n,2n)} |\nabla(Q - U)|^2 dx.$$

Plugging this into (11) and using the Cauchy-Schwarz inequality, we arrive at

$$\int_{\mathbb{R}^N} |\nabla Q - \nabla U|^2 \zeta_n dx \leq c \|\nabla Q - \nabla U\|_{L^2(\mathbb{R}^N)} \left(\int_{R(0,n,2n)} |\nabla(Q - U)|^2 dx \right)^{1/2}.$$

Letting now n go to $+\infty$ results in $\nabla Q = \nabla U$. Therefore the whole sequence ∇P_ε weakly converges to ∇U . In particular we infer that

$$\int_{\mathbb{R}^N} \nabla P_\varepsilon \cdot \nabla U dx \rightarrow \int_{\mathbb{R}^N} |\nabla U|^2 dx.$$

From the identity

$$\|\nabla P_\varepsilon - \nabla U\|_{L^2(\mathbb{R}^N)}^2 = \|\nabla P_\varepsilon\|_{L^2(\mathbb{R}^N)}^2 + \|\nabla U\|_{L^2(\mathbb{R}^N)}^2 - 2 \int_{\mathbb{R}^N} \nabla P_\varepsilon \cdot \nabla U dx$$

we derive

$$\limsup_{\varepsilon \rightarrow 0} \|\nabla P_\varepsilon - \nabla U\|_{L^2(\mathbb{R}^N)}^2 \leq \limsup_{\varepsilon \rightarrow 0} \|\nabla P_\varepsilon\|_{L^2(\mathbb{R}^N)}^2 - \|\nabla U\|_{L^2(\mathbb{R}^N)}^2 \leq 0.$$

We have shown that ∇P_ε strongly converges to ∇U in $L^2(\mathbb{R}^N)$.

Step 2. Using (9) we obtain

$$\begin{aligned} \int_{\varepsilon^{-1}(\Omega-z)} \sigma(y) \nabla(P_\varepsilon - U_\varepsilon)(y) \cdot \nabla \Phi(y) dy &= \int_{\varepsilon^{-1}(\Omega-z)} \sigma(y) \nabla P_\varepsilon(y) \cdot \nabla \Phi(y) dy \\ &\quad - (\alpha - \beta) \int_{\omega} \nabla u_0(z + \varepsilon y) \cdot \nabla \Phi(y) dy \quad \forall \Phi \in H_0^1(\varepsilon^{-1}(\Omega - z)). \end{aligned}$$

In view of (10) this rewrites as

$$\begin{aligned} \int_{\varepsilon^{-1}(\Omega-z)} \sigma(y) \nabla(P_\varepsilon - U_\varepsilon)(y) \cdot \nabla \Phi(y) dy &= \int_{\varepsilon^{-1}(\Omega-z)} \sigma(y) \nabla(P_\varepsilon - U)(y) \cdot \nabla \Phi(y) dy \\ &\quad - (\alpha - \beta) \int_{\omega} (\nabla u_0(z + \varepsilon y) - \nabla u_0(z)) \cdot \nabla \Phi(y) dy \quad \forall \Phi \in H_0^1(\varepsilon^{-1}(\Omega - z)). \end{aligned}$$

Choose $\Phi = P_\varepsilon - U_\varepsilon$. We obtain for some constant c

$$\begin{aligned} \|\nabla(P_\varepsilon - U_\varepsilon)\|_{L^2(\varepsilon^{-1}(\Omega-z))}^2 &\leq c(\|\nabla(P_\varepsilon - U)\|_{L^2(\varepsilon^{-1}(\Omega-z))} \|\nabla(P_\varepsilon - U_\varepsilon)\|_{L^2(\varepsilon^{-1}(\Omega-z))} \\ &\quad + \|\nabla u_0(z + \varepsilon y) - \nabla u_0(z)\|_{L^2(\omega)} \|\nabla(P_\varepsilon - U_\varepsilon)\|_{L^2(\varepsilon^{-1}(\Omega-z))}), \end{aligned}$$

leading to

$$\|\nabla(P_\varepsilon - U_\varepsilon)\|_{L^2(\varepsilon^{-1}(\Omega-z))} \leq c(\|\nabla(P_\varepsilon - U)\|_{L^2(\varepsilon^{-1}(\Omega-z))} + \|\nabla u_0(z + \varepsilon y) - \nabla u_0(z)\|_{L^2(\omega)}).$$

Using step 1 and the continuity assumption we arrive at $\|\nabla(P_\varepsilon - U_\varepsilon)\|_{L^2(\varepsilon^{-1}(\Omega-z))} \rightarrow 0$.

Conclusion. The proof is completed by combining step 1 and step 2. \square

Corollary 4.2 *Under the assumption of Proposition 4.1 we have*

$$\|\tilde{u}_\varepsilon\|_{H^1(\Omega)}^2 = O(\varepsilon^N),$$

and for any $R > 0$

$$\|\tilde{u}_\varepsilon\|_{H^1(\Omega \setminus B(z, R))}^2 = o(\varepsilon^N).$$

PROOF. By change of variables it is straightforward that

$$\|\nabla \tilde{u}_\varepsilon\|_{L^2(\Omega)}^2 = \varepsilon^N \|\nabla U_\varepsilon\|_{L^2(\varepsilon^{-1}(\Omega-z))}^2.$$

Proposition 4.1 yields that $\|\nabla U_\varepsilon\|_{L^2(\varepsilon^{-1}(\Omega-z))} = O(1)$, whereby $\|\tilde{u}_\varepsilon\|_{H^1(\Omega)}^2 = O(\varepsilon^N)$ by the Poincaré inequality. Let now $R > 0$. The same change of variables provides

$$\|\nabla \tilde{u}_\varepsilon\|_{L^2(\Omega \setminus B(z, R))}^2 = \varepsilon^N \|\nabla U_\varepsilon\|_{L^2(\varepsilon^{-1}(\Omega-z)) \setminus B(0, \varepsilon^{-1}R)}^2 = \varepsilon^N \|\nabla U_\varepsilon\|_{L^2(\mathbb{R}^N \setminus B(0, \varepsilon^{-1}R))}^2.$$

This can be rephrased as

$$\|\nabla \tilde{u}_\varepsilon\|_{L^2(\Omega \setminus B(z, R))}^2 = \varepsilon^N \int_{\mathbb{R}^N} (1 - \chi_{B(0, \frac{R}{\varepsilon})}(y)) |\nabla U_\varepsilon(y)|^2 dy.$$

Using Proposition 4.1 we infer $\|\nabla \tilde{u}_\varepsilon\|_{L^2(\Omega \setminus B(z, R))}^2 = o(\varepsilon^N)$. The claim is achieved by the Poincaré inequality. \square

4.3 Variation of the adjoint state

In view of Proposition 3.1 we define the adjoint state $v_\varepsilon \in H_0^1(\Omega)$ solution of

$$\int_{\Omega} \sigma_\varepsilon \nabla v_\varepsilon \cdot \nabla \varphi dx = - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0) \varphi dt \quad \forall \varphi \in H_0^1(\Omega).$$

In particular the unperturbed adjoint state v_0 satisfies

$$\int_{\Omega} \alpha \nabla v_0 \cdot \nabla \varphi dx = -dJ_0(u_0) \varphi \quad \forall \varphi \in H_0^1(\Omega). \quad (12)$$

As for the direct state we set $\tilde{v}_\varepsilon = v_\varepsilon - v_0$. We obtain by difference the equality, valid for all $\varphi \in H_0^1(\Omega)$,

$$\int_{\Omega} \sigma_\varepsilon \nabla \tilde{v}_\varepsilon \cdot \nabla \varphi dx = (\alpha - \beta) \int_{\omega_\varepsilon} \nabla v_0 \cdot \nabla \varphi dx - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0) \varphi dt + dJ_0(u_0) \varphi.$$

By (8) this rewrites

$$\int_{\Omega} \sigma_\varepsilon \nabla \tilde{v}_\varepsilon \cdot \nabla \varphi dx = (\alpha - \beta) \int_{\omega_\varepsilon} \nabla v_0 \cdot \nabla \varphi dx - \int_0^1 \left(d\hat{J}((tu_\varepsilon + (1-t)u_0)|_{\hat{\Omega}}) - d\hat{J}(u_0|_{\hat{\Omega}}) \right) \varphi|_{\hat{\Omega}} dt.$$

We will later justify that the latter integral can be disregarded, therefore we define $w_\varepsilon \in H_0^1(\Omega)$ solution of

$$\int_{\Omega} \sigma_\varepsilon \nabla w_\varepsilon \cdot \nabla \varphi dx = (\alpha - \beta) \int_{\omega_\varepsilon} \nabla v_0 \cdot \nabla \varphi dx \quad \forall \varphi \in H_0^1(\Omega).$$

In order to approximate this w_ε we proceed exactly as for the direct state. We define

$$W_\varepsilon(y) = \frac{1}{\varepsilon} w_\varepsilon(z + \varepsilon y), \quad y \in \varepsilon^{-1}(\Omega - z),$$

and $W \in X/\mathbb{R}$ solution of

$$\int_{\mathbb{R}^N} \sigma(y) \nabla W(y) \cdot \nabla \Phi(y) dy = (\alpha - \beta) \int_{\omega} \nabla v_0(z) \cdot \nabla \Phi(y) dy \quad \forall \Phi \in X/\mathbb{R}. \quad (13)$$

We obtain:

Proposition 4.3 *It holds $\nabla W_\varepsilon \rightarrow \nabla W$ in $L^2(\mathbb{R}^N)$ when $\varepsilon \searrow 0$, provided that ∇v_0 be continuous at point z .*

We now analyze the approximation of \tilde{v}_ε by w_ε .

Lemma 4.4 *It holds*

$$\|\tilde{v}_\varepsilon - w_\varepsilon\|_{H^1(\Omega)}^2 = o(\varepsilon^N).$$

PROOF. Set $e_\varepsilon = \tilde{v}_\varepsilon - w_\varepsilon$. It solves

$$\int_{\Omega} \sigma_\varepsilon \nabla e_\varepsilon \cdot \nabla \varphi dx = - \int_0^1 \left(d\hat{J}((tu_\varepsilon + (1-t)u_0)|_{\hat{\Omega}}) - d\hat{J}(u_0|_{\hat{\Omega}}) \right) \varphi|_{\hat{\Omega}} dt \quad \forall \varphi \in H_0^1(\Omega).$$

We choose $\varphi = e_\varepsilon$. Using that $d\hat{J}$ is Lipschitz we obtain

$$\|\nabla e_\varepsilon\|_{L^2(\Omega)}^2 \leq c \int_0^1 t \|u_\varepsilon - u_0\|_{H^1(\hat{\Omega})} \|e_\varepsilon\|_{H^1(\hat{\Omega})} dt.$$

Corollary 4.2 and the Poincaré inequality provide the desired estimate. \square

4.4 Variation of the Lagrangian

Following Proposition 3.1 and (7) we define the Lagrangian

$$\mathcal{L}_\varepsilon(u, v) = J_\varepsilon(u) + \int_{\Omega} \sigma_\varepsilon \nabla u \cdot \nabla v dx - \int_{\Omega} f v dx. \quad \forall u, v \in H_0^1(\Omega).$$

We need to estimate the variation

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = (\beta - \alpha) \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_\varepsilon dx,$$

which can be decomposed as

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = (\beta - \alpha) \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx + (\beta - \alpha) \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla w_\varepsilon dx + (\beta - \alpha) \int_{\omega_\varepsilon} \nabla u_0 \cdot (\nabla \tilde{v}_\varepsilon - \nabla w_\varepsilon) dx.$$

Lemma 4.5 *If ∇u_0 and ∇v_0 are continuous at point z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \varepsilon^N (\beta - \alpha) |\omega| \nabla u_0(z) \cdot \nabla v_0(z) + (\beta - \alpha) \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla w_\varepsilon dx + o(\varepsilon^N).$$

PROOF. We first estimate

$$\begin{aligned} \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx - \varepsilon^N |\omega| \nabla u_0(z) \cdot \nabla v_0(z) &= \int_{\omega_\varepsilon} (\nabla u_0 \cdot \nabla v_0 - \nabla u_0(z) \cdot \nabla v_0(z)) dx \\ &= \varepsilon^N \int_{\omega} (\nabla u_0(z + \varepsilon y) \cdot \nabla v_0(z + \varepsilon y) - \nabla u_0(z) \cdot \nabla v_0(z)) dy \\ &= o(\varepsilon^N). \end{aligned}$$

Secondly, the Cauchy-Schwarz inequality yields

$$\left| \int_{\omega_\varepsilon} \nabla u_0 \cdot (\nabla \tilde{v}_\varepsilon - \nabla w_\varepsilon) dx \right| \leq \|\nabla u_0\|_{L^2(\omega_\varepsilon)} \|\nabla \tilde{v}_\varepsilon - \nabla w_\varepsilon\|_{L^2(\omega_\varepsilon)} = O(\varepsilon^{N/2}) o(\varepsilon^{N/2}),$$

by Lemma 4.4. □

From the expression found in Lemma 4.5 we make a change of variables to obtain

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \varepsilon^N (\beta - \alpha) |\omega| \nabla u_0(z) \cdot \nabla v_0(z) + \varepsilon^N (\beta - \alpha) \int_{\omega} \nabla u_0(z + \varepsilon y) \cdot \nabla W_\varepsilon(y) dy + o(\varepsilon^N).$$

Lemma 4.6 *If ∇u_0 and ∇v_0 are continuous at point z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \varepsilon^N (\beta - \alpha) |\omega| \nabla u_0(z) \cdot \nabla v_0(z) + \varepsilon^N (\beta - \alpha) \int_{\omega} \nabla u_0(z) \cdot \nabla W(y) dy + o(\varepsilon^N).$$

PROOF. We have to show that

$$\lim_{\varepsilon \rightarrow 0} \int_{\omega} (\nabla u_0(z + \varepsilon y) \cdot \nabla W_\varepsilon(y) - \nabla u_0(z) \cdot \nabla W(y)) dy = 0.$$

It is an immediate consequence of Proposition 4.3, using

$$\begin{aligned} &\int_{\omega} (\nabla u_0(z + \varepsilon y) \cdot \nabla W_\varepsilon(y) - \nabla u_0(z) \cdot \nabla W(y)) dy \\ &= \int_{\omega} (\nabla u_0(z + \varepsilon y) - \nabla u_0(z)) \cdot \nabla W_\varepsilon(y) dy + \int_{\omega} \nabla u_0(z) \cdot (\nabla W_\varepsilon(y) - \nabla W(y)) dy \end{aligned}$$

and the Cauchy-Schwarz inequality. □

4.5 Polarization matrix

The definition (13) of W shows that W depends linearly on $\nabla v_0(z)$. Denoting by (e_1, \dots, e_N) the canonical basis of \mathbb{R}^N and by $\zeta_i \in X/\mathbb{R}$ the solution of

$$\int_{\mathbb{R}^N} \sigma(y) \nabla \zeta_i(y) \cdot \nabla \Phi(y) dy = (\alpha - \beta) \int_{\omega} e_i \cdot \nabla \Phi(y) dy \quad \forall \Phi \in X/\mathbb{R}, \quad (14)$$

we identify that $W(y) = \nabla v_0(z) \cdot \zeta(y)$. It follows that

$$\int_{\omega} \nabla u_0(z) \cdot \nabla W(y) dy = \nabla u_0(z) \cdot \left(\int_{\omega} D\zeta(y)^\top dy \right) \nabla v_0(z).$$

Choosing $\Phi = \zeta_j$ in (14) yields

$$(\alpha - \beta) \int_{\omega} e_i \cdot \nabla \zeta_j(y) dy = \int_{\mathbb{R}^N} \sigma(y) \nabla \zeta_i(y) \cdot \nabla \zeta_j(y) dy = (\alpha - \beta) \int_{\omega} e_j \cdot \nabla \zeta_i(y) dy.$$

This shows that

$$Q := \int_{\omega} D\zeta(y) dy = Q^\top.$$

We arrive at

$$|\omega| \nabla u_0(z) \cdot \nabla v_0(z) + \int_{\omega} \nabla u_0(z) \cdot \nabla W(y) dy = \nabla u_0(z) \cdot (|\omega| I + Q) \nabla v_0(z). \quad (15)$$

Definition 4.7 *We call polarization matrix the symmetric matrix*

$$\mathcal{P} = \left(\frac{\beta}{\alpha} - 1 \right) (|\omega| I + Q).$$

Note that (14) can be equivalently rewritten as

$$\int_{\mathbb{R}^N} \sigma(y) \nabla \zeta_i(y) \cdot \nabla \Phi(y) dy = (\alpha - \beta) \int_{\partial\omega} e_i \cdot n(y) \Phi(y) dy \quad \forall \Phi \in X/\mathbb{R}, \quad (16)$$

therefore the corresponding strong form reads

$$\begin{cases} \Delta \zeta_i = 0 \text{ in } \mathbb{R}^N \setminus \partial\omega \\ \beta \left(\frac{\partial \zeta_i}{\partial n} \right)_{\text{int}} - \alpha \left(\frac{\partial \zeta_i}{\partial n} \right)_{\text{ext}} = (\alpha - \beta) e_i \cdot n \text{ on } \partial\omega. \end{cases} \quad (17)$$

Let us now give an additional property of the polarization matrix.

Proposition 4.8 *The eigenvalues (λ_i) of the polarization matrix satisfy the inequality*

$$\lambda_i \leq \left(\frac{\beta}{\alpha} - 1 \right) |\omega|.$$

Moreover the polarization matrix is

- symmetric positive definite if $\beta > \alpha$,
- symmetric negative definite if $\beta < \alpha$.

1D	2D	3D
$\mathcal{P} = 2 \left(1 - \frac{\alpha}{\beta}\right)$	$\mathcal{P} = 2\pi \frac{\beta - \alpha}{\beta + \alpha} I$	$\mathcal{P} = 4\pi \frac{\beta - \alpha}{\beta + 2\alpha} I$

Table 2: Polarization matrix in the inclusion case for $\omega = B(0, 1)$

PROOF. Since \mathcal{P} is symmetric, let us choose an othogonal basis in which it is diagonal. Choosing $\Phi = \zeta_i$ in (14) entails

$$(\alpha - \beta) \int_{\omega} e_i \cdot \nabla \zeta_i(y) dy \geq 0.$$

We infer that

$$\mathcal{P} e_i \cdot e_i = \left(\frac{\beta}{\alpha} - 1\right) \left(|\omega| + \int_{\omega} e_i \cdot \nabla \zeta_i(y) dy\right) \leq \left(\frac{\beta}{\alpha} - 1\right) |\omega|.$$

Case $\beta < \alpha$. The above inequality directly shows that $\mathcal{P} e_i \cdot e_i < 0$, hence \mathcal{P} is symmetric negative definite.

Case $\beta > \alpha$. We write

$$\mathcal{P} e_i \cdot e_i = \left(\frac{\beta}{\alpha} - 1\right) \int_{\omega} (\nabla \zeta_i + e_i) \cdot e_i dy = \left(\frac{\beta}{\alpha} - 1\right) \int_{\omega} (|\nabla \zeta_i + e_i|^2 - (\nabla \zeta_i + e_i) \cdot \nabla \zeta_i) dy.$$

Using (14) we obtain

$$(\alpha - \beta) \int_{\omega} (\nabla \zeta_i + e_i) \cdot \nabla \zeta_i dy = (\alpha - \beta) \int_{\omega} |\nabla \zeta_i|^2 dy + \int_{\mathbb{R}^N} \sigma |\nabla \zeta_i|^2 dy = \alpha \int_{\mathbb{R}^N} |\nabla \zeta_i|^2 dy.$$

This yields

$$\mathcal{P} e_i \cdot e_i = \left(\frac{\beta}{\alpha} - 1\right) \int_{\omega} |\nabla \zeta_i + e_i|^2 dy + \int_{\mathbb{R}^N} |\nabla \zeta_i|^2 dy > 0,$$

hence \mathcal{P} is symmetric positive definite. \square

The notion of polarization matrix goes back at least to [47, 45]. Detailed properties and generalizations can be found in [2]. In case of a vector-valued state, a polarization tensor is involved. In the framework of linear elasticity it is called the elastic moments tensor.

We end this section by providing closed forms of the polarization matrix for ball-shaped inclusions, i.e., when ω is the unit ball $B(0, 1)$. The main step is to solve (17). The typical method is to "guess" a candidate solution, including free parameters, and to plug this ansatz in the system to fix the parameters. For instance, in the 2D case, we find the solution

$$\zeta_i(y) = \frac{\alpha - \beta}{\alpha + \beta} \times \begin{cases} e_i \cdot y & \text{in } \omega \\ \frac{e_i \cdot y}{|y|^2} & \text{in } \mathbb{R}^2 \setminus \bar{\omega}, \end{cases}$$

Results are gathered in Table 2.

4.6 Expression of the topological asymptotic expansion

Let us recapitulate our findings.

Theorem 4.9 *Consider a cost function of form (8). Let $v_0 \in H_0^1(\Omega)$ be the solution of (12). Then*

$$J_{\varepsilon}(u_{\varepsilon}) - J_0(u_0) = \varepsilon^N \alpha \nabla u_0(z) \cdot \mathcal{P} \nabla v_0(z) + o(\varepsilon^N),$$

where \mathcal{P} is the polarization matrix.

ball (2D, 3D)	straight crack (2D)	circular crack (3D)
$\mathcal{P} = -2\pi I$	$\mathcal{P} = -\pi\nu \otimes \nu$	$\mathcal{P} = -\frac{8}{3}\nu \otimes \nu$

Table 3: Polarization matrix in the Neumann case

PROOF. It follows from Proposition 3.1, Lemma 4.6, (15) and Definition 4.7. The regularity conditions for u_0 and v_0 are ensured by elliptic regularity. \square

As a consequence of Theorem 4.9 and Definition 1.1 we can set the topological derivative of a shape functional \mathcal{J} such that $\mathcal{J}(\Omega_\varepsilon) = J_\varepsilon(u_\varepsilon)$ as

$$d_T \mathcal{J}(\Omega, \omega, z) = \alpha \nabla u_0(z) \cdot \mathcal{P} \nabla v_0(z).$$

Remark 4.10 *Other types of cost functions may yield additional terms in the topological derivative, see e.g. [5, 11, 10, 27, 40].*

4.7 The Neumann case

The Neumann case ($\beta = 0$) can be analyzed along the same lines as the inclusion case, with some minor modifications. In particular, some volume integrals over the inclusion have to be replaced by boundary integrals using the Green formula. The function U is now defined in the space X_ω/\mathbb{R} , with

$$X_\omega = \{u \in L^2_{\text{loc}}(\mathbb{R}^N \setminus \omega) : \nabla u \in L^2(\mathbb{R}^N \setminus \bar{\omega})\},$$

by, instead of (10),

$$\int_{\mathbb{R}^N \setminus \omega} \nabla U(y) \cdot \nabla \Phi(y) dy = \int_{\partial\omega} \nabla u_0(z) \cdot n \Phi ds \quad \forall \Phi \in X_\omega/\mathbb{R}.$$

In order to use the semi-norm as a norm on the quotient space X_ω/\mathbb{R} , it is required that $\mathbb{R}^N \setminus \omega$ be connected. This is in principle a rather mild assumption, except in dimension 1 where it cannot hold. Actually, it is clear that in dimension 1 the cost function is likely to be discontinuous at $\varepsilon = 0$. This singularity can also be seen by observing that the polarization of the inclusion diverges when $\beta \rightarrow 0$, see Table 2. The same phenomenon occurs in higher dimension with higher order differential operators [11].

Eventually, in dimension $N \geq 2$ with $\mathbb{R}^N \setminus \omega$ connected, we arrive at the same result as in Theorem 4.9 with the polarization matrix defined through $\zeta_i \in X_\omega/\mathbb{R}$,

$$\int_{\mathbb{R}^N \setminus \omega} \nabla \zeta_i(y) \cdot \nabla \Phi(y) dy = \int_{\partial\omega} e_i \cdot n \Phi(y) dy \quad \forall \Phi \in X_\omega/\mathbb{R}, \quad Q := \int_{\partial\omega} \zeta \otimes n ds.$$

Note that (17) still holds, with $\beta = 0$. Therefore, the expressions given in Table 2 apply with $\beta = 0$.

An interesting extension is the case of a crack with Neumann boundary condition on each face. A particular case is the planar crack $\omega = \{x \in \mathbb{R}^N : x \cdot \nu = 0, |x| < 1\}$ of unit normal ν . The corresponding polarization matrix in dimensions 2 and 3 are reported in Table 3, see [9, 8, 15, 19].

4.8 Extensions

The topological derivative concept in the inclusion or Neumann cases has been developed in many directions. To give a non-exhaustive overview, let us mention the linear elasticity case [28, 21], the Maxwell equations [37], Stokes flows [16], nonlinear problems [7, 27], evolution problems [12, 17, 15], higher order differential equations [11, 14, 20], higher order topological derivatives [18, 19, 34, 43]; see also the monograph [40].

5 Dirichlet case in 2D

5.1 Problem formulation

Let Ω be an open and bounded subset of \mathbb{R}^2 and ω be a bounded, smooth open subset of \mathbb{R}^2 . Like in the previous section we consider a point $z \in \Omega$ and, for $\varepsilon \geq 0$ small enough, the perforated domain

$$\Omega_\varepsilon = \Omega \setminus \overline{\omega_\varepsilon}, \quad \omega_\varepsilon = z + \varepsilon\omega \subset \Omega.$$

We assume for convenience, but without loss of generality, that $0 \in \omega \subset\subset B(0, 1)$. We address the problem

$$\begin{cases} -\Delta u_\varepsilon = f & \text{in } \Omega_\varepsilon \\ u_\varepsilon = 0 & \text{on } \partial\Omega_\varepsilon. \end{cases} \quad (18)$$

It is assumed that $f \in L^2(\Omega)$ with $f = 0$ in a neighborhood of z . We will implicitly suppose that ε is small enough so that $f = 0$ in ω_ε . We denote by u_0 the unperturbed state, solution of

$$\begin{cases} -\Delta u_0 = f & \text{in } \Omega \\ u_0 = 0 & \text{on } \partial\Omega. \end{cases} \quad (19)$$

In order to develop the adjoint method of section 3 in a fixed space we extend u_ε by 0 inside ω_ε . We set

$$a_\varepsilon(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx \quad \forall u, v \in H_0^1(\Omega), \quad (20)$$

$$l_\varepsilon(v) = \int_{\Omega} \nabla u_\varepsilon \cdot \nabla v dx = \int_{\Omega_\varepsilon} \nabla u_\varepsilon \cdot \nabla v dx = \int_{\Omega_\varepsilon} f v dx - \int_{\partial\omega_\varepsilon} \frac{\partial u_\varepsilon}{\partial n} v ds \quad \forall v \in H_0^1(\Omega). \quad (21)$$

By convention the normal to $\partial\omega_\varepsilon$ is chosen outward to ω_ε . This construction ensures that $u_\varepsilon \in H_0^1(\Omega)$ satisfies

$$a_\varepsilon(u_\varepsilon, v) = l_\varepsilon(v) \quad \forall v \in H_0^1(\Omega).$$

We again consider a cost function of the form

$$J_\varepsilon(u) = \hat{J}(u|_{\hat{\Omega}}), \quad (22)$$

where $\hat{\Omega}$ is an open subset of Ω excluding a neighborhood of z and $\hat{J} : H^1(\hat{\Omega}) \rightarrow \mathbb{R}$ is Fréchet differentiable. We further assume that $d\hat{J}$ is Lipschitz continuous.

5.2 A preliminary estimate

We shall work with the weighted Sobolev space [3, 23]

$$W(\mathbb{R}^2) = \{u \in L_{\text{loc}}^2(\mathbb{R}^2) : wu \in L^2(\mathbb{R}^2), \nabla u \in L^2(\mathbb{R}^2)\},$$

with the weight function

$$w(x) = \frac{1}{\sqrt{1 + |x|^2} \log(2 + |x|)}.$$

It is a Hilbert space for the inner product

$$\langle u, v \rangle_{W(\mathbb{R}^2)} = \int_{\mathbb{R}^2} (w^2 uv + \nabla u \cdot \nabla v) dx.$$

We also define the subspace

$$W_0(\mathbb{R}^2 \setminus \bar{\omega}) = \{u \in W(\mathbb{R}^2 \setminus \bar{\omega}) : u = 0 \text{ on } \partial\omega\}.$$

We have the following standard Poincaré inequality, for which we sketch a proof for completeness.

Proposition 5.1 *There exists $c > 0$ such that*

$$\|u\|_{W(\mathbb{R}^2 \setminus \bar{\omega})} \leq c \|\nabla u\|_{L^2(\mathbb{R}^2 \setminus \bar{\omega})} \quad \forall u \in W_0(\mathbb{R}^2 \setminus \bar{\omega}).$$

PROOF. Step 1. Consider first a function $u \in \mathcal{C}_c^\infty(\mathbb{R}^2 \setminus \bar{B}(0, a))$, $a > 1$. For an arbitrary unit vector e we set $f(r) = u(re)$. Integration by parts yields

$$\int_a^{+\infty} \frac{1}{r \log^2 r} f(r)^2 dr = \int_a^{+\infty} \frac{2}{\log r} f(r) f'(r) dr,$$

whereby we obtain by the Cauchy-Schwarz inequality

$$\int_a^{+\infty} \frac{1}{r \log^2 r} f(r)^2 dr \leq 2 \left(\int_a^{+\infty} \frac{1}{r \log^2 r} f(r)^2 dr \right)^{1/2} \left(\int_a^{+\infty} r f'(r)^2 dr \right)^{1/2}.$$

This implies

$$\int_a^{+\infty} \frac{1}{r \log^2 r} f(r)^2 dr \leq 4 \int_a^{+\infty} r f'(r)^2 dr,$$

thence

$$\|u\|_{W(\mathbb{R}^2 \setminus \bar{B}(0, a))} \leq \sqrt{5} \|\nabla u\|_{L^2(\mathbb{R}^2 \setminus \bar{B}(0, a))}.$$

By a density argument, this holds for all $u \in W_0(\mathbb{R}^2 \setminus \bar{B}(0, a))$.

Step 2. Let now $u \in W_0(\mathbb{R}^2 \setminus \bar{\omega})$, and $a > 1$. Let $\theta \in \mathcal{C}_c^\infty(\mathbb{R}^2)$ such that $\theta = 1$ in $B(0, 2a)$ and $\theta = 0$ outside $B(0, 3a)$. By step 1 we have

$$\|(1 - \theta)u\|_{W(\mathbb{R}^2 \setminus \bar{\omega})} \leq c \|(1 - \theta)\nabla u - u\nabla\theta\|_{L^2(\mathbb{R}^2 \setminus \bar{\omega})} \leq c(\|\nabla u\|_{L^2(\mathbb{R}^2 \setminus \bar{\omega})} + \|u\|_{L^2(B(0, 3a) \setminus \bar{\omega})}).$$

The Poincaré inequality in $\{v \in H^1(B(0, 3a) \setminus \bar{\omega}) : v = 0 \text{ on } \partial\omega\}$ permits to conclude. \square

Lemma 5.2 *Let $\psi \in H^{1/2}(\partial\omega)$, $\psi_\varepsilon(x) = \psi((x - z)/\varepsilon)$ and $w_\varepsilon \in H^1(\Omega_\varepsilon)$ be the solution of*

$$\begin{cases} -\Delta w_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ w_\varepsilon = 0 & \text{on } \partial\Omega \\ w_\varepsilon = \psi_\varepsilon & \text{on } \partial\omega_\varepsilon. \end{cases}$$

Let $R > 0$ such that $B(z, R) \subset \Omega$. There exists a constant $c > 0$ independent of ε and ψ such that, for ε small enough,

$$\begin{aligned} \|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} &\leq c \|\psi\|_{H^{1/2}(\partial\omega)}, \\ \|w_\varepsilon\|_{H^1(\Omega \setminus B(z, R))} &\leq \frac{c}{\sqrt{-\log \varepsilon}} \|\psi\|_{H^{1/2}(\partial\omega)}. \end{aligned}$$

PROOF. We assume for convenience of notation that $z = 0$.

Step 1. We denote by $\Psi \in H^1(B(0, 1) \setminus \bar{\omega})$ a function such that $\Psi = \psi$ on $\partial\omega$ and $\Psi = 0$ on $\partial B(0, 1)$, obtained from standard lifting, then extended by 0 outside $B(0, 1)$. We set $\Psi_\varepsilon(x) = \Psi(x/\varepsilon)$ and $\tilde{w}_\varepsilon = w_\varepsilon - \Psi_\varepsilon$. We have from the weak formulation

$$\int_{\Omega_\varepsilon} \nabla w_\varepsilon \cdot \nabla \tilde{w}_\varepsilon dx = 0,$$

whereby

$$\|\nabla \tilde{w}_\varepsilon\|_{L^2(\Omega_\varepsilon)}^2 = - \int_{\Omega_\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla \tilde{w}_\varepsilon dx.$$

This entails $\|\nabla \tilde{w}_\varepsilon\|_{L^2(\Omega_\varepsilon)} \leq \|\nabla \Psi_\varepsilon\|_{L^2(\Omega_\varepsilon)}$ and subsequently $\|\nabla w_\varepsilon\|_{L^2(\Omega_\varepsilon)} \leq 2\|\nabla \Psi_\varepsilon\|_{L^2(\Omega_\varepsilon)}$. We infer by change of variables

$$\|\nabla w_\varepsilon\|_{L^2(\Omega_\varepsilon)} \leq 2\|\nabla \Psi\|_{L^2(B(0,1)\setminus\omega)} \leq c\|\psi\|_{H^{1/2}(\partial\omega)}.$$

We can also lift ψ inside ω by a function $\tilde{\psi}$, and setting $\tilde{\psi}_\varepsilon(x) = \tilde{\psi}(x/\varepsilon)$ we get

$$\|\nabla \tilde{\psi}_\varepsilon\|_{L^2(\omega_\varepsilon)} = \|\nabla \tilde{\psi}\|_{L^2(\omega)} \leq c\|\psi\|_{H^{1/2}(\partial\omega)}.$$

Extending w_ε by $\tilde{\psi}_\varepsilon$ in ω_ε and applying the Poincaré inequality in $H_0^1(\Omega)$ yields

$$\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq c\|\psi\|_{H^{1/2}(\partial\omega)}.$$

Step 2. To address the second claim we first focus on the problem

$$\begin{cases} -\Delta w_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ w_\varepsilon = 0 & \text{on } \partial\Omega \\ w_\varepsilon = \bar{\psi} & \text{on } \partial\omega_\varepsilon, \end{cases}$$

with $\bar{\psi} \in \mathbb{R}$ constant. The variational formulation yields

$$\|\nabla w_\varepsilon\|_{L^2(\Omega_\varepsilon)}^2 \leq \|\nabla v_\varepsilon\|_{L^2(\Omega_\varepsilon)}^2$$

for any $v_\varepsilon \in H^1(\Omega_\varepsilon)$ such that $v_\varepsilon = 0$ on $\partial\Omega$ and $v_\varepsilon = \bar{\psi}$ on $\partial\omega_\varepsilon$. We choose the following one, for some $\rho > 0$ such that $B(0, \rho) \subset \Omega$:

$$v_\varepsilon(x) = \begin{cases} \bar{\psi} & \text{if } |x| \leq \varepsilon \\ \bar{\psi} \frac{\log |x| - \log \rho}{\log \varepsilon - \log \rho} & \text{if } \varepsilon \leq |x| \leq \rho \\ 0 & \text{if } |x| \geq \rho. \end{cases}$$

This yields

$$\|\nabla v_\varepsilon\|_{L^2(\Omega_\varepsilon)}^2 = \int_\varepsilon^\rho \left(\frac{\bar{\psi}}{\log \varepsilon - \log \rho} \right)^2 \frac{1}{r^2} 2\pi r dr = 2\pi \frac{\bar{\psi}^2}{\log \rho - \log \varepsilon}.$$

It follows that

$$\|\nabla w_\varepsilon\|_{L^2(\Omega_\varepsilon)} \leq \left(\frac{2\pi}{\log \rho - \log \varepsilon} \right)^{1/2} |\bar{\psi}|.$$

The Poincaré inequality yields for ε small enough

$$\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq \frac{c}{\sqrt{-\log \varepsilon}} |\bar{\psi}|.$$

Step 3. We turn to the general case. By lifting, Proposition 5.1 and the Lax-Milgram theorem, there exists a unique $S \in W(\mathbb{R}^2 \setminus \bar{\omega})$ such that $S = \psi$ on $\partial\omega$ and

$$\int_{\mathbb{R}^2 \setminus \bar{\omega}} \nabla S \cdot \nabla \Phi dx = 0 \quad \forall \Phi \in W_0(\mathbb{R}^2 \setminus \bar{\omega}). \quad (23)$$

Obviously it holds $-\Delta S = 0$ in $\mathbb{R}^2 \setminus \bar{\omega}$ in the sense of distributions. Let ζ be a smooth function equal to 0 in $B(0, 1)$ and 1 outside $B(0, 2)$. Set $\hat{S} = \zeta S$ and

$$G = -\Delta \hat{S} = -\Delta \zeta S - 2\nabla \zeta \cdot \nabla S. \quad (24)$$

By construction G is supported in the ring $R(0, 1, 2)$, and it is smooth by elliptic regularity for S . Let now $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function equal to 1 in $B(0, 2)$ and 0 outside $B(0, 3)$ and set $\xi_\rho = \xi(x/\rho)$, $\rho > 1$. The Green formula yields

$$\int_{\mathbb{R}^2} G dx = \int_{\mathbb{R}^2} G \xi_\rho dx = \int_{\mathbb{R}^2} \nabla \hat{S} \cdot \nabla \xi_\rho dx = \int_{\mathbb{R}^2 \setminus \bar{B}(0, 2\rho)} \nabla \hat{S} \cdot \nabla \xi_\rho dx.$$

Applying the Cauchy-Schwarz inequality, using $\nabla \hat{S} \in L^2(\mathbb{R}^2)$, a change of variables, and letting ρ go to $+\infty$ results in

$$\int_{\mathbb{R}^2} G dx = 0. \quad (25)$$

We have for all $\Phi \in W(\mathbb{R}^2)$, using (24)

$$\begin{aligned} \int_{\mathbb{R}^2} G \Phi dx &= \int_{\mathbb{R}^2} \nabla \zeta \cdot \nabla (S \Phi) dx - 2 \int_{\mathbb{R}^2} \nabla \zeta \cdot \nabla S \Phi dx \\ &= \int_{\mathbb{R}^2} S \nabla \zeta \cdot \nabla \Phi dx - \int_{\mathbb{R}^2} \nabla \zeta \cdot \nabla S \Phi dx \\ &= \int_{\mathbb{R}^2} \nabla \hat{S} \cdot \nabla \Phi dx - \int_{\mathbb{R}^2} \zeta \nabla S \cdot \nabla \Phi dx - \int_{\mathbb{R}^2} \nabla \zeta \cdot \nabla S \Phi dx \\ &= \int_{\mathbb{R}^2} \nabla \hat{S} \cdot \nabla \Phi dx - \int_{\mathbb{R}^2} \nabla S \cdot \nabla (\zeta \Phi) dx. \end{aligned}$$

By (23) the latter integral vanishes, resulting in

$$\int_{\mathbb{R}^2} \nabla \hat{S} \cdot \nabla \Phi dx = \int_{\mathbb{R}^2} G \Phi dx \quad \forall \Phi \in W(\mathbb{R}^2). \quad (26)$$

Let

$$E(y) = \frac{-1}{2\pi} \log |y|$$

be the fundamental solution of the Laplacian and $\hat{S}_0 = G * E$. Since G is smooth and compactly supported and $E \in L^1_{\text{loc}}(\mathbb{R}^2)$, it follows that \hat{S}_0 is smooth. Using (25) we obtain the expressions

$$\hat{S}_0(x) = \int_{\mathbb{R}^2} G(y) (E(x-y) - E(x)) dy \quad \forall x \neq 0,$$

$$\nabla \hat{S}_0(x) = \int_{\mathbb{R}^2} G(y) (\nabla E(x-y) - \nabla E(x)) dy \quad \forall x \neq 0.$$

From the mean value theorem we infer that $|\hat{S}_0(x)| \leq c/|x|$ and $|\nabla \hat{S}_0(x)| \leq c/|x|^2$, implying that $\hat{S}_0 \in W(\mathbb{R}^2)$. Let $\Phi \in W(\mathbb{R}^2)$. We have

$$\begin{aligned} \int_{\mathbb{R}^2} \nabla \hat{S}_0 \cdot \nabla \Phi dx &= \lim_{\rho \rightarrow +\infty} \int_{\mathbb{R}^2} \xi_\rho \nabla \hat{S}_0 \cdot \nabla \Phi dx = \lim_{\rho \rightarrow +\infty} \left(\int_{\mathbb{R}^2} \nabla \hat{S}_0 \cdot \nabla (\xi_\rho \Phi) dx - \int_{\mathbb{R}^2} \nabla \hat{S}_0 \cdot \nabla \xi_\rho \Phi dx \right) \\ &= \int_{\mathbb{R}^2} G \Phi dx - \lim_{\rho \rightarrow +\infty} \int_{\mathbb{R}^2} \nabla \hat{S}_0 \cdot \nabla \xi_\rho \Phi dx = \int_{\mathbb{R}^2} G \Phi dx, \end{aligned}$$

using the decay properties of \hat{S}_0 and Φ . Comparing with (26), choosing $\Phi = \hat{S} - \hat{S}_0$, we obtain that $\hat{S} = \hat{S}_0 + \lambda$, for some $\lambda \in \mathbb{R}$. In particular we have the expression

$$\lambda = -\hat{S}_0(0),$$

showing that $|\lambda| \leq c\|\psi\|_{H^{1/2}(\partial\omega)}$. Denoting $S_0 = S - \lambda$ we have $S_0 = \hat{S} - \lambda = \hat{S}_0$ in $\mathbb{R}^2 \setminus B(0, 2)$, i.e.,

$$S_0(x) = \int_{B(0,2)} G(y) E(x-y) dy \quad \forall x \in \mathbb{R}^2 \setminus B(0, 2).$$

We now set

$$s_\varepsilon(x) = S_0\left(\frac{x}{\varepsilon}\right).$$

Using again (25) we get

$$s_\varepsilon(x) = \int_{B(0,2)} G(y) \left(E\left(\frac{x}{\varepsilon} - y\right) - E\left(\frac{x}{\varepsilon}\right) \right) dy \quad \forall x \in \mathbb{R}^2 \setminus B(0, 2\varepsilon).$$

The particular form of the fundamental solution leads to

$$s_\varepsilon(x) = \int_{B(0,2)} G(y) (E(x - \varepsilon y) - E(x)) dy \quad \forall x \in \mathbb{R}^2 \setminus B(0, 2\varepsilon).$$

The mean value theorem easily shows that

$$\|s_\varepsilon\|_{H^1(\Omega \setminus B(0,R))} \leq c\varepsilon \|G\|_{L^2(\mathbb{R}^2)} \leq c\varepsilon \|\psi\|_{H^{1/2}(\partial\omega)}.$$

We note that on $\partial\omega_\varepsilon$ we have $s_\varepsilon(x) = S_0(x/\varepsilon) = S(x/\varepsilon) - \lambda = \psi(x/\varepsilon) - \lambda$. We now define $r_\varepsilon = w_\varepsilon - s_\varepsilon$, solution of

$$\begin{cases} -\Delta r_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ r_\varepsilon = -s_\varepsilon & \text{on } \partial\Omega \\ r_\varepsilon = \lambda & \text{on } \partial\omega_\varepsilon. \end{cases}$$

We have by step 2 and a standard decomposition

$$\|r_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq c\|s_\varepsilon\|_{H^{1/2}(\partial\Omega)} + c \frac{|\lambda|}{\sqrt{-\log \varepsilon}} \leq \frac{c}{\sqrt{-\log \varepsilon}} \|\psi\|_{H^{1/2}(\partial\omega)}.$$

This completes the proof by $w_\varepsilon = s_\varepsilon + r_\varepsilon$. □

5.3 Variation of the direct state

Set $\tilde{u}_\varepsilon = u_\varepsilon - u_0$. It solves

$$\begin{cases} -\Delta \tilde{u}_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ \tilde{u}_\varepsilon = 0 & \text{on } \partial\Omega \\ \tilde{u}_\varepsilon = -u_0 & \text{on } \partial\omega_\varepsilon. \end{cases} \quad (27)$$

We define $h_\varepsilon \in C^\infty(\mathbb{R}^2 \setminus \{z\})$ and $r_\varepsilon \in H^1(\Omega)$ by

$$h_\varepsilon(x) = -\frac{\log|x-z|}{\log \varepsilon} u_0(z), \quad \begin{cases} -\Delta r_\varepsilon = 0 & \text{in } \Omega \\ r_\varepsilon = -h_\varepsilon & \text{on } \partial\Omega. \end{cases}$$

As for the fundamental solution we have $\Delta h_\varepsilon = 0$ in $\mathbb{R}^2 \setminus \{z\}$. We now set $e_\varepsilon = \tilde{u}_\varepsilon - h_\varepsilon - r_\varepsilon$, which solves

$$\begin{cases} -\Delta e_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ e_\varepsilon = 0 & \text{on } \partial\Omega \\ e_\varepsilon = -u_0 - h_\varepsilon - r_\varepsilon & \text{on } \partial\omega_\varepsilon. \end{cases} \quad (28)$$

Lemma 5.3 *If u_0 is C^1 in a neighborhood of z then*

$$\begin{aligned} \|r_\varepsilon\|_{H^1(\Omega)} &= O((- \log \varepsilon)^{-1}), \\ \|e_\varepsilon\|_{H^1(\Omega \setminus B(z, R))} &= O((- \log \varepsilon)^{-3/2}). \end{aligned}$$

PROOF. The first estimate is obvious since $\|h_\varepsilon\|_{H^{1/2}(\partial\Omega)} = O((- \log \varepsilon)^{-1})$ by construction.

Set

$$\psi_\varepsilon(x) = (-u_0 - h_\varepsilon - r_\varepsilon)(z + \varepsilon x).$$

We decompose as

$$\psi_\varepsilon(x) = \underbrace{[u_0(z) - u_0(z + \varepsilon x)]}_{p_\varepsilon(x)} - \underbrace{[u_0(z) + h_\varepsilon(z + \varepsilon x)]}_{q_\varepsilon(x)} - \underbrace{r_\varepsilon(z + \varepsilon x)}_{\hat{r}_\varepsilon(x)}.$$

By regularity of u_0 we have immediately $\|p_\varepsilon\|_{H^{1/2}(\partial\omega)} = O(\varepsilon)$. Next, from

$$q_\varepsilon(x) = u_0(z) \left(1 - \frac{\log |\varepsilon x|}{\log \varepsilon} \right) = -u_0(z) \frac{\log |x|}{\log \varepsilon},$$

we get $\|q_\varepsilon\|_{H^{1/2}(\partial\omega)} = O((- \log \varepsilon)^{-1})$. Lastly, a change of variables yields

$$\|\hat{r}_\varepsilon\|_{H^1(\omega)} \leq \|\nabla r_\varepsilon\|_{L^2(\omega_\varepsilon)} + \varepsilon^{-1} \|r_\varepsilon\|_{L^2(\omega_\varepsilon)} \leq \|\nabla r_\varepsilon\|_{L^2(\omega_\varepsilon)} + c \|r_\varepsilon\|_{L^\infty(\omega_\varepsilon)}.$$

By elliptic regularity we have $\|\nabla r_\varepsilon\|_{L^2(\omega_\varepsilon)} + \|r_\varepsilon\|_{L^\infty(\omega_\varepsilon)} \leq c \|h_\varepsilon\|_{H^{1/2}(\partial\Omega)}$, thus $\|\hat{r}_\varepsilon\|_{H^{1/2}(\partial\omega)} = O((- \log \varepsilon)^{-1})$. We conclude using Lemma 5.2. \square

We infer from Lemma 5.3 and the triangle inequality:

Lemma 5.4 *If u_0 is C^1 in a neighborhood of z then*

$$\|\tilde{u}_\varepsilon\|_{H^1(\Omega \setminus B(z, R))} = O((- \log \varepsilon)^{-1}).$$

5.4 Variation of the adjoint state

In view of Proposition 3.1 and (20)-(21), we define the adjoint state $v_\varepsilon \in H_0^1(\Omega)$ solution of

$$\int_{\Omega} \nabla v_\varepsilon \cdot \nabla \varphi dx = - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0) \varphi dt \quad \forall \varphi \in H_0^1(\Omega).$$

In particular the unperturbed adjoint state v_0 satisfies

$$\int_{\Omega} \nabla v_0 \cdot \nabla \varphi dx = -dJ_0(u_0) \varphi \quad \forall \varphi \in H_0^1(\Omega). \quad (29)$$

Lemma 5.5 *If u_0 is C^1 in a neighborhood of z then*

$$\|v_\varepsilon - v_0\|_{H^1(\Omega)} = O((- \log \varepsilon)^{-1}).$$

PROOF. Set $\tilde{v}_\varepsilon = v_\varepsilon - v_0$. We have

$$\int_{\Omega} \nabla \tilde{v}_\varepsilon \cdot \nabla \varphi dx = \int_0^1 (dJ_0(u_0) - dJ_\varepsilon(tu_\varepsilon + (1-t)u_0)) \varphi dt \quad \forall \varphi \in H_0^1(\Omega),$$

leading to

$$\int_{\Omega} \nabla \tilde{v}_\varepsilon \cdot \nabla \varphi dx = \int_0^1 (d\hat{J}(u_0|_{\hat{\Omega}}) - d\hat{J}((tu_\varepsilon + (1-t)u_0)|_{\hat{\Omega}})) \varphi|_{\hat{\Omega}} dt \quad \forall \varphi \in H_0^1(\Omega).$$

Choosing $\varphi = \tilde{v}_\varepsilon$ and using that $d\hat{J}$ is Lipschitz yields

$$\|\tilde{v}_\varepsilon\|_{H^1(\Omega)} \leq c \|\tilde{u}_\varepsilon\|_{H^1(\hat{\Omega})}.$$

The conclusion follows from Lemma 5.4. \square

5.5 Variation of the Lagrangian

In view of Proposition 3.1 we define the Lagrangian

$$\mathcal{L}_\varepsilon(u, v) = J_\varepsilon(u) + \int_{\Omega} \nabla u \cdot \nabla v dx - \int_{\Omega} f v + \int_{\partial\omega_\varepsilon} \frac{\partial u_\varepsilon}{\partial n} v ds. \quad (30)$$

This provides the variation

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \int_{\partial\omega_\varepsilon} \frac{\partial u_\varepsilon}{\partial n} v_\varepsilon ds.$$

Lemma 5.6 *If u_0, v_0 are of class \mathcal{C}^1 in a neighborhood of z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} v_0 ds + O((- \log \varepsilon)^{-2}).$$

PROOF. We work with the decomposition

$$\begin{aligned} (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) - \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} v_0 ds &= \int_{\partial\omega_\varepsilon} \frac{\partial u_0}{\partial n} v_\varepsilon ds + \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} (v_\varepsilon - v_0) ds \\ &= \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_\varepsilon dx - \int_{\Omega_\varepsilon} \nabla \tilde{u}_\varepsilon \cdot \nabla (v_\varepsilon - v_0) dx. \end{aligned}$$

Extending \tilde{u}_ε by $-u_0$ in ω_ε allows to write

$$\begin{aligned} (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) - \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} v_0 ds &= - \int_{\Omega} \nabla \tilde{u}_\varepsilon \cdot \nabla (v_\varepsilon - v_0) dx + \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx \\ &= \int_0^1 (d\hat{J}((tu_\varepsilon + (1-t)u_0)|_{\hat{\Omega}}) - d\hat{J}(u_0|_{\hat{\Omega}}))(\tilde{u}_\varepsilon)|_{\hat{\Omega}} dt \\ &\quad + \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx, \end{aligned}$$

where the last equality is obtained as in Lemma 5.5. We conclude using Lemma 5.4. \square

Lemma 5.7 *If u_0, v_0 are of class \mathcal{C}^1 in a neighborhood of z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \frac{2\pi}{-\log \varepsilon} u_0(z) v_0(z) + O((- \log \varepsilon)^{-3/2}).$$

PROOF. We decompose the expression found in Lemma 5.6 as

$$\begin{aligned} (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0 ds + \int_{\partial\omega_\varepsilon} \frac{\partial r_\varepsilon}{\partial n} v_0 ds + \int_{\partial\omega_\varepsilon} \frac{\partial e_\varepsilon}{\partial n} v_0 ds + O((- \log \varepsilon)^{-2}) \\ &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds \\ &\quad + \int_{\omega_\varepsilon} \nabla r_\varepsilon \cdot \nabla v_0 dx - \int_{\Omega_\varepsilon} \nabla e_\varepsilon \cdot \nabla v_0 dx + O((- \log \varepsilon)^{-2}). \end{aligned}$$

Let $\rho > 0$ such that $B(0, \rho) \subset \omega$. We define

$$\tilde{h}_\varepsilon(x) = \begin{cases} h_\varepsilon(x) & \text{if } |x - z| \geq \rho\varepsilon \\ -\left(\frac{\log \rho}{\log \varepsilon} + 1\right) u_0(z) & \text{if } |x - z| \leq \rho\varepsilon. \end{cases}$$

This truncation ensures that $\tilde{h}_\varepsilon \in H^1(\Omega)$. We extend e_ε by $-u_0 - \tilde{h}_\varepsilon - r_\varepsilon$ in ω_ε to write

$$\begin{aligned} (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds \\ &\quad - \int_{\Omega} \nabla e_\varepsilon \cdot \nabla v_0 dx - \int_{\omega_\varepsilon} \nabla(\tilde{h}_\varepsilon + u_0) \cdot \nabla v_0 dx + O((- \log \varepsilon)^{-2}) \\ &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds \\ &\quad + d\hat{J}(u_0|_{\hat{\Omega}}) e_\varepsilon|_{\hat{\Omega}} - \int_{\omega_\varepsilon} \nabla(\tilde{h}_\varepsilon + u_0) \cdot \nabla v_0 dx + O((- \log \varepsilon)^{-2}). \end{aligned}$$

By the definition of h_ε , Lemma 5.3 and the smoothness assumptions we arrive at

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + O((- \log \varepsilon)^{-3/2}).$$

The Green formula yields

$$\int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} ds = \int_{\partial B(z, \varepsilon)} \frac{\partial h_\varepsilon}{\partial n} ds = \frac{-2\pi}{\log \varepsilon} u_0(z),$$

which completes the proof. \square

5.6 Expression of the topological asymptotic expansion

Based on the previous results we arrive at the following counterpart to Theorem 4.9.

Theorem 5.8 *Consider a cost function of form (22). Let $v_0 \in H_0^1(\Omega)$ be the solution of (29). Then*

$$J_\varepsilon(u_\varepsilon) - J_0(u_0) = \frac{2\pi}{-\log \varepsilon} u_0(z) v_0(z) + o\left(\frac{1}{-\log \varepsilon}\right).$$

We observe that this expression does not depend on the shape of the hole. This is typical of the 2D case. We infer the topological derivative

$$d_T \mathcal{J}(\Omega, \omega, z) = u_0(z) v_0(z).$$

Lastly, we highlight in Fig. 2 the slow convergence of the cost function compared with the inclusion case.

6 Dirichlet case in 3D

We consider the same problem as in section 5.1, but now in dimension $N = 3$. We follow the same general scheme, but significant technical differences will show up.

6.1 Preliminary estimate

The weighted Sobolev spaces $W(\mathbb{R}^3)$ and $W_0(\mathbb{R}^3)$ are defined in the same way as their 2D counterparts, with the weight function

$$w(x) = \frac{1}{\sqrt{1 + |x|^2}}.$$

It leads to the same Poincaré inequality [3, 23]:

$$\|u\|_{W(\mathbb{R}^3 \setminus \bar{\omega})} \leq c \|\nabla u\|_{L^2(\mathbb{R}^3 \setminus \bar{\omega})} \quad \forall u \in W_0(\mathbb{R}^3 \setminus \bar{\omega}). \quad (31)$$




Figure 2: Comparison of the functions $f_1(\varepsilon) = \frac{1}{-\log \varepsilon}$ (in blue) and $f_2(\varepsilon) = \varepsilon^2$ (in red).

Lemma 6.1 *Let $\psi \in H^{1/2}(\partial\omega)$, $\psi_\varepsilon(x) = \psi((x - z)/\varepsilon)$ and $w_\varepsilon \in H^1(\Omega_\varepsilon)$ be the solution of*

$$\begin{cases} -\Delta w_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ w_\varepsilon = 0 & \text{on } \partial\Omega \\ w_\varepsilon = \psi_\varepsilon & \text{on } \partial\omega_\varepsilon. \end{cases}$$

Let $R > 0$ such that $B(z, R) \subset \Omega$. There exists a constant $c > 0$ independent of ε and ψ such that, for ε small enough,

$$\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq c\sqrt{\varepsilon}\|\psi\|_{H^{1/2}(\partial\omega)}, \quad (32)$$

$$\|w_\varepsilon\|_{H^1(\Omega \setminus B(z, R))} \leq c\varepsilon\|\psi\|_{H^{1/2}(\partial\omega)}. \quad (33)$$

PROOF. We assume here again for simplicity that $z = 0$. The proof of (32) follows exactly the same lines as in Lemma 5.2, hence we focus on (33). By lifting, (31) and the Lax-Milgram theorem, there exists a unique $S \in W(\mathbb{R}^3 \setminus \bar{\omega})$ such that $S = \psi$ on $\partial\omega$ and

$$\int_{\mathbb{R}^3 \setminus \bar{\omega}} \nabla S \cdot \nabla \Phi dx = 0 \quad \forall \Phi \in W_0(\mathbb{R}^3 \setminus \bar{\omega}). \quad (34)$$

Similarly to Lemma 5.2 we introduce a smooth function ζ equal to 0 in $B(0, 1)$ and 1 outside $B(0, 2)$, we set $\hat{S} = \zeta S$ and $\hat{S}_0 = G * E$, with $G = -\Delta \hat{S}$,

$$E(y) = \frac{1}{4\pi|y|}$$

the fundamental solution of the Laplacian. Here we do not have the counterpart to (25), nevertheless the definition directly shows that $\hat{S}_0 \in W(\mathbb{R}^3)$. It follows by the same argument as in Lemma 5.2 that $\hat{S} = \hat{S}_0 + \lambda$, for some $\lambda \in \mathbb{R}$, but here the inclusion $\hat{S} - \hat{S}_0 \in W(\mathbb{R}^3)$ implies $\lambda = 0$ since constants do not belong to $W(\mathbb{R}^3)$. We arrive at

$$S(x) = \int_{B(0, 2)} G(y) E(x - y) dy \quad \forall x \in \mathbb{R}^3 \setminus B(0, 2). \quad (35)$$

Setting

$$s_\varepsilon(x) = S\left(\frac{x}{\varepsilon}\right),$$

we get

$$s_\varepsilon(x) = \varepsilon \int_{B(0, 2)} G(y) E(x - \varepsilon y) dy \quad \forall x \in \mathbb{R}^3 \setminus B(0, 2\varepsilon).$$

This representation entails

$$\|s_\varepsilon\|_{H^1(\Omega \setminus B(0,R))} \leq c\varepsilon \|G\|_{L^2(\mathbb{R}^3)} \leq c\varepsilon \|\psi\|_{H^{1/2}(\partial\omega)}. \quad (36)$$

We now define $r_\varepsilon = w_\varepsilon - s_\varepsilon$, which is solution of

$$\begin{cases} -\Delta r_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ r_\varepsilon = -s_\varepsilon & \text{on } \partial\Omega \\ r_\varepsilon = 0 & \text{on } \partial\omega_\varepsilon. \end{cases}$$

The decomposition $w_\varepsilon = s_\varepsilon + r_\varepsilon$ combined with (36) yields (33). \square

6.2 Variation of the direct state

Similarly to section 5.3 we set $\tilde{u}_\varepsilon = u_\varepsilon - u_0$, which solves

$$\begin{cases} -\Delta \tilde{u}_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ \tilde{u}_\varepsilon = 0 & \text{on } \partial\Omega \\ \tilde{u}_\varepsilon = -u_0 & \text{on } \partial\omega_\varepsilon. \end{cases} \quad (37)$$

Now we consider the reference problem: find $U \in W(\mathbb{R}^3 \setminus \bar{\omega})$ such that

$$\begin{cases} -\Delta U = 0 & \text{in } \mathbb{R}^3 \setminus \bar{\omega} \\ U = 1 & \text{on } \partial\omega, \end{cases} \quad (38)$$

whose variational formulation is as in (34). We define $h_\varepsilon \in W(\mathbb{R}^3 \setminus \bar{\omega}_\varepsilon)$ and $r_\varepsilon \in H^1(\Omega)$ by

$$h_\varepsilon(x) = -u_0(z)U\left(\frac{x-z}{\varepsilon}\right), \quad \begin{cases} -\Delta r_\varepsilon = 0 & \text{in } \Omega \\ r_\varepsilon = -h_\varepsilon & \text{on } \partial\Omega. \end{cases}$$

We now set $e_\varepsilon = \tilde{u}_\varepsilon - h_\varepsilon - r_\varepsilon$. It solves

$$\begin{cases} -\Delta e_\varepsilon = 0 & \text{in } \Omega_\varepsilon \\ e_\varepsilon = 0 & \text{on } \partial\Omega \\ e_\varepsilon = -u_0 - h_\varepsilon - r_\varepsilon & \text{on } \partial\omega_\varepsilon. \end{cases} \quad (39)$$

Lemma 6.2 *If u_0 is \mathcal{C}^1 is a neighborhood of z then*

$$\|h_\varepsilon\|_{H^1(\Omega_\varepsilon \setminus B(z,R))} = O(\varepsilon),$$

$$\|r_\varepsilon\|_{H^1(\Omega)} = O(\varepsilon),$$

$$\|e_\varepsilon\|_{H^1(\Omega \setminus B(z,R))} = O(\varepsilon^{3/2}).$$

PROOF. The same representation as in (35) shows that $|U(x)| \leq c/|x|$ and $|\nabla U(x)| \leq c/|x|^2$, hence $\|h_\varepsilon\|_{H^1(\Omega \setminus B(z,R))} = O(\varepsilon)$, as in (36) whereby also $\|r_\varepsilon\|_{H^1(\Omega)} = O(\varepsilon)$. Set $\hat{r}_\varepsilon(x) = r_\varepsilon(z + \varepsilon x)$. A change of variables yields

$$\|\hat{r}_\varepsilon\|_{H^1(\omega)} \leq \varepsilon^{-1/2} \|\nabla r_\varepsilon\|_{L^2(\omega_\varepsilon)} + \varepsilon^{-3/2} \|r_\varepsilon\|_{L^2(\omega_\varepsilon)} \leq \varepsilon^{-1/2} \|\nabla r_\varepsilon\|_{L^2(\omega_\varepsilon)} + c\varepsilon^{-1/2} \|r_\varepsilon\|_{L^6(\Omega)} = O(\varepsilon^{1/2}),$$

where we have used the Sobolev embedding $H^1 \hookrightarrow L^6$. We obtain $\|(u_0 + h_\varepsilon + r_\varepsilon)(z + \varepsilon x)\|_{H^{1/2}(\partial\omega)} = \|(u_0(z + \varepsilon x) - u_0(z) + \hat{r}_\varepsilon(x))\|_{H^{1/2}(\partial\omega)} = O(\varepsilon^{1/2})$. We conclude by Lemma 6.1. \square

We deduce:

Lemma 6.3 *If u_0 is \mathcal{C}^1 is a neighborhood of z then*

$$\|\tilde{u}_\varepsilon\|_{H^1(\Omega \setminus B(z,R))} = O(\varepsilon).$$

6.3 Variation of the adjoint state

We again define the adjoint state $v_\varepsilon \in H_0^1(\Omega)$ solution of

$$\int_{\Omega} \nabla v_\varepsilon \cdot \nabla \varphi dx = - \int_0^1 dJ_\varepsilon(tu_\varepsilon + (1-t)u_0) \varphi dt \quad \forall \varphi \in H_0^1(\Omega).$$

Similarly to Lemma 5.5 we have:

Lemma 6.4 *If u_0 is C^1 in a neighborhood of z then*

$$\|v_\varepsilon - v_0\|_{H^1(\Omega)} = O(\varepsilon).$$

6.4 Variation of the Lagrangian

We consider the same Lagrangian as in (30).

Lemma 6.5 *If u_0, v_0 are of class C^1 in a neighborhood of z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} v_0 ds + O(\varepsilon^2).$$

PROOF. Like in Lemma 5.6 we have

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) - \int_{\partial\omega_\varepsilon} \frac{\partial \tilde{u}_\varepsilon}{\partial n} v_0 ds = \int_0^1 (d\hat{J}((tu_\varepsilon + (1-t)u_0)|_{\hat{\Omega}}) - d\hat{J}(u_0|_{\hat{\Omega}}))(\tilde{u}_\varepsilon)|_{\hat{\Omega}} dt + \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx.$$

We conclude using Lemma 6.3. □

The standard capacity of ω can be defined by (see e.g. [33])

$$\mathcal{K} = - \int_{\partial\omega} \frac{\partial U}{\partial n} ds = \int_{\mathbb{R}^3 \setminus \omega} |\nabla U|^2 dx. \quad (40)$$

Lemma 6.6 *If u_0, v_0 are of class C^1 in a neighborhood of z then*

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \mathcal{K}\varepsilon u_0(z) v_0(z) + O(\varepsilon^{3/2}).$$

PROOF. We decompose the expression found in Lemma 6.5 as

$$(\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) = \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds + \int_{\omega_\varepsilon} \nabla r_\varepsilon \cdot \nabla v_0 dx - \int_{\Omega_\varepsilon} \nabla e_\varepsilon \cdot \nabla v_0 dx + O(\varepsilon^2).$$

Inside ω_ε we extend h_ε by $-u_0(z)$ and e_ε by $-u_0 + u_0(z) - r_\varepsilon$. This permits to write

$$\begin{aligned} (\mathcal{L}_\varepsilon - \mathcal{L}_0)(u_0, v_\varepsilon) &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds \\ &\quad - \int_{\Omega} \nabla e_\varepsilon \cdot \nabla v_0 dx - \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx + O(\varepsilon^2) \\ &= \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} v_0(z) ds + \int_{\partial\omega_\varepsilon} \frac{\partial h_\varepsilon}{\partial n} (v_0 - v_0(z)) ds \\ &\quad + d\hat{J}(u_0|_{\hat{\Omega}}) e_\varepsilon|_{\hat{\Omega}} - \int_{\omega_\varepsilon} \nabla u_0 \cdot \nabla v_0 dx + O(\varepsilon^2). \end{aligned}$$

The claim follows from the definition of h_ε and Lemma 6.2. □

6.5 Expression of the topological asymptotic expansion

From the preceding findings we infer the 3D counterpart to Theorem 5.8.

Theorem 6.7 *Consider a cost function of form (22). Let $v_0 \in H_0^1(\Omega)$ be the solution of (29), \mathcal{K} be defined by (40). Then*

$$J_\varepsilon(u_\varepsilon) - J_0(u_0) = \mathcal{K}\varepsilon u_0(z)v_0(z) + o(\varepsilon).$$

We note that here, unlike in the 2D case, the expression of the topological asymptotic expansion depends on the shape of the hole, through the capacity \mathcal{K} . In the case of the ball $\omega = B(0, 1)$ the solution to (38) is found as

$$U(x) = \frac{1}{|x|},$$

leading to the capacity $\mathcal{K} = 4\pi$.

Remark 6.8 *Like in the inclusion case, more general cost functions may yield additional terms, see e.g. [29, 6].*

6.6 Extensions

Topological derivatives in the Dirichlet case has been comparatively less studied than in the inclusion / Neumann cases. Related contributions can be found in particular on the linear elasticity case [28], the Helmholtz equation [46], the Stokes problem [30], and semilinear problems [6, 35].

Acknowledgements

The author benefited from the supports of the chair "Modeling advanced polymers for innovative material solutions" led by the Ecole polytechnique and the Fondation de l'Ecole polytechnique and sponsored by Arkema, and of the project ANR-18-CE40-0013 SHAPO financed by the French Agence Nationale de la Recherche.

References

- [1] G. Allaire. *Conception optimale de structures*, volume 58 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Berlin, 2007. With the collaboration of Marc Schoenauer (INRIA) in the writing of Chapter 8.
- [2] H. Ammari and H. Kang. *Polarization and moment tensors*, volume 162 of *Applied Mathematical Sciences*. Springer, New York, 2007.
- [3] C. Amrouche, V. Girault, and J. Giroire. Weighted Sobolev spaces for Laplace's equation in \mathbf{R}^n . *J. Math. Pures Appl. (9)*, 73(6):579–606, 1994.
- [4] S. Amstutz. *Aspects théoriques et numériques en optimisation de forme topologique*. PhD thesis, Institut National des Sciences Appliquées, Toulouse, France, 2003.
- [5] S. Amstutz. Sensitivity analysis with respect to a local perturbation of the material property. *Asymptot. Anal.*, 49(1-2):87–108, 2006.
- [6] S. Amstutz. Topological sensitivity analysis for some nonlinear PDE system. *J. Math. Pures Appl. (9)*, 85(4):540–557, 2006.

- [7] S. Amstutz and A. Bonnafé. Topological derivatives for a class of quasilinear elliptic equations. *J. Math. Pures Appl. (9)*, 107(4):367–408, 2017.
- [8] S. Amstutz and N. Dominguez. Topological sensitivity in the context of ultrasonic nondestructive testing. *Engineering Analysis with Boundary Elements*, 32:936–947, 2008.
- [9] S. Amstutz, I. Horchani, and M. Masmoudi. Crack detection by the topological gradient method. *Control Cybernet.*, 34(1):81–101, 2005.
- [10] S. Amstutz, A. A. Novotny, and E. A. de Souza Neto. Topological derivative-based topology optimization of structures subject to Drucker-Prager stress constraints. *Comput. Methods Appl. Mech. Engrg.*, 233/236:123–136, 2012.
- [11] S. Amstutz, A. A. Novotny, and N. Van Goethem. Topological sensitivity analysis for elliptic differential operators of order $2m$. *J. Differential Equations*, 256(4):1735–1770, 2014.
- [12] S. Amstutz, T. Takahashi, and B. Vexler. Topological sensitivity analysis for time-dependent problems. *ESAIM Control Optim. Calc. Var.*, 14(3):427–455, 2008.
- [13] H. Attouch, G. Buttazzo, and G. Michaille. *Variational analysis in Sobolev and BV spaces*, volume 17 of *MOS-SIAM Series on Optimization*. Second edition, 2014.
- [14] G. Aubert and A. Drogoul. Topological gradient for a fourth order operator used in image analysis. *ESAIM Control Optim. Calc. Var.*, 21(4):1120–1149, 2015.
- [15] C. Bellis and M. Bonnet. Qualitative identification of cracks using 3D transient elastodynamic topological derivative: formulation and FE implementation. *Comput. Methods Appl. Mech. Engrg.*, 253:89–105, 2013.
- [16] A. Ben Abda, M. Hassine, M. Jaoua, and M. Masmoudi. Topological sensitivity analysis for the location of small cavities in Stokes flow. *SIAM J. Control Optim.*, 48(5):2871–2900, 2009/10.
- [17] M. Bonnet. Topological sensitivity for 3D elastodynamic and acoustic inverse scattering in the time domain. *Comput. Methods Appl. Mech. Engrg.*, 195(37-40):5239–5254, 2006.
- [18] M. Bonnet. Higher-order topological sensitivity for 2-D potential problems. Application to fast identification of inclusions. *Internat. J. Solids Structures*, 46(11-12):2275–2292, 2009.
- [19] M. Bonnet. Fast identification of cracks using higher-order topological sensitivity for 2-D potential problems. *Eng. Anal. Bound. Elem.*, 35(2):223–235, 2011.
- [20] M. Bonnet and R. Cornaggia. Higher order topological derivatives for three-dimensional anisotropic elasticity. *ESAIM Math. Model. Numer. Anal.*, 51(6):2069–2092, 2017.
- [21] M. Bonnet and G. Delgado. The topological derivative in anisotropic elasticity. *Quart. J. Mech. Appl. Math.*, 66(4):557–586, 2013.
- [22] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [23] R. Dautray and J.-L. Lions. *Analyse mathématique et calcul numérique pour les sciences et les techniques. Vol. 4*. Masson, Paris, 1988.

- [24] F. Demengel and G. Demengel. *Functional spaces for the theory of elliptic partial differential equations*. Universitext. Springer, London; EDP Sciences, Les Ulis, 2012. Translated from the 2007 French original by Reinie Ern .
- [25] J. Deny and J. L. Lions. Les espaces du type de Beppo Levi. *Ann. Inst. Fourier (Grenoble)*, 5:305–370, 1955.
- [26] H. Eschenauer, V. Kobelev, and A. Schumacher. Bubble method for topology and shape optimization of structures. *Structural optimization*, 8:42–51, 1994.
- [27] P. Gangl and K. Sturm. A simplified derivation technique of topological derivatives for quasi-linear transmission problems. *ESAIM Control Optim. Calc. Var.*, 26:Paper No. 106, 20, 2020.
- [28] S. Garreau, P. Guillaume, and M. Masmoudi. The topological asymptotic for PDE systems: the elasticity case. *SIAM J. Control Optim.*, 39(6):1756–1778, 2001.
- [29] P. Guillaume and K. Sid Idris. The topological asymptotic expansion for the Dirichlet problem. *SIAM J. Control Optim.*, 41(4):1042–1072, 2002.
- [30] P. Guillaume and K. Sid Idris. Topological sensitivity and shape optimization for the Stokes equations. *SIAM J. Control Optim.*, 43(1):1–31, 2004.
- [31] B. B. Guzina and M. Bonnet. Small-inclusion asymptotic of misfit functionals for inverse problems in acoustics. *Inverse Problems*, 22(5):1761–1785, 2006.
- [32] M. Hassine, S. Jan, and M. Masmoudi. From differential calculus to 0-1 topological optimization. *SIAM J. Control Optim.*, 45(6):1965–1987, 2007.
- [33] A. Henrot and M. Pierre. *Shape variation and optimization*, volume 28 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Z rich, 2018.
- [34] M. Hinterm ller, A. Laurain, and A. A. Novotny. Second-order topological expansion for electrical impedance tomography. *Adv. Comput. Math.*, 36(2):235–265, 2012.
- [35] M. Iguernane, S. A. Nazarov, J.-R. Roche, J. Sokolowski, and K. Szulc. Topological derivatives for semilinear elliptic equations. *Int. J. Appl. Math. Comput. Sci.*, 19(2):191–205, 2009.
- [36] M. Masmoudi. The topological asymptotic. In R. Glowinski, H. Karawada, and J. Periaux, editors, *Computational Methods for Control Applications*, pages 53–72. GAKUTO International Series, 2002.
- [37] M. Masmoudi, J. Pommier, and B. Samet. The topological asymptotic expansion for the Maxwell equations and some applications. *Inverse Problems*, 21(2):547–564, 2005.
- [38] F. Murat and J. Simon. Sur le contr le par un domaine g om trique. Technical Report 76 015, L. d’Analyse Num rique Univ. Paris 6, 1976.
- [39] A.-A. Novotny. *An lise de sensibilidade topol gica*. PhD thesis, Laborat rio Nacional de Computa  o Cient fica, Petr polis, Brazil, 2003.
- [40] A. A. Novotny and J. Sokolowski. *Topological derivatives in shape optimization*. Interaction of Mechanics and Mathematics. Springer, Heidelberg, 2013.
- [41] A. A. Novotny and J. Sokolowski. *An introduction to the topological derivative method*. Springer, 2020.

- [42] A. A. Novotny, J. Sokolowski, and A. Żochowski. Topological derivatives of shape functionals. Part I: Theory in singularly perturbed geometrical domains. *J. Optim. Theory Appl.*, 180(2):341–373, 2019.
- [43] A. A. Novotny, J. Sokolowski, and A. Żochowski. Topological derivatives of shape functionals. Part III: Second-order method and applications. *J. Optim. Theory Appl.*, 181(1):1–22, 2019.
- [44] C. Ortner and E. Süli. A note on linear elliptic systems on \mathbb{R}^d . Technical Report 1202:3970, ArXiv, 2012.
- [45] G. Pólya and G. Szegő. *Isoperimetric Inequalities in Mathematical Physics*. Annals of Mathematics Studies, No. 27. Princeton University Press, Princeton, N. J., 1951.
- [46] B. Samet, S. Amstutz, and M. Masmoudi. The topological asymptotic for the Helmholtz equation. *SIAM J. Control Optim.*, 42(5):1523–1544, 2003.
- [47] M. Schiffer and G. Szegő. Virtual mass and polarization. *Trans. Amer. Math. Soc.*, 67:130–205, 1949.
- [48] A. Schumacher. *Topologieoptimisierung von Bauteilstrukturen unter Verwendung von Topchpositionierungskriterien*. PhD thesis, Universität-Gesamthochschule-Siegen, 1995.
- [49] J. Sokolowski and A. Żochowski. On the topological derivative in shape optimization. *SIAM J. Control Optim.*, 37(4):1251–1272, 1999.
- [50] J. Sokolowski and J.-P. Zolésio. *Introduction to shape optimization*, volume 16 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 1992. Shape sensitivity analysis.
- [51] K. Sturm. Topological sensitivities via a Lagrangian approach for semilinear problems. *Nonlinearity*, 33(9):4310–4337, 2020.