



HAL
open science

Fundamental questions and applications of sclerochronology: Community-defined research priorities

Tamara Trofimova, Stella J. Alexandroff, Madelyn J. Mette, Elizabeth Tray, Paul G. Butler, Steven E. Campana, Elizabeth M. Harper, Andrew L. A. Johnson, John R. Morrongiello, Melita Peharda, et al.

► To cite this version:

Tamara Trofimova, Stella J. Alexandroff, Madelyn J. Mette, Elizabeth Tray, Paul G. Butler, et al.. Fundamental questions and applications of sclerochronology: Community-defined research priorities. *Estuarine, Coastal and Shelf Science*, 2020, 245, pp.106977. 10.1016/j.ecss.2020.106977. hal-03311943

HAL Id: hal-03311943

<https://hal.science/hal-03311943>

Submitted on 3 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Fundamental questions and applications of sclerochronology: Community-defined research priorities

Tamara Trofimova^{a,*}, Stella J. Alexandroff^{b,c}, Madelyn J. Mette^a, Elizabeth Tray^d, Paul G. Butler^c, Steven E. Campana^e, Elizabeth M. Harper^f, Andrew L.A. Johnson^g, John R. Morrongiello^h, Melita Pehardaⁱ, Bernd R. Schöne^j, Carin Andersson^a, C. Fred T. Andrus^k, Bryan A. Black^l, Meghan Burchell^m, Michael L. Carrollⁿ, Kristine L. DeLong^o, Bronwyn M. Gillanders^p, Peter Grønkjær^q, Daniel Killam^r, Amy L. Prendergast^s, David J. Reynolds^l, James D. Scourse^c, Kotaro Shirai^t, Julien Thébault^u, Clive Trueman^v, Niels de Winter^{w,x}

^a NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Norway

^b School of Ocean Sciences, Bangor University, Wales, UK

^c College of Life and Environmental Sciences, University of Exeter, UK

^d Marine and Freshwater Research Centre, Galway-Mayo Institute of Technology, Ireland

^e Life and Environmental Sciences, University of Iceland, Reykjavik, Iceland

^f University of Cambridge, UK

^g School of Environmental Sciences, University of Derby, UK

^h School of BioSciences, University of Melbourne, Australia

ⁱ Institute of Oceanography and Fisheries, Split, Croatia

^j Institute of Geosciences, University of Mainz, Mainz, Germany

^k University of Alabama, USA

^l Laboratory of Tree-Ring Research, University of Arizona, USA

^m Memorial University, Canada

ⁿ Akvaplan-niva, Tromsø, Norway

^o Department of Geography and Anthropology and the Coastal Studies Institute, Louisiana State University, USA

^p School of Biological Sciences, University of Adelaide, Australia

^q Department of Biology, Aarhus University, Denmark

^r Biosphere 2, University of Arizona, USA

^s School of Geography, University of Melbourne, Australia

^t Atmosphere and Ocean Research Institute, The University of Tokyo, Japan

^u Univ Brest, CNRS, IRD, Ifremer, LEMAR, F-29280, Plouzane, France

^v Ocean and Earth Science, University of Southampton, UK

^w Department of Earth Sciences, Utrecht University, Utrecht, the Netherlands

^x AMGC Research Group, Vrije Universiteit Brussel, Brussels, Belgium

A B S T R A C T

Horizon scanning is an increasingly common strategy to identify key research needs and frame future agendas in science. Here, we present the results of the first such exercise for the field of sclerochronology, thereby providing an overview of persistent and emergent research questions that should be addressed by future studies. Through online correspondence following the 5th International Sclerochronology Conference in 2019, participants submitted and rated questions that addressed either knowledge gaps or promising applications of sclerochronology. An initial list of 130 questions was compiled based on contributions of conference attendees and reviewed by expert panels formed during the conference. Herein, we present and discuss the 50 questions rated to be of the highest priority, determined through an online survey distributed to sclerochronology community members post the conference. The final list (1) includes important questions related to mechanisms of biological control over biomineralization, (2) highlights state of the art applications of sclerochronological methods and data for solving long-standing questions in other fields such as climate science and ecology, and (3) emphasizes the need for common standards for data management and analysis. Although research priorities are continually reassessed, our list provides a roadmap that can be used to motivate research efforts and advance sclerochronology toward new, and more powerful, applications.

* Corresponding author.

E-mail address: tatr@norce-research.no (T. Trofimova).

<https://doi.org/10.1016/j.ecss.2020.106977>

Received 1 February 2020; Received in revised form 15 July 2020; Accepted 4 August 2020

Available online 9 September 2020

0272-7714/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Sclerochronology is a rapidly developing field of research. While growth bands in hard tissues of some organisms have long been observed and studied (e.g., Pulteney, 1781; Maton, 1805; Isely, 1914; Ma, 1934; Davenport, 1938; Adams, 1940; Kohler, 1964; Clark, 1974; Jones, 1981, 1983), the term “sclerochronology” was first introduced to the published literature in the 1970s. Analogous to the long-established field of dendrochronology (e.g., Fritts et al., 1971), sclerochronology was originally defined as “the study of growth patterns in calcareous exoskeletons and shells” (Buddemeier et al., 1974) and was first applied to coral research (Buddemeier et al., 1974; Hudson et al., 1976). The term has since been broadened to include various terrestrial and aquatic taxa with growth patterns, whereby the most common examples are fish (e.g., Coulson et al., 2014; Martino et al., 2019), coralline algae (e.g., Halfar et al., 2011; Williams et al., 2017), gastropods (e.g., Surge et al., 2013; Prendergast and Schöne, 2017) and bivalves (e.g., Jones et al., 1989). The list of sclerochronological archives is continuously expanding as more species are being assessed for their utility in sclerochronological studies. The term was redefined during the First International Sclerochronology Conference held at St. Petersburg, FL, USA in 2007 as “... the study of physical and chemical variations in the accretionary hard tissues of organisms, and the temporal context in which they formed ...” (Oschmann, 2009).

Today, sclerochronology is an increasingly diverse and interdisciplinary field. Apart from utilizing a wide array of archives, sclerochronology employs a suite of morphological, geochemical, microstructural, and crystallographic techniques. The data provided by sclerochronological studies have shown clear application across a range of fields, including ecology (e.g., Rhoads and Pannella, 1970; Rhoads and Lutz, 1980; Black et al., 2018), geophysics (e.g., Wells, 1963; Rosenberg and Runcorn, 1975; Zachariassen et al., 2000), archaeology (e.g., Coutts, 1970; Andrus, 2011; Wang et al., 2013), climate reconstruction (e.g., Jones et al., 1989; Butler et al., 2010; Tierney et al., 2015), and environmental (e.g., Steinhardt et al., 2016) and fisheries (e.g., Campana et al., 2001) sciences. Crossdated sclerochronological records, in particular, can provide powerful archives of past spatiotemporal environmental variability on local to hemispheric scales (Black et al., 2019). Advances in sclerochronological methods continually open up new applications, indicating that the full potential of sclerochronology has yet to be realized.

The triennial International Sclerochronology Conference (ISC) and other regular meetings with a sclerochronology component have played an important role in the development of the field. Journal special issues associated with such meetings have provided regular overviews of the most recent results and demonstrations of the potential of sclerochronology (Schöne and Surge, 2005; Gröcke and Gillikin, 2008; Oschmann, 2009; Wanamaker et al., 2011; Schöne and Gillikin, 2013; Butler and Schöne, 2017; Prendergast et al., 2017; Gillikin et al., 2019). Although significant effort has been made to review and synthesize recent findings, the sclerochronological community faces a variety of challenges and opportunities to be addressed in future work.

Now, 46 years after the term “sclerochronology” first appeared in the literature, and more than a decade after the first ISC, we have reached a timely moment to evaluate existing challenges and the most promising research directions. Inspired by previous examples from other research fields (e.g., Sutherland et al., 2011; Seddon et al., 2013; Patiño et al., 2017), the coordinating authors (Trofimova, Alexandroff, Mette, and Tray) initiated this community effort at the 5th ISC held in Split, Croatia in June 2019. The aim of our project is to advance the field and support its progress by identifying key research needs and providing an overview of persistent and emergent research questions. Due to the connection to the 5th ISC, the main focus of this article is on the sclerochronology of invertebrates and fish.

2. Methodology

Our project employed a horizon-scanning approach to identify community-defined priority research directions (for details see Supplementary Material 1; Fig. 1), adapted from similar studies performed in other research fields (e.g., Sutherland et al., 2011; Seddon et al., 2013; Patiño et al., 2017). At the initial stage, the coordinators (first four authors in the author list), in collaboration with expert panels and with the input from the wider research community, collected and curated a list of questions addressing fundamental knowledge gaps (Section “Foundations”) and promising applications of sclerochronological methods (Section “Applications”). In total, 202 questions and statements were submitted. Based on these contributions, we formulated an initial list of 130 questions (see Supplementary Material 2) that met previously outlined criteria (see Supplementary Material 3; adapted from Sutherland et al., 2011). An anonymous survey was launched and distributed to the sclerochronology and paleoclimatology communities through email list servers and social media. Participants were asked to rate each question on a 5-point Likert scale (Zero/Low/Neutral/High/Top Priority) in response to “Considering how fundamental the question is for sclerochronology, what should its priority be for future research?”.

In total, 52 complete survey responses were submitted. The top 25 questions from each of the two categories (“Foundations” and “Applications”) were selected by calculating the percentage of total respondents rating the question as a priority (“High priority” or “Top priority”, without differentiation). The expert panels reviewed the questions that did not make the final list to retain those that addressed underexplored research directions with potential to widen the horizon of sclerochronology (presented in section 3.3).

2.1. Limitations of this study

As a horizon-scanning project, the present study relies heavily on the expertise, interests, and skills of the participants. Continual efforts to exchange input and feedback from a diverse array of sclerochronological expertise were made throughout the development of the project in an attempt to maintain a wide perspective on the field and reduce bias. Even so, the fundamental questions identified by this exercise cannot be wholly separated from the research interests of the participants. While the final list of questions presents a community-informed perspective on priorities within sclerochronology, the rankings were determined by a relatively low number of participants ($n = 52$) with potentially strong geographical, archive-based, and research-based bias (see Supplemental Material 4). Additional bias was introduced during the preparation stage, as questions were collected via input from participants in the 5th ISC. While the ISC invites participation of sclerochronologists from all fields and regions, some archives are more highly represented than others. This is particularly the case for fish otoliths and mollusk shells, as the research utilizing these archives dominated the scientific presentations at the conference. This bias was also evident in the survey, where otoliths and mollusks were the primary or secondary expertise of all participants. Similarly, the most common applications of sclerochronology presented at the ISC were related to scientific inquiries in (paleo-)ecology and climate science. In addition, the location of the conference and associated travel, as well as other expenses, are contributing factors to regional bias. Thus, the final list by no means reflects the true boundaries of the extended field covered by the term “sclerochronology”. The 130 identified questions (see Results and Supplementary Material 2) represent a wide-ranging, but far from exhaustive, overview of possible future research directions and priorities. As this project was initiated to stimulate discussion among researchers, encourage collaboration, and spur new ideas for scientific advances, the results presented here, despite the inevitable biases, reflect a unique community-based insight.

3. Results

Below, we present 50 priority questions identified by this study. The questions are divided into two categories that identify (1) fundamental knowledge gaps ("Foundations of sclerochronology", 25 questions) and (2) promising applications ("Applications of sclerochronology", 25 questions). In this section, we also discuss the general motivation and background behind the questions. Due to the high diversity of archives and proxies, techniques, and research topics within sclerochronology, we aimed to provide a general overview, fully acknowledging that there will be exceptions to the rule. The literature cited in our paper deliberately represents a mixture of seminal papers, highlights in the field, and unique applications spanning a range of research groups and archives. These references are meant to point the reader to useful or interesting examples, and by no means represent a comprehensive review of the state-of-the-art on a particular question. We have grouped the 50 questions into topics to avoid repetition in the discussion and provide an outline of the general themes evident in the collection. In addition, eight questions not included in the list of the leading 50 questions, but highlighted by the expert panels as cutting-edge ideas, are presented in the final section of the results.

3.1. Foundations of sclerochronology

This section presents the highest ranked questions addressing knowledge gaps in our understanding of sclerochronological archives. It covers a range of topics, including but not limited to physiology, biomineralization, interpretation of sclerochronological data, development of standards, and establishment and calibration of new proxies.

3.1.1. Biomineralization

1. How, and to what extent, do vital effects influence biomineral stable isotope composition, elemental distribution, and elemental concentration?
2. What controls the incorporation of trace and minor elements into biogenic carbonates and how do these processes affect distribution of different trace elements between crystal lattice and organic phases?
3. Are there differences in biomineralization processes across ontogeny and between species and/or populations that affect skeletal isotopic composition and elemental concentrations?
4. How might climate and environmental change (e.g., ocean acidification) be altering processes of biomineralization?

A sound understanding of biological mineralization is fundamental to sclerochronology and the establishment of geochemical proxies in




Fig. 1. Conceptual scheme illustrating the methods employed to identify fundamental questions and priority applications in sclerochronology.

various contexts. Yet, the exact mechanisms driving biomineralization are not fully characterized, which is reflected in the questions in this topic [Q1-4]. The hallmark of biomineralization, as opposed to its abiogenic counterpart, is the remarkable control that organisms can exert over mineral formation (Weiner and Dove, 2003). These so-called “vital effects” (Urey et al., 1951), including kinetic and taxonomic effects (Weiner and Dove, 2003), can obscure the environmental signal in geochemical proxies, and thus confound proxy interpretation. An understanding of the role of vital effects in biomineralization is a major challenge for sclerochronology, as highlighted by Questions 1–3. Whereas the relationships of some geochemical properties and environmental variables are well established for sclerochronological archives (e.g., stable oxygen isotope value of biogenic carbonate ($\delta^{18}\text{O}_\text{c}$); see Topic “3.1.5 Temperature reconstructions”), other properties are often difficult to interpret due to taxon-specific physiological effects. For example, previous studies have demonstrated that Mg to Ca ratios in coralline algae (e.g., Nürnberg et al., 1996; Kamenos et al., 2008), and Sr to Ca ratios in tropical shallow-water corals and sclerosponges (e.g., Beck et al., 1992; Rosenheim et al., 2004) are robust temperature recorders, but are currently still difficult to interpret in otoliths (e.g., Campana, 1999) and bivalves, specifically those with aragonitic shells (e.g., Zhao et al., 2017a; Gillikin et al., 2019). Stable carbon isotope values ($\delta^{13}\text{C}$) of coralline algae (Williams et al., 2011), corals (Swart et al., 2010; Dassié et al., 2013), and sclerosponges (Druffel and Benavides, 1986; Böhm et al., 1996) have been successfully used for environmental reconstructions, e.g., as a proxy for the $\delta^{13}\text{C}$ value of dissolved inorganic carbon in ambient water. However, vital effects on the $\delta^{13}\text{C}$ signature are suspected among corals, fish otoliths, and some mollusks (Kalish, 1991; Iacumin et al., 1992; McConnaughey et al., 1997; Lorrain et al., 2004; McConnaughey and Gillikin, 2008). Further characterization of the mechanisms controlling the isotope and element chemical variability in biominerals will improve the usability of proxies across various taxonomic groups and therefore is crucial for future sclerochronological studies [Q1].

While biomineralization processes differ among taxonomic groups, they can also vary through an individual’s lifetime, even within skeletal structural layers, and between individuals. Apart from environmental factors, the chemical composition, as well as microstructure, of biominerals is affected by genetics (Carter, 1980; Clarke et al., 2010; Norrie et al., 2019) and can vary throughout ontogeny (Marshall and McCulloch, 2002; Elliot et al., 2003; Gillikin et al., 2007; Nishida et al., 2011; Grammer et al., 2017; Reynolds et al., 2019). The role of these factors in biomineralization is not well understood in the broad context of archives and proxies, representing a significant knowledge gap for sclerochronology [Q3]. In the case of element proxies, additional complexity may result from the presence of non-lattice bound trace elements (e.g., Takesue et al., 2008). Mechanisms of elemental incorporation into biogenic carbonates, therefore, require a special focus in future research to improve application of trace element records in sclerochronology [Q2].

Development of biophysical models coupling metabolism, biomineral growth, and elemental and isotopic dynamics could provide much needed mechanistic insights into vital effects in terms of their effects on environmental proxies, and the use of elemental and isotopic compositions of biominerals as physiological tracers. This will require experimental and theoretical modeling work, but frameworks such as Dynamic Energy Budget theory provide a platform suitable for model development (e.g., Fablet et al., 2011; Spalding et al., 2017) [Q1-3].

Modern climate change and associated ocean acidification can pose major threats to marine calcifying organisms due to their potential effects on biomineralization. Changes in biomineralization can have consequences for the survival of species, as well as the applicability of proxies, and therefore require further research [Q4]. In the context of sclerochronological research, it is particularly important to understand natural variations in biomineralization across different physical and biotic stressors (e.g., Telesca et al., 2019; De Noia et al., 2020), and

physiological responses and adaptations leading to changes in biomineralization rate, mineralogy, and geochemistry of skeletal structures. Although the effects of ocean acidification on biomineralization are increasingly studied (e.g., Checkley et al., 2009; Ivanina et al., 2013; Fitzer et al., 2014; Milano et al., 2016; Cornwall et al., 2018; Cross et al., 2019), the results suggest variable responses among taxa (e.g., Zhao et al., 2018, 2020), likely depending on the degree of biological control over biomineralization and variable compensatory mechanisms and their energetic costs (Kleypas et al., 2005; Spalding et al., 2017; Melzner et al., 2020, and references therein). More research is needed to fully understand potential outcomes and identify possible patterns. Furthermore, growing concerns about the overall effects of acidification on marine ecosystems call for reliable proxies for past ocean acidification events. Boron isotopes ($\delta^{11}\text{B}$) and U to Ca ratios have recently shown potential as proxies for pH levels (Hönisch et al., 2012; Raddatz et al., 2014; Foster and Rae, 2016; Jurikova et al., 2019). However, more research is needed to evaluate the broad applicability of these proxies across sclerochronological archives, which is intimately linked to the understanding of biomineralization processes under changing environments.

3.1.2. Drivers of skeletal growth

5. What are the specific processes by which climate signals are translated into growth of calcified structures?
6. What determines the timing of the growth season and does it vary throughout ontogeny?
7. Are the growth/chemical responses to specific environmental drivers consistent/stationary over geologic time?
8. How can we predict the sclerochronological patterns (growth and/or chemical records) expected under differing combinations of movement, physiology, and environmental change?

Measurement of growth patterns and structures within skeletal archives is a standard procedure for sclerochronological work. Whereas the growth record serves as an age model to anchor geochemical or other proxies, individual or population-averaged growth variability is itself often a valuable environmental or ecological proxy (e.g., Halfar et al., 2011). For population-averaged records, the process of crossdating to produce absolutely dated growth increment width chronologies hinges on the assumption that common environmental drivers impart a shared growth response within a population (Black et al., 2019). However, questions remain regarding the mechanistic pathways leading to growth responses from environmental and biological drivers [Q5]. Combinations of food availability and quality, temperature, and light intensity are commonly identified as primary environmental drivers of year-to-year increment width variability that differ among sclerochronological archives (for a brief overview, see Schöne and Surge, 2005).

Structural properties of biominerals provide other promising proxies reflecting interactions between biology and environment (e.g., Füllenchbach et al., 2014; Milano et al., 2017; Höche et al., 2020). The process by which environmental signals are translated into microstructure variability also informs mechanisms of biomineralization, and is increasingly being studied (e.g., Nishida et al., 2015; Checa, 2018). More detailed insight into the archive-specific drivers behind increment width and microstructural variability will enable robust linkages between growth proxy records and environmental variability. It will also contribute to a better understanding of the synchrony or lack of synchrony among individuals in a population (Marali and Schöne, 2015; Muslic et al., 2013; Rountrey et al., 2014). Finally, the question whether drivers of growth are constant and stationary over geologic time should be considered [Q7]. Proxy records collected from sub-fossil material for which a precise calendar date cannot be attached (“floating” records) offer windows into past time intervals (e.g., Kilbourne et al., 2004; Scourse et al., 2006). However, the discussion of potentially variable

growth drivers in past time intervals or across the lifetime of an individual, and the resulting impact on proxy reconstructions, has received little attention.

A common approach to the determination of the timing of the growing season for sclerochronological archives is analysis of seasonal oxygen isotope ($\delta^{18}\text{O}_\text{C}$) profiles within annual increments (Weidman et al., 1994; Schöne and Surge, 2005; Mannino et al., 2008; Judd et al., 2018) or trace elemental ratios (e.g., corals, DeLong et al., 2011; coralline algae; Williams et al., 2014). Because many sclerochronological archives exhibit decreasing growth rate as they age (e.g., bivalve shells, fish otoliths), accurate determination of the full range of the growing season is best accomplished by sampling the wider, juvenile increments (Goodwin et al., 2003). Extrapolating these findings throughout the life of the animal, however, is problematic if there are ontogenetic effects on the duration and or timing of the growing season. Such effects are not consistent among species (Goodwin et al., 2003; Schöne et al., 2005), warranting further investigation [Q6].

The translation of climate and environmental signals into growth and geochemical signatures is further complicated in the case of mobile organisms. The environment experienced by an animal may vary across large-scale migrations as well as differing habitat utilization across its life history (e.g., Gillanders et al., 2015; Roberts et al., 2019). Disentangling interpretations of environmental change from interpretations of animal movement is a difficult task. Modeling provides a powerful tool to predict the effects of differing combinations of life history patterns, environmental change, and potentially adaptive drivers of growth, as discussed above [Q8]. Whereas research in this area has advanced in recent years (e.g., van der Sleen et al., 2018; Hobbs et al., 2019; Trueman et al., 2019), it is still recognized as a priority research question in the field.

3.1.3. Data standards

9. What common data standards should be adopted to improve our ability to compare sclerochronological datasets with each other and with other datasets?
10. What level of sample replication is required for geochemical records for sound estimation of uncertainty associated with inter-individual variability and ensuring comparability between records?

Variability in sclerochronological methods and reporting standards can affect the quality and comparability of datasets, representing a major challenge for the field [Q9, Q10]. Whereas the tree-ring community has agreed-upon methods and reporting standards for data sharing (e.g., International Tree-Ring Data Bank (ITRDB)), similar agreements are missing in the sclerochronological community. The issues with data reporting and sharing became apparent during the various PAGES2k projects that brought together publicly shared paleo data from a variety of archives to build global databases (e.g., PAGES2k Consortium, 2013; Tierney et al., 2015; Emile-Geay et al., 2017; Konecky et al., 2020). Recently, global efforts have been made to improve scientific data interoperability and reusability (Wilkinson et al., 2016; McKay and Emile-Geay, 2016). Initiatives to provide growth, geochemical, microstructural, or other proxy data in common formats have also begun within the sclerochronology community (Dassié et al., 2017; Khider et al., 2019; Tray et al., 2020). Collaboration among sclerochronology researchers to standardize data collection methods and data reporting, as well as standards for archiving physical samples, are needed to address these issues. Some studies comparing multiple, inter-species sclerochronology datasets with varying temporal resolutions provide examples of progress in this area (Matta et al., 2010; Ong et al., 2016; Peharda et al., 2018), however, further extensions of such work is a priority for future research.

At present, various statistical methods are used to account for inter-individual variability within sclerochronological datasets, one of the

most commonly used being mixed-effects models (Weisberg et al., 2010; Morrongiello and Thresher, 2015). Additionally, power analyses can be used to estimate appropriate sample sizes (Toft and Shea, 1983). Whereas these are acceptable methods for studies that address individual-level variation, there is less certainty about their applicability to geochemical data [Q10] (e.g., stable isotope and trace element records; but see Grammer et al., 2017 and Macdonald et al., 2019). Correlation coefficients paired with significance levels (i.e., *p*-values) are typically used to determine robustness of environmental reconstructions (e.g., Montagna et al., 2014 for seawater temperature). Still, there are arguments against using these metrics, due to the short length of marine instrumental records available for calibration (Crowley et al., 1999; Corrège, 2006; Finney et al., 2010) and problems with statistical inference (Wasserstein et al., 2019). In addition to correlation, coral replication studies have used expressed population signal (EPS), absolute differences, and root mean squared statistical tests to assess replication and reproducibility at the intra- and inter-coral colony levels and between species at the same location (DeLong et al., 2007, 2011; Wu et al., 2014; Dassié et al., 2014). Chronological uncertainty, especially in non-crossdated reconstructions, needs to be better understood and assessed in the various sclerochronological archives (e.g., Comboul et al., 2014). There is, therefore, a need to define and clarify the types of data and analyses that constitute a sclerochronological reconstruction, and to further develop and standardize statistical techniques to quantify and account for uncertainty.

3.1.4. Data analysis and interpretation

11. What methods can we use to better assess the leads, lags, and synchronicities in sclerochronological records across large spatial regions?
12. How can we disentangle the separate and combined effects of multiple causal factors in sclerochronological records?
13. How can common environmental signals be identified in multiple records which have different spatial and temporal scales and resolutions?
14. How can we disentangle multiscale spatial and temporal variability within sclerochronological records?
15. To what extent do variations in multiannual to multicentennial growth patterns represent a community/ecosystem response to changing environmental conditions?

Interpretation of environmental signals in sclerochronological records is not a trivial task, and this is reflected in the questions in this topic [Q11-15]. Linked with the issue of standardizing data sharing (see Topic "3.1.3 Data standards"), these questions highlight methods and strategies for sclerochronological data analysis that require further development and standards for sharing data that have been agreed upon by the community.

Sclerochronological archives provide high-resolution (e.g., daily, annual) environmental proxy data, which, provided that live-collected samples are used, are absolutely dated. This makes them uniquely suited for studies of spatiotemporal heterogeneity in the response of different components of the Earth system to forcing factors (e.g., Evans, 1972; Ohno, 1989; Black et al., 2014; Reynolds et al., 2016; Black et al., 2019). Therefore, the development of methods and strategies for the determination of leads and lags across different spatial scales is an important avenue for future research [Q11]. Methods used by tree-ring (e.g., Cook et al., 2004) and PAGES2k communities (e.g., Tierney et al., 2015; Atsawawanunt et al., 2018; Konecky et al., 2020) in compilation studies could be assessed and applied to sclerochronological reconstructions.

An understanding of the extent to which individual or population-averaged growth records represent a community or ecosystem response to a changing environment is crucial for interpretation of climate signals [Q12]. Sclerochronological records (e.g., growth records

and geochemical data) often encapsulate a response to a suite of environmental factors which, importantly, can act on different biological and temporal scales (e.g., [Morrongiello et al., 2019](#)). Disentangling these influences is of major importance for interpreting environmental signals at different resolutions and scales within one record, and for identifying common signals across multiple records [Q13-15]. To deconvolve this complexity, past studies have employed multiple linear regression (e.g., [Mette et al., 2016](#)), principal component analysis (e.g., [Black et al., 2014](#)), univariate and multivariate mixed-effects models (e.g., [Morrongiello et al., 2015](#); [Macdonald et al., 2019](#)), dynamic energy budget models (e.g., [Pecquerie et al., 2012](#)), and Bayesian hierarchical modeling (e.g., [Helser et al., 2012](#)). Further assessment of these tools and the adoption of new statistical techniques for time series analysis of sclerochronological records will undoubtedly improve the interpretation and impact of sclerochronological studies.

3.1.5. Temperature reconstructions

16. How can we improve estimates of past water isotopic composition to increase accuracy of temperature reconstructions?
17. Why do we often observe an offset between seawater temperature reconstructed from oxygen isotope values (using widely applied paleotemperature equations) and those measured *in situ*?
18. How can we determine if species-specific paleotemperature equations are a valid and necessary approach to increase the accuracy of paleotemperature reconstructions?
19. What are the limitations of using clumped-isotope paleothermometry to constrain isotopic paleotemperature estimates from fossil organisms?

Oxygen isotope values of biogenic carbonates ($\delta^{18}\text{O}_c$) have long been used to reconstruct temperatures ([Urey et al., 1947](#); [Epstein et al., 1953](#)). Sclerochronological archives allow the construction of highly resolved $\delta^{18}\text{O}_c$ records, which have important applications in many research fields, such as climatology, physiology, anthropology, paleoceanography, and ecology, among others. The applicability of $\delta^{18}\text{O}_c$ as a temperature proxy has been established for all sclerochronological archives, including scleractinian corals (e.g., [Weber and Woodhead, 1972](#)), fish otoliths (e.g., [Devereux, 1967](#)), mollusks (e.g., [Weidman et al., 1994](#)), and coralline algae (e.g., [Halfar et al., 2008](#)). The premise for using $\delta^{18}\text{O}_c$ to study temperature is that the fractionation of oxygen isotopes during biomineralization is temperature-dependent and in (near) equilibrium with the ambient water ([Grossman and Ku, 1986](#); [Kelemen et al., 2017](#); [Thorrold et al., 1997](#); [Weidman et al., 1994](#); but see [Smith et al., 2000](#)). The accuracy and reliability of this method depend on knowledge of (1) the $\delta^{18}\text{O}$ value of the ambient water ($\delta^{18}\text{O}_w$) and other site-specific physical and chemical properties [Q16-18], and (2) the specific biomineralization processes within the chosen archive [Q17, 18].

Question 16 raises a well-known and pertinent issue in $\delta^{18}\text{O}$ paleothermometry and its application in sclerochronological studies ([Prendergast and Stevens, 2014](#); [Yan et al., 2014](#)). Given that $\delta^{18}\text{O}_c$ is a function of $\delta^{18}\text{O}_w$ and temperature-driven fractionation, independent estimates or measurements of $\delta^{18}\text{O}_w$ are crucial. In studies of modern samples, $\delta^{18}\text{O}_w$ is often estimated using a region-specific relationship between salinity and $\delta^{18}\text{O}_w$ (i.e., mixing lines) ([LeGrande and Schmidt, 2006](#)). However, the accuracy of mixing lines in local studies needs to be scrutinized, especially at locations where freshwater input causes high $\delta^{18}\text{O}_w$ variability (e.g., [Wagner et al., 2011](#)). One approach to overcome the issue of unknown $\delta^{18}\text{O}_w$ is to use paired proxies to constrain temperature with the element-to-element ratio and solve for $\delta^{18}\text{O}_w$. Examples for such paired proxies in aragonitic sclerochronological archives are Sr/Ca and $\delta^{18}\text{O}$ in corals and sclerosponges (e.g., [McCulloch et al., 1994](#); [Gagan et al., 1998](#); [Ren et al., 2002](#); [Rosenheim et al., 2004](#)), $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in bivalve shells (e.g., [Reynolds et al., 2019](#)), and $\delta^{18}\text{O}$ in otoliths paired with $\delta^{18}\text{O}$ in bivalve shells ([Wang et al., 2011](#)), among others. [Carroll et al. \(2006\)](#) suggested another approach, using hydrogen

isotope values (δD) within the organic matrix of freshwater bivalves to independently estimate $\delta^{18}\text{O}_w$. Carbonate clumped isotope values (Δ_{47} , Δ_{48}) provide an avenue to circumvent this question altogether, as this method does not require an estimate for $\delta^{18}\text{O}_w$ ([Eiler, 2007](#); [Fiebig et al., 2019](#)). This is especially useful in studies where $\delta^{18}\text{O}_w$ uncertainty is high, for example in estuarine environments or deep-time marine settings (e.g., [Martin and Letolle, 1979](#); [de Winter et al., 2018](#)), or where kinetic effects are present. However, as Question 19 shows, limitations of the Δ_{47} value as a paleothermometer have yet to be fully assessed. One drawback is the requirement for relatively large sample sizes; conventional carbonate clumped isotope techniques require 3–7 mg of carbonate sample, while recent techniques have been able to lower the number to 14–20 replicates of 100 μg ([Meckler et al., 2014](#); [Müller et al., 2017](#)). This size requirement limits high-resolution temperature reconstruction, and most archives will have specific limitations due to the overall size of the calcified structure. Additionally, yet-to-be assessed taxon-specific vital effects may be a limiting factor ([Eiler, 2011](#)). Other limitations of clumped isotope paleothermometry are its time-consuming and demanding analytical methods and its sensitivity to diagenetic overprint ([Leutert et al., 2019](#)).

Offsets between the $\delta^{18}\text{O}_c$ -derived temperatures and measured *in situ* temperatures have been reported for different archives and locations (e.g., [Weber, 1970](#); [McConnaughey, 1989](#); [Bonitz et al., 2017](#); [Dunbar and Wefer, 1984](#); [Kelemen et al., 2017](#)). In response to this issue, species-specific or site-specific calibrations are often formulated [Q17, 18]. However, performing new and specific calibrations might lead to false conclusions, e.g., a steeper slope between $\delta^{18}\text{O}_c$ and temperature can mask the real temperature amplitude ([Waite and Swart, 2015](#)). It is therefore crucial to understand what causes the observed offsets and eliminate inaccurate $\delta^{18}\text{O}_w$ assumptions [Q16] or sampling methods as potential sources of error. Suspected sources of unexpected offsets or observed variability in $\delta^{18}\text{O}_c$ include (1) isotopic alteration caused by mechanical sampling or analytical methods ([Tobin et al., 2011](#); [Waite and Swart, 2015](#); but see [Foster et al., 2009](#)), (2) signal aliasing as a result of limited sampling resolution ([Goodwin et al., 2003](#); [DeLong et al., 2007](#); [Gagan et al., 2012](#)), (3) sampling imprecision considering layers of different architectural structures, which can influence $\delta^{18}\text{O}_c$ ([Leder et al., 1996](#); [DeLong et al., 2016](#); [Mette et al., 2018](#); [Trofimova et al., 2018](#)), and (4) cleaning methods altering primary mineralogy and $\delta^{18}\text{O}_c$ signal ([Boiseau et al., 1997](#); [Wierzbowski, 2007](#); [Holcomb et al., 2015](#); [Grottoli et al., 2005](#)), and vital effects ([McConnaughey, 1989](#)). One way to more accurately quantify $\delta^{18}\text{O}_c$ -temperature relationships is through experiments in controlled or closely monitored settings (e.g., [Wanamaker et al., 2007](#); [Ford et al., 2010](#); [Nishida et al., 2014, 2015](#); [Sakamoto et al., 2017](#)). It is crucial to improve our understanding of oxygen isotope dynamics in organisms and the biominerals they synthesize in the effort to explain offsets between reconstructed and *in situ* temperatures, while sampling methods and local context also have to be taken into account.

3.1.6. Archive-specific research

20. What not-yet-identified long-term sclerochronological archives exist, especially outside of the North Atlantic region?
21. What environmental parameters can be reconstructed from trace element concentrations and ratios within mollusk shells, and why are some trace element proxies unreliable?
22. Through which pathways are trace and minor elements transported into mollusk extrapallial fluid, and from where are they sourced (e.g., digested food, directly from water)?
23. Why is it that sometimes within the same population of bivalves, not all of the individual growth patterns from live-collected specimens crossmatch and how should we deal with such inter-individual variability?
24. What drives the formation of annual growth increments in fish otoliths?

25. How does inter-individual variation in growth patterns in fish affect long-term growth time series?

Newly identified archives or proxies are often sought to address geographic, environmental, and/or temporal gaps in earth system research (e.g., Peharda et al., 2016; Milano et al., 2017). Established, long-lived (>100 years), sclerochronological archives have limited geographic ranges, resulting in a high density of research focused in certain regions. This is particularly true in the North Atlantic Ocean, where studies using mollusk shell archives, especially *Arctica islandica*, dominate the literature (Steinhardt et al., 2016). The western tropical Atlantic and Pacific are hotspots for research using reef-building coral archives, representing another geographic bias in the availability of long-term proxy archives (Corrège, 2006). Some long-lived species are becoming well-established sclerochronological archives applicable in other regions (e.g., *Tridacna* sp., Jones et al., 1986; Elliot et al., 2009; Killam et al., 2020; geoducks, Strom et al., 2004; Black et al., 2009; coralline algae, Williams et al., 2017; deep sea corals, Robinson et al., 2014). However, to better address past climate and environmental research questions outside of the tropics and the North Atlantic, in particular, the search for long-lived archives from other regions is a priority [Q20].

Mollusk shells and fish otoliths are among the most frequently used sclerochronological archives. Several mollusk- and fish-specific issues were highlighted in the community survey. In particular, trace element concentrations, a reliable geochemical proxy among many archives, have been shown to be problematic or inconsistent within and across most mollusk species (see Topic “3.1.1. Biomineralization”). Little is known about the uptake of elements from the ambient water to the site of biomineralization, the transport mechanisms and pathways of elements within the body, nor the specific incorporation mechanisms of elements in the skeletal hard parts (Suzuki et al., 2009; Zhao et al., 2017b). Solving these questions will likely require heavy involvement from cell biologists and geneticists to improve our understanding of elemental proxies among molluscan species [Q21-22].

Synchronous growth is remarkably prevalent among mollusks (Jones et al., 1989; Weidman et al., 1994; Black et al., 2019), meaning it is unusual to encounter an individual that does not match the population growth pattern. When such individuals are identified, it is important to assess the quality of the material and clarity of the increment boundaries, and the geographic extent over which the samples were collected, as well as the experience of the worker with the particular population, to establish whether the shell truly exhibits a unique growth pattern. However, the reasons why some shells have unclear or irregular growth and do not easily crossmatch with the local population growth pattern are poorly understood [Q23] and could be biologically based. The extent to which difficult-to-crossdate individuals within a population is a problem has not been addressed within the literature, revealing an opportunity to improve understanding of molluscan growth records, and other crossdatable archives, as environmental proxies. While some research utilizes the varying strength of the common signal as an environmental record in itself (Marali and Schöne, 2015), more work is needed to assess the scope and implications of individual growth variability.

Fish otolith growth increments form with an annual periodicity in almost all fish species, even if the species lives in a relatively constant environment, such as the deep sea (Cailliet et al., 2001; Campana, 2005). Photoperiod, temperature, growth, sexual maturation, feeding, migration, and other processes have all been linked to annual increment formation (Campana and Thorrold, 2001; Grønkvær, 2016), but the ubiquity of these increments suggests that there is an innate physiological mechanism involved (e.g., circadian periodicity). Many analytical approaches to otolith sclerochronology average out individual-level variation in growth and focus on the mean population trends. As fish otolith sclerochronologies continue to develop, it is critical to understand the mechanisms driving otolith growth increment formation

[Q24]. Furthermore, between short- and long-lived species, attempts to crossdate individuals within a population produce varying results in terms of synchrony and environmental relationships (Rountrey et al., 2014). Understanding the occurrence and drivers behind individual growth variability [Q25] is a priority for advancing research using long-term growth series from otoliths (Morrongiello and Thresher, 2015; Morrongiello et al., 2019). For example, understanding temporal growth variability within otolith time series could aid fisheries management by improving stock discrimination methods (Denechaud et al., 2020) and assessing the impact of harvest on populations (Morrongiello et al., 2019).

3.2. Applications of sclerochronology

This section presents the highest-ranked questions addressing potential applications of established sclerochronological archives and techniques to outstanding questions in a wide range of research fields, including enhanced applications for climatological, oceanographic, ecological, and cultural studies.

3.2.1. Global climate

26. How spatially heterogeneous were climate and environmental conditions under “normal” past conditions (i.e., as opposed to extreme climate scenarios, such as the Little Ice Age, Last Glacial Maximum, Younger Dryas)?
27. How did seasonality vary in the past in the temperate climate zone?
28. Can we detect changes in variability in sclerochronological records that indicate an approach to a climate or environmental tipping point?
29. How did major climate changes affect the intrinsic variability of El Niño/Southern Oscillation (ENSO) in the past?
30. To what extent do sclerochronologies covary with tree-ring data, and what does that tell us about the coherence of climate variability over hemispheric scales through time?
31. How can we integrate tropical growth-increment data with mid- and upper-latitude sclerochronologies to explore tropical-extratropical teleconnections?
32. Which sclerochronological data are most suited for climate model assimilation?

Previous work has demonstrated successful applications of sclerochronological records to questions of global climate through paleoclimate reconstruction (Eakin and Grotto, 2006; Reynolds et al., 2018). The annual and often subannual resolution of sclerochronological records, especially when supported by a crossdated chronology, makes them uniquely suitable to address questions of seasonality, rapid climate change, tipping points, and lead-lag climate responses across the Earth system (Corrège, 2006; Butler and Schöne, 2017; Reynolds et al., 2016). Several key directions for sclerochronological climate research on these topics were identified as a priority for future research [Q26-29].

The rapidly expanding range of species, geographic, and temporal coverage represented in published sclerochronological records enables new avenues for research on long-term, high-resolution climate variability. It is now possible to address questions of climate variability across broad spatial areas to explore large-scale climate and teleconnections (e.g., northern and southern hemisphere water mass temperatures, Thresher et al., 2014; circumtropical SST, Tierney et al., 2015; Pacific and Atlantic ocean-atmosphere teleconnections, Wana-maker et al., 2019) across diverse time periods (e.g., late Holocene, Black et al., 2014; middle Eocene, Bougeois et al., 2014; Paleogene, Huyghe et al., 2015; late Cretaceous, de Winter et al., 2017). Because the geographic ranges of sclerochronological archives differ, large-scale climate reconstructions often require compilation of records sourced from different species (e.g., Reynolds et al., 2018). The comparability of

such records must be assessed, and limitations explored in order to apply both multispecies sclerochronology and multiproxy-based interpretations to questions of, for example, atmospheric and oceanic interactions [Q30] and tropical-extratropical teleconnections [Q31].

Additionally, much attention has been placed on how climate modeling interfaces with sclerochronology [Q32]. Many studies utilize data from climate models to inform proxy interpretations (e.g., Tindall et al., 2017; Trueman et al., 2019). However, with the exception of coral records (see Okazaki and Yoshimura, 2017), the use of sclerochronological records in large-scale proxy data assimilations, has rarely been accomplished despite its high potential value (Goosse, 2016; Pyrina et al., 2017). More work is needed to assess the quality and richness of detail provided by sclerochronological records that will enable their appropriate inclusion in paleoclimate reanalyses (see Schmidt et al., 2014). Efforts to maximize sampling resolution, improve measurement techniques, and provide a robust understanding of the mechanisms by which environmental signals are embedded within the archive will improve confidence in these modeling applications (Goosse, 2016; Butler and Schöne, 2017).

3.2.2. Paleooceanography

33. How can sclerochronological proxies be used to study historical changes in the extent of Arctic sea ice?
34. How can we use sclerochronological archives to monitor changes in the role of the oceans as a buffer for carbon emissions and heat?
35. How can we use sclerochronological archives to detect high resolution variability in strength of Atlantic Meridional Overturning Circulation (AMOC)?
36. How has the ^{14}C reservoir effect varied over time and at different temporal scales (e.g., subannual, annual, decadal)?
37. What can sclerochronological records tell us about the links between the marine carbon and nitrogen cycles in the past, especially during times of abrupt climate change?
38. How can we utilize both high resolution sclerochronological records and traditional paleoceanographic data (e.g., sediment core records) to produce spatial reconstructions of broad scale climate variability?
39. How can sclerochronological records from shelf seas be used as proxies for open-ocean conditions and what are the temporal and spatial limitations?

Questions specific to paleoceanography highlight motivations and research avenues similar to those presented in the topic "Global Climate" (see Topic "3.2.1. Global climate"). The wide range of established marine sclerochronological archives and proxies enables deeper understanding of past and present oceanographic processes at high temporal resolution. The questions in this topic [Q33-39] demonstrate some of the most prominent and promising applications of sclerochronological data for solving long-standing questions in paleoceanography.

Sclerochronological data have been successfully used to study past sea ice variability in the Arctic [Q33] (e.g., Halfar et al., 2013; Chan et al., 2017; Hetzinger et al., 2019), track local oceanic uptake of anthropogenic CO_2 [Q34] (e.g., Schöne et al., 2011; Williams et al., 2011; Dassié et al., 2013), estimate oceanic heat content and temperature variability [Q34] (Linsley et al., 2015), and assess changes in past oceanic circulation [Q35] (e.g., Wanamaker et al., 2012). The ability to combine radiocarbon (^{14}C) dating and independent sclerochronological age models (e.g., growth chronologies) has been used to reconstruct local ^{14}C reservoir changes through time [Q36] (e.g., Druffel and Griffin, 1993; Sherwood et al., 2008; Butler et al., 2009; Wanamaker et al., 2012; Hirabayashi et al., 2017). Furthermore, promising results have been obtained in studies on nitrogen isotope ($\delta^{15}\text{N}$) values in sclerochronological archives (e.g., Yamazaki et al., 2016; Sherwood et al., 2014; Gillikin et al., 2019, and references therein). There is great

potential for stable nitrogen and carbon isotope records to be coupled to enable detailed investigation of marine food web links in the past, as well as other aspects of the marine nitrogen and carbon cycles [Q37].

To fully address important questions on ocean history and broad-scale climate variability [e.g., Q33-37], spatiotemporal expansion of proxy data coverage is needed. Use of paleoceanographic data based on sediment cores in conjunction with diverse sclerochronological archives represents a promising avenue for future research [Q38]. While some work has been done in this area (e.g., Reynolds, et al., 2013), literature on methods and limitations for such work is sparse. Related to this issue, the temporal and spatial limitations of sclerochronological reconstructions of open-ocean conditions based on archives from shelf seas (e.g., bivalves and tropical corals) should be further investigated [Q39].

3.2.3. Paleoeology and human-environmental interactions

40. How can sclerochronological tools help us to decide which time period/condition provides an appropriate baseline for studies which require "natural", "pristine" or pre-human impact data on the environment?
41. How can sclerochronology be used to assess the anthropogenic impacts on overall ecosystem process and structure throughout the Holocene?
42. How can we use sclerochronological data to detect the first signs of human impact on the marine system through fishing and climate change?
43. How can sclerochronological records be used to assess changes in fish and shellfish populations due to harvesting?
44. In the context of global climate change, which aquatic ecosystems/environments experience ecological change first or to the greatest degree (e.g., open ocean, upwelling, subtidal, intertidal, estuarine, riverine, lacustrine)?
45. How can we use sclerochronology to distinguish variations in the effects of climate change on marine ecosystems at various spatial scales (e.g., local, regional, and global)?
46. How can we use sclerochronological archives to monitor the lag in the ecosystem response to climate change and other environmental change in the oceans?
47. How can we use sclerochronology to quantify the rate of recovery of marine ecological systems from natural or anthropogenic disturbances?
48. How are different classes of chemical pollutants presented in the sclerochronological record and can their temporal distribution be inferred?
49. How can sclerochronological records be used to study eutrophication dynamics in coastal ecosystems?
50. How can sclerochronological records be used to infer the frequency and intensity of hypoxia and anoxia events in the past?

It is abundantly clear that most, if not all, modern ecosystems are severely affected by human activities (IPCC, 2018). Quantifying and understanding the impact of human activities on these ecosystems in the past [Q40-43] is among the key challenges in paleoecology (Seddou et al., 2013). Climate change, pollution, and industrial fishing have been identified as major threats to aquatic biodiversity and ecosystem health. It is therefore important to monitor environmental change in order to inform the definition of goals and directives for environmental protection, health assessment, and restoration [Q44-50]. However, due to shifting baselines and scarce data, restoration targets are often based on images of ecosystems that have already been disturbed and are thus no longer pristine [Q40]. Sclerochronology provides powerful tools to extend and analyze such baselines by constructing highly resolved chronologies that span centuries or millennia.

Sclerochronological archives are often used to study the human impact on marine species and ecosystems in the past and present [Q41-43]. Fishing and shellfish harvesting, in particular, have been important

sources of food for humans since prehistoric times. Shell middens (i.e., anthropogenic sites of shellfish remains) are a particularly useful resource for studies of early human-environmental interactions, as they can be found on coastlines worldwide (except for Antarctica) and have been deposited throughout the Holocene and beyond (see [Erlandson, 2001](#)). Material obtained from shell middens has been used to study season of capture, resource management, measures of overharvesting, and environmental changes (see [Andrews et al., 2003](#); [Andrus, 2011](#); [Geffen et al., 2011](#); [Carré et al., 2019](#); [Butler et al., 2019](#)). Accurate age-structured information and growth rates of commercial species are of great importance in fisheries science and have been gathered extensively from fish otoliths, and also from mollusk shells or statoliths ([Campana and Thorrold, 2001](#); [Henry and Nixon, 2008](#); [Ezgeta-Balic et al., 2011](#); [Hollyman et al., 2018](#)). Otolith chemistry is also used to study the thermal life history of populations, or as a geochemical tracer to determine past locations and stock identity (e.g., [Campana, 1999](#) and references therein; [Wang et al., 2016](#)). Moreover, additional information on food-web dynamics can be gained from $\delta^{15}\text{N}$ composition in carbonate-bound organic material (e.g., [Gillikin et al., 2017](#); [Sirot et al., 2017](#)). Sclerochronology thus offers valuable insight into the impact of human activity and climate change on commercial species as well as the wider ecosystem. Further advances in this field to inform stakeholders and management are thus a priority research area.

Carbon dioxide emissions and climate change affect marine environments in complex ways through changes in temperature, mixing regimes, circulation patterns, oxygen solubility, and carbon chemistry. Responses of marine ecosystems are manifold, interlinked, and spatially and temporally heterogeneous [Q44-47]. While single-population chronologies provide limited insight into an ecosystem, a more holistic view can be achieved by comparing sclerochronological data from archives at different trophic levels or from different regions (e.g., [Black, 2009](#); [Reynolds et al., 2017](#)). This approach enables us to study leads and lags in response to climate change or environmental disturbances between different ecosystems, regions, or taxa. Another method used to study or predict ecosystem response to climate change is the use of sclerochronological data to parameterize forecasting models (e.g., [Morrongiello et al., 2012](#); [Barrow et al., 2018](#)). These and other methods (e.g., dynamic energy budget models) for investigating characteristics of ecosystem response to environmental change should be further explored.

The impacts of environmental pollution on aquatic ecosystems and mixing regimes is a matter of rising concern [Q48-50]. Some studies have applied sclerochronological techniques to monitor heavy metal pollution retrospectively (e.g., [Scott, 1990](#); [Gillikin et al., 2005](#); [Krause-Nehring et al., 2012](#); [Holland et al., 2014](#)), which confirms that sclerochronology can provide long-term and highly resolved records that are not obtainable by standard monitoring techniques. While elemental content alteration through diagenesis or biological control is a potential limitation of these methods, the applicability and advantages of sclerochronology in the field of biomonitoring are evident ([Schöne and Krause, 2016](#); [Steinhardt et al., 2016](#)). Another form of pollution is the anthropological input of nutrients into freshwater and coastal ecosystems, e.g., through agricultural runoff or wastewater, which fuels eutrophication [Q49, 50]. Sclerochronological studies on deep-water corals have shown that enrichment in skeletal ^{15}N is an indicator for terrestrial runoff (e.g., [Williams et al., 2007](#); [Prouty et al., 2014](#)). Similarly, bivalve shell $\delta^{15}\text{N}$ is increasingly used to assess human and animal waste input into waterways (e.g., [Black et al., 2017](#); [Thibault et al., 2020](#)). While this has been predominantly done via whole-shell analysis, time series $\delta^{15}\text{N}$ data have been published in other contexts ([Gillikin et al., 2017](#)), and the potential for sclerochronological studies is evident. Cultural eutrophication is relevant to many issues within public health and ecology, such as the concern for safe drinking water and the increasing development of hypoxic areas in oceans and lakes known as ‘dead zones’ ([Chislock et al., 2013](#); [Breitburg et al., 2018](#)). Previous studies highlight promising applications of sclerochronological methods

in this area, for example, shifts in Mn/Ca of cod otoliths have been used as a proxy for exposure to hypoxia in the Baltic Sea ([Limburg et al., 2011, 2015](#); [Limburg and Casini, 2018](#)). Given that they are (mostly) immobile and benthic bioaccumulators with worldwide distribution, bivalves might be particularly suitable to track the history of hypoxic and anoxic events (e.g., [Zhao et al., 2017c](#); [Murakami-Sugihara et al., 2019](#)). As dead zones are rapidly increasing worldwide, it would be very beneficial to develop sclerochronological applications to assess and monitor these phenomena.

3.3. Cutting edge sclerochronology

This section presents questions that were not ranked in the leading 50, but were nevertheless highlighted by the expert panels as potentially groundbreaking. These questions may have been downgraded because they are very specific to particular archives, or because they were perceived as high-risk with little chance of success. After discussion, they have been resurrected by the expert panels because they were thought to have the potential to broaden the horizons of sclerochronology, leading to highly novel applications.

A. Can we use material within the growth line to infer conditions outside the main growing season?

This question alludes to archives with varying growth rates, in particular, bivalves from mid- and high-latitude locations. Growth lines are formed during times of slow growth, often during autumn and winter months (see [Killam and Clapham, 2018](#)). Thus, growth lines potentially contain information on the environment outside of the main growing season. Attempts to use growth lines to study environmental and climate changes are absent within the literature, most likely due to the analytical challenges that arise from the fine scale of growth lines ([Shirai et al., 2014](#)). However, some research has demonstrated potential for shell Sr/Ca and Mg/Ca proxies to reveal environmental information near the growth line (e.g., [Schöne et al., 2013](#)), despite general challenges in elemental ratio proxies within bivalve shells (see Topic “3.1.6. Archive-specific research”). Thus, analytical techniques making use of growth lines in bivalves is a promising avenue for studying previously inaccessible seasons, as well as improving our understanding of biomineralization processes.

B. What approaches can we use to identify coeval shells for deep-time geological settings that will enable us to construct multicentennial crossmatched chronologies?

Crossdating allows construction of well-replicated, annually resolved, and exactly dated records that can span multiple centuries to millennia ([Black et al., 2016](#)). Also, software tools such as Shellcorr ([Scourse et al., 2006](#)) or CDendro (Cybis Dendrochronology) can assist in pattern matching among dead-collected specimens which are known to be roughly coeval. However, construction of deep-time chronologies by crossmatching shells requires the identification of fossil specimens with overlapping lifespans from accumulations and lags which may cover many thousands of years. The antiquity of these shells precludes rangefinder radiometric dating, and other methods need to be developed to identify coeval specimens in cases such as this. While stratigraphy or spatial proximity among specimens can provide time constraints in certain cases, these factors alone are not sufficient to guarantee contemporaneity — modern shell lags, for example, can contain specimens separated in time by several thousand years ([Butler et al., 2010](#)). The likelihood of contemporaneity might increase when fossils can be interpreted to have been rapidly buried in life position (e.g., bivalves with both valves still intact; [Lockwood and Work, 2006](#)) or in a calcified reef formation ([Greer et al., 2006](#); [Wu et al., 2017](#)). A taphonomic indicator of rapid burial is good preservation of the surface structure with no signs of grazing, boring, or postmortem microboring ([Vogel, 2000](#);

Lescinsky et al., 2011, and references therein). However, even where rapid burial in life position is assumed, time-averaging effects may complicate the search for coeval specimens; this is especially true for shelly fossils in siliciclastic environments, and to a lesser degree also for shells in carbonate sediments as well as for reef coral assemblages (Kidwell et al., 2005; Edinger et al., 2007). Developing sampling strategies in the fossil record could represent a breakthrough that enables us to investigate change at high resolution over extended periods in deep-time settings.

- C. What can sclerochronological records tell us about which seasons are represented by non-sclerochronological estimates of paleo-seawater temperature (e.g., from sediment core proxies)?

Temperature estimates based on sediment core proxies (i.e., marine microorganisms and their organic residues) are an important source of paleoceanographic data. Yet, interpretation of the climate signal is often complicated by the uncertainties related to the life cycle of biological sediment core archives (i.e., planktonic and benthic microorganisms). Ecological factors, such as the length and timing of the growing season, determine whether paleo-seawater temperature estimates represent an annual mean or an average over a certain season (typically summer). Sediment traps and core-top analysis in combination with instrumental data are typically used to calibrate proxy-based reconstructions. However, this approach cannot account for changes in the growing season through time and is not applicable to extinct species. The advantage of sclerochronology for providing seasonally resolved paleotemperature records opens a possibility for comparison with contemporary non-sclerochronological estimates (e.g., de Winter et al., 2018), thus providing the means for proxy calibration. The feasibility of this approach depends on our understanding of how sclerochronological records can be used to reconstruct past open-ocean conditions [Q39] typically reflected in sediment core records. To enable calibration of proxies based on planktonic species, sclerochronological records that reflect mixed-layer temperature dynamics have to be developed. Identification of suitable sclerochronological archives and development of new methods to solve these issues can lead to a novel application of sclerochronology and improve our understanding of past climate.

- D. How can we use sclerochronology to investigate potential latitudinal gradients in the response of marine biota to climate change, in terms of species die-off or range shifts?

An understanding of how biotas respond to climate change, to possibly predict future extinctions and species range shifts, are among the key challenges of ecology (Sutherland et al., 2013). Most of the research on this topic focuses on abundance and species distribution data, which is a coarse metric of change (Rombouts et al., 2012). For the impact to be detectable in such data, organisms have to die, stop breeding, or shift their geographic distribution. Sclerochronology provides promising tools for analyzing population metrics from stationary (e.g., bivalves and corals) and/or mobile (e.g., fish) organisms to infer latitudinal shifts in optimal conditions and/or fisheries regime shifts in population demography (Neuheimer et al., 2011; Morrongiello and Thresher, 2015). Using sclerochronology to identify sublethal impacts of changing environmental conditions can help to recognize potential range shifts or fisheries productivity changes before distributional change has happened.

- E. Can fisheries management advice be improved by combining traditional stock assessment techniques (e.g., otolith aging) with machine learning?

Advances in software and analytical tools have had profound impacts in the environmental sciences (Fielding, 1999). One such advancement is in the field of 'machine learning', which is a transformative tool across

disciplines (Malde et al., 2019). Image recognition is a common application of machine learning within the natural sciences. Machine learning methods have been utilized for mollusk shell identification (Zhang et al., 2019) and geometric morphometric analysis of gastropods (Doyle et al., 2018). Furthermore, fisheries scientists have applied machine learning techniques to fish age assessment through otolith image analysis with some success (Dub et al., 2013; Moen et al., 2018). Combining machine learning with long-term historical datasets of sclerochronology images, from any species, and their associated growth and ages, could (1) automate aging, (2) reduce human error, (3) improve predictions of population growth responses, and (4) identify anomalies. Large-scale incorporation of open-source otolith image recognition software could improve stock management advice for commercially important species.

- F. What proportion of the whole ecosystem extent does the environmental DNA (eDNA) in bivalve shells capture and how can the eDNA be used to reconstruct ecosystem change?

Technological advances in molecular ecology have led to the utilization of environmental DNA (eDNA) to detect the presence of certain taxa within aquatic environments. From a physical environmental sample (e.g., water, soil, shell), molecular markers within fragments of available eDNA are amplified and compared against a database to identify the presence of target species within the environment (Ardura et al., 2015). This tool is particularly useful for ecosystem monitoring. For example, eDNA can allow for early detection of invasive species, and the identification of vulnerable species. Sclerochronology and genetics are not traditionally paired together, but recently, the carbonate biominerals of fossilized marine mollusks have been found to contain eDNA (Der Sarkissian et al., 2017). More studies are needed to verify how successful bivalve eDNA is at reconstructing the full range of species present within the entire ecosystem, and how eDNA results may vary depending on environmental and physiological conditions. The ability to identify the timing of presence and absence of taxa within sclerochronological archives could revolutionize our understanding of ecosystem shifts during changes in global climate.

- G. How can possible effects of early human harvesting be separated from natural variability in marine fauna to better assess how changes in resources affected hunter-fisher-gatherers?

Material from middens can be used to reconstruct past climatic and environmental change as well as human behavior (see also Q40-43). Fish and shellfish have been an important source of food throughout human history. The impact of fishing, harvesting, and maricultures on marine ecosystems can be traced back to the early Holocene and beyond — for instance, in Europe, records of shellfish harvesting date back to over 450,000 years ago (Bailey and Milner, 2008). However, natural variability also influences ecosystem change, and not all environmental changes that we see in the paleorecord are anthropogenic. Thus, anthropogenic influences such as overharvesting or ancient maricultural constructions have to be separated from natural variability of ecosystems and populations. For example, regional differences in shellfish harvesting practices and underlying environmental and historical factors can be investigated by combining growth and $\delta^{18}\text{O}$ data in shells collected from archaeological shell middens in different environmental settings (Burchell et al., 2013b). Disentangling natural and anthropogenic signals recorded in middens using sclerochronological tools, would allow us to study not only how humans have impacted the environment, but in turn how environmental change has affected human societies and food security through time.

- H. How can data from sclerochronology be used to inform us about land claims by indigenous people?

Indigenous land rights are tightly linked with physical and economic safety, as well as mental and emotional well-being, and therefore of fundamental importance for the self-determination of indigenous groups. As discussed above, many studies have used midden material to study past human-environmental interactions. Season-of-capture studies provide insight into residential mobility and sedentism of hunter-fisher-gatherers in the past (e.g., Burchell et al., 2013a). In the Americas and Australia, as well as some regions in Asia and Africa, midden archives are almost always on indigenous land. Therefore, scientists should acknowledge that these archives are the result of indigenous labor, and seek conversation and exchange with indigenous groups. Given that middens can provide continuous records that span the last 11,000 years of human history (Tonello et al., 2019), sclerochronological techniques could be applied to prove long-term land use of indigenous groups based on ancient middens. Further applications and research questions should be developed by actively involving indigenous descent groups in the scientific process (see, e.g., Kaiser et al., 2019).

4. Discussion

4.1. Priority research questions

This collaborative project was conducted to reflect on the state-of-the-art in sclerochronology and to identify existing challenges and possible future developments. The questions identified herein as fundamental and priority to the field (see Supplementary Material 2 for the complete list) represent a snapshot in time describing the potential and challenges of sclerochronological research as perceived by a group of leading experts and community members following the 5th ISC. The link to the 5th ISC was an advantage for this endeavor, as the conference provided an overview of contemporary research results, facilitating the compilation of well-informed contributions.

We intended to highlight research addressing promising applications of sclerochronology, as well as gaps in our understanding of archives and methods used in the field. The questions presented and discussed in this paper reveal significant knowledge gaps in our understanding of biomineralization processes [Q1-4, 22], mechanisms driving the growth of skeletal structures [Q5-8, 23, 25], and challenges in geochemical proxy interpretations [Q16-19, 21]. They also emphasize the need to identify common standards for data management and analysis [Q9-15] and introduce fundamental questions related to the use of specific archives, such as fish otoliths [Q24, 25] and bivalves [Q21-23]. Even though many of the submitted questions suggest that current tools and methods require further development, many existing techniques are sufficient to address important scientific questions in other fields. Our results highlight the potential for applying sclerochronological data and methods to answer long-standing questions in other research fields, such as climate sciences [Q26-39] and (paleo-)ecology [Q40-50].

Identification of research priorities entails certain trade-offs. The feasibility of approaches and their potential impacts are often difficult to foresee and evaluate. This complicates identification of the research directions that are not-yet established and thus truly lie on the horizon. Although our list of questions provides an overview of possibilities and challenges in sclerochronology, considering all the limitations of this study, it should be treated with some degree of caution. The questions selected by the community are largely reflective of ongoing research. As the 5th ISC conference showed, sclerochronological tools have been used in many research fields. Nevertheless, the most common applications to date are related to climate and environmental sciences, which is reflected in both the research presented at the ISC and the questions submitted to the survey that forms the basis of this paper. However, these are not the limits of sclerochronology, and applications to social and anthropological sciences, for example, are feasible and promising [e.g., Questions G, H]. Thus, the range of questions presented here by no means portrays the full diversity and potential of the field and the list of questions will require further updates as the field develops.

4.2. Future outlook

The discussion arising from this project emphasizes important issues relevant to the future of the field. While previous advances in sample preparation, observation and analysis have moved the field significantly forward and enabled new and improved applications of sclerochronological research, our results show there is still room for technical and methodological development in the field (Section 3.1), as support and motivation for applications in the field (Section 3.2). The survey participants identified common standards for data management and analysis as high priority. Similarly, identification of common terminology within and across different branches of sclerochronological research would significantly improve communication and potential collaboration. A further promising avenue for sclerochronology is future collaboration with researchers utilizing similar archives (e.g., calcium phosphate biominerals), or researchers in the social sciences. Relatively low response rates and the bias towards certain disciplines and archives in the framework of the ISC indicate that the field requires consolidation and more collaborative work. Collaboration across different disciplines would undoubtedly widen the profile of sclerochronology among the scientific community. As the field matures, it should become possible to combine climatological, ecological, biogeochemical and archaeological applications of sclerochronology to create an integrated sclerochronological approach to the study of the Earth system, covering the physical and living systems and extending to human cultural history.

5. Conclusions

This is the first effort to identify priority and fundamental research questions in sclerochronology via horizon scanning. As the field grows and advances, we recognize that research priorities will have to be reassessed. The list of questions presented and discussed in this paper contains the highest-ranked research priorities and fundamental questions, as identified in a community-based survey. While this list should not be considered definitive, we hope that the results of this project will stimulate discussion and serve as a stepping stone to future collaborations and groundbreaking research.

CRedit authorship contribution statement

Tamara Trofimova: Project administration, Conceptualization, Methodology, Writing - original draft, Formal analysis. **Stella J. Alexandroff:** Conceptualization, Methodology, Writing - original draft, Formal analysis. **Madelyn J. Mette:** Conceptualization, Methodology, Writing - original draft, Formal analysis. **Elizabeth Tray:** Conceptualization, Methodology, Writing - original draft, Formal analysis. **Paul G. Butler:** Writing - review & editing, Supervision. **Steven E. Campana:** Writing - review & editing, Supervision. **Elizabeth M. Harper:** Writing - review & editing, Supervision. **Andrew L.A. Johnson:** Writing - review & editing, Supervision. **John R. Morrongiello:** Writing - review & editing, Supervision. **Melita Peharda:** Writing - review & editing, Supervision. **Bernd R. Schöne:** Writing - review & editing, Supervision. **Carin Andersson:** Writing - review & editing. **C. Fred T. Andrus:** Writing - review & editing. **Bryan A. Black:** Writing - review & editing. **Meghan Burchell:** Writing - review & editing. **Michael L. Carroll:** Writing - review & editing. **Kristine L. DeLong:** Writing - review & editing. **Bronwyn M. Gillanders:** Writing - review & editing. **Peter Grønkjær:** Writing - review & editing. **Daniel Killam:** Writing - review & editing. **Amy L. Prendergast:** Writing - review & editing. **David J. Reynolds:** Writing - review & editing. **James D. Scourse:** Writing - review & editing. **Kotaro Shirai:** Writing - review & editing. **Julien Thébault:** Writing - review & editing. **Clive Trueman:** Writing - review & editing. **Niels de Winter:** Writing - review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This project was initiated following discussions and suggestions within the PAGES ECN (Past Global Changes Early-Career Network), and this paper is therefore a product of PAGES and the PAGES ECN. We would like to thank the organizing committee of the 5th International Sclerochronology Conference, which provided a framework and platform to initiate the survey. We thank Stefania Milano for contribution to the initial list of questions. Many thanks to all survey participants without whose responses this manuscript would not have been possible. We thank three anonymous reviewers and the managing editor, Steve Mitchell, for significant comments which improved the manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecss.2020.106977>.

References

- Adams, L.A., 1940. Some characteristic otoliths of American ostariophys. *J. Morphol.* 66 (3), 497–527. <https://doi.org/10.1002/jmor.1050660307>.
- Andrews, A.H., Gobalet, K.W., Jones, T.L., 2003. Reliability assessment of season-of-capture determination from archaeological otoliths. *Bull. South Calif. Acad. Sci.* 102 (2), 66–78. https://digitalcommons.calpoly.edu/ssci_fac/14/.
- Andrus, C.F.T., 2011. Shell midden sclerochronology. *Quat. Sci. Rev.* 30 (21–22), 2892–2905. <https://doi.org/10.1016/j.quascirev.2011.07.016>.
- Ardura, A., Zaiko, A., Martínez, J.L., Samuiloviene, A., Semenova, A., García-Vázquez, E., 2015. eDNA and specific primers for early detection of invasive species - a case study on the bivalve *Rangia cuneata*, currently spreading in Europe. *Mar. Environ. Res.* 112 (B), 48–55. <https://doi.org/10.1016/j.marenvres.2015.09.013>.
- Atsawararant, K., Comas-Bru, L., Amirnezhad Mozhdzhi, S., Deininger, M., Harrison, S.P., Baker, A., Boyd Kaushal, N., Ahmad, S.M., Ait Brahim, Y., Arienzo, M., Bajo, P., Braun, K., Burstyn, Y., Chawchai, S., Duan, W., Hatvani, I.G., Hu, J., Kern, Z., Labuhn, L., Lachniet, M., Lechleiter, F.A., Lorrey, A., Pérez-Mejías, C., Pickering, R., Scroton, N., Sisal Working Group Members, 2018. The SISAL database: a global resource to document oxygen and carbon isotope records from speleothems. *Earth Syst. Sci. Data* 10, 1687–1713. <https://doi.org/10.5194/essd-2018-17>.
- Bailey, G., Milner, N.J., 2008. Molluscan archives from European prehistory. In: Antczak, A., Cipriani, R. (Eds.), *Early Human Impact on Megamolluscs, British Archaeological Series, International Series, vol. 1865*, pp. 111–134. Oxford, Archaeopress.
- Barrow, J., Ford, J., Day, R., Morrongiello, J., 2018. Environmental drivers of growth and predicted effects of climate change on a commercially important fish, *Platycephalus laevigatus*. *Mar. Ecol. Prog. Ser.* 598, 201–212. <https://doi.org/10.3354/meps12234>.
- Beck, J.W., Edwards, R.L., Ito, E., Taylor, F.W., Recy, J., Rougerie, F., Joanon, P., Henin, C., 1992. Sea-surface temperature from coral skeletal strontium/calcium ratios. *Science* 257 (5070), 644–647. <https://doi.org/10.1126/science.257.5070.644>.
- Black, B.A., Copenheaver, C.A., Frank, D.C., Stuckey, M.J., Kormanyos, R.E., 2009. Multi-proxy reconstructions of northeastern Pacific sea surface temperature data from trees and Pacific geoduck. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 278, 40–47. <https://doi.org/10.1016/j.palaeo.2009.04.010>.
- Black, B.A., Sydeman, W.J., Frank, D.C., Griffin, D., Stahle, D.W., García-Reyes, M., Rykaczewski, R.R., Bograd, S.J., Peterson, W.T., 2014. Six centuries of variability and extremes in a coupled marine-terrestrial ecosystem. *Science* 345 (6203), 1498–1502. <https://doi.org/10.1126/science.1253209>.
- Black, B.A., Griffin, D., van der Sleen, P., Wanamaker, A.D.J., Speer, J.H., Frank, D.C., Stahle, D.W., Pederson, N., Copenheaver, C.A., Trouet, V., Griffin, S., Gillanders, B.M., 2016. The value of crossdating to retain high-frequency variability, climate signals, and extreme events in environmental proxies. *Global Change Biol.* 22 (7), 2582–2595. <https://doi.org/10.1111/gcb.13256>.
- Black, H.D., Andrus, C.F.T., Lambert, W.J., Rick, T.C., Gillikin, D.P., 2017. $\delta^{15}\text{N}$ values in *Crassostrea virginica* shells provides early direct evidence for nitrogen loading to Chesapeake Bay. *Sci. Rep.* 7, 442–541. <https://doi.org/10.1038/srep44241>.
- Black, B.A., van der Sleen, P., Di Lorenzo, E., Griffin, D., Sydeman, W.J., Dunham, J.B., Rykaczewski, R.R., García-Reyes, M., Safeeq, M., Arismendi, I., Bograd, S.J., 2018. Rising synchrony controls western North American ecosystems. *Global Change Biol.* 24, 2305–2314. <https://doi.org/10.1111/gcb.14128>.
- Black, B.A., Andersson, C., Butler, P.G., Carroll, M.L., DeLong, K.L., Reynolds, D.J., Schöne, B.R., Scourse, J.D., van der Sleen, P., Wanamaker Jr., A.D., Witbaard, R., 2019. The revolution of crossdating in marine palaeoecology and palaeoclimatology. *Biol. Lett.* 15 <https://doi.org/10.1098/rsbl.2018.0665>.
- Böhm, F., Joachimski, M.M., Lehnert, H., Morgenroth, G., Kretschmer, W., Vacelet, J., Dullo, W.C., 1996. Carbon isotope records from extant Caribbean and South Pacific sponges: evolution of $\delta^{13}\text{C}$ in surface water DIC. *Earth Planet Sci. Lett.* 139 (1), 291–303. [https://doi.org/10.1016/0012-821X\(96\)00006-4](https://doi.org/10.1016/0012-821X(96)00006-4).
- Boiseau, M., Juillet-Leclerc, A., 1997. H₂O₂ treatment of recent coral aragonite: oxygen and carbon isotopic implications. *Chem. Geol.* 143, 171–180. [https://doi.org/10.1016/S0009-2541\(97\)00112-5](https://doi.org/10.1016/S0009-2541(97)00112-5).
- Bonitz, F.G.W., Andersson, C., Trofimova, T., Hátún, H., 2017. Links between phytoplankton dynamics and shell growth of *Arctica islandica* on the Faroe Shelf. *J. Mar. Syst.* 179, 72–87. <https://doi.org/10.1016/j.jmarsys.2017.11.005>.
- Bougeois, L., De Raféls, M., Reichert, G.-J., De Nooijer, L.J., Nicollin, F., Dupont-Nivet, G., 2014. A high resolution study of trace elements and stable isotopes in oyster shells to estimate Central Asian Middle Eocene seasonality. *Chem. Geol.* 363, 200–212. <https://doi.org/10.1016/j.chemgeo.2013.10.037>.
- Breitbart, D., Levin, L.A., Oshlies, A., Grégoire, M., Chavez, F.P., Conley, D.J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee, K., Jacinto, G.S., Limburg, K.E., Montes, I., Naqvi, S.W.A., Pitcher, G.C., Rabalais, N.N., Roman, M.R., Rose, K.A., Seibel, B.A., Telszewski, M., Yasuhara, M., Zhang, J., 2018. Declining oxygen in the global ocean and coastal waters. *Science* 359 (6374), eam7240. <https://doi.org/10.1126/science.aam7240>.
- Buddemeier, R.W., Maragos, J.E., Knutson, D.W., 1974. Radiographic studies of reef coral exoskeletons: rates and patterns of coral growth. *J. Exp. Mar. Biol. Ecol.* 14 (2), 179–199. [https://doi.org/10.1016/0022-0981\(74\)90024-0](https://doi.org/10.1016/0022-0981(74)90024-0).
- Burchell, M., Cannon, A., Hallmann, N., Schwarcz, H.P., Schöne, B.R., 2013a. Inter-site variability in the season of shellfish collection on the central coast of British Columbia. *J. Archaeol. Sci.* 40 (1), 626–636. <https://doi.org/10.1016/j.jas.2012.07.002>.
- Burchell, M., Hallmann, N., Martindale, A., Cannon, A., Schöne, B.R., 2013b. Seasonality and intensity of shellfish harvesting on the North coast of British Columbia. *J. I. Coast Archaeol.* 8, 152–169. <https://doi.org/10.1080/15564894.2013.787566>.
- Butler, P.G., Schöne, B.R., 2017. New research in the methods and applications of sclerochronology. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 465, 295–299. <https://doi.org/10.1016/j.palaeo.2016.11.013>.
- Butler, P.G., Scourse, J.D., Richardson, C.A., Wanamaker Jr., A.D., Bryant, C.L., Bennell, J.D., 2009. Continuous marine radiocarbon reservoir calibration and the 13C Suess effect in the Irish Sea: results from the first multi-centennial shell-based marine master chronology. *Earth Planet Sci. Lett.* 279 (3–4), 230–241. <https://doi.org/10.1016/j.epsl.2008.12.043>.
- Butler, P.G., Richardson, C.A., Scourse, J.D., Wanamaker Jr., A.D., Shammon, T.M., Bennell, J.D., 2010. Marine climate in the Irish Sea: analysis of a 489-year marine master chronology derived from growth increments in the shell of the clam *Arctica islandica*. *Quat. Sci. Rev.* 29 (13–14), 1614–1632. <https://doi.org/10.1016/j.quascirev.2009.07.010>.
- Butler, P.G., Freitas, P.S., Burchell, M., Chauvaud, L., 2019. Archaeology and Sclerochronology of marine bivalves. In: Smaal, A.C., Ferreira, J.G., Grant, J., Petersen, J.K., Strand, Ø. (Eds.), *Goods and Services of Marine Bivalves*. 413–444. Springer Open. <https://doi.org/10.1007/978-3-319-96776-9>.
- Cailliet, G., Andrews, A., Burton, E., Watters, D., Kline, D., Ferry-Graham, L., 2001. Age determination and validation studies of marine fishes: do deep-dwellers live longer? *Exp. Gerontol.* 36 (4–6), 739–764. [https://doi.org/10.1016/s0531-5565\(00\)00239-4](https://doi.org/10.1016/s0531-5565(00)00239-4).
- Campana, S.E., 1999. Chemistry and composition of fish otoliths: pathways, mechanisms and applications. *Mar. Ecol. Prog. Ser.* 188, 263–297. <https://doi.org/10.3354/meps188263>.
- Campana, S., 2005. Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *J. Fish. Biol.* 59 (2), 197–242. <https://doi.org/10.1111/j.1095-8649.2001.tb00127.x>.
- Campana, S.E., Thorrold, S.R., 2001. Otoliths, increments, and elements: keys to a comprehensive understanding of fish populations? *Can. J. Fish. Aquat. Sci.* 58, 30–38. <https://doi.org/10.1139/cjfas-58-1-30>.
- Carré, M., Azzoug, M., Zaharias, P., Camara, A., Cheddadi, R., Chevalier, M., Fiorillo, D., Gaye, A.T., Janicot, S., Khodri, M., Lazar, A., Lazareth, C.E., Mignot, J., Mitma García, N., Patris, N., Perrot, O., Wade, M., 2019. Modern drought conditions in western Sahel unprecedented in the past 1600 years. *Clim. Dynam.* 52, 1949–1964. <https://doi.org/10.1007/s00382-018-4311-3>.
- Carroll, M.L., Romanek, C., Paddock, L., 2006. The relationship between the hydrogen and oxygen isotopes of freshwater bivalve shells and their home streams. *Chem. Geol.* 234, 211–222. <https://doi.org/10.1016/j.chemgeo.2006.04.012>.
- Carter, J.G., 1980. Environmental and biological controls of bivalve shell mineralogy and microstructure. In: Rhoads, D.C., Lutz, R.A. (Eds.), *Skeletal Growth of Aquatic Organisms*. Plenum, New York.
- Chan, P., Halfar, J., Adey, W., Hetzinger, S., Zack, T., Moore, G.W.K., Hou, A., 2017. Multicentennial record of Labrador Sea primary productivity and sea-ice variability archived in coralline algal barium. *Nat. Commun.* 8 (1), 15543. <https://doi.org/10.1038/ncomms15543>.
- Checa, A., 2018. Physical and biological determinants of the fabrication of Molluscan shell microstructures. *Front. Marine Sci.* 5 (353) <https://doi.org/10.3389/fmars.2018.00353>.
- Checkley, D.M., Dickson, A.G., Takahashi, M., Radich, J.A., Eisenkolb, N., Asch, R., 2009. Elevated CO₂ enhances otolith growth in young fish. *Science* 324 (5935). <https://doi.org/10.1126/science.1169806>, 1683–1683.
- Chislock, M.F., Doster, E., Zitomer, R.A., Wilson, A.E., 2013. Eutrophication: causes, consequences, and controls in aquatic ecosystems. *Nat. Education Knowledge* 4 (4), 10.

- Clark II, G.R., 1974. Growth lines in invertebrate skeletons. *Annu. Rev. Earth Planet Sci.* 2, 77. <https://doi.org/10.1146/annurev.ea.02.050174.000453>.
- Clarke, L.M., Thorrold, S.R., Conover, D.O., 2010. Population differences in otolith chemistry have a genetic basis in *Menidia menidia*. *Can. J. Fish. Aquat. Sci.* 68 (1), 105–114. <https://doi.org/10.1139/F10-147>.
- Comboul, M., Emile-Geay, J., Evans, M.N., Mirnateghi, N., Cobb, K.M., Thompson, D.M., 2014. A probabilistic model of chronological errors in layer-counted climate proxies: applications to annually banded coral archives. *Clim. Past* 10, 825–841. <https://doi.org/10.5194/cp-10-825-2014>.
- Cook, E.R., Woodhouse, C.A., Eakin, C.M., Meko, D.M., Stahle, D.W., 2004. Long-term aridity changes in the western United States. *Science* 306, 1015–1018. <https://doi.org/10.1126/science.1102586>.
- Cornwall, C.E., Comeau, S., DeCarlo, T.M., Moore, B., D'Alexis, Q., McCulloch, M.T., 2018. Resistance of corals and coralline algae to ocean acidification: physiological control of calcification under natural pH variability. *Proc. Biol. Sci.* 285 (1884) <https://doi.org/10.1098/rspb.2018.1168>.
- Corrège, T., 2006. Sea surface temperature and salinity reconstruction from coral geochemical tracers. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 232, 408–428. <https://doi.org/10.1016/j.palaeo.2005.10.014>.
- Coulson, P.G., Black, B.A., Potter, I.C., Hall, N.G., 2014. Sclerochronological studies reveal that patterns of otolith growth of adults of two co-occurring species of *Platycephalidae* are synchronised by water temperature variations. *Marine Biol.* 161 (2), 383–393. <https://doi.org/10.1007/s00227-013-2343-0>.
- Coutts, P.J.F., 1970. Bivalve growth patterning as a method for seasonal dating in archaeology. *Nature* 226, 874.
- Cross, E.L., Harper, E.M., Peck, L.S., 2019. Thicker shells compensate extensive dissolution in brachiopods under future ocean acidification. *Environ. Sci. Technol.* 53 (9), 5016–5026. <https://doi.org/10.1021/acs.est.9b00714>.
- Crowley, T.J., Quinn, T.M., Hyde, W.T., 1999. Validation of coral temperature calibrations. *Paleoceanography* 14, 605–615. <https://doi.org/10.1029/1999PA000032>.
- Dassié, E.P., Lemley, G.M., Linsley, B.K., 2013. The Suess effect in Fiji coral $\delta^{13}\text{C}$ and its potential as a tracer of anthropogenic CO_2 uptake. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 370, 30–40. <https://doi.org/10.1016/j.palaeo.2012.11.012>.
- Dassié, E.P., Linsley, B.K., Corrège, T., Wu, H.C., Lemley, G.M., Howe, S., Cabioch, G., 2014. A Fiji multi-coral $\delta^{18}\text{O}$ composite approach to obtaining a more accurate reconstruction of the last two-centuries of the ocean-climate variability in the South Pacific Convergence Zone region. *Paleoceanography Paleoclimatol.* 29 (12), 1196–1213. <https://doi.org/10.1002/2013PA002591>.
- Dassié, E., DeLong, K., Williams, B., Abram, N., Brenner, L., Brahm, C., Cobb, K.M., Corrège, T., Dissard, D., Emile-Geay, J., Evangelista, H., Evans, M.N., Farmer, J., Felis, T., Gagan, M., Gillikin, D.P., Goodkin, N., Khodri, M., Lavagnino, A.C., LaVigne, M., Lazareth, C., Linsley, B., Lough, J., McGregor, H., Nurhati, I.S., Ouellette, G., Raymo, M., Rosenheim, B., Sandstrom, M., Schöne, B.R., Sifeddine, A., Stevenson, S., Thompson, D.M., Waite, A., Wanamaker, A., Wu, H., 2017. Saving our marine archives. *Earth and Space Sci.* News 98. <https://doi.org/10.1029/2017EO068159> 2017.
- Davenport, C.B., 1938. Growth lines in fossil pectens as indicators of past climates. *J. Paleontol.* 12 (5), 514–515.
- DeLong, K.L., Quinn, T.M., Taylor, F.W., 2007. Reconstructing twentieth-century sea surface temperature variability in the southwest Pacific: a replication study using multiple coral Sr/Ca records from New Caledonia. *Paleoceanography* 22, PA4212. <https://doi.org/10.1029/2007PA001444>.
- DeLong, K.L., Flannery, J.A., Maupin, C.R., Poore, R.Z., Quinn, T.M., 2011. A coral Sr/Ca calibration and replication study of two massive corals from the Gulf of Mexico. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 307, 1–4. <https://doi.org/10.1016/j.palaeo.2011.05.005>.
- DeLong, K.L., Maupin, C.R., Flannery, J.A., Quinn, T.M., Shen, C.-C., 2016. Refining temperature reconstructions with the Atlantic coral *Siderastrea siderea*. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 462, 1–15. <https://doi.org/10.1016/j.palaeo.2016.08.028>.
- Denechaud, C., Smolinski, S., Geffen, A.J., Godiksen, J.A., 2020. Long-term temporal stability of Northeast Arctic cod (*Gadus morhua*) otolith morphology. *ICES (Int. Counc. Explor. Sea) J. Mar. Sci.* 77 (3), 1043–1054. <https://doi.org/10.1093/icesjms/fsz259>.
- Devereux, I., 1967. Temperature measurements from oxygen isotope ratios of fish otoliths. *Science* 155, 1683–1685. <https://doi.org/10.1126/science.155.3770.1684>.
- Doyle, D., Gammell, M.P., Nash, R., 2018. Morphometric methods for the analysis and classification of gastropods: a comparison using *Littorina littorea*. *J. Molluscan Stud.* 84 (2), 190–197. <https://doi.org/10.1093/mollus/eyy010>.
- Druffel, E.R.M., Benavides, L.M., 1986. Input of excess CO_2 to the surface ocean based on $^{13}\text{C}/^{12}\text{C}$ ratios in a banded Jamaican sclerosponge. *Nature* 321 (6065), 58–61. <https://doi.org/10.1038/321058a0>.
- Druffel, E.R.M., Griffin, S., 1993. Large variations of surface ocean radiocarbon: evidence of circulation changes in the southwestern Pacific. *J. Geophys. Res.: Oceans* 98 (C11), 20249–20259. <https://doi.org/10.1029/93jc02113>.
- Dub, J.D., Redman, R.A., Wahl, D.H., Czesny, S.J., 2013. Utilizing random forest analysis with otolith mass and total fish length to obtain rapid and objective estimates of fish age. *Can. J. Fish. Aquat. Sci.* 70, 1396–1401. <https://doi.org/10.1139/cjfas-2013-0169>.
- Dunbar, R.B., Wefer, G., 1984. Stable isotope fractionation in benthic foraminifera from the Peruvian continental margin. *Mar. Geol.* 59, 215–225. [https://doi.org/10.1016/0025-3227\(84\)90094-X](https://doi.org/10.1016/0025-3227(84)90094-X).
- Eakin, M.C., Grotto, A.G., 2006. Coral reef records of past climatic change. In: Phinney, J.T., et al. (Eds.), *Coral Reefs and Climate Change: Science and Management, Coastal Estuarine Stud.*, vol. 61. AGU, Washington, D. C., pp. 33–54. <https://doi.org/10.1029/61CE04>.
- Edinger, E.N., Burr, G.S., Pandolfi, J.M., Ortiz, J.C., 2007. Age accuracy and resolution of Quaternary corals used as proxies for sea level. *Earth Planet Sci. Lett.* 253, 37–49. <https://doi.org/10.1016/j.epsl.2006.10.014>.
- Eiler, J.M., 2007. “Clumped-isotope” geochemistry — the study of naturally-occurring, multiply-substituted isotopologues. *Earth Planet Sci. Lett.* 262, 309–327. <https://doi.org/10.1016/j.epsl.2007.08.020>.
- Eiler, J.M., 2011. Paleoclimate reconstruction using carbonate clumped isotope thermometry. *Quat. Sci. Rev.* 30, 3575–3588. <https://doi.org/10.1016/j.quascirev.2011.09.001>.
- Elliot, M., deMenocal, P.B., Linsley, B.K., Howe, S.S., 2003. Environmental controls on the stable isotopic composition of *Mercenaria mercenaria*: potential application to paleoenvironmental studies. *G-cubed* 4 (7). <https://doi.org/10.1029/2002GC000425>.
- Elliot, M., Welsh, K., Chilcott, C., McCulloch, M., Chappell, J., Ayling, B., 2009. Profiles of trace elements and stable isotopes derived from giant long-lived *Tridacna gigas* bivalves: potential applications in paleoclimate studies. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 280, 132–142. <https://doi.org/10.1016/j.palaeo.2009.06.007>.
- Emile-Geay, J., McKay, N.P., Kaufman, D.S., von Gunten, L., Wang, J.G., Anchukaitis, K. J., Abram, N.J., Pages2k Consortium, 2017. A global multiproxy database for temperature reconstructions of the Common Era. *Scientific Data* 4, 170088. <https://doi.org/10.1038/sdata.2017.88>.
- Epstein, S., Buchsbaum, R., Lowenstam, A., Urey, H.C., 1953. Revised carbonate-water isotopic temperature scale. *Geol. Soc. Am. Bull.* 64, 1315–1326. [https://doi.org/10.1130/0016-7606\(1953\)64\[1315:RCITS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1953)64[1315:RCITS]2.0.CO;2).
- Erlandson, J.M., 2001. The archaeology of aquatic adaptations: paradigms for a new millennium. *J. Archaeol. Res.* 9 (4), 287–350. <https://doi.org/10.1023/A:1013062712695>.
- Evans, J.W., 1972. Tidal growth increments in the cockle *Clinocardium nuttalli*. *Science* 176, 416–417.
- Ezgeta-Balic, D., Peharda, M., Richardson, C.A., Kuzmanic, M., Vrgoc, N., Isajlovic, I., 2011. Age, growth, and population structure of the smooth clam *Callista chione* in the eastern Adriatic Sea. *Helgol. Mar. Res.* 65, 457–465. <https://doi.org/10.1007/s10152-010-0235-y>.
- Fablet, R., Pecquerie, L., de Pontual, H., Hoie, H., Millner, R., Mosegaard, H., Koojiman, S.A.L.M., 2011. Shedding light on fish otolith biomineralization using a bioenergetic approach. *PLoS One* 6, e27055. <https://doi.org/10.1371/journal.pone.0027055>.
- Fiebig, J., Bajnai, D., Löffler, N., Methner, K., Krsnik, E., Mulch, A., Hofmann, S., 2019. Combined high-precision $\Delta 48$ and $\Delta 47$ analysis of carbonates. *Chem. Geol.* 522, 186–191. <https://doi.org/10.1016/j.chemgeo.2019.05.019>.
- Fielding, A.H. (Ed.), 1999. *Machine Learning Methods for Ecological Applications*. <https://doi.org/10.1007/978-1-4615-5289-5>.
- Finney, B.P., Alheit, J., Emeis, K.-C., Field, D.B., Gutiérrez, D., Struck, U., 2010. Paleoenvironmental studies on variability in marine fish populations: a long-term perspective on the impacts of climatic change on marine ecosystems. *J. Mar. Syst.* 79, 316–326. <https://doi.org/10.1016/j.jmarsys.2008.12.010>.
- Fitzer, S.C., Phoenix, V.R., Cusack, M., Kamenos, N.A., 2014. Ocean acidification impacts mussel control on biomineralisation. *Sci. Rep.* 4 (1), 6218. <https://doi.org/10.1038/srep06218>.
- Ford, H.L., Schellenberg, S.A., Becker, B.J., Deutschman, D.L., Dyck, K.A., Koch, P.L., 2010. Evaluating the skeletal chemistry of *Mytilus californianus* as a temperature proxy: effects of microenvironment and ontogeny. *Paleoceanography* 25, 1–14. <https://doi.org/10.1029/2008PA001677>.
- Foster, G.L., Rae, J.W.B., 2016. Reconstructing ocean pH with boron isotopes in foraminifera. *Annu. Rev. Earth Planet Sci.* 44 (1), 207–237. <https://doi.org/10.1146/annurev-earth-060115-012226>.
- Foster, L.C., Allison, N., Finch, A.A., Andersson, C., Ninnemann, U.S., 2009. Controls on ^{18}O and ^{13}C profiles within the aragonite bivalve *Arctica islandica*. *Holocene* 19, 549–558. <https://doi.org/10.1177/0959683609104028>.
- Fritts, H.C., Blasing, T.J., Hayden, B.P., Kutzbach, J.E., 1971. Multivariate techniques for specifying tree-growth and climate relationships and for reconstructing anomalies in paleoclimate. *J. Appl. Meteorol.* 10 (5), 845–864. [https://doi.org/10.1175/1520-0450\(1971\)010<0845:Mtfstg>2.0.Co;2](https://doi.org/10.1175/1520-0450(1971)010<0845:Mtfstg>2.0.Co;2).
- Füllenbach, C.S., Schöne, B.R., Branscheid, R., 2014. Microstructures in shells of the freshwater gastropod *Viviparus viviparus*: a potential sensor for temperature change? *Acta Biomater.* 10 (9), 3911–3921. <https://doi.org/10.1016/j.actbio.2014.03.030>.
- Gagan, M.K., Ayliffe, L.K., Hopley, D., Cali, J.A., Mortimer, G.E., Chappell, J., McCulloch, M.T., Head, M.J., 1998. Temperature and surface-ocean water balance of the mid-Holocene tropical western Pacific. *Science* 279, 1014–1018. <https://doi.org/10.1126/science.279.5353.1014>.
- Gagan, M.K., Dunbar, G.B., Suzuki, A., 2012. The effect of skeletal mass accumulation in *Porites* on coral Sr/Ca and $\delta^{18}\text{O}$ paleothermometry. *Paleoceanography* 27, 1–16. <https://doi.org/10.1029/2011PA002215>.
- Geffen, A.J., Hoie, H., Folkvord, A., Hufthammer, A.K., Andersson, C., Ninnemann, U., Pedersen, R.B., Nedreaas, K., 2011. High-latitude climate variability and its effect on fisheries resources as revealed by fossil cod otoliths. *ICES (Int. Counc. Explor. Sea) J. Mar. Sci.* 68 (6), 1081–1089. <https://doi.org/10.1093/icesjms/fsr017>.
- Gillanders, B.M., Izzo, C., Doubleday, Z.A., Ye, Q., 2015. Partial migration: growth varies between resident and migratory fish. *Biol. Lett.* 11, 20140850. <https://doi.org/10.1098/rsbl.2014.0850>.
- Gillikin, D.P., Dehairs, F., Baeyens, W., Navez, J., Lorrain, A., Andre, L., 2005. Inter- and intra-annual variations of Pb/Ca ratios in clam shells (*Mercenaria mercenaria*): a record of anthropogenic lead pollution? *Mar. Pollut. Bull.* 50, 1530–1540. <https://doi.org/10.1016/j.marpolbul.2005.06.020>.

- Gillikin, D.P., Lorrain, A., Meng, L., Dehairs, F., 2007. A large metabolic carbon contribution to the $\delta^{13}\text{C}$ record in marine aragonitic bivalve shells. *Geochem. Cosmochim. Acta* 71 (12), 2936–2946. <https://doi.org/10.1016/j.gca.2007.04.003>.
- Gillikin, D.P., Lorrain, A., Jolivet, A., Kelemen, Z., Chauvaud, L., Bouillon, S., 2017. High-resolution nitrogen stable isotope sclerochronology of bivalve shell carbonate-bound organics. *Geochem. Cosmochim. Acta* 200, 55–66. <https://doi.org/10.1016/j.gca.2016.12.008>.
- Gillikin, D.P., Wanamaker Jr., A.D., Andrus, C.F.T., 2019. Chemical sclerochronology. *Chem. Geol.* 526, 1–6. <https://doi.org/10.1016/j.chemgeo.2019.06.016>.
- Goodwin, D.H., Schöne, B.R., Dettman, D.L., 2003. Resolution and fidelity of oxygen isotopes as paleotemperature proxies in bivalve mollusk shells: models and observations. *Palaios* 18, 110–125. [https://doi.org/10.1669/0883-1351\(2003\)18<110:RAFOOL>2.0.CO;2](https://doi.org/10.1669/0883-1351(2003)18<110:RAFOOL>2.0.CO;2).
- Goosse, H., 2016. An additional step towards comprehensive paleoclimatic reanalyses. *J. Adv. Model. Earth Syst.* 8, 1501–1503. <https://doi.org/10.1002/2016MS000739>.
- Grammer, G.L., Morrongiello, J.R., Izzo, C., Hawthorne, P.J., Middleton, J.F., Gillanders, B.M., 2017. Coupling biogeochemical tracers with fish growth reveals physiological and environmental controls on otolith chemistry. *Ecol. Monogr.* 87, 487–507. <https://doi.org/10.1002/ecm.1264>.
- Greer, L., Swart, P.K., 2006. Decadal cyclicity of regional mid-Holocene precipitation: evidence from Dominican coral proxies. *Paleoceanography* 21. <https://doi.org/10.1029/2005PA001166>.
- Gröcke, D.R., Gillikin, D.P., 2008. Advances in mollusk sclerochronology and sclerochemistry: tools for understanding climate and environment. *Geo Mar. Lett.* 28 (5–6), 265–268. <https://doi.org/10.1007/s00367-008-0108-4>.
- Grønkvær, P., 2016. Otoliths as individual indicators - a reappraisal of the link between fish physiology and otolith characteristics. *Mar. Freshw. Res.* 67, 881–888. <https://doi.org/10.1017/MF15155>.
- Grossman, E.L., Ku, T.-L., 1986. Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects. *Chem. Geol.: Isot. Geosci.* 59, 59–74. [https://doi.org/10.1016/0168-9622\(86\)90057-6](https://doi.org/10.1016/0168-9622(86)90057-6).
- Grottoli, A.G., Rodrigues, L.J., Matthews, K.A., Palardy, J.E., Gibb, O.T., 2005. Pre-treatment effect on coral skeletal $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. *Chem. Geol.* 221, 225–242. <https://doi.org/10.1016/j.chemgeo.2005.05.004>.
- Halfar, J., Steneck, R.S., Joachimski, M., Kronz, A., Wanamaker Jr., A.D., 2008. Coralline red algae as high-resolution climate recorders. *Geology* 36, 463–466. <https://doi.org/10.1130/G24635A.1>.
- Halfar, J., Hetzinger, S., Adey, W., Zack, T., Gamboa, G., Kunz, B., Williams, B., Jacob, D. E., 2011. Coralline algal growth-increment widths archive North Atlantic climate variability. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 302 (1–2), 71–80. <https://doi.org/10.1016/j.palaeo.2010.04.009>.
- Halfar, J., Adey, W.H., Kronz, A., Hetzinger, S., Edinger, E., Fitzhugh, W.W., 2013. Arctic sea-ice decline archived by multicentury annual-resolution record from crustose coralline algal proxy. *Proc. Natl. Acad. Sci. Unit. States Am.* 110 (49), 19737–19741. <https://doi.org/10.1073/pnas.1313775110>.
- Helser, T.E., Lai, H.L., Black, B.A., 2012. Bayesian hierarchical modeling of Pacific geoduck growth increment data and climate indices. *Ecol. Model.* 247, 210–220. <https://doi.org/10.1016/j.ecolmodel.2012.08.024>.
- Henry, K.M., Nixon, S.W., 2008. A half century assessment of hard clam, *Mercenaria mercenaria*, growth in Narragansett Bay, Rhode Island. *Estuar. Coast* 31, 755–766. <https://doi.org/10.1007/S12237-008-9060-9>.
- Hetzinger, S., Halfar, J., Zajacz, Z., Wisshak, M., 2019. Early start of 20th-century Arctic sea-ice decline recorded in Svalbard coralline algae. *Geology* 47 (10), 963–967. <https://doi.org/10.1130/g46507.1>.
- Hirabayashi, S., Yokoyama, Y., Suzuki, A., Miyairi, Y., Aze, T., 2017. Short-term fluctuations in regional radiocarbon reservoir age recorded in coral skeletons from the Ryukyu Islands in the north-western Pacific. *J. Quat. Sci.* 32 (1), 1–6. <https://doi.org/10.1002/jqs.2923>.
- Hobbs, J.A., Lewis, L.S., Willmes, M., Denney, C., Bush, E., 2019. Complex life histories discovered in a critically endangered fish. *Sci. Rep.* 9, 16772. <https://doi.org/10.1038/s41598-019-52273-8>.
- Höche, N., Peharda, M., Walliser, E.O., Schöne, B.R., 2020. Morphological variations of crossed-lamellar ultrastructures of *Glycymeris bimaculata* (Bivalvia) serve as a marine temperature proxy. *Estuar. Coast Shelf Sci.* 237, 106658. <https://doi.org/10.1016/j.ecss.2020.106658>.
- Holcomb, M., DeCarlo, T.M., Schoepf, V., Dissard, D., Tanaka, K., McCulloch, M., 2015. Cleaning and pre-treatment procedures for biogenic and synthetic calcium carbonate powders for determination of elemental and boron isotopic compositions. *Chem. Geol.* 398, 11–21. <https://doi.org/10.1016/j.chemgeo.2015.01.019>.
- Holland, H.A., Schöne, B.R., Marali, S., Jochum, K.P., 2014. History of bioavailable lead and iron in the Greater North Sea and Iceland during the last millennium a bivalve sclerochronological reconstruction. *Mar. Pollut. Bull.* 87, 104–116. <https://doi.org/10.1016/j.marpolbul.2014.08.005>.
- Hollyman, P.R., Leng, M.J., Chenery, S.R.N., Laptikhovskiy, V.V., Richardson, C.A., 2018. Statoliths of the whelk *Buccinum undatum*: a novel age determination tool. *Mar. Ecol. Prog. Ser.* 598, 261–272. <https://doi.org/10.3354/meps12119>.
- Hönisch, B., Ridgwell, A., Schmidt, D.N., Thomas, E., Gibbs, S.J., Sluijs, A., Zeebe, R., Kump, L., Martindale, R.C., Greene, S.E., Kiessling, W., Ries, J., Zachos, J.C., Royer, D.L., Barker, S., Marchitto, T.M., Moyer, R., Pelejero, C., Ziveri, P., Foster, G. L., Williams, B., 2012. The geological record of ocean acidification. *Science* 335 (6072), 1058–1063. <https://doi.org/10.1126/science.1208277>.
- Hudson, J.H., Shinn, E.A., Halley, R.B., Lidz, B., 1976. Sclerochronology: a tool for interpreting past environments. *Geology* 4 (6), 361–364.
- Huyghe, D., Lartaud, F., Emmanuel, L., Merle, D., Renard, M., 2015. Palaeogene climate evolution in the Paris Basin from oxygen stable isotope ($\delta^{18}\text{O}$) compositions of marine molluscs. *J. Geol. Soc.* 172, 576–587. <https://doi.org/10.1144/jgs2015-016>.
- Iacumin, P., Bianucci, G., Longinelli, A., 1992. Oxygen and carbon isotopic composition of fish otoliths. *Marine Biol.* 113 (4), 537–542. <https://doi.org/10.1007/bf00349696>.
- IPCC, 2018. Summary for Policymakers. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways. In: Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J.B.R., Chen, Y., Zhou, X., Gomis, M.I., Lonnoy, E., Maycock, T., Tignor, M., Waterfield, T. (Eds.), *The Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty* [Masson-Delmotte, V. World Meteorological Organization, Geneva, Switzerland, p. 32.
- Iseley, F.B., 1914. Experimental Study of the Growth and Migration of Fresh-Water Mussels. Govt. Washington. <https://doi.org/10.5962/bhl.title.42718> print. off.
- Ivanina, A.V., Dickinson, G.H., Matoo, O.B., Bagwe, R., Dickinson, A., Beniasi, E., Sokolova, I.M., 2013. Interactive effects of elevated temperature and CO₂ levels on energy metabolism and biomineralization of marine bivalves *Crassostrea virginica* and *Mercenaria mercenaria*. *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 166 (1), 101–111. <https://doi.org/10.1016/j.cbpa.2013.05.016>.
- Jones, D.S., 1981. Repeating layers in the molluscan shell are not always periodic. *J. Paleontol.* 55, 1076–1082.
- Jones, D.S., 1983. Sclerochronology: reading the record of the molluscan shell. *Am. Sci.* 71, 384–391.
- Jones, D.S., Williams, D.F., Romanek, C.S., 1986. Life history of symbiont-bearing giant clams from stable isotope profiles. *Science* 231, 46–48. <https://doi.org/10.1126/science.231.4733.46>.
- Jones, D.S., Arthur, M.A., Allard, D.J., 1989. Sclerochronological records of temperature and growth from shells of *Mercenaria mercenaria* from Narragansett Bay, Rhode Island. *Marine Biol.* 102, 225–234. <https://doi.org/10.1007/BF00428284>.
- Judd, E.J., Wilkinson, B.H., Ivany, L.C., 2018. The life and time of clams: derivation of intra-annual growth rates from high-resolution oxygen isotope profiles. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 490, 70–83. <https://doi.org/10.1016/j.palaeo.2017.09.034>.
- Jurikova, H., Liebetrau, V., Gutjahr, M., Rollion-Bard, C., Hu, M.Y., Krause, S., Henkel, D., Hiebenthal, C., Schmidt, M., Laudien, J., Eisenhauer, A., 2019. Boron isotope systematics of cultured brachiopods: response to acidification, vital effects and implications for palaeo-pH reconstruction. *Geochem. Cosmochim. Acta* 248, 370–386. <https://doi.org/10.1016/j.gca.2019.01.015>.
- Kaiser, B.A., Hoeberechts, M., Maxwell, K.H., Eerkes-Medrano, L., Hilmi, N., Safa, A., Horbel, C., Juniper, S.K., Roughan, M., Lowen, N.T., Short, K., Paruru, D., 2019. The importance of connected ocean monitoring knowledge systems and communities. *Front. Marine Sci.* 6, 1–17. <https://doi.org/10.3389/fmars.2019.00309>.
- Kalish, J.M., 1991. ^{13}C and ^{18}O isotopic disequilibria in fish otoliths: metabolic and kinetic effects. *Mar. Ecol. Prog. Ser.* 75 (2–3), 191–203. <https://doi.org/10.3354/meps075191>.
- Kamenos, N.A., Cusack, M., Moore, P.G., 2008. Red coralline algae are global paleothermometers with bi-weekly resolution. *Geochem. Cosmochim. Acta* 72, 771–779. <https://doi.org/10.1016/j.gca.2007.11.019>.
- Kelemen, Z., Gillikin, D.P., Graniero, L.E., Havel, H., Darchambeau, F., Borges, A.V., Yambélé, A., Bassirou, A., Bouillon, S., 2017. Calibration of hydroclimate proxies in freshwater bivalve shells from Central and West Africa. *Geochem. Cosmochim. Acta* 208, 41–62. <https://doi.org/10.1016/j.gca.2017.03.025>.
- Khider, D., Emile-Geay, J., McKay, N.P., Gil, Y., Garrijo, D., Ratnakar, V., Brewer, P., Bunn, A., Chevalier, M., Comas-Bru, L., Csank, A., Dassié, E., DeLong, K., Felis, T., Francus, P., Frappier, A., Gray, W., Goring, S., Jonkers, L., Kahle, M., Kaufman, D., Kehrwald, N.M., Martrat, B., McGregor, H., Richey, J., Schmittner, A., Sroxtton, N., Sutherland, E., Thirumalai, K., Allen, K., Arnaud, F., Axford, Y., Barrows, T.T., Bazin, L., Pilaar Birch, S.E., Bradley, E., Bregy, J., Capron, E., Cartapanis, O., Chiang, H.-W., Cobb, K., Debret, M., Dommaint, R., Du, J., Dyez, K., Emerick, S., Erb, M.P., Falster, G., Finsinger, W., Fortier, D., Gauthier, N., George, S., Grimm, E., Hertzberg, J., Hibbert, F., Hillman, A., Hobbs, W., Huber, M., Hughes, A.L.C., Jaccard, S., Ruan, J., Kienast, M., Konecky, B., Le Roux, G., Lyubchich, V., Novello, V.F., Olaka, L., Partin, J.W., Pearce, C., Phipps, S.J., Pignol, C., Piotrowska, N., Poli, M.-S., Prokopenko, A., Schwanck, F., Stepanek, C., Swann, G.E. A., Telford, R., Thomas, E., Thomas, Z., Truebe, S., von Gunten, L., Waite, A., Weitzel, N., Wilhelm, B., Williams, J., Williams, J.J., Winstrup, M., Zhao, N., Zhou, Y., 2019. PaCTS 1.0: a crowdsourced reporting standard for paleoclimate data. *Paleoceanography Paleoclimatol.* 34 (10), 1570–1596. <https://doi.org/10.1029/2019pa003632>.
- Kidwell, S.M., Best, M.M.R., Kaufman, D.S., 2005. Taphonomic trade-offs in tropical marine death assemblages: differential time averaging, shell loss, and probable bias in siliciclastic vs. carbonate facies. *Geology* 33, 729–732. <https://doi.org/10.1130/G21607.1>.
- Kilbourne, K.H., Quinn, T.M., Taylor, F.W., 2004. A fossil coral perspective on western tropical Pacific climate ~350 ka. *Paleoceanography* 19, PA1019. <https://doi.org/10.1029/2003PA000944>.
- Killam, D.E., Clapham, M.E., 2018. Identifying the ticks of bivalve shell clocks: seasonal growth in relation to temperature and food supply. *Palaios* 33 (5), 228–236. <https://doi.org/10.2110/palo.2017.072>.
- Killam, D.E., Thomas, R., Al-Naijjar, T., Clapham, M., 2020. Interspecific and intra-shell stable isotope variation among the Red Sea giant clams. *G-cubed* 21 (7). <https://doi.org/10.1029/2019GC008669> e2019GC008669.
- Kleypas, J.A., Feely, R.A., Fabry, V.J., Langdon, C., Sabine, C.L., Robbins, L.L., 2005. Impacts of Ocean Acidification on Coral Reefs and Other Marine Calcifiers: a Guide for Future Research (Paper presented at the Report of a workshop held).

- Kohler, A.C., 1964. Variations in the growth of Atlantic cod (*Gadus morhua* L.). J. Fish. Res. Board Can. <https://doi.org/10.1139/f64-007>. Canada 21: 57-100.
- Konecky, B.L., McKay, N.P., Churakova, O.V. (Sidorova), Comas-Bru, L., Dassié, E.P., DeLong, K.L., Falster, G.M., Fischer, M.J., Jones, M.D., Jonkers, L., Kaufman, D.S., Leduc, G., Managave, S.R., Martrat, B., Opel, T., Orsi, A.J., Partin, J.W., Sayani, H.R., Thomas, E.K., Thompson, D.M., Tyler, J.J., Abram, N.J., Atwood, A.R., Conroy, J.L., Kern, Z., Porter, T.J., Stevenson, S.L., von Gunten, L., the Iso2k Project Members, 2020. The Iso2k Database: A global compilation of paleo- $\delta^{18}\text{O}$ and $\delta^{2}\text{H}$ records to aid understanding of Common Era climate. in review Earth Syst. Sci. Data Discuss. <https://doi.org/10.5194/essd-2020-5>.
- Krause-Nehring, J., Brey, T., Thorrold, S.R., 2012. Centennial records of lead contamination in northern Atlantic bivalves (*Arctica islandica*). Mar. Pollut. Bull. 64, 233-240. <https://doi.org/10.1016/j.marpolbul.2011.11.028>.
- Leder, J.J., Swart, P.K., Szmant, A.M., Dodge, R.E., 1996. The origin of variations in the isotopic record of scleractinian corals: I. Oxygen. Geochim. Cosmochim. Acta 60 (15), 2857-2870. [https://doi.org/10.1016/0016-7037\(96\)00118-4](https://doi.org/10.1016/0016-7037(96)00118-4).
- LeGrande, A.N., Schmidt, G.A., 2006. Global gridded data set of the oxygen isotopic composition in seawater. Geophys. Res. Lett. 33, 1-5. <https://doi.org/10.1029/2006GL026011>.
- Lescinsky, H., Titus, B., Hubbard, D., 2011. Live coral cover in the fossil record: an example from Holocene reefs of the Dominican Republic. Coral Reefs 31, 335-346. <https://doi.org/10.1007/s00338-011-0863-y>.
- Leutert, T.J., Sexton, P.F., Tripathi, A., Piasecki, A., Ling, S.H., Meckler, A.N., 2019. Sensitivity of clumped isotope temperatures in fossil benthic and planktic foraminifera to diagenetic alteration. Geochim. Cosmochim. Acta 257, 354-372. <https://doi.org/10.1016/j.gca.2019.05.005>.
- Limburg, K.E., Casini, M., 2018. Effect of marine hypoxia on Baltic Sea cod *Gadus morhua*: evidence from otolith chemical proxies. Front. Marine Sci. 5 (482) <https://doi.org/10.3389/fmars.2018.00482>.
- Limburg, K.E., Olson, C., Walther, Y., Dale, D., Slomp, C.P., Høie, H., 2011. Tracking Baltic hypoxia and cod migration over millennia with natural tags. Proc. Natl. Acad. Sci. U.S.A. 108, E177-E182. <https://doi.org/10.1073/pnas.1100684108>.
- Limburg, K.E., Walther, B.D., Lu, Z., Jackman, G., Mohan, J., Walther, Y., Nissling, A., Weber, P.K., Schmitt, A.K., 2015. In search of the dead zone: use of otoliths for tracking fish exposure to hypoxia. J. Mar. Syst. 141, 167-178. <https://doi.org/10.1016/j.jmarsys.2014.02.014>.
- Linsley, B.K., Wu, H.C., Dassié, E.P., Schrag, D.P., 2015. Decadal changes in South Pacific sea surface temperatures and the relationship to the Pacific decadal oscillation and upper ocean heat content. Geophys. Res. Lett. 42 (7), 2358-2366. <https://doi.org/10.1002/2015gl063045>.
- Lockwood, R., Work, L.A., 2006. Quantifying taphonomic bias in Molluscan death assemblages from the upper Chesapeake bay: patterns of shell damage. Palaios 21, 442-450. <https://doi.org/10.2110/palo.2005.P05-083R>.
- Lorrain, A., Paulet, Y.-M., Chauvaud, L., Dunbar, R.B., Mucciarone, D., Fontugne, M., 2004. $\delta^{13}\text{C}$ variation in scallop shells: increasing metabolic carbon contribution with body size? Geochim. Cosmochim. Acta 68, 3509-3519. <https://doi.org/10.1016/j.gca.2004.01.025>.
- Ma, T.Y.H., 1934. On the seasonal change of growth in a reef coral, *Favia speciosa* (Dana), and the water-temperature of the Japanese seas during the latest geological times. Paper presented at the In: Proceedings of the Imperial Academy.
- Macdonald, J.I., Drysdale, R.N., Witt, R., Cságo, Z., Martensdóttir, G., 2019. Isolating the influence of ontogeny helps predict island-wide variability in fish otolith chemistry. Rev. Fish Biol. Fish. 30, 172-202. <https://doi.org/10.1007/s11160-019-09591-x>.
- Malde, K., Handegard, N.O., Eikvil, L., Salberg, A.B., July-August 2020. Machine intelligence and the data-driven future of marine science. ICES Journal of Marine Science 77 (4), 1274-1285. <https://doi.org/10.1093/icesjms/fsz057>.
- Mannino, M., Thomas, K., Leng, M., Sloane, H., 2008. Shell growth and oxygen isotopes in the topshell *Osilinus turbinatus*: resolving past inshore sea surface temperatures. Geo Mar. Lett. 28 (5-6), 309-325. <https://doi.org/10.1007/s00367-008-0107-5>.
- Marali, S., Schöne, B.R., 2015. Oceanographic control on shell growth of *Arctica islandica* (Bivalvia) in surface waters of Northeast Iceland - implications for paleoclimate reconstructions. Palaeogeogr. Palaeoclimatol. Palaeoecol. 420, 138-149. <https://doi.org/10.1016/j.palaeo.2014.12.016>.
- Marshall, J.F., McCulloch, M.T., 2002. An assessment of the Sr/Ca ratio in shallow water hermatypic corals as a proxy for sea surface temperature. Geochim. Cosmochim. Acta 66, 3263-3280. [https://doi.org/10.1016/S0016-7037\(02\)00926-2](https://doi.org/10.1016/S0016-7037(02)00926-2).
- Martin, J.M., Letolle, R., 1979. Oxygen 18 in estuaries. Nature 282, 292-294.
- Martino, J.C., Fowler, A.J., Doubleday, Z.A., Grammer, G.L., Gillanders, B.M., 2019. Using otolith chronologies to understand long-term trends and extrinsic drivers of growth in fisheries. Ecosphere 10 (1), e02553. <https://doi.org/10.1002/ecs2.2553>.
- Maton, W.G., 1805. A General View of the Writings of Linnaeus by Richard Pulteney, M. D. F.R.S, second ed. J. Mawman & R: Taylor, p. 595.
- Matta, M.E., Black, B.A., Wilderbuer, T.K., 2010. Climate-driven synchrony in otolith growth-increment chronologies for three Bering Sea flatfish species. Mar. Ecol. Prog. Ser. 413, 137-145. <https://doi.org/10.3354/meps08689>.
- McConnaughey, T., 1989. ^{13}C and ^{18}O isotopic disequilibrium in biological carbonates: II. In vitro simulation of kinetic isotope effects. Geochim. Cosmochim. Acta 53, 163-171. [https://doi.org/10.1016/0016-7037\(89\)90283-4](https://doi.org/10.1016/0016-7037(89)90283-4).
- McConnaughey, T.A., Gillikin, D.P., 2008. Carbon isotopes in mollusk shell carbonates. Geo Mar. Lett. 28 (5-6), 287-299. <https://doi.org/10.1007/s00367-008-0116-4>.
- McConnaughey, T.A., Burdett, J., Whelan, J.F., Paull, C.K., 1997. Carbon isotopes in biological carbonates: respiration and photosynthesis. Geochim. Cosmochim. Acta 61 (3), 611-622. [https://doi.org/10.1016/S0016-7037\(96\)000](https://doi.org/10.1016/S0016-7037(96)000).
- McCulloch, M.T., Gagan, M.K., Mortimer, G.E., Chivas, A.R., Isdale, P.J., 1994. A high-resolution Sr/Ca and $\delta^{18}\text{O}$ coral record from the great barrier reef, Australia and the 1982-1983 El Niño. Geochimica et Cosmochimica Acta 58, 2747-2754.
- McKay, N.P., Emile-Geay, J., 2016. Technical note: the Linked Paleo Data framework - a common tongue for paleoclimatology. Clim. Past 12, 1093-1100. <https://doi.org/10.5194/cp-12-1093-2016>.
- Meckler, A.N., Ziegler, M., Millán, M.I., Breitenbach, S.F.M., Bernasconi, S.M., 2014. Long-term performance of the Kiel carbonate device with a new correction scheme for clumped isotope measurements. RCM (Rapid Commun. Mass Spectrom.) 28, 1705-1715. <https://doi.org/10.1002/rcm.6949>.
- Melzner, F., Mark, F.C., Seibel, B.A., Tomanek, L., 2020. Ocean acidification and coastal marine invertebrates: tracking CO₂ effects from seawater to the cell. Annual Rev. Marine Sci. 12 (1) <https://doi.org/10.1146/annurev-marine-010419-010658>.
- Mette, M.J., Wanamaker Jr., A.D., Carroll, M.L., Ambrose Jr., W.G., Retelle, M.J., 2016. Linking large-scale climate variability with *Arctica islandica* shell growth and geochemistry in northern Norway. Limnol. Oceanogr. 61 (2), 748-764. <https://doi.org/10.1002/lno.10252>.
- Mette, M.J., Whitney, N.M., Ballew, J., Wanamaker Jr., A.D., 2018. Unexpected isotopic variability in biogenic aragonite: a user issue or proxy problem? Chem. Geol. 483, 286-294. <https://doi.org/10.1016/j.chemgeo.2018.02.027>.
- Milano, S., Schöne, B.R., Wang, S., Müller, W.E., 2016. Impact of high pCO₂ on shell structure of the bivalve *Cerastoderma edule*. Mar. Environ. Res. 119, 144-155. <https://doi.org/10.1016/j.marenvres.2016.06.002>.
- Milano, S., Schöne, B.R., Witbaard, R., 2017. Changes of shell microstructural characteristics of *Cerastoderma edule* (Bivalvia) - a novel proxy for water temperature. Palaeogeogr. Palaeoclimatol. Palaeoecol. 465 (B), 395-406. <https://doi.org/10.1016/j.palaeo.2015.09.051>.
- Moen, E., Handegard, N.O., Alken, V., Albert, O.T., Harbitz, A., Malde, K., 2018. Automatic interpretation of otoliths using deep learning. PLoS One 13 (12). <https://doi.org/10.1371/journal.pone.0204713>.
- Montagna, P., McCulloch, M., Douville, E., López Correa, M., Trotter, J., Rodolfo-Metalpa, R., Dissard, D., Ferrier-Pages, C., Frank, N., Friewald, A., Goldstein, S., Mazzoli, C., Reynaud, S., Ruggerberg, A., Russo, S., Taviani, M., 2014. Li/Mg systematics in scleractinian corals: calibration of the thermometer. Geochim. Cosmochim. Acta 132, 288-310. <https://doi.org/10.1016/j.gca.2014.02.005>.
- Morrongiello, J.R., Thresher, R.E., 2015. A statistical framework to explore ontogenetic growth variation among individuals and populations: a marine fish example. Ecol. Monogr. 85 (1), 93-115. <https://doi.org/10.1890/13-2355.1>.
- Morrongiello, J.R., Thresher, R.E., Smith, D.C., 2012. Aquatic biochronologies and climate change. Nat. Clim. Change 2, 849-857. <https://doi.org/10.1038/nclimate1616>.
- Morrongiello, J.R., Sweetman, P., Thresher, R.E., 2019. Fishing constrains phenotypic responses of marine fish to climate variability. J. Anim. Ecol. 88, 1645-1656. <https://doi.org/10.1111/1365-2656.12999>.
- Müller, I.A., Fernandez, A., Radke, J., van Dijk, J., Bowen, D., Schwieters, J., Bernasconi, S.M., 2017. Carbonate clumped isotope analyses with the long-integration dual-inlet (LIDI) workflow: scratching at the lower sample weight boundaries. Rapid Commun. Mass Spectrom. 31, 1057-1066. <https://doi.org/10.1002/rcm.7878>.
- Murakami-Sugihara, N., Shirai, K., Hori, M., Amano, Y., Fukuda, H., Obata, H., Tanaka, K., Mizukawa, K., Sano, Y., Takada, H., Ogawa, H., 2019. Mussel shell geochemical analyses reflect coastal environmental changes following the 2011 Tohoku tsunami. ACS Earth and Space Chem. 3, 1346-1352. <https://doi.org/10.1021/acsearthspacechem.9b00040>.
- Music, A., Flannery, J.A., Reich, C.D., Umberger, D.K., Smoak, J.M., Poore, R.Z., 2013. Linear Extension Rates of Massive Corals from the Dry Tortugas National Park (Drt), Florida. U.S. Geological Survey Open-File Report 2013-1121.
- Neuheimer, A.B., Thresher, R.E., Lyle, J.M., Semmens, J.M., 2011. Tolerance limit for fish growth exceeded by warming waters. Nat. Clim. Change 1, 110-113. <https://doi.org/10.1038/nclimate1084>.
- Nishida, K., Nakashima, R., Majima, R., Hikida, Y., 2011. Ontogenetic changes in shell microstructures in the cold seep-associated bivalve, *Conchocele bisecta* (Bivalvia: thyasiridae). Paleontol. Res. 15 (4), 193-212. <https://doi.org/10.2517/1342-8144-15.4.193>.
- Nishida, K., Ishikawa, K., Iguchi, A., Tanaka, Y., Sato, M., Ishimura, T., Inoue, M., Nakamura, T., Sakai, K., Suzuki, A., 2014. Skeletal oxygen and carbon isotope compositions of *Acropora* coral primary polyps experimentally cultured at different temperatures. G-cubed 15, 2840-2849. <https://doi.org/10.1002/2014GC005322>.
- Nishida, K., Suzuki, A., Isono, R., Hayashi, M., Watanabe, Y., Yamamoto, Y., Irie, T., Nojiri, Y., Mori, C., Sato, M., Sasaki, T., 2015. Thermal dependency of shell growth, microstructure, and stable isotopes in laboratory-reared *Scapharca broughtonii* (Mollusca: Bivalvia). G-cubed 16 (7), 2395-2408. <https://doi.org/10.1002/2014GC005634>.
- Nürnberg, D., Bijma, J., Hemleben, C., 1996. Assessing the reliability of magnesium in foraminiferal calcite as a proxy for water mass temperatures. Geochim. Cosmochim. Acta 60 (5), 803-814. [https://doi.org/10.1016/0016-7037\(95\)00446-7](https://doi.org/10.1016/0016-7037(95)00446-7).
- De Noia, M., Telesca, L., Vendrami, D.L.J., Gokalp, H.K., Charrier, G., Harper, E.M., Hoffman, J.I., 2020. Population genetic structure is unrelated to shell shape, thickness and organic content in European populations of the soft-shell clam *mya arenaria*. Genes 11 (3), 298. <https://doi.org/10.3390/genes11030298>.
- Norrie, C.R., Dunphy, B.J., Ragg, N.L.C., Lundquist, C.J., 2019. Comparative influence of genetics, ontogeny and the environment on elemental fingerprints in the shell of *Perna canaliculus*. Sci. Rep. 9 (1), 8533. <https://doi.org/10.1038/s41598-019-44947-0>.
- Ohno, T., 1989. Paleotidal characteristics determined by micro-growth patterns in bivalves. Paleontology 32, 237-263.

- Okazaki, A., Yoshimura, K., 2017. Development and evaluation of a system of proxy data assimilation for paleoclimate reconstruction. *Clim. Past* 13, 379–393. <https://doi.org/10.5194/cp-13-379-2017>.
- Ong, J.J.L., Rountrey, A.N., Zinke, J., Meeuwij, J.J., Grierson, P.F., O'Donnell, A.J., Newman, S.J., Wakefield, C.B., Meeuwij, J.J., Meekan, M.G., 2016. Evidence for climate-driven synchrony of marine and terrestrial ecosystems in northwest Australia. *Global Change Biol.* 22, 2776–2786. <https://doi.org/10.1111/gcb.13239>.
- Oschmann, W., 2009. Sclerochronology: editorial. *Int. J. Earth Sci.* 98 (1), 1–2. <https://doi.org/10.1007/s00531-008-0403-3>.
- Pages 2K Consortium, 2013. Continental-scale temperature variability during the past two millennia. *Nat. Geosci.* 6, 339–346. <https://doi.org/10.1038/ngeo1797>.
- Patiño, J., Whittaker, R.J., Borges, P.A.V., Fernández-Palacios, J.M., Ah-Peng, C., Araújo, M.B., Ávila, S.P., Cardoso, P., Cornuault, J., de Boer, E.J., de Nascimento, L., Gil, A., González-Castro, A., Gruner, D.S., Heleno, R., Hortal, J., Illera, J.C., Kaiser-Bunbury, C.N., Matthews, T.J., Papadopoulos, A., Pettorelli, N., Price, J.P., Santos, A.M.C., Steinbauer, M.J., Triantis, K.A., Valente, L., Vargas, P., Weigelt, P., Emerson, B.C., 2017. A roadmap for island biology: 50 fundamental questions after 50 years of the Theory of Island Biogeography. *J. Biogeogr.* 44, 963–983. <https://doi.org/10.1111/jbi.12986>.
- Pecquerie, L., Fablet, R., de Pontual, H., Bonhommeau, S., Alunno-Bruscia, M., Petitgas, P., Kooijman, S.A.L.M., 2012. Reconstructing individual food and growth histories from biogenic carbonates. *Mar. Ecol. Prog. Ser.* 447, 151–164. <https://doi.org/10.3354/meps09492>.
- Peharda, M., Black, B.A., Purroy, A., Mihanović, H., 2016. The bivalve *Glycymeris pilosa* as a multidecadal environmental archive for the Adriatic and Mediterranean Seas. *Mar. Environ. Res.* 119, 79–87. <https://doi.org/10.1016/j.marenvres.2016.05.022>.
- Peharda, M., Vilčić, I., Black, B.A., Markulin, K., Dunic, N., Dzoic, T., Mihanović, H., Gacić, M., Puljas, S., Waldman, R., 2018. Using bivalve chronologies for quantifying environmental drivers in a semi-enclosed temperate sea. *Sci. Rep.* 8 <https://doi.org/10.1038/s41598-018-23773-w>.
- Prendergast, A.L., Schöne, B.R., 2017. Oxygen isotopes from limpet shells: implications for palaeothermometry and seasonal shellfish foraging studies in the Mediterranean. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 484, 33–47. <https://doi.org/10.1016/j.palaeo.2017.03.007>.
- Prendergast, A.L., Stevens, R.E., 2014. Molluscs (Isotopes): Analyses in Environmental Archaeology. *The Encyclopedia of Global Archaeology*. Springer.
- Prendergast, A.L., Versteegh, E.A.A., Schöne, B.R., 2017. New research on the development of high-resolution palaeoenvironmental proxies from geochemical properties of biogenic carbonates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 484, 1–6. <https://doi.org/10.1016/j.palaeo.2017.05.032>.
- Prouty, N.G., Roark, E.B., Koenig, A.E., Demopoulos, A.W.J., Batista, F.C., Kocar, B.D., Selby, D., McCarthy, M.D., Mienis, F., Ross, S.W., 2014. Deep-sea coral record of human impact on watershed quality in the Mississippi River Basin. *Global Biogeochem. Cycles* 28, 29–43. <https://doi.org/10.1002/2013GB004754>.
- Pulteney, R., Payne, T., White, B., 1781. A General View of the Writings of Linnaeus, vol. 425. <https://doi.org/10.5962/bhl.title.96885>. Cambridge.
- Pyrina, M., Wagner, S., Zorita, E., 2017. Pseudo-proxy evaluation of climate field reconstruction methods of North Atlantic climate based on an annually resolved marine proxy network. *Clim. Past* 13, 1339–1354. <https://doi.org/10.5194/cp-13-1339-2017>.
- Raddatz, J., Rüggeberg, A., Flögel, S., Hathorne, E.C., Liebetrau, V., Eisenhauer, A., Dullo, W.-C., 2014. The influence of seawater pH on U/Ca ratios in the scleractinian cold-water coral *Lophelia pertusa*. *Biogeosciences* 11 (7), 1863–1871. <https://doi.org/10.5194/bg-11-1863-2014>.
- Ren, L., Linsley, B.K., Wellington, G.M., Schrag, D.P., Hoegh-Guldberg, O., 2002. Deconvolving the $\delta 180$ seawater component from subseasonal coral $\delta 180$ and Sr/Ca at Rarotonga in the southwestern subtropical Pacific for the period 1726 to 1997. *Geochem. Cosmochim. Acta* 67, 1609–1621.
- Reynolds, D.J., Butler, P.G., Williams, S.M., Scourse, J.D., Richardson, C.A., Wanamaker Jr., A.D., Austin, W.E.N., Cage, A.G., Sayer, M.D.J., 2013. A multiproxy reconstruction of Hebridean (NW Scotland) spring sea surface temperatures between AD 1805 and 2010. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 386, 275–285. <https://doi.org/10.1016/j.palaeo.2013.05.029>.
- Reynolds, D.J., Scourse, J.D., Halloran, P.R., Nederbragt, A.J., Wanamaker Jr., A.D., Butler, P.G., Richardson, C.A., Heinemeier, J., Eiríksson, J., Knudsen, K.L., Hall, I.R., 2016. Annually resolved North Atlantic marine climate over the last millennium. *Nat. Commun.* 7, 13502 <https://doi.org/10.1038/ncomms13502>.
- Reynolds, D.J., Richardson, C.A., Scourse, J.D., Butler, P.G., Hollyman, P., Roman-Gonzalez, A., Hall, I.R., 2017. Reconstructing North Atlantic marine climate variability using an absolutely-dated sclerochronological network. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 465, 333–346. <https://doi.org/10.1016/j.palaeo.2016.08.006>.
- Reynolds, D.J., Hall, I.R., Slater, S.M., Mette, M.J., Wanamaker Jr., A.D., Scourse, J.D., Garry, F.K., Halloran, P.R., 2018. Isolating and reconstructing key components of North Atlantic Ocean variability from a sclerochronological spatial network. *Paleoceanography Paleoclimatol.* 33, 1086–1098. <https://doi.org/10.1029/2018PA003366>.
- Reynolds, D.J., Hall, I.R., Slater, S.M., 2019. An integrated carbon and oxygen isotope approach to reconstructing past environmental variability in the northeast Atlantic Ocean. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 523, 48–61. <https://doi.org/10.1016/j.palaeo.2019.03.005>.
- Rhoads, D.C., Lutz, R.A., 1980. *Skeletal Growth of Aquatic Organisms: Biological Records of Environmental Change*. Plenum Press, New York and London, p. 750.
- Rhoads, D.C., Pannella, G., 1970. The use of molluscan shell growth patterns in ecology and paleoecology. *Lethaia* 3 (2), 143–161.
- Roberts, B.H., Morrongiello, J.R., King, A.J., Morgan, D.L., Saunders, T.M., Woodhead, J., Crook, D.A., 2019. Migration to freshwater increases growth rates in a facultatively catadromous tropical fish. *Oecologia* 191, 253–260. <https://doi.org/10.1007/s00442-019-04460-7>.
- Robinson, L.F., Adkins, J.F., Frank, N., Gagen, A.C., Prouty, N.G., Roark, E.B., van der Fliert, T., 2014. The geochemistry of deep-sea coral skeletons: a review of vital effects and applications for paleoceanography. *Deep-Sea Res.* II 99, 184–198. <https://doi.org/10.1016/j.dsr2.2013.06.005>.
- Rombouts, L., Beaugrand, G., Dauvin, J., 2012. Potential changes in benthic macrofaunal distributions from the English Channel simulated under climate change scenarios. *Estuar. Coast Shelf Sci.* 99, 153–161. <https://doi.org/10.1016/j.ecss.2011.12.026>.
- Rosenberg, G.D., Runcorn, S.K., 1975. *Growth Rhythms and the History of the Earth's Rotation*. John Wiley and Sons Ltd., New York, p. 576.
- Rosenheim, B.E., Swart, P.K., Thorold, S.R., Willenz, P., Berry, L., Latkoczy, C., 2004. High-resolution Sr/Ca records in sclerosponges calibrated to temperature in situ. *Geology* 32 (2), 145–148. <https://doi.org/10.1130/g20117.1>.
- Rountrey, A.N., Coulson, P.G., Meeuwij, J.A., Meekan, M., 2014. Water temperature and fish growth: otoliths predict growth patterns of a marine fish in a changing climate. *Global Change Biol.* 20, 2450–2458. <https://doi.org/10.1111/gcb.12617>.
- Sakamoto, T., Komatsu, K., Yoneda, M., Ishimura, T., Higuchi, T., Shirai, K., Kamimura, Y., Watanabe, C., Kawabata, A., 2017. Temperature dependence of $\delta 180$ in otolith of juvenile Japanese sardine: laboratory rearing experiment with micro-saline analysis. *Fish. Res.* 194, 55–59. <https://doi.org/10.1016/j.fishres.2017.05.004>.
- Der Sarkissian, C., Pichereau, V., Dupont, C., Ilsoe, P.C., Perrigault, M., Butler, P., Chauvaud, L., Eiríksson, J., Scourse, J., Paillard, C., Orlando, L., 2017. Ancient DNA analysis identifies marine mollusc shells as new metagenomic archives of the past. *Mol. Ecol. Resour.* 17 (5), 835–853. <https://doi.org/10.1111/1755-0998.12679>.
- Schmidt, G.A., Annan, J.D., Bartlein, P.J., Cook, B.I., Guiliardi, E., Hargreaves, J.C., Harrison, S.P., Kageyama, M., LeGrande, A.N., Konecky, B., Lovejoy, S., Mann, M.E., Masson-Delmotte, V., Risi, C., Thompson, D., Timmermann, A., Tremblay, L.-B., Yiou, P., 2014. Using palaeo-climate comparisons to constrain future projections in Cmp5. *Clim. Past* 10, 221–250. <https://doi.org/10.5194/cp-10-221-2014>.
- Schöne, B.R., Gillikin, D.P., 2013. Unravelling environmental histories from skeletal diaries - advances in sclerochronology. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 373, 1–5. <https://doi.org/10.1016/j.palaeo.2012.11.026>.
- Schöne, B.R., Krause Jr., R.A., 2016. Retrospective environmental biomonitoring - mussel Watch expanded. *Global Planet. Change* 14, 228–251. <https://doi.org/10.1016/j.gloplacha.2016.08.002>.
- Schöne, B.R., Surge, D., 2005. Looking back over skeletal diaries — high-resolution environmental reconstructions from accretionary hard parts of aquatic organisms. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 228 (1), 1–3. <https://doi.org/10.1016/j.palaeo.2005.03.043>.
- Schöne, B.R., Wanamaker Jr., A.D., Fiebig, J., Thébault, J., Kreutz, K., 2011. Annually resolved $\delta 13$ C shell chronologies of long-lived bivalve molluscs (*Arctica islandica*) reveal oceanic carbon dynamics in the temperate North Atlantic during recent centuries. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 302 (1–2), 31–42. <https://doi.org/10.1016/j.palaeo.2010.02.002>.
- Schöne, B.R., Radermacher, P., Zhang, Z., Jacob, D.E., 2013. Crystal fabrics and element impurities (Sr/Ca, Mg/Ca, and Ba/Ca) in shells of *Arctica islandica* - implications for paleoclimate reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 373, 50–59. <https://doi.org/10.1016/j.palaeo.2011.05.013>.
- Scott, P.J.B., 1990. Chronic pollution recorded in coral skeletons in Hong Kong. *J. Exp. Mar. Biol. Ecol.* 139 (1–2), 51–64. [https://doi.org/10.1016/0022-0981\(90\)90038-E](https://doi.org/10.1016/0022-0981(90)90038-E).
- Scourse, J., Richardson, C., Forsythe, G., Harris, I., Heinemeier, J., Fraser, N., Briffa, K., Jones, P., 2006. First cross-matched floating chronology from the marine fossil record: data from growth lines of the long-lived bivalve mollusc *Arctica islandica*. *Holocene* 16, 967–974. <https://doi.org/10.1177/095958360601987p>.
- Seddon, A.W.R., Mackay, A.W., Baker, A.G., Birks, H.J.B., Breman, E., Buck, C.E., Ellis, E. C., Froyd, C.A., Gill, J.L., Gillson, L., Johnson, E.A., Jones, V.J., Juggins, S., Macias-Fauria, M., Mills, K., Morris, J.L., Nogués-Bravo, D., Surangi, W., Roland, T.P., Tanentzap, A.J., Willis, K.J., Aberhan, M., Asperen, E. N. Van, Austin, W.E.N., Battarbee, R.W., Bhagwat, S., Belanger, C.L., Bennett, K.D., Birks, H.H., Ramsey, C. B., Brooks, S.J., de Bruyn, M., Butler, P.G., Chambers, F.M., Clarke, S.J., Davies, A.L., Dearing, J.A., Ezard, H.G., Feurdean, A., Flower, R.J., Gell, P., Hausmann, S., Liow, L.H., McGowan, S., Miller, J.H., Montoya, E., Morton, O., Nogué, S., Onoufriou, C., Boush, L.P., Rodriguez-Sanchez, F., Rose, N.L., Sayer, C.D., Shaw, H. E., Payne, R., Simpson, G., Sohar, K., Whitehouse, N.J., Williams, J.W., Witkowski, A., 2013. Looking forward through the past: identification of 50 priority research questions in palaeoecology. *J. Ecol.* 102, 256–267. <https://doi.org/10.1111/1365-2745.12195>.
- Sherwood, O., Edinger, E.N., Guilderson, T.P., Ghaleb, B., Risk, M.J., Scot, D.B., 2008. Late Holocene radiocarbon variability in Northwest Atlantic slope waters. *Earth Planet Sci. Lett.* 275, 146–153. <https://doi.org/10.1016/j.epsl.2008.08.019>.
- Sherwood, O.A., Guilderson, T.P., Batista, F.C., Schiff, J.T., McCarthy, M.D., 2014. Increasing subtropical north pacific ocean nitrogen fixation since the little ice age. *Nature* 505 (7481), 78–81. <https://doi.org/10.1038/nature12784>.
- Shirai, K., Schöne, B.R., Miyaji, T., Radermacher, P., Krause, R., Tanabe, K., 2014. Assessment of the mechanism of elemental incorporation into bivalve shells (*Arctica islandica*) based on elemental distribution at the microstructural scale. *Geochem. Cosmochim. Acta* 126, 307–320. <https://doi.org/10.1016/j.gca.2013.10.050>.
- Sirot, C., Grønckjær, P., Pedersen, J.B., Panfili, J., Zetina-Rejon, M., Tripp-Valdez, A., Ramos-Miranda, J., Flores-Hernandez, D., Sosa-Lopez, A., Darnaude, A.M., 2017. Using otolith organic matter to detect diet shifts in *Bardiella chrysoura*, during a period of environmental changes. *Mar. Ecol. Prog. Ser.* 575, 137–152. <https://doi.org/10.3354/meps12166>.

- van der Sleen, P., Stransky, C., Morrongiello, J.R., Haslob, H., Peharda, M., Black, B.A., 2018. Otolith increments in European plaice (*Pleuronectes platessa*) reveal temperature and density-dependent effects on growth. *ICES (Int. Coun. Explor. Sea) J. Mar. Sci.* 75, 1655–1663. <https://doi.org/10.1093/icesjms/fsy011>.
- Smith, J.E., Schwarcz, H.P., Risk, M.J., McConnaughey, T.A., Keller, N., 2000. Paleotemperatures from deep-sea corals: overcoming “vital effects. *Palaios* 15 (25). <https://doi.org/10.2307/3515589>.
- Spalding, C., Finnegan, S., Fischer, W.W., 2017. Energetic costs of calcification under ocean acidification. *Global Biogeochem. Cycles* 31 (5), 866–877. <https://doi.org/10.1002/2016GB005597>.
- Steinhardt, J., Butler, P.G., Carroll, M.L., Hartley, J., 2016. The application of long-lived bivalve sclerochronology in environmental baseline monitoring. *Front. Marine Sci.* 3 (176) <https://doi.org/10.3389/fmars.2016.00176>.
- Strom, A., Francis, R.C., Mantua, N.J., Miles, E.L., Peterson, D.L., 2004. North Pacific climate recorded in growth rings of geoduck clams: a new tool for paleoenvironmental reconstruction. *Geophys. Res. Lett.* 31 (6) <https://doi.org/10.1029/2004GL019440>.
- Surge, D., Wang, T., Gutiérrez-Zugasti, I., Kelley, P.H., 2013. Isotope sclerochronology and season of annual growth line formation in limpet shells (*Patella vulgata*) from warm- and cold-temperate zones in the eastern North Atlantic. *Palaios* 28, 386–393. <https://doi.org/10.2110/palo.2012.p12-038r>.
- Sutherland, W.J., Fleishman, E., Mascia, M.B., Pretty, J., Rudd, M.A., 2011. Methods for collaboratively identifying research priorities and emerging issues in science and policy. *Methods in Ecol. Evol.* 2, 238–247. <https://doi.org/10.1111/j.2041-210X.2010.00083.x>.
- Sutherland, W.J., Freckleton, R.P., Godfray, H.C.J., Beissinger, S.R., Benton, T., Cameron, D.D., Carmel, Y., Coomes, D.A., Coulson, T., Emmerson, M.C., Hails, R.S., Hays, G.C., Hodgson, D.J., Hutchings, M.J., Johnson, D., Jones, J.P., Keeling, G., Kokko, M.J., Kunin, H., Lambin, W.E., Lewis, X., Malhi, O.T., Mieszekowska, Y., Milner-Gulland, N., Norris, E.J., Whillimore, K., Purves, A.B., Reid, D.W., Reuman, J. M., Thompson, D.C., Travis, K., Turnbull, J.M.J., Wardle, D. A.L.A., Wiegand, T., 2013. Identification of 100 fundamental ecological questions. *J. Ecol.* 101, 58–67. <https://doi.org/10.1111/1365-2745.12025>.
- Suzuki, M., Saruwatari, K., Kogure, T., Yamamoto, Y., Nishimura, T., Kato, T., Nagasawa, H., 2009. An acidic matrix protein, Pif, is a key macromolecule for nacre formation. *Science* 325 (5946), 1388–1390. <https://www.jstor.org/stable/40301783>.
- Swart, P.K., Greer, L., Rosenheim, B.E., Moses, C.S., Waite, A.J., Winter, A., Dodge, R.E., Helmle, K., 2010. The ¹³C Suess effect in scleractinian corals mirror changes in the anthropogenic CO₂ inventory of the surface oceans. *Geophys. Res. Lett.* 37 (5), L05604 <https://doi.org/10.1029/2009GL041397>.
- Takesue, R.K., Bacon, C.R., Thompson, J.K., 2008. Influences of organic matter and calcification rate on trace elements in aragonitic estuarine bivalve shells. *Geochem. Cosmochim. Acta* 72, 5431–5445. <https://doi.org/10.1016/j.gca.2008.09.003>.
- Telesca, L., Peck, L.S., Sanders, T., Thyrring, J., Sejr, M.K., Harper, E.M., 2019. Biomineralization plasticity and environmental heterogeneity predict geographical resilience patterns of foundation species to future change. *Global Change Biol.* 25 (12), 4179–4193. <https://doi.org/10.1111/gcb.14758>.
- Thibault, M., Duprey, N., Gillikin, D.P., Thébaud, J., Douillet, P., Chauvaud, L., Amice, E., Munaron, J.M., Lorrain, A., 2020. Bivalve δ15N isoscapes provide a baseline for urban nitrogen footprint at the edge of a World Heritage coral reef. *Mar. Pollut. Bull.* 152, 110870. <https://doi.org/10.1016/j.marpolbul.2019.110870>.
- Thorrold, S.R., Campana, S.E., Jones, C.M., Swart, P.K., 1997. Factors determining δ13C and δ18O fractionation in aragonitic otoliths of marine fish. *Geochem. Cosmochim. Acta* 61, 2909–2919. [https://doi.org/10.1016/S0016-7037\(97\)00141-5](https://doi.org/10.1016/S0016-7037(97)00141-5).
- Thresher, R., Morrongiello, J., Sloyan, B.M., Krusic-Golub, K., Shephard, S., Minto, C., Nolan, C.P., Cerna, F., Cid, L., 2014. Parallel decadal variability of inferred water temperatures for Northern and Southern Hemisphere intermediate water masses. *Geophys. Res. Lett.* 41 (4), 1232–1237. <https://doi.org/10.1002/2013GL058638>.
- Tierney, J.E., Abram, N.J., Anchukaitis, K.J., Evans, M.N., Giry, C., Kilbourne, K.H., Saenger, C.P., Wu, H.C., Zinke, J., 2015. Tropical sea surface temperatures for the past four centuries reconstructed from coral archives. *Paleoceanography* 30, 226–252. <https://doi.org/10.1002/2014PA002717>.
- Tindall, J.C., Haywood, A.M., Thirumalai, K., 2017. Modeling the stable water isotope expression of El Niño in the Pliocene: implications for the interpretation of proxy data. *Paleoceanography* 32, 881–902. <https://doi.org/10.1002/2016PA003059>.
- Tobin, T.S., Schauer, A.J., Lewarch, E., 2011. Alteration of micromilled carbonate δ18O during Kiel Device analysis. *Rapid Commun. Mass Spectrom.* 25, 2149–2152. <https://doi.org/10.1002/rcm.5093>.
- Toft, C.A., Shea, P.J., 1983. Detecting community-wide patterns: estimating power strengths statistical inference. *Am. Nat.* 122, 618–625. <https://doi.org/10.1086/284162>.
- Toniello, G., Lepofsky, D., Lertzman-Lepofsky, G., Salomon, A.K., Rowell, K., 2019. 11,500 y of human-clam relationships provide long-term context for intertidal management in the Salish Sea, British Columbia. *Proc. Natl. Acad. Sci. Unit. States Am.* 116, 22106–22114. <https://doi.org/10.1073/pnas.1905921116>.
- Tray, E., Leadbetter, A., Meaney, W., Conway, A., Kelly, C., Ó Maoiléidigh, N., de Eyto, E., Moran, S., Brophy, D., 2020. An open-source database model and collections management system for fish scale and otolith archives. *Ecol. Inf.* 101115 <https://doi.org/10.1016/j.ecoinf.2020.101115>.
- Trofimova, T., Milano, S., Andersson, C., Bonitz, F.G.W., Schöne, B.R., 2018. Oxygen isotope composition of *Arctica islandica* aragonite in the context of shell architectural organization: implications for paleoclimate reconstructions. *G-cubed* 19, 453–470. <https://doi.org/10.1002/2017GC007239>.
- Trueman, C.N., Jackson, A.L., Chadwick, K.S., Coombs, E.J., Feyrer, L.J., Magozzi, S., Sabin, R.C., Cooper, N., 2019. Combining simulation modeling and stable isotope analyses to reconstruct the last known movements of one of Nature’s giants. *PeerJ* 7, e7912. <https://doi.org/10.7717/peerj.7912>.
- Urey, H.C., 1947. The thermodynamic properties of isotopic substances. *J. Chem. Soc.* 562–581, 0.
- Urey, H.C., Lowenstam, H.A., Epstein, S., McKinney, C.R., 1951. Measurement of paleotemperatures and temperatures of the upper cretaceous of England, Denmark, and the southeastern United States. *GSA Bull.* 62, 399–416. [https://doi.org/10.1130/0016-7606\(1951\)62\[399.MOPATO\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1951)62[399.MOPATO]2.0.CO;2).
- Vogel, K., Gektidis, M., Golubic, S., Kiene, W.E., Radtke, G., 2000. Experimental studies on microbial bioerosion at lee stocking island, Bahamas and one tree island, great barrier reef, Australia: implications for paleoecological reconstructions. *Lethaia* 33, 190–204. <https://doi.org/10.1080/00241160025100053>.
- Wanamaker Jr., A.D., Butler, P.G., Scourse, J.D., Heinemeier, J., Eiríksón, J., Knudsen, K.L., Richardson, C.A., 2012. Surface changes in the North Atlantic meridional overturning circulation during the last millennium. *Nat. Commun.* 3, 1–7. <https://doi.org/10.1038/ncomms1901>.
- Wagner, A.J., Slowey, N.C., 2011. Oxygen isotopes in seawater from the Texas-Louisiana shelf. *Bull. Mar. Sci.* 87, 1–12. <https://doi.org/10.5343/bms.2010.1004>.
- Waite, A.J., Swart, P.K., 2015. The inversion of aragonite to calcite during the sampling of skeletal archives: implications for proxy interpretation. *Rapid Commun. Mass Spectrom.* 29, 955–964. <https://doi.org/10.1002/rcm.7180>.
- Wanamaker Jr., A.D., Kreutz, K.J., Borns, H.W., Introne, D.S., Feindel, S., Funder, S., Rawson, P.D., Barber, B.J., 2007. Experimental determination of salinity, temperature, growth, and metabolic effects on shell isotope chemistry of *Mytilus edulis* collected from Maine and Greenland. *Paleoceanography* 22 (2). <https://doi.org/10.1029/2006PA001352>.
- Wanamaker Jr., A.D., Hetzinger, S., Halfar, J., 2011. Reconstructing mid- to high-latitude marine climate and ocean variability using bivalves, coralline algae, and marine sediment cores from the Northern Hemisphere. *Paleogeogr. Palaeoclimatol. Palaeoecol.* 302, 1–9. <https://doi.org/10.1016/j.palaeo.2010.12.024>.
- Wanamaker Jr., A.D., Griffin, S.M., Ummerhofer, C.C., Whitney, N.M., Black, B., Parfitt, R., Lower-Spies, E.E., Introne, D., Kreutz, K.J., 2019. Pacific climate influences on ocean conditions and extreme shell growth events in the Northwestern Atlantic (Gulf of Maine). *Clim. Dynam.* 52, 6339–6356. <https://doi.org/10.1007/s00382-018-4513-8>.
- Wang, T., Surge, D., Walker, K.J., 2011. Isotopic evidence for climate change during the Vandal Minimum from *Ariopsis felis* otoliths and *Mercenaria campechiensis* shells, southwest Florida, USA. *Holocene* 21, 1081–1091. <https://doi.org/10.1177/0959683611400458>.
- Wang, T., Surge, D., Walker, K.J., 2013. Seasonal climate change across the Roman Warm Period/Vandal Minimum transition using isotope sclerochronology in archaeological shells and otoliths, southwest Florida, USA. *Quat. Int.* 308–309, 230–241. <https://doi.org/10.1016/j.quaint.2012.11.013>.
- Wang, Y., Huang, J., Tang, X., Jin, X., Sun, Y., 2016. Stable isotopic composition of otoliths in identification of stock structure of small yellow croaker (*Larimichthys polyactis*) in China. *Acta Oceanol. Sin.* 35, 29–33. <https://doi.org/10.1007/s13131-016-0868-z>.
- Wasserstein, R.L., Schirm, A.L., Lazar, N.A., 2019. Moving to a world beyond “p < 0.05. *Am. Statistician* 73, 1–19. <https://doi.org/10.1080/00031305.2019.1583913>.
- Weber, J.N., Woodhead, P.M.J., 1970. Carbon and oxygen isotope fractionation in the skeletal carbonate of reef-building corals. *Chem. Geol.* 6, 93–117. [https://doi.org/10.1016/0009-2541\(70\)90009-4](https://doi.org/10.1016/0009-2541(70)90009-4).
- Weber, J.N., Woodhead, P.M.J., 1972. Temperature dependence of oxygen-18 concentration in reef coral carbonates, 1896–1977. *J. Geophys. Res.* 77, 463–473. <https://doi.org/10.1029/JC077i003p00463>.
- Weidman, C.R., Jones, G.A., Lohman, K.C., 1994. The long-lived mollusc *Arctica islandica*: a new paleoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern North Atlantic Ocean. *J. Geophys. Res.: Oceans* 99, 18305–18314. <https://doi.org/10.1029/94JC01882>.
- Weiner, S., Dove, P.M., 2003. An overview of biomineralization processes and the problem of the vital effect. *Rev. Mineral. Geochem.* 54, 1–29. <https://doi.org/10.2113/0540001>.
- Weisberg, S., Spangler, G., Richmond, L.S., 2010. Mixed effects models for fish growth. *Can. J. Fish. Aquat. Sci.* 67, 269–277. <https://doi.org/10.1139/F09-181>.
- Wells, J.W., 1963. Coral growth and geochronometry. *Nature* 197, 948–950. <https://doi.org/10.1038/197948a0>.
- Wierzbowski, H., 2007. Effects of pre-treatments and organic matter on oxygen and carbon isotope analyses of skeletal and inorganic calcium carbonate. *Int. J. Mass Spectrom.* 268, 16–29. <https://doi.org/10.1016/j.ijms.2007.08.002>.
- Wilkinson, M.D., Dumontier, M., Aalbersberg, L.J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.W., da Silva Santos, L.B., Bourne, P.E., Bouwman, J., Brookes, A.J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C.T., Finkers, R., Gonzalez-Beltran, A., Gray, A.J.G., Groth, P., Goble, C., Grethe, J.S., Heringa, J., ‘t Hoen, P.A.C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons, A., Packer, A.L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., Van Der Lei, J., Van Mulligen, E., Velterop, J., Waagmeester, A., Wittenberg, P., Wolstencroft, K., Zhao, J., Mons, B., 2016. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* 3, 1–9. <https://doi.org/10.1038/sdata.2016.18>.
- Williams, B., Risk, M.J., Ross, S.W., Sulak, K.J., 2007. Stable isotope data from deep-water Antipatharians: 400-year records from the southeastern coast of the United States of America. *Bull. Mar. Sci.* 81, 437–447.
- Williams, B., Halfar, J., Steneck, R.S., Wortmann, U.G., Hetzinger, S., Adey, W., Lebednik, P., Joachimski, M., 2011. Twentieth century 13C variability in surface water dissolved inorganic carbon recorded by coralline algae in the northern North

- Pacific Ocean and the Bering Sea. *Biogeosciences* 8, 165–174. <https://doi.org/10.5194/bg-8-165-2011>.
- Williams, B., Halfar, J., DeLong, K.L., Hetzinger, S., Steneck, R.S., Jacob, D.E., 2014. Multi-specimen and multi-site calibration of Aleutian coralline algal Mg/Ca to sea surface temperature. *Geochem. Cosmochim. Acta* 139, 190–204. <https://doi.org/10.1016/j.gca.2014.04.006>.
- Williams, B., Halfar, J., DeLong, K.L., Smith, E., Steneck, R., Lebednik, P.A., Jacob, D.E., Fietzke, J., Moore, G.W.K., 2017. North Pacific twentieth century decadal-scale variability is unique for the past 342 years. *Geophys. Res. Lett.* 44, 3761–3769. <https://doi.org/10.1002/2017GL073138>.
- de Winter, N.J., Goderis, S., Dehairs, F., Jagt, J.W., Fraaije, R.H., Van Malderen, S.J., Vanhaecke, F., Claeys, P., 2017. Tropical seasonality in the late Campanian (late Cretaceous): comparison between multiproxy records from three bivalve taxa from Oman. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 485, 740–760. <https://doi.org/10.1016/j.palaeo.2017.07.031>.
- de Winter, N.J., Vellekoop, J., Vorrsselmans, R., Golreihani, A., Soete, J., Petersen, SierraV., Meyer, KyleW., Casadio, S., Speijer, RobertP., Claeys, P., 2018. An assessment of latest Cretaceous *Pycnodonte vesicularis* (Lamarck, 1806) shells as records for palaeoseasonality: a multi-proxy investigation. *Clim. Past* 14, 725–749. <https://doi.org/10.5194/cp-14-725-2018>.
- Wu, H.C., Moreau, M., Linsley, B.K., Schrag, D.P., Corrège, T., 2014. Investigation of sea surface temperature changes from replicated coral Sr/Ca variations in the eastern equatorial Pacific (Clipperton Atoll) since 1874. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 412, 208–222. <https://doi.org/10.1016/j.palaeo.2014.07.039>.
- Wu, H.C., Felis, T., Scholz, D., Giry, C., Kölling, M., Jochum, K.P., Scheffers, S.R., 2017. Changes to Yucatán Peninsula precipitation associated with salinity and temperature extremes of the Caribbean Sea during the Maya civilization collapse. *Sci. Rep.* 7, 1–12. <https://doi.org/10.1038/s41598-017-15942-0>.
- Yamazaki, A., Watanabe, T., Tsunogai, U., Iwase, F., Yamano, H., 2016. A 150-year variation of the Kuroshio transport inferred from coral nitrogen isotope signature. *Paleoceanography* 31 (6), 838–846. <https://doi.org/10.1002/2015PA002880>.
- Yan, H., Chen, J., Xiao, J., 2014. A review on bivalve shell, a tool for reconstruction of paleo-climate and paleo-environment. *Chin. J. Geochem.* 33, 310–315. <https://doi.org/10.1007/s11631-014-0692-0>.
- Zachariassen, J., Sieh, K., Taylor, F.W., Hantoro, W.S., 2000. Modern vertical deformation above the sumatran subduction zone: paleogeodetic insights from coral microatolls. *Bull. Seismol. Soc. Am.* 90 (4), 897–913. <https://doi.org/10.1785/0119980016>.
- Zhang, Q., Zhou, J., He, J., Xiaodong, C., Zeng, S., Zhang, B., 2019. A shell dataset, for shell features extraction and recognition. *Scientific Data* 6, 226. <https://doi.org/10.1038/s41597-019-0230-3>.
- Zhao, L., Schöne, B.R., Mertz-Kraus, R., 2017a. Controls on strontium and barium incorporation into freshwater bivalve shells (*Corbicula fluminea*). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 465, 386–394. <https://doi.org/10.1016/j.palaeo.2015.11.040>.
- Zhao, L., Schöne, B.R., Mertz-Kraus, R., Yang, F., 2017b. Sodium provides unique insights into transgenerational effects of ocean acidification on bivalve shell formation. *Sci. Total Environ.* 577, 360–366. <https://doi.org/10.1016/j.scitotenv.2016.10.200>.
- Zhao, L., Walliser, E.O., Mertz-Kraus, R., Schöne, B.R., 2017c. Unionid shells (*Hyriopsis cumingii*) record manganese cycling at the sediment-water interface in a shallow eutrophic lake in China (Lake Taihu). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 484, 97–108. <https://doi.org/10.1016/j.palaeo.2017.03.010>.
- Zhao, L., Milano, S., Walliser, E.O., Schöne, B.R., 2018. Bivalve shell formation in a naturally CO₂-enriched habitat: unraveling the resilience mechanisms from elemental signatures. *Chemosphere* 203, 132–138. <https://doi.org/10.1016/j.chemosphere.2018.03.180>.
- Zhao, L., Milano, S., Tanaka, K., Liang, J., Deng, Y., Yang, F., Walliser, E.O., Schöne, B.R., 2020. Trace elemental alterations of bivalve shells following transgenerational exposure to ocean acidification: implications for geographical traceability and environmental reconstruction. *Sci. Total Environ.* 705, 135501. <https://doi.org/10.1016/j.scitotenv.2019.135501>.