



HAL
open science

Total Synthesis of Poisonous Aconitum Alkaloids Empowered by a Fragment Coupling Strategy

Bastien Nay

► **To cite this version:**

Bastien Nay. Total Synthesis of Poisonous Aconitum Alkaloids Empowered by a Fragment Coupling Strategy. ACS Central Science, 2021, 7, pp.1298-1299. 10.1021/acscentsci.1c00887. hal-03313910

HAL Id: hal-03313910

<https://hal.science/hal-03313910>

Submitted on 4 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Total Synthesis of Poisonous *Aconitum* Alkaloids Empowered by a Fragment Coupling StrategyCite This: <https://doi.org/10.1021/acscentsci.1c00887>

Read Online

ACCESS |

Metrics & More

Article Recommendations

Bastien Nay

Convergent total synthesis of highly complex bridged alkaloids through key fragment coupling.

It is no mean feat to synthesize such complex natural products as the C₁₉ diterpene alkaloids (C₁₉ DTAs), featuring unique polycyclic and bridged structures (Figure 1). In this issue of *ACS Central Science*, Reisman and




Figure 1. Structure of C₁₉ diterpene alkaloids and an *Aconitum* plant from which they can be isolated (Picture Credit: Peter H. Raven Library/Missouri Botanical Garden).

co-workers take on this considerable challenge, employing a fragment coupling strategy to synthesize (–)-talatisamine (1), (–)-liljestrandisine (2), and (–)-liljestrandinine (3).¹ A series of highly chemoselective transformations was used

to forge the polycyclic bridged skeleton of the natural products, one of which was structurally revised (2 vs 4).

The *Aconitum* plants are well-known for their ability to biosynthesize neurotoxic C₁₉ DTAs.² The plants themselves were used in medicinal preparations or as poison for hunting, due to the properties of the natural products to block cation channels in neurons, and to exhibit anti-inflammatory, antihypertensive, pain relief, or antiarrhythmic activities. The structure of these complex compounds results from peculiar biosynthetic transformations, especially a semipinacol rearrangement of the denudatine-type (5) into the aconitine-type (6) skeletons (Figure 2a).³ This reaction has inspired the synthetic strategy of many chemists, starting with Wiesner's total synthesis of talatisamine **1** in the 1970s.^{4–7} Besides, the retrosynthetic logic classically advises to unravel the complexity of bridged compounds by disconnecting the maximally bridged cycles.^{5,8}

In fact, Reisman et al. solved this synthetic puzzle in a distinctly different manner, by coupling together two complex fragments that were separately assembled (Figure 2b). Such convergent approaches are well-known to empower the total synthesis of complex structures by making the synthetic route more flexible.⁹ Fragment **A** was made from phenol thanks to a stereoselective *meta*-photocycloaddition, while fragment **B** was enantioselectively constructed from cyclopentenone. Both intermediates were coupled through a unique sequence of 1,2-addition followed by a high-yielding semipinacol rearrangement to deliver 4 g of intermediate 7, which allows the stereoselective formation of the C-11

In fact, Reisman et al. solved this synthetic puzzle in a distinctly different manner, by coupling together two complex fragments that were separately assembled.




Figure 2. (a) Biosynthesis of *Aconitum* diterpene alkaloids and (b) the fragment coupling approach proposed by Reisman et al.

quaternary center at the same time. Unfortunately, the later key step involving a straightforward challenging *N*-radical cascade failed to construct the complete carbocyclic core of the natural product, thereby imposing a modification of the initial strategy. To solve these difficulties, a stepwise approach was adopted, and numerous chemoselective reactions were applied to finally afford the natural products. The reader will surely thrive on the many clever transformations that were used for this work.

The reader will surely thrive on the many clever transformations that were used for this work.

Thus, complex C_{19} DTAs were stereoselectively delivered in about 30 steps (1–3 mg each). While this achievement is not to compete with the phytochemical extraction from the plant (100 mg of **2** were isolated from 8 kg of plant through a complex purification process),¹⁰ the preparative goal of total synthesis is by far not the only output that can be expected from such projects. First, challenging total syntheses closely correlate with the development of modern robust and reliable synthetic methods, many of which were developed and applied by Reisman and co-workers. Second, working on natural product synthesis leads to the preparation of complex synthetic intermediates that can be natural product analogues for additional prospects in biology. This is particularly relevant with such biologically active series. Third, the comparison of synthetic compounds with the natural products can occasionally lead to structure revision. This is a striking result of this new work: to revise the stereochemistry of liljestrandisine, initially reported as **2**,

into structure **4**. Fourth and to conclude this discussion, there is well-established evidence that the field of total synthesis participates in the education of the next generation of organic chemists. In that context, synthetic targets as complex as the C_{19} DTA natural products certainly offer one of the best playgrounds to stimulate our creativity and challenge our capabilities.

In that context, synthetic targets as complex as the C_{19} DTA natural products certainly offer one of the best playgrounds to stimulate our creativity and challenge our capabilities.

Author Information

Corresponding Author

Bastien Nay – Laboratoire de Synthèse Organique, Ecole Polytechnique, Palaiseau 91128, France; orcid.org/0000-0002-1209-1830; Email: bastien.nay@polytechnique.edu

Complete contact information is available at: <https://pubs.acs.org/10.1021/acscentsci.1c00887>

REFERENCES

- (1) Wong, A. R.; Fastuca, N. J.; Mak, V. W.; Kerkovius, J. K.; Stevenson, S. M.; Reisman, S. E. Total Syntheses of the C_{19} Diterpenoid Alkaloids (–)-Talatisamine, (–)-Liljestrandisine, and (–)-Liljestrandinine by a Fragment Coupling Approach. *ACS Cent. Sci.* **2021**, DOI: [10.1021/acscentsci.1c00540](https://doi.org/10.1021/acscentsci.1c00540).
- (2) Wang, F.-P.; Chen, Q.-H.; Liu, X.-Y. Diterpenoid Alkaloids. *Nat. Prod. Rep.* **2010**, *27*, 529–570.
- (3) Johnston, J. P.; Overton, K. H. A Laboratory Model for the Atisane → Aconane Conversion. *J. Chem. Soc., Perkin Trans. 1* **1972**, *1*, 1490–1500.
- (4) Wiesner, K. The Total Synthesis of Racemic Talatisamine. *Pure Appl. Chem.* **1975**, *41*, 93–112.
- (5) Marth, C. J.; Gallego, G. M.; Lee, J. C.; Lebold, T. P.; Kulyk, S.; Kou, K. G. M.; Qin, J.; Lilien, R.; Sarpong, R. Network-Analysis-Guided Synthesis of Weisaconitine D and Liljestrandinine. *Nature* **2015**, *528*, 493–498.
- (6) Nishiyama, Y.; Yokoshima, S.; Fukuyama, T. Total Synthesis of (–)-Cardiopetaline. *Org. Lett.* **2016**, *18*, 2359–2362.
- (7) Kamakura, D.; Todoroki, H.; Urabe, D.; Hagiwara, K.; Inoue, M. Total Synthesis of Talatisamine. *Angew. Chem., Int. Ed.* **2020**, *59*, 479–486.
- (8) Corey, E. J.; Cheng, X.-M. *The Logic of Chemical Synthesis*; Wiley, 1989.
- (9) Tomanik, M.; Hsu, I. T.; Herzon, S. B. Fragment Coupling Reactions in Total Synthesis That Form Carbon–Carbon Bonds via Carbanionic or Free Radical Intermediates. *Angew. Chem., Int. Ed.* **2021**, *60*, 1116–1150.
- (10) Xie, G.-B.; Wang, F.-P. Structure of Liljestrandisine—a New C_{19} -Diterpenoid Alkaloid from *Aconitum Liljestrandii*. *Chem. J. Chin. Univ.* **2004**, *25*, 482–483.