



HAL
open science

Knowledge and vaccination practices among family physicians in northeastern France regarding tick-borne encephalitis virus

Marie Dollat, Anne-Pauline Bellanger, Laurence Millon, Catherine Chirouze, Quentin Lepiller, Philippe Marguet

► To cite this version:

Marie Dollat, Anne-Pauline Bellanger, Laurence Millon, Catherine Chirouze, Quentin Lepiller, et al.. Knowledge and vaccination practices among family physicians in northeastern France regarding tick-borne encephalitis virus. *Ticks and Tick-borne Diseases*, 2021, 12 (5), pp.101774. 10.1016/j.ttbdis.2021.101774 . hal-03316054

HAL Id: hal-03316054

<https://hal.science/hal-03316054v1>

Submitted on 2 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Knowledge and vaccination practices among family physicians in northeastern France**
2 **regarding tick-borne encephalitis virus.**

3

4 Marie Dollat^a, Anne-Pauline Bellanger^{b*}, Laurence Millon^b, Catherine Chirouze^c, Quentin
5 Lepiller^d, Philippe Marguet^a

6

7 ^a Emergency Department, Pontarlier Regional Hospital, 25300 Pontarlier, France

8 ^b Department of Parasitology-Mycology, Besançon University Hospital, 25000 Besançon,
9 France

10 ^c Department of Infectious and Tropical Diseases, Besançon University Hospital, 25000
11 Besançon, France

12 ^d Department of Virology, Besançon University Hospital, 25000 Besançon, France

13

14 ***Corresponding author:** Anne-Pauline Bellanger, Department of Parasitology-Mycology,
15 Jean Minjoz University Hospital, 25030 Besançon, France.

16 Fax 33 (0)3 70 63 23 64. Tel 33 (0)3 70 63 23 51

17 E-mail address: apbellanger@chu-besancon.fr

18

19

20 **Declaration of interests:** The authors declare no conflict of interests.

21 **Running title:** Knowledge and vaccination practices regarding TBE

22 **Keywords:** family physicians, knowledge, practices, tick-borne encephalitis, vaccination

23 **Funding source:** No specific funding was obtained for this work

24

25

26 **Abstract**

27 Tick-borne encephalitis (TBE) cases have been emerging in Europe. The Franche-Comte area,
28 in northeastern France, borders Switzerland, but the two countries differ in their approach to
29 TBE surveillance and prevention. Because family physicians (FPs) are in direct contact with
30 the local population, at-risk of infected tick bites, they need to be well aware of TBE
31 epidemiology and management. An observational survey was performed in 2019 in order to
32 investigate Franche-Comte physicians' knowledge and vaccination practices with regard to
33 TBE. Standardized online questionnaires were sent to a list of FPs practicing in Franche-
34 Comte. The questionnaires included socio-demographic details, questions about TBE
35 knowledge, symptomatology and vaccination. The response rate was 14.7%. FPs practicing in
36 rural areas reported a significantly higher frequency of consultations for tick bites. While 81%
37 of FPs indicated that they had some knowledge about TBE, only 20% were at ease with its
38 clinical symptomatology. Thirty-one % of the FP participants performed TBE vaccinations. A
39 general lack of knowledge about TBE and its clinical symptoms was observed in this survey.
40 FPs play an essential role in screening and preventing TBE, especially those practicing in
41 rural areas and in areas bordering Switzerland.

42

43 **Introduction**

44 Tick-borne encephalitis (TBE) is due to a virus of the *Flaviviridae* family, which is
45 transmitted ~~immediately~~ during the tick blood meal of infected *Ixodes* spp. The *I. ricinus* tick
46 species is the primary vector of TBE in Europe, and its geographic expansion has been
47 documented both in latitude and in altitude (Suk and Semenza, 2014). This geographic
48 expansion is associated with an increase of high-risk areas and the emergence of TBE foci
49 (Michel et al., 2020). ~~Since TBE is a zoonosis, reservoirs of animals, including rodents, may~~
50 ~~explain this geographic extension (Michel et al., 2020).~~ Three subtypes of TBE virus are
51 circulating from Russia to Western Europe; the European subtype, present in France, was
52 reported as the less virulent one (Lehrer and Holbrook, 2011).

53 The case fatality rate of TBE is low (less than 2%) but there is a considerable proportion of
54 long-lasting sequelae (17-30%) (Bogovic et al., 2018; Velay et al., 2018). As no curative
55 treatment is available for this viral infection, prevention depends on avoiding tick bites and
56 receiving the TBE vaccination. Two licensed TBE vaccines are currently distributed in
57 Europe (Erber and Schmitt, 2018).

58 Out of approximately 3000 TBE cases reported in Europe each year, approximately 300 are
59 from Switzerland and only approximately 30 from France (Velay et al., 2019). In Switzerland,
60 TBE case notification has been mandatory for several years. The recent increase of cases
61 reported in 2018 led the Swiss authorities to promote vaccination for the entire Swiss
62 population, especially for people who regularly engage in outdoor activities (OFSP-Bulletin
63 16/19; [https://www.bag.admin.ch/bag/fr/home/das-bag/publikationen/periodika/bag-](https://www.bag.admin.ch/bag/fr/home/das-bag/publikationen/periodika/bag-bulletin.html)
64 [bulletin.html](https://www.bag.admin.ch/bag/fr/home/das-bag/publikationen/periodika/bag-bulletin.html)).

65 TBE cases have increased in France in recent years, and particularly in areas located in
66 northeastern regions bordering Germany (Beauté et al., 2018, Velay et al., 2018). In June

67 2020, the High Council for Public Health recommended that TBE case notification become
68 mandatory
69 (<https://www.hcsp.fr/explore.cgi/avisrapportsdomaine?clefr=856>).

70 Few TBE cases have thus far been reported in Franche-Comte, which is a heavily forested
71 area in northeastern France (<https://inventaire-forestier.ign.fr>), bordering Switzerland.
72 Serologic assays performed on non-vaccinated forestry workers in northeastern France
73 showed a TBE seroprevalence of about 3.4% (Thorin et al., 2008). Every day, frontier
74 workers, forestry workers and outdoor athletes cross the fictive line between Switzerland and
75 France.

76 Several other factors may also account for an increase in TBE cases: increased abundance of
77 ticks, increased range of *I. ricinus* tick, climate change, changes in human life-style such as
78 the general increase of leisure outdoor activities (Lehrer and Holbrook, 2011; Velay et al.,
79 2019).

80 Family physicians (FPs) are in direct contact with the local population and should be well
81 aware of the epidemiologic characteristics of infectious diseases present in the area where
82 they work. Indeed, FPs play a major role in promoting appropriate preventive measures. ~~and~~
83 ~~allowing early screening of the disease.~~

84 In this context, in 2019, an observational survey was undertaken in the Franche-Comte region
85 to assess family physicians' knowledge and vaccination practices for TBE.
86
87

88 **Methods**

89 Over a 12-week period from September to December 2019, standard online questionnaires
90 were sent to 544 FPs practicing in the Franche-Comte area. The questionnaire was designed
91 to be short to encourage participation. It was made up of 10 questions: questions requesting
92 sociodemographic details (number of years of practice, geographical area of practice, zip
93 code, estimated number of consultations for tick bites per year), perception of their general
94 knowledge about TBE, and questions concerning TBE vaccination practices (estimated
95 frequency and circumstances of TBE vaccinations performed).

96 The FPs were asked to participate on a voluntary and anonymous basis. The inclusion criteria
97 were: FPs working in Franche Comte with an e-mail address validated by the Regional Union
98 of Health Professionals and /or by the Department of General Medicine.

99 The online R software was used for statistical analysis. Qualitative variables were tested using
100 Chi-square test, with a p -value of less than 0.05 as the threshold of statistical significance.

101

102

103 **Results**

104 The response rate was 14.7% (80/544 questionnaires completed).

105 Most of participating FPs had been practicing for more than ten years (64%). A total of 77.5%
106 (62/80) FPs reported seeing 10 to 50 consultations for tick bites per year (Figure 1).

107 Interestingly, of the 15% (12/80) of FPs reporting more than 50 consultations per year for tick
108 bites (Figure 1), 83% were working in rural areas ($p < 0.0001$, Chi²) and 92% were practicing
109 in areas bordering Switzerland.

110 A total of 81% (65/80) of the participating FPs declared having some knowledge of TBE but
111 only 20% (16/80) were at ease with its clinical symptomatology.

112 FPs with more than 10 years of practice indicated a lack of knowledge of TBE more often
113 than the FPs with less experience (28% vs 3%, $p=0.0081$ Chi²).

114 A total of 31.3 % (25/80) reported that they performed the TBE vaccination. More precisely,
115 in 32% of the cases, patients received the vaccination before travelling to Central Europe, in
116 29% of the cases due to professional exposure, in 26% of the cases because of leisure outdoor
117 activities and in 13% of the cases before travelling to Central Asia. Reasons for not
118 performing TBE vaccinations are presented in Figure 2. A total of 68% (54/80) of the
119 participating FPs did not propose the TBE vaccination to their patients either because they
120 were not familiar with vaccine recommendations (46%) or with vaccine modalities (22%)
121 (Figure 2).

122

123

124

125

126 **Discussion**

127 The aim of this observational survey was to investigate the knowledge and vaccination
128 practices of FPs working in the Franche-Comte area of northeastern France. The response
129 rate of 14.7% was low, however, the sample was considered representative of the population
130 for the Franche-Comte area based on their geographical distribution (area with or without
131 common borders with Switzerland) and their age distribution (compared with the 2015-age
132 pyramid).

133 On the whole, our study highlighted family physicians' lack of knowledge about TBE and its
134 symptomatology. There was no significant correlation between better knowledge of TBE and
135 rural medical practices with a high number of consultations for tick bites. This observation is
136 alarming as rural areas in the northeastern part of France are particularly at risk of exposure to
137 infected ticks. In areas bordering Switzerland, FPs reported a higher rate of consultations for
138 tick bites (>50 per year), a finding which concurs with a French study performed from 2010
139 to 2012 (Serre et al., 2017). FPs performing TBE vaccinations followed the official
140 recommendation (professional exposure, travel in endemic areas) (Chrdle et al., 2016). ~~Those~~
141 ~~not performing TBE vaccinations (68%) were unfamiliar with vaccine recommendations or~~
142 ~~modalities.~~ FPs practicing in areas bordering Switzerland performed more TBE vaccinations.
143 The difference in epidemiology between France and Switzerland is mainly due to the fact that
144 TBE case notification has been mandatory for several years in Switzerland but not in France,
145 although case notification will soon become mandatory: (High Council for Public Health
146 recommendation June 2020 <https://www.hcsp.fr/explore.cgi/avisrapportsdomaine?clefr=856>).
147 The difference in of incidence ~~the number of TBE cases reported in~~ between Switzerland and
148 France could also be explained, at least in part, by eco-epidemiological factors ~~the~~
149 ~~geographical location of Switzerland and its proximity to Germany and Austria.~~ The tick
150 season lasts from March to November, and recent data from 2020 showed a general increase

176

177 **Conclusion**

178 A lack of knowledge about TBE and its symptomatology was observed in the present survey
179 of FPs practicing in Franche-Comte. Efforts should be made to provide up-to-date specialized
180 information about TBE to family physicians, especially those working in rural areas and areas
181 bordering Switzerland. For this reason, a flyer summarizing the results of the study and
182 including reminders about TBE epidemiology, clinical symptoms, prevention, and vaccination
183 was sent to all the participating FPs.

184

185 **References**

186

187 Beauté, J., Spiteri, G., Warns-Petit, E., Zeller, H., 2018. Tick-borne encephalitis in Europe,

188 2012 to 2016. *Euro Surveill.* 23,1800201. [https://dx.doi.org/10.2807/1560-](https://dx.doi.org/10.2807/1560-7917.ES.2018.23.45.1800201)

189 7917.ES.2018.23.45.1800201.

190 Bogovic, P., Stupica, D., Rojko, T., Kotric-Furlan, S., Avsic-Zupanc, T., Kastrin, A., Lusa, L.,

191 Strle, F., 2018. The long-term outcome of tick-borne encephalitis in Central Europe. *Ticks*

192 *Tick-borne Dis.* 9,369-378. <https://dx.doi.org/10.1016/j.ttbdis.2017.12.001>.

193 Chrdle, A., Chmelík, V., Růžek, D., 2016. Tick-borne encephalitis: what travelers should

194 know when visiting an endemic country. *Hum Vaccin Immunother.* 12:2694-2699.

195 <https://dx.doi.org/10.1080/21645515.2016.1218098>.

196 *Epid. Bull* 9-2021.

197 Erber, W., Schmitt, H.J., 2018. Self-reported tick-borne encephalitis vaccination coverage in

198 Europe: results from a cross-sectional study. *Ticks Tick Borne Dis.* 9:768-777.

199 <https://dx.doi.org/10.1016/j.ttbdis.2018.02.007>.

200 ~~Koch R. Institut, 2021: FSME: Risikogebiete in Deutschland. *Epid. Bull.* 3-20.~~

201 Lehrer, A.T., Holbrook, M.R., 2011. Tick-borne encephalitis vaccines. *J bioterror Biodef.*

202 *Suppl 1*-doi: 10.4172/2157-2526.S1-003.

203 Lipowski, D., Popiel, M., Perlejewski K., Nakamura S., Bukowska-Osko I., Rzedkiewicz

204 E., Dzieciatkowski, T., Milecka, A., Wenski, W., Ciszek, M., Debska-Slizien, A., Ignacak

205 E., Caraballo Cortes, K., Pawelczyk, A., Horban, A., Radkowski, M., Laskus, T., 2017. A

206 cluster of fatal tick-borne encephalitis virus infection in organ transplant setting. *J Infect Dis.*

207 215:896-901. <https://dx.doi.org/10.1093/infdis/jix040>.

208 Michel, F., Ziegler U., Fast, C., Eiden, M., Klaus, C., Dobler, G., Stiasny, K., Groschup,

209 M.H., 2020. Role of ducks in the transmission cycle of tick-borne encephalitis virus?

210 *Transbound Emerg Dis.* doi: 10.1111/tbed.13704.

211 Rubel, F., Brugger, K., 2020. Tick-borne encephalitis forecasts for Austria, Germany, and
212 Switzerland. *Ticks Tick Borne Dis.* 11:101437. doi: 10.1016/j.ttbdis.2020.101437.

213 Septfons, A., Goronflot, T., Jaulhac, B., Roussel, V., De Martino, S., Guerreiro, S., Launay,
214 T., Fournier, L., De Valk, H., Figoni, J., Blanchon, T., Couturier, E., 2019. Epidemiology of
215 Lyme borreliosis through two surveillance systems: the national Sentinelles GP network and
216 the national hospital discharge database, France, 2005 to 2016. *Euro Surveill.* 24:1800134.
217 doi: 10.2807/1560-7917.ES.2019.24.11.1800134

218 Serre, A., Tessier, S., Stoll, J., 2017. [Étude sur les maladies transmises par les tiques en
219 Franche Comté, 2010-2012]. In *VS Bourgogne Franche Comte.*

220 Suk, J.E., Semenza, J.C., 2014. From global to local : vector-borne disease in an
221 interconnected world. *Eur J Public Health.* 24:531-532. doi: 10.1093/eurpub/cku041.

222 Thorin, C., Rigaud, E., Capek, I., André-Fontaine, G., Oster, B., Gastinger, G., Abadia, G.,
223 2008. [Séroprévalence de la borréliose de Lyme et de l'encéphalite à tiques chez des
224 professionnels exposés dans le grand est de la France]. *Med Mal Infect.* 38:533-542. doi:
225 10.1016/j.medmal.2008.06.008.

226 Velay, A., Solis, M., Kack-Kack, W., Gantner, P., Maquart, M., Martinot, M., Augereau, O.,
227 De Briel, D., Kieffer, P., Lohmann, C., Poveda, J.D., Cart-Tanneur, E., Argemi, X., Leparc-
228 Goffart, I., De Martino, S., Jaulhac, B., Raguét, S., Wendling, M.J., Hansmann, Y., Fafi-
229 Kremer, S., 2018. A new hot spot for tick-borne encephalitis (TBE): A marked increase of
230 TBE cases in France in 2016. *Ticks Tick Borne Dis.* 9:120-125. [https://dx.doi.org/
231 10.1016/j.ttbdis.2017.09.015.](https://dx.doi.org/10.1016/j.ttbdis.2017.09.015)

232 Velay, A., Argemi, X., Wendling, M.J., Martinot, M., Hansmann, Y., Fati-Kremer, S., 2019.
233 [L'encéphalite à tique en France : qu'en savons nous aujourd'hui]. *Revue Francophone des
234 Laboratoires.* 513:34-43.

235

236 **Acknowledgement**

237 We thank Pamela Albert for her editorial assistance.

238

239 **Figure legends**

240 **Figure 1.** Distribution of the participants according to the number of patients consulting for

241 tick bites per year.

242 **Figure 2.** Reasons for not performing TBE vaccinations.


243

244

245

246

247



Reasons for not practicing TBE vaccination

