



HAL
open science

Ancient forest statistics provide centennial perspective over the status and dynamics of forest area in France

Timothée Audinot, Holger Wernsdörfer, Jean-Daniel Bontemps

► To cite this version:

Timothée Audinot, Holger Wernsdörfer, Jean-Daniel Bontemps. Ancient forest statistics provide centennial perspective over the status and dynamics of forest area in France. *Annals of Forest Science*, 2020, 77 (3), pp.1-24. 10.1007/s13595-020-00987-5 . hal-03317972

HAL Id: hal-03317972

<https://hal.science/hal-03317972v1>

Submitted on 9 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Ancient forest statistics provide centennial perspective over the status and dynamics of forest area in France

Timothée Audinot^{1,2} · Holger Wernsdörfer² · Jean-Daniel Bontemps¹

Received: 3 April 2020 / Accepted: 16 July 2020 / Published online: 5 August 2020
© INRAE and Springer-Verlag France SAS, part of Springer Nature 2020

Abstract

- **Key message** Centenary forest statistics informing major attributes of French forests were digitized, checked for consistency, and used to infer forest dynamics. Comparison to forest inventory data highlights increases in forest area and tree diversity, and substantial maturation of forests. Dataset access at <https://doi.org/10.5281/zenodo.3739458>
- **Context** The history of European forest dynamic remains fragmental. In France, the *Daubrée* statistics (1908) and agricultural statistics (1892, 1929) formed fundamental material to fill this gap.
- **Aims** Release, test, and summarize the digitalized dataset. Analyze long-term forest changes in forest area, composition, and structure.
- **Methods** Primary data on forest area across NUTS-3 geographic units, split by forest management and ownership categories and dominating tree species (*Daubrée*), were digitized and cross-compared. Centennial changes in forest attributes were assessed from modern forest inventory data.
- **Results** Cross-comparison revealed: (1) strong temporal consistency in forest changes over time, (2) systematic and interpretable biases in ownership/management categories between *Daubrée* and agricultural statistics. Strong shift from coppices to high forests, increased prevalence of private ownership, and constant proportion of broadleaf- and conifer-dominated forests were highlighted, with increased tree species diversity at country scale.
- **Conclusion** Ancient statistics are shown to play a major role in retrospective land-use and forest policy analysis.

Keywords Historical data · National forest inventory · Forest dynamic · Land-use change · Forest structure · Tree diversity

1 Introduction

As an outcome of past demographic transition and industrial revolution, most European countries have been experiencing a forest transition since the nineteenth century, consisting of a

strong regain of forest area (Mather 1992; Meyfroidt and Lambin 2011). While present forests' state has been accurately documented since 1990 with implementation of Forest Europe reporting (Forest Europe 2015) and the increasing implementation of statistical national forest inventory (NFI) programs across Europe (Tomppo et al. 2010), forest dynamics over more ancient time periods remains poorly covered (Mather et al. 1999; Meyfroidt and Lambin 2011). This has prevented from elaborating a clear view on the nature and dynamic of forest changes after the forest transition, including appraisal of areal, structural and compositional forest changes. Such long-term perspective is however of major concern to forest status and ecosystem conservation appraisal (Willis and Birks 2006; Foster et al. 2003).

Across European countries, forests of metropolitan France are those currently experiencing the strongest areal and growing stock changes (Forest Europe 2015). According to the French NFI, started in 1961, French forests have been increasing by almost 17% in area over 30 years (Denardou et al.

Handling Editor: Marco Ferretti

✉ Timothée Audinot
timothee.audinot@ign.fr

Holger Wernsdörfer
holger.wernsdorfer@agroparistech.fr

Jean-Daniel Bontemps
Jean-Daniel.Bontemps@ign.fr

¹ IGN, Laboratoire d'Inventaire Forestier, 14 rue Girardet, 54000 Nancy, France

² Université de Lorraine, AgroParisTech, INRAE, SILVA, F-54000 Nancy, France

2017). This increase results from both land abandonment (Chakir and Madignier 2006; Cavailhès and Normandin 1993) and forest extension ensuing from natural colonization, and active afforestation programs based on conifer species (afforestation of the Landes de Gascogne in Southwestern France by the mid-nineteenth century, David 1993, Buffault 1942; Restoration of Mountain Terrain (RTM) by the end nineteenth century, Saillet 1991; or afforestation supported by the National Forest Fund (FFN) after WWII, Betolaud 1972), with unquantified consequences on forest area and composition. Also, spontaneous forest maturation may have strongly modified the prevalence of forest management types (e.g., coppice or high forest abundance) together with the growing stock.

With this respect, ancient forest data are fairly available in France as compared with other countries (Mather et al. 1999). The first comprehensive survey of French forests is provided by *Daubrée* statistics of 1908 (Daubrée (Ministère de l’agriculture, FRANCE) 1912; Brenac 1984). Preliminary analysis of these data revealed an increase by +60% in the forested area over one century (Denardou et al. 2017). These ancient data were not yet based on survey sampling, and methods actually used remain unspecified. Also, no definition of forests based on dimensional and vegetation cover criteria (international forest definition of FAO 2000) was used at this remote period of time, making these figures uncertain as regards how forested land was distinguished from other types of vegetation and land uses.

In this respect, agricultural statistics provide valuable information for both the test of data consistency by cross-comparison approaches, and analyses of forest-area changes. Between 1862 and 1929, five decennial agricultural statistics have been produced by public authorities about every 10–30 years (1862, 1882, 1892, 1902, and 1929; Oberlé 2000). All of them include a section dedicated to forests. In addition to the *Daubrée* statistics of 1908, agricultural statistics from 1892 to 1929 were considered in the present study, with a view to cross-compare ancient statistics, and cover a period distant by around one century to present time for analyses of long-term forest changes.

The three statistical surveys describe forest area according to three categorical variables of interest: (1) the department administrative units, corresponding to the 90 NUTS-3 units of the nomenclature of the European Commission (European Parliament and Council of the European Union 2003) with an area of 3000 to 10,000 km² per unit; (2) forest management types, where at least coppices, high forests or coppice-with-standards forests are distinguished. Of note, forest management types primarily reflected objectives of forest planning rather than physical forest structure; (3) the legal category of forest ownership. In addition, *Daubrée* statistics provides areal statistics for major forest tree species of the country, making it possible to infer long-term compositional changes in

French forests by comparison with present NFI data that are available yearly on the whole French territory since 2005.

The first objective of this paper was to improve our knowledges about past forest dynamics based on ancient statistics. The underlying goals were: (1) to digitize the three ancient statistics, only found in paper or image format to date, in order to make them available for further analysis; (2) to analyze and cross-compare ancient statistics to evaluate their consistency; (3) to analyze forest dynamics over more than one century, according to forest management types, ownership categories, and forest tree species, where modern NFI were also used. All comparisons between statistics required harmonization of data variables and resolution.

2 Material and methods

2.1 Data description

2.1.1 Data overview

Daubrée statistics *Daubrée* statistics (names after its director) as the first comprehensive forest statistics in France was published in two volumes in 1912. Data were collected by the administration of “Eaux et Forêts” (“Water and Forests”, Corvol 2013) between 1904 and 1910 (Brenac 1984). Nevertheless, all data refer to 1908 (Brenac 1984; Oberlé 2000). Protocols for historical data acquisition remain fragmental. Forest data were collected from the archives of “Eaux et Forêts”, and from forest owners and associated organization. Forest areal data were provided by public local authorities and ensued from the French land registry (Corvol 2013; IGN 2013). The statistics provides forest information also at a resolution below the NUTS-3 level, i.e. by “canton” administrative units (LAU-1 of the EC, >3000 units in France; European Parliament and Council of the European Union 2013). While the latter resolution remains out of scope of the present study, it was, e. g., used in Dupouey and Hervé (2013) for spatial inference of forest changes.

One synoptic table of *Daubrée* statistics is reproduced in Annex 1.

Decennial agricultural statistics of 1892 and 1929 Decennial agricultural statistics (Ministère de l’Agriculture - Direction de l’Agriculture 1892; 1929) were intended to provide a comprehensive survey of the agricultural land of the national territory and its associated productions (Hitier 1899; Fromont 1938). A special section is dedicated to forests. We considered data from this section, only.

Decennial agricultural statistics relied on questionnaire surveys of municipalities (Oberlé 2000). Survey results were aggregated by the French administration prior to their publication (Musset 1938).

Agricultural statistics of 1892 (published in 1897 and acquired in 1892, Hitier 1899) and 1929 (published in 1936 and acquired between 1928 and 1933, Musset 1938) also provide information on forest area and harvested volume at the departmental scale.

Forest tables of both decennial agricultural statistics are reproduced in Annex 1.

Complementary data from the French national forest inventory NFI data were used to assess changes in both forest structure and composition over a one-century period.

Inventory data rely on an annual and spatially systematic sampling of forests (Robert et al. 2010; Vidal et al. 2016) and comply with the international FAO definition of forest (FAO 2000). With aggregation of successive years of inventory increasing data precision (Wernsdörfer et al. 2012) a 5-year aggregation of 2006-to-2010 inventories (median year 2008) distant by 100 years from the *Daubrée* statistics was computed.

2.1.2 Standard categorical variables

Daubrée statistics Forest area was split by ownership category and submission to the “Forest regime”, associated to legally-binding management rules defined by the forestry code of 1827 (Némoz-Rajot 1998, Annex 2) and isolating: State-owned forests, municipal forests, and forests of other territorial public bodies subjected to forest regime; municipal forests non subjected to the forest regime and private forests (never submitted to the forest regime). In addition to the three forest management types identified in all three statistics (see Introduction), two management types were specific of *Daubrée* statistics: (i) the transient regime of coppice conversion enforced by the forest law of 1827, and aiming at shifting to high forests for timber production (Degron 1998), and (ii) unproductive forest areas. The latter corresponded to marshes, bogs, rocks, and reforestation perimeters of forest protection programs. “Unproductive” hence needs to be understood in the literal meaning of not subjected to tree felling, as these areas are systematically encompassed for estimation of the afforestation rating, yet systematically associated with zero harvested volume.

Daubrée statistics included statistics on forest owners and annual harvested volume. All state variables available are summarized in Tables 1 and 2.

Decennial agricultural statistics of 1892 and 1929 Forest area was classified according to forest management type, ownership category, and forest regime submission (Tables 1 and 2). Both statistics split the area of high forests according to botanical classes (broadleaf- or conifer-dominated forests). Whereas no information on the method used to achieve this distinction (evaluation of class predominance), the

significance of compositional mixtures in French forests (Morneau et al. 2008) makes these figures uncertain.

Like in the *Daubrée* statistics, specific forest management types were identified. 1892 statistics includes category decree afforestation perimeters and unproductive coastal dunes solely found in forests subjected to the forest regime. This complies with the temporary or permanent acquisition by the State, of lands dedicated to landscape protection in coastal and mountain areas, as ordinated by successive RTM laws (Saillet 1991; Sonnier 1991). 1929 statistics includes a forest management type related to recently afforested area.

In addition, annual harvested volume is included in both decennial agricultural statistics (Table 2).

French national forest inventory We used the breakdown of the forest area according to ownership category and the forest regime submission (State-owned forests, municipal and other public forests subjected to forest regime, private and other public forests non-subjected to forest regime merged in the NFI process).

Forest area at a NUTS-3 resolution was further broken down by stand vertical structure (i.e., coppice, coppice-with-standards, and high forest). The definitions of these structures are provided in Annex 3.

2.1.3 Tree species composition

This data was available in *Daubrée* statistics and current forest inventory data. In the first one, forest area of a given NUTS-3 unit was split by forest tree species at a 10% resolution, in two broad ownership categories: (i) public forests subjected to the forest regime, (ii) forests not subjected to the forest regime (predominantly private forests). Tree species listed in *Daubrée* statistics varied across NUTS-3 units and were partially described by genus or common names, requesting technical harmonization (Annex 4).

In NFI data, plots (and associated estimated area) are described by a dominant species of the canopy layer of the census strata (trees with diameter ≥ 7.5 cm). 3.3% of total forest area has a non-identified dominant species (open forests).

2.1.4 Statistics for *Bas-Rhin*, *Haut-Rhin*, and *Moselle* units

Territories of three NUTS-3 units (*Bas-Rhin*, *Haut-Rhin* and *Moselle*) were annexed by Germany between 1870 and 1918, and their forests were not described in *Daubrée* statistics and in decennial agricultural statistics of 1892. Owing to the existence of the German statistics of 1907 (Ministerium für Elsaß-Lothringen 1909), we supplemented the statistics to cover the whole territory in *Daubrée* statistics.

German statistics provided information on the legal category of ownership (State-owned forests, municipal forests, and private forests). However, forest areas were not split by

Table 1 Description of variables found in ancient forest statistics and national forest inventory

Statistics	Data resolution	Tree species	Forest area split by	
			Ownership category	Forest management type (or structure type for NFI data)
Decennial agricultural statistics (1892)	86 NUTS-3	—	MPBNS	SC / CWS / HBF / HCF
			PNS	SC / CWS / HBF / HCF
			MPBS	SC / CC / CWS / HF / DPUD
			SFS	SC / CC / CWS / HF / DPUD
German statistics (1907) for <i>Bas-Rhin</i> , <i>Haut-Rhin</i> and <i>Moselle</i> units	3 NUTS-3 (<i>Bas-Rhin</i> , <i>Haut-Rhin</i> and <i>Moselle</i>)	—	M	—
			P	—
			SFS	—
<i>Daubrée</i> statistics (1908)	86 NUTS-3, LAU-1 and arrondissements of Paris	15 main tree species and two tree species botanic class (<i>various broadleaves</i> and <i>various conifers</i>)	MPBNS	SC / CC / CWS / HF / UA
			MPBS	SC / CC / CWS / HF / UA
			PNS	SC / CC / CWS / HF / UA
			SFS	SC / CC / CWS / HF / UA
Decennial agricultural statistics (1929)	89 NUTS-3	—	MNS	SC / CWS / HBF / HCF / RF
			MPBNS	SC / CWS / HBF / HCF / RF
			PBNS	SC / CWS / HBF / HCF / RF
			PNS	SC / CWS / HBF / HCF / RF
			S	SC / CWS / HBF / HCF / RF
NFI (2008 ¹)	90 NUTS-3	70 tree species	M	C / MCHF / EAHF / UAHF / NS
			P	C / MCHF / EAHF / UAHF / NS
			SF	C / MCHF / EAHF / UAHF / NS

—: not available variables. ¹ median year

Ownership categories: *S* forests subjected to the forest regime, *SF* state-owned forests, *SFS* state-owned forests subjected to the forest regime, *M* municipal forests (and other territorial public bodies' forests), *MPBS* municipal forests and other territorial public bodies' forests subjected to the forest regime, *MNS*: municipal forests not subjected to the forest regime, *MPBNS* municipal forests and other territorial public bodies' forests not subjected to the forest regime, *PBNS* other territorial public bodies' forests not subjected to the forest regime, *P* private forests, *PNS* private forests not subjected to the forest regime. Forest management types: *C* coppice, *SC* simple coppice, *CC* coppice conversion (into high forest), *CWS* coppice-with-standards, *MCHF* mixture of coppice and high-forest, *HF* high forest, *HBF* high broadleaved dominated forest, *HCF*, high coniferous dominated forest, *EAHF*, even-aged high forest, *UAHF* uneven-aged high forest, *DPUD* decreed perimeters and unproductive coastal dunes, *RF*: recently afforested forest area, *UA* unproductive area

management type. Thus, German data were used in relation with *Daubrée* statistics solely for information on the total forest area and on forest area split by ownership category.

Bas-Rhin, *Haut-Rhin*, and *Moselle* NUTS-3 units in the 1892 statistics could not be supplemented by German data.

Table 2 Summary of data found in ancient forest statistics and national forest inventory. We provide minimum and maximum values (in square brackets) at NUTS-3 unit scale. Also, we provide totals for the whole of France

	Decennial agricultural statistics (1892)	German statistics (1907) for <i>Bas-Rhin</i> , <i>Haut-Rhin</i> and <i>Moselle</i> units	<i>Daubrée</i> statistics (1908)	Decennial agricultural statistics (1929)	NFI (2008 ¹)
Total forest area (ha)	[2158–522,768] 9,520,000	[125,201–160,485] 445,000	[1810–516,608] 9890000	[2122–626,836] 10,675,000	[25,632–567,423] 16,155,000
Total annual harvested volume (m ³)	[10,596–1,774,267] 27,600,000	—	[2296–1,904,118] 23,500,000	[0–883,400] 4,700,000 ²	[20,070–4,649,477] 44,500,000
Number of private owners	—	—	[73–67,207] 1,530,000	[0–49,589] 1,210,000	—
Afforestation rate (%)	[3.5–56.1] 18.0	—	[3.3–55.4] 18.7	[3.0–66.9] 19.4	[5.1–62.5] 29.4

NFI national forest inventory. —:s not available data. ¹ median year. ² only for timber wood

2.2 Data entry and error correction

Ancient statistics were available in paper textbook format and image format. Areal data were given for each NUTS-3 unit (table rows) and intersections of forest management type and ownership categories (table columns). Marginal totals were also provided. Digitization of the three statistics was the first step of the present work (Audinot et al. 2020). Data from *Daubrée* statistics were digitized only at a NUTS-3 unit. Initial typing errors of different types were identified in ancient statistics by comparing row/column totals, and were corrected. They are summarized in Annex 5.

2.3 Harmonization of statistics

For the purpose of cross-comparison of forest area across the three ancient statistics, categorical variables were harmonized including tree species composition, forest management type, and ownership categories (Annex 4).

As an overview, forest areas by forest management type were found in equal proportions among the ancient statistics (Fig. 1), except for the other forests' areas found smaller, mainly in private forests (0% in 1892, 4.3% in 1908, and 2.1% in 1929, Table 3) in line with their heterogeneity. By contrast, high forests that formed a restricted category in ancient statistics are currently predominant (53%) in the total forest area (Fig. 1). Forest ownership categories were distributed in roughly the same proportions between the three ancient statistics (68% of the forest area was private in 1892, 66% in 1908, and 67% in 1929; Fig. 1).

Moreover, NUTS-3 units' boundaries were unstable between ancient statistics. The standardization of NUTS-3 units' boundaries and the analysis of their potential effects of the forest area changes are provided in Annex 6.

2.4 Data analysis

2.4.1 Principle

Forest area differences between successive statistics, including NFI data, were computed, with NUTS-3 units as the reference population. Forest area differences were analyzed at NUTS-3 level, as a whole, or conditionally to forest management type and ownership categories. Paired *t* tests were used to analyze forest area differences. Maps at NUTS-3 level were produced to detect spatial structures of forest changes.

2.4.2 Cross-comparison of ancient statistics

We analyzed forest area differences over two sets of successive statistics: the ancient periods 1892–1908 and 1908–1929. The following consistency criteria were considered:

- absence of erratic variations between forest area differences at successive periods (1892–1908 and 1908–1929 periods);
- absence of bias in the variables of interest documented in the three ancient statistics (forest management type and ownership category);
- accordance of forest area differences (direction and magnitude) with forest policies at play at both periods (1892–1908 and 1908–1929 periods), and with forest transition features.

2.4.3 Long-term changes in forest area

Forest area differences by NUTS-3 units, forest management types, and ownership categories were also analyzed over two long periods: 1892–1929 (ca. 40 years) and 1908–2008 (ca. 100 years). Over the 1908–2008 period, we analyzed forest area differences at two levels of tree-species composition including botanical classes and tree species.

A synoptic view of periods studied, corresponding statistics, and NUTS-3 populations considered is summarized in Table 4.

3 Results


3.1 Cross-comparison of ancient statistics (1892–1908 and 1908–1929 periods)

3.1.1 Cross-comparison of forest statistics at NUTS-3 units level

Positive forest area differences were found in many NUTS-3 units in 1908–1929 period (70%, Fig. 2), less in 1892–1908 period (49%, Fig. 2). However, the average difference in NUTS-3 forest area was positive and significant at both periods ($p < 0.05$, Table 5). Strong standard-deviations of forest area differences indicated large heterogeneity across NUTS-3 units (Table 5). Yet, differences lower than -2000 ha of forest at both periods were only observed for two NUTS-3 units (Pyrénées-Atlantiques and Gers).

An obvious spatial structure in positive/negative differences was identified (Fig. 2). Over the 1892–1908 period, most NUTS-3 units showing positive differences were situated in Eastern France, essentially formed of public forests, with negative differences located in Western and central France, where forests remain largely private. Over the 1908–1929 period, positive differences extended to northern France, with differences found to reverse in Central and Southern France. Differences in Western France turned null or positive (Fig. 2). Also, some southern NUTS-3 units showed negative

Fig. 1 French forest area split by forest management (a), ownership (b), and species composition (c) categories according to three ancient statistics (1892, 1908, and 1929) and national forest inventory data (2008). *Bas-Rhin*, *Haut-Rhin*, and *Moselle* NUTS-3 units were excluded as these were not documented in the *Daubr e* statistics (German annexation), Corsican NUTS-3, and those surrounding Paris were considered as a single unit. Data about tree species composition do not exist in 1892 and 1929 statistics



differences after positive differences in the previous period (for example forest area in Ari ge with + 5000 ha/− 60,000 ha over 1892–1908/1908–1929 periods, respectively). Such shifts suggested a bias in *Daubr e* statistics relative to other decennial agricultural statistics, corresponding to underestimation of private forests.

3.1.2 Cross-comparison of forest statistics by forest management type

The analysis of forest area differences by forest management types showed significant and inverse differences between coppice forests (\pm over 1892–1908/1908–1929 periods;

Table 3 Summary statistics for the French forest area according to the four data sources under study.

			Forest area ($\times 10^3$ ha)				
			Decennial agricultural statistics (1892)	German statistics(1907) for <i>Bas-Rhin</i> , <i>Haut-Rhin</i> and <i>Moselle</i> units	<i>Daubr�e</i> statistics (1908) ¹	Decennial agricultural statistics (1929)	NFI (2008 ²)
Forest management type by ownership category	Private forest	Coppice	3210	90 ³	2050	2840	1535
		Coppice-with-standards	1515		2430	1900	3420
		High forest	1790		1980	2315	5765
		Other forests	0		280	145	1275
	State-owned forest subjected to the forest regime	Coppice	20	155 ³	25	—	45
		Coppice-with-standards	410		405	—	235
		High forest	460		620	—	1135
	Other public forest subjected to the forest regime	Other forests	200		150	—	85
		Coppice	270	200 ³	260	370	190
		Coppice-with-standards	1000		1015	1370	610
	Public forest in total subjected to the forest regime	High forest	575		600	1585	1635
		Other forests	70		75	150	225
		Coppice	290	355 ³	285	370	235
		Coppice-with-standards	1410		1420	1370	845
			High forest	1035		1220	1585
		Other forests	270		225	150	310
Totals			9520	445	9890	10,675	16,155

¹ without *Bas-Rhin*, *Haut-Rhin*, and *Moselle* NUTS-3 units. ² median year. ³ breakdown by forest management type not provided in original data. —: not available data

NFI national forest inventory

$p < 0.001$) and coppice-with-standards forests (\pm over the same periods; $p < 0.001$, Table 5). Hence, 87% of the NUTS-3 units had negative differences over the 1892–1908 period in coppice forests, and conversely 71% of these units had positive differences over the “1908–1929” period Fig. 3). In coppice-with-standards forests, 76% of the NUTS-3 units


had positive differences over the 1892–1908 period, while 74% of these had negative differences over the 1908–1929 period, Fig. 3). *Daubr e* statistics thus systematically overestimated the area of coppice-with-standards forests at the expense of those of coppice forests, as compared to other decennials agricultural statistics.

Table 4 Statistics and population of NUTS-3 units used by study periods, The transition from 89 to 86 department administrative units (NUTS-3) is due to the annexation of the *Alsace-Moselle* region. *Gironde* was excluded from the whole paired *t* tests but it was kept in all other calculations (Annex 4)

Periods	Statistics used				Number of NUTS-3 used in analyzes			
	Decennial agricultural statistics (1892)	<i>Daubr�e</i> statistics (1908)	Decennial agricultural statistics (1929)	NFI (2008 ¹)	Whole French forest	Ownership category	Forest management type	Tree species
1892–1908	X	X			86	86	86	
1908–1929		X	X		89	89	86	
1892–1929	X		X		86	86	86	
1908–2008		X		X	89	89	86	86

NFI national forest inventory. ¹ median year. X: statistics used

Fig. 2 Forest area differences across NUTS-3 units of the territory over the 1892–1908 and 1908–1929 periods of the statistical cross-comparison. See Fig. 1 for details on NUTS-3 data



The large standard-deviations of forest area differences found in other forests' areas despite their small areas (Fig. 3 and Table 3) pointed out to the heterogeneous definition of this forest management type (Table 5). For example, other forests' areas in *Gironde* and *Landes* showed large differences over the one or two periods (+ 78,000 ha over the 1892–1908 period and – 73,000 ha over the 1908–1929 period in *Gironde*; + 1400 ha over the 1892–1908 period and + 58,000 ha over the 1908–1929 period in *Landes*) at a time of active afforestation, bound to explain these strong differences.

In contrast, means of forest area differences were much more constant in high forests (Table 5) and both positive and statistically significant ($p < 0.01$ for 1892–1908; $p < 0.05$ for 1908–1929, Table 5). Positive differences represented 56% of NUTS-3 units over the 1892–1908 period and 76% units over the 1908–1929 period (Fig. 3).

3.1.3 Cross-comparison of forest statistics by ownership category

Over 1892–1908 and 1908–1929 periods, a substantial fraction of NUTS-3 units showed positive forest area differences in public forests (64% of NUTS-3 units over the 1892–1908 period, 69% of NUTS-3 units over the 1908–1929 period, Fig. 3), however, statistically significant only in the first period ($p < 0.01$, Table 5). The strong increase in public forests area

over the 1892–1908 period was in agreement with the spatial structure identified in the Eastern half of France (Fig. 2) where there is a large share of public forest.

Fifty-five percent of NUTS-3 units in private forests also showed a negative forest area differences over the 1892–1908 period. Among them, 68% then had positive differences over the period 1908–1929. Means of forest area differences means between these NUTS-3 units showed inverted differences over the two periods (– 5200 ha/NUTS-3 units over the 1892–1908 period and + 4900 ha/NUTS-3 units over the 1908–1929 period), but these differences were statistically significant only for the 1908–1929 period ($p < 0.01$, Table 5). These results confirmed the underestimation of private forests areas in the *Daubr e* statistics, as inferred in 3.1.1.

Area of private forest in *Gironde* and *Landes* (southwestern France) last showed tremendous differences at a period of active afforestation (+ 110,000 ha in *Gironde* over the 1892–1908 period, + 110,000 ha in *Landes* over the 1908–1929 period).

3.2 Long-term changes in forest area (1892–1929 and 1908–2008 periods)

3.2.1 Historical differences in whole French forest area

Significantly positive forest area differences in total forest area were identified at both periods ($p < 0.001$ in

Table 5 Summary of forest area differences across NUTS-3 units over the four periods under study, Forest area differences calculated for the 1892–1908, 1908–1929, 1892–1929, and 1908–2008 periods according department administrative units (NUTS-3) population. Significance assessed from paired *t* tests

	1892–1908			1908–1929			1892–1929			1908–2008		
	FAD mean ($\times 10^3$ ha)	FAD standard-deviation ($\times 10^3$ ha)	Paired <i>t</i> test	FAD mean ($\times 10^3$ ha)	FAD standard-deviation ($\times 10^3$ ha)	Paired <i>t</i> test	FAD mean ($\times 10^3$ ha)	FAD standard-deviation ($\times 10^3$ ha)	Paired <i>t</i> test	FAD mean ($\times 10^3$ ha)	FAD standard-deviation ($\times 10^3$ ha)	Paired <i>t</i> test
Total forest area	+4.2	16.5	<0.05	+3.8	15.2	<0.05	+8.1	20.7	<0.001	+65.4	57.4	<10 ⁻¹⁰
Ownership category	+1.6	4.8	<0.01	-0.4	8.6	NS	+1.2	9.2	NS	+7.4	10.6	<10 ⁻⁸
Private forest	+2.6	15.9	NS	+4.2	14.2	<0.01	+6.8	20.3	<0.01	+58.0	56.2	<10 ⁻¹⁰
Coppice	-13.6	15.4	<10 ⁻¹⁰	+10.0	17.8	<10 ⁻⁵	-3.6	17.0	<0.1	-6.6	19.3	<0.01
Coppice-with-standards	+10.8	15.0	<10 ⁻⁸	-8.1	14.8	<10 ⁻⁵	+2.6	15.0	NS	+4.3	47.2	NS
High forest	+4.3	11.7	<0.01	+4.5	15.5	<0.05	+8.8	18.8	<10 ⁻⁴	+57.0	40.5	<10 ⁻¹⁰
Other forests	+2.8	9.1	<0.01	-2.5	13.2	<0.1	+0.3	10.5	NS	+14.7	12.5	<10 ⁻⁵

FAD forest area difference

1892–1929 period and $p < 10^{-10}$ in 1908–2008 period, Table 5), suggesting a continuous increase in forest area since 1892. The magnitude of the total forest area difference at 1892–1929 period was +700,000 ha, this corresponding to a relative increase by +7% and to an annual rate of change by +0.19% over the 1892–1929 period. However, large standard-deviations of forest area differences indicated that not all NUTS-3 units were subjected to positive differences over this period (30% of NUTS-3 units showed negative differences). Also, the total forest area differences in 1908–2008 period was around +5,800,000 ha corresponding to a relative difference by +36% and to an annual rate of change by +0.45%, thus foot-printing an acceleration in forest areal increases between 1892 and 1929 and 1908–2008 periods ($p < 10^{-7}$).

3.2.2 Historical differences in forest area by forest management type

Forest area differences at both periods showed a decline in coppice forests on almost the whole territory, together with a large increase in high-forests over the 1908–2008 period, indicating a massive change in the structure of these forests over the period. In 1892, the area of coppice forests was of 3,500,000 ha. About one century later (2008), coppice forests have reached 1,700,000 ha, as documented by NFI data. Assuming that ancient coppice forests have fully evolved into high forests, and since the total increase in forest area is 5,800,000 ha, this suggested that an additional +4,100,000 ha of high forests had another origin than coppice conversion since 1908. High forests, therefore, have formed the majority of afforested or returning forests since 1908 (Fig. 1 and Fig. 4).

Forest area differences in coppice-with-standard forests showed a North-South division over the 1908–2008 period (Fig. 4), with a marked decrease in Northern and Northeastern France (-1,200,000 ha), where the area of high forests has increased (+2,500,000 ha), and a conversely marked increase in South France and Brittany (+1,600,000 ha) where coppice forests area has decreased (-360,000 ha). Maps of forest area differences therefore suggested similar maturation in forest structure (coppice-coppice-with-standards-high forest) across the whole territory, though with a much longer apparent delay in southern France.

3.2.3 Historical differences in forest area by ownership category

Increase in private forests area was found substantial (corresponding to an annual rate of change +0.27%/+




Fig. 3 Comparison of successive 1892–1908 and 1908–1929 periods forest area differences at a NUTS-3 level across forest management and ownership categories, See Fig. 1 for details on NUTS-3 data, FAD, forest

area differences. **a** FAD in public forests. **b**: FAD in private forests. **c** FAD in coppice forests, **d** FAD in coppice-with-standards forests, **e** FAD in high forests, **f** FAD for other forests (see Annex 4 of the manuscript)

0.56% over 1892–1929/1908–2008 periods, respectively) and concerned 97% of NUTS-3 units over the

1908–2008 period. In 1908, the area of private forests was of 6,800,000 ha. About one century later (2008),




Fig. 4 Changes in forest area classified by forest management type across NUTS-3 units of the territory over the 1892–1929 and 1908–2008 periods, See Fig. 1 for details on NUTS-3 data

private forests area has almost doubled, reaching 12,000,000 ha. In private forests, mean of forest area differences was very strongly significant over the 1908–2008 period ($p < 10^{-10}$, Table 5). As a consequence, whereas two-thirds of French forests were private in ancient statistics, three-fourths of the forest area is nowadays private (Fig. 1).


Public forests also revealed significant differences over the 1908–2008 period (+ 700,000 ha, $p < 10^{-8}$, annual rate of change by +0.17%, Table 5). The slight increase over 1892–1929 period was not statistically

significant ($p = 0.21$, annual rate of change by +0.39%, Table 5).

3.3 Long-term changes in forest composition

Broadleaf- and conifer-dominated forests showed significant positive differences over the 1908–2008 period (Fig. 5) with extension of broadleaved forests twice as intense in magnitude as in conifer forests (+ 3,600,000 ha, $p < 10^{-10}$ and + 1,600,000 ha, $p < 10^{-7}$, respectively), yet favored in afforestation

Fig. 5 Changes in broadleaf- and conifer-dominated forest area over the 1908–2008 period across NUTS-3 units of the territory, See Fig. 1 for details on NUTS-3 data



programs. Consistently, various broadleaves also progressed three times faster than various conifers (Table 6 in Annex 4). These relative changes were strikingly similar (+34/+38% in broadleaved/conifer forests), making the proportion of these forests unchanged over one century (Fig. 1).

Tree species (without various broadleaves and various conifers) showed a significant positive differences over 100 years (+3,900,000 ha or 67% of total extension, $p < 10^{-10}$), indicating that species identified in *Daubr e* statistics had been playing a role in forest extension.


Significant positive areal changes were detected in all tree species/species groups but hornbeam (*Carpinus betulus* L.) and cork oak (*Quercus suber* L.) showing negative and non-significant negative changes (Fig. 6 and Table 6 in Annex 4), two species that were important contributors to coppice production systems in France.

White oaks showed sharp significant positive areal changes (Fig. 6 and Table 6 in Annex 4) in a large part of territory (81% of NUTS-3 units). White oaks privileged in both coppice-with-standards and high forests took advantage of changes in the structure of the French forest.

Norway spruce (*Picea abies* Karst.) and pine species used in afforestation programs showed areal increases by +40 to +70% larger than those of, e.g., Silver fir (*Abies alba* Mill.), a coniferous tree species regenerated naturally (Fig. 6 and Table 6 in Annex 4). Also, European larch (*Larix decidua* Mill.) privileged in landscape protection programs of the nineteenth century still showed progression after *Daubr e* statistics (Fig. 6 and Table 6 in Annex 4).

In summary, the strong absolute progression of broadleaved-dominated forests was emphasized, with afforested coniferous tree species and coppice broadleaved species found at extreme magnitudes.

Fig. 6 Changes in forest area classified by dominating tree species/tree species groups of the French forests over the 1908–2008 period, See Fig. 1 for details on NUTS-3 data. Significance of forest area differences was assessed by paired *t* tests across the NUTS-3 population. Individual tree species not identified in the tree species groups white oaks, pines, various broadleaves and various conifers, (*): *p* value < 0.1. ***p* value < 0.05. ****p* value < 0.01. *****p* value < 0.001



4 Discussion

4.1 Statistical value of ancient statistics

This study is based on historical sources of forest information: *Daubrée* statistics (1908) completed by a German forest statistic (1907) for *Bas-Rhin*, *Haut-Rhin*, and *Moselle* NUTS-3 units and decennial agricultural statistics of 1892 and 1929. Their supplementations with modern NFI data offered an extended overview on broad long-term forest changes, and in this sense, also increase the value of modern data.

The lack of a clear forest definition and of a specific description of methods used at that time to produce ancient statistics questions their consistency. Also, the absence of foundations in survey sampling at that time

period (Frayer and Furnival 1999) and of subsequent error estimation makes these statistics of lower statistical value than modern forest inventory. However, over a long time period, changes in the forest can turn strong enough for these statistics' uncertainties to appear much lower in magnitude than the main shifts detected (Table 5). This suggests their practical use to identify broad patterns and the footprints of past forest policies and natural processes.

Also, despite the reduced availability of information technologies at that time (calculator, computer), only few errors were detected in all of the ancient statistics in comparison to the total number of digits entered (Annex 5). The rate of errors in primary unit figures however remains out of reach.

4.2 Cross-comparison of ancient statistics

4.2.1 Potential systematic biases in *Daubrée* statistics

Forest area differences across NUTS-3 units showed systematic positive shifts in forest area at both periods (1892–1908 and 1908–1929, Table 5). All successive statistics suggested an increase in the total forest area, in line with current knowledge of the French forest transition (Cinotti 1996; Denardou et al. 2017). Yet, many of these units experienced successive increases/decreases in area over the two periods (notably NUTS-3 units of Southern France, Fig. 2), suggesting that either *Daubrée* statistics is biased positively, or decennial agricultural statistics are so, negatively. Over-estimation of forest area in *Daubrée* statistics may have several causes: a lower attention paid to forests of lower economic interest like Mediterranean forests of southern France or for mountain forest areas, also poorly described in administrative documents (Alps, Pyrenees, and Corsica). Negative forest area differences on 1892–1929 period followed by positives differences on 1908–1929 period affected Massif Central (Fig. 2), mainly private (Dodane 2009) and as such prone to lower attention in *Daubrée* statistics. These regional heterogeneities are discussed in Denardou et al. (2017).

4.2.2 Interplay between political will and past forest reporting

Cross-comparisons suggested a relative bias in the forest area of coppice/coppice-with-standards in *Daubrée* statistics with respect to other decennial agricultural statistics (Fig. 3). Over the 1892–1908 period, coppice forests regressed by 1,200,000 ha and coppice-with-standards forests increased by 930,000 ha. Over the 1908–1929 period, however, coppice forests increased by 860,000 ha and coppice-with-standards forests decreased by 700,000 ha. This over-estimation of coppice-with-standards (including coppice conversion) pattern in *Daubrée* statistics may find root in the coppice conversion policy, hindered by political and economic issues, which promoted a climate of suspicion between the State and the forest administration (Degron 1998). Such context is bound to have led to over-estimations of the conversion effort for better agreement with policy objectives (Corvol 2013). Moreover, the very short study period (1892–1908–1929) with respect to the pace of coppice conversion into high forest makes this evaluation highly dependent on survey processes.

The moderate negative bias in private forests in *Daubrée* statistics (Fig. 3) could stem from forest owners' distrust towards State policies. The difficult

economic context of that period (Degron 1998) pushed the State to reflect on the privatization of public forests (Dabat 1920), or to interfere in the management of private forests (Corvol 2013), encouraging private owners to voluntarily underestimate the extent of their forests.

4.3 Increases in the forest area

4.3.1 Returning forests

Forest area differences over a longer time scale (1892–1929 and 1908–2008 periods) highlighted the sharp significant increase in forest area (+5,800,000 in whole French forest area and an average +65,000 ha/NUTS-3 unit over the 1908–2008 period; $p < 10^{-10}$, Table 5) and particularly in private forests (+5,100,000 in whole French forest area and +58,000/NUTS-3 unit over the 1908–2008 period; $p < 10^{-10}$, Table 5). Abandonment of agriculture is alleged to be one of the main drivers of this increase (Chakir and Madignier 2006; Cavailhès and Normandin 1993) and remains acute in rural areas (Denardou et al. 2017).

4.3.2 Afforestation programs

Major afforestation programs have taken place during 19th and 20th centuries including the landscape protection of the RTM program (1860 and 1882, Larrère et al. 1980) which led to planting of 170,000 ha by 1914 (Métaillié 1999), and the much more extended FFN program (1946–1999) intended to foster the French forest industry and which reached a net 1.2 million hectares (Dodane 2009). This total afforestation (1400,000 ha) only represented 24% of total forest area increase (5,800,000 ha) over the 1908–2008 period room for undocumented secondary afforestation programs (e.g., *national lottery* program in 1920, Bazire and Gadant 1991), or more largely to natural expansion processes.

4.4 Structural and compositional changes in forests

4.4.1 Structural forest changes

Regarding forest management, the shift between coppice and high forest area over both long periods 1892–1929 and 1908–2008 is obvious (Fig. 4)—while at different stages depending on regions—and it allows concluding that forest conversion (1908–2008 period: –600,000 ha in coppice, $p < 0.01$ and +4,900,000 ha in high forest, $p < 10^{-10}$; Table 5) over the territory (Fig. 4) has been a very long-lasting and still ongoing process. Of importance, such process is bound to also have also dramatically increased the forest growing stock (Pignard 2000; Forest Europe 2015).

4.4.2 Compositional forest changes

The two main afforestation policies (section 4.3.2) mainly concerned conifer plantations. For example, the area of Norway spruce plantations in the Massif Central has increased from 360,000 to 610,000 ha over 20 years (Bazire and Gadant 1991). This massive afforestation has supported the idea, widespread in France and Europe (Moriniaux 1999) of an increase in conifer prevalence, at the expense of broadleaves. Whereas we confirmed the significant expansion of conifer-dominated forests (+ 1,600,000 ha, $p < 10^{-7}$), we also showed a parallel massive expansion of broadleaf-dominated forests (+ 3,600,000 ha, $p < 10^{-14}$) over the 1908–2008 period (Fig. 5). Hence, the proportion of broadleaf- and conifer-dominated forests has not changed in a century (Fig. 1), conflicting the view of conifer progression across the territory.

The significant sharp increase in *white oaks* areas (+ 1,500,000 ha over the 1908–2008 period, Fig. 6 and Table 6 in Annex 4) has a dynamic interpretation, as oak species are found in both coppice-with-standards forests and high forests owing to their ability to grow from sprouts (Boppe 1886), and their wood quality has led to favor them in the conversion of coppice. The non-significant small decrease of hornbeam in the 1908–2008 period (– 35,000 ha, Fig. 6 and Table 6 in Annex 4), a species predominantly found in coppice and coppice-with-standards forests due to its ability to sprout (Boppe 1886), is in accordance with conversion in favor of oaks. In addition, maintaining an understory of hornbeam benefits to the oak silviculture (Office National des Forêts 2010).

The significant increases of area in Norway spruce (+ 340,000 ha, Fig. 6 and Table 6 in Annex 4) and in European larch (+ 40,000 ha, Fig. 6 and Table 6 in Annex 4) reflect afforestation conducted by the FFN (Bazire and Gadant 1991; Bouchon 1984). Parallel significant increases in chestnut (*Castanea sativa* Mill., + 520,000 ha), *various broadleaves* (+ 1,000,000 ha), and *various conifers* (+ 320,000 ha) areas (Fig. 6 and Table 6 in Annex 4) can also reflect an effect of tree species diversification in afforestation programs (de Rochebouet 1987).

Note that declining cork oak areas can be explained by a decline in the demand for oak cork (Bugalho et al. 2011). Also, the mortality of cork oak has increased in the Mediterranean region, due to the increased intensity of droughts (Costa et al. 2010).

Statistics over this 1908–2008 period therefore suggested that intensive afforestation programs (RTM, FFN) did not develop at the expense of tree species diversity (Fig. 6 and Table 6 in Annex 4), but instead allowed for a tree species diversity increase at country scale, by introducing new tree species while maintaining and even increasing prevalence of the others. The illustration that of forest transition can increase tree diversity as a country scale is major.

5 Conclusion

- Cross-comparison of the three ancient French forest statistics showed that their exploitation was effective. Specific inconsistencies between the three statistics highlighted directional biases in forest management types and ownership categories, namely in *Daubrée* statistics, and must be dealt with when exploiting the data.
- These ancient statistics can support many fields of research. Examples include the impacts of the two World Wars on forests, the evolution of tree biodiversity, and spatial analyses of land-use change.
- The study of forest-area changes over more than one a century, using the three ancient statistics and NFI data (2008) highlighted features of the French forest expansion, found mainly in private forests, and characterized a slow conversion of coppices into high forests on the whole territory. Moreover, a statistically significant increases in the area of all major tree species found in the early twentieth century, with only minor areal decreases in two species common in coppice systems (hornbeam and cork oak) together with a clear effect of conifer afforestation programs (Restoration of Mountain Terrain and National Forest Fund) were identified, highlighting the major outcome of a long-term increase in major tree diversity across the whole territory.

Acknowledgments We are particularly grateful to Anaïs Denardou-Tisserand for her contribution to data analysis.

Authors' Contributions Conceptualization: Jean-Daniel Bontemps; Methodology: Jean-Daniel Bontemps, Timothée

Audinot; Validation: Holger Wernsdörfer, Jean-Daniel Bontemps; Formal Analysis: Timothée Audinot, Jean-Daniel Bontemps; Investigation: Timothée Audinot, Jean-Daniel Bontemps,

Holger Wernsdörfer; Data curation: Jean-Daniel Bontemps; Writing – original draft:

Timothée Audinot, Jean-Daniel Bontemps, Holger Wernsdörfer; Supervision: Jean-Daniel Bontemps, Holger Wernsdörfer; Project Administration: Jean-Daniel Bontemps, Holger Wernsdörfer; Funding acquisition: Jean-Daniel Bontemps

Funding information This study and PhD project are funded by Institut National de l'Information Géographique et Forestière (IGN). The Laboratory of Forest Inventory and UMR SILVA are supported by grants overseen by the French National Research Agency (ANR) as part of the "Investissements d'Avenir" program (ANR-11-LABX-0002-01, Lab of Excellence ARBRE).

Data availability The datasets generated and analyzed during the current study are available in the Zenodo repository, <https://doi.org/10.5281/zenodo.3739458>

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Annex 1: Overview of forest data tables in the three ancient statistics as images

DÉPARTEMENTS.	SURFACE TOTALE.	RÉPARTITION DE LA SURFACE TOTALE.					PRODUCTION ANNUELLE.		
		SURFACES IMPRODUCTIVES (rochers, marais, tourbières, etc.).	TAILLIS.			FUTAIE.	BOIS D'ŒUVRE.	BOIS DE FEU.	TOTAL.
			SIMPLE, sarlé ou fureté.	SOUS FUTAIE.	en CONVERSION.				
hectares.	hectares.	hectares.	hectares.	hectares.	hectares.	mètres cubes.	mètres cubes.	mètres cubes.	
Ain.....	69,873	3,382	29,375	29,051	726	7,339	21,095	115,730	136,825
Aisne.....	74,999	1,453	10,155	61,103	149	2,139	60,176	160,316	220,492
Allier.....	53,866	1,198	6,241	43,316	23	3,088	18,596	145,001	163,597
Alpes (Basses-).....	67,606	1,977	53,643	1,813	—	10,173	5,374	44,181	49,555
Alpes (Hautes-).....	27,438	1,511	18,698	63	11	7,155	4,207	12,136	16,343
Alpes-Maritimes.....	43,499	5,558	10,527	2,784	—	24,630	5,483	23,210	28,693
Ardèche.....	76,269	1,018	45,098	2,279	—	27,874	41,130	58,195	99,325
Ardennes.....	78,474	1,127	15,313	50,849	—	11,185	75,450	172,818	248,268
Ariège.....	61,399	159	21,477	34,584	839	4,340	8,207	55,704	63,911
Aube.....	94,783	2,895	5,551	57,792	44	28,501	51,154	184,175	235,329
Aude.....	35,339	1,904	24,859	1,317	—	7,529	19,353	49,136	68,489
Aveyron.....	68,679	2,170	31,474	10,088	7,212	17,735	10,261	88,594	98,855
Belfort (Territoire de).....	8,019	95	2,451	4,021	264	1,188	2,667	21,908	24,575
Bouches-du-Rhône.....	60,539	4,350	18,513	—	—	37,676	16,459	40,227	57,276
Calvados.....	34,800	2,487	7,976	21,029	71	3,237	3,941	77,681	81,622
Cantal.....	48,595	1,939	14,126	7,241	715	24,574	19,431	38,427	57,858
Charente.....	78,052	4,860	28,441	42,635	—	2,116	8,984	180,955	189,939
Charente-Inférieure.....	72,036	2,487	36,409	21,851	—	11,289	20,406	227,451	247,857
Cher.....	103,913	132	14,750	84,644	42	4,345	40,203	278,035	318,238
Corrèze.....	103,366	2,285	20,714	9,900	107	70,360	5,413	45,951	51,364
Corse.....	29,989	2,176	4,578	5,463	4	17,768	12,264	21,070	34,334
Côte-d'Or.....	116,411	403	2,863	103,981	25	9,139	54,894	290,357	345,251
Côtes-du-Nord.....	32,387	1,168	17,164	9,297	7	4,741	8,905	86,711	95,616
Creuse.....	30,589	445	10,554	13,042	871	5,677	12,363	79,997	92,360
Dordogne.....	255,672	689	104,506	80,199	—	70,278	163,304	327,339	490,643
Doubs.....	40,985	178	8,200	14,057	2,101	16,449	44,421	75,532	119,953
Drôme.....	123,485	5,077	106,959	511	1,349	9,589	7,614	118,487	126,101
Eure.....	98,707	1,140	19,153	71,247	—	7,167	23,302	193,821	217,123
Eure-et-Loir.....	54,273	556	20,396	32,147	74	1,100	8,767	104,080	112,847
Finistère.....	25,926	17	22,734	—	—	3,175	4,509	71,561	76,070
Gard.....	90,715	1,057	68,474	14,857	—	6,327	17,987	70,057	88,044
Garonne (Haute-).....	49,953	45	9,246	37,129	1,245	2,288	3,878	96,316	100,194
Gers.....	46,045	196	13,151	29,836	—	2,862	7,763	83,226	91,189
Gironde.....	407,299	74,657	41,523	34,655	—	256,464	594,168	790,630	1,384,798
Hérault.....	62,107	7,886	52,840	—	—	1,381	1,331	29,010	30,341
Ille-et-Vilaine.....	36,861	1,188	14,656	15,982	—	5,035	10,470	92,760	103,230
Indre.....	67,191	10	—	65,230	—	1,951	11,988	181,694	193,682
Indre-et-Loire.....	86,873	5,065	60,750	17,762	227	3,069	6,325	157,625	163,950
Isère.....	119,987	4,364	72,957	18,903	6,000	17,763	47,527	202,667	250,194
Jura.....	57,692	7,475	8,700	23,322	5,364	12,831	40,825	95,984	136,809
Landes.....	433,275	1,255	23,091	16,757	—	392,172	1,186,846	436,839	1,623,685
Loir-et-Cher.....	129,564	5,889	21,646	73,435	6,180	22,414	49,219	238,571	287,790
Loire.....	50,811	632	10,374	8,796	100	30,909	147,075	56,543	203,618

Fig. 7 Overview of a data table presented in *Daubrée* statistics at a NUTS-3 level

VI. -- BOIS ET RÉGIME FORESTIER.

BOIS SOUMIS AU RÉGIME FORESTIER.

COMMUNES ET ÉTABLISSEMENTS PUBLICS.

BOIS DE L'ÉTAT.

DÉPARTEMENTS.	COMMUNES ET ÉTABLISSEMENTS PUBLICS.										BOIS DE L'ÉTAT.																					
	SUPERFICIL.					PRODUCTION LIGNIERE TOTALE ANUELLE.					LIGES, ÉCORCES, RÉSINES.					PRODUCTION LIGNIERE TOTALE ANUELLE.					ÉCORCES, LIÈGES, RÉSINES.											
	Tailles		Futaie		En quiniaux.	Tailles		Futaie		En quiniaux.	Tailles		Futaie		En quiniaux.	Tailles		Futaie		En quiniaux.	Tailles		Futaie		En quiniaux.							
	hectares.	m. c.	m. c.	m. c.	quint.	hectares.	m. c.	m. c.	quint.	hectares.	m. c.	m. c.	quint.	hectares.	m. c.	m. c.	quint.	hectares.	m. c.	m. c.	quint.	hectares.	m. c.	m. c.	quint.							
1	638	6,585	22,543	413	16,948	9,619	37,925	388	98,618	777	113	539	2,545	301	3,172	14,791	8	113	539	2,545	301	3,172	14,791	8	113	539	2,545	301	3,172	14,791		
2	38	3,723	12,303	109	4,323	16,753	1,438	568	1,438	568	31	8,200	13,010	52,393	13,652	65,204	31	31	8,200	13,010	52,393	13,652	65,204	31	31	8,200	13,010	52,393	13,652	65,204		
3	4,126	5,121	12,303	133	28,375	8,432	7,300	36	8,906	476	168	268	21,005	55	831	70,403	104,000	104,000	104,000	22,138	1	1,205	1,910	118	118	1,100	3	70,403	70,403	70,403		
4	8,108	5,991	912	69,077	13,910	20,747	880	29,650	29	2,650	1	1,255	4,069	350	76,227	9,450	75,000	75,000	75,000	3,171	3	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171		
5	4,258	8,299	402	33,793	10,388	880	703	29,065	29	2,650	1	1,255	4,069	350	76,227	9,450	75,000	75,000	75,000	3,171	3	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	
6	770	8,231	31,335	1,404	10,388	3,113	313	29,065	29	2,650	1	1,255	4,069	350	76,227	9,450	75,000	75,000	75,000	3,171	3	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	
7	808	12,207	31,335	388	38,932	78,856	313	29,065	29	2,650	1	1,255	4,069	350	76,227	9,450	75,000	75,000	75,000	3,171	3	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171	3,171
8	388	300	24,331	694	2,047	914	8,553	6	1,592	180	40,100	4,836	182	840	30,300	21,138	10,270	10,270	10,270	3,716	6	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	
9	2,665	3,350	3,046	99	6,829	2,007	4,680	4,420	176	1,592	40,100	4,836	182	840	30,300	21,138	10,270	10,270	10,270	3,716	6	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	3,716	
10	609	6,128	1,202	9,048	8,911	1,479	12,321	12,321	12,321	3,113	15,000	27	2,523	130	101	692	5,866	131	109	56	6	6	6	6	6	6	6	6	6	6	6	
11	133	11,200		9,048	8,911	1,479	12,321	12,321	12,321	3,113	15,000	27	2,523	130	101	692	5,866	131	109	56	6	6	6	6	6	6	6	6	6	6	6	
12	99	1,158	6,681	2,431	4,128	2,010	6,428	1,440	5,091		69	1,583	1,832		4,832																	
13	204	204					728	1,440	5,091		69	1,583	1,832		4,832																	
14	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
15	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
16	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
17	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
18	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
19	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
20	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
21	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
22	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
23	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
24	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
25	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
26	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
27	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
28	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
29	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
30	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
31	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
32	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
33	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
34	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
35	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
36	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
37	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
38	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
39	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
40	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
41	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
42	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
43	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
44	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	
45	294	294					728	1,440	5,091		69	1,583	1,832		4,832																	

Fig. 8 Overview of a forest data table presented in decennial agricultural statistics of 1892 at a NUTS-3 level

XVI. Bois et forêts (surface). [Suite.]

1	BOIS NON SOUMIS AU RÉGIME FORESTIER APPARTENANT À DES SOCIÉTÉS COMMERCIALES									BOIS NON SOUMIS AU RÉGIME FORESTIER APPARTENANT À DES PARTICULIERS.												
	DEPARTEMENTS.				OF CIVILES.					RECEMENT.				FUTAIE					TOTAL.			
	TABLES		FUTAIE		TABLES			RECEMENT		TABLES		TABLES			RECEMENT		TOTAL.			TOTAL.		
simples.		de feuillus.		de feuillus.			plantés.		simples.		de feuillus.			plantés.		Total.			Total.			
hectares.		hectares.		hectares.			hectares.		hectares.		hectares.			hectares.		hectares.			hectares.			
3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18							
1	182	86	268	158		126	31,630	25,056	59,135	3,555	6,576	10,111	307	60,563	97,217							
2	1,889	1,613	3,502	476		3,978	46,301	22,393	68,597	965	570	702	792	68,869	114,732							
3	151	27	181	5		186	5,787	13,298	19,085	3,215	131	3,379	347	52,811	4,400							
4	29	20	10	25		40	47,415	7,245	51,660	3,258	8,880	12,118		66,578	8,680							
5	130	15	115	430		67	14,859	1,719	16,578	2,763	4,106	6,869	295	29,652	5,316							
6	86	86	101	1,229		575	10,750	2,152	12,878	5,857	13,728	19,215		32,093	2,512							
7	2,423	100	2,523	500		2,773	62,580	5,000	67,280	7,059	31,432	92,111	1,012	88,011	19,759							
8	763	763	1,526	173		936	29,903	5,000	34,903	1,151	3,397	4,718		56,083	22,589							
9	101	617	718	122		692	17,211	56,026	73,237	1,369	35,691	37,290	1,631	112,111	22,511							
10	1,385	813	2,198	1,240		2,629	25,232	2,508	27,710	3,212	3,923	7,135	265	35,100	5,251							
11	921	968	1,911	2,679		4,390	18,058	10,088	28,146	13,215	3,713	16,958	7,212	67,715	6,540							
12							33,457	35,515	70,972	7,212	26,976	49,130		11,740	13							
13							18,058	3,216	21,451	1,650	25,146	26,976		34,156	3,521							
14							22,817	8,608	31,425	1,289	1,328	2,617	114	51,300	16,011							
15							18,000	7,300	25,300	13,000	13,000	26,000	230	86,530	30,200							
16							366	32,309	32,675	81,100	2,000	5,290	33	78,786	30,095							
17							35	55,301	55,336	61,865	1,556	15,392	55	113,901	4,846							
18							1,830	25,763	27,593	102,007	3,277	8,025	58	105,550	27,000							
19							195	34,500	34,695	35,700	22,250	37,900	8,300	27,809	1,100							
20							4,578	5,463	10,041	5,924	11,811	17,768		121,818	6,005							
21							9,114	91,758	103,872	755	16,118	16,903	1,673	30,640	1,800							
22							13,901	8,573	22,473	4,099	4,097	8,166		38,778	6,293							
23							291	13,901	14,192	22,473	4,099	6,048	2,117	94,818	41,921							
24							15	55,746	8,837	30,583	3,598	2,550	1,754	36,093	9,910							
25							1,116	5,575	12,511	18,119	1,934	16,878	152	118,281	17,903							
26							878	108,181	500	108,681	6,320	5,680	2,365	92,953	9,001							
27							6,682	51,297	32,115	83,422	5,010	2,018	7,028	55,904	11,000							
28							70	39,082	8,500	37,612	6,250	776	7,026	25,130	1,575							
29							963	16,660	3,970	21,430	1,970	1,860	3,590	61,073	8,828							
30							5,271	51,372	2,896	57,168	2,010	3,883	1,02	49,555	9,856							
31							163	33,458	11,910	18,368	867	320	1,187	39,635	10,875							
32							10,750	25,885	36,635	2,200	800	3,000		380,358	19,928							
33							33,753	20,985	54,738	21,200	983,369	310,698	8,940	67,352	4,369							
34							110	61,153	1,936	63,071	3,399	477	5	32,778	4,282							
35							3,762	22,990	6,920	29,880	1,301	2,378	3,882	16	32,778							
36							3,109	10,397	62,112	2,183	1,166	3,319	38	73,506	6,269							
37							91	61,496	17,030	78,526	4,180	3,787	7,967	100	86,393							
38							5,211	51,301	17,870	72,071	5,141	14,668	19,809	355	92,235							
39							3,702	6,363	11,131	17,197	605	4,597	5,292	21,396	15,000							
40							10,590	16,966	1,229	18,295	30,796	453,112	58,569	229,916	22,600							
41							80,179	14,826	95,005	5,001	32,340	27,905	1,159	121,360	10,615							
42							32	18,397	57,628	24,355	500	33,880	37,886	62,121	16,788							
43							35	10,115	1,825	11,940	5,151	37,688	61,009	474	75,423							
44							15,803	7,891	23,694	1,563	1,047	3,210	35	59,539	8,000							
45							5	39,317	38,874	78,191	6,744	15,056	629	100,629	12,810							
46							88,836	8,187	97,023	8,619	12,268	11,268	331	109,622	21,202							
47							22,111	21,081	33,192	3,350	31,414	37,794	3,009	81,295	13,927							
48							9,793	809	10,602	4,974	16,900	21,874	2,830	35,326	7,000							

Fig. 9 Overview of a forest data table presented in decennial agricultural statistics of 1929 at a NUTS-3 level

Annex 2: The Forest regime

During the forest transition (dated in the nineteenth century, Mather 1992), a forest law was set in 1827 to protect and restore forest resources, yet property rights embodied in the Declaration of Human Rights of 1789 prevented State's intervention in private ownership, making the forest law limited to public forests in its application, a principle termed the Forest Regime (Némoz-Rajot 1998). All public forests are subjected to the Forest Regime, while private forests are not. Forest regime aims to plan and supervise forest management, to arrange the sale of forest products and ensure the monitoring of forest ownership (Némoz-Rajot 1998). The Forest Regime's definition is as follows: "The forest regime includes all the management rules defined by the Forestry Code and implemented by the National Forestry Commission. It is applicable to the State-owned forests, to the local authorities and to public institutions of public interest. Furthermore, it concerns a number of non-wooded areas such as the perimeters of restoration of mountain land or the zones of the coastal dunes (Aquitaine, Charente-Maritime and Vendée, Nord-Pas-de-Calais)." (INSEE 2016).

Annex 3: Stand structure definition in French national forest inventory (NFI)

Stand structure is observed in the field, and measured by the relative cover rates (crown canopy projection) of vertical high forest and coppice layers. Structure is described only in stands whose relative cover rate is > 40%. Forests with a relative cover rate < 40% are considered to be open forests with no clear structure (6.8% of total forest area). As soon as the relative cover rate of coppice is < 25%, the NFI plot is described as a high-forest. With a relative cover rate of coppice and high-forest is $\geq 25\%$, plot is described as coppice-withstandards. Otherwise, the plot is described as coppice (relative cover rate of high-forest $\leq 25\%$). However, the structure of some forests could not be defined (3.0% of total forest area).

Annex 4: Harmonization of statistics

Forest tree species harmonization between *Daubrée* statistics and national forest inventory (NFI) data

Table 6 summarizes the 15 main tree species and the 2 generic categories (various broadleaves and various conifers) described in *Daubrée* statistics. A harmonization was intended for comparison between *Daubrée* and NFI statistics.

A tree species described in *Daubrée* statistics could be found with two different names. This is the case of *Chêne vert* and *Chêne yeuse*, which correspond to holm oak

(*Quercus ilex* L.). This distinction proves to come from the difference of species terminology according to the region, as both are never found in a same NUTS-3 unit. Therefore, terminology was merged.

Some tree species groups did not show a single species, but a grouping of species depending on the composition (various broadleaves and various conifers), or two mixed species ("Fir and Spruce" and "Maritime Pine and Scots Pine" (*Pinus pinaster* Aiton. and *Pinus sylvestris* L., respectively) observed only in two NUTS-3 units, *Ain* and *Finistère*). In the first case, we did not change the original tree species group (becoming various broadleaves and various conifers). In the second case, we observed that when one of the mixed categories was present, the individual categories of the same species were systematically absent. We concluded that the distinction of species in mixed groups had not been made and that the corresponding areas did not refer to mixed-species areas. Thus, for a given category of two species, we conventionally attributed half of the area divided for two of the group to each of the species (for two NUTS-3 units).

The *Daubrée* statistics also refers to a tree species category termed oak in some NUTS-3 units. The absence of pedunculate and pubescent oak tree species (*Quercus robur* L. and *Quercus pubescens* Willd., respectively) in the statistics, and knowledge about their presence at the time of *Daubrée* (Boppe 1889) suggested that the oak category may comprise the area of these two species.

We identify that when the generic oak category was present in a NUTS-3 unit, the other oak tree species identified by *Daubrée* statistics (holm oak, cork oak, sessile oak (*Quercus petraea* Matt. Liebl)) were not present. And conversely, when one of the three oak tree species was present (holm oak, cork oak, sessile oak), the generic oak category was absent. We concluded that for each NUTS-3, oaks could either be described as an individual oak tree species, or only described as oak.

For our analyses, we merged terminology of the generic oak category and sessile oak tree species into a white oaks tree species group.

All the pines of *Daubrée* statistics and NFI data have been grouped in a generic pines tree species group.

Note, for three NUTS-3 units (Côte-d'Armor, Dordogne and Lot), the area of forests subjected to forest regime was null in *Daubrée* statistics.

Concerning NFI, all the species that were not inventoried by *Daubrée* statistics were placed in the two tree species groups of various broadleaves and various conifers. The *white oaks* tree species group was created by adding areas associated with the NFI data of sessile, pedunculated and pubescent oaks. Holm oak and cork oak were analyzed as individual species. Other oaks were placed in various broadleaves tree species group.

Table 6 Harmonized tree species categories and summary statistics for differences in forest areas between *Daubr e* statistics (1908) and national forest inventory data (2008¹) across NUTS-3 units. Forests area differences according department administrative units (NUTS-3) population

were calculated over the 1908–2008 period. Significance of the forest area differences were analyzed using a paired *t* test. Tree species not specified in oak, various broadleaves, and various conifers, yet deduced for the oak group (see Annex 4)

<i>Daubr�e</i> statistics' tree species Seminal terminology / Latin name	Generic translation used	Harmonization	FAD mean ($\times 10^3$ ha)	FAD standard-deviation ($\times 10^3$ ha)	Total change ($\times 10^3$ ha)	Annual rate of change (%)	Paired <i>t</i> test	
Ch�ene	<i>Quercus robur</i> L. <i>Quercus pubescens</i> Willd. <i>Quercus petraea</i> Matt. Liebl	Oak	White oaks	+ 18	21	+ 1550	+ 0.37	< 10 ⁻¹⁰
Ch�ene rouvre	<i>Quercus petraea</i> Matt. Liebl	Sessile oak						
Ch�ene vert	<i>Quercus ilex</i> L.	Holm oak	Holm oak	+ 2.7	39.3	+ 450	+ 0.97	< 0.05
Ch�ene yeuse	<i>Quercus suber</i> L.	Cork oak	Cork oak	- 0.2	6.5	- 60	- 0.46	NS
Ch�ene li�ge								
H�tre	<i>Fagus sylvatica</i> L.	Beech	Beech	+ 2.8	9.7	+ 190	+ 0.15	< 0.05
Ch�taignier	<i>Castanea sativa</i> Mill.	Chestnut	Chestnut	+ 7.4	8.2	+ 520	+ 1.20	< 10 ⁻⁹
Charme	<i>Carpinus betulus</i> L.	Hornbeam	Hornbeam	- 0.6	8.4	- 40	- 0.06	NS
Sapin	<i>Abies alba</i>	Silver fir	Silver fir	+ 3.7	7	+ 200	+ 0.48	< 0.001
Epic�ea	<i>Picea abies</i> Karst.	Norway spruce	Norway spruce	+ 6.1	8.7	+ 350	+ 0.96	< 10 ⁻⁵
Pin d'Alep	<i>Pinus halepensis</i> Mill.	Allepo pine	<i>Pines</i>	+ 8.4	23.5	+ 710	+ 0.34	< 0.01
Pin sylvestre	<i>Pinus sylvestris</i> L. <i>Pinus pinaster</i> Aiton.	Scots pine Maritime pine						
Pin maritime	<i>Pinus nigra</i> subsp. <i>laricio</i> Poir. Mair	Laricio pine Mountain pine						
Pin laricio	<i>Pinus uncinata</i>							
Pin � crochets	Ramond ex DC.							
M�l�ze	<i>Larix decidua</i> Mill.	Larch	Larch	+ 1.9	4.2	+ 40	+ 0.49	< 0.05
Feuillus divers		<i>Various broadleaves</i>	<i>Various broadleaves</i>	+ 11.7	15.1	+ 1000	+ 0.57	< 10 ⁻⁹
R�sineux divers		<i>Various conifers</i>	<i>Various conifers</i>	+ 3.9	10.6	+ 320	+ 0.58	< 0.01

FAD forest area difference. NFI national forest inventory. ¹: median year

Harmonization of forest management types and ownership categories

We aggregated ownership categories of the three ancient statistics into three/two groups depending on our analyses: (1) State-owned forest, other public forest and private forest; or (2) public forest and private forest (Table 7).

Forest management types were aggregated into four types: coppice, coppice-with-standards, high forest, and other forests' areas (Table 8). Other forests' areas category corresponds

to the specific marginal forest management types of ancient statistics. It includes unproductive forest areas of *Daubr e* statistics, decreed perimeters, and unproductive coastal dunes areas of 1892 statistics and recently afforested areas of 1929 statistics. Areas belonging to this forest management type are encompassed in the calculation of the afforestation rates in the three ancient statistics, and were therefore considered as well as other management type. Areas with no forest structure in NFI data, i.e. with a relatively low forest cover, were placed in such other forests' areas.

Table 7 Harmonization of forest ownership categories across ancient statistics and national forest inventory data

Statistics	Ownership category				
Decennial agricultural statistics (1892)	State-owned forests subjected to the forest regime	Municipal forests and other territorial public bodies' forests subjected to the forest regime	Private forests	Municipal forests and other territorial public bodies' forests not subjected to the forest regime	
German statistics (1907) for <i>Bas-Rhin</i> , <i>Haut-Rhin</i> and <i>Moselle</i> units	State-owned forests	Municipal forests (other public forest)	Private forests		
<i>Daubrée</i> statistics (1908)	State-owned forests subjected to the forest regime	Municipal forests and other territorial public bodies' forests subjected to the forest regime	Private forests	Municipal forests and other territorial public bodies' forests not subjected to the forest regime	
Decennial agricultural statistics (1929)		Forests subjected to the forest regime (including State-owned forests and other public forests)	Private forests	Municipal forests not subjected to the forest regime	Other territorial public bodies' forests not subjected to the forest regime
NFI (2008 ¹)	State-owned forests subjected to the forest regime	Municipal forests and other public forest subjected to the forest regime	Private forests	other territorial public bodies' forests not subjected to the forest regime	
Harmonization	State-owned forests subjected to the forest regime	Other public forests subjected to the forest regime	Private forests	not subjected to the forest regime	
	Alternate aggregation: Public forests				

NFI national forest inventory. ¹ median year

Table 8 Harmonization of forest management types across ancient statistics and national forest inventory data. While these data were merged, national forest inventory data primarily describes the physical forest structure of forest plots, and the three ancient statistics describe forest management types

Statistics	Forest management type					
Decennial agricultural statistics (1892)	Simple coppice	Coppice conversion	Coppice-with-standards	High broadleaved forest	High coniferous forest	Decreed afforestation perimeters and unproductive coastal dunes
<i>Daubrée</i> statistics (1908)	Simple coppice	Coppice conversion	Coppice-with-standards	High forest		Unproductive forest area
Decennial agricultural statistics (1929)	Simple coppice	Coppice-with-standards		High broadleaved forest	High coniferous forest	Recently afforested area
NFI (2008 ¹)	Coppice	Mixture of coppice and high forest		Even-aged and uneven-aged high forest		Undefined forest structure
Harmonization	Coppice	Coppice-with-standards		High forest		Other forests

¹ median year

Annex 5: Data entry and error correction

In the three ancient statistics, digitization errors of different types were identified by comparing row/column totals:

- miscalculation or typographical error on total statistics (input error when compiling total or printing error, 31 errors found);
- typographical errors on basic figures detected by double-check on horizontal and vertical totals (input error when compiling figures or printing error, 6 errors found);
- missing number (1 error found).

A miscalculation or typographical error on a total was detected when, for a given row and column, only one total was equal to that recomputed. When both row/column totals did not coincide with those recomputed, a typographical error on the basic figure at their intersection or missing number was inferred. Once the error was identified, it was possible to correct it.

All detected errors were corrected. However, primary typographical potential errors on basic figures, thus consistent with both row/column totals could not be detected and corrected.

These errors—as calculated from area and production data—actually represent a small share of the total number of statistics entered (0.005% and 0.003% of figures respectively in decennial agricultural statistics of 1892 and of 1929 and 0.002% in *Daubrée* statistics). The three statistics further showed many errors in the published afforestation rates due to the handling of decimals. These rates were also recomputed.

Annex 6: Department administrative units

Standardization of department administrative units' boundaries

Ancient department administrative units were not systematically identical to current NUTS-3 units and had to be harmonized. The broadest department administrative units (NUTS-3) scale compatible with all four statistics (three ancient ones and NFI) was selected.

Some department administrative units names in the three ancient statistics changed over time (for example: *Charente-inférieure* becoming *Charente-maritime*), and others were sub-divided (*Seine-et-Oise* turned into *Yvelines*, *Essonne* and *Val-d'Oise*; *Seine* turned into *Paris*, *Hauts-de-Seine*, *Seine-Saint-Denis* and *Val-de-Marne*). The latter units were considered as a single one in our analyses. The Corsica region was not separated in two department administrative units in the ancient statistics. We thus considered the island as one single unit for the whole study.

Potential effect of changes in department administrative units' area on forest differences investigated

Differences in total department administrative units' areas across the three ancient statistics were detected. For example, *Alpes-Maritimes* lost 17,900 ha between 1892 and 1908 while *Gironde* gained 24,000 ha over the same period. We investigated the potential average effect of these differences on between statistics forest areal changes using univariate regression analysis, and did not find any significant relationship across the 1908–1929 period ($p = 0.5$, R -squared = 0.005) and across 1908–2008 period ($p = 0.7$, R -squared = 0.002). However, we found a significant positive association across 1892–1908 period ($p < 0.01$, R -squared = 0.096) and in the 1892–1929 period ($p < 0.05$, R -squared = 0.051). This relationship yet turning non-significant without *Gironde* department administrative unit ($p = 0.7/p = 0.2$, R -squared = 0.001/ R -squared = 0.018 in the 1892–1908/1892–1929 periods, respectively). In 1817, a prefectural decree ordered the afforestation of 100,000 ha of coastal dunes in *Gironde* and *Landes* (Buffault 1942). The coastal dunes have an area of 124,000 ha (Favenec 1999), and in 1876, they were afforested by over 80,000 ha (Buffault 1942). *Gironde* had strongest area differences (+ 100,000 ha) and strongest forest area differences with *Landes* (+ 100,000 ha for both) over the 1892–1929 period. Consequently, the absence or presence of coastal dunes depending to their status (afforested or non-afforested) in department administrative units provided by the statistics may explain a part of this sharp increase. However, *Landes* showed small area differences over the 1892–1929 period (+ 4000 ha), suggesting an existence of another possible origin of area differences in *Gironde*. As a result, *Gironde* was excluded from the whole paired t tests in our analyses. However, it was kept in all other computations (totals, standard-deviations, maximum, and minimum).

References

- Audinot T, Wernsdörfer H, Bontemps JD (2020) Ancient forest statistics (Daubrée statistics 1908, decennial agricultural statistics of 1892 and 1929) describing French forest area by department administrative units, ownership and forest management types. [dataset]. Zenodo repository. V2. <https://doi.org/10.5281/zenodo.3739458>
- Bazire P, Gadant J (1991) La Forêt en France. La Documentation française, Paris
- Betolaud Y (1972) Un quart de siècle au service de la forêt française. *Revue Forestière Française* 24 n° spécial Le Fonds forestier national:501–504
- Boppe L (1889) *Traité de sylviculture*. Berger-Levrault et Cie, Paris and Nancy
- Bouchon J (1984) Importance des plantations de Douglas et Épicéa en France. *Revue forestière française* 36:254–258
- Brenac L (1984) Connaissances statistiques des forêts françaises avant l'Inventaire forestier national. *Revue Forestière Française* 36:7–90

- Buffault P (1942) Histoire des dunes maritimes de la Gascogne. Editions Delmas, Bordeaux
- Bugalho MN, Caldeira MC, Pereira JS, Aronson J, Pausas JG (2011) Mediterranean cork oak savannas require human use to sustain biodiversity and ecosystem services. *Front Ecol Environ* 9:278–286
- Cavaillès J, Normandin D (1993) Déprise agricole et boisement: état des lieux, enjeux et perspectives dans le cadre de la réforme de la PAC. *Revue forestière française* 45:465–482
- Chakir R, Madignier AC (2006) Analyse des changements d'occupation des sols en France entre 1992 et 2003. *Économie* 296:59–68
- Cinotti B (1996) Evolution des surfaces boisées en France: proposition de reconstitution depuis le début du XIXe siècle. *Revue forestière française* 48:547–562
- Corvol A (2013) Lucien Daubrée: 1912 et la statistique forestière. Proceedings of the conference: 1912-2012, de la statistique Daubrée à l'inventaire forestier de l'IGN: un siècle d'expansion des forêts françaises, Nancy. <http://www2.agroparistech.fr/podcast/Lucien-Daubree-1912-et-la-statistique-forestiere.html>. Accessed 19 Dec 2019
- Costa A, Pereira H, Madeira M (2010) Analysis of spatial patterns of oak decline in cork oak woodlands in Mediterranean conditions. *Ann For Sci* 67:204. <https://doi.org/10.1051/forest/2009097>
- Dabat L (1920) La Guerre et les forêts françaises - programme forestier. Imprimerie nationale, Paris
- Daubrée L (Ministère de l'agriculture, FRANCE) (1912) Statistique et atlas des forêts de France par Lucien Daubrée d'après les renseignements fournis par les agents des Eaux et Forêts. Eaux et forêts - Direction générale, Imprimerie nationale, Paris, 2 vol.
- David R (1993) L'assainissement des Landes de Gascogne et leur mise en valeur par le boisement. *Comptes Rendus de l'Académie des Sciences (La Vie des Sciences)* 10:235–259
- de Rochebouet H (1987) Le reboisement. *Revue forestière française* 39 no. spécial Le Fonds forestier national a 40 ans:34–42
- Degron R (1998) La conversion des forêts domaniales feuillues françaises (1860-1888): une grande vague brisée. *Revue forestière française* 50:71–84
- Denardou A, Hervé JC, Dupouey JL, Bir J, Audinot T, Bontemps JD (2017) L'expansion séculaire des forêts françaises est dominée par l'accroissement du stock sur pied et ne sature pas dans le temps. *Revue Forestière Française* 69:319–339
- Dodane C (2009) Les nouvelles forêts du Massif Central: enjeux sociétaux et territoriaux. Ces hommes qui plantaient des résineux pour éviter la friche. Doctoral dissertation. Ecole normale supérieure Lettres et Sciences humaines, Lyon
- Dupouey JL, Hervé J-C (2013) Comparaison entre les forêts de 1912 et les forêts actuelles. Proceedings of the conference: 1912-2012, de la statistique Daubrée à l'inventaire forestier de l'IGN: un siècle d'expansion des forêts françaises, Nancy. http://www2.agroparistech.fr/podcast/Comparaison-entre-les-forets-de-1912-et-les-forets-actuelles.html?debut_pagi=%40939. Accessed 30 Jan 2020
- European Parliament, Council of the European Union (2003) Regulation (EC) No 1059/2003 of the European Parliament and of the Council of 26 May 2003 on the establishment of a common classification of territorial units for statistics (NUTS). <https://eur-lex.europa.eu/eli/reg/2003/1059/oj>. Accessed 13 Mar 2020
- European Parliament, Council of the European Union (2013) regulation (EU) no 1305/2013 of the European Parliament and of the council of 17 December 2013 on support for rural development by the European agricultural Fund for Rural Development (EAFRD) and repealing council regulation (EC) no 1698/2005. <https://eur-lex.europa.eu/eli/reg/2013/1305/oj>. Accessed 13 Mar 2020
- FAO (2000) Global Forest Resources Assessment 2000. FAO forestry paper 140
- Favennec J (1999) Aménagement des forêts littorales: cas des forêts dunaires du littoral atlantique français. *Revue Forestière française* 51 n° spécial L'aménagement forestier hier, aujourd'hui, demain: 217–229
- Forest Europe (2015) State of Europe's forests 2015 report – forest Europe. <https://foresteurope.org/state-europes-forests-2015-report/>. Accessed 20 Apr 2016
- Foster D, Swanson F, Aber J, Burke I, Brokaw N, Tilman D, Knapp A (2003) The importance of land-use legacies to ecology and conservation. *BioScience* 53:77–88. [https://doi.org/10.1641/0006-3568\(2003\)053\[0077:TIOLUL\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0077:TIOLUL]2.0.CO;2)
- Fraye WE, Furnival GM (1999) Forest survey sampling designs: a history. *J For* 97:4–10
- Fromont P (1938) Statistique agricole de la France - Résultats généraux de l'enquête de 1929. *Revue D'économie Politique* 52:215–217
- Hitier H (1899) La statistique agricole de la France. *Annales de Géographie* 8:350–357
- IGN (2013) Un siècle d'expansion des forêts françaises - De la statistique Daubrée à l'inventaire forestier de l'IGN. L'IF 31
- INSEE (2016) Forest regime. <https://www.insee.fr/en/metadonnees/definition/c1176>. Accessed 30 Jan 2020
- Larrère R, Brun A, Kalaora B, Nougarede O, Poupardin D (1980) Reboisement des montagnes et systèmes agraires. *Revue Forestière Française* no. spécial Société et forêts:20–36
- Mather AS (1992) The Forest transition. *Area* 24:367–379
- Mather AS, Fairbairn J, Needle CL (1999) The course and drivers of the forest transition: the case of France. *J Rural Stud* 15:65–90. [https://doi.org/10.1016/S0743-0167\(98\)00023-0](https://doi.org/10.1016/S0743-0167(98)00023-0)
- Métaillé JP (1999) Los bosques de Francia en los siglos XIX y XX: Bosques nuevos y nuevas investigaciones. *Historia Agraria* 18:33–55
- Meyfroidt P, Lambin EF (2011) Global forest transition: prospects for an end to deforestation. *Annu Rev Environ Resour* 36:343–371. <https://doi.org/10.1146/annurev-environ-090710-143732>
- Ministère de l'Agriculture Direction de l'Agriculture (1892) Statistique agricole décennale de la France. 1ère Edition, Imprimerie nationale, Paris
- Ministère de l'Agriculture Direction de l'Agriculture (1929) Statistique agricole décennale de la France. 1ère Edition, Imprimerie nationale, Paris
- Ministerium für Elsass-Lothringen (1909) Beiträge zur Forststatistik von Elsass-Lothringen. Strasbourg, 26
- Moriniaux V (1999) Les Français face à l'enrésinement, XVIe-XXe siècles. *Annales de Géographie*, no. 609-610:660–663
- Morneau F, Duprez C, Hervé JC (2008) Les forêts mélangées en France métropolitaine. Caractérisation à partir des résultats de l'Inventaire forestier national *Revue forestière française* 60:107–120
- Musset R (1938) La statistique agricole de la France de 1929. *Annales de géographie* 47:67–70
- Némoz-Rajot H (1998) Le Régime forestier: une mosaïque moderne et évolutive. *Revue forestière française* 50:10–19
- Oberlé Y (2000) Les statistiques forestières - Catalogue des sources de données anciennes. Office National des Forêts, Collection dossiers forestiers 1ère édition
- Office National des Forêts (2010) Sylviculture des chênaies dans les forêts publiques françaises. Rendez-vous techniques 5 hors-série
- Pignard (2000) Évolution récente des forêts françaises : surface, volume sur pied, productivité. *Revue forestière française* 52 n° spécial Conséquence des changements climatiques pour la forêt et la sylviculture:27–36
- Robert N, Vidal C, Colin A, Hervé JC, Hamza N, Cluzeau C (2010) France. In: Tomppo E, Gschwanter T, Lawrence M, McRoberts RE, Gabler K, Schadauer K, Vidal C, Lanz A, Ståhl G, Cienciala E, Chirici G, Winter S, Bastrup-Birk A, Tomter S, Kändler G, McCormick M. National Forest Inventories: pathways for common reporting. Springer, Heidelberg, pp 207–221

- Saillet B (1991) De l'utilité des boisements pour la restauration et la conservation des terrains en montagne. *Revue forestière française* 43:119–130
- Sonnier J (1991) Analyse du rôle de protection des forêts domaniales de montagne. *Revue forestière française* 43:131–145
- Tomppo E, Gschwantner T, Lawrence M, McRoberts RE, Gabler K, Schadauer K, Vidal C, Lanz A, Ståhl G, Cienciala E, Chirici G, Winter S, Bastrup-Birk A, Tomter S, Kändler G, McCormick M (2010) National Forest Inventories: pathways for common reporting. Springer, Heidelberg
- Vidal C, Alberdi I, Redmond J, Vestman M, Lanz A, Schadauer K (2016) The role of European National Forest Inventories for international forestry reporting. *Ann For Sci* 73:793–806. <https://doi.org/10.1007/s13595-016-0545-6>
- Wernsdörfer H, Colin A, Bontemps JD, Chevalier H, Pignard G, Cauria S, Leban JM, Hervé JC, Fournier M (2012) Large-scale dynamics of a heterogeneous forest resource are driven jointly by geographically varying growth conditions, tree species composition and stand structure. *Ann For Sci* 69:829–844. <https://doi.org/10.1007/s13595-012-0196-1>
- Willis KJ, Birks HJB (2006) What is natural? The need for a long-term perspective in biodiversity conservation. *Science* 314:1261–1265. <https://doi.org/10.1126/science.1122667>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.