



HAL
open science

A multiple time renewal equation for neural assemblies with elapsed time model

Nicolás Torres, Benoît Perthame, Delphine Salort

► **To cite this version:**

Nicolás Torres, Benoît Perthame, Delphine Salort. A multiple time renewal equation for neural assemblies with elapsed time model. 2021. hal-03324280v1

HAL Id: hal-03324280

<https://hal.science/hal-03324280v1>

Preprint submitted on 23 Aug 2021 (v1), last revised 14 Apr 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A multiple time renewal equation for neural assemblies with elapsed time model

Nicolás Torres^{*a}, Benoît Perthame^{†a}, and Delphine Salort^{‡b}

^aSorbonne Université, CNRS, Laboratoire Jacques-Louis Lions, UMR 7598. 4 Place Jussieu, 75005 Paris, France.

^bSorbonne Université, CNRS, Laboratoire de Biologie Computationnelle et Quantitative, UMR 7238. 4 Place Jussieu, 75005 Paris, France.

August 2021

Abstract

We introduce and study an extension of the classical elapsed time equation in the context of neuron populations that are described by the elapsed time since the last discharge, i.e., the refractory period. In this extension we incorporate the elapsed time since the penultimate discharge and we obtain a more complex system of integro-differential equations. For this new system we prove convergence to stationary state by means of Doeblin's theory in the case of weak non-linearities in an appropriate functional setting, inspired by the case of the classical elapsed time equation. Moreover, we present some numerical simulations to observe how different firing rates can give different types of behaviors and to contrast them with theoretical results of both classical and extended models.

2010 *Mathematics Subject Classification.* 35B40, 35F20, 35R09, 92B20

Keywords and phrases. Structured equations; Renewal Equation; Mathematical neuroscience; Neural networks; Doeblin theory.

1 Introduction

In the study and modelling of neural processes, population density models have proved to be a useful approach to understand brain phenomena at different scales. Among these models we mention for example the well-known integrate-and-fire model which describes the dynamics of the membrane

*Corresponding author. Email : nicolas.torres_escorza@sorbonne-universite.fr

†Email : benoit.perthame@sorbonne-universite.fr

‡Email : delphine.salort@sorbonne-universite.fr

potential and has been studied by several authors like Carrillo et al. [3, 6], Perthame et al. [1, 26, 25] and Zhou et al. [16] in different variants and approaches. For a connection with stochastic processes, see . Another class of population-based model is the elapsed time model, which has attracted the attention of many researchers. In this model we consider a neural network where neurons are described by their refractory period as the key variable, i.e. the elapsed time since the last discharge. After receiving some stimulation, neurons spike and interact with other neurons leading them to spike as well.

Like the integrate-and-fire model, the elapsed time model is closely related to the limit of stochastic processes at microscopic scale and the connection with Poisson processes was established in the work of Chevalier et al. [7]. Other important works on spiking neurons include Brunel [2], Gerstner et al. [14], Ly et al. in [17] and Pham et al. [27]. A recent survey is available by Schwalger et al. [28]. Moreover, the elapsed time model has been studied from a mathematical and analytical point of view by several authors using different techniques such as Cañizo et al. in [5], Kang et al. [15], Mischler et al. in [20, 19] and the pioneer works of Pakdaman et al. in [21, 22, 23]. The relation between integrate-and-fire and the elapsed time model was studied in Dumont et al. [9, 10].

Different extensions of the elapsed time model have been considered by incorporating new variables such as spatial dependence and a connectivity kernel in Salort et al. [29] or a leaky memory variable in Fonte et al. [11]. The aim of the present work is to extend the classical elapsed time model by taking into account the elapsed time since the penultimate discharge in addition to the last one. We extend the ideas of Cañizo et al. in [5] on the application of Doeblin's theory for the exponential convergence to the steady state for the classical elapsed time equation. For a reference on Doeblin's theory, see for example [13] and for further extensions see [4]. This theory on Markov semi-groups is an alternative to the classical entropy methods used to prove convergence to the steady state, as it was developed in the works [18, 24]. Moreover, convergence for the integrate-and-fire model has been proved in Perthame et al. [25] and Dumont et al. [8], by means of Doeblin's theory.

The extended model is described as follows. Let $n = n(t, s, a)$ the probability density of finding a neuron at time t , such that the elapsed times since its last and penultimate discharge are respectively s and a . For simplicity, we simply call s as the first elapsed time and a as the second one. Moreover, we assume that for all $t \geq 0$ the domain of definition of n in the elapsed time variables is contained in the domain

$$\mathcal{D} := \{(s, a) \in \mathbb{R}^2 : 0 \leq s \leq a\}.$$

Neural dynamics are modelled through the following nonlinear renewal system

$$\begin{cases} \partial_t n + \partial_s n + \partial_a n + p(s, a, X(t))n = 0 & t > 0, a > s > 0, \\ n(t, s = 0, a) = N(t, a) := \int_0^\infty p(a, u, X(t))n(t, a, u) du & t > 0, a > 0, \\ X(t) = \int_0^\infty N(t, a) da & t > 0, \\ n(t = 0, s, a) = n_0(s, a) & a > s > 0. \end{cases} \quad (1)$$

As in the classical elapsed time model the function $p: \mathcal{D} \times \mathbb{R} \rightarrow \mathbb{R}$ is the firing rate of neurons, which depends on the total activity $X(t)$. Furthermore, for the firing rate function p , we assume that there exist $\sigma, p_0, p_\infty > 0$ such that

$$p_0 \mathbb{1}_{\{a > s \geq \sigma\}} \leq p(s, a, X) \leq p_\infty. \quad (2)$$

Thus, we get

$$0 \leq X(t) \leq p_\infty, \quad \forall t \geq 0. \quad (3)$$

We assume for simplicity that $p \in W^{1,\infty}(\mathcal{D} \times \mathbb{R})$, although most of the theoretical results are also valid for firing rates with simple jump discontinuities and the behaviour of solutions does not depend on this regularity assumption as we show in the numerical simulations. Furthermore, we say that the network is inhibitory if p is decreasing with respect to the total activity X and excitatory if p is increasing. If in addition $\|\partial_X p\|_\infty$ is small, we say that System (1) is under a weak interconnection regime.

The function $N(t, a)$ represents the flux discharging neurons conditioned to elapsed time since penultimate discharge, so that the total activity $X(t)$ corresponds to integrate with respect to all penultimate times. The boundary condition of n at $s = 0$ states that the second elapsed time resets to the first elapsed time.

Finally, we assume that the initial data $n_0 \in L^1(\mathcal{D})$ is a probability density so that System (1) formally verifies

$$\iint n(t, s, a) da ds = \iint n_0(s, a) da ds = 1, \quad n(t, s, a) \geq 0 \quad \forall t \geq 0. \quad (4)$$

The multiple time renewal equation has been investigated in Fournier et al. [12] in the linear case, where a non-expanding distance was introduced via a coupling argument.

In the particular case when p does not depend on a , the probability density $m(t, s) := \int_s^\infty n(t, s, a) da$ satisfies the equation

$$\begin{cases} \partial_t m + \partial_s m + p(s, X(t))m = 0 & t > 0, s > 0, \\ m(t, s = 0) = X(t) = \int_0^\infty p(u, X(t))m(t, u) du & t > 0, \\ m(t = 0, s) = \int_s^\infty n_0(s, a) da & s > 0. \end{cases} \quad (5)$$

In other words, the probability with respect to the last elapsed time is a solution of the classical elapsed time equation. If in addition we consider a the firing rate of the form

$$p = \varphi(X(t))\mathbb{1}_{\{s > \sigma\}},$$

with $\varphi \in W^{1,\infty}(\mathbb{R})$ strictly positive and $\sigma > 0$ a constant, we know from Caceres et al. [29] that the total activity $X(t)$ satisfies the integral equation

$$\int_{t-\sigma}^t X(s) ds + \frac{X(t)}{\varphi(X(t))} = 1, \quad \forall t \geq \sigma. \quad (6)$$

Moreover, we know that the solutions of this integral equation may have different behaviors like periodic solution and jump discontinuities. This gives us an idea of possible asymptotic behaviors that solutions of System (1) may exhibit.

The article is organized as follows. In Section 2 we prove that System (1) is well-posed in a suitable space for weak non-linearities. Starting with the asymptotic analysis for the linear case, we prove in Section 3 the existence of a stationary state and exponential convergence via Doeblin's theory. For the non-linear problem in the case of weak interconnections, we show in Section 4 the uniqueness of the steady state and in Section 5 we prove the exponential convergence via a perturbation argument. Finally in Section 6 we present some examples of numerical simulations for different initial data and firing rates.

2 Well-posedness for weak non-linearities

We prove that System (1) is well-posed under the weak interconnection regime. In order to do so, we start by studying an auxiliary linear problem where total activity is fixed and then we proceed to prove well-posedness of system (1) via a fixed point argument by contraction.

2.1 The linear problem

Given $X \in \mathcal{C}_b[0, \infty)$, we consider the following linear problem

$$\begin{cases} \partial_t n + \partial_s n + \partial_a n + p(s, a, X(t))n = 0 & t > 0, a > s > 0, \\ n(t, s = 0, a) = N(t, a) := \int_0^\infty p(a, u, X(t))n(t, a, u) du & t > 0, a > 0, \\ n(t = 0, s, x) = n_0(s, x) \geq 0 & a > s > 0. \end{cases} \quad (7)$$

We look for weak solutions satisfying $n \in \mathcal{C}_b([0, \infty), L^1(\mathcal{D}))$, so that $N \in \mathcal{C}_b([0, \infty), L^1(0, \infty))$ and $X \in \mathcal{C}_b[0, \infty)$.

Lemma 1. *Assume that $n_0 \in L^1(\mathcal{D})$ is a probability density and $p \in W^{1, \infty}((0, \infty) \times \mathbb{R})$ satisfies (2). Then for a given $X \in \mathcal{C}_b[0, \infty)$, Equation (7) has a unique weak solution $n \in \mathcal{C}_b([0, \infty), L^1(\mathcal{D}))$ with $N \in \mathcal{C}_b([0, \infty), L^1(0, \infty))$ and $X \in \mathcal{C}_b[0, \infty)$. Moreover n is non-negative and verifies the property (4).*

In particular this lemma proves the property (4) for the non-linear System (1).

Proof. From the method of characteristics, we start by noticing that a solution of the linear System (7) satisfies the following fixed point equation

$$\begin{aligned} n(t, s, x) = \Psi[n](t, s, x) := & n_0(s - t, a - t) e^{-\int_0^t p(t' + s - t, t' + a - t, X(t')) dt'} \mathbb{1}_{\{a > s > t\}} \\ & + N(t - s, a - s) e^{-\int_0^s p(s' + a - s, X(s' + t - s)) ds'} \mathbb{1}_{\{t, a > s\}}, \end{aligned} \quad (8)$$

with $N(t, a) = \int_0^\infty p(a, u, X(t))n(t, a, u) du$ depending on n .

Let $T > 0$ and $\mathcal{X}_T := \{n \in \mathcal{C}_b([0, T], L^1(\mathcal{D})) : n(0) = n_0\}$, it readily follows that Ψ maps $\mathcal{X}_T \rightarrow \mathcal{X}_T$. We prove by the Picard contraction theorem that Ψ has a unique fixed point in \mathcal{X}_T for $T > 0$ small enough, i.e., there exists a unique weak solution of (7) defined on $[0, T]$. Consider $n_1, n_2 \in \mathcal{X}_T$, we compute

$$\begin{aligned} \iint |\Psi[n_1] - \Psi[n_2]|(t, s, a) ds da & \leq \int_0^t \int_s^\infty |N_1 - N_2|(t - s, a - s) da ds \\ & \leq T \sup_{t \in [0, T]} \int_0^\infty |N_1 - N_2|(t, a) da \\ & \leq T p_\infty \sup_{t \in [0, T]} \|n_1(t, s, a) - n_2(t, s, a)\|_{L^1(\mathcal{D})}, \end{aligned} \quad (9)$$

thus for $T < \frac{1}{p_\infty}$, we have proved that Ψ is a contraction and there exists a unique $n \in \mathcal{X}_T$ such that $\Psi[n] = n$. Since the choice of T is independent of n_0 , we can reiterate this argument to get a unique

solution of (7), which is defined for all $t \geq 0$. Moreover this argument is also valid in $(\mathcal{M}(\mathcal{D}), \|\cdot\|_{TV})$, the space of regular measures on \mathcal{D} with the norm of the total variation.

Next we prove the mass conservation property. For all $t \geq 0$, consider $\mathcal{S}_t: \mathcal{M}(\mathcal{D}) \rightarrow \mathcal{M}(\mathcal{D})$ the semi-group given by

$$\mathcal{S}_t[f](s, a) = f(s - t, a - t) \mathbb{1}_{\{a > s > t\}},$$

whose infinitesimal generator is the operator $\mathcal{L}f = -\partial_s f - \partial_a f$. From Duhamel's formula, the solution of the fixed point problem (8) also verifies the following equality

$$n(t, s, a) = \mathcal{S}_t[n_0](s, a) + \int_0^t \mathcal{S}_{t-\tau}[\delta_{\{s=0\}}(s, a)N(\tau, a)] d\tau - \int_0^t \mathcal{S}_{t-\tau}[p(s, a, X(\tau))n(\tau, s, a)] d\tau, \quad (10)$$

where $\delta_{\{s=0\}}(s, a)$ is the measure along the line $\{(0, a) : a \geq 0\}$. This formula is translated as

$$\begin{aligned} n(t, s, x) &= n_0(s - t, a - t) \mathbb{1}_{\{a > s > t\}} + N(t - s, a - s) \mathbb{1}_{\{t, a > s\}} \\ &\quad - \int_0^t p(s - t + \tau, a - t + \tau, X(\tau))n(\tau, s - t + \tau, a - t + \tau) \mathbb{1}_{\{a > s > t - \tau\}} d\tau, \end{aligned} \quad (11)$$

and we get the mass conservation property by integrating with respect to (s, a) on the domain \mathcal{D} .

Finally, since n_0 is non-negative then Ψ preserves positivity, so by uniqueness of fixed point the corresponding solution n must be non-negative. \square

2.2 The non-linear problem

We are now ready to prove that System (1) is well-posed in the case of weak interconnection.

Theorem 1 (Well-posedness for weak interconnections). *Assume that $n_0 \in L^1(\mathcal{D})$ is a probability density and that $p \in W^{1,\infty}(\mathcal{D} \times \mathbb{R})$ satisfies (2). Then for*

$$\|\partial_X p\|_\infty < 1,$$

System (1) has a unique solution with $n \in \mathcal{C}_b([0, \infty), L^1(\mathcal{D}))$, $N \in \mathcal{C}_b([0, \infty), L^1(0, \infty))$ and $X \in \mathcal{C}_b[0, \infty)$. Moreover n verifies Condition (4) for all $t > 0$.

Proof. Consider $T > 0$. We fix a function $X \in \mathcal{C}_b[0, \infty)$ and define the functions $n \in \mathcal{C}_b([0, \infty), L^1(\mathcal{D}))$ and $N \in \mathcal{C}_b([0, \infty), L^1(0, \infty))$ which are solutions of System (7) by Lemma 1. Furthermore, the solution of this linear equation satisfies (4).

So we have a solution of System (1) defined on $[0, T]$ if X satisfies for all $0 \leq t \leq T$ and $x \in \Omega$, the following fixed point condition

$$X(t) = \mathcal{T}[X](t) := \int_0^\infty N[X](t, a) da. \quad (12)$$

We prove that \mathcal{T} defines for all $T > 0$ an operator that maps $\mathcal{X}_T \rightarrow \mathcal{X}_T$ with $\mathcal{X}_T := \mathcal{C}_b([0, T])$. First, we observe the following estimate

$$\left| \int N(t, a) da \right| \leq p_\infty, \quad \forall t \in [0, T], \quad (13)$$

and it is immediate that $\mathcal{T}[X] \in \mathcal{X}_T$.

We now prove that for T small enough, \mathcal{T} is a contraction. Let $X_1, X_2 \in \mathcal{X}_T$ with their respective solutions $(n_1, N_1), (n_2, N_2)$ of System (7). For the difference between N_1 and N_2 we have

$$\begin{aligned} \int |N_1 - N_2|(t, a) da &\leq \iint |p(a, u, X_1) n_1(t, a, u) - p(a, u, X_2) n_2(t, a, u)| du da \\ &\leq \iint |p(a, u, X_1) - p(a, u, X_2)| n_1 du da + \iint p(a, u, X_2) |n_1 - n_2|(t, a, u) du da \\ &\leq \|\partial_X p\|_\infty \|X_1 - X_2\|_\infty + p_\infty \|n_1 - n_2\|_{L^1(\mathcal{D})}. \end{aligned} \quad (14)$$

Now we have to estimate the difference between n_1 and n_2 . From (11) and estimate (14), we get

$$\|n_1 - n_2\|_{L^1(\mathcal{D})} \leq 2T \|\partial_X p\|_\infty \|X_1 - X_2\|_\infty + 2Tp_\infty \|n_1 - n_2\|_{L^1(\mathcal{D})}.$$

Then, for $T < \frac{1}{2p_\infty}$ we obtain

$$\|n_1 - n_2\|_{L^1(\mathcal{D})} \leq \frac{2T \|\partial_X p\|_\infty}{1 - 2Tp_\infty} \|X_1 - X_2\|_\infty. \quad (15)$$

Finally by using again estimate (14), the operator \mathcal{T} satisfies

$$\|\mathcal{T}[X_1] - \mathcal{T}[X_2]\|_\infty \leq \|\partial_X p\|_\infty \left(1 + \frac{2Tp_\infty}{1 - 2Tp_\infty}\right) \|X_1 - X_2\|_\infty \quad (16)$$

Hence for $\|\partial_X p\|_\infty < 1$ and T small enough, \mathcal{T} is a contraction.

From Picard's fixed point we get a unique $X \in \mathcal{X}_T$ such that $\mathcal{T}[X] = X$, and this implies the existence of a unique solution of (1) defined on $[0, T]$. Since estimate (13) is uniform in T , we can iterate this argument to get a unique solution of (1) defined for all $t > 0$.

Furthermore, we conclude from this construction that the non-linear System (1) satisfies (4) like the linear System (7). \square

3 Asymptotic behavior for the linear case

In order to study the behavior of System (1), we start by studying the case when $X \geq 0$ is a fixed constant. Thus we consider the linear problem given by

$$\begin{cases} \partial_t n + \partial_s n + \partial_a n + p(s, a, X)n = 0 & t > 0, a > s > 0, \\ n(t, s = 0, a) = N(t, a) := \int_0^\infty p(a, u, X)n(t, a, u) du & t > 0, a > 0, \\ n(t = 0, s, a) = n_0(s, a) & a > s > 0. \end{cases} \quad (17)$$

To determine the behavior of System (17), we consider $n_X(s, a)$ as the solution of the steady-state problem given by

$$\begin{cases} \partial_s n + \partial_a n + p(s, a, X)n = 0 & a > s > 0, \\ n(s = 0, a) = N(a) := \int_0^\infty p(a, u, X)n(a, u) du & t > 0, a > 0, \end{cases} \quad (18)$$

and we need to remind the main concepts on Markov semi-groups and Doeblin's theorem.

Definition 1. Let $(\mathcal{X}, \mathcal{A})$ be a measure space and $P_t: \mathcal{M}(\mathcal{X}) \rightarrow \mathcal{M}(\mathcal{X})$ be a linear semi-group acting on the space of measures $(\mathcal{M}(\mathcal{X}), \|\cdot\|_{TV})$. We say that P_t is a Markov semi-group if $P_t\mu \geq 0$ for all $\mu \geq 0$ and $\int_X P_t\mu = \int_X \mu$ for all $\mu \in \mathcal{M}(\mathcal{X})$. In other words, (P_t) preserves the subset of probability measures $\mathcal{P}(\mathcal{X})$.

Definition 2. Let $P_t: \mathcal{M}(\mathcal{X}) \rightarrow \mathcal{M}(\mathcal{X})$ be a Markov semi-group. We say that (P_t) satisfies Doeblin's condition if there exists $t_0 > 0$, $\alpha \in (0, 1)$ and $\nu \in \mathcal{P}(\mathcal{X})$ such that

$$P_{t_0}\mu \geq \alpha\nu \quad \forall \mu \in \mathcal{P}(\mathcal{X}).$$

Theorem 2 (Doeblin's Theorem). Let $P_t: \mathcal{M}(\mathcal{X}) \rightarrow \mathcal{M}(\mathcal{X})$ be a Markov semi-group that satisfies Doeblin's condition. Then the semigroup has a unique equilibrium $\mu^* \in \mathcal{P}(\mathcal{X})$. Moreover, for all $\mu \in \mathcal{M}(\mathcal{X})$ we have

$$\|P_t\mu - \langle \mu \rangle \mu^*\|_{\mathcal{M}(\mathcal{X})} \leq \frac{1}{1-\alpha} e^{-\lambda t} \|\mu - \langle \mu \rangle \mu^*\|_{\mathcal{M}(\mathcal{X})} \quad \forall t \geq 0,$$

with $\langle \mu \rangle = \int_X \mu$ and $\lambda = -\frac{\ln(1-\alpha)}{t_0} > 0$.

For a proof of Doeblin's Theorem, see for example [13].

From Lemma 7, the solution of the linear problem (17) determines a Markov semi-group acting on $L^1(\mathcal{D})$. By means of Doeblin's theory, the solutions of linear Equation (17) converge exponentially to a unique steady state, as we assert in the following theorem.

Theorem 3. Let $n_0 \in L^1(\mathcal{D})$ be a probability density and assume that p smooth satisfies Assumption (2). Then for a fixed $X > 0$, there exists a unique stationary solution $n_X(s, a) \in L^1(\mathcal{D})$ of the linear Equation (17) satisfying $\iint n_X(s, a) da ds = 1$. Moreover, the corresponding solution of Equation (17) satisfies

$$\|n(t) - n_X\|_{L^1_{s,a}} \leq \frac{1}{1-\alpha} e^{-\lambda t} \|n_0 - n_X\|_{L^1_{s,a}} \quad \forall t \geq 0,$$

with $\alpha = \frac{1}{2} p_0^2 \sigma^2 e^{-3p_\infty \sigma}$ and $\lambda = -\frac{\log(1-\alpha)}{3\sigma} > 0$.

In order to obtain the result, we show that after some time the solution of the linear problem is uniformly bounded from below for all probability densities. Thus from Doeblin's theorem we get the exponential convergence to equilibrium.

Lemma 2. Assume (4) and (2). Let $n(t, s, a)$ be a solution of (17), then there exists $t_0 > 0$, $\alpha \in (0, 1)$ and a probability density $\nu \in L^1$ such that

$$n(t_0, s, a) \geq \alpha\nu(s, a).$$

Proof. The main idea of the proof is to control the mass transported along the lines of direction (1, 1). Firstly, we observe the transport of the initial data n_0 . From Assumption (2) and the characteristics Formula (8) the following inequality holds

$$\int_t^\infty \int_s^\infty n(t, s, a) da ds \geq e^{-p_\infty t}, \quad \forall t \geq \sigma. \quad (19)$$

Secondly, we see the mass that returns at $s = 0$. From (19) we get for all $t \geq \sigma$

$$\int_t^\infty n(t, s = 0, a) da = \int_t^\infty N(t, a) da \geq p_0 \int_t^\infty \int_a^\infty n(t, a, u) du da \geq p_0 e^{-p_\infty t}. \quad (20)$$




Figure 1: First reduction of dimension. For a $t \geq \sigma$, all points in \mathcal{D} are transported to the red region, which has a total mass of at least $e^{-p_\infty t}$. Then a mass of at least $p_0 e^{-p_\infty t}$ returns to the green line.

This means that we reduced by one dimension the problem of finding the uniform lower bound. For $t \geq \sigma$ the mass of the region $\{(s, a) : a > s > t\}$ concentrates in the line $\{(0, a) : a \geq t\}$, as we see in Figure 1.

Thirdly, in order to control the point values of $n(t, s, a)$, we regard the values of $N(t, a)$. Observe that from Formula (8) we have

$$\begin{aligned} n(t, s, a) &\geq N(t - s, a - s) e^{-\int_0^s p(s', s' + a - s, X) ds'} \mathbb{1}_{\{t, a > s\}} \\ &\geq N(t - s, a - s) e^{-p_\infty s} \mathbb{1}_{\{t, a > s\}}, \end{aligned} \quad (21)$$

thus for $a \geq \sigma$ and $t - a > \sigma$, we obtain by using again Assumption (2) that

$$\begin{aligned} N(t, a) &\geq p_0 \int_a^\infty n(t, a, u) du \\ &\geq p_0 e^{-p_\infty a} \int_a^\infty N(t - a, u - a) du = p_0 e^{-p_\infty a} \int_0^\infty N(t - a, u) du \\ &\geq p_0 e^{-p_\infty a} \int_{t-a}^\infty N(t - a, u) du \geq p_0^2 e^{-p_\infty t}. \end{aligned} \quad (22)$$

This means we reduced the problem of finding the uniform lower bound by one dimension again, as we see in Figure 2.

Finally, once we have estimated $N(t, a)$ from below, we come back to estimate (21) to conclude that for $a - s \geq \sigma$ and $t - a > \sigma$ we have

$$\begin{aligned} n(t, a, s) &\geq N(t - s, a - s) e^{-p_\infty s} \mathbb{1}_{\{t, a > s\}} \\ &\geq p_0^2 e^{-p_\infty t} \mathbb{1}_{\{t - a, a - s > \sigma\}}, \end{aligned} \quad (23)$$

so that we can choose $t = 3\sigma$ and conclude that

$$n(3\sigma, a, s) \geq p_0^2 e^{-3p_\infty \sigma} \mathbb{1}_{\{2\sigma > a > s + \sigma\}}.$$




Figure 2: Second reduction of dimension. For $t \in [\sigma, 2\sigma]$ the green lines are transported to the region where $s \geq \sigma$ and their mass is of at least $p_0 e^{-2p_\infty \sigma}$. Then the mass of each green line is concentrated in the orange points, whose values are at least $p_0^2 e^{-2p_\infty \sigma}$.

Therefore we get the desired result with $t_0 = 3\sigma$, $\alpha = \frac{1}{2} \sigma^2 p_0^2 e^{-3p_\infty \sigma} \in (0, 1)$ and ν given by

$$\nu(s, a) = \frac{2}{\sigma^2} \mathbb{1}_{\{2\sigma > a > s + \sigma\}},$$

whose support is contained in orange region of Figure 3.




Figure 3: Finally for $t \in [2\sigma, 3\sigma]$ the orange dots are transported to region where $s \geq \sigma$, which allows to construct a minorization function for Doeblin's Theorem.

□

From Lemma 2 the hypothesis of Doeblin's theorem are verified and Theorem 3 readily follows.

Concerning the conditioned activity N in System (18), we conclude from Theorem 3 that for X fixed, there is a unique stationary $N_X \in L^1(0, \infty)$ determined by the method of characteristics through the

formula

$$n_X(s, a) = N_X(a - s) \exp\left(-\int_0^s p(s', a - s + s', X) ds'\right), \quad a > s. \quad (24)$$

Replacing this expression in the boundary condition at $s = 0$, we obtain the following integral equation for $N_X(a)$

$$N_X(a) = \mathcal{T}_X[N_X](a), \quad (25)$$

with $\mathcal{T}_X: L^1(0, \infty) \rightarrow L^1(0, \infty)$ given by

$$\begin{aligned} \mathcal{T}_X[N](a) &:= \int_0^\infty p(a, u + a, X) \exp\left(-\int_0^a p(s', u + s', X) ds'\right) N(u) du \\ &= -\frac{\partial}{\partial a} \int_0^\infty \exp\left(-\int_0^a p(s', u + s', X) ds'\right) N(u) du. \end{aligned}$$

Moreover, by integrating Equation (24) we get

$$\int_0^\infty \int_0^\infty N_X(a) \exp\left(-\int_0^s p(s', a + s', X) ds'\right) da ds = 1. \quad (26)$$

Therefore we conclude that finding a function $N \in L^1(0, \infty)$ satisfying Equation (25) and Condition (26) is equivalent to finding a steady-state $n_X(s, a)$ in Equation (18). The integral Equation (25) will play an important role in the analysis of the non-linear System (1), thus we prove the following two lemmas on the operator \mathcal{T}_X that will be useful in the sequel.

Lemma 3. *Assume that p is Lipschitz. For each $X > 0$ the operator \mathcal{T}_X is compact and it satisfies that $\dim \ker(I - \mathcal{T}_X) = 1$, which is generated by a strictly positive function, and*

$$\text{ran}(I - \mathcal{T}_X) = \left\{ f \in L^1(0, \infty) : \int_0^\infty f(x) dx = 0 \right\}.$$

Proof. The first step is to prove that \mathcal{T}_X is a compact operator. This means we have to prove that the set $A = \{\mathcal{T}_X[f] : \|f\|_1 \leq 1\}$ is relatively compact in L^1 .

First observe that $\|\mathcal{T}_X[f]\|_1 \leq p_\infty$ for all u with $\|f\|_1 \leq 1$, so A is bounded.

Second, we prove that

$$\int_r^\infty |\mathcal{T}_X[f](a)| da \rightarrow 0 \quad \text{uniformly when } r \rightarrow \infty.$$

Indeed for $r > \sigma$ we have

$$\begin{aligned} \int_r^\infty |\mathcal{T}_X[f](a)| da &\leq p_\infty \int_r^\infty \int_0^\infty |f(u)| e^{-\int_0^a p(s', u + s', X) ds'} du da \\ &\leq p_\infty \int_r^\infty \int_\sigma^\infty |f(u)| e^{-p_0(a-\sigma)} du da \\ &\leq p_\infty e^{p_0\sigma} \int_\sigma^\infty |f(u)| du \int_r^\infty e^{-p_0 a} da \\ &\leq p_\infty e^{p_0\sigma} \frac{e^{-p_0 r}}{p_0} \rightarrow 0. \end{aligned}$$

Now we prove the equicontinuity property. Observe that

$$\begin{aligned} \frac{d}{da} \mathcal{T}_X[f](a) &= \int_0^\infty (\partial_s p + \partial_a p)(a, u+a, X) e^{-\int_0^a p(s', u+s', X) ds'} f(u) du \\ &\quad - \int_0^\infty p(a, u+a, X)^2 e^{-\int_0^a p(s', u+s', X) ds'} f(u) du, \end{aligned}$$

thus for f with $\|f\|_1 \leq 1$ we have

$$\int_0^\infty \left| \frac{d}{da} \mathcal{T}_X[f](a) \right| da \leq \|\nabla p\|_\infty + p_\infty^2.$$

Therefore by the Kolmogorov-Frechet theorem we conclude that A is relatively compact so the operator \mathcal{T}_X is.

Furthermore, since n_X is the unique steady state of Equation (17) that is a probability density, from the linearity we deduce that any other function in $\ker(I - \mathcal{T}_X)$ is a multiple of N_X and thus $\dim \ker(I - \mathcal{T}_X) = 1$.

Next, we proceed to determine $\text{ran}(I - \mathcal{T}_X)$. Observe that adjoint operator $\mathcal{T}_X^*: L^\infty \rightarrow L^\infty$ is given by

$$\mathcal{T}_X^*[g](a) = \int_0^\infty p(u, u+a, X) \exp\left(-\int_0^u p(s', a+s', X) ds'\right) g(u) du,$$

and from Fredholm's alternative we get $\dim \ker(I - \mathcal{T}_X^*) = \dim \ker(I - \mathcal{T}_X) = 1$. Since $\mathcal{T}_X^*[g] \equiv 0$ for any constant function, we deduce that $\ker(I - \mathcal{T}_X^*)$ is the subspace of constant functions. Finally from orthogonality conditions we conclude that

$$\text{ran}(I - \mathcal{T}_X) = \left\{ f \in L^1(0, \infty) : \int_0^\infty f(x) dx = 0 \right\}.$$

□

A direct consequence of Lemma 3 is the following result

Lemma 4. *Assume that p is smooth respect to variable X , then $N_X(a)$ is also smooth with respect to X .*

Proof. Define the $F: L^1(0, \infty) \times (0, \infty) \rightarrow \text{ran}(I - \mathcal{T}_X) \times \mathbb{R}$ given by

$$F(N, X) = \left((I - \mathcal{T}_X)[N], \iint N(a) e^{-\int_0^a p(s', u+s', X) ds'} da ds - 1 \right),$$

so that for each X we have $F(N_X(a), X) = 0$. Observe that $D_N F$ is given by

$$D_N F[h] = \left((I - \mathcal{T}_X)[h], \iint h(a) e^{-\int_0^a p(s', u+s', X) ds'} da ds \right)$$

Thus by Lemma 3 this operator is an isomorphism and from the implicit function theorem we conclude that $N_X(a)$ depends smoothly on X . □

Remark 1. The lower bound condition (2) on the firing rate p is important to verify the existence of a steady state for System (1) and Doeblin's condition. For example, when we consider $X > 0$ and

$$p(s, a, X) = \mathbb{1}_{\{a-s > X\}},$$

then there is no steady state of the linear Equation (17), besides the zero solution. Indeed, from Equation (25) we deduce that the discharging flux N should satisfy

$$N(a) = e^{-a} \int_X^\infty N(u) du,$$

whose unique non-negative solution in $L^1(0, \infty)$ is $N \equiv 0$.

4 Steady states

Consider $n^* = n^*(s, a)$ with support in the set $\{s \leq a\}$. We are interested in the stationary solutions of the non-linear System (1) given by

$$\begin{cases} \partial_s n + \partial_a n + p(s, a, X)n = 0 & a > s > 0, \\ n(s = 0, a) = N(a) := \int_0^\infty p(a, u, X)n(a, u) du & a > 0, \\ X = \int_0^\infty N(a) da, \\ \iint n(s, a) da ds = 1, \quad n(s, a) \geq 0. \end{cases} \quad (27)$$

We define N_X as the respective conditional activity in terms of X . In order to have a steady state of the non-linear Problem (1), we must find $X > 0$ such that

$$X = \Phi(X) := \int_0^\infty N_X(a) da. \quad (28)$$

In the general case this equation has always a solution since the right-hand side is uniformly bounded thanks to estimate (3) and $N_X(a)$ depends continuously on X . By using the properties of the operator \mathcal{T}_X , we prove that under the weak interconnections regime the non-linear System (1) has a unique steady state.

Theorem 4. Assume (4) and that p smooth satisfies Assumption (2). Then for $\|\partial_X p\|_\infty$ small enough, System (1) has a unique steady state (n^*, N^*, X^*) .

Proof. The goal is to prove that Φ is a contraction in order to obtain a unique fixed point. In order to estimate $\partial_X N$ we make use of the implicit function theorem. By differentiating Equation (26) we get

$$\iint \partial_X(N_X) e^{-\int_0^s p(s', a+s', X) ds'} da ds = \iint N_X e^{-\int_0^s p(s', a+s', X) ds'} \left(\int_0^s \partial_X p(s', u' + s', X) ds' \right) da ds. \quad (29)$$

Furthermore, if we differentiate with respect to X the Equation (25), we get

$$\begin{aligned} \partial_X N_X(a) &= \int_0^\infty \left(-\partial_a e^{-\int_0^a p(s', u+s', X) ds'} \right) \partial_X N_X(u) du \\ &\quad + \int_0^\infty \left(-\partial_a \partial_X e^{-\int_0^a p(s', u+s', X) ds'} \right) N_X(u) du, \end{aligned}$$

i.e. $\partial_X N_X(a)$ satisfies the equation

$$\begin{aligned} (I - \mathcal{T}_X)[\partial_X N](a) &= \int_0^\infty p(a, u + a, X) e^{-\int_0^a p(s', u+s', X) ds'} N_X(u) du \\ &\quad - \int_0^\infty p(a, u + a, X) \left(\int_0^a \partial_X p(s', u + s', X) ds' \right) e^{-\int_0^a p(s', u+s', X) ds'} N_X(u) du. \end{aligned} \quad (30)$$

By using the implicit function theorem and the Condition (29) we can define an inverse of $I - \mathcal{T}_X$ which depends continuously on X . Observe that $\|(I - \mathcal{T}_X)^{-1}\|$ is uniformly bounded on X in the operator norm, since X is uniformly bounded. Thus, for the function Φ we get

$$\begin{aligned} |\Phi'(X)| &= \left| \int \partial_X N_X(a) da \right| \\ &\leq \|(I - \mathcal{T}_X)^{-1}\| \|\partial_X p\|_\infty \left(\iint (1 + p_\infty a) e^{-\int_0^a p(s', u+s', X) ds} N_X(u) da du \right) \\ &\leq C \|\partial_X p\|_\infty \left(\iint (1 + p_\infty a) e^{-p_0 a} N_X(u) da du \right) \\ &\leq C \|\partial_X p\|_\infty p_\infty \left(\int (1 + p_\infty a) e^{-p_0 a} da \right), \end{aligned}$$

so that for $\|\partial_X p\|_\infty$ small enough Φ is a contraction and we conclude the result. \square

5 Convergence to equilibrium

After studying the linear case, we are now ready to prove convergence to the steady state under the weak interconnection regime, i.e. $\|\partial_X p\|_\infty$ small enough, by a perturbation argument.

Theorem 5 (Convergence to equilibrium). *Assume that $n_0 \in L^1(\mathcal{D})$ satisfies Assumption (4) and that p Lipschitz satisfies Assumption (2). Let (n^*, N^*, X^*) be the corresponding stationary state of System (1). Then for $\|\partial_X p\|_\infty$ small enough, there exist $C, \lambda > 0$ such that the solution n of System (1) satisfies*

$$\|n(t) - n^*\|_{L^1_{s,a}} \leq C e^{-\lambda t} \|n_0 - n^*\|_{L^1_{s,a}}, \quad \forall t \geq 0.$$

Moreover $\|N(t) - N^*\|_{L^1}$ and $|X(t) - X^*|$ converge exponentially to 0 when $t \rightarrow \infty$.

Proof. Observe that n satisfies the evolution equation

$$\partial_t n = \mathcal{L}_X[n] := -\partial_s n - \partial_a n - p(s, a, X(t))n + \delta_{\{s=0\}}(s, a) \int_0^\infty p(a, u, X(t))n(t, a, u) du,$$

where $\delta_{\{s=0\}}(s, a)$ is the measure along the line $\{(0, a) : a \geq 0\}$. We can rewrite the evolution equation as

$$\partial_t n = \mathcal{L}_{X^*}[n] + (\mathcal{L}_X[n] - \mathcal{L}_{X^*}[n]) = \mathcal{L}_{X^*}[n] + h. \quad (31)$$

with $h(t, s, a)$ given by

$$h = (p(s, a, X^*) - p(s, a, X(t)))n(t, s, a) + \delta_{\{s=0\}}(s, a) \int_0^\infty (p(a, u, X(t)) - p(a, u, X^*))n(t, a, u) du. \quad (32)$$

Let $P_t: L^1(\mathcal{D}) \rightarrow L^1(\mathcal{D})$ be the linear semi-group associated to operator \mathcal{L}_{X^*} . As in the proof of Lemma 1, P_t extends to space $(\mathcal{M}(\mathcal{D}), \|\cdot\|_{TV})$ in order to be able to evaluate at the measure h . Since $P_t n^* = n^*$ for all $t \geq 0$, we get that n satisfies

$$n - n^* = P_t(n_0 - n^*) + \int_0^t P_{t-\tau} h(\tau, s, a) d\tau, \quad (33)$$

so we need find an estimate for the function h . Observe that we have the following inequalities:

$$\begin{aligned} \|h(t)\|_{L^1_{s,a}} &\leq 2\|\partial_X p\|_\infty |X(t) - X^*|, \\ |X(t) - X^*| &\leq \|N(t) - N^*\|_1, \\ \|N(t) - N^*\|_1 &\leq \|\frac{\partial p}{\partial X}\|_\infty |X(t) - X^*| + p_\infty \|n(t) - n^*\|_{L^1_{s,a}}, \end{aligned}$$

and since $\|\partial_X p\|_\infty < 1$ we get

$$\begin{aligned} \|h(t)\|_{L^1_{s,a}} &\leq \frac{2p_\infty \|\partial_X p\|_\infty}{1 - \|\partial_X p\|_\infty} \|n(t) - n^*\|_{L^1_{s,a}}, \\ |X(t) - X^*| &\leq \frac{p_\infty}{1 - \|\partial_X p\|_\infty} \|n(t) - n^*\|_{L^1_{s,a}}, \\ \|N(t) - N^*\|_1 &\leq p_\infty \left(\frac{\|\partial_X p\|_\infty}{1 - \|\partial_X p\|_\infty} + 1 \right) \|n(t) - n^*\|_{L^1_{s,a}}, \end{aligned}$$

thus by taking norm in Equality (33) and applying Doeblin's Theorem we obtain

$$\begin{aligned} \|n(t) - n^*\|_{L^1_{s,a}} &\leq \|P_t(n_0 - n^*)\|_{L^1_{s,a}} + \int_0^t \|P_{t-\tau} h(\tau)\|_{L^1_{s,a}} d\tau \\ &\leq \frac{e^{-\lambda t}}{1 - \alpha} \|n_0 - n^*\|_{L^1_{s,a}} + \frac{1}{1 - \alpha} \int_0^t e^{-\lambda(t-\tau)} \|h(\tau)\|_{L^1_{s,a}} d\tau \\ &\leq \frac{e^{-\lambda t}}{1 - \alpha} \|n_0 - n^*\|_{L^1_{s,a}} + C \int_0^t e^{-\lambda(t-\tau)} \|n(\tau) - n^*\|_{L^1_{s,a}} d\tau, \end{aligned}$$

with $C := \frac{1}{1-\alpha} \frac{2p_\infty \|\partial_X p\|_\infty}{1 - \|\partial_X p\|_\infty}$. By using Gronwall's inequality with respect to the function $e^{\lambda t} \|n(t) - n^*\|_{L^1_{s,a}}$ we conclude

$$\|n(t) - n^*\|_{L^1_{s,a}} \leq \frac{e^{-(\lambda-C)t}}{1 - \alpha} \|n_0 - n^*\|_{L^1_{s,a}},$$

so that for $\|\partial_X p\|_\infty$ small enough we have $C < \lambda$ and we deduce the exponential convergence of $n(t, \cdot, \cdot)$, $N(t, \cdot)$ and $X(t)$ when $t \rightarrow \infty$. \square

6 Numerical Simulations

In order to illustrate the theoretical long time results and other possible behaviors of System (1), we present numerical simulations for different firing rates and initial data. The numerical illustrations below are obtained by solving the equation (1) with a classical first-order upwind scheme.

We focus in displaying the discharging flux $N(t, a)$ and the total activity $X(t)$ since these two elements determine the general behavior of system (1).

6.1 Example 1: Convergence to equilibrium

For our first example, we choose as initial data $n_0(s, a) = e^{-a}$ and the firing rate is given by

$$p = \mathbb{1}_{\{s > X\}} + \mathbb{1}_{\{s-a > X\}},$$

which corresponds to an inhibitory case since p is decreasing with respect to X . Moreover, this particular form of p is decomposed as the sum of two simple threshold functions with the first one depending only on the first elapsed time and the second one depending on the difference between the last two discharges.




Figure 4: **Example 1.** Case $n_0(s, a) = e^{-a}$ and $p = \mathbb{1}_{\{s > X\}} + \mathbb{1}_{\{s-a > X\}}$.

In this case the solution simply converges to a steady state, as we see in Figure 4 for the discharging flux N and the total activity X . From Equation (25) we note that the discharging flux at equilibrium N^* has a jump discontinuity at X^* , which is consistent with the numerical solution. This convergence is compatible with Theorem 5.

6.2 Example 2: Jump discontinuities

We now consider the initial data $n_0(s, a) = 2 \cdot \mathbb{1}_{\{2 > a > s+1\}}$ and the firing rate is given

$$p = \mathbb{1}_{\{s > e^{-X}\}} + \mathbb{1}_{\{s-a > e^{-X}\}},$$

which corresponds to an excitatory case since p is increasing with respect to X .

Like the previous example the solution converges to the steady state, but the total activity X shows three jump discontinuities as we see in Figure 5. The multiple jump discontinuities are consequence of the contribution of the term depending on the difference between the two elapsed times. Furthermore, solutions convergent to the steady state that present a single jump discontinuity were already observed in Caceres et al. [29] for the classical elapsed time model. The phenomenon of multiple jumps discontinuities in Figure 5 is an extension for the case of Equation (1).




Figure 5: **Example 2.** Case $n_0(s, a) = 2 \cdot \mathbb{1}_{\{2>a>s+1\}}$ and $p = \mathbb{1}_{\{s>e^{-x}\}} + \mathbb{1}_{\{s-a>e^{-x}\}}$.

6.3 Example 3: Periodic solutions and stabilization

Next, we choose initial data $n_0(s, a) = \frac{1}{2}e^{-(a-1)}\mathbb{1}_{\{a>\max(s,1)\}}$ and the firing rate is given by

$$p = \varphi(X)\mathbb{1}_{\{s>1\}}, \quad \varphi(u) = \frac{10u^2}{u^2 + 1} + 0.5,$$

which corresponds to an excitatory case since $\varphi'(u) > 0$. Since p does not depend on a , we take advantage by solving the classical elapsed time Equation (5) after integrating with respect to a , as we remarked in the introduction.




Figure 6: **Example 3.1.** Case $n_0(s, a) = \frac{1}{2}e^{-(a-1)}\mathbb{1}_{\{a>\max(s,1)\}}$ and $p = \varphi(X)\mathbb{1}_{\{s>1\}}$.

For these data, both the discharging flux N and the total activity X are asymptotic to a periodic

pattern as we see in Figure 6. Similar examples on periodic solutions were found in Caceres et al. [29] in the classical elapsed time model for the same type of firing rates.

However, when we incorporate the effects of the difference between the elapsed times the periodic regime changes. For the same initial data and

$$p = \varphi(X)\mathbb{1}_{\{s>1\}} + \mathbb{1}_{\{s-a>X\}},$$

we observe in Figure 7 that, with the term depending on the difference between the two elapsed times, the solution of System (1) converges to the steady state.




Figure 7: **Example 3.2.** Case $n_0(s, a) = \frac{1}{2} e^{-(a-1)} \mathbb{1}_{\{a>\max(s,1)\}}$ and $p = \varphi(X)\mathbb{1}_{\{s>1\}} + \mathbb{1}_{\{s-a>X\}}$.

7 Perspectives

By means of Doeblin's theory applied to a more complex problem than the classical elapsed time model, we managed to understand the dynamics of System (1) for weak non-linearities by adapting the ideas of Cañizo et al. [5]. However, aspects like well-posedness and the asymptotic behavior for strong interconnections are still an open problem as in the classical elapsed time model.

Concerning the strongly inhibitory case, it remains pending to prove uniqueness of the steady state. Whilst in the classical elapsed time model this problem is reduced to a simple equation, for the model with two elapsed times we have to prove uniqueness for the integral Equation (28). Moreover, we conjecture in the general case that the speed of convergence to a steady state must be exponential like it is expected for the classical elapsed equation.

With respect to the existence of periodic solutions, we still have to find or construct a non-trivial example relying on dynamics for two elapsed times. The only examples we have found so far are adaptations of solutions of the classical elapsed time equation that were obtained in Caceres et al. [29] and these types of solutions presents jump discontinuities, making them difficult to analyze. Furthermore, it remains as an open problem to find continuous periodic solutions.

Acknowledgements

NT has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 754362. BP has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 740623). DS has received support from ANR ChaMaNe No: ANR-19-CE40-0024.



References

- [1] Delphine Salort Benoît Perthame. On a voltage-conductance kinetic system for integrate & fire neural networks. *Kinetic & Related Models*, 6(4):841–864, 2013.
- [2] Nicolas Brunel. Dynamics of sparsely connected networks of excitatory and inhibitory spiking neurons. *Journal of computational neuroscience*, 8(3):183–208, 2000.
- [3] María J Cáceres, José A Carrillo, and Benoît Perthame. Analysis of nonlinear noisy integrate & fire neuron models: blow-up and steady states. *The Journal of Mathematical Neuroscience*, 1(1):7, 2011.
- [4] José A Cañizo, Pierre Gabriel, and Havva Yoldaş. Spectral gap for the growth-fragmentation equation via harris's theorem. *arXiv preprint arXiv:2004.08343*, 2020.
- [5] José A. Cañizo and Havva Yoldaş. Asymptotic behaviour of neuron population models structured by elapsed-time. *Nonlinearity*, 32(2):464, 2019.
- [6] José Antonio Carrillo, Benoît Perthame, Delphine Salort, and Didier Smets. Qualitative properties of solutions for the noisy integrate and fire model in computational neuroscience. *Nonlinearity*, 28(9):3365, 2015.
- [7] Julien Chevallier, María José Cáceres, Marie Doumic, and Patricia Reynaud-Bouret. Microscopic approach of a time elapsed neural model. *Mathematical Models and Methods in Applied Sciences*, 25(14):2669–2719, 2015.
- [8] Grégory Dumont and Pierre Gabriel. The mean-field equation of a leaky integrate-and-fire neural network: measure solutions and steady states. *Nonlinearity*, 33(12):6381, 2020.
- [9] Grégory Dumont, Jacques Henry, and Carmen Oana Tarniceriu. Noisy threshold in neuronal models: connections with the noisy leaky integrate-and-fire model. *J. Math. Biol.*, 73(6-7):1413–1436, 2016.
- [10] Grégory Dumont, Jacques Henry, and Carmen Oana Tarniceriu. A theoretical connection between the noisy leaky integrate-and-fire and the escape rate models: the non-autonomous case. *Math. Model. Nat. Phenom.*, 15:Paper No. 59, 20, 2020.

- [11] Claudia Fonte and Valentin Schmutz. Long time behavior of an age and leaky memory-structured neuronal population equation. *arXiv preprint arXiv:2106.11110*, 2021.
- [12] Nicolas Fournier and Benoît Perthame. A non-expanding transport distance for some structured equations. *arXiv preprint arXiv:2102.04092*, 2021.
- [13] Pierre Gabriel. Measure solutions to the conservative renewal equation. *ESAIM: Proceedings and Surveys*, 62:68–78, 2018.
- [14] Wulfram Gerstner and Werner M. Kistler. *Spiking neuron models: Single neurons, populations, plasticity*. Cambridge university press, 2002.
- [15] Moon-Jin Kang, Benoît Perthame, and Delphine Salort. Dynamics of time elapsed inhomogeneous neuron network model. *Comptes Rendus Mathématique*, 353(12):1111–1115, 2015.
- [16] Jian-guo Liu, Ziheng Wang, Yuan Zhang, and Zhennan Zhou. Rigorous justification of the fokker-planck equations of neural networks based on an iteration perspective. *arXiv preprint arXiv:2005.08285*, 2020.
- [17] Cheng Ly and Daniel Tranchina. Spike train statistics and dynamics with synaptic input from any renewal process: a population density approach. *Neural Computation*, 21(2):360–396, 2009.
- [18] Philippe Michel, Stéphane Mischler, and Benoît Perthame. General relative entropy inequality: an illustration on growth models. *Journal de mathématiques pures et appliquées*, 84(9):1235–1260, 2005.
- [19] Stéphane Mischler, Cristobal Quiñinao, and Qilong Weng. Weak and strong connectivity regimes for a general time elapsed neuron network model. *Journal of Statistical Physics*, 173(1):77–98, 2018.
- [20] Stéphane Mischler and Qilong Weng. Relaxation in time elapsed neuron network models in the weak connectivity regime. *Acta Applicandae Mathematicae*, 157(1):45–74, 2018.
- [21] Khashayar Pakdaman, Benoît Perthame, and Delphine Salort. Dynamics of a structured neuron population. *Nonlinearity*, 23(1):55–75, 2010.
- [22] Khashayar Pakdaman, Benoît Perthame, and Delphine Salort. Relaxation and self-sustained oscillations in the time elapsed neuron network model. *SIAM J. Appl. Math.*, 73(3):1260–1279, 2013.
- [23] Khashayar Pakdaman, Benoît Perthame, and Delphine Salort. Adaptation and fatigue model for neuron networks and large time asymptotics in a nonlinear fragmentation equation. *J. Math. Neurosci.*, 4:Art. 14, 26, 2014.
- [24] Benoît Perthame. *Transport equations in biology*. Springer Science & Business Media, 2006.
- [25] Benoît Perthame and Delphine Salort. Derivation of a voltage density equation from a voltage-conductance kinetic model for networks of integrate-and-fire neurons. *Communications in Mathematical Sciences*, 17(5):1193–1211, 2019.
- [26] Benoît Perthame, Delphine Salort, and Gilles Wainrib. Distributed synaptic weights in a lif neural network and learning rules. *Physica D: Nonlinear Phenomena*, 353:20–30, 2017.

- [27] Joël Pham, Khashayar Pakdaman, Jean Champagnat, and Jean-François Vibert. Activity in sparsely connected excitatory neural networks: effect of connectivity. *Neural Networks*, 11(3):415–434, 1998.
- [28] Tilo Schwalger and Anton V. Chizhov. Mind the last spike—firing rate models for mesoscopic populations of spiking neurons. *Current opinion in neurobiology*, 58:155–166, 2019.
- [29] Nicolás Torres, María J Cáceres, Benoît Perthame, and Delphine Salort. An elapsed time model for strongly coupled inhibitory and excitatory neural networks. *Physica D: Nonlinear Phenomena*, page 132977, 2021.