



HAL
open science

Draft Genome Sequences of the Black Truffles *Tuber brumale* Vittad. and *Tuber indicum* Cook & Masee

Emmanuelle Morin, Claude Murat, Nicolas Cichocki, Herminia de La Varga,
Annegret Kohler, Jianping Xu, Igor Grigoriev, Francis Martin

► To cite this version:

Emmanuelle Morin, Claude Murat, Nicolas Cichocki, Herminia de La Varga, Annegret Kohler, et al.. Draft Genome Sequences of the Black Truffles *Tuber brumale* Vittad. and *Tuber indicum* Cook & Masee. *Microbiology Resource Announcements*, 2021, 10 (4), 10.1128/MRA.00799-20 . hal-03324444

HAL Id: hal-03324444

<https://hal.science/hal-03324444>

Submitted on 18 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.




L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Draft Genome Sequences of the Black Truffles *Tuber brumale* Vittad. and *Tuber indicum* Cook & Masee

 Emmanuelle Morin,^a Claude Murat,^a Nicolas Cichocki,^a Herminia De la Varga,^{a*} Annegret Kohler,^a  Jianping Xu,^b Igor V. Grigoriev,^c  Francis M. Martin^a

^aUniversité de Lorraine, INRAE, IAM, Nancy, France

^bMcMaster University, Department of Biology, Hamilton, Canada

^cDOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, California, USA

ABSTRACT *Tuber brumale* and *Tuber indicum* (Pezizomycetes) are two edible black truffles establishing ectomycorrhizal symbiosis with trees and shrubs. *T. brumale* is ubiquitous in Europe, and *T. indicum* is mainly found in China. Here, we present the draft genome sequences of *T. brumale* and *T. indicum*.

The black truffles *Tuber brumale* Vittadini and *Tuber indicum* Cook & Masee are ectomycorrhizal ascomycetes. *T. brumale* is widespread in Europe, except in the boreal and Arctic regions (1). This species often competes with the Périgord black truffle (*Tuber melanosporum*) in truffle orchards (2). *T. indicum* is found mainly in the Chinese provinces of Yunnan and Sichuan (3). The two species belong to the *Melanosporum* phylogenetic clade (4) and have morphological features similar to those of *T. melanosporum*, making their distinctions sometimes difficult (5). Together with the published genome sequences of *T. aestivum*, *T. borchii*, *T. magnatum*, and *T. melanosporum* (6–8), these newly sequenced genomes will allow a better understanding of the evolution, biology, and ecology of truffles.

For genome and RNA sequencing, a *T. brumale* fruiting body was harvested in Lozère (Occitanie, France) in March 2014, and a *T. indicum* fruiting body was purchased at a French market in 2013. For both species, genomic DNA (gDNA) was extracted from 2 g of fruiting body by using a modified cetyl trimethylammonium bromide (CTAB) protocol (9). Total RNA was extracted using the RNeasy plant minikit (Qiagen) as described earlier (6). The gDNA and the Illumina TruSeq Nano kit were used to construct paired-end libraries (2 × 100 bp for both and 2 × 125 bp for *T. brumale*) as well as mate pair libraries (with insert sizes of 3 and 8 kbp) using the Illumina Nextera mate pair kit. In addition, paired-end libraries (2 × 100 bp and 2 × 125 bp) were generated from total RNA using the Illumina TruSeq stranded mRNA kit. Sequencing was performed at the GeT-PlaGe sequencing facility (Toulouse, France) using the Illumina HiSeq 2500 platform. The raw Illumina reads were trimmed of adapter sequences and low-quality bases using Trimmomatic v.0.32 (10) with the following parameters: TRAILING:20, LEADING:20, SLIDINGWINDOW:4:20, and MINLEN:70. Assembly of the genomes was carried out using ALLPATHS-LG v.46154 (11) and GapCloser v.1.12.6 (12). The genome assemblies were then annotated using the Joint Genome Institute (JGI) annotation pipeline (13, 14).

The sequencing data statistics are shown in Table 1. The genome sizes of *T. brumale* and *T. indicum* are in the range of other truffle species, from 97.18 to 192 Mb (6–8).

RepeatScout v.1.0.5 (15) was used to identify *de novo* repetitive DNA in the genome assemblies as reported by Peter et al. (16). RepeatMasker v.4.0.9 (17) was used to estimate the repeat element coverage in the genomes. Transposable elements constitute 61.5% and 47.1% of the *T. brumale* and *T. indicum* genomes, respectively. Default parameters were used for all software except where otherwise noted.

Citation Morin E, Murat C, Cichocki N, De la Varga H, Kohler A, Xu J, Grigoriev IV, Martin FM. 2021. Draft genome sequences of the black truffles *Tuber brumale* Vittad. and *Tuber indicum* Cook & Masee. *Microbiol Resour Annot* 10:e00799-20. <https://doi.org/10.1128/MRA.00799-20>.

Editor Antonis Rokas, Vanderbilt University

Copyright © 2021 Morin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Emmanuelle Morin, emmanuelle.morin@inrae.fr, or Francis M. Martin, francis.martin@inrae.fr.

* Present address: Herminia De la Varga, Fertinagro Biotech, S.L., R+D+i Department, Teruel, Spain.

Received 5 August 2020

Accepted 11 January 2021

Published 28 January 2021

TABLE 1 Genomic features and raw data of *Tuber brumale* and *Tuber indicum*

Organism	Source	No. of reads	Draft genome size (Mb)	No. of scaffolds	N_{50} (bp)	G+C content (%)	Mean coverage (×)	SRA accession no.	GenBank accession no.	BioProject accession no.
<i>Tuber brumale</i>	DNA	136,348,163	171.44	1,475	336,267	46.46	131.63	SRR12018987, SRR12018988, SRR12018989	JACCEG000000000	PRJNA633036
		135,457,273						SRR12018993, SRR12018994, SRR12018995		
	RNA	128,865,953						SRR12018990, SRR12018991, SRR12018992		
<i>Tuber indicum</i>	DNA	558,521,206	110.49	734	538,733	47.41	239.48	SRR12104989, SRR12104990, SRR12104991	JACCEH000000000	PRJNA633038
	RNA	86,652,466						SRR12104986, SRR12104987, SRR12104988		

A total of 12,380 protein-coding genes for *T. brumale* and 11,870 protein-coding genes for *T. indicum* were predicted. The number of protein-coding genes is also in the range of other truffle species, from 9,344 to 12,346 protein-coding genes (6–8).

Data availability. The draft whole-genome shotgun projects were deposited in DDBJ/ENA/GenBank. The SRA and GenBank accession numbers for *T. brumale* and *T. indicum* are listed in Table 1. The genome assemblies and annotations are also available at the JGI-DOE Mycocosm portal (13) (https://mycocosm.jgi.doe.gov/Tubbr1_1 and https://mycocosm.jgi.doe.gov/Tubin1_1).

ACKNOWLEDGMENTS

This research was supported by the Laboratory of Excellence ARBRE (ANR-11-LABX-0002-01), the Region Lorraine, and the European Regional Development Fund. The project was also funded by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, and supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231 within the framework of the Mycorrhizal Genomics Initiative (CSP no. 305), Metatranscriptomics of Forest Soil Ecosystems project (CSP no. 570), and the 1000 Fungal Genomes Project (CSP no. 1974).

We are grateful to Jean-Yves Magaud for providing the *T. brumale* ascocarp and to François Le Tacon for purchasing the *T. indicum* ascocarps. We thank Alan Kuo, Sajeet Haridas, and Stephen Mondo from JGI for their advice on gene annotation.

REFERENCES

- Merényi Z, Varga T, Bratek Z. 2016. *Tuber brumale*: a controversial tuber species, p 49–68. In Zambonelli A, Iotti M, Murat C (ed), True truffle (*Tuber* spp.) in the world. Springer International Publishing, Cham, Switzerland.
- Olivier J-M, Savignac J-C, Sourzat P. 2012. Truffe et trufficulture. FANLAC Editions, Périgueux, France.
- Chen J, Murat C, Oviatt P, Wang Y, Le Tacon F. 2016. The black truffles *Tuber melanosporum* and *Tuber indicum*, p 19–32. In Zambonelli A, Iotti M, Murat C (ed), True truffle (*Tuber* spp.) in the world. Springer International Publishing, Cham, Switzerland.
- Bonito G, Smith ME, Nowak M, Healy RA, Guevara G, Cázares E, Kinoshita A, Nouhra ER, Dominguez LS, Tedersoo L, Murat C, Wang Y, Moreno BA, Pfister DA, Nara K, Zambonelli A, Trappe JM, Vilgalys R. 2013. Historical biogeography and diversification of truffles in the Tuberales and their newly identified southern hemisphere sister lineage. PLoS One 81:e52765. <https://doi.org/10.1371/journal.pone.0052765>.
- Murat C. 2015. Forty years of inoculating seedlings with truffle fungi: past and future perspectives. Mycorrhiza 25:77–81. <https://doi.org/10.1007/s00572-014-0593-4>.
- Murat C, Payen T, Noel B, Kuo A, Morin E, Chen J, Kohler A, Krizsán K, Balestrini R, Da Silva C, Montanini B, Hainaut M, Levati E, Barry KW, Belfiori B, Cichocki N, Clum A, Dockter RB, Fauchery L, Guy J, Iotti M, Le Tacon F, Lindquist EA, Lipzen A, Malagnac F, Mello A, Molinier V, Miyauchi S, Poulain J, Riccioni C, Rubini A, Sitrit Y, Splivallo R, Traeger S, Wang M, Žifčáková L, Wipf D, Zambonelli A, Paolucci F, Nowrousian M, Ottonello S, Baldrian P, Spatafora JW, Henrissat B, Nagy LG, Aury J-M, Wincker P, Grigoriev IV, Bonfante P, Martin FM. 2018. Pezizomycetes genomes reveal the molecular basis of ectomycorrhizal truffle lifestyle. Nat Ecol Evol 2:1956–1965. <https://doi.org/10.1038/s41559-018-0710-4>.
- Murat C, Kuo A, Barry KW, Clum A, Dockter RB, Fauchery L, Iotti M, Kohler A, LaButti K, Lindquist EA, Lipzen A, Morin E, Wang M, Grigoriev IV, Zambonelli A, Martin FM. 2018. Draft genome sequence of *Tuber borchii* Vittad., a whitish edible truffle. Genome Announc 6:e00537-18. <https://doi.org/10.1128/genomeA.00537-18>.
- Martin F, Kohler A, Murat C, Balestrini R, Coutinho PM, Jaillon O, Montanini B, Morin E, Noel B, Percudani R, Porcel B, Rubini A, Amicucci A, Amselem J, Anthonard V, Arcioni S, Artiguenave F, Aury J-M, Ballario P, Bolchi A, Brenna

- A, Brun A, Buée M, Cantarel B, Chevalier G, Couloux A, Da Silva C, Denoeud F, Duplessis S, Ghignone S, Hilselberger B, Iotti M, Marçais B, Mello A, Miranda M, Pacioni G, Quesneville H, Riccioni C, Ruotolo R, Splivallo R, Stocchi V, Tisserant E, Viscomi AR, Zambonelli A, Zampieri E, Henrissat B, Lebrun M-H, Paolucci F, Bonfante P, Ottonello S, et al. 2010. Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* 464:1033–1038. <https://doi.org/10.1038/nature08867>.
9. Kohler A, Kuo A, Nagy LG, Morin E, Barry KW, Buscot F, Canbäck B, Choi C, Cichocki N, Clum A, Colpaert J, Copeland A, Costa MD, Doré J, Floudas D, Gay G, Girlanda M, Henrissat B, Herrmann S, Hess J, Högberg N, Johansson T, Khouja H-R, LaButti K, Lahrmann U, Levasseur A, Lindquist EA, Lipzen A, Marmeisse R, Martino E, Murat C, Ngan CY, Nehls U, Plett JM, Pringle A, Ohm RA, Perotto S, Peter M, Riley R, Rineau F, Ruytinx J, Salamov A, Shah F, Sun H, Tarkka M, Tritt A, Veneault-Fourrey C, Zuccaro A, Tunlid A, Grigoriev IV, Mycorrhizal Genomics Initiative Consortium, et al. 2015. Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nat Genet* 47:410–415. <https://doi.org/10.1038/ng.3223>.
 10. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
 11. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A* 108:1513–1518. <https://doi.org/10.1073/pnas.1017351108>.
 12. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu S-M, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam T-W, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience* 1:18. <https://doi.org/10.1186/2047-217X-1-18>.
 13. Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otilar R, Riley R, Salamov A, Zhao X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids Res* 42:D699–D704. <https://doi.org/10.1093/nar/gkt1183>.
 14. Kuo A, Bushnell B, Grigoriev IV. 2014. Fungal genomics: sequencing and annotation, p 1–52. *In* Martin F (ed), *Advances in botanical research: fungi*, vol 70. Elsevier Academic Press, Cambridge, MA.
 15. Price AL, Jones NC, Pevzner PA. 2005. De novo identification of repeat families in large genomes. *Bioinformatics* 21:i351–i358. <https://doi.org/10.1093/bioinformatics/bti1018>.
 16. Peter M, Kohler A, Ohm RA, Kuo A, Krützmann J, Morin E, Arend M, Barry KW, Binder M, Choi C, Clum A, Copeland A, Grisel N, Haridas S, Kipfer T, LaButti K, Lindquist E, Lipzen A, Maire R, Meier B, Mihaltcheva S, Molinier V, Murat C, Pöggeler S, Quandt CA, Sperisen C, Tritt A, Tisserant E, Crous PW, Henrissat B, Nehls U, Egli S, Spatafora JW, Grigoriev IV, Martin FM. 2016. Ectomycorrhizal ecology is imprinted in the genome of the dominant symbiotic fungus *Cenococcum geophilum*. *Nat Commun* 7:12662. <https://doi.org/10.1038/ncomms12662>.
 17. Smit AFA, Hubley R, Green P. 2015. RepeatMasker Open 4.0. <http://www.repeatmasker.org>.