



HAL
open science

Luminescent lanthanide small molecules, polymetallic dendrimer complexes, MOF and metallacrowns for biological imaging: from fundamental research to applications

S. Petoud, V. Eliseeva S., I. Martinic, Franck Suzenet, L. Pecoraro V.

► To cite this version:

S. Petoud, V. Eliseeva S., I. Martinic, Franck Suzenet, L. Pecoraro V.. Luminescent lanthanide small molecules, polymetallic dendrimer complexes, MOF and metallacrowns for biological imaging: from fundamental research to applications. ICP 2017 - 28th International Conference on Photochemistry, Jul 2017, Strasbourg, France. hal-03326196

HAL Id: hal-03326196

<https://hal.science/hal-03326196v1>

Submitted on 3 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Luminescent Lanthanide Small Molecules, Polymetallic Dendrimer Complexes, MOF and Metallacrowns for Biological Imaging: from Fundamental Research to Applications

Stéphane Petoud¹, Svetlana V. Eliseeva², Ivana Martinić², Franck Suzenet³,
Vincent L. Pecoraro⁴

¹ Department of Inorganic Chemistry, University of Geneva, CH1211 Geneva 4, Switzerland

² Centre de Biophysique Moléculaire, CNRS UPR 4301, F-45071 Orléans Cedex 2, France

³ Institut de Chimie Organique et Analytique - ICOA UMR7311, F-45067 Orléans Cedex 2,
France

⁴ Department of Chemistry, Willard H. Dow Laboratories, University of Michigan, Ann Arbor,
Michigan 48109, USA

E-mail: stephane.petoud.petoud@inserm.fr

Fluorescence and luminescence based detection techniques possess important advantages for bioanalytical applications and biologic imaging: high sensitivity, high cellular resolution, portability, versatility and low costs of instrumentation. A common characteristic of biologic analytes present in biological systems is their presence in small quantities among complex matrices such as blood, cells, tissue and organs. These matrices emit a significant background fluorescence (autofluorescence), limiting detection sensitivity.

The luminescence of lanthanide cations has several complementary advantages over the fluorescence of organic fluorophores and semiconductor nanocrystals, such as sharp emission bands for spectral discrimination from background emission, long luminescence lifetimes for temporal discrimination and strong resistance to photobleaching. In addition, several lanthanides emit near-infrared (NIR) photons that can cross deeply tissues for non-invasive investigations and that result in improved detection sensitivity due to the absence of native NIR luminescence from tissues and cells (autofluorescence).

The main requirement to generate lanthanide emission is to sensitize them with an appropriate chromophore (“antenna effect”). The choice of this antenna allows the tuning of the excitation wavelength of the resulting complexes. We will discuss in this paper different strategies and aspects of fundamental and applied research to create imaging agents suitable for optical imaging.

We will discuss different systems that we have created in the last few years including Small Molecules, Polymetallic Dendrimer Complexes, MOF and Metallacrowns

Funding: This research has received funding from the European Community's Marie Curie Seventh Framework Programme (FP7/2007-2013: n° 316906 (ITN Luminet) and n° 611488 (IRSES Metallacrowns)) and in part by La Ligue Contre le Cancer, La Région Centre, Cancéropôle Grand Ouest, Agence Nationale de la Recherche (Lumiphage ANR-13-BSV5-009, NIRA ANR-13-BS08-0011) and the US National Science Foundation (CH-1361779).

References:

- [1] I. Martinić, S.V. Eliseeva, S. Petoud, *J. Lumin.* DOI : 10.1016/j.jlumin.2016.09.058
- [2] C. Y. Chow, S. V. Eliseeva, E. R. Trivedi, T. N. Nguyen, J. W. Kampf, S. Petoud, V. L. Pecoraro, *J. Amer. Chem. Soc.* 2016, 138, 5100.
- [3] A. Foucault-Collet, C. M. Shade, I. Nazarenko, S. Petoud, S. V. Eliseeva, *Angew. Chem.*, 2014, 53, 2927.
- [4] A. Foucault-Collet, K. A. Gogick, K. A. White, S. Villette, A. Pallier, G. Collet, C. Kieda, T. Lib, S. J. Geib, N. L. Rosi*, S. Petoud*, *Proc. Natl. Acad. Sci. USA*, 2013, 110, 17199.
- [5] K. A. White, D. A. Chengelis, K. A. Gogick, J. Stehman, N. L. Rosi, S. Petoud, *J. Am. Chem. Soc.*, 2009, 131, 18069-18071.

Figure 1. a) Schematic Representation of a Polymetallic Dendrimer Complex b) Structure of a Near-infrared Emitting Metallacrown c) Structure of a luminescent nanoMOF

