



HAL
open science

Expedited NMR Assignment of Small- to Medium-Sized Molecules with Improved HSQC–CLIP–COSY Experiments

Tamás Gyöngyösi, István Timári, Davy Sinnaeve, Burkhard Luy, Katalin E. Kövér

► **To cite this version:**

Tamás Gyöngyösi, István Timári, Davy Sinnaeve, Burkhard Luy, Katalin E. Kövér. Expedited NMR Assignment of Small- to Medium-Sized Molecules with Improved HSQC–CLIP–COSY Experiments. *Analytical Chemistry*, 2021, 93 (6), pp.3096-3102. <10.1021/acs.analchem.0c04124>. <hal-03328594>

HAL Id: hal-03328594

<https://hal.science/hal-03328594v1>

Submitted on 30 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

This document is the unedited Author's version of a Submitted Work that was subsequently accepted for publication in *Analytical Chemistry*, copyright © American Chemical Society after peer review. To access the final edited and published work see: <https://doi.org/10.1021/acs.analchem.0c04124>

Expedited NMR Assignment of Small to Medium-Sized Molecules with Improved HSQC-CLIP-COSY Experiments

Tamás Gyöngyösi,^{a,b} István Timári,^{a,*} Davy Sinnaeve,^{c,d} Burkhard Luy,^e Katalin E. Kövér^{a,b,*}

^aDepartment of Inorganic and Analytical Chemistry, Faculty of Science and Technology, University of Debrecen, Egyetem tér 1, H-4032 Debrecen, Hungary

^bMTA-DE Molecular Recognition and Interaction Research Group, University of Debrecen, Egyetem tér 1, H-4032 Debrecen, Hungary

^cUniv. Lille, Inserm, Institut Pasteur de Lille, CHU Lille, U1167 – Labex DISTALZ – RID-AGE – Risk Factors and Molecular Determinants of Aging-Related Diseases, 59000 Lille, France

^dCNRS, ERL9002 – Integrative Structural Biology, F-59000 Lille, France

^eInstitute of Organic Chemistry and Institute for Biological Interfaces 4 – Magnetic Resonance, Karlsruhe Institute of Technology (KIT), Fritz-Haber-Weg 6, 76131 Karlsruhe, Germany

*Corresponding authors:

Dr. István Timári, email: timari.istvan@science.unideb.hu

Prof. Katalin E. Kövér, email: kover@science.unideb.hu

Abstract

Resonance assignment is a pivotal step for any NMR analysis, such as structure elucidation or investigation of protein-ligand interactions. Both ¹H-¹³C HSQC and ¹H-¹H COSY 2D experiments are invaluable for ¹H NMR assignment, by extending the high signal dispersion of ¹³C chemical shifts onto the ¹H resonances, and by providing a high amount of through-bond ¹H-¹H connectivity information, respectively. The recently introduced HSQC-CLIP-COSY method combines these two experiments, providing COSY correlations along the high-resolution ¹³C dimension with clean in-phase multiplets. However, two experiments need to be recorded to unambiguously identify COSY cross-peaks. Here, we propose novel variants of the HSQC-CLIP-COSY pulse sequence that edit cross-peak signs so that direct HSQC responses can be distinguished from COSY relay peaks, and/or the multiplicities of the ¹³C nuclei are reflected, allowing assignment of all peaks in just a single experiment. The advanced HSQC-CLIP-COSY variants have the potential to accelerate and simplify the NMR structure elucidation process of both synthetic and natural products, and to become valuable tools for high-throughput computer-assisted structure determination.

Introduction

Structure verification and elucidation are day-to-day tasks at research institutes, pharmaceutical and other chemical industries. Nuclear magnetic resonance (NMR) spectroscopy is proven to be a central analytical tool for these purposes.¹ Resonance assignment is a fundamental step of any NMR investigation surrounding characterization of molecular structure and dynamics or protein-ligand interaction. NMR resonance assignment and structure elucidation of small and medium-sized molecules typically rest on a key set of classical 2D experiments including correlation spectroscopy (COSY),^{2,3} total correlation spectroscopy (TOCSY),⁴ nuclear Overhauser effect spectroscopy (NOESY)⁵ / rotating-frame Overhauser effect spectroscopy (ROESY),⁶ heteronuclear single quantum correlation (HSQC)⁷ and heteronuclear multiple bond correlation (HMBC)⁸ methods. The growing number of regulatory requirements in, for example, drug development increases the amount of individual experiments that must be recorded, creating a demand for approaches that provide a maximum information in the shortest possible time. For instance, NOAH-type (NMR by Ordered Acquisition using ¹H-detection) experiments,⁹⁻¹² which allow sequential recording of up to four 2D spectra with only one relaxation delay employed in the combined pulse sequence, offer significant time savings compared to the conventional data recording. Also considerable progress has been made in the development of non-uniform sampling (NUS) methods¹³⁻¹⁶ for the reconstruction of multidimensional NMR spectra with high digital resolution using only a fraction of increments along the indirect dimension(s), cutting down the measurement time substantially. Furthermore, the resolution of ¹H-¹H COSY, NOESY or TOCSY experiments can be improved so that unambiguous spectral interpretation is greatly facilitated within a single spectrum. Pure shift methods enhance spectral resolution up to an order of magnitude by removing splittings from homonuclear couplings,¹⁷⁻²⁴ but these experiments often come with experimental restrictions or with extended measurement times relative to their parent experiment. An alternative strategy for reducing spectral crowding is to replace the indirect ¹H dimension with a ¹³C dimension, which features a much wider dispersion in chemical shifts. For instance, HSQC-TOCSY²⁵ and HSQC-NOESY²⁶ experiments extend the parent HSQC experiment with a TOCSY or NOESY mixing step, revealing both one-bond ¹H-¹³C correlations and subsequent ¹H-¹H connections. Multiple quantum based heteronuclear experiments, such as H2BC,^{27,28} H2OBC²⁹ and HMQC-COSY,³⁰ were also introduced for the identification of neighboring protons using both ¹H and ¹³C chemical shifts for distinction. However, an HSQC-based experiment would be preferred in many cases, as these generally achieve higher resolution in the ¹³C dimension. We recently developed the HSQC-CLIP-COSY method,³¹ which limits the ¹H-¹H correlations to those that are directly *J*-coupled, rendering the step-by-step assignment walk straightforward relative to the HSQC-TOCSY²⁵ experiment. Our method³¹ is based on the CLIP-

COSY experiment,³² which provides clean in-phase multiplets with full absorption mode lineshapes and enhanced cross-peak sensitivity relative to the standard COSY experiment for protons with small *J*-couplings. However, the original HSQC-CLIP-COSY experiment³¹ has one major limitation, namely, a supplemental HSQC spectrum needs to be recorded for unambiguous distinction between the direct ¹³C–¹H correlations and COSY-type ¹³C–¹H–¹H peaks of the HSQC-CLIP-COSY spectrum.

Here, we show how the power of the HSQC-CLIP-COSY method can be further boosted by introducing simple heteronuclear spin-echo based building blocks into the original pulse sequence. The amended variants of HSQC-CLIP-COSY experiment allow differentiation of the direct and COSY responses and provide valuable carbon multiplicity information. This makes it possible to obtain unambiguously all one-bond ¹H–¹³C correlations, ¹H–¹H COSY correlations, and ¹³C multiplicities from just a single 2D experiment (i.e., one single 2D spectrum), significantly reducing the NMR measurement and analysis time needed for structure elucidation.

Experimental section

All experiments were performed on an Avance NEO 700 (¹H: 700.25 MHz; ¹³C: 176.08 MHz) spectrometer (Bruker BioSpin GmbH, Rheinstetten, Germany) equipped with a 5 mm TCI prodigy probe. Data were processed with TopSpin 3.5 or 4.0.5 (Bruker Biospin GmbH, Karlsruhe, Germany). The pulse sequence specific experimental details are given in the corresponding figure legends.

For testing the proposed edited HSQC-CLIP-COSY heteronuclear correlation experiments, samples of 60 mg heparin-analog trisaccharide dissolved in 550 μl D₂O and 7.55 mg octapeptide (amino acid sequence: A I K L S T V G) dissolved in 550 μl d₆-DMSO were used. For the measurements the nominal temperature was 298 K.

Results and discussion

Figure 1 illustrates the general scheme of the proposed editing options of the HSQC-CLIP-COSY experiment. On the one hand, the sign of signal amplitudes in the correlation spectra can be edited according to direct HSQC correlations versus COSY peaks (route I in Figure 1). The sign encoding of the direct and relay peaks facilitates tracking of the connectivity network of protonated carbons, making the assignment of both ¹H and ¹³C resonances possible from a single spectrum without the necessity of recording an additional HSQC spectrum. On the other hand, even and odd

multiplicity of ^{13}C nuclei can be also distinguished on the basis of pertinent sign characteristics. Thus, peaks arising from CH_2 groups will appear as negative signals whereas those arising from CH and CH_3 groups will all be positive in the spectrum (route II/A). Optionally, ^{13}C multiplicity editing can be achieved by a recently proposed, alternative way,³³ where correlation peaks from CH_2 groups will appear as mirror images with reversed frequency offset and with negative amplitude in the spectra (route II/B). This latter approach changes the dispersion of the correlation peaks, reduces signal overlaps in severely crowded regions when the CH_2 correlation peaks can be reflected in a sparse spectral region. This also fully resolves potential sign interpretation ambiguities of combined (I and II) editing. The approach can be very useful in the case of peptides and also smaller proteins,³³ as will be demonstrated in the present article.




Figure 1. Schematic overview of the editing options of the HSQC-CLIP-COSY experiment. There are two major routes, which can be applied separately or simultaneously, depending on the complexity of the molecular structure investigated. On the one hand, signal amplitudes in the correlation spectra can be edited according to direct HSQC correlations versus COSY peaks. On the other hand, even and odd multiplicities of ^{13}C nuclei can be distinguished based on cross-peak signs (II/A), or by reflecting even multiplicities along F_1 (II/B).³³

The pulse scheme of the novel, edited variants of the HSQC-CLIP-COSY experiment is shown in Figure 2 and is briefly discussed in the following. Initially, standard HSQC and CLIP-COSY sequences are merged to form the HSQC-CLIP-COSY as introduced and described in a recent publication of our group.³¹ Importantly, for refocusing the antiphase proton magnetization generated in the first $^1\text{H} \rightarrow ^{13}\text{C} \rightarrow ^1\text{H}$ polarization transfer block of the sequence, a suitably positioned carbon 180° pulse is employed during the perfect-echo based COSY mixing segment. The HSQC vs. COSY response editing block of $(^1J_{\text{CH}})^{-1}$ duration (labeled with I in Figure 2) is inserted at the end of the original HSQC-CLIP-COSY sequence prior to ^1H detection. This spin-echo block allows reversing the sign of amplitude of direct (HSQC-type) peaks with respect to the relay (COSY-type) peaks due

to the $^1J_{\text{CH}}$ coupling evolution of the directly ^{13}C -coupled proton magnetization. Furthermore, to achieve the desired carbon multiplicity editing, an additional heteronuclear spin-echo block either with simultaneous ^1H and ^{13}C 180° pulses or with a modified BIRD $^{(r,X)}$ pulse applied in the center of the $(^1J_{\text{CH}})^{-1}$ echo³³ is incorporated into the pulse sequence after the t_1 -evolution period. It is worth noting that the two different types of editing – HSQC vs. COSY peaks and ^{13}C multiplicity – can be employed either separately or together, as illustrated in Figure 1. To avoid signal overlap and/or accidental cancellation of signals with opposite phases due to the editing, high digital resolution in F_1 is recommended, which can be optimally achieved by applying non-uniform sampling (NUS) without increasing the measurement time.

Regarding previously published methods, the HMQC-based H2OBC experiment²⁹ provides similar spectral information to our HSQC-CLIP-COSY method. However, the H2OBC being a constant time (CT) experiment offers only limited resolution in F_1 , while in HSQC-CLIP-COSY there is no such limit, so ultra-high ^{13}C resolution can be reached allowing separation of closely resonating overlapping signals. Moreover, in H2OBC/2BOB,²⁹ the direct and COSY cross-peaks have a 90° phase difference, which adversely affects spectral resolution in F_2 . In contrast, all peaks in HSQC-CLIP-COSY possess pure absorption lineshapes. COSY correlations are also fully in-phase, offering increased sensitivity when homonuclear splittings are small, as discussed in an earlier publication.³² Such favorable peak-shape characteristics of HSQC-CLIP-COSY also provide the best possible condition for automatic peak-picking procedures, central to computer-assisted structure elucidation.

Overall, the improved, edited variants of the HSQC-CLIP-COSY experiment allow differentiation of direct and COSY responses, and can provide useful carbon multiplicity information with only minor penalties in sensitivity from the use of more pulses, mismatch of echo-delays with respect to the pertinent $^1J(\text{XH})$ value and resonance offset effects of carbon pulses. To minimize the latter adverse effect, utilization of adiabatic,³⁴⁻³⁸ composite adiabatic,³⁹ or optimal control^{40,41} broadband ^{13}C pulses is recommended, particularly at higher magnetic fields.




Figure 2. Pulse sequence scheme for HSQC-CLIP-COSY experiments. Narrow and wide filled bars correspond to 90° and 180° pulses, respectively, with phase x unless indicated otherwise. Broadband adiabatic inversion pulses (CHIRP) and shaped refocusing pulses applied to (composite CHIRP or BURBOP-180) ^{13}C are shown as open and filled half-ellipses. Phases are $\phi_1 = x$; $\phi_2 = y$; $\phi_3 = x, -x$; $\phi_4 = (x)_2, (-x)_2$; $\phi_5 = (x)_4, (-x)_4$; $\phi_6 = (-x)_2, (-y)_2$; $\phi_7 = -y$; $\phi_8 = -x$; $\phi_9 = x, -x$; $\phi_{\text{rec}} = x, -x, -x, x$. The minimum number of transients required by phase cycling is 2. Delays are set as follows: Δ is typically set to 20 – 25 ms, allowing sufficient transfer of magnetization between protons coupled with couplings ranging from 2 up to 14 Hz, and complying the preferred perfect-echo duration for in-phase coherence transfer between coupled spins; $\tau' = 1 / ({}^1J_{\text{XH}} \times 4) - t_{\pi(x)2}$. In case of classical editing route (A) coherence order selection and echo-antiecho phase sensitive detection in the ^{13}C dimension are achieved with gradient pulses G₂ and G₆ in the ratio 80:20.1 for ^{13}C . While in route (B) coherence order selection and phase sensitive detection are achieved according to States-TPPI protocol. Purging gradient pulses G₁, G₃, G₅ are set to 17%, 31%, -17.9% of maximum gradient strength (50 G/cm), typically with 1 ms duration followed by a recovery delay of 100 μs . Weak magnetic field gradient (G₄) used under the frequency-swept CHIRP pulse is adjusted for -7.5% of maximum gradient strength.

The heteronuclear correlation-based assignment strategy utilizing the HSQC *vs.* COSY response and the classical ^{13}C multiplicity editing is illustrated on the example of a heparin-analog trisaccharide. Based on the sign of correlation peaks – that is color-coded as positive-black and negative-red in the edited HSQC-CLIP-COSY spectrum shown in Figure 3 – the directly coupled ^{13}C - ^1H (HSQC) pairs (black) and the indirectly coupled ^{13}C - ^1H - ^1H spin triples (red) can be easily and unambiguously identified, which makes a step-by-step walk along the spin system of each sugar residue possible. Note that D6, F6 and F7 CH₂-carbon correlation peaks show opposite sign pattern – red for HSQC- and black for COSY-type correlations – due to ^{13}C multiplicity editing. As a result, the ^1H - ^1H coupled spin partners can be simply traced along the rows, while walking along the columns

identifies the neighboring carbons. In Figure 3 this assignment walk is indicated by dotted and solid lines for each residue of the trisaccharide. Thus, the HSQC-CLIP-COSY experiment delivering both HSQC-type one-bond heteronuclear correlation and COSY-type homonuclear connectivity information in one single spectrum allows rapid and accurate spin system identification in small to medium-sized molecules.




Figure 3. Edited HSQC-CLIP-COSY spectrum of a heparin-analog trisaccharide.⁴² The assignment walks of D, E and F residues are labeled by colored solid and dotted lines along ^{13}C and ^1H dimension, respectively. All ring proton resonances of the trisaccharide can be unambiguously assigned by simply walking along the color-coded cross peaks of directly (black) and indirectly (red) coupled spins. For CH_2 -correlations (D6, F6, F7) the color pattern is opposite due to carbon multiplicity editing. OMe label on the spectrum indicates the HSQC peaks of O-Methyl groups. Spectrum was recorded with spectral widths of 6.93 (120.0) ppm in ^1H (^{13}C) dimensions, a relaxation delay of 1.5 s, 1024 t_1 increments using 25% NUS, 2048 data points acquired in ^1H dimension and 2 scans per increment with an overall experimental time of 16 min.

In cases when the chemical shift distribution of resonances is small, and hence spectra contain severely crowded regions of carbons with different multiplicities (e.g. $\text{H}_\beta/\text{C}_\beta$, $\text{H}_\gamma/\text{C}_\gamma$ and $\text{H}_\delta/\text{C}_\delta$ regions of oligopeptides and proteins), the carbon-multiplicity editing approach, as proposed by Sakhaii and Bermel,³³ can be utilized as an additional, powerful tool to resolve the overlapping resonances. Inserting the modified, switchable (ON/OFF) $\text{BIRD}^{(r,X)}$ pulse cluster in the multiplicity editing module (Figure 2), the sign of carbon chemical shift evolution of the CH_2 groups is reversed relative to the CH/CH_3 groups, so that the corresponding CH_2 cross-peak positions are reflected relative to

the ^{13}C carrier frequency. When the latter is chosen judiciously, this repositions the CH_2 peaks into an empty region of the spectrum, avoiding overlap and cancellation between CH_2 and the oppositely-phased CH/CH_3 peaks. Hence, improved signal dispersion can be achieved without the need of increasing digital resolution in the F_1 dimension of the spectrum. In the doubly-edited HSQC-CLIP-COSY spectrum, positive peaks (black) represent direct CH and CH_3 connectivities or COSY-type relay peaks originating from CH_2 moieties, while negative peaks (red) represent direct CH_2 correlations or relay peaks arising from CH and CH_3 moieties. In addition, applying the BIRD-type carbon multiplicity editing,³³ correlation peaks from CH_2 groups appear as mirror images in the corresponding low-field (large ^{13}C chemical shift) region of the spectrum. The real ^{13}C chemical shifts (δ_{CH_2}) of the reflected (mirror image) CH_2 peaks can be easily assessed from the values of the measured chemical shift ($\delta_{\text{CH}_2 \text{ measured}}$) and the ^{13}C carrier offset frequency ($\delta_{^{13}\text{C} \text{ offset}}$) used for the measurement with the relation $\delta_{\text{CH}_2} = 2 \times \delta_{^{13}\text{C} \text{ offset}} - \delta_{\text{CH}_2 \text{ measured}}$.

The proposed doubly-edited variant of HSQC-CLIP-COSY experiment relying on the frequency reversing multiplicity editing block provides an efficient tool for the identification of amino acid type in oligopeptides. The characteristic HSQC-CLIP-COSY peak patterns of α -amino acids with different spin topology in their side chains are shown in Figure 4 and Figures S1-S20 with illustration of the assignment walk along the spin system. Based on the peak pattern observable in the doubly-edited HSQC-CLIP-COSY spectrum, four types of amino acid topologies can be easily assigned and distinguished. Namely, Gly with CH_2 in α position gives only two peaks, the direct ($\text{C}\alpha\text{-H}\alpha$) and relay ($\text{C}\alpha\text{-H}\alpha\text{-HN}$) peaks at ‘mirror image’ carbon chemical shift in the spectrum (Figures 4/a and S1). The second group of amino acids with isolated CH_2 in β position (such as Asp, Asn, Cys, Ser, Phe, Tyr, Trp, His) can be easily identified following the assignment walk shown in Figures 4/b and S2-S9. Starting from the direct ($\text{C}\alpha\text{-H}\alpha$) peak and ‘walking down’ along the column, the neighboring $\text{C}\beta(\text{CH}_2)$ can be found in the large carbon chemical shift (mirror image) region of the spectrum. Since $\text{CH}_2\text{-}\beta$ protons have only $\text{H}\alpha$ as coupling partner, the row of $\text{C}\beta$ contains only one relay peak $\text{C}\beta\text{-H}\beta\text{-H}\alpha$. Alternatively, the amino acids with $\text{C}\beta(\text{CH}_2)$ followed by protonated $\text{C}\gamma$ in their side chain (such as Pro, Leu, Met, Glu, Gln, Arg, Lys) show additional relay peaks $\text{C}\beta\text{-H}\beta\text{-H}\gamma$ in the row of their $\text{C}\beta$ carbon (Figures 4/c and S10-S16). The assignment walk along the rest of the side chain – as demonstrated in Figures 4/c and S10-S16 – allows differentiation of the amino acids with different side chain topology. In the last group of amino acids (Ala, Val, Thr, Ile) with CH/CH_3 in β position the assignment walk starting from the direct ($\text{C}\alpha\text{-H}\alpha$) peak goes upward along the column, locating the adjoining $\text{C}\beta$ in the high field (aliphatic) region of the spectrum (Figures 4/d and S17-S20). Again, completing the assignment walk along the side chain, the specific amino acids of the group can be easily identified.




Figure 4. Characteristic doubly-edited HSQC-CLIP-COSY peak patterns of representative α -amino acids with different spin topology in their side chains. a) Glycine (containing only C_{α}); b) Phenylalanine (containing isolated CH_2 in β position); c) Proline (containing CH_2 in β position followed by protonated C_{γ}); d) Valine (containing CH/CH_3 in β position). Arrows on the spectra indicate the assignment walk along the spin systems.

The utility of this peak pattern-based assignment protocol is demonstrated on the example of an octapeptide containing at least one specific amino acid from each amino acid type sub-group. Accomplishing the assignment walk in the doubly-edited HSQC-CLIP-COSY spectrum as shown in Figure 5 – where different colors are used for coding the different amino acid types – a complete and unambiguous assignment of all 1H and protonated ^{13}C resonances of individual amino acids could be attained. The mandatory sequential assignment of residues within bio-oligomers can next be accomplished by 2D ROESY, utilizing the shortrange 1H – 1H distances between neighboring residues, or by 2D HMBC experiment, observing the interresidue three-bond 1H – ^{13}C connectivities as shown in Figures S21-S22.

Taken together, these examples demonstrate how the enhanced spectral information content of the edited HSQC-CLIP-COSY provides an ideal tool for quick and unambiguous assignment of 1H and protonated ^{13}C resonances of small to medium-sized molecules, such as (oligo)saccharides and peptides.




Figure 5. Doubly-edited HSQC-CLIP-COSY spectrum of an oligopeptide (amino acid sequence: A I K L S T V G), which contains at least one specific amino acid from each amino acid type subgroup. The different colors are used for coding the different amino acid types as described in the text. Starting from the direct C_α - H_α peaks (region is indicated by black box), all amino acid residues can be easily identified. The spectrum was recorded with spectral widths of 10.2 (190.0) ppm in ^1H (^{13}C) dimensions, a relaxation delay of 1.7 s, 1024 t_1 increments, 2048 total data points acquired in the ^1H dimension, 4 scans per increment.

Conclusions

Edited HSQC-CLIP-COSY experiments have been developed for making the ^1H and ^{13}C NMR assignment of small to medium-sized molecules simpler and faster. Sign editing of direct HSQC responses vs. CLIP-COSY peaks provides a straightforward approach for tracking down the

backbone of protonated ^{13}C nuclei step-by-step. This edited HSQC-CLIP-COSY spectrum offers the information content of HSQC and COSY spectra integrated in one data set, obviating the need for recording of separate experiments. The information content of an HSQC-CLIP-COSY spectrum can be further boosted by introduction of carbon multiplicity-editing. To simplify overcrowded spectral regions, the sign of carbon frequency of CH_2 signals can be also reversed utilizing the recently proposed BIRD^(r,X) scheme for multiplicity editing. We believe that the proposed edited HSQC-CLIP-COSY experiments, providing high-quality, sign- and/or frequency-coded, well-resolved and pure absorption spectra, hold great promise to significantly accelerate and simplify NMR assignment of bio-oligomers and other types of synthetic molecules or natural products. Due to the high information content, excellent resolution and line shape characteristics of the resulting spectra, the edited HSQC-CLIP-COSY experiments have the potential to be valuable and indispensable for computer-assisted structure elucidation.

Associated Content

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/XXX>.

Additional NMR spectra about the characteristic HSQC-CLIP-COSY peak pattern of α -amino acids, sequential assignment of the trisaccharide and the peptide, pulse sequence codes for Bruker spectrometers (PDF).

Notes

The authors declare no competing financial interest.

Acknowledgements

This research was supported by the National Research, Development and Innovation Office of Hungary (grant numbers: NKFI/OTKA NN 128368 (to T.G. and K.E.K.) and NKFI/OTKA PD 135034 (to I.T.)) and co-financed by the European Regional Development Fund (projects GINOP-2.3.3-15-2016-00004 and GINOP-2.3.2-15-2016-00008). The research of I.T. was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (BO/00372/20/7) and the ÚNKP-20-5-DE-262 New National Excellence Program of the Ministry for Innovation and

Technology from the source of the National Research, Development and Innovation Fund. B.L. acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG LU 835/13-1) and the HGF program BIFTM (IBG-4, 47.02.04).

The authors thank Prof. Anikó Borbás, Dr. Mihály Herczeg and Prof. Gábor Tóth for providing the heparin-analog trisaccharide and the oligopeptide, respectively.

Keywords: CLIP-COSY, HSQC, NMR spectroscopy, peptide, structure elucidation

References

- (1) Claridge, T. D. W. In *High-Resolution NMR Techniques in Organic Chemistry (Third Edition)*, Claridge, T. D. W., Ed.; Elsevier: Boston, 2016, pp 1-541.
- (2) Aue, W. P.; Bartholdi, E.; Ernst, R. R. Two-dimensional spectroscopy. Application to nuclear magnetic resonance *J. Chem. Phys.* **1976**, *64*, 2229-2246.
- (3) Piantini, U.; Sorensen, O. W.; Ernst, R. R. Multiple quantum filters for elucidating NMR coupling networks *J. Am. Chem. Soc.* **1982**, *104*, 6800-6801.
- (4) Braunschweiler, L.; Ernst, R. R. Coherence transfer by isotropic mixing: Application to proton correlation spectroscopy *J. Magn. Reson.* **1983**, *53*, 521-528.
- (5) Jeener, J.; Meier, B. H.; Bachmann, P.; Ernst, R. R. Investigation of exchange processes by two-dimensional NMR spectroscopy *J. Chem. Phys.* **1979**, *71*, 4546-4553.
- (6) Bothner-By, A. A.; Stephens, R. L.; Lee, J.; Warren, C. D.; Jeanloz, R. W. Structure determination of a tetrasaccharide: transient nuclear Overhauser effects in the rotating frame *J. Am. Chem. Soc.* **1984**, *106*, 811-813.
- (7) Bodenhausen, G.; Ruben, D. J. Natural abundance nitrogen-15 NMR by enhanced heteronuclear spectroscopy *Chem. Phys. Lett.* **1980**, *69*, 185-189.
- (8) Bax, A.; Summers, M. F. Proton and carbon-13 assignments from sensitivity-enhanced detection of heteronuclear multiple-bond connectivity by 2D multiple quantum NMR *J. Am. Chem. Soc.* **1986**, *108*, 2093-2094.
- (9) Kupče, Ě.; Claridge, T. D. W. NOAH: NMR Supersequences for Small Molecule Analysis and Structure Elucidation *Angew. Chem. Int. Ed.* **2017**, *56*, 11779-11783.
- (10) Kupče, Ě.; Claridge, T. D. W. Molecular structure from a single NMR supersequence *Chem. Commun.* **2018**, *54*, 7139-7142.
- (11) Motiram-Corral, K.; Pérez-Trujillo, M.; Nolis, P.; Parella, T. Implementing one-shot multiple-FID acquisition into homonuclear and heteronuclear NMR experiments *Chem. Commun.* **2018**, *54*, 13507-13510.
- (12) Nagy, T. M.; Gyöngyösi, T.; Kövér, K. E.; Sørensen, O. W. BANGO SEA XLOC/HMBC-H2OBC: complete heteronuclear correlation within minutes from one NMR pulse sequence *Chem. Commun.* **2019**, *55*, 12208-12211.
- (13) Kazimierczuk, K.; Orekhov, V. Y. Accelerated NMR Spectroscopy by Using Compressed Sensing *Angew. Chem. Int. Ed.* **2011**, *50*, 5556-5559.
- (14) Orekhov, V. Y.; Jaravine, V. A. Analysis of non-uniformly sampled spectra with multi-dimensional decomposition *Prog. Nucl. Magn. Reson. Spectrosc.* **2011**, *59*, 271-292.
- (15) Mobli, M.; Hoch, J. C. Nonuniform sampling and non-Fourier signal processing methods in multidimensional NMR *Prog. Nucl. Magn. Reson. Spectrosc.* **2014**, *83*, 21-41.

- (16) Li, D.; Hansen, A. L.; Bruschiweiler-Li, L.; Brüschiweiler, R. Non-Uniform and Absolute Minimal Sampling for High-Throughput Multidimensional NMR Applications *Chem. Eur. J.* **2018**, *24*, 11535-11544.
- (17) Aguilar, J. A.; Faulkner, S.; Nilsson, M.; Morris, G. A. Pure shift ¹H NMR: a resolution of the resolution problem? *Angew. Chem. Int. Ed.* **2010**, *49*, 3901-3903.
- (18) Morris, G. A.; Aguilar, J. A.; Evans, R.; Haiber, S.; Nilsson, M. True chemical shift correlation maps: a TOCSY experiment with pure shifts in both dimensions *J. Am. Chem. Soc.* **2010**, *132*, 12770-12772.
- (19) Aguilar, J. A.; Colbourne, A. A.; Cassani, J.; Nilsson, M.; Morris, G. A. Decoupling two-dimensional NMR spectroscopy in both dimensions: pure shift NOESY and COSY *Angew. Chem. Int. Ed.* **2012**, *51*, 6460-6463.
- (20) Meyer, N. H.; Zangger, K. Simplifying proton NMR spectra by instant homonuclear broadband decoupling *Angew. Chem. Int. Ed.* **2013**, *52*, 7143-7146.
- (21) Foroozandeh, M.; Adams, R. W.; Nilsson, M.; Morris, G. A. Ultrahigh-Resolution Total Correlation NMR Spectroscopy *J. Am. Chem. Soc.* **2014**, *136*, 11867-11869.
- (22) Castañar, L.; Parella, T. Broadband ¹H homodecoupled NMR experiments: recent developments, methods and applications *Magn. Reson. Chem.* **2015**, *53*, 399-426.
- (23) Zangger, K. Pure shift NMR *Prog. Nucl. Magn. Reson. Spectrosc.* **2015**, *86-87*, 1-20.
- (24) Kakita, V. M. R.; Hosur, R. V. Non-Uniform-Sampling Ultrahigh Resolution TOCSY NMR: Analysis of Complex Mixtures at Microgram Levels *ChemPhysChem* **2016**, *17*, 2304-2308.
- (25) Lerner, L.; Bax, A. Sensitivity-enhanced two-dimensional heteronuclear relayed coherence transfer NMR spectroscopy *J. Magn. Reson.* **1986**, *69*, 375-380.
- (26) Majumdar, A.; Zuiderweg, E. R. P. Improved ¹³C-Resolved HSQC-NOESY Spectra in H₂O, Using Pulsed Field Gradients *J. Magn. Reson., Ser B* **1993**, *102*, 242-244.
- (27) Nyberg, N. T.; Duus, J. Ø.; Sørensen, O. W. Heteronuclear Two-Bond Correlation: Suppressing Heteronuclear Three-Bond or Higher NMR Correlations while Enhancing Two-Bond Correlations Even for Vanishing 2J_{CH} *J. Am. Chem. Soc.* **2005**, *127*, 6154-6155.
- (28) Nyberg, N. T.; Duus, J. Ø.; Sørensen, O. W. Editing of H₂BC NMR spectra *Magn. Reson. Chem.* **2005**, *43*, 971-974.
- (29) Kupče, Ě.; Sørensen, O. W. 2BOB – extracting an H₂BC and an HSQC-type spectrum from the same data set, and H₂OBC – a fast experiment delineating the protonated ¹³C backbone *Magn. Reson. Chem.* **2017**, *55*, 515-518.
- (30) Hu, K.; Westler, W. M.; Markley, J. L. Two-dimensional concurrent HMQC-COSY as an approach for small molecule chemical shift assignment and compound identification *J. Biomol. NMR* **2011**, *49*, 291-296.
- (31) Gyöngyösi, T.; Timári, I.; Haller, J.; Koos, M. R. M.; Luy, B.; Kövér, K. E. Boosting the NMR Assignment of Carbohydrates with Clean In-Phase Correlation Experiments *ChemPlusChem* **2018**, *83*, 53-60.
- (32) Koos, M. R. M.; Kummerlöwe, G.; Kaltschnee, L.; Thiele, C. M.; Luy, B. CLIP-COSY: A Clean In-Phase Experiment for the Rapid Acquisition of COSY-type Correlations *Angew. Chem. Int. Ed.* **2016**, *55*, 7655-7659.
- (33) Sakhaei, P.; Bermel, W. A different approach to multiplicity-edited heteronuclear single quantum correlation spectroscopy *J. Magn. Reson.* **2015**, *259*, 82-86.
- (34) Böhlen, J.-M.; Rey, M.; Bodenhausen, G. Refocusing with chirped pulses for broadband excitation without phase dispersion *J. Magn. Reson.* **1989**, *84*, 191-197.
- (35) Böhlen, J.-M.; Burghardt, I.; Rey, M.; Bodenhausen, G. Frequency-modulated “Chirp” pulses for broadband inversion recovery in magnetic resonance *J. Magn. Reson.* **1990**, *90*, 183-191.
- (36) Ogura, K.; Terasawa, H.; Inagaki, F. Fully¹³C-Refocused Multidimensional¹³C-Edited Pulse Schemes Using Broadband Shaped Inversion and Refocusing Pulses *J. Magn. Reson., Ser B* **1996**, *112*, 63-68.
- (37) Kupče, Ě.; Freeman, R. Compensation for Spin–Spin Coupling Effects during Adiabatic Pulses *J. Magn. Reson.* **1997**, *127*, 36-48.

- (38) Zwahlen, C.; Legault, P.; Vincent, S. J. F.; Greenblatt, J.; Konrat, R.; Kay, L. E. Methods for Measurement of Intermolecular NOEs by Multinuclear NMR Spectroscopy: Application to a Bacteriophage λ N-Peptide/boxB RNA Complex *J. Am. Chem. Soc.* **1997**, *119*, 6711-6721.
- (39) Degraaf, R. A.; Luo, Y.; Terpstra, M.; Merkle, H.; Garwood, M. A New Localization Method Using an Adiabatic Pulse, BIR-4 *J. Magn. Reson., Ser B* **1995**, *106*, 245-252.
- (40) Kobzar, K.; Ehni, S.; Skinner, T. E.; Glaser, S. J.; Luy, B. Exploring the limits of broadband 90° and 180° universal rotation pulses *J. Magn. Reson.* **2012**, *225*, 142-160.
- (41) Skinner, T. E.; Gershenson, N. I.; Nimbalkar, M.; Bermel, W.; Luy, B.; Glaser, S. J. New strategies for designing robust universal rotation pulses: Application to broadband refocusing at low power *J. Magn. Reson.* **2012**, *216*, 78-87.
- (42) Lázár, L.; Mező, E.; Herczeg, M.; Lipták, A.; Antus, S.; Borbás, A. Synthesis of the non-reducing end trisaccharide of the antithrombin-binding domain of heparin and its bioisosteric sulfonic acid analogues *Tetrahedron* **2012**, *68*, 7386-7399.

For Table of Contents Only

