



Linking personality and health to use of space in the gray partridge: perspectives for management.

Clément Harmange, Vincent Bretagnolle, Olivier Pays, Mathieu Sarasa,
Jérôme Moreau

► To cite this version:

Clément Harmange, Vincent Bretagnolle, Olivier Pays, Mathieu Sarasa, Jérôme Moreau. Linking personality and health to use of space in the gray partridge: perspectives for management.. European Journal of Wildlife Research, 2021, 67 (5), pp.81. 10.1007/s10344-021-01526-x . hal-03328949

HAL Id: hal-03328949

<https://hal.science/hal-03328949>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Linking Personality and Health to Use of Space in the Gray Partridge: Perspectives for Management

Clément Harmange^{1,2,3}, Vincent Bretagnolle^{2,4}, Olivier Pays^{1,3}, Mathieu Sarasa^{5,6}, Jérôme Moreau^{2,7}

¹LETG-Angers, UMR 6554 CNRS, Université d'Angers, Angers, France.

²Centre d'Etudes Biologiques de Chizé, UMR 7372, CNRS and La Rochelle Université, F-79360 Beauvoir-sur-Niort, France.

³REHABS International Research Laboratory, CNRS-Université Lyon 1-Nelson Mandela University, George Campus, Madiba drive 6531 George, South Africa.

⁴LTSE « Zone Atelier Plaine & Val de Sèvre », CNRS, 79360, Beauvoir sur Niort, France.

⁵BEOPS, 1 Esplanade Compans Caffarelli, 31000 Toulouse, France.

⁶Fédération Nationale des Chasseurs, 13 rue du Général Leclerc, 92136 Issy-les-Moulineaux cedex, France.

⁷UMR CNRS 6282 Biogéosciences, Université Bourgogne Franche-Comté, 6 Boulevard Gabriel, 21000 Dijon, France.

Corresponding author:

Clément Harmange, LETG-Angers, UMR 6554 CNRS, Université d'Angers, 49000 Angers, France, c.harmange.pro@gmail.com, ORCID: 0000-0001-5207-021X

ORCID Vincent Bretagnolle: 0000-0002-2320-7755

ORCID Olivier Pays: 0000-0001-8268-1804

ORCID Mathieu Sarasa: 0000-0001-9067-7522

ORCID Jérôme Moreau: 0000-0001-9984-0998

ABSTRACT

Behavioral adjustments are of critical importance for the survival of animals when exposed to risk and resource variations in their environment. Moving to safer habitats allows prey to reduce the risk of mortality triggered by predation. This response has been well studied at the population level to assist in identifying key habitat components for wildlife conservation. However, while individuals may differ in their response to risk and thus in their survival expectancy, the determinants of individual adjustments in space and habitat use are largely unknown in management and conservation studies. Using GPS monitoring, we investigated the relationship between individual traits (morphometry, personality, health status) and adjustments in use of space relative to

risky features (hedgerows, roads/tracks, woodlands, built-up areas) with regard to predation risk. We used the gray partridge (*Perdix perdix*) as a study model (captive-reared birds released into the farmlands). During the diel cycle, predation risk mainly occurs by carnivores at night whereas human hunting pressure occurs during the day. We found that personality and health status had an effect on adjustments in use of space as response to risky features. At night, when carnivore activity is expected to be high, bolder individuals were farther from hedgerows relative to the day. Similarly, individuals in good health were located farther from hedgerows, roads/tracks, and woodland at night relative to the day. We discuss the need for better consideration of individual traits in management and conservation programs, with a particular focus on recovery programs that rely on translocation of captive-reared individuals.

KEYWORDS: Behavioral adjustment, Gray partridge, Health status, Personality, Space use, Wildlife management

DECLARATIONS

Funding: This work was supported by the Fédération Nationale des Chasseurs (grant number FNC-PSN-PR15-2014) and the French National Center for Scientific Research (CNRS)

Conflicts of interest/Competing interests: The authors have no conflicts of interest to declare that are relevant to the content of this article.

Availability of data and material: The authors agree to archive the data associated with this manuscript should the manuscript be accepted. The following statement will be added in the manuscript upon acceptance “The datasets analysed during the current study are available in the Open Science Framework repository, <https://dx.doi.org/10.17605/OSF.IO/G6PQV>”.

Code availability: The authors agree to archive the code associated with this manuscript should the manuscript be accepted. The code will be added to the same repository as the data.

Ethics approval: Field studies did not involve endangered or protected species. Permission to handle the study birds was given by the authority Préfecture Départementale des Deux-Sèvres (Number 2017/1). All applicable French guidelines for the ethical use of animals in research were followed.

ACKNOWLEDGMENTS

We owe special thanks for the valuable assistance received from T. Rinaud, A. Villers, M. Souchet, C. Palma, and N. Chabaud in the release, monitoring and recapture of the partridges, and to T. Rinaud, M. Crépin, and F.

61 Derouin Tochon for preparation and analyses of blood smears and partridge coloration. We thank all the
62 fieldworkers who contributed to data collection, especially R. Bonnet, P. Lagrange, J-B. Perrotin, G. Lemonnier,
63 J. Léauté, P. Harmange, T. Joubin, and M. Saillard. We also thank N. Aebischer, E. Bro, D. Fortin, and R.
64 Julliard for their valuable comments on the manuscript. Finally, we thank the Fédération Départementale 79 and
65 the Communal Hunting Society of Marigny for their support. This work was supported by the Fédération
66 Nationale des Chasseurs (grant number FNC-PSN-PR15-2014) and the French National Center for Scientific
67 Research (CNRS).

INTRODUCTION

Animals adjust their behavior in response to the spatial and temporal variations in resource availability and predation risk (Lima and Dill 1990; Johnson et al. 1992; Lima and Bednekoff 1999). Adjustments in space and habitat use are among the most commonly studied responses that allow prey to reduce their immediate mortality risk from predators, by moving toward safer areas or habitats (Laundré et al. 2001; Fortin et al. 2005; Tolon et al. 2009; Lone et al. 2015). However, these responses may become particularly complex in human-dominated landscapes, where human disturbance may conflict with risks imposed by non-human predators (Lone et al. 2014, 2017; Bonnot et al. 2020). For example, some prey living in heterogeneous landscapes, such as the European rabbit (*Oryctolagus cuniculus*) and the roe deer (*Capreolus capreolus*), shift their behavior patterns during the day and night in response to variation in risk. They use open habitats that offer better detection and escape from nocturnal carnivores at night, and shift to the opposite more vegetated habitats during the day, which provide shelter from human disturbance and/or diurnal predators (Moreno et al. 1996; Bonnot et al. 2014; Lone et al. 2017). Similar patterns have also been found in gamebirds, such as the gray partridge (Harmange et al. 2021).

Individuals within a population may however show significant variation in their perception of, and response to dangerous situations, depending on their phenotypes (Sih and Del Giudice 2012; Bonnot et al. 2014). This may result in between-individual differences in survival (Ciuti et al. 2012; Lone et al. 2015). However, little is known about the between-individual differences in how space is used and what determines these differences. Individual traits may influence the movement-related decision-making of individuals or their ability to move, thus affecting the use of space by several mechanisms (Nathan et al. 2008). First, the internal state of animals (i.e., the physiological and neurological states that affect motivation and readiness to move, Nathan et al. 2008) depends on personality traits that influence their propensity to take risks, and then their fitness (Both et al. 2005; Réale et al. 2007; Smith and Blumstein 2008; Sih et al. 2012). Personality has been defined as correlated behavioral traits within individuals, usually related to boldness, aggressiveness, activity and exploratory tendency, and is commonly expressed as a bold/shy gradient (Sih and Del Giudice 2012). Bolder individuals are often more aggressive and often engage in risk-taking behaviors, such as exploring novel environments or returning to disturbed areas faster than shy individuals (Verbeek et al. 1994; Van Oers et al. 2004a, b), thereby increasing their mortality risk (Van Oers et al. 2004a; Smith and Blumstein 2008). Secondly, morphometric traits may constrain motion capacity (i.e., the biomechanical properties that affect the ability to execute movement, Nathan et al. 2008). For instance, increased wing loading (or body mass) in birds impairs flight

abilities such as maneuverability (Dietz et al. 2007; van den Hout et al. 2010), as well as take-off velocity and angle (Kullberg et al. 1996), which alters escape behavior (Gosler et al. 1995; Lindström et al. 2003; Møller 2015). Thirdly, health status may interfere with navigation capacity (i.e., the ability to orient and select where and/or when to move, Nathan et al. 2008). Illness and increased allocation of energy to immune function in infected individuals, may reduce the locomotion abilities (Moore 2002; Lindström et al. 2003). Despite calls to integrate such individual variability into conservation and management schemes (Watters and Meehan 2007; Smith and Blumstein 2008; Caro and Sherman 2012; Berger-Tal and Saltz 2014), this has been seldom incorporated to date (see Merrick and Koprowski 2017).

It is of particular interest to address these issues in game species, for which management strategies often rely on the translocation of captive-reared animals (Griffith et al. 1989; Laikre et al. 2006; Sokos et al. 2008; Champagnon et al. 2012a). Animals raised in captivity usually show low survival expectancy after release, which is mainly ascribed to predation. Indeed, mortality rates one month after release can be as high as 81% in *Phasianus colchicus* (Hessler et al. 1970), 63-94% in *Alectoris rufa* (Gortázar et al. 2000), 63-95% in *Alectoris graeca* (Dessi-Fulgheri et al. 2001), or 73-90% in *Perdix perdix* (Rymesova et al. 2013). The captive-reared origin of individuals has been viewed as a major cause of failures in restoring self-sustainable populations in game species (Griffith et al. 1989; Sokos et al. 2008), because such animals often undergo multiple alterations in traits related to anti-predator behavior, such as morphology and/or physiology, which affect their ability to escape from predators (Putaala and Hissa 1995; Putaala et al. 1997; van Heezik et al. 1999; Rantanen et al. 2010; Champagnon et al. 2012b).

The gray partridge (*Perdix perdix*) is an iconic gamebird living in farmland landscape in Europe. However, agricultural intensification has resulted in reduced reproductive success for this species (Potts 1986; Kuijper et al. 2009; Gée et al. 2018). Populations have now reached such a low abundance in many areas and even gone locally extinct (Aebischer and Kavanagh 1997; Chamberlain and Fuller 2000; Comolet-Tirman et al. 2015; Harmange et al. 2019), that population persistence relies on releasing millions of captive-reared birds per year (e.g. 2 million individuals per year in France in Bro 2016; see also Buner et al. 2011; Bro and Crosnier 2012). In most cases, these programs have failed to help population recovery (Bro and Mayot 2006; Buner 2009; Jenny 2015; Harmange et al. 2019). Increased predation and shooting pressure have been identified as two of the main proximal causes for the decline of partridges over the last decades (Kuijper et al. 2009; Aebischer and Ewald 2012). However, these act in contrasting ways: predation mainly occurs at night on birds located closer to roads or tracks, while birds that are killed by hunters during the day are located in areas farther from hedgerows (i.e.,

in more open areas) during daytime (Harmange et al. 2021). Such contrasting pressures should result in a partitioning in the use of space between day and night (Moreno et al. 1996; Monterroso et al. 2013, 2014; Lone et al. 2014, 2017), but whether, and how individual traits may affect behavioral adjustments remains unknown in the context of management (see Merrick and Koprowski 2017).

In this study, we investigated the relationship between individual traits (morphometry, personality, and health status) and individual response to risk across the diel cycle. We did this by monitoring released partridges equipped with GPS devices, allowing to correlate individual traits to use of space. Although naïve at release, we already showed that captive-reared partridges are able to adjust to some extent their use of space in ways consistent with a reduction of risk from nocturnal carnivores at night, and of human-induced disturbance and/or raptor predation risk during daytime (Harmange et al. 2021). In particular, landscape features such as woodlands and built-up areas, which are predator reservoirs, and hedgerows and roads, which are predator corridors (Pereira and Rodríguez 2010; Krauze-Gryz et al. 2012; Bischof et al. 2019; Pita et al. 2020), present a high risk to partridges. We therefore assume that distance to risky landscape features may be a reasonable proxy of risk-taking behavior, and used these distances to estimate to what extent individual partridges adjust their use of space in response to the spatial variation in risks between day and night. We thus contrasted the average distance to risk-related features between the night and day by calculating difference in the mean nocturnal less the mean diurnal distances (following Harmange et al. 2021). A positive difference would indicate high levels of behavioral adjustment, that is avoidance of features with high risk at night, and greater proximity to shelter during daytime, in response to human disturbance and diurnal birds of prey (Moreno et al. 1996; Lone et al. 2017), or food-rich habitats (e.g. roadside habitats: Hopwood 2008; von der Lippe et al. 2013). We predicted that bolder individuals would adjust better to such contrasting environmental pressures because of their capacity for rapid exploration (Verbeek et al. 1994; Van Oers et al. 2004a; Fucikova et al. 2009), their fast decision-making ability in a risky situation (Burns and Rodd 2008; Sih and Del Giudice 2012), their ability to return faster to a place after a disturbance (Van Oers et al. 2004a, b; Quinn and Cresswell 2005), and their ability to adopt routine behaviors (Groothuis and Carere 2005). Boldness was measured through behaviors indicative of birds fearfulness when handled (Edelaar et al. 2012; Garcia-Longoria et al. 2014). Because high investment in the immune function is often associated with reduction in the locomotion abilities, birds with low health conditions were expected to be less mobile and thus to adjust their use of space less than healthy birds. Finally, individuals with low wing loading were expected to adjust better because of their better flying ability. As the gray partridge

is a gregarious species in winter (Beani and Dessi-Fulgheri 1986), we did not expect a strong difference in spatial adjustment between sexes.

MATERIAL AND METHODS

Study area

The study was conducted on the Long-Term Socio-Ecological Research (LTSER) platform “Zone Atelier Plaine & Val de Sèvre”, *Département des Deux-Sèvres*, in Western France. Study area covers 435 km², which is characterized by intensive cereal production. Land-cover use has been surveyed annually since 1994, and is represented by winter cereals (41.5%, mean value calculated over the years 2009–2016), sunflower (10.4%), corn (9.6%), rape (8.3%), meadows (13.5%), woodlands (2.9%) and built-up areas (9.8%) (Bretagnolle et al. 2018). The densities of hedgerows and roads/tracks are respectively 34 m/ha and 50 m/ha. In the study area, the main predators of partridges are mammalian carnivores including the red fox *Vulpes vulpes*, mustelids and the domestic cat *Felis catus*, and to a lesser extent, raptors mainly represented by the hen harrier *Circus cyaneus*.

The population density of gray partridge was very low during the years in which the study was conducted (Harmange et al. 2019). Important annual releases of captive-reared birds are carried out by hunter associations (around 250 birds per commune, see Harmange et al. 2019). Hunting activity occurs twice a week on Thursdays and Sundays in the LTSER area (Casas et al. 2009), during daylight hours (from 1 hour before sunrise to 1 hour after sunset). While the hunting season for partridges starts in mid-September and ends in mid-November, disturbance from hunting extends for longer because of shooting of other game species such as the brown hare (*Lepus europaeus*), and the common pheasant (*Phasianus colchicus*). Although shooting of the gray partridge had been suspended for three years at the release site, shooting of the red-legged partridge and other game species was still allowed.

GPS Monitoring

Five-month old captive-reared gray partridges were released into the area of the Communal Hunting Society of Marigny. Released birds were raised in a commercial game farm following a standard game rearing method (Sokos et al. 2008): the rearing farm receives 1-day-old chicks from an egg production farm (around 30 generations bred in captivity). Chicks are then artificially reared using brooders, without parents. Three sessions of releases were carried out (mid-December 2016, mid-September 2017, and late October 2017; see Table 1), following the same protocol: partridges were transported by car from the rearing farm to the release sites

(transportation time of ca. 40 minutes), in cardboard boxes of 8 individuals, separating the males and females. Flocks of 12–16 individuals were then released (with the same proportion of each sex), 6–8 of whom were equipped with a GPS device. The birds were left in acclimation pens of approximately 5 m² for five days. Prior to placement in acclimation pens, birds were weighed using a spring scale (Pesola 500 g, accuracy: ± 5 g), measured (wing length, using a ruler, accuracy: ± 1 mm), sexed, ringed (with colored and numbered metal gamebird rings), and fitted using GPS loggers (CatLog-S2, Catnip Technologies Ltd, Hong Kong). These loggers were placed on the breast of the bird using a necklace. Other birds were fitted using better performing but more expensive GPS devices allowing for remote data transmission (5 units of UvA-BiTS GPS-transmitters, www.uva-bits.nl Bouten et al. 2013 in 2016 ; and 2 units of Milsar's GsmRadioTag-S9, www.milsar.com, in 2017). These GPS transmitters were fitted onto the back using a Teflon harness (Table 1). The average handling time was 3–4 minutes per bird. The GPS loggers (CatLog-S2) were turned on at release and locations recorded as one fix every 15 minutes during daytime, and one fix per hour at night. GPS transmitters were initially programmed to record one fix per 10 minutes during daytime and one fix every 30 minutes at night, and then remotely reprogrammed to a finer temporal resolution (up to one fix every five minutes during daytime and one fix every 15 minutes at night), when the battery charge allowed. Permission to handle the study birds was given by the authority Préfecture Départementale des Deux-Sèvres (Number 2017/1). All applicable French guidelines for the ethical use of animals in research were followed.

Data Collection

The birds were recaptured to allow for retrieval of the GPS loggers (CatLog-S2) for downloading of data (Table 1). We used cages (height 30 * width 40 * depth 30 cm) positioned near release sites and close to locations where recent partridge presence was reported by stakeholders or fieldworkers. Recaptures began one month after release and were carried out monthly for 3–5 days. The recaptured birds were handled and measured directly at the capture site (no transport). Once the assessment of the measures described below was performed, birds were refitted with a GPS device with a fully charged battery, and released. The total handling time at recapture was 12–13 minutes per bird.

Individual Traits

Morphometric traits were measured again on individuals recaptured in the cages (n = 16). The ratio of body mass to wing length was determined as a proxy for *wing loading*. Then, we used the handling of the birds as an

experimental procedure allowing to measure their reaction to a stressful or risky situation, by assessing the following seven personality traits indicative of birds fearfulness (Møller et al. 2011; Edelaar et al. 2012; Garcia-Longoria et al. 2014): *flight initiation distance (FID)*, *wriggle score*, *feather loss*, *distress call*, *tonic immobility*, *alarm call* and *escape flight distance* (see Online Resource 1, Table S1 for details). All traits were assessed by the same operator to standardize the recapture protocol and limit any bias in measurements.

Finally, two proxies of health status were measured. The white cell composition (i.e., lymphocytes, monocytes, eosinophils, heterophils) was assessed from blood smears (see Online Resource 2 for protocol details; see also Hõrak et al. 2004; Biard et al. 2015). The red coloration of the carotenoid-based red skin ornament located behind the eye of the gray partridge was analyzed from photographs (see Online Resource 3 for protocol details). Carotenoids are involved in a trade-off between investment in ornamental signaling color and immune function (Møller et al. 2000; Hõrak et al. 2001). Coloration of the carotenoid-based ornaments has thus been described as a reliable proxy of health status (Zuk et al. 1990; Hõrak et al. 2001, 2004; Svobodová et al. 2013). Infected individuals are thus expected to have a less intensive red color because of an increased allocation of carotenoids to the immune response (see Svobodová et al. 2013 for example in gray partridge).

Statistical Analyses

Computation of personality and health status variables

Two principal component analyses (PCA) were performed from the sixteen recaptured individuals: one on the personality traits (flight initiation distance, wriggle score, feather loss, distress call, tonic immobility, alarm call and escape flight distance) and a second on the health status variables (lymphocytes, monocytes, eosinophils, heterophils, and red coloration), to obtain synthetic variables (i.e., the first two axes of each PCA). As only one measure of the health status variables was available for each bird, the data did not account for intra-individual variability. A single measure of personality was available for birds that were recaptured only once (4 of 16 individuals). Each personality trait was thus averaged prior to PCA for those birds that were caught several times.

Adjustment in use of space

As GPS data often contains erroneous or inaccurate fixes (see Lewis et al. 2007; Bjørneraas et al. 2010), a filter was applied to the data before analysis, that resulted in a reduction of approximately 4% of the dataset and a mean GPS location error of 15 m after filtering (see Harmange et al. 2021). We calculated the distances between

each GPS fix (mean number of fixes per bird = 3322; min = 339; max = 11346) and the nearest hedgerow, road/track, woodland, and building for the sixteen individuals. Then, for each individual, each monitoring day (mean duration = 55 days \pm 32 SD), the mean distance between the bird and the four risky features were calculated during the day- and night-time periods. Day- and night-time periods used to compute these distances were defined as follows: from 75 minutes after dawn to 45 minutes before dusk, and from 75 minutes after dusk to 45 minutes before dawn, respectively (see Harmange et al. 2021). The time deviations from dawn and dusk were computed using the *suncalc* package (Agafonkin and Thieurmél 2018). To study the influence of individual traits on the adjustments in use of space in response to risk variation, synthetic dependent variables capturing the magnitude and the sense of the adjustment were needed. Diel adjustments in use of space were thus computed for each individual as the difference in the mean night- less the mean day-time ($\Delta_{\text{night} - \text{day}}$) distances to each risky feature, averaged over the days monitored in the study and thus resulting in one synthetic value per bird.

Then, we tested the relationship between individual traits (i.e., sex, morphometry, personality, and health status) and the four dependent variables measuring the $\Delta_{\text{night} - \text{day}}$ mean distances to the four risky features, using linear models with recaptured partridges as sampling units (n = 16). The independent variables used were *sex*, *wing loading*, *personality* (the first two axes of the PCA represented personality traits), and *health status* (the first two axes of the PCA represented immunity parameters and coloration of the red skin ornament). Models were selected based on Akaike's information criterion corrected for a small sample size (AICc) (Burnham and Anderson 2002). All candidate models derived from a full (including all independent variables) to a null model were considered in the model selection. We calculated the Akaike model weights (ω_i) as support for each model and contrasted weights of each competing model with the best model (i.e., the model with the lowest AICc value) by calculating the evidence ratio (Burnham and Anderson 2002). Competing models with a difference in AICc (ΔAICc) <2 were considered equivalent. As many candidate models were tested (i.e., 64 models for each of the four dependent variables), only the top four of the most competitive models (ΔAICc with the best models <4) and the null models are shown in the results.

All analyses were performed using R software (R Core Team 2019).

RESULTS

Correlation in Individual Traits

The PCA synthetizing personality traits accounted for 55.6% of the behavioral variability of individuals in the first two dimensions (Fig. 1a; see Table S2 in Online Resource 1 for contributions). The first dimension of the

PCA (PC1-personality, accounting for 33.9% of total variance; Table S2) was positively correlated with alarm calls displayed at release (PCA correlation coefficient between axis and variable $r = 0.83$) and the wriggle score ($r = 0.60$), and was negatively correlated with escape distance ($r = -0.71$), tonic immobility ($r = -0.59$) and flight initiation distance ($r = -0.51$). These results suggest a gradient in boldness in individuals: the higher the PC1-personality score, the bolder the behavior of the individual (Fig. 1a). PC2-personality (21.7% of total variance; Table S2) was positively correlated with distress calls displayed when handled ($r = 0.77$) and escape distance ($r = 0.60$).

The health status of birds was also assessed using a PCA that accounted for 77.4% of the inter-individual variability in the first two dimensions (Fig. 1b, see Table S3 in Online Resource 1 for contributions). PC1-health status (54.6% of total variance, Table S3) was positively correlated with the proportion of monocytes ($r = 0.89$) and heterophils ($r = 0.85$), and negatively correlated with lymphocytes ($r = -0.85$) and eosinophils ($r = -0.69$), hence representing a gradient of acute immune response (Fig. 1b). The PC2-health status (22.8% of the total variance, Table S3) was positively correlated with the red coloration of carotenoid-based skin ornament ($r = 0.94$).

No correlation was detected between the principal components of personality traits (PC1 and PC2-personality) and health status (PC1 and PC2-health status) (Table S4 in Online Resource 1).

Individual Traits and Adjustments in Use of Space

Model selection showed that individual traits affected the adjustments in use of space relative to risky features (Table 2). Analyses showed that $\Delta_{\text{night} - \text{day}}$ distances to hedgerows were positively related to PC1-personality (Table 3): bolder individuals (high PC1-personality score) showed significant greater $\Delta_{\text{night} - \text{day}}$ distances to risky hedgerows (Fig. 2a), suggesting that the birds stayed farther from risky features at night than in the day. Based on model estimates (Fig. 2a), bolder individuals moved 127 m (± 7 SE) farther from hedgerows at night than in the day, while shyer individuals moved only 69 m (± 12) farther away. $\Delta_{\text{night} - \text{day}}$ distances to hedgerows, roads/tracks, and woodlands were significantly related to PC1-health status (Table 3). Individuals in good health moved at a distance of 124 m (± 8) farther from hedgerows, 173 m (± 32) farther from woodlands and 85 m (± 18) farther from roads at night than during the day (Fig. 2b-d). In contrast, individuals with poorer health moved at only 83 m (± 11) farther from hedgerows at night than during the day, did not change their distance from woodlands day or night, and were closer to roads (by 50 to 110 m) at night than during the day (Fig. 2b-d).

No influence was detected for PC2- personality, PC2-health status, sex and wing loading (mean value = 2.44 g/mm \pm 0.14 SD) on adjustments in use of space, nor any effect of any individual traits on the adjustment in distances to built-up areas.

DISCUSSION

We found inter-individual variability in personality and health status parameters, indicating that individuals in our study population actually differed in their boldness traits and health conditions. Moreover, personality and health status affected adjustments in the use of space in relation to risky habitat features.

Inter-individual Variability in Personality and Health Status

Personality traits were reliably identified using the PC1-personality values and varied along a boldness gradient: some individuals appeared to be more aggressive or impulsive as suggested by their strong wriggle behavior when handled, and by the alarm calls they displayed once released (Koolhaas et al. 2007). They also demonstrated higher risk-taking behaviors (lower flight initiation and shorter escape flight distances), consistent with their low tonic immobility, suggesting lower levels of fear and stress (Møller et al. 2011; Edelaar et al. 2012). Such associations between personality traits, shown to be repeatable and heritable both in wild-caught and hand-reared birds (e.g. in the great tit [*Parus major*]: Verbeek et al. 1994, 1996; Van Oers et al. 2004a, b), are commonly observed in birds (Sih and Del Giudice 2012; Hall et al. 2015), and may have implications for their survival and reproduction (Smith and Blumstein 2008; Arroyo et al. 2017).

Individuals also vary in their immune parameters. Some individuals had higher proportions of heterophils and monocytes, and a low proportion of lymphocytes, suggesting a poor health status. Heterophils are the main granulated leukocytes involved in the acute inflammatory response in Galliformes (Harmon 1998), and are known to increase after a coccidian infection in other birds, such as in the greenfinch (*Chloris chloris*) (Hörak et al. 2004). In birds, their high proportion relative to lymphocytes is also indicative of a high level of long-term stress (Müller et al. 2011; see also Svobodová et al. 2013 for gray partridge). These results are consistent with high proportions of monocytes, which are mainly linked to acute inflammatory responses or chronic diseases (e.g. from bacterial and parasitic infection; Weiss and Wardrop 2010). This suggests that the PC1-health status reliably predicts the intensity of the immune response in these birds. Although a correlation between health status and personality traits of birds may have been expected (Barber and Dingemanse 2010), no such correlation was found here (see Klueen et al. 2014 for similar results in the blue tit *Cyanistes caeruleus*).

Influence on Adjustments in Use of Space

We showed that birds varied in their adjustment in use of space relative to risky features and that individual traits (personality and health status) may play a role in determining these diel adjustments (see Dunn et al. 2011; and Bonnot et al. 2014 for other examples on the influence of such individual traits on the use of space, in the great tit and roe deer, respectively). Indeed, we found that individuals that were apparently bolder, were located farther from hedgerows at night than during the day. This suggests that bold individuals adjust their use of space in a pattern consistent with avoiding nocturnal carnivores near such habitats, and/or human activity in open areas during the day. This has also been previously documented in several studies on prey species inhabiting agricultural landscapes such as the European rabbit and the roe deer (Moreno et al. 1996; Lone et al. 2017). This could result from the ability of bolder individuals to quickly return to an environment after a disturbance (Van Oers et al. 2004b; Quinn and Cresswell 2005), and to develop routine behaviors (Groothuis and Carere 2005; Carere et al. 2010), favoring greater adjustments in terms of the diel cycle. Overall, these results are supported by Sih et al. (2004), who reported that highly aggressive animals actively cope with environmental challenges, such as the contrasting risks imposed by predators and humans in agricultural landscapes (Moreno et al. 1996; Lone et al. 2017). They are also consistent with the findings reported by Arroyo et al. (2017), who showed that long-term human disturbance can affect the composition of populations of the Montagu's harrier *Circus pygargus*, by increasing the proportion of bold individuals. However, it should be noted that the data were collected in an area where the shooting of gray partridges had been suspended (though shooting on other game species was allowed). Therefore, bold individuals might not have survived here without this shooting suspension. Indeed, hunters are reported to harvest particularly bold individuals, as observed in the common pheasant and other game species (Ciuti et al. 2012; Madden and Whiteside 2014), as a result of their escape strategy. Bold individuals are more likely to attempt to escape by flying over hunters and are thus more likely to be shot than shy birds adopting more discrete behaviors such as freezing, running, or flying low to the ground (Robertson et al. 1993; Quinn and Cresswell 2005; Madden and Whiteside 2014).

Birds that appeared to be in good health conditions (with low levels of heterophils and monocytes) were located farther from three of the four tested risky features (hedgerows, roads/tracks, and woodlands) at night than during the day. This suggests that healthy individuals might be more inclined to avoid carnivore predation risk at night and human activity in open areas during the day. The gray partridge achieves diel adjustments in use of

space by commuting flights between diurnal and nocturnal sites and habitats (Harmange et al. 2021). The energetic cost of short flights is high (see Tatner and Bryant 1986; Nudds and Bryant 2000) and birds in poor health may thus invest more energy in fighting infection, rather than in costly adjustments in use of space (Lindström et al. 2003; Dunn et al. 2011; van Dijk et al. 2015). For example, Kollias et al. (2004) showed that, within 4 days of an infection administered experimentally, the house finch (*Haemorrhous mexicanus*) demonstrated a 50% reduction in motor activity. The interplay between parasitism and personality or cognition (Kortet et al. 2010; Ducatez et al. 2020) could also explain such behavior, since individuals with higher exploratory behavior and boldness would be more exposed to parasites and subjected to infection (Wilson et al. 1993; Boyer et al. 2010). Despite the absence of a correlation between personality and health in the present study, we cannot exclude the possibility that such interplay may operate over time (see Kortet et al. 2010; Barber and Dingemanse 2010).

Our results are however based on 16 recaptured birds out of the 129 GPS-equipped birds. Of the remaining 113 individuals, 91 disappeared few days after release, 7 were predated (fresh carcasses with predator traces), 3 were shot inadvertently despite shooting suspension, 1 was hit by a vehicle, 4 died from an undetermined cause, and 7 were recaptured but had missing data (e.g. unexploitable GPS data). The low recapture rate is consistent with the high mortality rates commonly observed in gamebirds within the first month after of release (Sokos et al. 2008; Rymesova et al. 2013; Madden et al. 2018). Therefore, individual traits were measured on the birds that survived long enough to be recaptured. This implies that (1) our data did not allow us to test or control for intra-individual variability as several individuals were recaptured only once; (2) the sample population may already have been subjected to selection, thus reducing the inter-individual variability in the individual traits and in the behavioral adjustments measured. Further research involving more individuals and based on longitudinal monitoring and measurements of birds, including pre-release assessment of individual traits, should confirm our findings and help to appropriately address the overall inter-individual variability within the released populations and assess its influence on survival.

Management Implications

Particular attention should be paid to management and conservation programs that rely on the translocation of captive-bred animals (McDougall et al. 2006; Smith and Blumstein 2008). Captive-bred animals often represent the only opportunity to restore self-sustainable populations, e.g. in conservation reintroductions or restocking game populations (Fischer and Lindenmayer 2000; Sokos et al. 2008; Jule et al. 2008), but the quality of such

individuals is in the same time the greatest challenge of translocation success (Griffith et al. 1989; Sokos et al. 2008; Champagnon et al. 2012a). Several generations of captivity and inbreeding may contribute to the impoverishment of the variability of immunogenetics (Ewen et al. 2012; Keller et al. 2012) and heritable personality traits, such as risk-taking and exploratory behaviors (Dingemanse et al. 2002; Van Oers et al. 2004b). This jeopardizes the ability of individuals to adapt to novel environmental conditions, and risks the recovery of these vulnerable populations (Ewen et al. 2012; Merrick and Koprowski 2017).

Our results emphasize the need for managers to examine and select founder populations meticulously, promoting appropriate management of individual traits (see McDougall et al. 2006; Smith and Blumstein 2008), particularly those related to health and behavior (see also WPA and IUCN 2009; IUCN 2013). This will help to reduce the immunological and behavioral naivety of released individuals (Faria et al. 2010; Tetzlaff et al. 2019). We suggest that effective population management through translocation requires focus on quality (i.e. by favoring individual traits that improve response to risk and thus survival expectancy), rather than merely on the quantity, of released individuals. This includes favoring the translocation of wild-caught individuals when possible (Sokos et al. 2008), or alternatively captive-reared populations promoting parental care that reduces the level of fear of animals compared to artificial rearing (Santilli and Bagliacca 2019), anti-predator training that promotes elicitation of motor behaviors in response to predators (Gaudioso et al. 2011; Sánchez-García et al. 2016), minimal human contact increasing the escape and fear-related behaviors to humans (Zaccaroni et al. 2007), controlled pre-exposure to pathogens (Faria et al. 2010) and the maintenance of high diversity of immunogenetics improving the response to infection within the captive population (Charpentier et al. 2008).

REFERENCES

- Aebischer NJ, Ewald JA (2012) The grey partridge in the UK: population status, research, policy and prospects. *Anim Biodivers Conserv* 35:353–362
- Aebischer NJ, Kavanagh B (1997) Grey partridge. In: Hagemeyer WJM, Blair MJ (eds) *The EBCC Atlas of European breeding birds, their distribution and abundance*. T&AD Poyser, London, pp 212–213
- Agafonkin V, Thieurmél B (2018) suncalc: Compute Sun Position, Sunlight Phases, Moon Position and Lunar Phases. R package version 0.4.
- Arroyo B, Mougeot F, Bretagnolle V (2017) Individual variation in behavioural responsiveness to humans leads to differences in breeding success and long-term population phenotypic changes. *Ecol Lett* 20:317–325. <https://doi.org/10.1111/ele.12729>

426 Barber I, Dingemanse NJ (2010) Parasitism and the evolutionary ecology of animal personality. *Philos Trans R*
 427 *Soc Lond B Biol Sci* 365:4077–4088. <https://doi.org/10.1098/rstb.2010.0182>
 428 Beani L, Dessì-Fulgheri F (1986) Gregarious behaviour in a group of captive grey partridges (*Perdix perdix*). *Ital*
 429 *J Zool* 53:69–72.
 430 Berger-Tal O, Saltz D (2014) Using the movement patterns of reintroduced animals to improve reintroduction
 431 success. *Curr Zool* 60:515–526
 432 Biard C, Monceau K, Motreuil S, Moreau J (2015) Interpreting immunological indices: The importance of taking
 433 parasite community into account. An example in blackbirds *Turdus merula*. *Methods Ecol Evol* 6:960–
 434 972.
 435 Bischof R, Gjevestad JGO, Ordiz A, et al (2019) High frequency GPS bursts and path-level analysis reveal linear
 436 feature tracking by red foxes. *Sci Rep* 9:1–13
 437 Bjørneraas K, Van Moorter B, Rolandsen CM, Herfindal I (2010) Screening global positioning system location
 438 data for errors using animal movement characteristics. *J Wildl Manag* 74:1361–1366
 439 Bonnot N, Couriot O, Berger A, et al (2020) Fear of the dark? Contrasting impacts of humans vs lynx on diel
 440 activity of roe deer across Europe. *J Anim Ecol* 89:132–145
 441 Bonnot N, Verheyden H, Blanchard P, et al (2014) Interindividual variability in habitat use: evidence for a risk
 442 management syndrome in roe deer? *Behav Ecol* 26:105–114
 443 Both C, Dingemanse NJ, Drent PJ, Tinbergen JM (2005) Pairs of extreme avian personalities have highest
 444 reproductive success. *J Anim Ecol* 74:667–674
 445 Bouten W, Baaij EW, Shamoun-Baranes J, Camphuysen KCJ (2013) A flexible GPS tracking system for
 446 studying bird behaviour at multiple scales. *J Ornithol* 154:571–580. [https://doi.org/10.1007/s10336-](https://doi.org/10.1007/s10336-012-0908-1)
 447 [012-0908-1](https://doi.org/10.1007/s10336-012-0908-1)
 448 Boyer N, Réale D, Marmet J, Pisanu B, Chapuis JL (2010) Personality, space use and tick load in an introduced
 449 population of Siberian chipmunks *Tamias sibiricus*. *J Anim Ecol* 79:538–547.
 450 Bretagnolle V, Berthet E, Gross N, et al (2018) Towards sustainable and multifunctional agriculture in farmland
 451 landscapes: Lessons from the integrative approach of a French LTSE platform. *Sci Total Environ*
 452 627:822–834
 453 Bro, E. 2016. La Perdrix grise. Biologie, écologie, gestion et conservation. Biotope, Mèze
 454 Bro E, Crosnier A (2012) Grey Partridges *Perdix perdix* in France in 2008: distribution, abundance, and
 455 population change. *Bird study* 59:320–326.

456 Bro E, Mayot P (2006) Opérations de reconstitution des populations de perdrix grises et de perdrix rouges en
 457 France. Bilan quantitatif et acquis techniques. Faune Sauvage 274:6–24

458 Buner F (2009) How to re-introduce gray partridges-conclusions from a releasing project in Switzerland. In:
 459 Cederbaum SB, Faircloth BC, Terhune TM, Thompson JJ, Carroll JP (eds) Gamebird 2006: Quail VI
 460 and Perdix XII. Warnell School of Forestry and Natural Resources, 31 May–4 June 2006, Athens, pp
 461 391–395

462 Buner F, Browne SJ, Aebischer NJ (2011) Experimental assessment of release methods for the re-establishment
 463 of a red-listed galliform, the grey partridge (*Perdix perdix*). Biol Conserv 144:593–601.

464 Burnham KP, Anderson DR (2002) Model selection and multi-model inference : A practical information-
 465 theoretic approach. Model selection and multimodel inference, 2nd ed. Springer, New York, USA.

466 Burns JG, Rodd FH (2008) Hastiness, brain size and predation regime affect the performance of wild guppies in
 467 a spatial memory task. Anim Behav 76:911–922. <https://doi.org/10.1016/j.anbehav.2008.02.017>

468 Carere C, Caramaschi D, Fawcett TW (2010) Covariation between personalities and individual differences in
 469 coping with stress: Converging evidence and hypotheses. Curr Zool 56:728–740

470 Caro T, Sherman PW (2012) Vanishing behaviors. Conserv Lett 5:159–166. [https://doi.org/10.1111/j.1755-](https://doi.org/10.1111/j.1755-263X.2012.00224.x)
 471 [263X.2012.00224.x](https://doi.org/10.1111/j.1755-263X.2012.00224.x)

472 Casas F, Mougeot F, Viñuela J, Bretagnolle V (2009) Effects of hunting on the behaviour and spatial distribution
 473 of farmland birds: importance of hunting-free refuges in agricultural areas. Anim Conserv 12:346–354.
 474 <https://doi.org/10.1111/j.1469-1795.2009.00259.x>

475 Chamberlain DE, Fuller RJ (2000) Local extinctions and changes in species richness of lowland farmland birds
 476 in England and Wales in relation to recent changes in agricultural land-use. Agric Ecosyst Environ
 477 78:1–17

478 Champagnon J, Elmberg J, Guillemain M, et al (2012a) Conspecifics can be aliens too: A review of effects of
 479 restocking practices in vertebrates. J Nat Conserv 20:231–241.
 480 <https://doi.org/10.1016/j.jnc.2012.02.002>

481 Champagnon J, Guillemain M, Elmberg J, et al (2012b) Low survival after release into the wild: assessing “the
 482 burden of captivity” on Mallard physiology and behaviour. Eur J Wildl Res 58:255–267.
 483 <https://doi.org/10.1007/s10344-011-0573-3>

484 Charpentier MJ, Williams CV, Drea CM (2008) Inbreeding depression in ring-tailed lemurs (*Lemur catta*):
485 genetic diversity predicts parasitism, immunocompetence, and survivorship. *Conserv Genet* 9:1605–
486 1615.

487 Ciuti S, Muhly TB, Paton DG, et al (2012) Human selection of elk behavioural traits in a landscape of fear. *Proc*
488 *R Soc Lond B Biol Sci* 279:4407–4416

489 Comolet-Tirman J, Siblet J-P, Witté I, et al (2015) Statuts et tendances des populations d’oiseaux nicheurs de
490 France. *Bilan Simpl Prem Rapp Natl Au Titre Dir Oiseaux Alauda* 83:35–76

491 Dessì-Fulgheri F, Dondini G, Paganin V, Vergari S, Beani L (2001) Factors influencing spatial behaviour and
492 survival of released rock partridges (*Alectoris graeca*). *Game and Wildlife Science* 18:305–317.

493 Dietz MW, Piersma T, Hedenström A, Brugge M (2007) Intraspecific variation in avian pectoral muscle mass:
494 constraints on maintaining manoeuvrability with increasing body mass. *Funct Ecol* 21:317–326

495 Dingemanse NJ, Both C, Drent PJ, et al (2002) Repeatability and heritability of exploratory behaviour in great
496 tits from the wild. *Anim Behav* 64:929–938. <https://doi.org/10.1006/anbe.2002.2006>

497 Ducatez S, Lefebvre L, Sayol F, et al (2020) Host Cognition and Parasitism in Birds: A Review of the Main
498 Mechanisms. *Front Ecol Evol* 8:. <https://doi.org/10.3389/fevo.2020.00102>

499 Dunn JC, Cole EF, Quinn JL (2011) Personality and parasites: sex-dependent associations between avian malaria
500 infection and multiple behavioural traits. *Behav Ecol Sociobiol* 65:1459–1471.
501 <https://doi.org/10.1007/s00265-011-1156-8>

502 Edelaar P, Serrano D, Carrete M, et al (2012) Tonic immobility is a measure of boldness toward predators: an
503 application of Bayesian structural equation modeling. *Behav Ecol* 23:619–626

504 Ewen JG, Acevedo-Whitehouse K, Alley MR, et al (2012) Empirical consideration of parasites and health in
505 reintroduction. *Reintroduction Biol Integrating Sci Manag* 290–335

506 Faria PJ, van Oosterhout C, Cable J (2010) Optimal release strategies for captive-bred animals in reintroduction
507 programs: Experimental infections using the guppy as a model organism. *Biol Conserv* 143:35–41.
508 <https://doi.org/10.1016/j.biocon.2009.06.002>

509 Fischer J, Lindenmayer DB (2000) An assessment of the published results of animal relocations. *Biol Conserv*
510 96:1–11

511 Fortin D, Beyer HL, Boyce MS, et al (2005) Wolves influence elk movements: behavior shapes a trophic
512 cascade in Yellowstone National Park. *Ecology* 86:1320–1330

513 Fucikova E, Drent PJ, Smits N, Van Oers K (2009) Handling stress as a measurement of personality in great tit
514 nestlings (*Parus major*). *Ethology* 115:366–374.

515 Garcia-Longoria L, Garamszegi LZ, Møller AP (2014) Host escape behavior and blood parasite infections in
516 birds. *Behav Ecol* 25:890–900

517 Gaudioso VR, Sánchez-Garcí C, Pérez JA, et al (2011) Does early antipredator training increase the suitability of
518 captive red-legged partridges (*Alectoris rufa*) for releasing? *Poult Sci* 90:1900–1908.
519 <https://doi.org/10.3382/ps.2011-01430>

520 Gée A, Sarasa M, Pays O (2018) Long-term variation of demographic parameters in four small game species in
521 Europe: opportunities and limits to test for a global pattern. *Anim Biodivers Conserv* 41:33–60

522 Gortázar C, Villafuerte R, Martín M (2000) Success of traditional restocking of red-legged partridge for hunting
523 purposes in areas of low density of northeast Spain aragón. *Z Für Jagdwiss* 46:23–30.
524 <https://doi.org/10.1007/BF02240661>

525 Gosler AG, Greenwood JJ, Perrins C (1995) Predation risk and the cost of being fat. *Nature* 377:621–623

526 Griffith B, Scott JM, Carpenter JW, Reed C (1989) Translocation as a species conservation tool: status and
527 strategy. *Science* 245:477–480

528 Groothuis TGG, Carere C (2005) Avian personalities: characterization and epigenesis. *Neurosci Biobehav R*
529 29:137–150.

530 Hall ML, van Asten T, Katsis AC, et al (2015) Animal personality and pace-of-life syndromes: do fast-exploring
531 fairy-wrens die young? *Front Ecol Evol* 3:1–14. <https://doi.org/10.3389/fevo.2015.00028>

532 Harmange C, Bretagnolle V, Chabaud N, et al (2021) Diel cycle in a farmland bird is shaped by contrasting
533 predation and human pressures. *Biol J Linn Soc* in press.

534 Harmange C, Bretagnolle V, Sarasa M, Pays O (2019) Changes in habitat selection patterns of the gray partridge
535 *Perdix perdix* in relation to agricultural landscape dynamics over the past two decades. *Ecol Evol*
536 9:5236–5247.

537 Harmon BG (1998) Avian heterophils in inflammation and disease resistance. *Poult Sci* 77:972–977

538 Hessler E, Tester JR, Siniff DB, Nelson MM (1970) A biotelemetry study of survival of pen-reared pheasants
539 released in selected habitats. *J Wildl Manag* 34:267–274

540 Hopwood JL (2008) The contribution of roadside grassland restorations to native bee conservation. *Biol Conserv*
541 141:2632–2640. <https://doi.org/10.1016/j.biocon.2008.07.026>

542 Hõrak P, Ots I, Vellau H, et al (2001) Carotenoid-based plumage coloration reflects hemoparasite infection and
 543 local survival in breeding great tits. *Oecologia* 126:166–173
 544 Hõrak P, Saks L, Karu U, et al (2004) How coccidian parasites affect health and appearance of greenfinches. *J*
 545 *Anim Ecol* 73:935–947
 546 IUCN S (2013) Guidelines for reintroductions and other conservation translocations. Gland Switz Camb UK
 547 IUCNSSC Re-Introd Spec Group
 548 Jenny M (2015) Chancen und Grenzen der Wiederansiedlung – Erfahrungen aus dem Rebhuhnprojekt. *Hotspot*
 549 31:6–7
 550 Johnson AR, Wiens JA, Milne BT, Crist TO (1992) Animal movements and population dynamics in
 551 heterogeneous landscapes. *Landsc Ecol* 7:63–75
 552 Jule KR, Leaver LA, Lea SEG (2008) The effects of captive experience on reintroduction survival in carnivores:
 553 A review and analysis. *Biol Conserv* 141:355–363. <https://doi.org/10.1016/j.biocon.2007.11.007>
 554 Keller LF, Biebach I, Ewing SR, Hoeck PE (2012) The genetics of reintroductions: inbreeding and genetic drift.
 555 *Reintroduction Biol Integrating Sci Manag* 9:360
 556 Klueen E, Siitari H, Brommer JE (2014) Testing for between individual correlations of personality and
 557 physiological traits in a wild bird. *Behav Ecol Sociobiol* 68:205–213. [https://doi.org/10.1007/s00265-](https://doi.org/10.1007/s00265-013-1635-1)
 558 013-1635-1
 559 Kollias GV, Sydenstricker KV, Kollias HW, Ley DH, Hosseini PR, Connolly V, Dhondt AA (2004)
 560 Experimental infection of house finches with *Mycoplasma gallisepticum*. *J Wildlife Dis* 40:79–86.
 561 Koolhaas JM, De Boer SF, Buwalda B, Van Reenen K (2007) Individual variation in coping with stress: a
 562 multidimensional approach of ultimate and proximate mechanisms. *Brain Behav Evol* 70:218–226
 563 Kortet R, Hedrick AV, Vainikka A (2010) Parasitism, predation and the evolution of animal personalities. *Ecol*
 564 *Lett* 13:1449–1458
 565 Krauze-Gryz D, Gryz JB, Goszczyński J, Chylarecki P, Zmihorski M (2012) The good, the bad, and the ugly:
 566 space use and intraguild interactions among three opportunistic predators—cat (*Felis catus*), dog (*Canis*
 567 *lupus familiaris*), and red fox (*Vulpes vulpes*)—under human pressure. *Can J Zool* 90:1402–1413.
 568 Kuijper DPJ, Oosterveld E, Wymenga E (2009) Decline and potential recovery of the European grey partridge
 569 (*Perdix perdix*) population—a review. *Eur J Wildlife Res* 55:455–463.
 570 Kullberg C, Fransson T, Jakobsson S (1996) Impaired predator evasion in fat blackcaps (*Sylvia atricapilla*). *Proc.*
 571 *R. Soc. Lond. B* 263:1671–1675.

572 Laikre L, Palmé A, Josefsson M, et al (2006) Release of alien populations in Sweden. *AMBIO J Hum Environ*
573 35:255–261

574 Laundré JW, Hernández L, Altendorf KB (2001) Wolves, elk, and bison: reestablishing the "landscape of fear"
575 in Yellowstone National Park, USA. *Can J Zool* 79:1401–1409

576 Lewis JS, Rachlow JL, Garton EO, Vierling LA (2007) Effects of habitat on GPS collar performance: using data
577 screening to reduce location error. *J Appl Ecol* 44:663–671. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2664.2007.01286.x)
578 2664.2007.01286.x

579 Lima SL, Bednekoff PA (1999) Temporal variation in danger drives antipredator behavior: the predation risk
580 allocation hypothesis. *Am Nat* 153:649–659

581 Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: a review and prospectus. *Can J*
582 *Zool* 68:619–640

583 Lindström KM, Van der Veen IT, Legault BA, Lundström JO (2003) Activity and predator escape performance
584 of common greenfinches *Carduelis chloris* infected with sindbis virus. *Ardea* 91:103–111.

585 Lone K, Loe LE, Gobakken T, et al (2014) Living and dying in a multi-predator landscape of fear: roe deer are
586 squeezed by contrasting pattern of predation risk imposed by lynx and humans. *Oikos* 123:641–651.
587 <https://doi.org/10.1111/j.1600-0706.2013.00938.x>

588 Lone K, Loe LE, Meisingset EL, et al (2015) An adaptive behavioural response to hunting: surviving male red
589 deer shift habitat at the onset of the hunting season. *Anim Behav* 102:127–138

590 Lone K, Mysterud A, Gobakken T, et al (2017) Temporal variation in habitat selection breaks the catch-22 of
591 spatially contrasting predation risk from multiple predators. *Oikos* 126:624–632.
592 <https://doi.org/10.1111/oik.03486>

593 Madden JR, Hall A, Whiteside MA (2018) Why do many pheasants released in the UK die, and how can we best
594 reduce their natural mortality? *Eur J Wildl Res* 64:40. <https://doi.org/10.1007/s10344-018-1199-5>

595 Madden JR, Whiteside MA (2014) Selection on behavioural traits during ‘unselective’ harvesting means that shy
596 pheasants better survive a hunting season. *Anim Behav* 87:129–135

597 McDougall PT, Réale D, Sol D, Reader SM (2006) Wildlife conservation and animal temperament: causes and
598 consequences of evolutionary change for captive, reintroduced, and wild populations. *Anim Conserv*
599 9:39–48

600 Merrick MJ, Koprowski JL (2017) Should we consider individual behavior differences in applied wildlife
601 conservation studies? *Biol Conserv* 209:34–44. <https://doi.org/10.1016/j.biocon.2017.01.021>

602 Møller AP (2015) Birds. In: Cooper WEJ, Blumstein DT (eds) Escaping from predators: An integrative view of
 603 escape decisions and refuge use. Cambridge University Press, Cambridge, pp 88–112
 604 Møller AP, Biard C, Blount JD, et al (2000) Carotenoid-dependent signals: indicators of foraging efficiency,
 605 immunocompetence or detoxification ability? *Poult Avian Biol Rev* 11:137–160
 606 Møller AP, Christiansen SS, Mousseau TA (2011) Sexual signals, risk of predation and escape behavior. *Behav*
 607 *Ecol* 22:800–807. <https://doi.org/10.1093/beheco/arr046>
 608 Monterroso P, Alves PC, Ferreras P (2013) Catch me if you can: diel activity patterns of mammalian prey and
 609 predators. *Ethology* 119:1044–1056
 610 Monterroso P, Alves PC, Ferreras P (2014) Plasticity in circadian activity patterns of mesocarnivores in
 611 Southwestern Europe: implications for species coexistence. *Behav Ecol Sociobiol* 68:1403–1417.
 612 <https://doi.org/10.1007/s00265-014-1748-1>
 613 Moore J (2002) Parasites and the behavior of animals. Oxford University Press, Oxford
 614 Moreno S, Delibes M, Villafuerte R (1996) Cover is safe during the day but dangerous at night: the use of
 615 vegetation by European wild rabbits. *Can J Zool* 74:1656–1660
 616 Müller C, Jenni-Eiermann S, Jenni L (2011) Heterophils/Lymphocytes-ratio and circulating corticosterone do
 617 not indicate the same stress imposed on Eurasian kestrel nestlings. *Funct Ecol* 25:566–576
 618 Nathan R, Getz WM, Revilla E, et al (2008) A movement ecology paradigm for unifying organismal movement
 619 research. *Proc Natl Acad Sci* 105:19052–19059
 620 Nudds RL, Bryant DM (2000) The energetic cost of short flights in birds. *J Exp Biol* 203:1561–1572
 621 Pereira M, Rodríguez A (2010) Conservation value of linear woody remnants for two forest carnivores in a
 622 Mediterranean agricultural landscape. *J Appl Ecol* 47:611–620
 623 Pita R, Morgado R, Moreira F, et al (2020) Roads, forestry plantations and hedgerows affect badger occupancy
 624 in intensive Mediterranean farmland. *Agric Ecosyst Environ* 289:106721.
 625 <https://doi.org/10.1016/j.agee.2019.106721>
 626 Potts G (1986) The partridge: pesticides, predation and conservation. Collins, London
 627 Putaala A, Hissa R (1995) Effects of hand-rearing on physiology and anatomy in the grey partridge. *Wildl Biol*
 628 1:27–31
 629 Putaala A, Oksa J, Rintamaki H, Hissa R (1997) Effects of hand-rearing and radiotransmitters on flight of gray
 630 partridge. *J Wildl Manag* 61:1345–1351. <https://doi.org/10.2307/3802136>

631 Quinn JL, Cresswell W (2005) Personality, Anti-Predation Behaviour and Behavioural Plasticity in the
 632 Chaffinch *Fringilla coelebs*. *Behaviour* 142:1377–1402.
 633 R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical
 634 Computing, Vienna, Austria
 635 Rantanen EM, Buner F, Riordan P, Sotherton NW, Macdonald DW (2010) Vigilance, time budgets and
 636 predation risk in reintroduced captive-bred grey partridges *Perdix perdix*. *Appl Anim Behav Sci*
 637 127:43–50.
 638 Réale D, Reader SM, Sol D, et al (2007) Integrating animal temperament within ecology and evolution. *Biol Rev*
 639 82:291–318
 640 Robertson PA, Wise DR, Blake KA (1993) Flying ability of different pheasant strains. *J Wildl Manag* 778–782
 641 Rymesova D, Tomasek O, Salek M (2013) Differences in mortality rates, dispersal distances and breeding
 642 success of commercially reared and wild grey partridges in the Czech agricultural landscape. *Eur J*
 643 *Wildl Res* 59:147–158. <https://doi.org/10.1007/s10344-012-0659-6>
 644 Sánchez-García C, Alonso ME, Tizado EJ, et al (2016) Anti-predator behaviour of adult red-legged partridge
 645 (*Alectoris rufa*) tutors improves the defensive responses of farm-reared broods. *Br Poult Sci* 57:306–
 646 316
 647 Santilli F, Bagliacca M (2019) Fear and behavior of young pheasants reared with or without parent figure. *Avian*
 648 *Biol Res* 12:23–27. <https://doi.org/10.1177/1758155919826765>
 649 Sih A, Bell A, Johnson JC (2004) Behavioral syndromes: an ecological and evolutionary overview. *Trends Ecol*
 650 *Evol* 19:372–378
 651 Sih A, Cote J, Evans M, et al (2012) Ecological implications of behavioural syndromes. *Ecol Lett* 15:278–289
 652 Sih A, Del Giudice M (2012) Linking behavioural syndromes and cognition: a behavioural ecology perspective.
 653 *Philos Trans R Soc B Biol Sci* 367:2762–2772
 654 Smith BR, Blumstein DT (2008) Fitness consequences of personality: a meta-analysis. *Behav Ecol* 19:448–455.
 655 <https://doi.org/10.1093/beheco/arm144>
 656 Sokos CK, Birtsas PK, Tsachalidis EP (2008) The aims of galliforms release and choice of techniques. *Wildl*
 657 *Biol* 14:412–422
 658 Svobodová J, Gabrielová B, Synek P, Marsik P, Vaněk T, Albrecht T, Vinkler M (2013) The health signalling of
 659 ornamental traits in the Grey Partridge (*Perdix perdix*). *J Ornithol* 154:717–725.

660 Tatner P, Bryant DM (1986) Flight Cost of a Small Passerine Measured Using Doubly Labeled Water:
 661 Implications for Energetics Studies. *The Auk* 103:169–180.

662 Tetzlaff SJ, Sperry JH, DeGregorio BA (2019) Effects of antipredator training, environmental enrichment, and
 663 soft release on wildlife translocations: A review and meta-analysis. *Biol Conserv* 236:324–331.
 664 <https://doi.org/10.1016/j.biocon.2019.05.054>

665 Tolon V, Dray S, Loison A, et al (2009) Responding to spatial and temporal variations in predation risk: space
 666 use of a game species in a changing landscape of fear. *Can J Zool* 87:1129–1137

667 van den Hout PJ, Mathot KJ, Maas LRM, Piersma T (2010) Predator escape tactics in birds: linking ecology and
 668 aerodynamics. *Behav Ecol* 21:16–25. <https://doi.org/10.1093/beheco/arp146>

669 van Dijk JG, Kleyheeg E, Soons MB, Nolet BA, Fouchier RA, Klaassen M (2015) Weak negative associations
 670 between avian influenza virus infection and movement behaviour in a key host species, the mallard
 671 *Anas platyrhynchos*. *Oikos* 124:1293–1303.

672 van Heezik Y, Seddon PJ, Maloney RF (1999) Helping reintroduced houbara bustards avoid predation: effective
 673 anti-predator training and the predictive value of pre-release behaviour. *Anim Conserv* 2:155–163

674 Van Oers K, de Jong G, Drent PJ, van Noordwijk AJ (2004a) A genetic analysis of avian personality traits:
 675 correlated, response to artificial selection. *Behav Genet* 34:611–619

676 Van Oers K, Drent PJ, Goede P de, Noordwijk AJ van (2004b) Realized heritability and repeatability of risk-
 677 taking behaviour in relation to avian personalities. *Proc R Soc Lond B Biol Sci* 271:65–73.
 678 <https://doi.org/10.1098/rspb.2003.2518>

679 Verbeek ME, Boon A, Drent PJ (1996) Exploration, aggressive behaviour and dominance in pair-wise
 680 confrontations of juvenile male great tits. *Behaviour* 133:945–963

681 Verbeek ME, Drent PJ, Wiepkema PR (1994) Consistent individual differences in early exploratory behaviour of
 682 male great tits. *Anim Behav* 48:1113–1121

683 von der Lippe M, Bullock JM, Kowarik I, et al (2013) Human-Mediated Dispersal of Seeds by the Airflow of
 684 Vehicles. *PLoS ONE* 8:e52733. <https://doi.org/10.1371/journal.pone.0052733>

685 Watters JV, Meehan CL (2007) Different strokes: can managing behavioral types increase post-release success?
 686 *Appl Anim Behav Sci* 102:364–379

687 Weiss, DJ, Wardrop KJ (2010) Schalm's veterinary hematology. John Wiley & Sons, New York

688 Wilson DS, Coleman K, Clark AB, Biederman L (1993) Shy-bold continuum in pumpkinseed sunfish (*Lepomis*
 689 *gibbosus*): An ecological study of a psychological trait. *J Comp Psychol* 107:250.

690 WPA, IUCN (WORLD PHEASANT ASSOCIATION & IUCN/SSC REINTRODUCTION SPECIALIST
691 GROUP) (2009) Guidelines for the Reintroduction of Galliformes for Conservation Purposes. IUCN,
692 Gland, pp 86

693 Zaccaroni M, Ciuffreda M, Paganin M, Beani L (2007) Does an early aversive experience to humans modify
694 antipredator behaviour in adult Rock partridges? *Ethol Ecol Evol* 19:193–200.
695 <https://doi.org/10.1080/08927014.2007.9522561>

696 Zuk M, Thornhill R, Ligon JD, Johnson K (1990) Parasites and mate choice in red jungle fowl. *Am Zool*
697 30:235–244

Figure captions

Fig. 1 Correlation circles of principal component analyses synthetizing the measures of the partridge personality (a), and health status from immune parameters (b). The inertia of each dimension is reported on the axes. Apparent gradients of boldness (a) and intensity of the immune response (b), suggested by the first axes of PCAs, were added for clarity. Details about personality metrics (Table S1) and variable contributions (Tables S2, S3) are available in Online resource 1

Fig. 2 Effect of personality traits (a) and health status (b-d) on the diel adjustments in use of space by the gray partridge, measured as the difference in distance to risky features (i.e., hedgerows, roads/tracks, and woodlands), between night and day ($\Delta_{\text{night} - \text{day}}$). Predicted values (\pm SE) derived from linear models (see Table 3) are represented with solid lines with gray envelopes. The apparent gradients of boldness (a) and intensity of the immune response (b-d) suggested by the first axes of PCAs were added to improve clarity




Fig. 1




Fig. 2

Table 1 Summary of the total number of released birds, the number of GPS-equipped birds, and the number of recaptured GPS-equipped birds on which individual traits were measured, per release session. The total number of birds are given first, with number of males in brackets.

	Period of release			Total
	2016/12	2017/09	2017/10	
N of birds	131 (66)	98 (48)	81 (40)	310 (154)
N of GPS-equipped birds	63 (57)	36 (17)	30 (15)	129 (89)
N of GPS-equipped birds recaptured	6 (1)	0	10 (6)	16 (7)

Table 2 Statistical support for models testing the effects of individual traits (morphometry, personality (PC1), health status (PC1) and sex) on the adjustments in use of space in relation to risky features (hedgerows, roads / tracks, woodlands, built-up areas). The selection process was based on the Akaike's Information Criterion corrected for small sample size (AICc). The number of parameters (K) is presented together with model weights (ω_i), log likelihood values (LL) and differences in AICc ($\Delta AICc$) in respect of the model with the highest statistical support. Personality (PC1- and PC2-perso) and health status (PC1- and PC2-health) correspond to the scores on the first two axes of the two principal component analyses previously performed. Models selected are represented in bold. Only the top four and the null models are shown (see Table 3 for estimates).

Dependent variables	Candidate models	K	AICc	$\Delta AICc$	ω_i	LL	Evidence ratio
(a) $\Delta_{\text{night-day}}$ distance to hedgerows	PC1-perso+PC1-health	4	148.18	0	0.35	-68.27	
	PC1-perso+PC1-health+PC2-health	5	150.44	2.27	0.11	-67.22	3.11
	Sex+PC1-perso+PC1-health	5	150.91	2.73	0.09	-67.46	3.92
	PC1-perso	3	150.94	2.77	0.09	-71.47	3.99
	NULL	2	158.25	10.07	0	-76.66	153.70
(b) $\Delta_{\text{night-day}}$ distance to roads	PC1-health	3	173.22	0	0.46	-82.61	
	Sex+PC1-health	4	176.13	2.91	0.11	-82.25	4.28
	PC1-perso+PC1-health	4	176.74	3.52	0.08	-82.55	5.81
	PC1-health+PC2-perso	4	176.77	3.55	0.08	-82.57	5.90
	NULL	2	187.45	14.23	0	-91.26	1230.28
(c) $\Delta_{\text{night-day}}$ distance to woodlands	PC1-health	3	191.73	0	0.28	-91.86	
	Wing load+PC1-health	4	192.91	1.18	0.15	-90.64	1.80
	PC1-health+PC2-health	4	193.66	1.93	0.10	-91.01	2.62
	PC1-perso+PC1-health	4	194.46	2.73	0.07	-91.41	3.92
	NULL	2	196.81	5.08	0.02	-95.94	12.68
(d) $\Delta_{\text{night-day}}$ distance to buildings	PC1-health	3	205.36	0	0.16	-98.68	
	Wing load+PC1-health	4	206.10	0.75	0.11	-97.23	1.45
	NULL	2	206.43	1.07	0.10	-100.75	1.71
	PC1-health+PC2-perso	4	207.62	2.26	0.05	-97.99	3.10
	Sex	3	207.86	2.50	0.05	-99.93	3.49

Table 3 Estimates of the effect of individual traits on the adjustments in use of space in relation to risky features (a: hedgerows, b: roads / tracks, c: woodlands), from linear models previously selected (see Table 2).

Dependent variable	Independent variables	Estimate (\pm SE)	P value	Adj. R²
$\Delta_{\text{night} - \text{day}}$ distances to hedges				0.60
	(intercept)	109.03 (4.79)	<0.01	
	Health status	-7.43 (2.94)	0.03	
	Personality	11.77 (3.15)	<0.01	
$\Delta_{\text{night} - \text{day}}$ distances to roads				0.64
	(intercept)	14.13 (11.30)	0.23	
	Health status	-35.73 (6.84)	<0.01	
$\Delta_{\text{night} - \text{day}}$ distances to woodlands				0.36
	(intercept)	98.25 (20.15)	<0.01	
	Health status	-37.22 (12.19)	<0.01	

Online Resource 1. Additional information and results

Table S1 Behavioral metrics assessed as measures of bird personality (see Møller et al. 2011).

Behavioral metric	Assessment	Mechanism
Flight initiation distance (FID)	Distance at which a bird placed in a cage in an open area composed of bare ground near the capture site, reacted with a take-off attempt to a straight-line approach of the field operator (after 2-min of acclimation). All FID were assessed under clear weather conditions in daylight, by the same operator to standardize the approach speed, the operator dress, and thus limit the detectability bias.	Individuals with shorter FID (i.e., they escape when the predator is closer) are bolder and have a higher risk of being predated (Møller et al. 2008, 2010, Holtmann et al. 2017).
Wriggle score	This scored how much a bird moved during handling: 0 – no movement, 1 – moved rarely, 2 – regularly, 3 – continuously	Individuals that wriggle more are more likely to escape from a predator when captured (Møller et al. 2011).
Feather loss	This scored whether the bird lost feathers during handling (1), or not (0).	Feather loss can advantage a bird caught by a predator, as predator could lose their grip as a result (Møller et al. 2006).
Distress call	This scored whether the bird produced fear screams during handling (1), or not (0).	Birds emitting fear screams could increase their probability of escape once captured, attracting the attention of secondary predators (Møller and Nielsen 2010).
Tonic immobility	Before release, the bird was placed on its back in our flat hand, with a second hand restraining it until it was still. The time between when the second hand was removed, and the bird flew away was recorded. The trial was stopped after a delay of more than 30 s.	The longer the bird stays in the hand, the higher its level of fear (Møller et al. 2011, Edelaar et al. 2012)
Alarm call	This scored whether the bird emitted an alarm call leaving our hand after release (1), or not (0).	Alarm calls could play a role in distracting a predator, thereby decreasing the predation susceptibility of conspecifics, and are expected to be produced by bolder individuals, placing them in a greater immediate risk (Charnov and Krebs 1975, Platzen and Magrath 2004).
Escape flight distance	This is the flight distance achieved post-handling release.	Individuals with a shorter escape flight distance (also called distance fled) have a higher risk of being caught by a predator (Cooper and Blumstein 2015, Tätté et al. 2018).

Table S2 Contributions (%) of variables to the first three dimensions of the principal component analysis carried out on personality traits (see Fig. 1a). The eigen values and the percentage of total variance explained by each dimension is given in brackets.

Variable	Dimension 1 (2.37; 33.91%)	Dimension 2 (1.52; 21.73%)	Dimension 3 (1.28; 18.24%)
Flight initiation distance	11.13	12.45	16.40
Wriggle score	15.15	15.34	6.74
Feather loss	4.33	8.65	49.99
Distress call	4.54	39.31	1.40
Tonic immobility	14.42	0.29	10.15
Alarm call	29.34	0.08	7.66
Escape distance	21.08	23.89	7.67

Table S3 Contributions (%) of variables to the first three dimensions of the principal component analysis carried out on health status (immune parameters, see Fig. 1b). The eigen values and the

Variable	Dimension 1 (2.73; 54.58%)	Dimension 2 (1.14; 22.79%)	Dimension 3 (0.91; 18.12%)
Red coloration	0.20	77.52	8.97
Heterophils	26.70	12.16	6.05
Eosinophils	17.37	0.88	54.44
Monocytes	29.04	8.35	3.54
Lymphocytes	26.69	1.10	26.99

percentage of total variance explained by each dimension is given in brackets.

Table S4 Matrix of Spearman's correlation coefficients (p-values) between the first two axes of each principal component analysis carried out on personality (PC1- and PC2-perso) and health status (PC1 and PC2-health).

	PC1-perso	PC2-perso	PC1-health	PC2-health
PC1-perso	-	-	-	-
PC2-perso	0.121 (0.66)	-	-	-
PC1-health	-0.197 (0.46)	0.006 (0.99)	-	-
PC2-health	-0.253 (0.34)	0.003 (1.00)	0.006 (0.99)	-

References

- Charnov EL, Krebs JR (1975) The evolution of alarm calls: altruism or manipulation? *Am Nat* 109:107–112.
- Cooper WE, Blumstein DT (2015) *Escaping from predators: an integrative view of escape decisions*. Cambridge University Press, Cambridge
- Edelaar P, Serrano D, Carrete M, Blas J, Potti J, Tella JL (2012) Tonic immobility is a measure of boldness toward predators: an application of Bayesian structural equation modeling. *Behav Ecol* 23:619–626.
- Holtmann B, Santos ES, Lara CE, Nakagawa S (2017) Personality-matching habitat choice, rather than behavioural plasticity, is a likely driver of a phenotype–environment covariance. *Proc R Soc Lond B Biol Sci* 284:1–9.
- Møller AP, Christiansen SS, Mousseau TA (2011) Sexual signals, risk of predation and escape behavior. *Behav Ecol* 22:800–807.
- Møller AP, Erritzøe J, Nielsen JT (2010) Causes of interspecific variation in susceptibility to cat predation on birds. *Chinese Birds* 1:97–111.
- Møller AP, Nielsen JT (2010) Fear screams and adaptation to avoid imminent death: effects of genetic variation and predation. *Ethol Ecol Evol* 22:183–202.
- Møller AP, Nielsen JT, Erritzøe J (2006). Losing the last feather: feather loss as an antipredator adaptation in birds. *Behav Ecol* 17:1046–1056.
- Møller AP, Nielsen JT, Garamzegi LZ (2008) Risk taking by singing males. *Behavioral Ecology* 19:41–53.
- Platzen D, Magrath RD (2004) Parental alarm calls suppress nestling vocalization. *Proc R Soc Lond B Biol Sci* 271:1271–1276.
- Tätte K, Møller AP, R Mänd (2018) Towards an integrated view of escape decisions in birds: relation between flight initiation distance and distance fled. *Anim Behav* 136:75–86.

Harmange C*, Bretagnol V, Pays O, Sarasa M, Moreau J. Linking Personality and Health to Use of Space in the Gray Partridge: Perspectives for Management. *European Journal of Wildlife Research*.

*Corresponding author: Clément Harmange, LETG-Angers, UMR 6554 CNRS, Université d'Angers, 49000 Angers, France, c.harmange.pro@gmail.com.

Online Resource 2. Protocol for the calculation of white cell proportions from gray partridge blood smears

A blood smear was collected from each recaptured bird, to determine the health status of the birds through white cell counts (Zuk et al. 1990, Hůrak et al. 2001, Svobodová et al. 2013, Biard et al. 2015). Birds were placed on their backs, with one wing spread, and a drop of blood was collected through a capillary from the brachial vein and deposited on a glass slide. Another slide was used as a spreader; it was placed at a 45° angle, pushed back towards the blood drop, and then pushed across the slide to spread the drop all along the slide.

Blood smears were fixed in the laboratory by submergence in a methanol bath for five to seven minutes. The smears were stained with a Giemsa solution diluted at 1/20 for 40 minutes. Finally, smear slices were rinsed several times with neutral pH water and left to air dry (Biard et al. 2010).

The white cell count was determined using immersion oil under a microscope with a 1000 × magnification. The number of leukocytes (distinguishing the heterophils, monocytes, lymphocytes, eosinophils, basophils; Campbell and Dein 1984, Weiss and Wardrop 2010) were reported, counted per reference number of 10000 erythrocytes. If the total number of leukocytes was lower than 100, the counting was continued until the number of leukocytes reached 100. The number of each cell type needed to reach this level of leukocyte count was also reported.

References

- Biard C, Monceau K, Motreuil S, Moreau J (2015) Interpreting immunological indices: The importance of taking parasite community into account. An example in blackbirds *Turdus merula*. *Methods Ecol Evol* 6:960–972.
- Biard C, Saulnier N, Gaillard M, Moreau J (2010) Carotenoid-based bill colour is an integrative signal of multiple parasite infection in blackbird. *Naturwissenschaften* 97:987–995.
- Campbell TW, Dein FJ (1984) Avian hematology. The basics. *The Veterinary Clinics of North America. Small Animal Practice* 14:223–248.
- Hůrak P, Ots I, Vellau H, Spottiswoode C, Møller AP (2001) Carotenoid-based plumage coloration reflects hemoparasite infection and local survival in breeding great tits. *Oecologia* 126:166–173.
- Svobodová J, Gabrielová B, Synek P, Marsik P, Vaněk T, Albrecht T, Vinkler M (2013) The health signalling of ornamental traits in the Grey Partridge (*Perdix perdix*). *J Ornithol* 154:717–725.
- Weiss DJ, Wardrop KJ (2010) Schalm's veterinary hematology. John Wiley & Sons, New York
- Zuk M, Thornhill R, Ligon JD, Johnson K (1990) Parasites and mate choice in red jungle fowl. *Am Zool* 30:235–244.

Harmange C*, Bretagnolle V, Pays O, Sarasa M, Moreau J. Linking Personality and Health to Use of Space in the Gray Partridge: Perspectives for Management. *European Journal of Wildlife Research*.

*Corresponding author: Clément Harmange, LETG-Angers, UMR 6554 CNRS, Université d'Angers, 49000 Angers, France, c.harmange.pro@gmail.com.

Online Resource 3. Protocol for the colorimetry calculation of carotenoid-based red skin ornament behind the eye of gray partridge

Each recaptured bird was photographed (Fujifilm FinePix S4000) to assess the coloration of the carotenoid-based red skin ornament behind the eye, as an indicator of health status (Svobodová et al. 2013). Birds were placed outside, out of direct sun exposure, and their right-side profile was photographed from a distance of approximately 50 cm from the camera lens. The same gray standard reference was placed near birds in each picture (thus, in the same exposure and at the same distance from the lens), to standardize coloration measurements between pictures.

Pictures were analyzed using Adobe Photoshop® software, following the method of Lopez-Antia et al. (2013). An area of 0.20*0.20 cm was selected on the red skin ornament patch. The mean intensity values of red, green and blue (or RGB components) were calculated for all pixels contained within this area (Lopez-Antia et al. 2013). These values were corrected using the average RGB values from an area of the same dimension (0.20*0.20 cm) and in the same light exposure conditions on the gray reference to standardize color measurements between pictures (Lopez-Antia et al. 2013).

Lastly, the intensity of the carotenoid-based red skin ornament was calculated as the R value divided by the average values of R, G and B (see Lopez-Antia et al. 2013).

References

- Lopez-Antia A, Ortiz-Santaliestra ME, Mougeot F, Mateo R (2013) Experimental exposure of red-legged partridges (*Alectoris rufa*) to seeds coated with imidacloprid, thiram and difenoconazole. *Ecotoxicology* 22:125–138.
- Svobodová J, Gabrielová B, Synek P, Marsik P, Vaněk T, Albrecht T, Vinkler M (2013) The health signalling of ornamental traits in the Grey Partridge (*Perdix perdix*). *J Ornithol* 154:717–725.

Harmange C*, Bretagnolle V, Pays O, Sarasa M, Moreau J. Linking Personality and Health to Use of Space in the Gray Partridge: Perspectives for Management. *European Journal of Wildlife Research*.

*Corresponding author: Clément Harmange, LETG-Angers, UMR 6554 CNRS, Université d'Angers, 49000 Angers, France, c.harmange.pro@gmail.com.