



Wearable Inertial Measurement Unit to Accelerometer-Based Training Monotony and Strain during a Soccer Season: A within-Group Study for Starters and Non-Starters

Hadi Nobari, Mustafa Sogut, Rafael Oliveira, Jorge Perez-Gomez, Katsuhiko Suzuki, Hassane Zouhal

► To cite this version:

Hadi Nobari, Mustafa Sogut, Rafael Oliveira, Jorge Perez-Gomez, Katsuhiko Suzuki, et al.. Wearable Inertial Measurement Unit to Accelerometer-Based Training Monotony and Strain during a Soccer Season: A within-Group Study for Starters and Non-Starters. International Journal of Environmental Research and Public Health, 2021, 18 (15), pp.8007. <10.3390/ijerph18158007>. <hal-03331922>

HAL Id: hal-03331922

<https://hal.science/hal-03331922v1>

Submitted on 2 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Article

Wearable Inertial Measurement Unit to Accelerometer-Based Training Monotony and Strain during a Soccer Season: A within-Group Study for Starters and Non-Starters

Hadi Nobari ^{1,2,3,4,*} , Mustafa Sögüt ⁵ , Rafael Oliveira ^{6,7,8} , Jorge Pérez-Gómez ^{2,*} , Katsuhiko Suzuki ^{9,*} and Hassane Zouhal ^{10,*}

- ¹ Department of Physical Education and Sports, University of Granada, 18010 Granada, Spain
 - ² HEME Research Group, Faculty of Sport Sciences, University of Extremadura, 10003 Cáceres, Spain
 - ³ Department of Exercise Physiology, Faculty of Sport Sciences, University of Isfahan, Isfahan 81746-7344, Iran
 - ⁴ Sports Scientist, Sepahan Football Club, Isfahan 81887-78473, Iran
 - ⁵ Department of Physical Education and Sports, Faculty of Education, Middle East Technical University, 06800 Ankara, Turkey; msogut@metu.edu.tr
 - ⁶ ESDRM-IPS-Sports Science School of Rio Maior—Polytechnic Institute of Santarém, 2040-413 Rio Maior, Portugal; rafaeloliveira@esdrm.ipsantarém.pt
 - ⁷ Research Center in Sport Sciences, Health Sciences and Human Development, Quinta de Prados, Edifício Ciências de Desporto, 5001-801 Vila Real, Portugal
 - ⁸ Life Quality Research Centre, Complexo Andaluz, Apartado, 2040-413 Rio Maior, Portugal
 - ⁹ Faculty of Sport Sciences, Waseda University, Tokorozawa 359-1192, Japan
 - ¹⁰ Movement, Sport, Health and Sciences Laboratory (M2S), University of Rennes 2, F-35000 Rennes, France
- * Correspondence: hadi.nobari1@gmail.com (H.N.); jorgepg100@gmail.com (J.P.-G.); katsu.suzu@waseda.jp (K.S.); hassane.zouhal@univ-rennes2.fr (H.Z.)



Citation: Nobari, H.; Sögüt, M.; Oliveira, R.; Pérez-Gómez, J.; Suzuki, K.; Zouhal, H. Wearable Inertial Measurement Unit to Accelerometer-Based Training Monotony and Strain during a Soccer Season: A within-Group Study for Starters and Non-Starters. *Int. J. Environ. Res. Public Health* **2021**, *18*, 8007. <https://doi.org/10.3390/ijerph18158007>

Academic Editor: Paul B. Tchounwou

Received: 17 June 2021

Accepted: 26 July 2021

Published: 28 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Abstract: The purpose of this study was to analyze the intragroup differences in weekly training monotony (TM) and training strain (TS) between starter and non-starter male professional soccer players at accelerometry based variables throughout the periods of a season. TM and TS of different accelerations and decelerations zones for twenty-one players were followed for forty-eight weeks. Regardless of group, players obtained the highest mean TM (starters = 3.3 ± 0.6 , non-starters = 2.2 ± 1.1 , in arbitrary unit, AU) and TS (starters = 1288.9 ± 265.2 , non-starters = 765.4 ± 547.5 , AU) scores in the pre-season for accelerations at Zone 1 ($<2 \text{ m/s}^2$). The results also indicated that both groups exhibited similar TM and TS scores in accelerations at Zones 2 ($2 \text{ to } 4 \text{ m/s}^2$) and 3 ($>4 \text{ m/s}^2$) across the entire season. While the starters showed the highest TM and TS scores at deceleration Zone 1 ($<-2 \text{ m/s}^2$) in the end-season, the non-starters exhibited the highest scores at the deceleration Zone 1 in pre-season. It seems that in pre-season, coaches applied higher levels of training with greater emphasis on deceleration for non-starters. This tendency was reduced over time for non-starters, while starters presented higher values of deceleration Zone 1. These results highlight the variations in TM and TS across the different periods of a full season according to match starting status among professional soccer players, and the results suggest that non-starter players should receive higher levels of load to compensate for non-participation in matches throughout a soccer season.

Keywords: football; external training load; WIMU; GPS; acceleration; deceleration



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Monitoring training load in professional soccer is one of the primary focuses of coaching staff in order to analyze individual training requirements, to optimize physical fitness, and to minimize the risk of injury [1,2]. The process of training load monitoring can be categorized as either internal or external with respect to the load units [3]. Internal training load is related to the psychophysiological stress imposed on players during training and competition, and it is usually measured using ratings of perceived exertion, blood lactate, oxygen consumption, and heart rate. External training load includes measures of

training load data derived from advanced technological devices such as global positioning system (GPS) and electronic tracking systems [4].

Other commonly used training load measures are training monotony (TM) and training strain (TS). The TM is a measure of day-to-day variations in training load, and it is calculated as the mean daily training load divided by the weekly standard deviation load, while TS is a product of TM and weekly training load [5]. More specifically, rather than performing an equal daily training load throughout the week, interspersing low and high loading days can help maintain lower or moderate monotony and strain [6]. Furthermore, quantifying weekly TM and TS may prevent overtraining syndrome and negative health consequences [7,8].

Previous studies have documented the seasonal changes in training load, TM, and TS in professional and collegiate soccer players [9–12]. Fessi et al. [11] revealed that exposure to higher pre-season training loads resulted in higher TM, TS, and a lower psychophysical state during the in-season. In other recent studies, match starting status-related differences in accumulated workload, TM, and TS were investigated, and greater values were reported for starting soccer players compared to their non-starting counterparts throughout the season [10,13]. The discrepancy in accumulated training and match workloads between starter and non-starter players may also have consequences on physical fitness levels. The findings of the several earlier examinations demonstrated the associations of playing time status with seasonal changes in body fat percentages, sprinting ability, and muscular strength capacity [14,15]. For example, a significant increase in body fat percentage was reported for non-starters, while significant decrements were found in sprinting ability and vertical jump performance for starters throughout a season [14].

Nevertheless, there is scarce information regarding the intragroup differences in starting and non-starting soccer players with respect to the weekly TM and TS at different speed zones and across the periods of a full season. Recently, and in opposition to the previous information, the study from Oliveira et al. [16] found that both starters and non-starters presented small differences. The same authors suggested that training workload adjustments applied over the season helped to reduce differences between player status.

Therefore, the purpose of this study was to determine the within-group variations in TM and TS at different speed zones during a professional soccer season in both starters and non-starters. The present study is a complement to a previous study that compared starter versus non-starter players across a full season [13]. The seasonal changes according to the players' playing time status may help coaching staff to design optimal training programs.

2. Materials and Methods

2.1. Participants

This study is the second stage in a two-stage investigation conducted on twenty-one professional soccer players (age, 28.3 ± 3.8 yrs; height, 181.2 ± 7.1 cm; and body mass, 74.5 ± 7.7 kg) of one team competing in the Iranian Persian Gulf Pro League were evaluated for 48 weeks during a full season [13]. Previously, the data from the same players were used to determine the differences between starter and non-starter players on weekly TM and TS values across the full soccer season [13]. Specifically, the main focus of this second-stage study is to understand the within-group variations with respect to playing time status. During the full season, there were 44 matches, 200 training sessions, 14,127 min of play and sessions, of which 3960 min are related to participating in match-play; 7 weeks were congested (i.e., two or more matches within seven days), and 30 weeks were non-congested.

The inclusion criteria were to have a minimum three training sessions per week. The two exclusion criteria of this study were (i) the absence of a player for two weeks, resulting in the player being removed from the study, and (ii) goalkeepers were excluded from the study. Prior to commencing the study, ethical approval was granted from the research ethics committee of the University of Isfahan (IR.UI.REC.1399.064). All players were informed of the purpose of the study before providing signed informed consent. All stages of this study

were conducted according to conditions for human studies as outlined in the declaration of Helsinki.

2.2. Experimental Approach to the Problem

The study included a full season of a professional soccer team for 48 weeks (W) in the Persian Gulf Pro League and a knockout tournament during the 2018–2019 season. The 48 weeks of the full-season were divided into four periods: pre-season, W1 to W5; early-season, W6 to W19; mid-season, W20 to W35; and end-season, W36 to W48, in order to analyze within-group differences between starting and non-starting players in terms of accelerometer variables obtained using GPS along the season periods.

The criterion for dividing players into two groups based on previous studies was 60 min of play in weekly matches [13,17]. If a competition was not held during a week, the division criterion was based on the total training time for each group per week [13,18,19]. Based on this criterion, 10 players were placed in the starter group, and 11 players were placed in the non-starter group.

2.3. External Monitoring Measures

The microelectromechanical system (MEMS) used in this study was a GPSPORTS systems Pty Ltd., model: SPI High-Performance Unit (HPU), made in Australia. It was used in all training and competition sessions during the full season for all players. This tool is based on tracking and SPI HPU and includes GPS position with 15 Hz; Accelerometer: 100 Hz, G Tri-Axial-Track impacts. Mag: 50 Hz, three-axis; Water resistance and data transmission: Infrared and weighs 56 g. The validity and reliability of the device for the variables used in previous studies have been confirmed [20]. A coefficient of variation = 0.90% showed the accuracy of the MEMS used in the present study to track high-sprinting velocities [21].

2.4. Data Collection by Wearable Inertial Measurement Unit

Data were exported from the GPS as described by previous studies [13,18,19,22]. The accelerations at zone (AcZ) and decelerations at zone (DcZ) for each level used from the output data for this study are as follows: AcZ1 ($<2 \text{ m/s}^2$); AcZ2 (2 to 4 m/s^2); AcZ3 ($>4 \text{ m/s}^2$); DcZ1 ($<-2 \text{ m/s}^2$); DcZ2 (-2 to -4 m/s^2); and DcZ3 ($>-4 \text{ m/s}^2$) [13,23]. Each of these variables was calculated based on acute training load means the total load in a week of body load [24,25]. Afterward, weekly training monotony (wTM = the average weekly acute training load) was divided by the standard deviation of that week [26,27], and the weekly training strain (wTS = obtained from the weekly acute training load multiplying of the wTM) was used [13,27,28].

2.5. Statistical Analysis

The Statistical Package for the Social Sciences (SPSS, version 25.0; IBM SPSS Inc., Chicago, IL, USA) was used for statistical procedures and analyses. Data are presented as mean and standard deviation (SD). Kolmogorov–Smirnov and Levene's tests were executed to check the normality and homogeneity of the data, respectively. Inferential tests were the conducted. Repeated measures analysis of variance (ANOVA) was applied to analyze within-group changes across the different periods of the season for all dependent variables for both starting and non-starting soccer players. Bonferroni post hoc tests were also executed to determine pairwise comparison outcomes. These tests are based on the linearly independent pairwise comparisons among the estimated marginal means. Significant differences were considered for $p \leq 0.05$. Partial eta squared (η^2) was calculated as the effect size of the repeated measures of ANOVA. Moreover, Hedge's g effect size (95% confidence interval) was calculated to determine the magnitude of the pairwise comparisons. Hopkins' thresholds for the Cohen d effect size statistics were used [29] as follow: ≤ 0.2 , trivial; >0.2 , small; >0.6 , moderate; >1.2 , large; >2.0 , very large; and >4.0 , nearly perfect.

3. Results

Intragroup differences for binary comparisons between season periods in TM based on Ac at zone (TM_{AcZ1}) and TS based on De at zone (TS_{DeZ1}) and each level for both the starters and non-starters are shown in Tables 1–6. Table 1 illustrates the intragroup comparisons of TM_{AcZ1} and TS_{AcZ1} between the different periods of the season for both starters and non-starters. The outcomes revealed significant differences between season periods for starters ($p < 0.001$, $\eta^2 = 0.759$) and non-starters ($p < 0.001$, $\eta^2 = 0.393$) at TM_{AcZ1} as well as for TS_{AcZ1} between starters ($p < 0.001$, $\eta^2 = 0.717$) and non-starters ($p = 0.029$, $\eta^2 = 0.405$). The dual sets TM_{AcZ1} and TS_{AcZ1} are shown in Table 1 and Figure 1a,b.

Table 1. Intragroup differences for binary comparisons between season periods in TM and TS in AcZ1 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Groups	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM _{AcZ1} (AU)	Pre-season	Starters: 3.25 ± 0.62 Non-starters: 2.18 ± 1.14	PreS vs. EarS	Starters	0.002	2.60 [1.41 to 3.79]
				Non-starters	0.016	1.24 [0.33 to 2.16]
	Early-season	Starters: 1.87 ± 0.36 Non-starters: 1.13 ± 0.13	PreS vs. MidS	Starters	0.010	2.13 [1.03 to 3.23]
				Non-starters	0.016	1.19 [0.28 to 2.10]
			PreS vs. EndS	Starters	0.320	1.04 [0.11 to 1.98]
				Non-starters	0.020	1.42 [0.49 to 2.36]
	Mid-season	Starters: 2.12 ± 0.37 Non-starters: 1.15 ± 0.27	EarS vs. MidS	Starters	0.143	−0.64 [−1.54 to 0.26]
				Non-starters	>0.999	−0.10 [−0.94 to 0.74]
			EarS vs. EndS	Starters	0.003	−0.95 [−1.88 to −0.03]
				Non-starters	>0.999	1.15 [0.25 to 2.05]
	End-season	Starters: 2.48 ± 0.79 Non-starters: 0.98 ± 0.13	MidS vs. EndS	Starters	0.021	−0.57 [−1.46 to 0.33]
				Non-starters	0.679	0.78 [−0.09 to 1.65]
TS _{AcZ1} (AU)	Pre-season	Starters: 1288.97 ± 265.15 Non-starters: 765.44 ± 547.51	PreS vs. EarS	Starters	0.002	2.76 [1.54 to 3.98]
				Non-starters	0.022	1.13 [0.23 to 2.03]
	Early-season	Starters: 659.98 ± 158.04 Non-starters: 306.50 ± 63.24	PreS vs. MidS	Starters	0.003	2.68 [1.47 to 3.88]
				Non-starters	0.013	1.18 [0.27 to 2.09]
			PreS vs. EndS	Starters	0.039	1.74 [0.71 to 2.76]
				Non-starters	0.013	1.36 [0.43 to 2.29]
	Mid-season	Starters: 683.62 ± 153.10 Non-starters: 282.55 ± 101.83	EarS vs. MidS	Starters	>0.999	−0.15 [−1.02 to 0.73]
				Non-starters	>0.999	0.27 [−0.57 to 1.11]
			EarS vs. EndS	Starters	0.114	−0.56 [−1.45 to 0.33]
				Non-starters	0.515	1.55 [0.59 to 2.50]
	End-season	Starters: 793.44 ± 281.32 Non-starters: 216.63 ± 47.49	MidS vs. EndS	Starters	0.047	−0.46 [−1.35 to 0.42]
				Non-starters	0.463	0.80 [−0.47 to 1.67]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, pre-season period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ1}, weekly average training monotony based on number of accelerations at Zone 1 (<2 m·s^{−2}); TS_{AcZ1}, weekly average training strain based on number of accelerations at Zone 1 (<2 m·s^{−2}); *p*, *p*-value at alpha level 0.05; Hedge's *g* (95% CI), Hedge's *g* effect size magnitude with 95% confidence interval. Significant differences ($p \leq 0.05$) are highlighted in bold.

Table 2 and Figure 1c,d present the intragroup comparisons of TM_{AcZ2} and TS_{AcZ2} between the different periods of the season for both the starters and non-starters. According to Table 2, TM_{AcZ2} analysis for the starters ($p = 0.031$, $\eta^2 = 0.397$) showed significant results. However, there were no significant changes in TM_{AcZ2} ($p = 0.280$, $\eta^2 = 0.197$) in the non-starters nor in TS_{AcZ2} for the starters ($p = 0.102$, $\eta^2 = 0.299$) and non-starters ($p = 0.054$, $\eta^2 = 0.354$).

Table 2. Intragroup differences for binary comparisons between season periods in TM and TS in AcZ2 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Groups	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM _{AcZ2} (AU)	Pre-season	Starters: 1.82 ± 0.26 Non-starters: 1.19 ± 0.39	PreS vs. EarS	Starters	>0.999	0.18 [−0.70 to 1.06]
				Non-starters	>0.999	0.36 [−0.48 to 1.20]
	Early-season	Starters: 1.76 ± 0.36 Non-starters: 1.09 ± 0.12	PreS vs. MidS	Starters	>0.999	−0.20 [−1.08 to 0.68]
				Non-starters	>0.999	0.24 [−0.59 to 1.08]
	Mid-season	Starters: 1.87 ± 0.27 Non-starters: 1.11 ± 0.27	PreS vs. EndS	Starters	0.365	−0.63 [−1.53 to 0.27]
				Non-starters	0.387	0.90 [0.02 to 1.77]
	End-season	Starters: 2.10 ± 0.55 Non-starters: 0.93 ± 0.10	EarS vs. MidS	Starters	0.980	−0.34 [−1.22 to 0.54]
				Non-starters	>0.999	−0.11 [−0.94 to 0.73]
			EarS vs. EndS	Starters	0.013	−0.70 [−1.61 to 0.20]
				Non-starters	0.630	1.32 [0.40 to 2.24]
			MidS vs. EndS	Starters	0.174	−0.51 [−1.40 to 0.39]
				Non-starters	0.389	0.85 [−0.03 to 1.72]
TS _{AcZ2} (AU)	Pre-season	Starters: 209.71 ± 38.04 Non-starters: 112.24 ± 71.12	PreS vs. EarS	Starters	>0.999	0.45 [−0.44 to 1.33]
				Non-starters	>0.999	0.26 [−0.58 to 1.10]
	Early-season	Starters: 187.65 ± 55.12 Non-starters: 97.81 ± 23.62	PreS vs. MidS	Starters	>0.999	0.51 [−0.38 to 1.40]
				Non-starters	0.704	0.44 [−0.40 to 1.28]
	Mid-season	Starters: 188.62 ± 41.45 Non-starters: 87.19 ± 31.90	PreS vs. EndS	Starters	>0.999	0.04 [−0.83 to 0.92]
				Non-starters	0.114	0.89 [0.02 to 1.77]
	End-season		EarS vs. MidS	Starters	>0.999	−0.02 [−0.90 to 0.86]
				Non-starters	>0.999	0.36 [−0.48 to 1.21]
			EarS vs. EndS	Starters	0.870	−0.35 [−1.23 to 0.54]
				Non-starters	0.094	1.67 [0.70 to 2.64]
		Starters: 207.53 ± 54.59 Non-starters: 64.75 ± 12.88	MidS vs. EndS	Starters	0.227	−0.37 [−1.26 to 0.51]
				Non-starters	0.072	0.89 [0.01 to 1.76]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, preseason period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ2}, weekly average training monotony based on number of accelerations at Zone 2 (2 to 4 m·s^{−2}); TS_{AcZ2}, weekly average training strain based on number of accelerations at Zone 2 (2 to 4 m·s^{−2}); *p*, *p*-value at alpha level 0.05; Hedges's *g* (95% CI), Hedges's *g* effect size magnitude with 95% confidence interval. Significant differences (*p* ≤ 0.05) are highlighted in bold.

Table 3 and Figure 1e,f demonstrate the intragroup comparisons of TM_{AcZ3} and TS_{AcZ3} between the different periods of the season for both starters and non-starters. The results indicated that there were no meaningful differences in the starters and non-starters (*p* = 0.114, η^2 = 0.288 and *p* = 0.095, η^2 = 0.305, respectively). TS_{AcZ3} analysis showed that there were significant differences between the starters and non-starters (*p* = 0.017, η^2 = 0.441 and *p* = 0.019, η^2 = 0.435, respectively).

The intragroup comparisons of TM_{DcZ1} and TS_{DcZ1} between the different periods of the season for both the starters and non-starters are represented in Table 4 and Figure 2a,b. Analysis of TM_{DcZ1} for the starters (*p* = 0.002, η^2 = 0.582) demonstrated meaningful results but did not show significant differences in the non-starters (*p* = 0.136, η^2 = 0.272). TS_{DcZ1} analyses showed that this was significant for both groups (*p* = 0.018, η^2 = 0.437).

The intragroup comparisons of TM_{DcZ2} and TS_{DcZ2} between the different periods of the season for both the starters and non-starters are represented in Table 5 and Figure 1c,d. The results of the TM_{DcZ2} showed that starters had significant changes (*p* = 0.031, η^2 = 0.397), while no changes were observed in the non-starters (*p* = 0.224, η^2 = 0.221). The TS_{DcZ2} workload displayed meaningful changes for the non-starters (*p* = 0.011, η^2 = 0.469), while no changes (*p* = 0.789, η^2 = 0.058) were shown for the starters.

Table 3. Intragroup differences for binary comparisons between season periods in TM and TS in AcZ3 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Group	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM _{AcZ3} (AU)	Pre-season	Starters: 0.94 ± 0.10 Non-starters: 0.70 ± 0.13	PreS vs. EarS	Starters	>0.999	−0.05 [−0.93 to 0.83]
				Non-starters	0.161	−0.84 [−1.74 to 0.03]
	Early-season	Starters: 0.95 ± 0.14 Non-starters: 0.80 ± 0.11	PreS vs. MidS	Starters	>0.999	−0.39 [−1.28 to 0.49]
				Non-starters	0.678	−0.51 [−1.36 to 0.34]
	Mid-season	Starters: 0.98 ± 0.09 Non-starters: 0.78 ± 0.18	PreS vs. EndS	Starters	0.083	−0.83 [−1.74 to 0.08]
				Non-starters	>0.999	−0.03 [−0.87 to 0.80]
	End-season	Starters: 1.05 ± 0.14 Non-starters: 0.70 ± 0.09	EarS vs. MidS	Starters	>0.999	−0.27 [−1.15 to 0.61]
				Non-starters	>0.999	0.13 [−0.70 to 0.97]
			EarS vs. EndS	Starters	0.258	−0.68 [−1.58 to 0.22]
				Non-starters	0.241	0.96 [0.08 to 1.84]
			MidS vs. EndS	Starters	>0.999	−0.53 [−1.42 to 0.36]
				Non-starters	0.767	0.53 [−0.32 to 1.38]
TS _{AcZ3} (AU)	Pre-season	Starters: 8.09 ± 2.40 Non-starters: 4.86 ± 2.59	PreS vs. EarS	Starters	0.158	−0.71 [−1.96 to 0.19]
				Non-starters	0.056	−0.93 [−1.81 to −0.05]
	Early-season	Starters: 10.17 ± 3.14 Non-starters: 7.24 ± 2.31	PreS vs. MidS	Starters	0.221	−0.73 [−1.63 to 0.18]
				Non-starters	0.432	−0.50 [−1.38 to 0.35]
	Mid-season	Starters: 9.65 ± 1.60 Non-starters: 6.12 ± 2.25	PreS vs. EndS	Starters	0.008	−1.02 [−1.95 to −0.09]
				Non-starters	>0.999	0.24 [−0.60 to 1.08]
	End-season	Starters: 10.61 ± 2.31 Non-starters: 4.35 ± 1.23	EarS vs. MidS	Starters	>0.999	0.20 [−0.68 to 1.08]
				Non-starters	0.857	0.47 [−0.37 to 1.32]
			EarS vs. EndS	Starters	>0.999	−0.15 [−1.03 to 0.72]
				Non-starters	0.011	1.50 [0.56 to 2.45]
			MidS vs. EndS	Starters	>0.999	−0.46 [−1.35 to 0.43]
				Non-starters	0.118	0.93 [0.05 to 1.82]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, preseason period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ3}, weekly average training monotony based on number of accelerations at Zone 2 (>4 m·s^{−2}); TS_{AcZ3}, weekly average training strain based on number of accelerations at Zone 2 (>4 m·s^{−2}); *p*, *p*-value at alpha level 0.05; Hedges's *g* (95% CI), Hedges's *g* effect size magnitude with 95% confidence interval. Significant differences (*p* ≤ 0.05) are highlighted in bold.

Table 4. Intragroup differences for binary comparisons between season periods in TM and TS in DcZ1 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Groups	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM _{DcZ1} (AU)	Pre-season	Starters: 1.82 ± 0.19 Non-starters: 1.31 ± 0.44	PreS vs. EarS	Starters	>0.999	0.07 [−0.81 to 0.95]
				Non-starters	0.626	0.64 [−0.22 to 1.50]
	Early-season	Starters: 1.80 ± 0.33 Non-starters: 1.09 ± 0.16	PreS vs. MidS	Starters	>0.999	−0.65 [−1.55 to 0.25]
				Non-starters	0.176	0.68 [−0.18 to 1.54]
	Mid-season	Starters: 1.98 ± 0.27 Non-starters: 1.06 ± 0.22	PreS vs. EndS	Starters	0.045	−0.99 [−1.98 to −0.06]
				Non-starters	0.202	1.10 [0.21 to 2.00]
	End-season	Starters: 2.33 ± 0.67 Non-starters: 0.94 ± 0.12	EarS vs. MidS	Starters	0.161	−0.57 [−1.46 to 0.33]
				Non-starters	>0.999	0.14 [−0.70 to 0.97]
			EarS vs. EndS	Starters	0.001	−0.96 [−1.88 to −0.03]
				Non-starters	0.966	1.03 [0.14 to 1.92]
			MidS vs. EndS	Starters	0.042	−0.66 [−1.56 to 0.26]
				Non-starters	>0.999	0.67 [−0.19 to 1.53]

Table 4. Cont.

Variables	Period	Mean (SD)	COMPARATIVE	Groups	<i>p</i>	Hedge's <i>g</i> (95% CI)
TS _{DcZ1} (AU)	Pre-season	Starters: 349.37 ± 42.58 Non-starters: 208.15 ± 116.22	PreS vs. EarS	Starters	>0.999	0.54 [−0.35 to 1.43]
				Non-starters	0.415	0.64 [−0.22 to 1.49]
	Early-season	Starters: 315.25 ± 74.37 Non-starters: 151.32 ± 35.13	PreS vs. MidS	Starters	>0.999	0.69 [−0.21 to 1.60]
				Non-starters	0.030	0.95 [0.07 to 1.83]
			PreS vs. EndS	Starters	>0.999	−0.26 [−1.15 to 0.62]
				Non-starters	0.023	1.21 [0.30 to 2.12]
	Mid-season	Starters: 310.75 ± 62.22 Non-starters: 122.72 ± 38.47	EarS vs. MidS	Starters	>0.999	0.06 [−0.81 to 0.94]
				Non-starters	0.591	0.75 [−0.12 to 1.61]
	End-season	Starters: 373.32 ± 114.88 Non-starters: 103.09 ± 22.17	EarS vs. EndS	Starters	0.059	−0.57 [−1.47 to 0.32]
				Non-starters	0.132	1.58 [0.62 to 2.54]
			MidS vs. EndS	Starters	0.018	−0.65 [−1.54 to 0.25]
				Non-starters	>0.999	0.60 [−0.25 to 1.46]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, preseason period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ1}, weekly average training monotony based on number of decelerations at Zone 1 ($< -2 \text{ m} \cdot \text{s}^{-2}$); TS_{AcZ1}, weekly average training strain based on number of decelerations at Zone 1 ($< -2 \text{ m} \cdot \text{s}^{-2}$); *p*, *p*-value at alpha level 0.05; Hedges's *g* (95% CI), Hedges's *g* effect size magnitude with 95% confidence interval. Significant differences ($p \leq 0.05$) are highlighted in bold.

Table 5. Intragroup differences for binary comparisons between season periods in TM and TS in DcZ2 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Groups	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM _{DcZ2} (AU)	Pre-season	Starters: 1.56 ± 0.19 Non-starters: 1.07 ± 0.32	PreS vs. EarS	Starters	<0.999	0.03 [−0.85 to 0.91]
				Non-starters	<0.999	0.40 [−0.44 to 1.24]
	Early-season	Starters: 1.56 ± 0.26 Non-starters: 0.97 ± 0.11	PreS vs. MidS	Starters	<0.999	−0.38 [−1.26 to 0.50]
				Non-starters	<0.999	0.22 [−0.62 to 1.05]
			PreS vs. EndS	Starters	0.467	−0.60 [−1.50 to 0.30]
				Non-starters	0.238	0.77 [−0.09 to 1.64]
	Mid-season	Starters: 1.65 ± 0.24 Non-starters: 1.01 ± 0.17	EarS vs. MidS	Starters	0.725	−0.35 [−1.24 to 0.53]
				Non-starters	<0.999	−0.27 [−1.11 to 0.57]
	End-season	Starters: 1.73 ± 0.33 Non-starters: 0.88 ± 0.12	EarS vs. EndS	Starters	0.093	−0.57 [−1.46 to 0.33]
				Non-starters	0.942	0.78 [−0.08 to 1.65]
			MidS vs. EndS	Starters	<0.999	−0.28 [−1.16 to 0.60]
				Non-starters	0.561	0.87 [−0.001 to 1.75]
TS _{DcZ2} (AU)	Pre-season	Starters: 94.91 ± 24.53 Non-starters: 50.88 ± 29.47	PreS vs. EarS	Starters	<0.999	0.32 [−0.56 to 1.21]
				Non-starters	<0.999	0.24 [−0.60 to 1.08]
	Early-season	Starters: 87.25 ± 20.56 Non-starters: 45.25 ± 13.33	PreS vs. MidS	Starters	<0.999	0.39 [−0.50 to 1.27]
				Non-starters	0.795	0.53 [−0.32 to 1.38]
			PreS vs. EndS	Starters	<0.999	0.26 [−0.62 to 1.14]
				Non-starters	0.055	0.99 [0.10 to 1.87]
	Mid-season	Starters: 86.08 ± 18.99 Non-starters: 38.33 ± 12.96	EarS vs. MidS	Starters	<0.999	0.06 [−0.82 to 0.93]
				Non-starters	0.655	0.51 [−0.34 to 1.36]
	End-season	Starters: 89.15 ± 17.18 Non-starters: 28.96 ± 6.29	EarS vs. EndS	Starters	<0.999	−0.10 [−0.97 to 0.78]
				Non-starters	0.019	1.50 [0.56 to 2.45]
			MidS vs. EndS	Starters	<0.999	−0.16 [−1.04 to 0.72]
				Non-starters	0.332	0.88 [0.01 to 1.76]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, preseason period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ2}, weekly average training monotony based on number of decelerations at Zone 2 (-2 to $-4 \text{ m} \cdot \text{s}^{-2}$); TS_{AcZ2}, weekly average training strain based on number of decelerations at Zone 2 (-2 to $-4 \text{ m} \cdot \text{s}^{-2}$); *p*, *p*-value at alpha level 0.05; Hedges's *g* (95% CI), Hedges's *g* effect size magnitude with 95% confidence interval. Significant differences ($p \leq 0.05$) are highlighted in bold.




Figure 1. Pairwise comparisons between season periods in TM and TS in all AcZ for both non- and starter players. Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; (a): TM_{AcZ1} , weekly average training monotony based on number of accelerations at Zone 1 ($<2 \text{ m}\cdot\text{s}^{-2}$); (b): TS_{AcZ1} , weekly average training strain based on number of accelerations at Zone 1 ($<2 \text{ m}\cdot\text{s}^{-2}$); (c): TM_{AcZ2} , weekly average training monotony based on number of accelerations at Zone 2 ($2 \text{ to } 4 \text{ m}\cdot\text{s}^{-2}$); (d): TS_{AcZ2} , weekly average training strain based on number of accelerations at Zone 2 ($2 \text{ to } 4 \text{ m}\cdot\text{s}^{-2}$); (e): TM_{AcZ3} , weekly average training monotony based on number of accelerations at Zone 2 ($>4 \text{ m}\cdot\text{s}^{-2}$); (f): TS_{AcZ3} , weekly average training strain based on number of accelerations at Zone 2 ($>4 \text{ m}\cdot\text{s}^{-2}$).

Table 6. Intragroup differences for binary comparisons between season periods in TM and TS in DcZ3 for both non- and starter players.

Variables	Period	Mean (SD)	COMPARATIVE	Group	<i>p</i>	Hedge's <i>g</i> (95% CI)
TM_{DcZ3} (AU)	Pre-season	Starters: 1.03 ± 0.11 Non-starters: 0.80 ± 0.18	PreS vs. EarS	Starters	<0.999	$-0.42 [-1.30 \text{ to } 0.47]$
				Non-starters	<0.999	$-0.04 [-0.88 \text{ to } 0.80]$
	Early-season	Starters: 1.08 ± 0.09 Non-starters: 0.81 ± 0.12	PreS vs. MidS	Starters	0.357	$-0.69 [-1.59 \text{ to } 0.21]$
				Non-starters	<0.999	$0.13 [-0.70 \text{ to } 0.97]$
	Mid-season	Starters: 1.14 ± 0.19 Non-starters: 0.78 ± 0.11	PreS vs. EndS	Starters	0.652	$-0.56 [-1.45 \text{ to } 0.34]$
				Non-starters	<0.999	$0.41 [-0.44 \text{ to } 1.25]$
	End-season	Starters: 1.11 ± 0.16 Non-starters: 0.74 ± 0.07	EarS vs. MidS	Starters	0.960	$-0.43 [-1.32 \text{ to } 0.45]$
				Non-starters	<0.999	$0.23 [-0.61 \text{ to } 1.06]$
			EarS vs. EndS	Starters	<0.999	$-0.26 [-1.14 \text{ to } 0.62]$
				Non-starters	0.726	$0.64 [-0.22 \text{ to } 1.50]$
			MidS vs. EndS	Starters	<0.999	$0.18 [-0.70 \text{ to } 1.05]$
				Non-starters	<0.999	$0.38 [-0.47 \text{ to } 1.22]$

Table 6. Cont.

Variables	Period	Mean (SD)	COMPARATIVE	Group	<i>p</i>	Hedge's <i>g</i> (95% CI)
TS _{DcZ3} (AU)	Pre-season	Starters: 15.76 ± 4.63 Non-starters: 10.77 ± 5.99	PreS vs. EarS	Starters	<0.999	−0.50 [−1.39 to 0.39]
				Non-starters	<0.999	0.03 [−0.81 to 0.86]
	Early-season	Starters: 17.88 ± 3.47 Non-starters: 10.64 ± 4.03	PreS vs. MidS	Starters	<0.999	−0.31 [−1.19 to 0.57]
				Non-starters	0.917	0.58 [−0.27 to 1.44]
	Mid-season	Starters: 17.36 ± 5.33 Non-starters: 7.98 ± 2.54	PreS vs. EndS	Starters	<0.999	−0.35 [−1.23 to 0.53]
				Non-starters	0.180	0.89 [0.01 to 1.76]
	End-season	Starters: 17.38 ± 4.22 Non-starters: 6.77 ± 1.30	EarS vs. MidS	Starters	<0.999	0.11 [−0.77 to 0.99]
				Non-starters	0.562	0.76 [−0.11 to 1.62]
			EarS vs. EndS	Starters	<0.999	0.12 [−0.75 to 1.00]
				Non-starters	0.026	1.24 [0.33 to 2.15]
			MidS vs. EndS	Starters	<0.999	−0.004 [−0.88 to 0.87]
				Non-starters	<0.999	0.58 [−0.28 to 1.43]

Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; PreS, preseason period; EarS, early-season period; MidS, mid-season period; EndS, end-season period; TM_{AcZ3}, weekly average training monotony based on number of decelerations at Zone 2 (>−4 m·s^{−2}); TS_{AcZ3}, weekly average training strain based on number of decelerations at Zone 2 (>−4 m·s^{−2}); *p*, *p*-value at alpha level 0.05; Hedges's *g* (95% CI), Hedges's *g* effect size magnitude with 95% confidence interval. Significant differences (*p* ≤ 0.05) are highlighted in bold.




Figure 2. Pairwise comparisons between season periods in TM and TS in all DcZ for both non- and starter players. Abbreviations: TM, training monotony; TS, training strain; AU, arbitrary units; (a): TM_{AcZ1}, weekly average training monotony based on number of decelerations at Zone 1 (<2 m·s^{−2}); (b): TS_{AcZ1}, weekly average training strain based on number of decelerations at Zone 1 (<2 m·s^{−2}); (c): TM_{AcZ2}, weekly average training monotony based on number of decelerations at Zone 2 (2 to 4 m·s^{−2}); (d): TS_{AcZ2}, weekly average training strain based on number of decelerations at Zone 2 (2 to 4 m·s^{−2}); (e): TM_{AcZ3}, weekly average training monotony based on number of decelerations at Zone 2 (>4 m·s^{−2}); (f): TS_{AcZ3}, weekly average training strain based on number of decelerations at Zone 2 (>4 m·s^{−2}).

The intragroup comparisons of TM_{DcZ3} and TS_{DcZ3} between the different periods of the season for both the starters and non-starters are represented in Table 6 and Figure 1e,f. There were no significant changes observed in training workload TM_{DcZ3} in any group. However, non-starters ($p = 0.034$, $\eta p^2 = 0.392$) had significant changes in TS_{DcZ3} .

4. Discussion

This study is the second part of a previously reported research study [13]. Thus, within-group differences in TM and TS in starters and non-starter soccer players at different speed zones across the periods of the full season were examined. The results showed that both starters and non-starters obtained the highest mean TM and TS scores in the pre-season and the highest accelerations at Zone 1. Moreover, the highest TM and TS scores were exhibited in the end-season and the pre-season decelerations at Zone 1 for starter and non-starter players, respectively. Regardless of match starting status, players had similar TM and TS scores at other speed zones and periods.

To the best of the authors' knowledge, there is no previous study on in-season intra-group variations in TM and TS among starters and non-starter soccer players at different speed zones. This makes it difficult to compare the results of the present study with those of earlier reports. However, the results are partly in line with the findings of Fessi et al. [11]. The authors of that study investigated changes in several psychophysical parameters such as the stress, affective valence, quantity of perceived internal training load, quality of sleep, fatigue, and muscle soreness of seventeen professional male soccer players during pre- and in-season periods. Their results demonstrated significantly higher monotony and strain values during the pre-season [11]. Similar results were also observed in a recent study by Clemente et al. [30]. In fact, these authors [30] studied the weekly monotony and strain measures throughout a professional soccer season. They monitored nineteen male players for forty-five weeks by using a GPS device. As a result, they noted greater monotony and strain scores during the pre-season compared to scores exhibited during the season. The higher monotony and strain values during the pre-season might be attributed to a higher perception of effort by the players as a consequence of emphasizing technical/tactical elements and greater training intensity during this period [31,32].

Overall, the results revealed that both starter and non-starter players had greater mean monotony and strain values when they performed acceleration and deceleration at speed Zone 1. In other words, there is an increase in TM and TS at slower speed activities. In another study, Fessi et al. [33] examined the influences of reducing training load on TM and TS in professional male soccer players. They compared weekly training load, monotony, strain, and physical performance indicators of players over seventeen standard and seven taper weeks. Consequently, they found lower monotony and strain values in taper weeks compared to standard weeks. It seems that more research is needed to provide a better understanding of the associations between the level of accelerations and decelerations and the psychophysical perceptions of players.

There are some limitations of the present study that should be addressed. The main limitation is that the data were collected from one professional soccer team, and thus, the study was conducted on small sample size. Furthermore, the study lacks information regarding other important parameters such as physical and technical performance indicators and injury prevalence. Additional studies are warranted to analyze the interactions between the measures of training load, functional capacities, and injury incidence in a larger group of players. For example, analyzing possible inverse relationships between accumulating training load and various physical fitness parameters during the different periods of the season in both male and female professional soccer players will be the topic of future research.

5. Conclusions

In conclusion, this study is the first to provide information on the intragroup differences in wearable inertial measurement unit derived accelerations, decelerations, and

monotony and strain values at different acceleration and deceleration speed zones both starter and non-starter professional soccer players during the different periods of a season. The findings of the study highlighted the variations in TM and TS with regard to the match starting status of players and certain periods of the season. More specifically, regardless of match starting status, at lower acceleration speed zones, the highest mean monotony and strain values were observed during the initial weeks of the season. On the other hand, at lower deceleration speed zones, the highest monotony and strain values were found during the end-season and pre-season for both starter and non-starter players, respectively. Comparisons of within-group values for both starter and non-starter players showed that they had similar monotony and strain values at other speed zones during periods of the season. It is suggested that coaches monitor these two training load indices in order to design optimal individual training programs. Moreover, measuring monotony and strain according to the playing time status of the players may not only help them to facilitate physical and technical development but may also provide valuable information to minimize potential overtraining and injury. Differences in weekly physical loads due to unequal match playing time may be reduced with an adjusted training programs for non-starters.

Author Contributions: Conceptualization, H.N. and J.P.-G.; methodology, H.N., R.O., M.S., K.S. and J.P.-G.; formal analysis, H.N. and H.Z.; investigation, H.N., R.O., H.Z. and J.P.-G.; writing—original draft preparation, H.N., M.S. and J.P.-G.; writing—review and editing, H.N., R.O., M.S., J.P.-G., R.O., K.S. and H.Z. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Ethics Committee of the University of Isfahan.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments: We would like to thank Manuchehr Rezaei, who was responsible for the Sepahan team GPS, for collecting the GPS data for this project. The authors acknowledge Sepahan F.C. for their cooperation, which will increase the level of football science knowledge.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jaspers, A.; Brink, M.S.; Probst, S.G.M.; Frencken, W.G.P.; Helsen, W.F. Relationships Between Training Load Indicators and Training Outcomes in Professional Soccer. *Sports Med.* **2017**, *47*, 533–544. [\[CrossRef\]](#)
2. Oliveira, R.; Brito, J.P.; Loureiro, N.; Padinha, V.; Nobari, H.; Mendes, B. Will Next Match Location Influence External and Internal Training Load of a Top-Class Elite Professional European Soccer Team? *Int. J. Environ. Res. Public Health* **2021**, *18*, 5229. [\[CrossRef\]](#)
3. Halson, S.L. Monitoring Training Load to Understand Fatigue in Athletes. *Sports Med.* **2014**, *44*, S139–S147. [\[CrossRef\]](#)
4. Bourdon, P.C.; Cardinale, M.; Murray, A.; Gastin, P.; Kellmann, M.; Varley, M.C.; Gabbett, T.J.; Coutts, A.J.; Burgess, D.J.; Gregson, W.; et al. Monitoring Athlete Training Loads: Consensus Statement. *Int. J. Sport Physiol. Perform.* **2017**, *12*, 161–170. [\[CrossRef\]](#)
5. Foster, C. Monitoring training in athletes with reference to overtraining syndrome. *Med. Sci. Sports Exerc.* **1998**, *30*, 1164–1168. [\[CrossRef\]](#)
6. Comyns, T.; Flanagan, E.P. Applications of the Session Rating of Perceived Exertion System in Professional Rugby Union. *Strength Cond. J.* **2013**, *35*, 78–85. [\[CrossRef\]](#)
7. Delecroix, B.; McCall, A.; Dawson, B.; Berthoin, S.; Dupont, G. Workload monotony, strain and non-contact injury incidence in professional football players. *Sci. Med. Footb.* **2019**, *3*, 105–108. [\[CrossRef\]](#)
8. Meeusen, R.; Duclos, M.; Foster, C.; Fry, A.; Gleeson, M.; Nieman, D.; Raglin, J.; Rietjens, G.; Steinacker, J.; Urhausen, A. Prevention, diagnosis and treatment of the overtraining syndrome: Joint consensus statement of the European College of Sport Science (ECSS) and the American College of Sports Medicine (ACSM). *Eur. J. Sport Sci.* **2013**, *13*, 1–24. [\[CrossRef\]](#)
9. Clemente, F.M.; Clark, C.; Castillo, D.; Sarmiento, H.; Nikolaidis, P.T.; Rosemann, T.; Knechtle, B. Variations of training load, monotony, and strain and dose-response relationships with maximal aerobic speed, maximal oxygen uptake, and isokinetic strength in professional soccer players. *PLoS ONE* **2019**, *14*, e0225522. [\[CrossRef\]](#) [\[PubMed\]](#)

10. Curtis, R.M.; Huggins, R.A.; Benjamin, C.L.; Sekiguchi, Y.; Arent, S.M.; Armwold, B.C.; Pullara, J.M.; West, C.A.; Casa, D.J. Seasonal Accumulated Workloads in Collegiate Men's Soccer: A Comparison of Starters and Reserves. *J. Strength Cond. Res.* **2019**. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Fessi, M.S.; Nouira, S.; Dellal, A.; Owen, A.; Elloumi, M.; Moalla, W. Changes of the psychophysical state and feeling of wellness of professional soccer players during pre-season and in-season periods. *Res. Sports Med.* **2016**, *24*, 375–386. [\[CrossRef\]](#)
12. Morales, J.; Roman, V.; Yanezi, A.; Solana-Tramunt, M.; Alamo, J.; Figuls, A. Physiological and Psychological Changes at the End of the Soccer Season in Elite Female Athletes. *J. Hum. Kinet.* **2019**, *66*, 99–109. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Nobari, H.; Oliveira, R.; Clemente, F.M.; Adsuar, J.C.; Perez-Gomez, J.; Carlos-Vivas, J.; Brito, J.P. Comparisons of Accelerometer Variables Training Monotony and Strain of Starters and Non-Starters: A Full-Season Study in Professional Soccer Players. *Int. J. Environ. Res. Public Health* **2020**, *17*, 6547. [\[CrossRef\]](#)
14. Kraemer, W.J.; French, D.N.; Paxton, N.J.; Hakkinen, K.; Volek, J.S.; Sebastianelli, W.J.; Putukian, M.; Newton, R.U.; Rubin, M.R.; Gomez, A.L.; et al. Changes in exercise performance and hormonal concentrations over a big ten soccer season in starters and nonstarters. *J. Strength Cond. Res.* **2004**, *18*, 121–128. [\[PubMed\]](#)
15. Silva, J.R.; Magalhaes, J.F.; Ascensao, A.A.; Oliveira, E.M.; Seabra, A.F.; Rebelo, A.N. Individual Match Playing Time during the Season Affects Fitness-Related Parameters of Male Professional Soccer Players. *J. Strength Cond. Res.* **2011**, *25*, 2729–2739. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Oliveira, R.; Palucci Vieira, L.H.; Martins, A.; Brito, J.P.; Nalha, M.; Mendes, B.; Clemente, F.M. In-Season Internal and External Workload Variations between Starters and Non-Starters-A Case Study of a Top Elite European Soccer Team. *Medicina* **2021**, *57*, 645. [\[CrossRef\]](#)
17. Nobari, H.; Castillo, D.; Clemente, F.M.; Carlos-Vivas, J.; Pérez-Gómez, J. Acute, chronic and acute/chronic ratio between starters and non-starters professional soccer players across a competitive season. *Proc. Inst. Mech. Eng. Part P J. Sports Eng. Technol.* **2021**. [\[CrossRef\]](#)
18. Los Arcos, A.; Mendez-Villanueva, A.; Martinez-Santos, R. In-season training periodization of professional soccer players. *Biol. Sport* **2017**, *34*, 149–155. [\[CrossRef\]](#)
19. Nobari, H.; Praca, G.M.; Clemente, F.M.; Perez-Gomez, J.; Vivas, J.C.; Ahmadi, M. Comparisons of new body load and metabolic power average workload indices between starters and non-starters: A full-season study in professional soccer players. *Proc. Inst. Mech. Eng. Part P J. Sports Eng. Technol.* **2021**, *235*, 105–113. [\[CrossRef\]](#)
20. Williams, J.; Tessaro, E. Validity and reliability of a 15 Hz GPS Device for court-based sports movements. *Sport Perform. Sci. Rep.* **2018**, *29*, 1–4.
21. Scott, B.R.; Lockie, R.G.; Knight, T.J.; Clark, A.C.; Janse de Jonge, X.A. A comparison of methods to quantify the in-season training load of professional soccer players. *Int. J. Sports Physiol. Perform.* **2013**, *8*, 195–202. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Nobari, H.; Khalili, S.M.; Oliveira, R.; Castillo-Rodríguez, A.; Pérez-Gómez, J.; Ardigo, L.P. Comparison of Official and Friendly Matches through Acceleration, Deceleration and Metabolic Power Measures: A Full-Season Study in Professional Soccer Players. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5980. [\[CrossRef\]](#)
23. Akenhead, R.; French, D.; Thompson, K.G.; Hayes, P.R. The acceleration dependent validity and reliability of 10 Hz GPS. *J. Sci. Med. Sport* **2014**, *17*, 562–566. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Nobari, H.; Tubagi Polito, L.F.; Clemente, F.M.; Pérez-Gómez, J.; Ahmadi, M.; Garcia-Gordillo, M.; Silva, A.F.; Adsuar, J.C. Relationships Between Training Workload Parameters with Variations in Anaerobic Power and Change of Direction Status in Elite Youth Soccer Players. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7934. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Nobari, H.; Alves, A.R.; Clemente, F.M.; Pérez-Gómez, J.; Clark, C.C.T.; Granacher, U.; Zouhal, H. Associations Between Variations in Accumulated Workload and Physiological Variables in Young Male Soccer Players Over the Course of a Season. *Front. Physiol.* **2021**, *12*, 638180. [\[CrossRef\]](#)
26. Nobari, H.; Barjaste, A.; Haghighi, H.; Clemente, F.M.; Carlos-Vivas, J.; Perez-Gomez, J. Quantification of training and match load in elite youth soccer players: A full-season study. *J. Sports Med. Phys. Fit.* **2021**. [\[CrossRef\]](#)
27. Nobari, H.; Aquino, R.; Clemente, F.M.; Khalafi, M.; Adsuar, J.C.; Pérez-Gómez, J. Description of acute and chronic load, training monotony and strain over a season and its relationships with well-being status: A study in elite under-16 soccer players. *Physiol. Behav.* **2020**, *225*, 113117. [\[CrossRef\]](#) [\[PubMed\]](#)
28. Nobari, H.; Vahabdelshad, R.; Pérez-Gómez, J.; Ardigo, L.P. Variations of Training Workload in Micro- and Meso-Cycles Based on Position in Elite Young Soccer Players: A Competition Season Study. *Front. Physiol.* **2021**, *12*, 668145. [\[CrossRef\]](#)
29. Hopkins, W.G.; Marshall, S.W.; Batterham, A.M.; Hanin, J. Progressive Statistics for Studies in Sports Medicine and Exercise Science. *Med. Sci. Sports Exerc.* **2009**, *41*, 3–12. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Clemente, F.M.; Silva, R.; Ramirez-Campillo, R.; Afonso, J.; Mendes, B.; Chen, Y.S. Accelerometry-based variables in professional soccer players: Comparisons between periods of the season and playing positions. *Biol. Sport* **2020**, *37*, 389–403. [\[CrossRef\]](#)
31. Jeong, T.S.; Reilly, T.; Morton, J.; Bae, S.W.; Drust, B. Quantification of the physiological loading of one week of “pre-season” and one week of “in-season” training in professional soccer players. *J. Sports Sci.* **2011**, *29*, 1161–1166. [\[CrossRef\]](#)
32. Rabelo, F.N.; Pasquarelli, B.N.; Goncalves, B.; Matzenbacher, F.; Campos, F.A.D.; Sampaio, J.; Nakamura, F.Y. Monitoring the Intended and Perceived Training Load of a Professional Futsal Team over 45 Weeks: A Case Study. *J. Strength Cond. Res.* **2016**, *30*, 134–140. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Fessi, M.S.; Zarrouk, N.; Di Salvo, V.; Filetti, C.; Barker, A.R.; Moalla, W. Effects of tapering on physical match activities in professional soccer players. *J. Sports Sci.* **2016**, *34*, 2189–2194. [\[CrossRef\]](#) [\[PubMed\]](#)