



Møller operators and Hadamard states for Dirac fields with MIT boundary conditions

Nicoló Drago, Nicolas Ginoux, Simone Murro

► To cite this version:

Nicoló Drago, Nicolas Ginoux, Simone Murro. Møller operators and Hadamard states for Dirac fields with MIT boundary conditions. 2021. hal-03333715v1

HAL Id: hal-03333715

<https://hal.science/hal-03333715v1>

Preprint submitted on 3 Sep 2021 (v1), last revised 17 Nov 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MØLLER OPERATORS AND HADAMARD STATES FOR DIRAC FIELDS WITH MIT BOUNDARY CONDITIONS

by

Nicoló Drago

Dipartimento di Matematica, Università di Trento, 38050 Povo (TN), Italy

email: nicolo.drago@unitn.it

Nicolas Ginoux

Université de Lorraine, CNRS, IECL, F-57000 Metz, France

email: nicolas.ginoux@univ-lorraine.fr

Simone Murro

Laboratoire de Mathématiques d'Orsay, Université Paris-Saclay, 91405 Orsay, France

email: simone.murro@u-psud.fr

Abstract

The aim of this paper is to prove the existence of Hadamard states for Dirac fields coupled with MIT boundary conditions on any globally hyperbolic manifold with timelike boundary. This is achieved by introducing a geometric Møller operator which implements a unitary isomorphism between the spaces of L^2 -initial data of particular symmetric systems we call weakly-hyperbolic and which are coupled with admissible boundary conditions. In particular, we show that for Dirac fields with MIT boundary conditions, this isomorphism can be lifted to a $*$ -isomorphism between the algebras of Dirac fields and that any Hadamard state can be pulled back along this $*$ -isomorphism preserving the singular structure of its two-point distribution.

Keywords: Møller operators, deformation arguments, Cauchy problem, symmetric weakly-hyperbolic systems, algebraic quantum field theory, Hadamard states, globally hyperbolic manifolds with timelike boundary.

MSC 2020: Primary 35L50, 58J45, 35Q41; Secondary 53C27, 53C50, 81T05.

Contents

1	Introduction	2
2	Møller operators for symmetric weakly-hyperbolic systems	4
2.1	Globally hyperbolic manifolds	4
2.2	Hyperbolic Friedrichs systems of constant characteristic	5
2.3	Admissible boundary conditions	6
2.4	Well-posedness of the Cauchy problem	7
2.5	Symmetric weakly-hyperbolic systems	8

2.6	Møller operators on manifolds with timelike boundary	11
2.7	Conservation of positive definite Hermitian scalar products	17
3	The algebraic approach to quantum Dirac fields	20
3.1	The Dirac operator	20
3.2	Self-adjoint admissible boundary conditions	21
3.3	Linear isometry between spinor bundles	22
3.4	The algebra of Dirac fields with MIT boundary condition	25
3.5	Hadamard states	26

1 Introduction

The initial value problem for a symmetric hyperbolic system on a Lorentzian manifold M is a classical problem which has been exhaustively studied in many contexts. If the underlying background is *globally hyperbolic*, a complete answer is known: In [3] it has been shown that the Cauchy problem is well-posed for any smooth initial data. Even if there exists a plethora of models in physics where globally hyperbolic spacetimes have been used as a background, there also exist many applications which require a manifold with non-empty boundary. Indeed, recent developments in quantum field theory focused their attention on manifolds with timelike boundary [12, 78], e.g. anti-de Sitter spacetime [28, 29] and BTZ spacetime [15]. Moreover, experimental setups for studying the Casimir effect enclose (quantum) fields between walls, which may be mathematically described by introducing timelike boundaries [32, 74]. Also moving walls in a spatial set-up correspond to a timelike boundary in the Lorentzian manifold. For the class of globally hyperbolic manifolds with timelike boundary, the Cauchy problem was investigated by two of us. In particular, we showed in [52] that the Cauchy problem for any symmetric hyperbolic system coupled with an admissible boundary condition is well-posed and the unique solution propagates with at most the speed of light. As a byproduct the existence of a causal propagator is guaranteed. This operator plays a pivotal role in the algebraic approach to quantum field theory since it allows to construct an algebra of observables in a covariant manner, see e.g. [13, 50] for textbooks, [5, 46] for recent reviews and [14, 16, 25–28, 30–32, 42, 43] for some applications. In order to complete the quantization of a free field theory, it is necessary to define an (algebraic) state, *i.e.* a positive functional on the algebra of the observables. Clearly not any state can be considered physically meaningful and, on globally hyperbolic spacetimes with empty boundary, only those satisfying the so-called Hadamard condition are regarded as states of physical interest.

Indeed, within this setting, Hadamard states guarantee the possibility of constructing Wick polynomials following a local and covariant scheme [55, 58], granting also the finiteness of the quantum fluctuations of such Wick polynomials e.g. [41]. Let us remark that in globally hyperbolic spacetimes with empty boundary, the ground state and any KMS states satisfy the Hadamard condition, see e.g. [68, 72]. In close analogy, in the presence of a timelike boundary a generalization of the Hadamard condition has been proposed in [77]. Once the Hadamard condition is assumed, several natural questions arise. The most important one concerns the existence of such states, a problem which was answered positively for free field theories on globally hyperbolic spacetimes with empty boundary (except linearised gravity for which the method cannot be applied – see for example [11]) in [48] by means of a spacetime deformation technique. In detail, once a globally hyperbolic manifold (M, g) is assigned, the key point is to find a (ultra-)static globally hyperbolic metric g_0 on M as well as a Lorentzian metric g_χ interpolating between g and g_0 which is globally hyperbolic. This is not an easy task, since the convex combination of two given globally hyperbolic metric is not in general globally hyperbolic. If the boundary is not empty, the situation gets worse. This is first due to the need for a boundary condition for Cauchy problem. Furthermore, if the interpolating metric is not explicit, then it is not clear how to construct that boundary condition. Hence the arguments used in [48] cannot be applied in a straightforward manner and a new proof has to be thought out.

The aim of this paper is to provide a geometric proof of the existence of Hadamard states for Dirac field with a boundary condition dubbed *MIT boundary condition*. Let us recall that the MIT boundary condition is a local boundary condition which was introduced for the first time in [21] in order to reproduce the confinement of quark in a finite region of space: “Dirac waves” are indeed reflected on the boundary, see also [20, 54] for the description of hadronic states, like baryons and mesons. The MIT boundary condition has been used more recently for many other applications, like the computation of the Casimir energy in a three-dimensional rectangular box [44, 45, 74] in order to construct an integral representation for the Dirac propagator in Kerr Black Hole Geometry and finally also in [59] to prove the asymptotic completeness for linear massive Dirac fields on the Schwarzschild Anti-de Sitter spacetime. A summary of our main result is the following:

Theorem 1.1. *Let $(M, g := -\beta^2 dt^2 + h(t))$ be a globally hyperbolic spin spacetime with timelike boundary and let D be the Dirac operator coupled with MIT boundary condition —cf. Equation (3.3). If there exists an ultrastatic manifold $(M, g_u := -dt^2 + h)$ such that $\beta^{-2}h(t) \leq C^2(t)h$, and if a propagation of singularities result holds for D with MIT boundary conditions —see Theorem 3.16 for the precise statement— then there exists a state for the algebra of Dirac fields with MIT boundary conditions that satisfies the Hadamard condition as per Definition 3.15.*

Remarks 1.2.

1. Let us remark that numerous physical globally hyperbolic spacetimes satisfy the condition of Theorem 1.1, as for example cosmological spacetimes – e.g. de Sitter spacetime or FRWL universe –, geometric blackholes – *i.e.* asymptotically (spacelike) flat spacetimes – and also any globally hyperbolic spacetime with compact Cauchy hypersurface.
2. The propagation of singularities has already been proved in a series of papers, see *e.g.* [9, 29, 49, 60, 63, 64, 75, 76] and we expect that a similar result would apply in our setting as well.

Our strategy to prove the existence of Hadamard states is as follows. In Section 2.5 we introduce the class of symmetric weakly-hyperbolic operators (*cf.* Definition 2.14) extending that of symmetric hyperbolic ones. We show that the Cauchy problem is well-posed for that generalized category of operators (*cf.* Theorem 2.19). Then in Section 2.6 we construct a Møller operator, *i.e.* a geometric map which compares the space of solutions of two given symmetric weakly-hyperbolic systems coupled with admissible boundary conditions on (possibly different) globally hyperbolic manifolds with timelike boundary (*cf.* Theorem 2.27). In particular Section 2.7, we show that this geometric map can be constructed to preserve the natural scalar product defined on the space of solution (*cf.* Proposition 2.33). In Section 3 we specialize ourself to the case of Dirac operators: after introducing the classical Dirac operator and the MIT boundary conditions 3.1 and 3.2 respectively, we construct an isomorphism between spinor bundles defined on different Lorentzian manifolds, see Section 3.3. In Section 3.4 we construct the algebras of Dirac fields and we promote the unitary map between the spaces of solutions of the Dirac equation to a $*$ -isomorphism between the algebras of Dirac fields (*cf.* Theorem 3.11). Finally in Section 3.5 we discuss and prove the existence of Hadamard states for Dirac fields with MIT boundary conditions.

Acknowledgements

We would like to thank Matteo Capoferri, Claudio Dappiaggi, Christian Gérard, Valter Moretti, Miguel Sánchez and Michał Wrochna for helpful discussions related to the topic of this paper.

Funding S.M. is supported by the the DFG research grant MU 4559/1-1 “Hadamard States in Linearized Quantum Gravity”.

Notation and convention

- The symbol \mathbb{K} denotes one of the elements of the set $\{\mathbb{R}, \mathbb{C}\}$.
- $M := (M, g)$ is a globally hyperbolic manifold with timelike boundary ∂M and we adopt the convention that g has the signature $(-, +, \dots, +)$. If g is a Lorentzian metric such that (M, g) is globally hyperbolic, then we shall write $g \in \mathcal{GH}_M$.
- For two Lorentzian metrics g, g' , $g \leq g'$ means that any causal tangent vector for g is causal for g' .
- $t : M \rightarrow \mathbb{R}$ is a Cauchy temporal function and $M_T := t^{-1}(t_0, t_1)$ is a time strip.
- \mathbf{n} is the outward unit normal vector to ∂M .
- $\flat : TM \rightarrow T^*M$ and $\sharp : T^*M \rightarrow TM$ are the musical isomorphisms.
- E is a \mathbb{K} -vector bundle over M with N -dimensional fibers, denoted by E_p for $p \in M$, and endowed with a Hermitian fiber metric $\prec \cdot | \cdot \succ_p$.
- $\Gamma_c(E), \Gamma_{sc}(E)$ *resp.* $\Gamma(E)$ denote the spaces of compactly supported, spacelike compactly supported *resp.* smooth sections of E .
- S is a symmetric weakly-hyperbolic system of constant characteristic coupled with principal symbol denoted by σ_S and B is admissible boundary space for S .
- When M is a Lorentzian spin manifold, we denote with SM the spinor bundle over M and with D the classical Dirac operator.

2 Møller operators for symmetric weakly-hyperbolic systems

The aim of this section is to construct a geometric map, named Møller operators, to compare the solution spaces of two symmetric weakly-hyperbolic operators coupled with admissible boundary conditions on possibly different (though related) globally hyperbolic manifolds with timelike boundary. To this end, we shall first recall the theory of symmetric hyperbolic systems on globally hyperbolic manifolds with timelike boundary. Then, after showing the well-posedness of the Cauchy problem for weakly hyperbolic systems, we shall construct a family of Møller operators depending on the choice of an arbitrary smooth function f . By setting suitably such a function, we shall show that the resulting Møller operator is actually a unitary map between the spaces of initial data endowed with a naturally defined positive scalar product. Our goal is achieved with the help of [52, 66].

2.1 Globally hyperbolic manifolds

Let M be a connected oriented smooth manifold with boundary. We assume M to be endowed with a smooth Lorentzian metric g for which M becomes time-oriented. Here and in the following we shall assume that the boundary is timelike, *i.e.* the pullback of g with respect to the natural inclusion $\iota : \partial M \rightarrow M$ defines a Lorentzian metric ι^*g on the boundary. In the class of Lorentzian manifolds with timelike boundary, those called globally hyperbolic provide a suitable background where to analyze the Cauchy problem for hyperbolic operators.

Definition 2.1. [1, Definition 2.14] A *globally hyperbolic manifold with timelike boundary* is an $(n + 1)$ -dimensional, oriented, time-oriented, smooth Lorentzian manifold M with timelike boundary ∂M such that

- (i) M is causal, *i.e.* there are no closed causal curves;

- (ii) for all points $p, q \in M$, the subset $J^+(p) \cap J^-(q)$ of M is compact, where $J^+(p)$ (*resp.* $J^-(p)$) denotes the causal future (*resp.* past) of p (*resp.* q) in M .

Remark 2.2. In case of an empty boundary, this definition agrees with the standard one, see e.g. [10, Section 3.2] or [6, Section 1.3].

Recently, Aké, Flores and Sánchez gave a characterization of globally hyperbolic manifolds with timelike boundary:

Theorem 2.3 ([1], Theorem 1.1). *Any globally hyperbolic manifold with timelike boundary admits a Cauchy temporal function $t: M \rightarrow \mathbb{R}$ with gradient tangent to ∂M . This implies that M splits into $\mathbb{R} \times \Sigma$ with metric*

$$g = -\beta^2 dt^2 \oplus h(t),$$

where $\beta: \mathbb{R} \times \Sigma \rightarrow \mathbb{R}$ is a smooth positive function, $h(t)$ is a Riemannian metric on each slice $\Sigma_t := \{t\} \times \Sigma$ varying smoothly with t , and these slices are spacelike Cauchy hypersurfaces with boundary $\partial\Sigma_t := \{t\} \times \partial\Sigma$, namely achronal sets intersected exactly once by every inextendible timelike curve.

2.2 Hyperbolic Friedrichs systems of constant characteristic

Let now $E \rightarrow M$ be a Hermitian vector bundle over a globally hyperbolic manifold with timelike boundary M , namely a \mathbb{K} -vector bundle with finite rank N endowed with a positive definite Riemannian or Hermitian fiber metric $\prec \cdot | \cdot \succ_p: E_p \times E_p \rightarrow \mathbb{K}$.

Definition 2.4. A linear differential operator $S: \Gamma(E) \rightarrow \Gamma(E)$ of first order is called a *symmetric system* over M if

- (S) The principal symbol $\sigma_S(\xi): E_p \rightarrow E_p$ is Hermitian with respect to $\prec \cdot | \cdot \succ_p$ for every $\xi \in T_p^*M$ and for every $p \in M$.

Additionally, we say that S is *hyperbolic* respectively *positive* if (H) *resp.* (P) holds, where conditions (H) and (P) are defined as follows:

- (H) For every future-directed timelike covector $\tau \in T_p^*M$, the bilinear form $\prec \sigma_S(\tau) \cdot | \cdot \succ_p$ is positive definite on E_p for every $p \in M$;
- (P) For any Cauchy hypersurface $\Sigma \subset M$, the quadratic form $\phi \mapsto \prec \Re(S^\dagger + S)\phi | \phi \succ$ on Σ is uniformly bounded from below by a positive scalar multiple c_t of the quadratic form $\phi \mapsto \prec \phi | \phi \succ$ which depends continuously on t .

As shown in [52, Section 7.2], the class of symmetric positive systems includes not only hyperbolic PDEs, but also parabolic PDEs, e.g. any diffusion-reaction system with linear reaction term. The latter ones are fundamental in modeling a wide range of phenomena in physics, biology, social sciences, see e.g. [33, 39, 40] but their Cauchy problem is in general not well-posed. The situation improves if we impose that the bilinear or sesquilinear form $\prec \sigma_S(dt) \cdot | \cdot \succ$ is positive definite on E_p for every $p \in M$. This motivates the following definition.

Definition 2.5. We call *hyperbolic Friedrichs system*, any symmetric system S which is hyperbolic or is positive and for which the sesquilinear form $\prec \sigma_S(dt) \cdot | \cdot \succ_p$ is positive definite on E_p , for every $p \in M$. Furthermore, we say that S is *of constant characteristic* if $\dim \ker \sigma_S(n^b)$ is constant. In particular, if $\sigma_S(n^b)$ has maximal rank equal to $\text{rk}(E) = N$ everywhere on ∂M we say that S is *nowhere characteristic*.

Remark 2.6. Notice that, if a system S is hyperbolic with respect to a metric g then it is also hyperbolic with respect to any metric in the conformal class of g . Indeed, conformal changes

preserve each type of covector. Furthermore, Condition (H) implies that for any spacelike covector $\xi \in T_p^*M$ such that $\tau := dt + \xi$ is timelike future-directed,

$$\prec \sigma_S(dt) \cdot | \cdot \succ_p + \prec \sigma_S(\xi) \cdot | \cdot \succ_p = \prec \sigma_S(dt + \xi) \cdot | \cdot \succ_p > 0$$

Therefore, a symmetric system is hyperbolic if and only if:

(H') For every spacelike covector $\xi \in T_p^*M$ such that $dt + \xi$ is a future-directed timelike covector, the bilinear form verifies

$$\prec \sigma_S(\xi) \cdot | \cdot \succ_p > - \prec \sigma_S(dt) \cdot | \cdot \succ_p .$$

2.3 Admissible boundary conditions

In order to discuss the Cauchy problem for a symmetric hyperbolic system, we have to impose suitable boundary conditions. In this paper we shall focus on a class introduced by Friedrichs and Lax-Phillips respectively in [47, 61], dubbed admissible boundary conditions.

Definition 2.7. A smooth linear bundle map $G_B: E|_{\partial M} \rightarrow E|_{\partial M}$ is said to be an *admissible boundary condition* for a symmetric hyperbolic system S if it holds:

- (i) The pointwise kernel B of G_B is a smooth subbundle of $E|_{\partial M}$;
- (ii) The quadratic form $\Psi \mapsto \prec \sigma_S(n^b)\Psi | \Psi \succ_p$ is positive semidefinite on B ;
- (iii) The rank of B is equal to the number of pointwise non-negative eigenvalues of $\sigma_S(n^b)$ counting multiplicity.

The bundle B is called the *boundary space* associated to the boundary condition G_B . Let us remark that Condition (iii) is equivalent to B being *maximal* with respect to property (ii), namely no smooth vector subbundle B' of E exists that properly contains B and such that, for every $\Phi' \in B'$, it holds

$$\prec \sigma_S(n^b)\Phi' | \Phi' \succ \geq 0 .$$

With the next lemma, we shall see that admissible boundary conditions are “stable” under conformal transformations, namely if B is an admissible boundary space for a system on a globally hyperbolic manifold (M, g) , then it is also admissible for the same system on $(M, \Omega^2 g)$, where Ω is a positive smooth function on M .

Lemma 2.8. *Let (M, g) be a globally hyperbolic spacetime with timelike boundary and let B be a boundary space for a hyperbolic Friedrichs system of constant characteristic S . Then B is admissible w.r.t g if and only if it is admissible w.r.t. $\Omega^2 g$, for any positive $\Omega \in C^\infty(M)$.*

Proof. Let denote with n and \tilde{n} the normal vector w.r.t g and $\Omega^2 g$. Since $\tilde{n} = \Omega^{-1}n$, we get $\sigma_S(n) = \Omega \sigma_S(\tilde{n})$. This guarantees conditions (i)–(iii) in Definition 2.7 to be satisfied. \square

Once an admissible boundary condition B is fixed, the *adjoint boundary condition* G_B^\dagger is defined as the pointwise orthogonal projection (with respect to $\prec \cdot | \cdot \succ$) onto $\sigma_S(n^b)(B)$, namely

$$B^\dagger := \ker(G_B^\dagger) = \left(\sigma_S(n^b)(B) \right)^\perp . \quad (2.1)$$

Definition 2.9. An admissible boundary condition B is *self-adjoint* if and only if $B = B^\dagger$.

Remark 2.10. It immediately follows by the definition of a self-adjoint boundary condition that for any $\Psi, \Phi \in B$, it holds

$$\prec \sigma_S(n^b)\Psi | \Phi \succ = 0 .$$

2.4 Well-posedness of the Cauchy problem

Let $t: M \rightarrow \mathbb{R}$ be a Cauchy temporal function with gradient tangent to the boundary, as in Theorem 2.3, and write a symmetric system as

$$S = \sigma_S(dt)\nabla_{\partial_t} - H$$

where H is a first-order linear differential operator which differentiates only in the directions that are tangent to Σ and where ∇ is any fixed metric connection for $\prec \cdot | \cdot \succ$. Let $G_B: E|_{\partial M} \rightarrow E|_{\partial M}$ be an admissible boundary condition for S , in particular its kernel defines the admissible boundary space B .

Definition 2.11. Let S be symmetric system S with positive definite $\prec \sigma_S(dt) \cdot | \cdot \succ$ and let $t_0 \in \mathbb{R}$. We say that $\mathfrak{h} \in \Gamma(E|_{\Sigma_{t_0}})$ and $\mathfrak{f} \in \Gamma(E)$ fulfils the *compatibility condition* of order $k \geq 0$ at time $t_0 \in \mathbb{R}$ if the following condition is satisfied:

$$\sum_{j=0}^k \binom{k}{j} \left(\nabla_{\partial_t}^j G_B \right) \mathfrak{h}_{k-j} \Big|_{\partial \Sigma_{t_0}} = 0, \quad (2.2)$$

where the sequence $(\mathfrak{h}_k)_k$ of sections of $E|_{\partial \Sigma_{t_0}}$ is defined inductively by $\mathfrak{h}_0 := \mathfrak{h}$ and

$$\mathfrak{h}_k := \sum_{j=0}^{k-1} \binom{k-1}{j} H_j \mathfrak{h}_{k-1-j} \Big|_{\partial \Sigma_{t_0}} + \nabla_{\partial_t}^{k-1} (\sigma_S^{-1}(dt)\mathfrak{f}) \Big|_{\partial \Sigma_{t_0}} \quad \text{for all } k \geq 1,$$

where $H_j := [\nabla_{\partial_t}, H_{j-1}]$ and $H_0 := \sigma_S(dt)^{-1}H$.

Roughly speaking, Equation (2.2) provides a sufficient and necessary condition to ensure C^k -regularity for the solution of the Cauchy problem (2.3) once Cauchy data are given on Σ_{t_0} . We recall one of the main results of [52], see [52, Theorem 1.2]:

Theorem 2.12 (Smooth solutions for hyperbolic Friedrichs systems). *Let M be a globally hyperbolic manifold with timelike boundary and let S be a hyperbolic Friedrichs system of constant characteristic. Assume B to be an admissible boundary space for S . Then, for every $\mathfrak{f} \in \Gamma_c(E)$ and $h \in \Gamma_c(E|_{\Sigma_{t_0}})$ satisfying the compatibility condition (2.2) up to any order, there exists a unique $\Psi \in \Gamma(E)$ satisfying the Cauchy problem*

$$\begin{cases} S\Psi = \mathfrak{f} \\ \Psi|_{\Sigma_{t_0}} = \mathfrak{h} \\ \Psi|_{\partial M} \in B \end{cases} \quad (2.3)$$

and the map $(\mathfrak{f}, \mathfrak{h}) \mapsto \Psi$ sending a pair $(\mathfrak{f}, \mathfrak{h}) \in \Gamma_c(E) \times \Gamma_c(E|_{\Sigma_{t_0}})$ to the solution $\Psi \in \Gamma_{sc}(E)$ of (2.3) is continuous.

The assignment $U_{S,t}: D(U_{S,t}) \subset \Gamma_c(E|_{\Sigma_t}) \ni \mathfrak{h} \rightarrow U_{S,t}\mathfrak{h} := \Psi \in \Gamma_{sc}(E)$ of a (unique) solution Ψ to any smooth initial data $\mathfrak{h} \in D(U_{S,t})$ is known as a *Cauchy evolution operator* —here $D(U_{S,t})$ is made by sections $\mathfrak{h} \in \Gamma_c(E|_{\Sigma_t})$ fulfilling the compatibility conditions (2.2) with $\mathfrak{f} = 0$. For later convenience we shall denote by $\rho_t: \Gamma(E) \rightarrow \Gamma(E|_{\Sigma_t})$ the restriction map for smooth sections: Notice that, ρ_t is a right-inverse for $U_{S,t}$. As shown in [16–19], on globally hyperbolic manifolds with empty boundary and compact Cauchy hypersurfaces, the evolution operator can be realized as a Fourier integral operator. As a matter of fact, the Fourier integral representation of the propagator contains the information on how singularities propagates in the manifold. As we shall see in Section 3.5, this is of fundamental importance in proving the existence of Hadamard states for a free quantum field theory on a curved spacetime.

We conclude this section with the following result:

Corollary 2.13. *Let M be a globally hyperbolic manifold with timelike boundary and let B be an admissible boundary space for a symmetric hyperbolic system of constant characteristic S . Then the Cauchy problem for S on (M, g) is well-posed if and only if it is well-posed on $(M, \Omega^2 g)$ for any positive $\Omega \in C^\infty(M)$.*

Proof. Our claim follows immediately by Remark 2.6 and Lemma 2.8. \square

2.5 Symmetric weakly-hyperbolic systems

We conclude this section by showing that the Cauchy problem for a symmetric system S is well-posed also if we assume that the principal symbol $\sigma_S(\xi)$ acts pointwise in a positive definite way only for a subset of future-directed timelike covectors ξ . We begin with the following definition.

Definition 2.14. A symmetric system of constant characteristic over M is *weakly-hyperbolic* if

(wH) There exists a positive smooth function $\alpha = \alpha(t) > 0$ such that for any future-directed timelike covector τ of the form $\tau = dt + \xi \in T_p^*M$ it holds

$$\prec \sigma_S(dt + \alpha\xi) \cdot | \cdot \succ_p > 0.$$

Remarks 2.15.

1. The idea behind the Definition 2.14 is to ‘shrink’ the light cone of the dual metric g^\sharp in the cotangent bundle, so that the condition (H) in Definition 2.4 has to be checked for a smaller class of future-directed timelike covectors (cf. Figure 1). Mind that, in the cotangent bundle, the causal future/past of g_α is not allowed to shrink too much along any Σ_t . Clearly any symmetric hyperbolic system is a symmetric weakly-hyperbolic system,




Figure 1: The future light cones of g^\sharp and g_α^\sharp in the cotangent bundle T^*M .

just take $\alpha = 1$ in Definition 2.14 (cf. Remark 2.6).

2. Assuming the quadratic form $\prec \sigma_S(dt) \cdot | \cdot \succ$ to be pointwise positive definite, there exists at each $p \in M$ an $\alpha(p) > 0$ such that $\prec \sigma_S(\xi) \cdot | \cdot \succ$ is positive definite for every $\xi \in J_\alpha^+(p)$, where $J_\alpha^+(p) \subset T_p^*M$ is the causal future in T_p^*M w.r.t. the metric $g_\alpha := -\beta^2 dt^2 \oplus \alpha(p)^2 h(t)$ defined on $T_p M$, which is hence only defined at p .

The requirement of (wH) is that α only depends on $t(p)$, i.e., that $\alpha|_{\Sigma_t}$ is constant. Mind that this is not equivalent to the quadratic form $\prec \sigma_S(dt) \cdot | \cdot \succ$ being uniformly positive definite on Σ_t , the linearized Euler equation being a counter-example, see e.g. [36].

The following lemma shows that symmetric weakly-hyperbolic systems are not so far from symmetric hyperbolic systems.

Lemma 2.16. *Let S be any symmetric weakly-hyperbolic system on a globally hyperbolic manifold $(M, g) = (\mathbb{R} \times \Sigma, -\beta^2 dt^2 \oplus h(t))$ with or without any timelike boundary. Let $\alpha \in C^\infty(\mathbb{R}, (0, \infty))$ be the function given by definition (wH). Then $(M, g_\alpha) := (\mathbb{R} \times \Sigma, -\beta^2 dt^2 \oplus \alpha^2 h(t))$ is globally hyperbolic and S is symmetric hyperbolic on (M, g_α) .*

Proof. Let $p \in M$ and let $\xi \in T_p^*M$ be g_α -timelike, that is, $g_\alpha^\sharp(\xi, \xi) < 0$ where $g_\alpha^\sharp = -\beta^{-2} \partial_t^{\otimes 2} \oplus \alpha^{-2} h(t)^\sharp$. Then there exists unique $\lambda > 0$ and $\check{\xi} \in T_{\pi_\Sigma(p)}^* \Sigma$ such that $\xi = \lambda \cdot (dt + \alpha \check{\xi})$ —here $\pi_\Sigma: M = \mathbb{R} \times \Sigma \rightarrow \Sigma$ is the standard projection. Condition $g_\alpha^\sharp(\xi, \xi) < 0$ is then equivalent to $g^\sharp(dt + \check{\xi}, dt + \check{\xi}) = g_\alpha^\sharp(dt + \alpha \check{\xi}, dt + \alpha \check{\xi}) < 0$, that is, to $dt + \check{\xi}$ being g -timelike. Then condition (wH) implies that $\sigma_S(dt + \alpha \check{\xi}) = \lambda^{-1} \sigma_S(\xi)$ is positive definite. This shows that S is symmetric hyperbolic on (M, g_α) . Therefore, only the global hyperbolicity of g_α on M remains to be proven. For this, it suffices to show that $\beta^{-2} g_\alpha = -dt^2 \oplus \beta^{-2} \alpha^2 h(t)$ is globally hyperbolic when restricted to any subset of the form $(a, b) \times \Sigma$, with real $a < b$. But since for all such a, b there exists a positive constant α_0 such that $\alpha(t) \geq \alpha_0 > 0$ for all $t \in [a, b]$, we have $\beta^{-2} g_\alpha \leq \beta^{-2} g_{\alpha_0}$ on $[a, b] \times \Sigma$, where $g_{\alpha_0} := -\beta^2 dt^2 \oplus \alpha_0^2 h(t)$. Therefore, it suffices to show that $\beta^{-2} g_{\alpha_0} = -dt^2 \oplus \beta^{-2} \alpha_0^2 h(t)$ is globally hyperbolic on $(a, b) \times \Sigma$. But fixing any $t_0 \in (a, b)$ and any inextensible $\beta^{-2} g_{\alpha_0}$ -timelike curve (which is C^0 and piecewise C^1) $\gamma = (\gamma_0, \hat{\gamma})$ in $(a, b) \times \Sigma$, the curve $\tilde{\gamma} := (\alpha_0^{-1} \gamma_0, \hat{\gamma})$ is $\beta^{-2} g$ -timelike and still inextensible, therefore it meets $\{t_0\} \times \Sigma$ exactly once. This shows $\beta^{-2} g_{\alpha_0}$ and therefore $\beta^{-2} g_\alpha$ (hence g_α) to be globally hyperbolic on $(a, b) \times \Sigma$. This finishes the proof. \square

Example 2.17. Let $M = \mathbb{R} \times \Sigma$ be a globally hyperbolic manifold and let \mathbb{R} be trivial line bundle over M . Then any future directed timelike vector field $X \in \Gamma(TM)$ defines an operator $S := \nabla_X$ which is a symmetric weakly-hyperbolic system if and only if the projection $g(X, \partial_t)^{-1} X_{\Sigma_t}$ is bounded along Σ_t , where X_{Σ_t} denotes the projection of X on Σ_t . This applies in particular for $X = \partial_t + v$, $v \in \Gamma(\Sigma)$, the resulting transport equation being known with the name of Vlasov equation once applied in kinetic theory.

The definition 2.7 of admissible boundary condition can be straightforwardly generalized for a symmetric weakly-hyperbolic system S . The resulting connection with standard hyperbolic systems is described by the following lemma.

Lemma 2.18. *Let B be an admissible boundary space for a symmetric weakly-hyperbolic system S over a globally hyperbolic manifold (M, g) . Then B is admissible for S over (M, g_α) , where g_α is chosen as in Lemma 2.16. Furthermore, if S is of constant characteristic on (M, g) then it is also of constant characteristic of (M, g_α) .*

Proof. To verify our claim is it enough to notice that the unit normal vectors to ∂M are the same up to a strictly positive smooth function. This is due to the choice of a Cauchy temporal function whose gradient is tangent to the boundary. \square

Combining Lemmas 2.16 and 2.18, we can conclude that the Cauchy problem for a symmetric weakly-hyperbolic system S is well-posed. Indeed, these two lemmas guarantee that any smooth solution propagates with at most speed of light (w.r.t. g_α). Therefore, the Cauchy problem can be equivalently reformulated in terms of a Cauchy problem for a symmetric positive system with $\sigma_S(dt) > 0$ in a globally hyperbolic manifold with compact Cauchy surfaces. We summarize our results in the following theorem and we leave the details to the reader.

Theorem 2.19 (Smooth solutions for symmetric weakly-hyperbolic systems). *Let M be a globally hyperbolic spacetime with timelike boundary and let S be a symmetric weakly-hyperbolic system of constant characteristic. Assume G_B to be an admissible boundary condition for S . Then, for every $f \in \Gamma_c(E)$ and $h \in \Gamma_c(E|_{\Sigma_{t_0}})$ satisfying the compatibility condition (2.2) up to any order,*

there exists a unique $\Psi \in \Gamma(\mathbf{E})$ satisfying the Cauchy problem

$$\begin{cases} S\Psi = \mathfrak{f} \\ \Psi|_{\Sigma_{t_0}} = \mathfrak{h} \\ \Psi|_{\partial\mathbf{M}} \in \mathbf{B} \end{cases} \quad (2.4)$$

and the map $(\mathfrak{f}, \mathfrak{h}) \mapsto \Psi$ sending a pair $(\mathfrak{f}, \mathfrak{h}) \in \Gamma_c(\mathbf{E}) \times \Gamma_c(\mathbf{E}|_{\partial\mathbf{M}})$ to the solution $\Psi \in \Gamma_{sc}(\mathbf{E})$ of (2.4), is continuous.

As usual, as a byproduct of the well-posedness of the Cauchy problem, we get the existence of Green operators.

Proposition 2.20. *A symmetric weakly-hyperbolic system of constant characteristic on a globally hyperbolic manifold with timelike boundary coupled with an admissible boundary condition is Green-hyperbolic, i.e. , there exist linear maps, called advanced/retarded Green operator respectively, $G^\pm: \Gamma_c(\mathbf{E}) \rightarrow \Gamma_{sc, \mathbf{B}}(\mathbf{E})$ satisfying*

(i) $S \circ G^\pm \mathfrak{f} = \mathfrak{f}$ for all $\mathfrak{f} \in \Gamma_c(\mathbf{E})$ and $G^\pm \circ S \mathfrak{f} = \mathfrak{f}$ for all $\mathfrak{f} \in \Gamma_{c, \mathbf{B}}(\mathbf{E})$;

(ii) $\text{supp}(G^\pm \mathfrak{f}) \subset J_\alpha^\pm(\text{supp } \mathfrak{f})$ for all $\mathfrak{f} \in \Gamma_c(\mathbf{E})$,

where J_α^\pm denote the causal future (+) and past (−) w.r.t. g_α chosen as in Lemma 2.16 and $\Gamma_{\sharp, \mathbf{B}}(\mathbf{E}) \subset \Gamma_\sharp(\mathbf{E})$, $\sharp \in \{sc, c\}$ denotes the space of smooth sections on \mathbf{E} (with \sharp support property) which fulfil the \mathbf{B} -boundary conditions.

Moreover, let $G := G^+ - G^-: \Gamma_c(\mathbf{E}) \rightarrow \Gamma_{sc, \mathbf{B}}(\mathbf{E})$ be the causal propagator associated with S and \mathbf{B} . Then there exists a short exact sequence

$$0 \rightarrow \Gamma_{c, \mathbf{B}}(\mathbf{E}) \xrightarrow{S} \Gamma_c(\mathbf{E}) \xrightarrow{G} \Gamma_{sc, \mathbf{B}}(\mathbf{E}) \xrightarrow{S} \Gamma_{sc}(\mathbf{E}) \rightarrow 0.$$

In particular the solution space $\text{Sol}_{sc, \mathbf{B}}(S) := \Gamma_{sc, \mathbf{B}}(\mathbf{E}) \cap \ker S$ is characterized as

$$\text{Sol}_{sc, \mathbf{B}}(S) = G\Gamma_c(\mathbf{E}) \simeq \Gamma_c(\mathbf{E})/S\Gamma_{c, \mathbf{B}}(\mathbf{E}). \quad (2.5)$$

Proof. Properties (i) and (ii) are satisfied by definition of G^\pm . As a straightforward consequence, $GS = 0$ on $\Gamma_{c, \mathbf{B}}(\mathbf{E})$ and $SG = 0$ on $\Gamma_c(\mathbf{E})$, therefore the sequence is a complex.

The injectivity of $S|_{\Gamma_{c, \mathbf{B}}(\mathbf{E})}$ immediately follows from property (i) since $Su = 0$ for a $u \in \Gamma_{c, \mathbf{B}}(\mathbf{E})$ yields $u = G^+Su = 0$.

To show that $\ker(G) \subset S(\Gamma_{c, \mathbf{B}}(\mathbf{E}))$, let $u \in \Gamma_c(\mathbf{E})$ with $Gu = 0$. Then $G^+u = -G^-u$, so that $\text{supp } G^+u \subset J_\alpha^+(\text{supp } u) \cap J_\alpha^-(\text{supp } u)$ must be compact by property (ii). Therefore $G^+u \in \Gamma_{c, \mathbf{B}}(\mathbf{E})$ and satisfies $SG^+u = u$ by property (i), from which $u \in S(\Gamma_{c, \mathbf{B}}(\mathbf{E}))$ follows.

To prove that $\ker(S|_{\Gamma_{sc, \mathbf{B}}(\mathbf{E})}) \subset G(\Gamma_c(\mathbf{E}))$, let $u \in \Gamma_{sc, \mathbf{B}}(\mathbf{E})$ be such that $Su = 0$. By definition, there exists a compact subset K of \mathbf{M} such that $\text{supp } u \subset J_\alpha^+(K) \cup J_\alpha^-(K)$. Up to enlarging K a little, we may assume that $\text{supp } u \subset I_\alpha^+(K) \cup I_\alpha^-(K)$, where I_α^+ and I_α^- denote the chronological future and past w.r.t. g_α respectively. Let $\{\chi_+, \chi_-\}$ be a partition of unity subordinated to the open covering $\{I_\alpha^+(K), I_\alpha^-(K)\}$ of $I_\alpha^+(K) \cup I_\alpha^-(K)$. Let $u_\pm := \chi_\pm u$. Then $u = u_+ + u_-$, where each u_\pm is smooth with $\text{supp } u_\pm \subset I_\alpha^\pm(K)$. Furthermore, because u_\pm is obtained by pointwise multiplication of u by a real number, we have $u_\pm|_{\partial\mathbf{M}} \in \mathbf{B}$. Let $v := Su_+ (= -Su_-)$. Then v is smooth with support contained in $J_\alpha^+(K) \cap J_\alpha^-(K)$, therefore $\text{supp } v$ is compact. We check that $Gv = u$ in the weak – and therefore also in the strong – sense. For that, we need the following fact: if $(G^\pm)^*$ denotes the formal adjoint of G^\pm , then actually

$$(G^\pm)^* = G_\dagger^\mp$$

holds, where G_\dagger^\mp are the Green operators for S^\dagger . Recall that, if \mathbf{B} is an admissible boundary condition for S , then so is \mathbf{B}^\dagger for S^\dagger . Moreover, S^\dagger becomes a symmetric weakly hyperbolic

system on M with reversed time orientation, in particular S^\dagger has unique advanced and retarded Green operators as well.

To check that $(G^\pm)^* = G_\mp^\mp$, let φ, ψ be arbitrary in $\Gamma_c(E)$. Because of $\text{supp } G^\pm \varphi \cap \text{supp } G_\mp^\mp \psi$ being compact, we may perform the following partial integration on M :

$$\begin{aligned} \int_M \prec G^\pm \varphi | \psi \succ \text{vol}_M &= \int_M \prec G^\pm \varphi | S^\dagger G_\mp^\mp \psi \succ \text{vol}_M \\ &= \int_M \prec S G^\pm \varphi | G_\mp^\mp \psi \succ \text{vol}_M - \int_{\partial M} \prec \sigma_S(n^b) G^\pm \varphi | G_\mp^\mp \psi \succ \text{vol}_{\partial M} \\ &= \int_M \prec \varphi | G_\mp^\mp \psi \succ \text{vol}_M, \end{aligned}$$

where the boundary term vanishes by $G^\pm \varphi|_{\partial M} \in B$ and $G_\mp^\mp \psi|_{\partial M} \in B^\dagger$. This shows $(G^\pm)^* = G_\mp^\mp$. Now given any $\psi \in \Gamma_c(E)$, we have

$$\begin{aligned} \int_M \prec G^\pm v | \psi \succ \text{vol}_M &= \int_M \prec v | (G^\pm)^* \psi \succ \text{vol}_M \\ &= \int_M \prec v | G_\mp^\mp \psi \succ \text{vol}_M \\ &= \pm \int_M \prec S u_\pm | G_\mp^\mp \psi \succ \text{vol}_M \\ &= \pm \int_M \prec u_\pm | S^\dagger G_\mp^\mp \psi \succ \text{vol}_M \\ &= \pm \int_M \prec u_\pm | \psi \succ \text{vol}_M, \end{aligned}$$

where we have used that $u_\pm|_{\partial M} \in B$ and $G_\mp^\mp \psi|_{\partial M} \in B^\dagger$. This proves that $G^\pm v = \pm u_\pm$ and therefore $Gv = u_+ + u_- = u$, as we claimed.

It remains to establish the surjectivity of $S: \Gamma_{sc,B}(E) \rightarrow \Gamma_{sc}(E)$. Let $f \in \Gamma_{sc}(E)$ be given and $K \subset M$ be compact such that $\text{supp } f \subset J_\alpha^+(K) \cup J_\alpha^-(K)$. As above, up to enlarging K we may assume that $f = f_+ + f_-$, where $f_\pm \in \Gamma_{sc}(E)$ with $\text{supp } f_\pm \subset J_\alpha^\pm(K)$. By Theorem 2.3 the spacetime M can be identified with $\mathbb{R} \times \Sigma$, where Σ a smooth spacelike Cauchy hypersurface of M . For each $n \in \mathbb{N}$ we let $M_{(-n,n)} := (-n, n) \times \Sigma$, where $\Sigma \simeq \{0\} \times \Sigma$. Let χ_n be a smooth function with timelike compact support such that $\chi_n|_{M_{(-n,n)}} = 1$. Then $\chi_n f_+$ lies in $\Gamma_c(E)$ and we may consider $u_n := G^+ \chi_n f_+ \in \Gamma_{sc,B}(E)$. Now letting $u^+(x) := u_n^+(x)$ for every $x \in M_{(-n,n)}$ defines a smooth section of E on M with $Su^+ = f_+$, for if e.g. $m > n$ then $v := u_m^+ - u_n^+$ is a smooth spacelike compact section of E satisfying $Sv = 0$ on $M_{(-n,n)}$ as well as $v|_{M \setminus J^+(\text{supp } (f_+))} = 0$ and $v|_{\partial M_{(-n,n)}} \in B$, so that $v = 0$ on $M_{(-n,n)}$ by uniqueness of the solution of the Cauchy problem. The support of u^+ must be contained in $J_\alpha^+(K)$ since this is the case for the support of each u_n^+ . Analogously, there exists a $u^- \in \Gamma_{sc,B}(E)$ with $Su^- = f_-$ and therefore $S(u^+ + u^-) = f$. This proves the surjectivity of $S: \Gamma_{sc,B}(E) \rightarrow \Gamma_{sc}(E)$ and concludes the proof of Proposition 2.20. \square

For further details we refer to [52, Proposition 5.1], [24, Proposition 20] and [23, Proposition 36].

2.6 Møller operators on manifolds with timelike boundary

In [66] a geometric process was realized to compare solutions of symmetric hyperbolic systems on different globally hyperbolic manifolds $M_0 := (M, g_0)$ and $M := (M, g_1)$ with empty boundary, provided that M_0 and M_1 admit the same Cauchy temporal function and $g_1 \leq g_0$, namely the set of timelike vectors for g_1 is contained in the one for g_0 . This was achieved via the construction of a family of so-called Møller operators [22, 35, 56]. The aim of this paper is to generalize that

construction on manifolds with timelike boundary, where the assumption $g_1 \leq g_0$ is adapted to the situation.

Let us introduce the following setup:

Setup 2.21.

- (i) $M_0 = (M, g_0)$ and $M_1 = (M, g_1)$ are globally hyperbolic manifolds with timelike boundary and with the same Cauchy temporal function $t: M \rightarrow \mathbb{R}$. Moreover, by realizing $(M, g_i) = (\mathbb{R} \times \Sigma, -\beta_i^2 dt^2 \oplus h_i(t))$ for $i = 0, 1$ —cf. Theorem 2.3— we shall assume that there exists a smooth positive function $C = C(t) > 0$ such that

$$\beta_1^{-2} h_1(t) \leq C(t)^2 \beta_0^{-2} h_0(t)$$

holds along Σ_t , for every $t \in \mathbb{R}$;

- (ii) E_1 (resp. E_0) is a \mathbb{K} -vector bundle over M_1 (resp. M_0) with finite rank and endowed with a nondegenerate bilinear or sesquilinear fiber metric $\prec \cdot | \cdot \succ_1$ (resp. $\prec \cdot | \cdot \succ_0$);
- (iii) $\kappa_{1,0}: E_0 \rightarrow E_1$ is a fiberwise linear isometry of vector bundles with inverse $\kappa_{0,1}: E_1 \rightarrow E_0$. With a slight abuse of notation we shall also denote by $\kappa_{1,0}: \Gamma(E_0) \rightarrow \Gamma(E_1)$ the corresponding linear map between sections defined by $[\kappa_{1,0}u](x) = \kappa_{1,0}u(x)$ for all $u \in \Gamma(E_0)$ and $x \in M$. Finally for any positive $f \in C^\infty(M)$, we set $\kappa_{1,0}^f := f \kappa_{1,0}: \Gamma(E_0) \rightarrow \Gamma(E_1)$ with inverse $\kappa_{0,1}^f := f^{-1} \kappa_{0,1}$;
- (iv) S_1 (resp. S_0) is a symmetric weakly-hyperbolic system with admissible boundary space B_1 (resp. B_0). Moreover we shall assume that $\dim \ker \sigma_{S_1}(\xi)$ is constant for any nonzero spacelike covector $\xi \in T^*M_1$;
- (v) Let $S_{0,1}^f: \Gamma(E_1) \rightarrow \Gamma(E_1)$ be the operator defined by $S_{0,1}^f := \kappa_{1,0}^f S_0 \kappa_{0,1}^f$. We assume there exists a linear isometry $\wp_{1,0}: T^*M_0 \rightarrow T^*M_1$ such that $\sigma_{S_{0,1}^f}(\xi) = \sigma_{S_1}(\wp_{1,0}\xi)$ for every $\xi \in T^*M_1$.

Remarks 2.22.

1. The assumption (i) in the Setup 2.21 can be equivalently rephrased as follows. Consider the metric $g_C := -dt^2 \oplus C(t)^2 \beta_1^{-2} h_1(t)$ which is globally hyperbolic on account of Lemma 2.16. Then we get the following two situations: for any vector $v \in TM$, we get

$$(C \leq 1) \quad g_C(v, v) \leq g_0(v, v) \text{ and } g_C(v, v) \leq g_1(v, v), \text{ which implies that } J_{g_C}^\pm \subset J_{g_0}^\pm \text{ and } J_{g_C}^\pm \subset J_{g_1}^\pm.$$


$$(C > 1) \quad g_1(v, v) \leq g_C(v, v) \leq g_0(v, v), \text{ which implies that } J_{g_1}^\pm \subset J_{g_C}^\pm \subset J_{g_0}^\pm.$$

2. Using assumption (iv), assumption (v) implies both that $\dim \ker \sigma_{S_0}(\xi)$ is constant for any nonzero spacelike covector ξ and that $\wp_{1,0}$ is time-orientation preserving.

Remark 2.23 (The principal symbol of $S_{0,1} := S_{0,1}^1$). For later convenience we shall compute the principal symbol $\sigma_{S_{0,1}}$ of $S_{0,1}$ in a slightly more general framework than the one depicted above. (Or maybe this is not at all necessary and we set $\zeta = \text{id}_M$ from the beginning.) Let $E_0 \rightarrow M_0$ and $E_1 \rightarrow M_1$ be two vector bundles such that there exists a diffeomorphism $\zeta_{1,0}: M_0 \rightarrow M_1$ with (inverse $\zeta_{0,1}$) which is lifted to a vector bundle isomorphism $\kappa_{1,0}: E_0 \rightarrow E_1$. With a slight abuse of notation we shall denote with $\kappa_{1,0}: \Gamma(E_0) \rightarrow \Gamma(E_1)$ the associated map of vector bundles defined by

$$(\kappa_{1,0}u_0)(x_1) := \kappa_{1,0}(u_0(\zeta_{0,1}x_1)).$$

for all $u_0 \in \Gamma(E_0)$ and $x_1 \in M_1$. Notice that $\kappa_{1,0}(fu_0) = (\zeta_{0,1}^* f) \kappa_{1,0}u_0$ for all $u_0 \in \Gamma(E_0)$ and $f \in C^\infty(M_0)$ where $\zeta_{0,1}^*: C^\infty(M_0) \rightarrow C^\infty(M_1)$. Moreover, $\kappa_{1,0}: \Gamma(E_0) \rightarrow \Gamma(E_1)$ is invertible with inverse $\kappa_{0,1}$.



(a) $0 < C \leq 1$

(b) $C > 1$

Figure 2: Future light cones of g_0 and g_1 satisfying $\beta_1^{-2}h_1(t) \leq C(t)^2\beta_0^{-2}h_0(t)$.

The principal symbol of $S_{0,1} := \kappa_{1,0}S_0\kappa_{0,1}$ is obtained as follow. For all $u_1 \in E_1|_{x_1}$ and $\xi_1 \in T_{x_1}^*M_1$, let $\tilde{u}_1 \in \Gamma(E_1)$ and $f \in C^\infty(M_1)$ be such that $\tilde{u}_1(x_1) = u_1$ and $df(x_1) = \xi_1$. Then we have

$$\begin{aligned}
\sigma_{S_{0,1}}(\xi_1)u_1 &= [\kappa_{1,0}S_0\kappa_{0,1}, f_1]\tilde{u}_1|_{x_1} = \kappa_{1,0}S_0\kappa_{0,1}f_1\tilde{u}_1 - f_1\kappa_{1,0}S_0\kappa_{0,1}\tilde{u}_1 \\
&= \kappa_{1,0}S_0(\zeta_{1,0}^*f_1\kappa_{0,1}\tilde{u}_1) - f_1\kappa_{1,0}S_0\kappa_{0,1}\tilde{u}_1 \\
&= \kappa_{1,0}[S_0, \zeta_{1,0}^*f_1]\kappa_{0,1}\tilde{u}_1 \\
&= \kappa_{0,1}\sigma_{S_0}(d(\zeta_{1,0}^*f_1))\kappa_{0,1}\tilde{u}_1|_{x_1} \\
&= \kappa_{0,1}\sigma_{S_0}((d\zeta_{1,0}^*df_1)\kappa_{0,1}\tilde{u}_1|_{x_1}) \\
&= \kappa_{0,1}\sigma_{S_0}((d\zeta_{1,0}^*\xi_1)\kappa_{0,1}\tilde{u}_1|_{x_1}),
\end{aligned}$$

where $(d\zeta_{1,0})^*: T^*M_1 \rightarrow T^*M_0$. Overall we have

$$\sigma_{S_{0,1}}(\xi_1) = \kappa_{1,0}\sigma_{S_0}[(d\zeta_{1,0})^*\xi_1]\kappa_{0,1}.$$

Similarly to the case of an empty boundary, the construction of a family of Møller operators requires to control the Cauchy problem for the operator $S_{0,1}^f$. With the next proposition, we shall show that the $S_{0,1}^f$ is actually symmetric weakly-hyperbolic over M_1 .

Proposition 2.24. *Assume the Setup 2.21. Then the operator $S_{0,1}^f$ is a symmetric weakly-hyperbolic system of constant characteristic on M_1 and $\kappa_{1,0}^f(B_0) = \kappa_{1,0}(B_0)$ is an admissible boundary space for $S_{0,1}^f$.*

Proof. On account of Remark 2.23 — with $\zeta_{0,1} = \text{id}_M$ — we have that, for every $\xi \in T^*M$, $\sigma_{S_{0,1}^f}(\xi) = \kappa_{1,0}^f\sigma_{S_0}(\xi)\kappa_{0,1}^f = \kappa_{1,0}\sigma_{S_0}(\xi)\kappa_{0,1}$. Since S_0 is symmetric and $\kappa_{1,0}$ is a fiberwise linear isometry by assumption, $S_{0,1}^f$ clearly satisfies property (S) in Definition 2.4. Moreover, because S_0 has constant characteristic and \mathbf{n}_1^b is a pointwise positive scalar multiple of \mathbf{n}_0^b , the operator $S_{0,1}^f$ has constant characteristic. Because B_0 is an admissible boundary condition for S_0 , the subbundle $\kappa_{1,0}(B_0) = \kappa_{1,0}^f(B_0)$ must be an admissible boundary space for $S_{0,1}^f$. We shall next verify property (wH) in Definition 2.14. To this end let $g_{0,\alpha_0} = -\beta_0^2 dt^2 \oplus \alpha_0^2 h_0(t)$ be the globally hyperbolic metric chosen accordingly with Lemma 2.16. Then S_0 is a symmetric hyperbolic system and $\sigma_{S_0}(\tau)$ is fiberwise positive definite for any future-directed (w.r.t. g_{0,α_0}) covector τ . Since any conformal transformation does not change the set of future-directed covectors, the operator S_0 is hyperbolic w.r.t. $\bar{g}_0 := \beta_0^{-2}g_{0,\alpha_0} = -dt^2 \oplus \alpha_0^2\beta_0^{-2}h_0(t)$. We shall now prove that $S_{0,1}^f$ is symmetric

hyperbolic with respect to the metric $\bar{g}_1 := -dt^2 \oplus \alpha_0^2 C(t)^{-2} \beta_1^{-2} h_1(t)$. For that, let $\tau = dt + \xi$ be \bar{g}_1 -timelike future directed: On account of the assumption $\beta_1^{-2} h_1(t) \leq C^2(t) \beta_0^{-2} h_0(t)$ we find

$$\bar{g}_0^\sharp(dt + \xi, dt + \xi) \leq \bar{g}_1^\sharp(dt + \xi, dt + \xi) < 0,$$

so that $dt + \xi$ is \bar{g}_0 -timelike future directed —notice that $\bar{g}_0^\sharp = \partial_t^{\otimes 2} \oplus \alpha_0^{-2} C(t)^2 \beta_0^2 h_0(t)^\sharp$ and similarly for \bar{g}_1 (cf. Figure 3).




Figure 3: The future light cones of g_1^\sharp and g_0^\sharp in the cotangent bundle T^*M .

It follows that $\sigma_{S_0}(dt + \xi) > 0$ and therefore $\sigma_{S_{0,1}^f}(dt + \xi) > 0$ as well. This shows that $S_{0,1}^f$ is symmetric hyperbolic with respect to \bar{g}_1 and therefore the same holds true for $g_{1,\alpha_1} := \beta_1^2 \bar{g}_1 = -\beta_1^2 dt^2 \oplus \alpha_1^2 h_1(t)$, where $\alpha_1^2 := \alpha_0^2 C(t)^{-2} > 0$ on account of the hypothesis on C and α_0 . This proves that $S_{0,1}^f$ is weakly-hyperbolic with respect to g_1 as per Definition 2.14. \square

Note that the existence of a linear isometry $\wp_{1,0}$ from assumption (v) is not required in the proof of Proposition 2.24.

So far, we considered a setting where the operators S_0, S_1 , though being defined on different bundles, can be compared through $\kappa_{1,0}$. As a matter of fact the next step in the construction of a Møller operator intertwining S_0 and S_1 is to build an intertwining operator between $S_{0,1}^f$ and S_1 . To this avail, we shall first consider an interpolating operator $S_{\chi,1}^f$ defined by $S_{\chi,1}^f := (1 - \chi)S_{0,1}^f + \chi S_1$, where $\chi \in C^\infty(M)$ is an arbitrary smooth function with $0 \leq \chi \leq 1$.

The following proposition ensures that $S_{\chi,1}^f$ is a symmetric weakly-hyperbolic system of constant characteristic as soon as $\wp_{1,0}(\mathbf{n}_1^b)$ is not any pointwise negative scalar multiple of \mathbf{n}_1^b .

Proposition 2.25. *Assume the Setup 2.21 and that $\wp_{1,0}(\mathbf{n}_1^b) \neq \mu \mathbf{n}_1^b$ for any $\mu < 0$. Then for any $\chi \in C^\infty(M, [0, 1])$, the operator defined by*

$$S_{\chi,1}^f := (1 - \chi) S_{0,1}^f + \chi S_1 + \frac{1}{2} \left(\sigma_{S_1} - \sigma_{S_{0,1}^f} \right) (d\chi) : \Gamma(E_1) \rightarrow \Gamma(E_1) \quad (2.6)$$

is a symmetric weakly-hyperbolic system of constant characteristic over M_1 .

Proof. By Remark 2.23 it follows that

$$\sigma_{S_{\chi,1}^f}(\xi) = (1 - \chi) \sigma_{S_{0,1}^f}(\xi) + \chi \sigma_{S_1}(\xi).$$

Therefore, $S_{\chi,1}^f$ is a symmetric system. We shall next notice that a convex combination of weakly-hyperbolic system is still weakly-hyperbolic. As a matter of fact if $\alpha_{0,1}, \alpha_1 \in (0, 1]$ denotes the positive functions of Definition 2.14 associated with $S_{0,1}^f$ and S_1 , it follows that

$$\prec \sigma_{S_{\chi,1}^f}(dt + \alpha \xi) \cdot |\cdot \succ_p > 0,$$

where $\alpha = \min\{\alpha_{0,1}, \alpha_1\}$ for every future-directed g_1 -timelike covector $\tau = dt + \xi$.

To conclude our proof, we shall show that $S_{0,1}^f$ is of constant characteristic. To this end, we consider

$$\sigma_{S_{\chi,1}^f}(\mathbf{n}_1^b) = (1 - \chi)\sigma_{S_{0,1}}(\mathbf{n}_1^b) + \chi\sigma_{S_1}(\mathbf{n}_1^b) = \sigma_{S_1}((1 - \chi)\wp_{1,0}\mathbf{n}_1^b + \chi\mathbf{n}_1^b).$$

Since we have by assumption that $\wp_{1,0}\mathbf{n}_1^b \neq \mu\mathbf{n}_1^b$ for any $\mu < 0$, the covector $(1 - \chi)\wp_{1,0}\mathbf{n}_1^b + \chi\mathbf{n}_1^b$ is non-vanishing along ∂M , which implies that $(1 - \chi)\wp_{1,0}\mathbf{n}_1^b + \chi\mathbf{n}_1^b$ is a nonzero spacelike covector. In particular, by assumption (iv) in Setup 2.21, $S_{\chi,1}^f$ is of constant characteristic. \square

Remarks 2.26.

1. Note that the zero-order operator $V := \frac{1}{2}(\sigma_{S_1} - \sigma_{S_{0,1}^f})(d\chi)$ is a Hermitian operator which vanishes on every open subset where χ is constant — in particular on both the chronological past of Σ_- and the chronological future of Σ_+ . The zero-order operator V does not play any role in the proof of Theorem 2.27. However, whenever S_1, S_0 are formally skew-adjoint, the presence of V ensures that $S_{\chi,1}^f$ is formally skew-adjoint provided a suitable choice of f is made — cf. Proposition 2.32 for the precise statement.
2. The Assumption (iv) in Setup 2.21 is needed in order to ensure that $S_{\chi,1}^f$ is of constant characteristic, and it can be dropped if $\wp_{1,0}\mathbf{n}_1^b = \mathbf{n}_1^b = \mathbf{n}_0^b$.

Building on Proposition 2.25, we now prove the main result of this paper.

Theorem 2.27. *Assume the Setup 2.21 and that $\wp_{1,0}(\mathbf{n}_1^b) \neq \mu\mathbf{n}_1^b$ for any $\mu < 0$. Consider two Cauchy hypersurfaces $\Sigma^\pm \subset M_1$ such that $\Sigma_+ \subset J_{g_1}^+(\Sigma_-)$ — where $J_{g_1}^\pm$ denote the causal cones w.r.t. g_1 — and let $\chi \in C^\infty(M_1, [0, 1])$ be non-decreasing along any future-oriented timelike curve such that*

$$\chi|_{J_{g_1}^+(\Sigma_+)} = 1, \quad \text{and} \quad \chi|_{J_{g_1}^-(\Sigma_-)} = 0.$$

Finally let B_χ be an admissible boundary space for $S_{\chi,1}^f$ such that — cf. Lemma 2.29 —

$$B_\chi = \begin{cases} B_{0,1} := \kappa_{1,0}^f(B_0) & \text{where } \chi = 0 \\ B_1 & \text{where } \chi = 1 \end{cases}.$$

Then the Cauchy problem for $S_{\chi,1}^f$ with B_χ -boundary conditions is well-posed. Moreover, let $U_{S_{\chi,1}^f, \pm}^f: D(U_{S_{\chi,1}^f, \pm}^f) \subset \Gamma_c(E_1|_{\Sigma_\pm}) \rightarrow \Gamma_{sc}(E_1)$ be the Cauchy evolution operator associated with $S_{\chi,1}^f$ and initial data on Σ_\pm and let $\rho_\pm: \Gamma_{sc}(E_1) \rightarrow \Gamma_c(E_1|_{\Sigma_\pm})$ be the standard restriction maps.

Then the Møller operator $R_{0,1} = U_{S_{1,+}} \circ \rho_+ \circ U_{S_{\chi,1}^f, -}^f \circ \rho_- \circ \kappa_{1,0}^f$, implements a non-canonical isomorphism between the spaces of solutions of S_0 and S_1 given by

$$\text{Sol}_{B_\alpha}(S_\alpha) := \{\Psi_\alpha \in \Gamma(E_\alpha) \mid S_\alpha \Psi_\alpha = 0 \text{ and } \Psi_\alpha|_{\partial M} \in B_\alpha\} \quad \text{for } \alpha = 0, 1.$$

Proof. Since B_χ is an admissible boundary space, the Cauchy evolution operators and the Cauchy data map are well-defined on account of Theorem 2.19. Furthermore, for any $\Psi_0 \in \text{Sol}_{B_0}(S_0)$ we have $\rho_- \kappa_{1,0}^f \Psi_0 \in D(U_{S_{\chi,1}^f, -}^f)$ because of B_χ coincide with $\kappa_{1,0}^f(B_0)$ on Σ_- . Therefore $U_{S_{\chi,1}^f, -}^f \rho_- \kappa_{1,0}^f \Psi_0$ is well defined. For a similar reason, for any $\Psi \in \text{Sol}(S_{\chi,1}^f)$ we have $\rho_+ \Psi \in D(U_{S_{1,+}})$. It follows that R is well-defined.

To conclude our proof, it is enough to notice that the Møller operator is a composition of isomorphisms. As such the inverse $R_{1,0}^{-1}$ of $R_{1,0}$ can be computed explicitly as $R_{1,0}^{-1} = \kappa_{0,1}^f \circ U_{S_{0,1}^f, -}^f \circ \rho_- \circ U_{S_{\chi,1}^f, +}^f \circ \rho_+$. \square

Example 2.28. Let (M, g) be a globally hyperbolic spacetime with timelike boundary and let S and \bar{S} be symmetric weakly-hyperbolic systems of constant characteristic which differ by a zero order term, *i.e.* $S - \bar{S} = V$, for $V \in \Gamma(\text{End}(E))$. It follows that any admissible boundary condition B for S is also an admissible boundary condition for \bar{S} . Therefore we can chose $B_\chi = B$.

We conclude this section by showing that for any pair of admissible boundary conditions B, B' for a given symmetric weakly-hyperbolic system there exists an interpolating admissible boundary condition B_χ . This applies in particular to Theorem 2.27 for the choices $V = E_1$, $W_0 = \kappa_{1,0}^f(B_0)$, $W_1 = B_1$.

Lemma 2.29. *Let $V \rightarrow M$ be any smooth vector bundle over any smooth manifold M . Let q be any smooth quadratic form on V . Assume $k := \dim \ker(q)$, the numbers n_+ of positive and n_- of negative pointwise eigenvalues of q to be constant on M . Let $W_0, W_1 \rightarrow M$ be any $n_+ + k$ -dimensional subbundles of V such that $q|_{W_i} \geq 0$ holds pointwise for both $i = 0, 1$.*

Then there exists a smooth map $\phi: [0, 1] \times W_0 \rightarrow V$ such that, for every $t \in [0, 1]$, $\phi_t := \phi(t, \cdot)$ is a linear and injective vector-bundle-map, $q|_{\phi_t(W_0)} \geq 0$ holds pointwise, and $\phi_0 = \text{Id}_{W_0}$ as well as $\phi_1(W_0) = W_1$ are satisfied.

Proof. If we can find a smooth subbundle W'_0 of V such that $W_0 \oplus W'_0 = W_1 \oplus W'_0 = V$ and $q|_{W'_0} \leq 0$ pointwise, then the map ϕ can be constructed as follows. Let π_{W_0} (*resp.* $\pi_{W'_0}$) be the pointwise linear projection onto W_0 with kernel W'_0 (*resp.* onto W'_0 with kernel W_0). Then $\pi_{W_0|W_1}: W_1 \rightarrow W_0$ is an isomorphism by $W_1 \cap W'_0 = \{0\}$ and equality of space dimensions. Let $F := \pi_{W'_0} \circ (\pi_{W_0|W_1})^{-1}: W_0 \rightarrow W'_0$. Observe that, for every $v \in W_0$, we can write

$$v + F(v) = \pi_{W_0} \left((\pi_{W_0|W_1})^{-1}(v) \right) + \pi_{W'_0} \left((\pi_{W_0|W_1})^{-1}(v) \right) = \left(\pi_{W_0|W_1} \right)^{-1}(v),$$

so that $v + F(v) \in W_1$. Now define $\phi: [0, 1] \times W_0 \rightarrow V$ by $\phi(t, v) := v + tF(v)$ for all $(t, v) \in [0, 1] \times W_0$. Clearly ϕ is smooth, $\phi_t = \phi(t, \cdot)$ is a linear injective vector-bundle-map for every $t \in [0, 1]$ because of $W_0 \cap W'_0 = \{0\}$ and obviously $\phi_0 = \text{Id}_{W_0}$ and $\phi_1(W_0) = W_1$ hold by the above observation. Moreover, for any $(t, v) \in [0, 1] \times W_0$,

$$q(v + tF(v), v + tF(v)) = q(v, v) + 2g(v, F(v))t + q(F(v), F(v))t^2.$$

Since the r.h.s of the last identity is a degree-2-polynomial in t , it is non-negative on $[0, 1]$ as soon as it is for $t = 0$ and $t = 1$ and $g(F(v), F(v)) \leq 0$. Therefore, if $q|_{W'_0} \leq 0$, then $q|_{\phi_t(W_0)} \geq 0$.

To construct W'_0 , we make use of the following fact:

Lemma: *Let A be any smooth section of $\text{End}(V)$. If $x \mapsto \dim \ker(A(x))$ is constant on M , then $\ker(A) \rightarrow M$ defines a smooth vector subbundle of V .*

Proof: Fix any Euclidean *resp.* Hermitian inner product on V and let $k := \dim \ker(A(x))$ for all $x \in U$. For any $x \in U$, we have $\ker(A(x)) = \text{ran}(A(x)^*)^\perp$, where $A(x)^*$ is the adjoint of $A(x)$ w.r.t. the chosen inner product on V . Now $\text{ran}(A^*) \rightarrow M$ defines a smooth subbundle of V . Namely it defines an $(n - k)$ -dimensional vector subspace of V at each point of M ; moreover, for any $x_0 \in M$, there exists an open neighbourhood U of x_0 in M and a family of smooth sections v_1, \dots, v_{n-k} of $V|_U$ such that $\{A(x)^*v_1(x), \dots, A(x)^*v_{n-k}(x)\}$ is a family of linearly independent vectors and therefore a basis of $\text{ran}(A(x)^*)$ for any $x \in U$. This shows $\text{ran}(A^*) \rightarrow M$ to be a smooth subbundle of V . As a straightforward consequence, its pointwise orthogonal complement must be a smooth subbundle as well. This proves our claim. \checkmark

It can be deduced from the claim that $\ker(q) \rightarrow M$ defines a smooth subbundle of V . Therefore there exists a smooth supplementary subbundle W to $\ker(q)$. The restriction of q to W defines a smooth nondegenerate quadratic form. Its signature is also constant, namely it is (n_+, n_-) . By e.g. [67, Theorem C.1.4], the bundle W can therefore be split as $W = W_+ \oplus W_-$, where W_\pm are smooth subbundles of W of rank n_\pm and on which q restricts pointwise as a positive-

resp. negative-definite quadratic form. On the whole, we obtain the smooth splitting $V = \ker(q) \oplus W_+ \oplus W_-$. Now $W'_0 := W_-$ does the job since automatically $W_0 \cap W_- = W_1 \cap W_- = \{0\}$ by the fact that $q|_{W_-}$ is pointwise negative definite. This concludes the proof of Lemma 2.29. \square

To apply Lemma 2.29, consider $q := \prec \sigma_{S_1}(\mathbf{n}^b) \cdot | \cdot \succ$ on $V := E_{1|\partial M}$ as well as $W_0 := B_{0,1}$ and $W_1 := B_1$. Then the map $\hat{\phi}$ realizing the interpolation of the boundary conditions is defined by

$$\hat{\phi}: B_{0,1} \rightarrow E_{1|\partial M}, \quad v \mapsto \phi(\chi(\pi(v)), v),$$

where $\pi: E_{1|\partial M} \rightarrow \partial M$ is the footpoint map.

Remark 2.30. Note also that Lemma 2.29 can be applied to the situation where a stronger condition as condition (iv) on the operator S_1 is assumed, namely that the numbers k, n_+, n_- of vanishing, positive resp. negative eigenvalues of $\sigma_{S_1}(\xi)$ are constant along ∂M whenever ξ is a nonvanishing covector in $T^*\Sigma|_{\partial\Sigma}$; then we need not assume any longer that $S_{0,1} = S_1$. This applies for instance to the Dirac operators associated to two different globally hyperbolic metrics g_0, g_1 and where the boundary condition is the MIT one, see Section 3.2 below.

2.7 Conservation of positive definite Hermitian scalar products

Consider now the pre-Hilbert space given by

$$\text{Sol}_{sc,B}(S) = \{\Psi \in \Gamma_{sc}(E) \mid S\Psi = 0, \Psi|_{\partial M} \in B\}$$

where $(\cdot | \cdot)$ is the positive definite Hermitian form defined by

$$(\cdot | \cdot) = \int_{\Sigma} \prec \cdot | \sigma_S(\mathbf{n}^b) \cdot \succ \text{vol}_{\Sigma}, \quad (2.7)$$

where $\mathbf{n} = -\frac{1}{\beta}\partial_t$ is the past-directed unit normal vector to Σ while $\mathbf{n}^b = g(\mathbf{n}, \cdot) = \beta dt$. With the next lemma, we shall show that if S is skew-adjoint then the scalar product (2.7) does not depend on the choice of the Cauchy hypersurface $\Sigma \subset M$.

Lemma 2.31. *Let $\Sigma \subset M$ be a smooth spacelike Cauchy hypersurface with its past-oriented unit normal vector field \mathbf{n} and its induced volume element vol_{Σ} . Furthermore, let S be a formally skew-adjoint, symmetric weakly-hyperbolic system of constant characteristic with self-adjoint admissible boundary condition, i.e. $B = B^\dagger$ —cf. Equation (2.1). Then*

$$(\cdot | \cdot): \text{Sol}_{sc,B}(S) \times \text{Sol}_{sc,B}(S) \rightarrow \mathbb{C} \quad (\Psi | \Phi) = \int_{\Sigma} \prec \Psi | \sigma_S(\mathbf{n}^b) \Phi \succ \text{vol}_{\Sigma},$$

where \mathbf{n}^b denotes here the future-directed unit conormal, yields a positive definite Hermitian scalar product which does not depend on the choice of Σ .

Proof. The proof virtually coincides with the one of [5, Lemma 3.17]. First note that $\text{supp}(\Psi) \cap \Sigma$ is compact since $\text{supp}(\Psi)$ is spacelike compact, so that the integral is well-defined. Let Σ' be any other smooth spacelike Cauchy hypersurface. Without loss of generality we may assume that $\Sigma \cap \Sigma' = \emptyset$, otherwise a third Cauchy hypersurface lying in the common pasts of Σ and Σ' has to be chosen, see proof of [5, Lemma 3.17]. Let $M_T = t^{-1}(\tau, \tau')$ be the time strip such that $t^{-1}(\tau) = \Sigma$ and $t^{-1}(\tau') = \Sigma'$. Its boundary is $\partial M_T = (\partial M \cap M_T) \cup \Sigma \cup \Sigma'$. By the Green identity [52, Lemma 2.11] we have

$$\int_{M_T} (\prec S\Psi | \Phi \succ - \prec \Psi | S^\dagger \Phi \succ) \text{vol}_{M_T} = \int_{\partial M_T} \prec \Psi | \sigma_S(\mathbf{n}^b) \Phi \succ \text{vol}_{\partial M_T}$$

for any $\Psi, \Phi \in \text{Sol}_{sc, B}(S)$. Since S is assumed to be skew-adjoint, the left-hand side of the latter equality vanishes identically. Moreover, since $B = B^\dagger$ also $\prec \Psi | \sigma_S(n^b) \Phi \succ$ vanishes identically at $\partial M \cap M_T$. Therefore we can conclude

$$0 = \int_{\Sigma'} \prec \Psi | \sigma_S(n^b) \Phi \succ \text{vol}_{\Sigma'} - \int_{\Sigma} \prec \Psi | \sigma_S(n^b) \Phi \succ \text{vol}_{\Sigma}.$$

This concludes our proof. \square

With the next lemma, we shall show that there exists a choice of f which makes $S_{\chi,1}^f: \Gamma_{sc, B_\chi}(E_1) \rightarrow \Gamma_{sc}(E_1)$ formally skew-adjoint on $\Gamma_{sc, B_\chi}(E_1)$, provided that B_χ is a self-adjoint boundary condition and S_0 (resp. S_1) are formally skew-adjoint with respect to the pairing $(\cdot | \cdot)_0$ (resp. $(\cdot | \cdot)_1$).

Proposition 2.32. *Assume the setup of Theorem 2.27, that S_0 and S_1 are formally skew-adjoint with respect to the pairings $(\cdot | \cdot)_0$ and $(\cdot | \cdot)_1$ respectively. Finally assume that B_χ is self-adjoint boundary condition for $S_{\chi,1}^f$. If $f \in C^\infty(M)$ is the positive smooth function such that*

$$\text{vol}_{M_0} = f^2 \text{vol}_{M_1}$$

on M , where vol_{M_0} (resp. vol_{M_1}) is the volume form of the metric g_0 (resp. g_1) on M , then $S_{\chi,1}^f$ is a skew-adjoint operator in the Hilbert space \mathcal{H}_1 .

Proof. First we compute the formal adjoint of $S_{0,1}^f$ on (M, g_1) . Let $\Psi_1, \Phi_1 \in \Gamma_c(E_1)$ be such that their supports do not meet ∂M . Since by assumption $f^2 \text{vol}_{M_1} = \text{vol}_{M_0}$ and $S_0^\dagger = -S_0$, we have

$$\begin{aligned} \int_M \prec S_{0,1}^f \Psi_1 | \Phi_1 \succ_1 \text{vol}_{M_1} &= \int_M \prec \kappa_{1,0}^f S_0 \kappa_{0,1}^f \Psi_1 | \Phi_1 \succ_1 \text{vol}_{M_1} \\ &= \int_M \prec \kappa_{1,0}^f S_0 \kappa_{0,1}^f \Psi_1 | \kappa_{1,0}^f \kappa_{0,1}^f \Phi_1 \succ_1 \text{vol}_{M_1} \\ &= \int_M f^2 \prec \kappa_{1,0} S_0 \kappa_{0,1}^f \Psi_1 | \kappa_{1,0} \kappa_{0,1}^f \Phi_1 \succ_1 \text{vol}_{M_1} \\ &= \int_M f^2 \prec S_0 \kappa_{0,1}^f \Psi_1 | \kappa_{0,1}^f \Phi_1 \succ_0 \text{vol}_{M_1} \\ &= \int_M \prec S_0 \kappa_{0,1}^f \Psi_1 | \kappa_{0,1}^f \Phi_1 \succ_0 \text{vol}_{M_0} \\ &= - \int_M \prec \kappa_{0,1}^f \Psi_1 | S_0^\dagger \kappa_{0,1}^f \Phi_1 \succ_0 \text{vol}_{M_0} \\ &= - \int_M \prec \kappa_{0,1}^f \Psi_1 | \kappa_{0,1}^f \kappa_{1,0}^f S_0^\dagger \kappa_{0,1}^f \Phi_1 \succ_0 \text{vol}_{M_0} \\ &= - \int_M f^{-2} \prec \Psi_1 | \kappa_{1,0}^f S_0^\dagger \kappa_{0,1}^f \Phi_1 \succ_1 \text{vol}_{M_0} \\ &= - \int_M \prec \Psi_1 | \kappa_{1,0}^f S_0^\dagger \kappa_{0,1}^f \Phi_1 \succ_1 \text{vol}_{M_1} \\ &= - \int_M \prec \Psi_1 | S_{0,1}^f \Phi_1 \succ_1 \text{vol}_{M_1}, \end{aligned}$$

that is, $(S_{0,1}^f)^\dagger = -S_{0,1}^f$ on (M, g_1) . As a consequence,

$$\begin{aligned} ((1-\chi)S_{0,1}^f + \chi S_1)^\dagger &= (1-\chi)(S_{0,1}^f)^\dagger - \sigma_{S_{0,1}^f}(d(1-\chi)) + \chi S_1^\dagger - \sigma_{S_1}(d\chi) \\ &= -(1-\chi)S_{0,1}^f + \sigma_{S_{0,1}^f}(d\chi) - \chi S_1 - \sigma_{S_1}(d\chi) \\ &= -(1-\chi)S_{0,1}^f - \chi S_1 - 2V, \end{aligned}$$

where V is the zero-order operator defined as above by $V := \frac{1}{2}[\sigma_{S_1}(d\chi) - \sigma_{S_{0,1}^f}(d\chi)]$. Since V is a Hermitian operator it follows that $S_{\chi,1}^f = (1-\chi)S_{0,1}^f + \chi S_1 + V$ is formally skew-adjoint. \square

Building on Lemma 2.31 and Proposition 2.32, we can show that $R_{1,0}$ is a unitary map between $\text{Sol}_{sc,B_0}(S_0)$ and $\text{Sol}_{sc,B_1}(S_1)$.

Proposition 2.33. *Assume the setup of Theorem 2.27, that S_0 and S_1 are formally skew-adjoint and that B_χ is a self-adjoint boundary condition for $S_{\chi,1}^f$. Let $\Sigma_1 \subset J^+(\Sigma_+)$ and $\Sigma_0 \subset J^-(\Sigma_-)$ be fixed spacelike Cauchy hypersurfaces of M (w.r.t. g_0 or g_1 , it makes no difference). Let $f \in C^\infty(M)$ be the positive smooth function such that*

$$\text{vol}_{M_0} = f^2 \text{vol}_{M_1}$$

on M , where vol_{M_0} (resp. vol_{M_1}) is the volume form of the metric g_0 (resp. g_1) on M . Then the Møller operator $R_{1,0}: \text{Sol}_{sc,B_0}(S_0) \rightarrow \text{Sol}_{sc,B_1}(S_1)$ is a unitary map once $\text{Sol}_{sc,B_0}(S_0)$ (resp. $\text{Sol}_{sc,B_1}(S_1)$) is equipped with the scalar product defined in Equation (2.7) associated with S_0 (resp. with S_1).

Proof. Let be $\Psi_0, \Phi_0 \in \text{Sol}_{sc,B_0}(S_0)$ and $\Psi_1 := R_{1,0}(\Psi_0), \Phi_1 := R_{1,0}(\Phi_0) \in \text{Sol}_{sc,B_1}(S_1)$, where the Møller operator $R_{1,0}$ is defined using the interpolating operator $S_{\chi,1}^f$ instead of $S_{\chi,1}^f$. We also denote by $\Psi_{\chi,1}$ (resp. $\Phi_{\chi,1}$) the smooth section with spacelike compact support in $\ker(S_{\chi,1}^f)$ on M with $\Psi_{\chi,1}|_{\Sigma_-} = \kappa_{1,0}^f \Psi_0|_{\Sigma_-}$ (resp. $\Phi_{\chi,1}|_{\Sigma_-} = \kappa_{1,0}^f \Phi_0|_{\Sigma_-}$). By Lemma 2.31 and definition of f , we have $\mathbf{n}_0^b \otimes \text{vol}_{\Sigma_-,g_0} = f^2 \mathbf{n}_1^b \otimes \text{vol}_{\Sigma_-,g_1}$ along Σ_- and therefore

$$\begin{aligned} \int_{\Sigma_0} \prec \sigma_{S_0}(\mathbf{n}_0^b) \Psi_0 | \Phi_0 \succ_0 \text{vol}_{\Sigma_0,g_0} &= \int_{\Sigma_-} \prec \sigma_{S_0}(\mathbf{n}_0^b) \Psi_0 | \Phi_0 \succ_0 \text{vol}_{\Sigma_-,g_0} \\ &= \int_{\Sigma_-} f^{-2} \prec \kappa_{1,0}^f \sigma_{S_0}(\mathbf{n}_0^b) \kappa_{0,1}^f \Psi_0 | \kappa_{1,0}^f \Phi_0 \succ_1 \text{vol}_{\Sigma_-,g_0} \\ &= \int_{\Sigma_-} f^{-2} \prec \sigma_{S_{0,1}^f}(\mathbf{n}_0^b) \Psi_{\chi,1} | \Phi_{\chi,1} \succ_1 \text{vol}_{\Sigma_-,g_0} \\ &= \int_{\Sigma_-} \prec \sigma_{S_{0,1}^f}(\mathbf{n}_1^b) \Psi_{\chi,1} | \Phi_{\chi,1} \succ_1 \text{vol}_{\Sigma_-,g_1} \\ &= \int_{\Sigma_-} \prec \sigma_{S_{\chi,1}^f}(\mathbf{n}_1^b) \Psi_{\chi,1} | \Phi_{\chi,1} \succ_1 \text{vol}_{\Sigma_-,g_1} \\ &= \int_{\Sigma_+} \prec \sigma_{S_{\chi,1}^f}(\mathbf{n}_1^b) \Psi_{\chi,1} | \Phi_{\chi,1} \succ_1 \text{vol}_{\Sigma_+,g_1} \\ &= \int_{\Sigma_+} \prec \sigma_{S_1}(\mathbf{n}_1^b) \Psi_1 | \Phi_1 \succ_1 \text{vol}_{\Sigma_+,g_1}, \end{aligned}$$

which concludes the proof of Proposition 2.33. \square

Definition 2.34. We call *unitary Møller operator* the operator $R_{1,0}$ defined in accordance with Proposition 2.33.

We conclude this section with the following Remark.

Remark 2.35. The unitary Møller operator $R_{1,0}: \text{Sol}_{sc,B_0}(S_0) \rightarrow \text{Sol}_{sc,B_1}(S_1)$ can be seen as the composition of two unitary Møller operators

$$\begin{aligned} R_{\chi,0}: \text{Sol}_{sc,B_0}(S_0) &\rightarrow \text{Sol}_{sc,B_\chi}(S_{\chi,1}^f) & R_{\chi,0} &:= U_{S_{\chi,1}^f,-} \circ \rho_- \circ \kappa_{1,0}^f, \\ R_{1,\chi}: \text{Sol}_{sc,B_\chi}(S_{\chi,1}^f) &\rightarrow \text{Sol}_{sc,B_1}(S_1) & R_{1,\chi} &:= U_{S_1,+} \circ \rho_+. \end{aligned}$$

3 The algebraic approach to quantum Dirac fields

In this section we shall compare the quantization of the Dirac field on two different (yet related) globally hyperbolic spacetimes with timelike boundary. To this end, we shall benefit from [22, 38, 66], where a class of Møller operator was introduced in order to construct unitary equivalent quantum field theories, together of the results of the previous Sections 2.5-2.6.

As a first step we introduce the relevant geometrical objects, showing how they fit within the framework introduced in Section 2.6. In particular we shall apply Theorem 2.27 and Proposition 2.33 for the case of the Dirac operator with MIT boundary conditions —*cf.* Equation (3.3).

3.1 The Dirac operator

Let (M, g) be a globally hyperbolic manifold and assume to have a spin structure *i.e.* a twofold covering map from the $\text{Spin}_0(1, n)$ -principal bundle P_{Spin_0} to the bundle of positively-oriented tangent frames P_{SO^+} of M such that the following diagram is commutative:

$$\begin{array}{ccc} P_{\text{Spin}_0} \times \text{Spin}_0(1, n) & \longrightarrow & P_{\text{Spin}_0} \\ \downarrow & & \downarrow \\ P_{\text{SO}^+} \times \text{SO}(1, n) & \longrightarrow & P_{\text{SO}^+} \longrightarrow M. \end{array}$$

Remark 3.1. Note that unlike differential forms, the definition of a spin structure depends on the metric of the underlying manifold.

The existence of spin structures is related to the topology of M . A sufficient (but not necessary) condition for the existence of a spin structure is the parallelizability of the manifold. Therefore, since any 3-dimensional orientable manifold is parallelizable, it follows by Theorem 2.3 that any 4-dimensional globally hyperbolic manifold admits a spin structure. Given a fixed spin structure, one can use the spinor representation to construct the spinor bundle, *i.e.* the complex vector bundle

$$\mathbb{S}M := \text{Spin}_0(1, n) \times_{\rho} \mathbb{C}^N$$

where $\rho : \text{Spin}_0(1, n) \rightarrow \text{Aut}(\mathbb{C}^N)$ is the complex $\text{Spin}_0(1, n)$ representation and $N := 2^{\lfloor \frac{n+1}{2} \rfloor}$. The spinor bundle comes together with the following structures:

- a natural $\text{Spin}_0(1, n)$ -invariant indefinite fiber metrics

$$\prec \cdot | \cdot \succ_p : \mathbb{S}_p M \times \mathbb{S}_p M \rightarrow \mathbb{C};$$

- a *Clifford multiplication*, *i.e.* a fiber-preserving map

$$\gamma : TM \rightarrow \text{End}(\mathbb{S}M);$$

which satisfies for all $p \in M$, $u, v \in T_p M$ and $\psi, \phi \in \mathbb{S}_p M$

$$\gamma(u)\gamma(v) + \gamma(v)\gamma(u) = -2g(u, v)\text{Id}_{\mathbb{S}_p M} \quad \text{and} \quad \prec \gamma(u)\psi | \phi \succ_p = \prec \psi | \gamma(u)\phi \succ_p.$$

Using the spin product (3.1), we denote as *adjunction map*, the complex anti-linear vector bundle isomorphism by

$$\Upsilon_p : \mathbb{S}_p M_g \rightarrow \mathbb{S}_p^* M_g \quad \psi \mapsto \prec \psi | \cdot \succ, \quad (3.1)$$

where $\mathbb{S}_p^* M_g$ is the so-called *cospinor bundle*, *i.e.* the dual bundle of $\mathbb{S}_p M_g$.

Definition 3.2. The (*classical*) Dirac operator D is the operator defined as the composition of the metric connection $\nabla^{\mathbb{S}}$ on $\mathbb{S}M$, obtained as a lift of the Levi-Civita connection on TM , and the Clifford multiplication:

$$D = \gamma \circ \nabla^{\mathbb{S}M} : \Gamma(\mathbb{S}M) \rightarrow \Gamma(\mathbb{S}M).$$

In local coordinates and with a trivialization of the spinor bundle \mathbb{SM} , the Dirac operator reads as

$$D\psi = \sum_{\mu=0}^n \varepsilon_\mu \gamma(e_\mu) \nabla_{e_\mu}^{\mathbb{SM}} \psi,$$

where $\{e_\mu\}$ is a local Lorentzian-orthonormal frame of \mathbf{TM} and $\varepsilon_\mu = g(e_\mu, e_\mu) = \pm 1$.

Proposition 3.3 ([52, Proposition 6.2]). *The classical Dirac operator D on globally hyperbolic spin manifolds \mathbf{M} with timelike boundary is a nowhere characteristic symmetric hyperbolic system.*

It follows, by Theorem 2.12, that the Cauchy problem for the Dirac operator on globally hyperbolic spacetimes with empty boundary is well-posed, therefore, it admits a Cauchy evolution operator $U_t: \Gamma_c(\mathbb{SM}|_{\Sigma_t}) \rightarrow \Gamma_{sc}(\mathbb{SM})$. Remarkably, as shown by Capoferri and Vassilliev [18], the Cauchy evolution operator for Dirac fields on Cauchy-compact ultrastatic manifolds (with empty boundary) can be realized as a Fourier integral operator. As a matter of fact, the Fourier integral representation of the propagator contains the information on how singularities propagate in the manifolds. For this reason, it would be desirable to extend their techniques to more general globally hyperbolic manifolds with possibly not empty boundary.

3.2 Self-adjoint admissible boundary conditions

The aim of this section is to recast the boundary conditions for the Dirac operator which are self-adjoint and admissible in the sense of Definition 2.7. Let us remark that not all physical interesting boundary conditions for Dirac fields enter in this class of boundary condition. Indeed there exists physically interesting non-local boundary conditions, like the so-called APS boundary condition, which guarantees that the Cauchy problem is well-posed [37], but they are not admissible (since any admissible boundary condition is a local condition). For further details on self-adjoint admissible boundary conditions for the Dirac fields we refer to [52, Section 6.1.1] and [53, Remark 3.19].

The first example of self-adjoint admissible boundary conditions are the so-called *chiral boundary conditions*. They are defined as follows: let \mathcal{G} be a chirality operator on \mathbb{SM} , *i.e.* a parallel involutive antiunitary (with respect to $\prec \cdot | \cdot \succ$) endomorphism-field of \mathbb{SM} that anti-commutes with Clifford multiplication by vectors. Then the so-called *chirality* boundary spaces \mathbf{B}_{CHI} are defined as the range of the maps

$$\pi_{\text{CHI}}^\pm := \frac{1}{2} (\text{Id} \pm \gamma(\mathbf{n})\mathcal{G}), \quad (3.2)$$

where $\gamma(\mathbf{n})$ denotes Clifford multiplication for the outward-pointing unit normal along $\partial\mathbf{M}$. It is not difficult to check that the range of π_{CHI} has dimension $2^{\lfloor \frac{n}{2} \rfloor - 1}$, which is the number of nonnegative eigenvalues of the endomorphism $\sigma_D(\mathbf{n}^b)$, and

$$\prec \sigma_D(\mathbf{n}^b) \pi_{\text{CHI}}^\pm(\psi) | \pi_{\text{CHI}}^\pm \psi \succ = 0.$$

Furthermore, since

$$\pi_{\text{CHI}}^\pm \pi_{\text{CHI}}^\pm = \pi_{\text{CHI}}^\pm, \quad \pi_{\text{CHI}}^\mp \pi_{\text{CHI}}^\pm = 0, \quad \pi_{\text{CHI}}^+ + \pi_{\text{CHI}}^- = \text{Id},$$

it can be easily verified that the boundary conditions are self-adjoint.

The second example of self-adjoint boundary conditions is the so-called *MIT boundary conditions*. The boundary space \mathbf{B}_{MIT} is defined as the range of

$$\pi_{\text{MIT}}^\pm := \frac{1}{2} (\text{Id} \pm \imath \gamma(\mathbf{n})), \quad (3.3)$$

where $\gamma(\mathbf{n})$ is again the Lorentzian Clifford multiplication for the outward-pointing unit normal vector along $\partial\mathbf{M}$. Similarly to the chiral boundary conditions, the range of π_{CHI} has dimension $2^{\lfloor \frac{n}{2} \rfloor - 1}$,

$$\prec \sigma_{\mathbf{D}}(\mathbf{n}^\flat) \pi_{\text{CHI}}^\pm(\psi) | \pi_{\text{CHI}}^\pm \psi \succ = 0$$

and we have

$$\pi_{\text{MIT}}^\pm \pi_{\text{MIT}}^\pm = \pi_{\text{MIT}}^\pm, \quad \pi_{\text{MIT}}^\mp \pi_{\text{MIT}}^\pm = 0, \quad \pi_{\text{MIT}}^+ + \pi_{\text{MIT}}^- = \text{Id}.$$

3.3 Linear isometry between spinor bundles

We shall now apply the results obtained in the Sections 2.6 to compare the solution spaces associated with pairs of Dirac operators $\mathbf{D}_0, \mathbf{D}_1$ defined using different metrics $g_0, g_1 \in \mathcal{GH}_{\mathbf{M}}$ and equipped with admissible self-adjoint boundary conditions. In what follows $g_0, g_1 \in \mathcal{GH}_{\mathbf{M}}$ are assumed to fulfil assumption (i) of Setup 2.21.

As already underlined in Remark 3.1, the space of spinors depends on the metric of the underlying manifold \mathbf{M}_α . Therefore, an identification between spaces of sections of spinor bundles for different metrics is needed to construct a unitary Møller operator. This can be achieved by following [4, Section 5].

Consider a family of Lorentzian spin manifolds $\mathbf{M}_\lambda := (\mathbf{M}, g_\lambda)$ with a common Cauchy temporal function, where $g_\lambda \in \mathcal{GH}_{\mathbf{M}}$ for any $\lambda \in \mathbb{R}$. For a given nonempty interval \mathbf{I} in \mathbb{R} let \mathbf{Z} be the Lorentzian manifold

$$\mathbf{Z} = \mathbf{I} \times \mathbf{M} \quad g_{\mathbf{Z}} = d\lambda^2 + g_\lambda.$$

On \mathbf{Z} there exists a globally defined vector field which we denote as $e_\lambda := \frac{\partial}{\partial \lambda}$. For any λ , the spin structures on \mathbf{Z} and $\mathbf{M}_\lambda \simeq \{\lambda\} \times \mathbf{M}$ are in one-to-one correspondence: Any spin structure on \mathbf{Z} can be restricted to a spin structure on \mathbf{M}_λ and a spin structure on \mathbf{M}_λ it can be pulled back on \mathbf{Z} – see [4, Section 3 and 5]. Actually, the spinor bundle \mathbf{SM}_λ on each globally hyperbolic spin manifold \mathbf{M}_λ can be identified with the restriction of the spinor bundle \mathbf{SZ} on \mathbf{M}_λ , in particular $\mathbf{SM}_\lambda \simeq \mathbf{SZ}|_{\mathbf{M}_\lambda}$ if n is even, while $\mathbf{SM}_\lambda \simeq \mathbb{S}^+ \mathbf{Z}|_{\mathbf{M}_\lambda} \simeq \mathbb{S}^- \mathbf{Z}|_{\mathbf{M}_\lambda}$ if n is odd. Equivalently we may identify

$$\mathbf{SZ}|_{\mathbf{M}_\lambda} = \begin{cases} \mathbf{SM}_\lambda & \text{if } n \text{ is even,} \\ \mathbf{SZ}|_{\mathbf{M}_\lambda} \oplus \mathbf{SZ}|_{\mathbf{M}_\lambda} & \text{if } n \text{ is odd.} \end{cases} \quad (3.4)$$

By denoting with $\gamma_{\mathbf{Z}}$ (*resp.* γ_λ) the Clifford multiplication on \mathbf{SZ} (*resp.* on \mathbf{SM}_λ), the family of Clifford multiplications γ_λ satisfies

$$\gamma_\lambda(v)\psi = \gamma_{\mathbf{Z}}(e_\lambda)\gamma_{\mathbf{Z}}(v)\psi \quad \text{if } n \text{ is even,} \quad (3.5)$$

$$\gamma_\lambda(v)(\psi_+ + \psi_-) = \gamma_{\mathbf{Z}}(e_\lambda)\gamma_{\mathbf{Z}}(v)(\psi_+ - \psi_-) \quad \text{if } n \text{ is odd,} \quad (3.6)$$

where in the second case $\psi = \psi_+ + \psi_- \in \mathbf{SZ}|_{\mathbf{M}_\lambda} \oplus \mathbf{SZ}|_{\mathbf{M}_\lambda}$ and each component ψ_\pm is identified with an element in $\mathbb{S}^\pm \mathbf{Z}|_{\mathbf{M}_\lambda}$.

Lemma 3.4 ([66, Lemma 3.7]). *Let \mathbf{Z} be the Lorentzian spin manifold given by*

$$\mathbf{Z} = \mathbf{I} \times \mathbf{M} \quad g_{\mathbf{Z}} = d\lambda^2 + g_\lambda,$$

where $(\mathbf{M}, g_\lambda) := \mathbf{M}_\lambda$ is a family of Lorentzian spin manifolds with a common Cauchy temporal function, and denote with \mathbf{SM}_λ be the spinor bundle over \mathbf{M}_λ . For any $p \in \mathbf{M}_\lambda$, the map

$$\kappa_{1,0}: \mathbb{S}_p \mathbf{M}_0 \rightarrow \mathbb{S}_p \mathbf{M}_1. \quad (3.7)$$

defined by the parallel translation on \mathbf{Z} along the curve $\lambda \mapsto (\lambda, p)$ is a linear isometry and preserves the Clifford multiplication, i.e. for any $v \in \Gamma(\mathbf{TM})$ and any $\Psi_0 \in \Gamma(\mathbf{SM}_0)$ it holds

$$\gamma_1(\wp_{1,0}v)(\kappa_{1,0}\Psi_0) = \kappa_{1,0}(\gamma_0(v)\Psi_0),$$

where $\wp_{1,0}: \mathbf{TM}_0 \rightarrow \mathbf{TM}_1$ is the parallel transport along the curve $\lambda \mapsto (\lambda, p)$.

Remark 3.5. Let us remark, that for any couple of Lorentzian metric g_0 and g_1 admitting a common Cauchy temporal function, there always exists a path of Lorentzian metric g_λ connecting g_0 to g_1 , e.g. $g_\lambda = \lambda g_1 + (1 - \lambda)g_0$ where $\lambda \in [0, 1]$. For more details we refer to [65, 66].

Lemma 3.4 provides an isomorphism $\kappa_{1,0} : \mathbb{SM} \rightarrow \mathbb{SM}$ with the same properties introduced in the Setup 2.21. We shall denote by $D_{0,1}^f$ the intertwining Dirac operator as in Proposition 2.24. Similarly $D_{\chi,1}^f$ shall denote the operator interpolating between $D_{0,1}^f$ and D_1 . Here and in what follows f is chosen as per Proposition 2.33.

Remark 3.6. Keeping the notation of Remark 2.23 and Lemma 3.4, the diffeomorphism $\zeta : \mathbb{M} \rightarrow \mathbb{M}$ is simply the identity Id . Since $\sigma_{D_0}(\xi) = \gamma_0(\xi^{\sharp_0})$, where \sharp_0 denotes the musical isomorphism with respect to g_0 , we find

$$\sigma_{D_{0,1}^f}(\xi_1) = \kappa_{0,1}^f \sigma_{D_0}(\xi_1) \kappa_{0,1}^f = \kappa_{1,0}^f \gamma_0(\xi_1^{\sharp_0}) \kappa_{0,1}^f = \gamma_1(\wp_{1,0} \xi_1^{\sharp_0}) = \sigma_{D_1}((\wp_{1,0} \xi_1^{\sharp_0})^{\flat_1}) = \sigma_{D_1}(\wp_{1,0} \xi_1),$$

where $\sharp_1 := \flat_1^{-1}$ is the musical isomorphism associated with g_1 . In the last equality we used that, for $\xi \in T_x^* \mathbb{M}$ and $X \in T_x \mathbb{M}$ we have

$$\begin{aligned} (\wp_{1,0} \xi^{\sharp_0})^{\flat_1}(X)|_x &= g_1(\wp_{1,0} \xi^{\sharp_0}, X)|_x = g_Z(\wp_{1,0} \xi^{\sharp_0}, X)|_{(1,x)} = g_Z(\xi^{\sharp_0}, \wp_{0,1} X)|_{(0,x)} \\ &= g_0(\xi^{\sharp_0}, \wp_{0,1} X)|_x = \xi(\wp_{0,1} X)|_x = [\wp_{0,1} \xi](X)|_x, \end{aligned}$$

where, with a slight abuse of notation, we denoted $\wp_{0,1} \xi$ the parallel transport of the 1-form ξ along the curve $\lambda \rightarrow (\lambda, x)$ within \mathbb{Z} : The latter coincides with $\wp_{0,1}^* \xi$, being $\wp_{0,1} : TM_1 \rightarrow TM_0$.

We are almost in position to apply Theorem 2.27 and Proposition 2.33. In the next lemma we shall prove that the assumption in Theorem 2.27 that the parallel transport of \mathbf{n}_1^{\flat} is not proportional to $\mu \mathbf{n}_1^{\flat}$ for any $\mu < 0$, is always satisfied, provided $g_\lambda = (1 - \lambda)g_0 + \lambda g_1$ for all $\lambda \in [0, 1]$.

Lemma 3.7. *Let (\mathbb{M}, g_0) and (\mathbb{M}, g_1) be globally hyperbolic manifolds with timelike boundary split as $(\mathbb{M}, g_i) = (\mathbb{R} \times \Sigma, -\beta_i^2 dt^2 \oplus h_i(t))$ for both $i = 0, 1$. Consider the manifold $\mathbb{Z} := [0, 1] \times \mathbb{M}$ endowed with the metric $g_Z := d\lambda^2 \oplus g_\lambda$, where*

$$g_\lambda := (1 - \lambda)g_0 + \lambda g_1 = -\beta_\lambda dt^2 \oplus h_\lambda(t),$$

where $\beta_\lambda := (1 - \lambda)\beta_0 + \lambda\beta_1$ and $h_\lambda(t) = (1 - \lambda)h_0(t) + \lambda h_1(t)$. Then $h_1(\wp_{1,0}(\mathbf{n}_1), \mathbf{n}_1) > 0$ along $\partial \mathbb{M}$, where $\wp_{1,0}$ is the parallel transport in (\mathbb{Z}, \hat{g}) along $[0, 1] \rightarrow \mathbb{Z}$, $\lambda \mapsto (\lambda, p)$, for any $p \in \partial \mathbb{M}$.

Proof. Note that, by definition of both g_i and of g_Z , we have $\nabla_{\partial_\lambda}^Z \partial_\lambda = \nabla_{\partial_\lambda}^Z \beta_\lambda^{-1} \partial_t = 0$, so that, for any $\lambda_0 \in [0, 1]$, the parallel transport along $[0, \lambda_0] \rightarrow \mathbb{Z}$, $\lambda \mapsto (\lambda, p)$ preserves $T\Sigma$. Writing $p = (t, x)$, we fix a pointwise h_0 -o.n.b. of $T_x \Sigma$ in which $h_1 = h_1(t)$ is diagonal i.e., there exist $\mu_1, \dots, \mu_n > 0$ such that $h_1(e_i, e_j) = \mu_i \delta_{ij}$ for all $1 \leq i, j \leq n$. This basis $(e_i)_{1 \leq i \leq n}$ is extended constantly in λ along $\lambda \mapsto (\lambda, p)$. Splitting $\wp_{\lambda,0} \mathbf{n}_1 = \sum_{j=1}^n \alpha_j e_j$, where $\alpha_j = h_0(\wp_{\lambda,0} \mathbf{n}_1, e_j)$, we have

$$\begin{aligned} 0 &= \nabla_{\partial_\lambda}^Z (\wp_{\lambda,0} \mathbf{n}_1) \\ &= \sum_{j=1}^n (\partial_\lambda \alpha_j) e_j + \alpha_j \nabla_{\partial_\lambda}^Z e_j \\ &= \sum_{j=1}^n (\partial_\lambda \alpha_j) e_j + \alpha_j \left(\underbrace{[\partial_\lambda, e_j]}_0 + \frac{1}{2} h_\lambda^{-1} \partial_\lambda h_\lambda(e_j, \cdot) \right), \end{aligned}$$

so that, denoting by $Y(\lambda) := \begin{pmatrix} \alpha_1(\lambda) \\ \vdots \\ \alpha_n(\lambda) \end{pmatrix}$ and identifying h_λ (as a homomorphism $T\Sigma \rightarrow T^*\Sigma$) and $\partial_\lambda h_\lambda$ (as symmetric 2-tensor on $T\Sigma$) with their respective matrices H_λ and $\partial_\lambda H_\lambda$ in the

bases $(e_j)_{1 \leq j \leq n}$ and $(e_j^*)_{1 \leq j \leq n}$ respectively, the vector-valued function Y must satisfy the linear first-order ODE

$$Y'(\lambda) + \frac{1}{2}H_\lambda^{-1}\partial_\lambda H_\lambda \cdot Y(\lambda) = 0 \quad (3.8)$$

on $[0, 1]$. In case $[H_\lambda, \partial_\lambda H_\lambda] = 0$ is fulfilled for all λ , equation (3.8) can be solved explicitly, namely $Y(\lambda) = H_\lambda^{-\frac{1}{2}} \cdot Y(0)$ for all $\lambda \in [0, 1]$ is the solution with initial condition $Y(0) \in \mathbb{R}^n$. But with $h_\lambda = (1 - \lambda)h_0 + \lambda h_1$, we have $H_\lambda = (1 - \lambda)I_n + \lambda \text{diag}(\mu_1, \dots, \mu_n)$, so that $\partial_\lambda H_\lambda = \text{diag}(\mu_1, \dots, \mu_n) - I_n$ and therefore $[H_\lambda, \partial_\lambda H_\lambda] = 0$ holds for all $\lambda \in [0, 1]$. This implies that

$$Y(\lambda) = H_\lambda^{-\frac{1}{2}} \cdot Y(0) = \text{diag}\left((1 - \lambda + \lambda\mu_1)^{-\frac{1}{2}}, \dots, (1 - \lambda + \lambda\mu_n)^{-\frac{1}{2}}\right) \cdot Y(0)$$

holds for all $\lambda \in [0, 1]$. As a consequence, $Y(1) = \text{diag}\left(\mu_1^{-\frac{1}{2}}, \dots, \mu_n^{-\frac{1}{2}}\right) \cdot Y(0)$, from which

$$h_1(\wp_{1,0}\mathbf{n}_1, \mathbf{n}_1) = H_1(Y(1), Y(0)) = \sum_{j=1}^n \mu_j^{\frac{1}{2}} \alpha_j(0)^2 > 0$$

and the claim follows. \square

We conclude this section by stating Theorem 2.27 and Proposition 2.33 for the particular case of MIT boundary conditions.

Proposition 3.8. *Let assume $g_0, g_1 \in \mathcal{GH}_M$ fulfils (i) in Setup 2.21. Let M_0 (resp. M_1) be a globally hyperbolic spin manifold with timelike boundary and let D_0 (resp. D_1) be a classical Dirac operator coupled with MIT boundary condition B_{MIT_0} (resp. B_{MIT_1}). Then the boundary space defined by*

$$B_\chi = \ker M_\chi := \ker \left(\gamma_1(v) - \imath \|v\|_1 \right)$$

is a self-adjoint boundary space for the operator

$$D_{\chi,1}^f := (1 - \chi) \kappa_{1,0}^f D_0 \kappa_{0,1}^f + \chi D_1 + \frac{1}{2} \left(\sigma_{D_1} + \sigma_{D_{0,1}^f} \right) (d\chi),$$

where $v = \chi \mathbf{n}_1 + (1 - \chi) \wp_{1,0} \mathbf{n}_1$ and $\|v\|_1 = \sqrt{g_1(v, v)}$.

Therefore, letting $\text{Sol}_{sc, MIT}(D_i) := \{\Psi \in \mathbb{SM}_i \mid D_i \Psi = 0, \Psi|_{\partial M} \in B_{MIT}\}$, there exists a unitary isomorphism (Møller operator) $R_{1,0}: \text{Sol}_{sc, MIT}(D_0) \rightarrow \text{Sol}_{sc, MIT}(D_1)$ where $\text{Sol}_{sc, MIT}(D_0)$ (resp. $\text{Sol}_{sc, MIT}(D_1)$) is equipped with the scalar product defined in Equation (2.7) associated with D_0 (resp. with D_1).

Proof. The last part of the statement is nothing but Theorem 2.27 together with Proposition 2.33. Thus, it remains to prove the first part. We begin by noticing that when $\chi = 0$ then B_χ reduces to $\ker(\gamma_1(\wp_{1,0}\mathbf{n}_1) - \imath) = \kappa_{1,0}(B_{MIT_0})$ on account of Remark 3.6, while when $\chi = 1$ B_χ reads as B_{MIT_1} . To conclude we need to show that B_χ is a self-adjoint admissible boundary condition for $D_{\chi,1}^f$. To this end, we first notice that the bilinear form $\prec \gamma_1(v)\Psi \mid \Psi \succ_q$ satisfies simultaneously

$$\begin{aligned} \prec \gamma_1(v)\Psi \mid \Psi \succ_q &= \prec \Psi \mid \gamma_1(v)\Psi \succ_q = \overline{\prec \gamma_1(v)\Psi \mid \Psi \succ_q} \\ \prec \gamma_1(v)\Psi \mid \Psi \succ_q &= \prec \imath \|v\| \Psi \mid \Psi \succ_q = \prec \Psi \mid -\imath \|v\| \Psi \succ_q = - \prec \Psi \mid \gamma_1(v)\Psi \succ_q = - \overline{\prec \gamma_1(v)\Psi \mid \Psi \succ_q} \end{aligned}$$

which implies that $\prec \gamma_1(v)\Psi \mid \Psi \succ_q = \prec \gamma_{D_{\chi,1}^f}(\mathbf{n}_1^b)\Psi \mid \Psi \succ_q = 0$. Furthermore the range of the projector $\pi = \frac{1}{2}(\text{Id} - \imath \|v\|_1^{-1} \gamma_1(v))$ which is equal to B_χ , has dimension $2^{\lfloor \frac{n}{2} - 1 \rfloor}$, which is exactly the number of nonnegative eigenvalues of $\gamma_{D_{\chi,1}^f}(\mathbf{n}_1^b)$. This concludes our proof. \square

Remarks 3.9.

1. Since, for any nonzero spacelike covector v on M , the operator $\sigma_D(v)$ has vanishing kernel and $\pm|v|$ as nonvanishing eigenvalues, each with multiplicity $2^{\lfloor \frac{n+1}{2} \rfloor - 1}$, the existence of an interpolating B_χ between B_{MIT_0} and B_{MIT_1} for D_0 and D_1 respectively follows from Lemma 2.29, see Remark 2.30 above.
2. As a consequence, a Møller operator can be constructed between the solution spaces to D_0 and D_1 , where the solutions of D_0 satisfy the chiral boundary conditions and D_1 the MIT ones. The reason is, such a Møller operator exists for a fixed Dirac operator and between both mentioned boundary conditions, and another Møller operator exists between the MIT boundary conditions for the two different Dirac operators D_0 and D_1 . Therefore, a Møller operator can be obtained composing three Møller operators obtained as described above.

3.4 The algebra of Dirac fields with MIT boundary condition

In this section we shall exploit Proposition 3.8 to compare the quantization of the Dirac field with MIT boundary condition on M_0 and M_1 . With this purpose we shall briefly recall the quantization procedure from the algebraic point of view.

In [28, 38, 66], the quantization of a free field theory is realized as a two-step procedure. On the one hand, the physical system classically described by $\text{Sol}_{sc, MIT}(D)$ is quantized by introducing a unital $*$ -algebra \mathfrak{A} , whose elements are interpreted as observables for the system under investigation. In a second stage, the description of possible physical states of the system is described through the choice of a suitable subclass of linear, positive and normalized functionals $\omega: \mathfrak{A} \rightarrow \mathbb{C}$.

By extending the analogous definition for a spacetime without boundary, we shall now introduce the $*$ -algebra \mathfrak{A} associated with the space $\text{Sol}_{sc, MIT}(D)$ of solutions with spatially compact support of the Dirac operator D coupled with MIT boundary conditions and endowed with the positive definite Hermitian scalar product (2.7).

To this avail we shall profit of the results and definition already present in the literature, see [2]. For later convenience let $\text{Sol}_{sc, MIT}^\oplus$ be the Hilbert space obtained by completion of

$$\text{Sol}_{sc, MIT}(D) \oplus \Upsilon \text{Sol}_{sc, MIT}(D),$$

equipped with the natural scalar product $(\cdot, \cdot)_{\text{Sol}_{sc, MIT}^\oplus}$ induced by $\text{Sol}_{sc, MIT}(D)$ —cf. Equation (2.7)— in particular $(\psi_1 | \psi_2) = \int_\Sigma \prec \psi_1 | \gamma(-\beta^{-1} \partial_t) \psi_2 \succ \text{vol}_\Sigma$. Moreover, let $\Gamma: \text{Sol}_{sc, MIT}^\oplus \rightarrow \text{Sol}_{sc, MIT}^\oplus$ be the antilinear involution defined by $\Gamma(\psi_1 \oplus \Upsilon \psi_2) := (-\psi_2) \oplus \Upsilon \psi_1$ where $\Upsilon: SM \rightarrow S^*M$ has been defined in Equation (3.1).

Definition 3.10. The *algebra of Dirac fields with MIT boundary condition* is the unital, complex $*$ -algebra \mathfrak{A} freely generated by the abstract elements $\Xi(\psi)$, $1_\mathfrak{A}$, with $\psi \in \text{Sol}_{sc, MIT}^\oplus$, together with the following relations for all $\psi, \phi \in \text{Sol}_{sc, MIT}^\oplus$ and $\alpha, \beta \in \mathbb{C}$:

- (i) Linearity: $\Xi(\alpha\psi + \beta\phi) = \alpha\Xi(\psi) + \beta\Xi(\phi)$
- (ii) Hermiticity: $\Xi(\psi)^* = \Xi(\Gamma\psi)$
- (iii) Canonical anti-commutation relations (CARs):

$$\Xi(\psi) \cdot \Xi(\phi) + \Xi(\phi) \cdot \Xi(\psi) = 0 \quad \text{and} \quad \Xi(\psi) \cdot \Xi(\phi)^* + \Xi(\phi)^* \cdot \Xi(\psi) = (\psi | \phi) 1_\mathfrak{A}.$$

As a matter of fact \mathfrak{A} can be completed in a unique way into a C^* -algebra [2] the C^* -norm being induced by the natural Hilbert structure of $\text{Sol}_{sc, MIT}^\oplus$. Occasionally we shall implicitly regard \mathfrak{A} as a C^* -algebra.

Recollecting the results of the previous sections we have the following:

Theorem 3.11. Assume that $g_0, g_1 \in \mathcal{GH}_M$ fulfils (i) in the Setup 2.21 and let \mathfrak{A}_α be the algebra of Dirac fields with MIT boundary condition on M_α . Then there exists a $*$ -isomorphism $\mathfrak{A}_{1,0}: \mathfrak{A}_0 \rightarrow \mathfrak{A}_1$.

Proof. Let $\Upsilon_\alpha : \mathbb{S}M_\alpha \rightarrow \mathbb{S}^*M_\alpha$ the adjunction map defined in (3.1) between the spinor and cospinor bundle over M_α and set $R_{1,0}^\Upsilon := \Upsilon_1 R_{1,0} \Upsilon_0^{-1}$. Then $R_{1,0}^\Upsilon$ implements an isomorphism between $\Upsilon_0 \text{Sol}_{sc, \text{MIT}}(D_0)$ and $\Upsilon_1 \text{Sol}_{sc, \text{MIT}}(D_1)$. On account of Proposition 3.8 $R_{1,0}^\oplus := R_{1,0} \oplus R_{1,0}^\Upsilon : \text{Sol}_{sc, \text{MIT}}^\oplus \rightarrow \text{Sol}_{sc, \text{MIT}}^\oplus$ is a unitary isomorphism. By direct inspection, the linear map $\mathfrak{R}_{1,0} : \mathfrak{A}_0 \rightarrow \mathfrak{A}_1$ defined by $\mathfrak{R}_{1,0} \Xi(\psi) := \Xi(R_{1,0}^\oplus \psi)$ extends to the seen $*$ -isomorphism. \square

Remark 3.12. The algebra of Dirac fields with MIT boundary condition cannot be considered as an algebra of observables, since observables are required to commute at spacelike separations and \mathfrak{A} does not fulfil such requirement. A good candidate as algebra of observables is the subalgebra $\mathfrak{A}_{\text{obs}} \subset \mathfrak{A}$ consisting of elements which are even, *i.e.* invariant by replacement $\Xi(\psi) \mapsto -\Xi(\psi)$, and invariant under the action of $\text{Spin}_0(1, n)$ (extended to \mathfrak{A}). For further details we refer to [26].

3.5 Hadamard states

In this section we study (algebraic) states and their interplay with the $*$ -isomorphism $\mathfrak{R}_{1,0}$.

Definition 3.13. Given a complex $*$ -algebra \mathfrak{A} we call (*algebraic*) *state* any linear functional from \mathfrak{A} into \mathbb{C} that is positive, *i.e.* $\omega(\mathfrak{a}^* \mathfrak{a}) \geq 0$ for any $\mathfrak{a} \in \mathfrak{A}$, and normalized, *i.e.* $\omega(1_{\mathfrak{A}}) = 1$.

Due to the natural grading on the algebra of Dirac fields with MIT boundary conditions \mathfrak{A} , it suffices to define ω on the monomials. Among all states, the so-called quasi-free states play a distinguished role.

Definition 3.14. A state ω on \mathfrak{A} is *quasifree* if it satisfies

$$\omega(\Xi(\psi_1) \cdots \Xi(\psi_n)) = \begin{cases} 0 & n \text{ odd} \\ \sum_{\sigma \in S'_n} (-1)^{\text{sign}(\sigma)} \prod_{i=1}^{n/2} \omega(\Xi(\psi_{\sigma(2i-1)}) \Xi(\psi_{\sigma(2i)})) & n \text{ even} \end{cases},$$

where S'_n denotes the set of ordered permutations of n elements.

As shown in [2, Lemma 3.2], for any quasi-free state ω on the C^* -algebra \mathfrak{A} there exists a bounded operator $Q_\omega \in B(\text{Sol}_{sc, \text{MIT}}^\oplus)$ on $\text{Sol}_{sc, \text{MIT}}^\oplus$ such that $0 \leq Q_\omega = Q_\omega^* \leq 1$, $Q_\omega + \Gamma Q_\omega \Gamma = \text{Id}_{\text{Sol}_{sc, \text{MIT}}^\oplus}$ and

$$\omega(\Xi(\psi_1)^* \Xi(\psi_2)) = (\psi_1, Q_\omega \psi_2)_{\text{Sol}_{sc, \text{MIT}}^\oplus}. \quad (3.9)$$

From a different perspective, we can realize $\omega(\Xi(\psi_1)^* \Xi(\psi_2))$ in terms of distributions. This turns out to be quite useful when looking for physically relevant states. To this avail we observe that, an application of Proposition 2.20 leads to $\text{Sol}_{sc, \text{MIT}}^\oplus \simeq \left(\Gamma_c(\mathbb{S}M) / D\Gamma_{c, \text{MIT}}(\mathbb{S}M) \right)^{\oplus 2}$ —cf. Equation (2.5)— the isomorphism being given by $\left(\Gamma_c(\mathbb{S}M) / D\Gamma_{c, \text{MIT}}(\mathbb{S}M) \right)^{\oplus 2} \ni ([f_1], [f_2]) \rightarrow \mathbf{G}f_1 \oplus \Gamma \mathbf{G}f_2 \in \text{Sol}_{sc, \text{MIT}}^\oplus$. In particular we can endow $\Gamma_c(\mathbb{S}M)$ with the standard locally convex topology which induces a locally convex topology on the quotient $\Gamma_c(\mathbb{S}M) / D\Gamma_{c, \text{MIT}}(\mathbb{S}M)$. With this choices the map $\left(\Gamma_c(\mathbb{S}M) / D\Gamma_{c, \text{MIT}}(\mathbb{S}M) \right)^{\oplus 2} \rightarrow \text{Sol}_{sc, \text{MIT}}^\oplus$ turns out to be continuous, so that to any quasi-free state we may associate its 2-point distributions $\omega^{(2)} \in \Gamma_c((\mathbb{S}M \oplus \mathbb{S}M)^2)'$ defined by

$$\omega^{(2)}(f_1, f_2) := \omega(\Xi(\psi_{f_1})^* \Xi(\psi_{f_2})).$$

where $\psi_f \in \text{Sol}_{sc, \text{MIT}}^\oplus$ is the element associated with $[f] \in \left[\Gamma_c(\mathbb{S}M) / D\Gamma_{c, \text{MIT}}(\mathbb{S}M) \right]^{\oplus 2}$. In particular, we have that the 2-point distribution is a solution of the Dirac equation with MIT boundary conditions, meaning that

$$\omega^{(2)}(f_1, (D \oplus D)f_1) = 0 \quad \forall f_1, f_2 \in \Gamma_{c, \text{MIT}}(\mathbb{S}M \oplus \mathbb{S}M). \quad (3.10)$$

Notice that, due to the CAR relations, Equation (3.10) cannot be strengthened to hold true for all $f_1, \dots, f_n \in \Gamma_c(\mathbb{SM} \oplus \mathbb{SM})$.

A widely accepted criterion to select physically relevant states is the renowned *Hadamard condition* [57, 69–71]. On a globally hyperbolic spacetime with empty boundary, the latter allows to construct Wick polynomials in a local and covariant fashion, moreover, it guarantees the finiteness of the fluctuations of such Wick polynomials [41].

At a technical level, the Hadamard condition characterizes the wave front set $\text{WF}(\omega^{(2)}) \subseteq \mathbb{T}^*\mathbb{M}^2$ of the 2-point function of a quasi-free state —generalization to non-quasi free states are possible [73]. Such a microlocal characterization is also possible for the case of a globally hyperbolic manifold with timelike boundary: therein the Hadamard condition has been formulated in [77] for the case of asymptotically Anti-de Sitter spacetimes and then exploited in [29] for a wider class of boundary conditions. In these situations the proper replacement for $\text{WF}(\omega^{(2)})$ is given by $\text{WF}_b(\omega^{(2)}) \subset {}^b\mathbb{T}^*\mathbb{M}^2 \setminus \{0\}$, where WF_b stands for the b -wave front set [62].

Definition 3.15. Let (\mathbb{M}, g) be a globally hyperbolic spin manifold with timelike boundary. A bidistribution $\omega^{(2)} \in \Gamma_c(\mathbb{SM} \oplus \mathbb{SM})'$ is called of *Hadamard form* if it has the following b -wave front set

$$\text{WF}_b(\omega^{(2)}) = \{(x, y, k_x, -k_y) \in \mathbb{T}^*(\mathbb{M} \times \mathbb{M}) \setminus \{0\} \mid (x, k_x) \sim (y, k_y), k_x \triangleright 0\},$$

where \sim entails that (x, k_x) and (y, k_y) are connected by a generalized broken bicharacteristic while that $k_x \triangleright 0$ means that the covector k_x at $x \in \mathbb{M}$ is future pointing. Since we deal with vector-valued distributions, the standard convention for the wave front set is to take the union of the wave front set of its components in an arbitrary but fixed local frame.

For further details on Hadamard states on globally hyperbolic manifolds with empty boundary we refer to [50, 51, 58], while on globally hyperbolic manifolds with timelike boundary, we refer to [29, 49, 77].

With the next theorem, we show that the pull-back of a quasifree state along the isomorphism $\mathfrak{R}_{1,0} : \mathfrak{A}_0 \rightarrow \mathfrak{A}_1$ induced by the unitary Møller operator R for D preserves the singularity structure of the two-point distribution $\omega^{(2)}$.

Theorem 3.16. Assume that $g_0, g_1 \in \mathcal{GH}_{\mathbb{M}}$ fulfil (i) in the Setup 2.21. Assume furthermore that a propagation of singularities theorem holds true for D with MIT boundary condition, namely for any $u \in \text{Sol}_{\text{MIT}}(D)$, $\text{WF}_b(u)$ is the union of maximally extended generalized broken bicharacteristics. Denote with \mathfrak{A}_α , $\alpha = 0, 1$, the algebras of Dirac fields with MIT boundary condition on \mathbb{M}_α and let $\omega_\alpha : \mathfrak{A}_\alpha \rightarrow \mathbb{C}$ be quasifree states satisfying

$$\omega_0 = \omega_1 \circ \mathfrak{R}_{1,0} : \mathfrak{A}_0 \rightarrow \mathbb{C}$$

with $\mathfrak{R}_{1,0}$ is the isomorphism induced by R as per Theorem 3.11. Then, if ω_1 is a Hadamard state as per Definition 3.15, then so is ω_0 .

Proof. Since $\mathfrak{R}_{1,0}$ preserves the grading of \mathfrak{A}_0 , \mathfrak{A}_1 , ω_0 inherits the property of being a quasifree state from ω_1 . In particular the two-point function $\omega_0^{(2)}$ satisfies

$$\omega_0^{(2)}(f_0, g_0) = \omega_0(\Xi(\psi_{f_0})^* \Xi(\psi_{g_0})) = \omega_1^{(2)}\left(\Xi(R_{1,0}^\oplus \psi_{f_0})^* \Xi(R_{1,0}^\oplus \psi_{g_0})\right).$$

We shall now prove that ω_1 fulfils the Hadamard condition. To this avail we first observe that $R_{1,0}$ can in fact be decomposed as $R_{1,0} = R_{1,\chi} \circ R_{\chi,0}$ (cf. Remark 2.35). With reference to Theorem 2.27, we have $R_{\chi,0} := U_{D_{\chi,1,-}^f} \circ \rho_- \circ \kappa_{1,0}^f$ whereas $R_{1,\chi} := U_{D_{1,+}} \circ \rho_+$. Let us consider $\mathfrak{R}_{1,\chi} : \mathfrak{A}_\chi \rightarrow \mathfrak{A}_1$, where $\mathfrak{R}_{1,\chi}$ is the $*$ -isomorphism defined as per Theorem 3.11 with \mathfrak{A}_0 replaced with \mathfrak{A}_χ . Moreover let $\omega_\chi := \omega_1 \circ \mathfrak{R}_{1,\chi}$. With reference to Theorem 2.27, let $f_1, f_2 \in \Gamma_c(\mathbb{SM} \oplus \mathbb{SM})$

be with support contained in a neighbourhood of Σ_+ . Then

$$\begin{aligned}
\omega_\chi^{(2)}(f_1, f_2) &= \omega_\chi(\Xi(G_\chi f_1)^* \Xi(G_\chi f_2)) \quad \text{def. } \omega_\chi^{(2)} \\
&= \omega_1(\Xi(R_{1,\chi}^\oplus G_\chi f_1)^* \Xi(R_{1,\chi}^\oplus G_\chi f_2)) \quad \text{def. } \mathfrak{R}_{1,\chi} \\
&= (R_{1,\chi} G_\chi f_1, Q_{\omega_1} R_{1,\chi} G_\chi f_2)_{\text{Sol}_{sc, \text{MIT}}(D_1)} \quad \text{Eq. (3.9)} \\
&= (\rho_- R_{1,\chi}^\oplus G_\chi f_1, Q_{\omega_1} \rho_- R_{1,\chi}^\oplus G_\chi f_2)_{\Sigma_+} \quad \text{choice of } \Sigma_+ \\
&= (\rho_- G_\chi f_1, Q_{\omega_1} \rho_- G_\chi f_2)_{\Sigma_+} \quad \rho_- R_{1,\chi} = \rho_- \\
&= (\rho_- G_1 f_1, Q_{\omega_1} \rho_- G_1 f_2)_{\Sigma_+} \\
&= \omega_1^{(2)}(f_1, f_2),
\end{aligned}$$

where we exploited the fact that, when computing $(\cdot, \cdot)_{\text{Sol}_{sc, \text{MIT}}^\oplus}$, we may choose Σ arbitrarily. In the second to last equation we used that $G_\chi f|_{\Sigma_+} = G_1 f|_{\Sigma_+}$ for f supported in a small enough neighbourhood of Σ_+ . This shows that $\omega_\chi^{(2)}$ coincides with $\omega_1^{(2)}$ in a neighbourhood of Σ_+ and therefore fulfils the Hadamard condition therein. By the assumed propagation of singularities, it follows that $\omega_\chi^{(2)}$ fulfils the Hadamard condition on M .

By observing that $\omega_1 = \omega_\chi \circ \mathfrak{R}_{\chi,0}$ and proceeding with a similar argument we have that ω_1 fulfils the Hadamard condition. \square

Remark 3.17. We expect the propagation of singularities to hold true, as there are already positive results in this direction, see *e.g.* [29, 49, 63, 64, 76] for the scalar wave equation, [60, 75] for first order systems, and [9] for the Dirac-Coulomb system. We postpone its investigation to a forthcoming paper.

We have finally all the tools to prove the existence of Hadamard states.

Proof of Theorem 1.1. On account of Proposition 3.8, there exists a unitary Møller operator $R_{1,0}$ which is a unitary map between the $\text{Sol}_{sc, \text{MIT}}(D_0)$ and $\text{Sol}_{sc, \text{MIT}}(D_1)$. As proved in Theorem 3.11, the action of $R_{1,0}$ can be lifted to a $*$ -isomorphism $\mathfrak{R}_{1,0} : \mathfrak{A} \rightarrow \mathfrak{A}_u$ between the algebra of Dirac field on (M, g) and (M, g_u) respectively. Hence for any Hadamard state ω_H on \mathfrak{A}_u , the state defined by

$$\omega = \omega_H \circ \mathfrak{R}_{1,0} : \mathfrak{A} \rightarrow \mathbb{C},$$

is also a Hadamard state on account of Theorem 3.16.

It remains to show that there exists a Hadamard state ω_H for \mathfrak{A}_u . For that, we shall define a quasi-free state by identifying a suitable operator Q_ω and then exploiting Equation (3.9). In order to construct the desired Q_ω , let us write the Dirac equation as $D = \sigma(dt)\partial_t + L$, where L differentiates only in the tangential part of Σ . Since we coupled D with self-adjoint boundary condition, it follows that L is skew-adjoint. As a consequence we may define the self-adjoint operator $H = iL$. To obtain a pure, quasi-free state it is enough to define the operator

$$Q_\omega := P_+(H) \oplus \text{Id}_{\mathcal{H}} - \Upsilon P_+(H) \Upsilon^{-1}$$

where $P_+(H)$ is the spectral projection in the positive spectrum of H . It is not difficult to see, that on globally hyperbolic ultrastatic manifolds with empty boundary, the associated quasi-free state is of the Hadamard form, since it provides the canonical frequencies splitting. This concludes our proof. \square

Remark 3.18. The main drawback of the definition of the Møller $*$ -isomorphism \mathfrak{R} , used in Theorem 3.16, is the lack of any control on the action of the group of $*$ -automorphism induced by the isometry group of M on ω_2 . Let us remark, that the study of invariant states is a well-established research topic (*cf.* [7, 8]). Indeed, the type of factor can be inferred by analyzing which and how many states are invariant. From a more physical perspective instead, invariant states can represent equilibrium states in statistical mechanics *e.g.* KMS-states or ground states.

The previous remark leads us to the following open question: Under which conditions it is possible to perform an adiabatic limit, namely when is $\lim_{\chi \rightarrow 1} \omega_1$ well-defined?

A priori we expect that there is no positive answer in all possible scenarios, since it is known that certain free-field theories, e.g., the massless and minimally coupled (scalar or Dirac) field on four-dimensional de Sitter spacetime, do not possess a ground state, even though their massive counterpart does. (Notice that this is not a no-go Theorem, but at least an indication that, in these situation, the map $\omega \rightarrow \omega \circ \mathfrak{A}$ cannot be expected to preserve the ground state property.) A partial investigation in this direction has been carried on in [22, 34] for the case of a scalar field theory on globally hyperbolic spacetimes with empty boundary. In this situation it has been shown that, under suitable hypotheses the adiabatic limit can be performed preserving the invariance property under time translation but spoiling in general the ground state or KMS property.

Since our results depends only on the principal symbol of the Dirac operator, we conclude our paper with the following corollary.

Corollary 3.19. *Let (M, g) be a globally hyperbolic spin spacetime with timelike boundary and such that there exists an ultrastatic manifold $(M, g_u := -dt^2 + h)$ such that $\beta^{-2}h(t) \leq C^2(t)h$. Let $D_V = D + V$ be the Dirac operator coupled with a external skew-symmetric potential $V \in \text{End}(\mathcal{SM})$ and with the MIT boundary condition. Then there exists a state for the algebra of Dirac fields with MIT boundary conditions which satisfies the Hadamard condition.*

References

- [1] L. Aké, J. L. Flores and M. Sánchez, *Structure of globally hyperbolic spacetimes with timelike boundary*. Rev. Mat. Iberoam. **37**, 45–94 (2021).
- [2] H. Araki, *On quasifree states of CAR and Bogoliubov automorphisms*. Publ. Res. Inst. Math. Sci. Kyoto **6**, 385 (1971).
- [3] C. Bär, *Green-hyperbolic operators on globally hyperbolic spacetimes*. Comm. Math. Phys. **333**, 1585–1615 (2015).
- [4] C. Bär, P. Gauduchon and A. Moroianu, *Generalized cylinders in semi-Riemannian and spin geometry*. Math. Z. **249**, 545–580 (2005).
- [5] C. Bär and N. Ginoux, *Classical and quantum fields on Lorentzian manifolds*. in: C. Bär, J. Lohkamp and M. Schwarz (eds.), Global Differential Geometry, 359–400, Springer-Verlag Berlin Heidelberg (2012).
- [6] C. Bär, N. Ginoux and F. Pfäffle, *Wave equations on Lorentzian Manifolds and Quantization*. ESI Lectures in Mathematics and Physics (2007).
- [7] F. Bambozzi and S. Murro, *On the uniqueness of invariant states*. Adv. Math. **376** 107445 (2021).
- [8] F. Bambozzi, S. Murro and N. Pinamonti, *Invariant states on noncommutative tori*. Int. Math. Res. Not. **2021**, 3299–3313 (2021).
- [9] D. Baskin and J. Wunsch, *Diffraction for the Dirac-Coulomb propagator*. arXiv:2011.08890 [math.AP] (2020).
- [10] J.K. Beem, P.E. Ehrlich and K.L. Easley, *Global Lorentzian geometry*. Marcel Dekker, New York (1996).

- [11] M. Benini, C. Dappiaggi and S. Murro, *Radiative observables for linearized gravity on asymptotically flat spacetimes and their boundary induced states*. J. Math. Phys. **55**, 082301 (2014).
- [12] M. Benini, C. Dappiaggi and A. Schenkel, *Algebraic quantum field theory on spacetimes with timelike boundary*. Ann. Henri Poincaré **19**, 2401–2433 (2018).
- [13] R. Brunetti, C. Dappiaggi, K. Fredenhagen and J. Yngvason, *Advances in algebraic quantum field theory*. Springer (2015)
- [14] R. Brunetti, K. Fredenhagen and N. Pinamonti, *Algebraic approach to Bose Einstein Condensation in relativistic Quantum Field Theory. Spontaneous symmetry breaking and the Goldstone Theorem*, Ann. Henri Poincaré volume **22**, 951–1000 (2021).
- [15] F. Bussola, C. Dappiaggi, H.R.C. Ferreira and I. Khavkine, *Ground state for a massive scalar field in the BTZ spacetime with Robin boundary conditions*. Phys. Rev. D **96**, 105016 (2017).
- [16] M. Capoferri, C. Dappiaggi and N. Drago, *Global wave parametrices on globally hyperbolic spacetimes*. J. Math. Anal. App. **490**, 124316 (2020).
- [17] M. Capoferri, M. Levitin and D. Vassiliev, *Geometric wave propagator on Riemannian manifolds*. to appear in Comm. Anal. Geom, arXiv:1902.06982 [math.AP] (2019).
- [18] M. Capoferri and D. Vassiliev, *Global propagator for the massless Dirac operator and spectral asymptotics*. arXiv:2004.06351 [math.AP], (2020).
- [19] M. Capoferri and D. Vassiliev, *Invariant subspaces of elliptic systems II: spectral theory*. to appear in J. Spectr. Theory, arXiv:2103.14334 [math.AP] (2020).
- [20] A. Chodos, R.L. Jaffe, K. Johnson and C.B. Thorn, *Baryon Structure in the Bag Theory*. Phys. Rev. D **10**, 2599 (1974).
- [21] A. Chodos, R.L. Jaffe, K. Johnson, C.B. Thorn and V.F. Weisskopf, *New extended model of hadrons*. Phys. Rev. D **9**, 3471 (1974).
- [22] C. Dappiaggi and N. Drago, *Constructing Hadamard States via an Extended Møller Operator*. Lett. Math. Phys. **106**, 1587–1615 (2016).
- [23] C. Dappiaggi, N. Drago, H. Ferreira, *Fundamental solutions for the wave operator on static Lorentzian manifolds with timelike boundary*, Lett Math Phys **109**: 2157 (2019).
- [24] C. Dappiaggi, N. Drago, R. Longhi, *On Maxwell’s Equations on Globally Hyperbolic Spacetimes with Timelike Boundary*. Ann. Henri Poincaré **21**, 2367–2409 (2020).
- [25] C. Dappiaggi, F. Finster, S. Murro and E. Radici, *The Fermionic Signature Operator in De Sitter Spacetime*. J. Math. Anal. Appl. **485**, 123808 (2020).
- [26] C. Dappiaggi, T.-P. Hack and N. Pinamonti *The extended algebra of observables for Dirac fields and the trace anomaly of their stress-energy tensor*. Rev. Math. Phys. **21**, 1241–1312 (2009).
- [27] C. Dappiaggi, T.-P. Hack and K. Sanders, *Electromagnetism, local covariance, the Aharonov-Bohm effect and Gauss’ law*. Commun. Math. Phys. **328**, 625 (2014).
- [28] C. Dappiaggi and A. Marta, *A generalization of the propagation of singularities theorem on asymptotically anti-de Sitter spacetimes*. arXiv:2006.00560 (2020).
- [29] C. Dappiaggi and A. Marta, *Fundamental solutions and Hadamard states for a scalar field with arbitrary boundary conditions on an asymptotically AdS spacetimes*. Math. Phys., Anal. Geom. **24** 28 (2021).

- [30] C. Dappiaggi, V. Moretti and N. Pinamonti, *Rigorous construction and Hadamard property of the Unruh state in Schwarzschild spacetime*. Adv. Theor. Math. Phys. **15** 355 (2011)
- [31] C. Dappiaggi, S. Murro and A. Schenkel, *Non-existence of natural states for Abelian Chern-Simons theory*. J. Geom. Phys. **116**, 119–123 (2017).
- [32] C. Dappiaggi, G. Nosari and N. Pinamonti, *The Casimir effect from the point of view of algebraic quantum field theory*. Math. Phys. Anal. Geom. **19**, 12 (2016).
- [33] M. Di Francesco, S. Fagioli and E. Radici, *Deterministic particle approximation for nonlocal transport equations with nonlinear mobility*. J. Differential Equations **266**, 2830–2868 (2019).
- [34] N. Drago and C. Gérard, *On the adiabatic limit of Hadamard states*. Lett. Math. Phys. **107**, 1409–1438 (2017).
- [35] N. Drago, T-P. Hack and N. Pinamonti, *The generalized principle of Perturbative Agreement and the thermal mass*. Ann. Henri Poincaré **18**, 807 (2017).
- [36] N. Drago, N. Ginoux and S. Murro, *On the Cauchy problem for hyperbolic Friedrichs systems with additional constraints*. in preparation.
- [37] N. Drago, N. Große and S. Murro, *The Cauchy problem of the Lorentzian Dirac operator with APS boundary conditions*. arXiv:2104.00585 [math.AP] (2021).
- [38] N. Drago and S. Murro, *A new class of Fermionic Projectors: Møller operators and mass oscillation properties*. Lett. Math. Phys. **107**, 2433–2451 (2017).
- [39] S. Fagioli and E. Radici, *Solutions to aggregation-diffusion equations with nonlinear mobility constructed via a deterministic particle approximation*. Math. Models Methods Appl. Sci. **28** (9), 1801–1829 (2018).
- [40] S. Fagioli and E. Radici, *Opinion formation systems via deterministic particles approximation*. Kinetic & Related Models, **14** (1), 45–76 (2021).
- [41] C. J. Fewster, R. Verch, *The necessity of the Hadamard condition*, Class. Quantum Grav. **30**, 235027 (2013).
- [42] F. Finster, S. Murro and C. Röken, *The Fermionic Projector in a Time-Dependent External Potential: Mass Oscillation Property and Hadamard States*. J. Math. Phys. **57**, 072303 (2016).
- [43] F. Finster, S. Murro and C. Röken, *The Fermionic Signature Operator and Quantum States in Rindler Space-Time*. J. Math. Anal. Appl. **454**, 385 (2017).
- [44] F. Finster and C. Röken, *Self-adjointness of the Dirac Hamiltonian for a class of non-uniformly elliptic boundary value problems*. Annals of Mathematical Sciences and Applications **1** , 301 (2016).
- [45] F. Finster and C. Röken, *An integral spectral representation for the massive Dirac propagator in Kerr geometry in Eddington-Finkelstein-type coordinates*. Adv. Theor. Math. Phys. **22**, no. 1, 47–92 (2018).
- [46] K. Fredenhagen and K. Rejzner, *Quantum field theory on curved spacetimes: Axiomatic framework and examples*. J. Math. Phys. **57**, 031101 (2016).
- [47] K.O. Friedrichs, *Symmetric positive linear differential equations*. Comm. Pure Appl. Math. **11**, 333–418 (1958).

- [48] S.A. Fulling, F.J. Narcowich and R.M. Wald, *Singularity structure of the two-point function in quantum field theory in curved spacetime, II*. Annals of Physics **136** (1981), 243.
- [49] O. Gannot and M. Wrochna, *Propagation of singularities on AdS spacetimes for general boundary conditions and the holographic Hadamard condition*. to appear in J. Inst. Math. Jussieu.
- [50] C. Gérard, *Microlocal Analysis of Quantum Fields on Curved Spacetimes*. ESI Lectures in Mathematics and Physics (2019).
- [51] C. Gérard, T. Stoskopf, *Hadamard states for quantized Dirac fields on Lorentzian manifolds of bounded geometry*, arXiv:2108.11630 [math.AP] (2021)
- [52] N. Ginoux and S. Murro, *On the Cauchy problem for Friedrichs systems on globally hyperbolic manifolds with timelike boundary*. arXiv:2007.02544 [math.AP] (2020).
- [53] N. Große and S. Murro, *The well-posedness of the Cauchy problem for the Dirac operator on globally hyperbolic manifolds with timelike boundary*. Documenta Math. **25**, 737–765 (2020).
- [54] R.W. Haymaker and T. Goldman, *Bag boundary conditions for confinement in the $q\bar{q}$ relative coordinate*. Phys. Rev. D **24**, 743 (1981).
- [55] S. Hollands and R. M. Wald, *Existence of local covariant time ordered products of quantum fields in curved spacetime*. Commun. Math. Phys. **231**, 309 (2002).
- [56] S. Hollands and R. M. Wald, *Conservation of the stress tensor in perturbative interacting quantum field theory in curved spacetimes* Rev. Math. Phys. **17**, 227 (2005).
- [57] B.S. Kay, R.M. Wald, *Theorems on the uniqueness and thermal properties of stationary, non-singular, quasifree states on spacetimes with a bifurcate Killing horizon*, Phys. Rep. 207(2), 49-136 (1991).
- [58] I. Khavkine and V. Moretti, *Algebraic QFT in Curved Spacetime and quasifree Hadamard states: an introduction*. Advances in Algebraic Quantum Field Theory. Springer International Publishing, 2015.
- [59] G. Idelon-Riton, *Scattering theory for the Dirac equation on the Schwarzschild-anti-de Sitter space-time*. Adv. Theor. Math. Phys. 22, no.4, 1007–1069 (2018).
- [60] V.Ya. Ivrii, *Microlocal Analysis, Sharp Spectral Asymptotics and Applications I*. Springer (2019).
- [61] P.D. Lax and R.S. Phillips, *Local Boundary Conditions for Dissipative Symmetric Linear Differential Operators*, Comm. Pure Appl. Math. **13**, 427–455, (1960).
- [62] R.B. Melrose, *The Atiyah-Patodi-Singer Index Theorem*, volume 4 of Research Notes in Mathematics. A K Peters Ltd., Wellesley, MA, (1993).
- [63] R.B. Melrose and J. Sjöstrand, *Singularities of boundary value problems. i*. Comm. Pure Appl. Math. **31** (5), 593–617 (1978).
- [64] R.B. Melrose and J. Sjöstrand, *Singularities of boundary value problems. ii*. Comm. Pure Appl. Math. **35** (2), 129–168 (1982).
- [65] V. Moretti, S. Murro and D. Volpe, *Paracausal deformations of Lorentzian metrics and Møller isomorphisms in algebraic quantum field theories*. in preparation.
- [66] S. Murro and D. Volpe, *Intertwining operators for symmetric hyperbolic systems on globally hyperbolic manifolds*. Ann. Glob. Anal. Geom. **59**, 1–25 (2021).

- [67] M. Nardmann, *Pseudo-Riemannian metrics with prescribed scalar curvature*, PhD thesis, Universität Leipzig, 2004.
- [68] W. Pusz and S.L. Woronowicz, *Passive States and KMS States for General Quantum Systems*. Comm. Math. Phys. **58**, 273–290 (1978).
- [69] M. J. Radzikowski, *Microlocal approach to the Hadamard condition in quantum field theory on curved space-time*, Commun. Math. Phys. **179**, 529–553 (1996).
- [70] M. J. Radzikowski (with an Appendix by Rainer Verch), *A Local-to-Global Singularity Theorem for Quantum Field Theory on Curved Space-Time*, Commun. Math. Phys. **180**, 1–22 (1996).
- [71] H. Sahlmann and R. Verch, *Microlocal spectrum condition and Hadamard form for vectorvalued quantum fields in curved space-time*, Rev. Math. Phys. **13**, 1203–1246 (2001).
- [72] H. Sahlmann and R. Verch, *Passivity and Microlocal Spectrum Condition*. Comm. Math. Phys. **241**, 705–731 (2000).
- [73] K. Sanders, *Equivalence of the (Generalised) Hadamard and Microlocal Spectrum Condition for (Generalised) Free Fields in Curved Spacetime*, Commun. Math. Phys. **295**, 485–501 (2010).
- [74] A. Seyedzahedi, R. Saghian and S.S. Gousheh, *Fermionic Casimir energy in a three-dimensional box*. Phys. Rev. A **82**, 032517 (2010).
- [75] M.E. Taylor, *Reflection of Singularities of Solutions to Systems of Differential Equations*, Comm. Pure Appl. Math. **23** 457–478 (1975).
- [76] A. Vasy, *Propagation of singularities for the wave equation on manifolds with corners.*, Ann. Math. **168** 749–812, (2008).
- [77] M. Wrochna, *The holographic Hadamard condition on asymptotically Anti-de Sitter spacetimes*. Lett. Math. Phys. **107**, 2291 (2017).
- [78] J. Zahn, *Generalized Wentzell Boundary Conditions and Quantum Field Theory*. Ann. Henri Poincaré, **19**, 163187 (2018).