



**HAL**  
open science

# Contrasting Winter Versus Summer Microbial Communities and Metabolic Functions in a Permafrost Thaw Lake

Adrien Vigneron, Connie Lovejoy, Perrine Cruaud, Dimitri Kalenitchenko, Alexander Culley, Warwick F Vincent

► **To cite this version:**

Adrien Vigneron, Connie Lovejoy, Perrine Cruaud, Dimitri Kalenitchenko, Alexander Culley, et al.. Contrasting Winter Versus Summer Microbial Communities and Metabolic Functions in a Permafrost Thaw Lake. *Frontiers in Microbiology*, 2019, 10, 10.3389/fmicb.2019.01656 . hal-03333742

**HAL Id: hal-03333742**

**<https://hal.science/hal-03333742>**

Submitted on 3 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Contrasting Winter Versus Summer Microbial Communities and Metabolic Functions in a Permafrost Thaw Lake

Adrien Vigneron<sup>1,2,3\*</sup>, Connie Lovejoy<sup>1,2,3,4</sup>, Perrine Cruaud<sup>3,5</sup>, Dimitri Kalenitchenko<sup>1,3,4</sup>, Alexander Culley<sup>2,3,5</sup> and Warwick F. Vincent<sup>1,2,3</sup>

<sup>1</sup> Département de Biologie, Université Laval, Quebec, QC, Canada, <sup>2</sup> Centre d'Études Nordiques, Takuvik Joint International Laboratory, Université Laval, Quebec, QC, Canada, <sup>3</sup> Institut de Biologie Intégrative et des Systèmes, Université Laval, Quebec, QC, Canada, <sup>4</sup> Québec Océan, Université Laval, Quebec, QC, Canada, <sup>5</sup> Département de Biochimie, de Microbiologie et de Bio-Informatique, Université Laval, Quebec, QC, Canada

## OPEN ACCESS

### Edited by:

Peter Dunfield,  
University of Calgary, Canada

### Reviewed by:

Maggie C. Y. Lau,  
Institute of Deep-Sea Science  
and Engineering (CAS), China  
Lyle Whyte,  
McGill University, Canada

### \*Correspondence:

Adrien Vigneron  
avignero@gmail.com

### Specialty section:

This article was submitted to  
Extreme Microbiology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 21 February 2019

**Accepted:** 04 July 2019

**Published:** 16 July 2019

### Citation:

Vigneron A, Lovejoy C, Cruaud P,  
Kalenitchenko D, Culley A and  
Vincent WF (2019) Contrasting Winter  
Versus Summer Microbial  
Communities and Metabolic  
Functions in a Permafrost Thaw Lake.  
*Front. Microbiol.* 10:1656.  
doi: 10.3389/fmicb.2019.01656

Permafrost thawing results in the formation of thermokarst lakes, which are biogeochemical hotspots in northern landscapes and strong emitters of greenhouse gasses to the atmosphere. Most studies of thermokarst lakes have been in summer, despite the predominance of winter and ice-cover over much of the year, and the microbial ecology of these waters under ice remains poorly understood. Here we first compared the summer versus winter microbiomes of a subarctic thermokarst lake using DNA- and RNA-based 16S rRNA amplicon sequencing and qPCR. We then applied comparative metagenomics and used genomic bin reconstruction to compare the two seasons for changes in potential metabolic functions in the thermokarst lake microbiome. In summer, the microbial community was dominated by Actinobacteria and Betaproteobacteria, with phototrophic and aerobic pathways consistent with the utilization of labile and photodegraded substrates. The microbial community was strikingly different in winter, with dominance of methanogens, Planctomycetes, Chloroflexi and Deltaproteobacteria, along with various taxa of the Patescibacteria/Candidate Phyla Radiation (Parcubacteria, Microgenomates, Omnitrophica, Aminicenantes). The latter group was underestimated or absent in the amplicon survey, but accounted for about a third of the metagenomic reads. The winter lineages were associated with multiple reductive metabolic processes, fermentations and pathways for the mobilization and degradation of complex organic matter, along with a strong potential for syntrophy or cross-feeding. The results imply that the summer community represents a transient stage of the annual cycle, and that carbon dioxide and methane production continue through the prolonged season of ice cover via a taxonomically distinct winter community and diverse mechanisms of permafrost carbon transformation.

**Keywords:** MAGs, microbial diversity, metagenomes, methane, permafrost, thermokarst, winter limnology

## INTRODUCTION

Northern landscapes are experiencing rapid change due to climate warming, with large scale thawing and erosion of permafrost soils in many regions (Oliva and Fritz, 2018). Around half of global soil carbon is estimated to be stored in permafrost (Schoor et al., 2015). This organic carbon is derived from the remnants of plants, animals and microbes that accumulated during warmer periods over the history of the Earth (Jansson and Taş, 2014; Schoor et al., 2015). Accelerated thawing is beginning to unlock these reserves along with indigenous permafrost microorganisms that were preserved in a dormant state (Graham et al., 2011; Mackelprang et al., 2017), and this old organic matter is now progressively decomposed by microbial activities. Shotgun metagenomic analysis of the microbiome of thawed subarctic soils has indicated a broad enzymatic capability to break down complex polymers that are found in permafrost (Woodcroft et al., 2018). The net result is the release of greenhouse gases to the atmosphere (N<sub>2</sub>O, CO<sub>2</sub> and CH<sub>4</sub>), and a positive feedback to climate warming (Schoor et al., 2015; Matveev et al., 2016). Decomposition and greenhouse gas emission rates from northern soils are variable (Parmentier et al., 2017), and some of the highest CH<sub>4</sub> emissions of the permafrost region have been measured in thermokarst lakes and ponds (Matveev et al., 2016). These open waters form by thawing of ice-rich permafrost, which causes surface collapse and the creation of basins that fill with meltwater and precipitation. These lakes represent a quantitatively important type of aquatic ecosystem in Arctic and subarctic regions (Bouchard et al., 2014; Farquharson et al., 2016) and are viewed as biogeochemical hotspots in the permafrost landscape (Deshpande et al., 2016). Calculations suggest that the rapid thaw of sediments beneath thermokarst lakes by the latent heat of the water body could more than double the rate of carbon release from these abundant northern lakes by the end of the century (Walter Anthony et al., 2018).

Microbial communities in thermokarst lakes are reported to be similar to communities in other freshwaters, but enriched in methane cycling microorganisms (Crevecoeur et al., 2015, 2016; Comte et al., 2016) and uncultured bacterial taxa that are poorly resolved with standard primer sets (Wurzbacher et al., 2017). Thermokarst lake studies have been mostly carried out in summer, and microbial community composition during this open water period is strongly influenced by atmospheric exchange, meteorological conditions and exposure to sunlight, as in other freshwater ecosystems (Bertilsson et al., 2013). In contrast, all of these factors are absent or strongly reduced as soon as freeze-up occurs each winter. During this period that covers 8 or more months of the year in the North, ice and snow cover acts as an insulating barrier to atmospheric exchange and sunlight, compounded by short day lengths in winter. The ice cover leads to a rapid depletion of oxygen under the ice (Deshpande et al., 2017), while maintaining temperatures above freezing in the water column below the ice.

Methane production in winter is common in ice-covered lakes in temperate regions (Ricão Canelhas et al., 2016; Denfeld et al., 2018), and methane bubbles can become trapped in the ice of northern lakes (Langer et al., 2015). Even in temperate systems,

the microbial community composition and metabolism under ice have been largely neglected compared to ice-free periods, despite evidence that the microbial community structure can differ substantially from summer to winter (Bertilsson et al., 2013). However, temperate lakes may not be an accurate guide to the functioning of thermokarst ecosystems. The summer primary production that occurs during a much shorter season of ice-free conditions in Arctic regions may not be sufficient to support the overwintering microbial communities, potentially leading to minimal biological activities and a rundown of microbial biomass by the end of the winter. Conversely, the abundance of terrestrially derived permafrost organic carbon (Wauthy et al., 2018), coupled with the prolonged anoxic conditions during winter (Deshpande et al., 2017), could be sufficient for an alternate community to develop during this time of year.

The aim of the present study was to compare the winter versus summer microbial communities of a thermokarst lake, and to evaluate two competing hypotheses: dormancy or downshift of the microbial community in winter toward lower population size and activity, versus a functional reassembly of the community without net loss of abundance or activity. We sampled a subarctic thermokarst lake, firstly during open water conditions in late summer, and then in late winter after several months of ice cover. To address the question of how microbiome structure and function differ between the two seasons, we characterized the microbial communities in summer and winter by DNA- and RNA-based 16S rRNA amplicons and quantitative PCR (qPCR), followed by construction of shotgun metagenomes. The analysis included evaluation of the metabolic pathways potentially involved in the transformation of permafrost organic carbon, and their seasonal persistence or change.

## MATERIALS AND METHODS

### Site Description

The Sasapimakwananisikw (SAS) River Valley is located in the sporadic permafrost zone of subarctic Quebec, around 10 km southwest of the village Whapmagoostui-Kuujuarapik, Quebec, Canada. The region experiences contrasting seasons, with a winter minimum air temperature of  $-42^{\circ}\text{C}$  and a summer maximum of  $33^{\circ}\text{C}$  (CEN, 2017). The SAS valley contains numerous palsas, which are raised permafrost mounds in the peatland, and associated thermokarst lakes and ponds of different ages that formed over the last 60 years due to rapid permafrost thawing (Payette et al., 2004; Vincent et al., 2017). Lake SAS2A ( $55^{\circ}13.160' \text{ N}$ ,  $77^{\circ}41.806' \text{ W}$ ) is a circular thermokarst lake surrounded by semi-aquatic plants (notably *Carex aquatilis* and *Sphagnum* mosses) with an area of  $196 \text{ m}^2$  and a maximum depth of 2.8 m. Ice and snow conditions were monitored using automated cameras and sensors installed on the edge of the lake. In 2015–2016, the lake surface froze in mid-October 2015 and ice persisted until the end of May 2016 (details in Matveev, 2018 and CEN, 2017). The maximal snow (0.5 m) and ice (0.6 m) depth coincided with the time of our winter sampling in mid-March. Environmental data for lake SAS2A, including details on geochemistry (methane, carbon dioxide, dissolved

organic carbon, sulfate and hydrogen sulfide concentrations) and limnological profiles (oxygen, pH, temperature, conductivity) measured at our two sampling times, are given in Matveev (2018) and Matveev et al. (2019), and are briefly described below.


In summer 2015, lake SAS2A was darkly colored, with high concentrations of dissolved organic carbon ( $13.7 \text{ mg DOC L}^{-1}$ ). This colored dissolved organic matter (CDOM) absorbs solar radiation, with <1% of the solar irradiance at below 0.7 m (Matveev et al., 2016) (Figure 1). The lake was slightly acidic (pH 6) in summer, with a drop in temperature, from  $15^\circ\text{C}$  at the surface to  $6^\circ\text{C}$  below 0.75 m. Although the surface waters are episodically mixed by wind, the lake is usually stratified throughout summer, with oxygen concentrations around  $4.13 \text{ mg L}^{-1}$  at the surface and values below the detection limit from 1 m to the water column bottom (Matveev, 2018). The lake is known to have high emission rates of  $\text{CH}_4$  (Matveev et al., 2016). Methane concentrations increased with depth, ranging from  $2.5 \mu\text{M}$  at the surface to  $300 \mu\text{M}$  at the bottom of the water column with  $20 \mu\text{M}$  of methane at 0.5 m (Matveev et al., 2019). The  $\text{CH}_4$  emitted during summer has been  $^{14}\text{C}$ -dated at 1360 years before present, indicating breakdown of old permafrost-derived soil organic carbon (Matveev et al., 2016). Sulfate concentrations were low at the surface at around  $1.46 \text{ nM}$ , and below detection throughout the rest of the water column (Figure 1).

During winter 2015–2016, lake SAS2A was covered with ice and snow. At the time of sampling, water temperatures increased slightly with depth, from  $0^\circ\text{C}$  immediately beneath the ice to  $3.5^\circ\text{C}$  at the bottom of the lake. The lake was more acidic in winter (pH 5), and oxygen was below detection and methane concentrations high (around  $200 \mu\text{M}$ )

throughout the water column (Figure 1). Under-ice sulfate and hydrogen sulfide concentrations were up to  $0.50$  and  $10.79 \text{ nM}$  respectively (J. Canario, University of Lisbon, personal communication). DOC and total nitrogen concentrations were higher in winter than summer, up to  $18.3$  and  $1.2 \text{ mg L}^{-1}$ , respectively (Figure 1; Matveev, 2018). Comparison of the upper water columns of three lakes (SAS2A and nearby SAS2B and SAS2C) indicated that SAS2A is likely to be representative of the humic-rich thermokarst waters throughout the valley, and that the chemical differences, including mean pH, methane and oxygen concentrations, are large and significant between seasons (all *t*-tests:  $p < 0.003$ ; full profile data available in Matveev et al., 2019).

## Sample Collection and Nucleic Acid Extraction

Lake SAS2A was sampled in summer (24 August 2015) and winter (18 March 2016), at the same time as the limnological and geochemical characterization of the lake as shown in Figure 1 and described above. Sampling was with a 3-L Limnos Water sampler (KC, Denmark) that was previously washed with 10% HCl, rinsed with sterile MilliQ water, maintained closed until sampling and then rinsed three times with the lake water prior to sample collection. For the summer sampling, water was collected at 0.5 m below the surface (measured from the middle of the sampler), corresponding to the metalimnion/hypolimnion transition zone (oxycline), at three mid-lake sites. For the winter samples, snow cover (0.5 m) was removed and three independent 24 cm-diameter holes were drilled through the ice (0.6 m thickness) near the middle of the lake. The Limnos Water sampler was



lowered into the ice holes and tripped immediately below the ice. For both the summer and winter campaigns, approximately 300 mL of each triplicate water sample were filtered through separate 0.22  $\mu\text{m}$  Sterivex filters and then stored below  $-50^{\circ}\text{C}$  until nucleic acid extraction.

Nucleic acids (DNA and RNA) were extracted from the same Sterivex filters using Qiagen Allprep DNA/RNA Mini Kit with modifications as described in Cruaud et al. (2017). The Sterivex cartridges were opened and the membrane filters were cut into small pieces before the lysis steps. All steps of the nucleic acid extractions, from the opening of the filters to the nucleic acid resuspension in autoclaved, filtered and UV-treated water, were carried out in a sterile laminar flow cabinet. The DNA extracts were stored at  $-20^{\circ}\text{C}$  until library preparation. For RNA extracts, two additional DNase steps (DNase I, Ambion, Foster City, CA, United States) were carried out to remove any trace of carried over DNA. The absence of DNA contamination was confirmed by amplification of 16S rRNA genes with bacterial primers using the RNA extracts (undiluted and diluted ten times) as template, with no product detected after 35 PCR cycles. The RNA was then immediately converted to cDNA using a High-Capacity cDNA Reverse Transcription kit (Applied Biosystems, Foster City, CA, United States) and stored as cDNA at  $-20^{\circ}\text{C}$  until library preparation.

## Quantitative Polymerase Chain Reaction (qPCR) Analysis

The number of Bacteria and Archaea 16S rRNA genes in the samples from the two seasons were estimated using quantitative PCR (qPCR), with primers Bact1369f/Bact1492r and Arc787f/Arc1059r, respectively (Vigneron et al., 2013). Quantification was performed in triplicate with a range of template concentrations (0.5, 1, 1.5 ng of DNA) to compensate for any PCR inhibition. Amplification reactions were carried out in a 7500 Fast Real-Time system (Applied Biosystems) in a final volume of 25  $\mu\text{L}$  using Brilliant III Ultra-Fast Master Mix (Agilent, Santa Clara, CA, United States), 0.5  $\mu\text{M}$  of each primer and 5  $\mu\text{l}$  of DNA template. qPCR conditions were as follows: 40 cycles of denaturation at  $95^{\circ}\text{C}$  for 15 s then annealing and extension at  $60^{\circ}\text{C}$  for 60 s. Standard curves were prepared in triplicate with dilutions ranging from 0.001 to 100 nM of DNA extracted from *Methylobacterium methanica* (DSM25384) and *Methanosarcina acetivorans* (DSM2834), corresponding to  $10^2$  to  $10^6$  16S rRNA gene copies per reaction. The  $R^2$  of standard curves obtained by qPCR were above 0.99, PCR efficiencies were above 90.5%, and melting curves showed no trace of non-specific amplifications. The qPCR results were expressed in terms of 16S rRNA gene numbers per milliliter of lake water.

## Illumina MiSeq Amplicon Library Preparation, Sequencing and Analysis

Microbial community composition of lake SAS2A in the two seasons was determined by high throughput sequencing of bacterial and archaeal 16S rRNA (cDNA) and 16S rRNA genes (DNA) using primers targeting the bacterial V4-V5

region (S-D-Bact-0516-a-S-18/S-D-Bact-0907-a-A-20; 460 bp product) (Klindworth et al., 2013) and the archaeal V1-V3 region (A27F/Arc518R; 500 bp), respectively (Teske and Sorensen, 2007). All PCR reactions were carried out following Vigneron et al. (2017). Amplicons were sequenced using an Illumina MiSeq v3 kit at the IBIS/Laval University, Plateforme d'Analyses Génomiques (Québec, QC). Reads were assembled into single paired-end sequences and curated as detailed in a GitHub repository<sup>1</sup>. Taxonomic affiliations of the reads were determined with Mothur (Schloss et al., 2009) using BLAST against Silva database release 128 as reference (Pruesse et al., 2007).

## Metagenomic Library Preparation, Sequencing and Analysis

Three metagenomes were constructed for winter and summer samples (total of 6 metagenomes) using a Nextera XT Library Kit (Illumina, San Diego, CA, United States) according to the manufacturer's recommendations at the CGEB - Integrated Microbiome Resource, Dalhousie University. Sequencing was performed using four lanes of an Illumina NextSeq system at the CGEB, Dalhousie University. Barcode and adapter sequences were removed from the metagenome data on-instrument using Illumina's MiSeq Reporter software and the sequence data were exported as FASTQ files. Datasets were quality filtered using the Trimmomatic tool (Bolger et al., 2014), with default settings. Paired-end joining was done using FLASH2 (Magoč and Salzberg, 2011). The 16S rRNA genes (on average  $n = 30795$  per sample) were isolated from metagenomic reads using REAGO 1.1 (Yuan et al., 2015), and taxonomic assignments were performed as for the 16S rRNA gene amplicons.

Each metagenome was assembled separately from paired-end reads after quality filtering using IDBA\_UD (Peng et al., 2012). Assembled contigs and mapping files (BAM files) were uploaded to the Department of Energy Joint Genome Institute (DOE-JGI) IMG/MER analysis pipeline (Markowitz et al., 2009) for gene calling and functional annotation. To account for differences in sequencing depth between samples, metagenomes were normalized by multiple rarefaction (100 iterations) to the size of the smallest dataset (218503 genes). For metagenome assembled genome reconstruction, all quality filtered sequences were pooled and co-assembled using MEGAHIT (Li et al., 2015), which was computationally less expensive than other options and appropriate for our dataset. Read coverage of the contig was carried out using bwa-mem<sup>2</sup>, then contig binning was done using MetaBAT-1 and -2 (Kang et al., 2015) and Maxbin 2.0 (Wu et al., 2016) with contigs longer than 3000 bp. In addition, binning of the single sample-assembled contigs (longer than 3000 bp) was also performed using Vizbin (Laczny et al., 2015). The multiple binning tools increased the number of bins. Redundant bins obtained from the 4 binning algorithms were identified and removed using CheckM (Parks et al., 2015). The completeness and

<sup>1</sup>[https://github.com/CruaudPe/MiSeq\\_Multigenique](https://github.com/CruaudPe/MiSeq_Multigenique)

<sup>2</sup><http://bio-bwa.sourceforge.net>

contamination level of the combined genomic bins were then evaluated using CheckM (Parks et al., 2015). Only bins with a contamination level under 5% and completeness above 50% were analyzed. Genetic composition of genomic bins was then explored using KEGG (Kanehisa et al., 2008) and MetaCyc (Caspi et al., 2014) based on genes identified by IMG/MER in the co-assembly. The results were manually checked and pathways were considered as present when 75% of the genes involved in the pathway were detected. The presence of specific pathways was also determined by detection of key genes (in brackets in **Figure 4**). The presence of genes for carbohydrate-active enzymes was investigated using dbCAN (Yin et al., 2012) and the CAZy database (Cantarel et al., 2009). Assembled metagenome data are available in IMG/MR under the following accession numbers: 3300022653, 3300022650 and 3300022591-4. Raw sequences were deposited in the NCBI public database under Bioproject PRJNA515027 and bin files are available in FigShare (doi: 10.6084/m9.figshare.8132546).

## Statistical Analyses

Statistical analyses of the data set [Student *t*-test, NPMANOVA, similarity percentages breakdown procedure (SIMPER)], Bray-Curtis-based dissimilarity index calculations and correlation-based clustering were carried out according to recommendations of the Guide to Statistical Analysis in Microbial Ecology (Buttigieg and Ramette, 2014), using PAST software (Hammer et al., 2001). Correlation matrices were calculated using microbial lineage relative proportions, then clustering of microbial lineages shown in **Figure 2** was determined by UPGMA based on Euclidian distances generated from these correlation matrices.

## RESULTS

### Microbial Community Composition

The microbial community composition and abundance in lake SAS2A was determined by 16S rRNA (cDNA), rRNA gene (DNA) amplicon sequencing, quantitative PCR and metagenomic shotgun sequencing. Overall, there was no significant difference in the total concentration of 16S rRNA genes (Bacteria plus Archaea) between summer and winter (*t*-test,  $p = 0.85$ ; **Figure 2A**). These values averaged around  $4.8 \pm 3 \times 10^5$  16S rRNA genes  $\text{mL}^{-1}$  for summer samples and for  $4.3 \pm 3 \times 10^5$  16S rRNA genes  $\text{mL}^{-1}$  for winter samples. However, the concentrations of archaeal gene copies were significantly greater (15 times higher; *t*-test,  $p < 0.001$ ) in winter than in summer, with up to  $6 \times 10^4$  16S rRNA genes  $\text{mL}^{-1}$  (**Figure 2B**). In summer, Archaea accounted for only 1% of the total prokaryotes in both the qPCR and metagenomic analyses, but 7% (qPCR) and 17% (metagenomes) in winter.


The winter archaeal community, determined from the 16S rRNA genes in the metagenomes, was mainly dominated by Methanomicrobiales (50% of the Archaea), Methanosarcinales (14%) and other co-varying potential methanogenic lineages and Bathyarchaeota (**Figure 2B**). In contrast, the archaeal community in summer had a lower contribution of Methanomicrobiales (22%), and a larger contribution of

Woesarchaeota (34% of the Archaea), and Miscellaneous Euryarchaeal Group (MEG, 11%).

Analysis of metagenomic 16S rRNA genes showed that bacterial community composition also differed between winter and summer samples (Bray Curtis dissimilarity index: 73.27, NPMANOVA:  $p = 0.05$ ). The summer bacterial community was dominated by Betaproteobacterial members of the Burkholderiales (27% of the bacterial 16S rRNA genes), *Comamonadaceae* (12%) and *Methylophilaceae* (3%); Actinobacteria (17%), Parcubacteria (10%) and other co-varying lineages of the Gammaproteobacteria including Methylococcales (**Figure 2C**). The relative proportions of these lineages were statistically greater in summer samples (*t*-tests,  $p < 0.005$ ; bottom cluster in **Figure 2C**). In contrast, in winter samples, members of the Planctomycetes (15%), Chloroflexi (14% with *Anaerolineae*, *Dehalococcoidia* and *Chloroflexi*), Microgenomates (9%), Deltaproteobacteria as Syntrophobacterales (6.4%), and Omnitrophica (5%) had a significantly greater representation compared to summer samples (*t*-tests,  $p < 0.05$ ; **Figure 2C**). Although the relative proportions of the Parcubacteria and Microgenomates were significantly different between seasons, the two phyla represented a major component of the bacterial community in all samples (**Figure 2C**). Finally, amplicon sequencing of the 16S rRNA gene using DNA and RNA templates showed that most of the detected lineages were potentially active, or at least maintained ribosomes in their cells in both summer and winter seasons (**Figure 2C**, **gray** and **pink** rectangles next to the heatmap).

### Overall Microbial Metabolic Potential

The shotgun metagenomes from summer ( $n = 3$ ) and winter ( $n = 3$ ) samples were used to explore the metabolic potential of the permafrost thaw lake. Up to  $8.15 \times 10^5$  genes with predicted functions were identified per sample (average:  $7.16 \times 10^5$  genes). These genes were distributed into at least 6500 different molecular-level functions (KEGG Orthology). The metagenomic datasets were normalized by multiple rarefaction (see Methods section) then compared. The metabolic potential for degradation of complex organic molecules (e.g., aromatic compounds, glycans, starch, cellulose, hydrocarbons) was present in all samples (**Figure 3**). In contrast, different potential energetic pathways were detected between winter and summer samples, notably in the methane and sulfur cycles (**Figure 3**). In the summer samples, which were collected from the oxycline, the potential for oxygen respiration and aerobic metabolism (sulfide, thiosulfate, methane and methanol oxidation) was predominant. The potential for light utilization through photosynthesis (*chlG*) and proton pumping with proteorhodopsin (*pro*) was also evident (**Figure 3**). In contrast, in the winter sampling, when no oxygen was detected in the water at the time of collection, a considerable potential for reductive metabolism was identified. The winter microbial community harbored the potential for sulfate (*dsrB*), sulfite (*asrB*), thiosulfate (*phsA*), dimethyl-sulfate (*dmsA*), polysulfide (*hydB*), nitrate (*narG*), nitrite (*nrfA*), selenate (*xdhD*), arsenate (*arrA*), trimethylamine oxide (*torZ*) and chelated



**FIGURE 2 | (A)** Bacterial and archaeal abundance determined by quantitative PCR in all summer (S1, S2, S3) and winter (W1, W2, W3) samples. **(B)** Archaeal community composition in summer (S1, S2, S3) and winter (W1, W2, W3) samples. MGBD: Marine Benthic Group B; MEG: Miscellaneous Euryarchaeotal Group. **(C)** Bacterial community composition in summer (S1, S2, S3) and winter (W1, W2, W3) samples. Heat maps of the relative proportion of microbial lineages are based on 16S rRNA genes from metagenomic dataset. Clustering of the microbial lineage relative proportion (constructed by UPGMA with Euclidian distance on the correlation matrix generated using microbial lineage relative proportions) gather lineages with similar relative proportion in the samples. Gray rectangles (detected by PCR on DNA template) and purple rectangles (detected by PCR on cDNA template) indicates the detection of a given lineage by 16S rRNA (gene) amplicon sequencing whereas lineages with yellow squares were not detected by amplicon sequencing (details in **Supplementary Figure 1**). Differences in the relative proportion between summer and winter samples were evaluated by *t*-tests; the significance of the test is presented as: -, not-significant; \**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.0005. The number of genomic bins reconstructed per lineage is indicated by the green circles.


iron (*feR*) reductions along with methanogenesis (*mcrA*) and dehalogenation (*dhaA*, **Figure 3**).

## Genomes Assembled From Community Metagenomes

Genomic bins corresponding to representative microbial lineages in the permafrost thaw lake were assembled, and metabolic

capabilities were inferred from the genomic bins with >50% completeness and contamination levels of <5% (**Figure 4**). The recovered genomic bins exhibited contrasting metabolic potentials, which reflected their taxonomic affiliation and the seasonal environmental conditions of the thermokarst lake.

Genomic bins recovered from the summer samples were largely affiliated to taxa within the phylum Actinobacteria with




15 bins, including the genera *Actinobacteria*, *Rhodoluna* and *Ca. Planktophilia* and *Ca. Nanopelagicus* (Figure 4). The potential for bacteriorhodopsin phototrophy and oxygen respiration was identified in most of the *Actinobacteria* genomic bins. Different types of cytochromes for oxygen respiration (types 1, cbb3 and bd) were detected. The potential for thiosulfate oxidation through thiosulfate:quinone oxidoreductase was also widespread in the *Actinobacteria* bins (Figure 4). Regarding carbon metabolism, the potential for degradation of cyanophycin and complex glycans was identified along with the pentose phosphate pathway, glycolysis and citrate cycle, which permit carbohydrate assimilation (Figure 4). In addition, the potential for anaplerotic CO<sub>2</sub> fixation (phosphoenolpyruvate carboxylase and pyruvate carboxylase genes), ethanol fermentation, thiosulfate and arsenate reduction was also present in most of the *Actinobacteria* genomic bins. Three large genomic bins affiliated to *Betaproteobacteria* were also recovered (*Burkholderia* and *Limnochabitans*). Numerous oxidative metabolic functions were identified in these bins, including oxygen respiration, thiosulfate oxidation through the Sox pathway and metal oxidation via multiheme cytochrome C. A strong metabolic potential for degradation of various organic compounds, including urea, hydrocarbons and aromatic compounds was also identified (Figure 4). Most catabolic pathways (pentose phosphate pathway, glycolysis, Entner-Doudoroff pathway, acetate kinase pathway, citrate cycle) were also detected

in these lineages. Genomic bins affiliated to *Methylobacter* (Gammaproteobacteria) and *Cyanobacteria* were also assembled. The capacity for thiosulfate reduction as well as methane oxidation was detected in the *Methylobacter* genomic bin, and photosynthetic functions were evident in the *Cyanobacteria*, as expected (Figure 4).

Genomic bins assembled from the winter samples were related to 13 different taxa, representing the dominant winter microbial lineages (Figures 2B,C, 4). The potential for hydrogen, acetate and pyruvate metabolism was widespread in the winter community. Planctomycetes genomic bins (*n* = 5) harbored a strong potential for organic matter degradation (cellulose, starch, xylan and pectin) associated with numerous genes for carbohydrate degradation (up to 192 carbohydrate-active enzymes (CAZy) genes per bin; Figure 4). The potential for ethanol fermentation, acetogenesis and organohalide and arsenate respiration was also detected in most Planctomycetes bins (Figure 4). Six genomic bins affiliated to *Chloroflexi* (*Chloroflexi* and *Anaerolineae*) were assembled. Different metabolic potentials were identified within this group. *Anaerolineae* bins showed the genetic potential for micro-oxygen and thiosulfate respiration, organohalide respiration and lactate fermentation coupled with the potential to degrade acid phenol, cyanophycin, glycans, cellulose and starch. Two other *Chloroflexi* bins had the potential for sulfate, arsenate and organohalide reduction, hydrogenotrophy, and various fermentations coupled





to genes involved in hydrocarbon and aromatic compound degradations. In addition, another Chloroflexi bin showed the potential for anoxygenic phototrophy (Figure 4). Such potential for light utilization was also detected in two bins affiliated to Chlorobi (*Pelodiction*), where this potential was associated with sulfur and thiosulfate oxidation and CO<sub>2</sub> fixation through the Calvin-Benson-Bassham cycle (Figure 4). The potential for thiosulfate reduction, hydrogenotrophy and acetogenesis was identified in two Aminicenantes genomic bins. A strong potential for complex carbon substrate degradation

(130 CAZY genes per bin) was also present in Aminicenantes genomic bins, along with the potential for multiple fermentation pathways (Figure 4). Acetothelia related bins contained genes for acetogenesis and formate fermentation, but a limited potential for organic carbon degradation (Figure 4). Two genomic bins related to *Desulfobacterium* lineages showed the potential for sulfate, thiosulfate, DMSO, arsenate and organohalide reduction, numerous fermentations and degradation of hydrocarbons and aromatic compounds (Figure 4). The Bacteroidetes (Bacteroidetes and Flavobacteriales) bins harbored the potential

for sulfate, sulfur and selenate reduction and micro-oxygen respiration, suggesting a facultative aerobic lifestyle. Finally, many genes for carbohydrate degrading enzymes were identified in all of these genomic bins (114 CAZy genes, **Figure 4**). Single bins of Omniphilica, Verrucomicrobia, *Gallionella* were also assembled. Archaea were also represented with a genomic bin related to the Methanomicrobiales, with genes for hydrogenotrophic ( $H_2:CO_2$ ) methanogenesis (**Figure 4**).

Finally, eight genomic bins were recovered from the Patescibacteria/candidate phyla radiation (*Parcubacteria*, *Ca. Moranbacteria*, *Ca. Dependientiae* and *Microgenomates*). These represented a substantial proportion of both winter and summer microbial communities, but had a limited number of genes with known functions. With the exception of a gene coding for a thiosulfate:quinone oxidoreductase in a *Parcubacteria* bin, no respiratory pathways were identified. The potential for simple carbohydrate assimilation through glycolysis or pentose phosphate pathway was detected, as well as the potential for lactate fermentation in *Parcubacteria* and *Ca. Dependientiae*. A few genes for carbohydrate degradation enzymes were also present in all Patescibacteria/CPR genomic bins (average of 30 CAZy genes; **Figure 4**).

## DISCUSSION

In this study, we compared the winter and summer microbial communities of a thermokarst lake to evaluate whether the winter season induces dormancy of the microbial community or, in contrast, is the time of functional reassembly into a winter-active community. Contrary to the hypothesis of a microbial downshift or dormancy in winter, the qPCR comparison of the ice-covered and ice-free periods indicated no statistical change in total prokaryotic abundances between seasons, and in fact there was an order of magnitude increase in archaeal abundance (**Figure 2A**). Furthermore, the amplicon sequencing based on cDNA indicated that with few exceptions, all detected microbial lineages were potentially active or ready to respond to favorable environmental conditions (pink rectangles in **Figures 2B,C**, details in **Supplementary Figure 1**). For the two seasons, a large panoply of microorganisms was identified by the 16S rRNA gene and metagenomic sequencing, including Actinobacteria, Chloroflexi, Planctomycetes, Proteobacteria, candidate phyla, Methanomicrobiales and Woesearchaeota (**Figures 2B,2C**). Additional phyla, notably Microgenomates, Aminicenantes and Acetothermia, were solely identified by PCR-free metagenomic sequencing (yellow squares in **Figure 2**). These latter phyla collectively represented a third of the thermokarst lake microbial community in both summer and winter samples, yet were not detected in previous reports based on 16S rRNA gene amplicons (Crevecoeur et al., 2015; Wurzbacher et al., 2017). This may be due to the presence of introns in their 16S rRNA genes that precluded PCR amplification with conventional 16S rRNA gene primer sets (Brown et al., 2015). These results highlight the value of a metagenomic approach for thermokarst lake community composition analysis, and they show that SAS2A is a rich microbial habitat containing diverse taxa

that likely maintain a substantial biomass and productivity throughout the year.

## Seasonal Contrasts

Consistent with the hypothesis of an alternate community developing under the ice, the clustering of the microbial lineages based on their relative abundance in the metagenomic dataset indicated a fundamental shift in community composition between summer and winter (NPMANOVA:  $p = 0.05$ ; **Figure 2**). This major shift in taxonomic composition was accompanied by large differences in potential functioning of the community, with contrasting, and sometimes opposing, metabolic pathways between seasons (**Figure 3**).

The summer microbial community was dominated by bacteria (99% of the 16S rRNA genes from qPCR and metagenomes) with predominant members of the Actinobacteria and Betaproteobacteria lineages (**Figure 2**), as previously reported in other thermokarst lakes (Comte et al., 2016) and freshwater ecosystems in general (Newton et al., 2006, 2007). The recovery of multiple Actinobacteria genomic bins (**Figure 4**) would be consistent with micro-diversification in genomic composition, suggesting a strong diversity within this lineage. This diversity would facilitate rapid responses to environmental fluctuations (Neuenschwander et al., 2017) that would arise from near-surface stratification and periodic mixing in the thermokarst lake environment during summer. In contrast, the genomic bins of Betaproteobacteria suggested a different ecological strategy to that observed in Actinobacteria, with multiple pathways of energy acquisition, suggesting a metabolic versatility to cope with environmental fluctuations. Consistent with ample light and oxygen in the upper water column during summer (**Figure 1**), the overall metabolic capacities of the microbial community were centered on phototrophy, and aerobic metabolism (oxygen respiration and oxidation of methane, methanol, thiosulfate and  $H_2S$ ) coupled to the degradation of particulate and photodegraded organic matter (*mhpA* in **Figure 3**). Methane, methanol,  $H_2S$  and thiosulfate oxidations are likely fuelled by the release of methane and other metabolic products from the sediments and the anoxic layer of the water column. This potential for methane oxidation and the larger  $CO_2$  emissions reported in summer supports the notion that the greenhouse gas balance of lake SAS2A is oriented toward net  $CO_2$  production during open water conditions (Laurion et al., 2010), and that the microbial community in the upper water column acts as a biofilter that dampens summer methane emissions, as previously suggested (Crevecoeur et al., 2017).

In the winter season, ice cover forms a diffusive barrier with the atmosphere, leading to the rapid (~10 days after ice-cover formation) depletion of oxygen and anoxia in the water column (**Figure 1**; Deshpande et al., 2017). Consequently, reduced compounds produced in the anoxic bottom of the lake are not recycled by the oxidative processes observed during summer. The microbial community composition under the ice shifted to members of the Planctomycetes, Chloroflexi, Deltaproteobacteria, Candidate phyla and Methanomicrobiales (**Figures 2A,B**). The metabolic capacities of the winter community also reflected this seasonal shift, with various

anaerobic pathways such as methanogenesis, consistent with higher methane concentrations in the water and the reduction of potential oxidants (Figure 3). Genomic bin reconstruction indicated that nitrate, sulfate, iron, arsenate, and selenate can be reduced by the winter community (Figure 4), potentially leading to a strongly reduced environment that would further allow the methanogens to flourish (Figure 2). These reduction pathways appeared to be coupled to complex organic matter degradation, fermentations, and the production and consumption of hydrogen, formate and acetate (Figures 3, 4). This suggests that syntrophy, defined as a metabolic interaction between dependent partners, and complementary metabolism (facultative cross-feeding) between methanogens and their putative syntrophic and fermentative partners (*Syntrophobacteraceae*, Chloroflexi; de Bok et al., 2004; Liang et al., 2015) are likely to be major ecological strategies of the winter microbial community. The net effect of such syntrophy and cross-feeding would be an enhanced capacity to degrade recalcitrant organic matter in the isolated and reduced environmental conditions of the ice-covered thermokarst lake. These results, coupled with the winter chemical conditions indicate a seasonal partitioning of the biogeochemical cycles in the thermokarst lake, where methane and potentially reduced molecules accumulate under ice, leading to a massive release of greenhouse gasses when the ice breaks up in spring (Deshpande et al., 2017). However, aerobic methanotrophic lineages, such as members of the *Methylococcaceae* were detected in winter (Figure 2C), suggesting that they might survive winter using alternate metabolic pathways such as thiosulfate or arsenate reduction and fermentations, as identified in their genomic bins (*Methylobacter* bin in Figure 4). This overwintering population of methanotrophs could potentially moderate atmospheric methane emissions from the lake in spring.

These results indicate that the thermokarst lake community has a high potential for rapid adjustments in response to episodic and seasonal changes in environmental conditions such as oxygen, temperature and mixing. An important consequence of such flexibility in composition and function is that analyses in summer of community structure, metagenomic features and microbial activities such as gas production cannot be considered representative of the full annual cycle. They relate to a specific, transient stage of 4 months per year, while the overall community structure and metabolism of thermokarst lakes may be mostly defined by microbial characteristics during the prolonged winter season. These results also imply that accurate estimates of annual carbon fluxes and biogenic gas emissions from thermokarst lakes will require greater attention during the ice-cover and break-up periods.

## Winter Pathways of Permafrost Organic Carbon Degradation

Thermokarst lakes are enriched in terrestrial organic carbon, with up to 96% of the carbon of the DOM originating from terrestrial (thawed permafrost) origin and negligible contribution from phytoplankton and macrophytes (Wauthy et al., 2018). The high DOM concentrations measured in the present study during winter imply an abundance of permafrost-derived organic

substrates for microbial degradation processes under the ice. Our metagenomic results indicate that the winter community had the potential for mobilizing and converting this complex organic matter to more labile carbon sources. For example, genes coding for hydrolytic arylsulfatases (*aslA* in Figure 3), which hydrolyse sulfate ester and release sulfate and phenols (Kertesz, 2000), or hydrolytic dehalogenases (*dhaA*, *had*), which cleave halogen groups from carbon substrates (Koudelakova et al., 2013), were abundant in the winter metagenomes and occurred in numerous genomic bins (Figures 3, 4). Sulfate released by sulfatase activity might be used by some members of the Deltaproteobacteria, Chloroflexi and Bacteroidetes that have the metabolic potential for sulfate and sulfite reduction (*dsr* and anaerobic sulfite reductase gene *asr* in Figure 4), contributing to the high concentrations of H<sub>2</sub>S measured in the water. The production of H<sub>2</sub>S and fermentation products such as fatty acids and CO<sub>2</sub> likely explain the observed acidification of the water during winter. This acidification has important consequences, since it would also enhance the mobilization of labile carbon, iron, arsenate and selenate from colloidal and complexed organic matter (Audry et al., 2011; Pokrovsky et al., 2011). Metagenomic analysis showed a strong potential for the respiration of arsenate and selenate (*arrA* and *xdhD* in Figures 3, 4), suggesting that acidic mobilization of such elements could provide an additional energy source for permafrost-degrading microorganisms.

The main winter lineages (Aminicantantes, Bacteroidetes, Chloroflexi, Deltaproteobacteria, Planctomycetes and Verrucomicrobia) had a strong catabolic potential centered on the degradation of aromatic carbon and other plant detritus, which constitute the main organic matter of permafrost soils (Jansson and Taş, 2014). The catabolic potential of the reconstructed bins related to these lineages (average of 90 CAZy genes per bin) far exceeded the catabolic potential detected in bins associated with the summer community members (average of 42 CAZy genes per bin) and is consistent with the known properties of cultured representatives (Yamada et al., 2005; Wagner and Horn, 2006; Meckenstock et al., 2016) and previous genomic investigations (Hug et al., 2013; Tran et al., 2018; Woodcroft et al., 2018). However, different substrate specificities (cellulose, xylan, pectin, starch, esters, aromatic compounds) were detected among these winter lineages (Figure 4), suggesting multiple ecological niches and nutritional diversification. Considerable potential for the fermentation of catabolic products associated with mechanisms for hydrogen, formate and acetate transfer and utilization was identified in the microbial community, suggesting that syntrophic degradation and co-metabolism represent major pathways for the decomposition of permafrost organic carbon. This is also supported by the high *in situ* concentrations of fermentation intermediates identified in Arctic and subarctic peat (Tveit et al., 2015). These fermentative lineages are likely coupled with the methanogens (Figure 2) that consume hydrogen, as identified in a genomic bin of the Methanomicrobiales or acetate for the members of the Methanosarcinales (Jetten et al., 1992). This high potential for methanogenesis coupled with syntrophic degradation of the permafrost carbon is consistent with artificial thawing experiments (Tveit et al., 2015) and explains the elevated

methane concentrations measured throughout the SAS2A water column in winter (Matveev et al., 2019).

In combination, our results indicate that catabolic diversification, fermentation pathways, syntrophic methanogenesis, and mobilization of recalcitrant organic matter and electron acceptors underpin the decomposition of permafrost-derived organic matter in lake SAS2A, and probably in many other thermokarst lakes. The detection of these processes implies a need for additional work. For example, although candidate phyla (*Parcubacteria*, *Microgenomates*) represented up to 15% of the total microbial community (Figure 2) and were detected throughout the year, suggesting a major role in the ecosystem, the metabolic functions of these taxa remain unclear. Genomic bin reconstruction highlighted that most of the enzymatic potential of these lineages differed from other bacterial lineages since only few genes were identified, as previously reported (Brown et al., 2015). The number of genes coding for carbohydrate degrading enzymes (CAZy in Figure 4) was low in *Patescibacteria*/CPR lineages, however, a potential for chitin or starch degradation was detected in the most complete genomic bins (Figure 4), indicating a possible role in permafrost carbon degradation.

## CONCLUSION

Previous work on thermokarst lakes has identified their importance for microbial activity and greenhouse gas production in northern landscapes, however, most studies to date have focused on the summer open water period. The present observations show that the winter microbiome differs greatly from that in summer in terms of both microbial composition and metabolic functions. The major shifts induced by prolonged ice cover in the winter (8 months per year) result in the seasonal partitioning of biogeochemical cycles, and favor the winter accumulation of methane and other reduced compounds. These results underscore the need for sampling thermokarst lakes in all seasons. They draw attention to the importance of winter at northern high latitudes, and show how this can be a period of diverse microbial pathways for the mobilization of organic matter and energy supply, notably fermentation processes, cross-feeding and syntrophic methanogenesis. Our observations imply that throughout much of the year, thermokarst lakes operate

as cold, ice-capped methanogenic brews fuelled by permafrost-derived organic carbon.

## DATA AVAILABILITY

The datasets generated for this study can be found in NCBI, PRJNA515027.

## AUTHOR CONTRIBUTIONS

AV, CL, AC, and WV designed the research. AV and PC analyzed the data with contributions from DK. AV, CL, PC, and WV led the writing of the manuscript. All authors provided comments on draft versions of the manuscript.

## FUNDING

This work was supported by the program Sentinel North financed by the Canada First Research Excellence Fund (CFREF); the Natural Sciences and Engineering Research Council of Canada (NSERC); the CNR (Italy) – Université Laval UMI-MicroMeNu program; the Fonds de Recherche du Québec-Nature et Technologies (FQRNT); and the Network of Centres of Excellence ArcticNet.

## ACKNOWLEDGMENTS

We thank Isabelle Laurion for coordinating the winter sampling activities, Alex Matveev, Isabelle Laurion, and João Canario for geochemical data and help in the field. We also thank Alice Lévesque for sampling, the Whapmagoostui-Kuujuuarapik CEN station and Canadian Helicopter Ltd., for logistic support, and Calcul Quebec for computing resources and technical support.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2019.01656/full#supplementary-material>

## REFERENCES

- Audry, S., Pokrovsky, O. S., Shirokova, L. S., Kirpotin, S. N., and Dupré, B. (2011). Organic matter mineralization and trace element post-depositional redistribution in Western Siberia thermokarst lake sediments. *Biogeosciences* 8, 3341–3358. doi: 10.5194/bg-8-3341-2011
- Bertilsson, S., Burgin, A., Carey, C. C., Fey, S. B., Grossart, H.-P., Grubisic, L. M., et al. (2013). The under-ice microbiome of seasonally frozen lakes. *Limnol. Oceanogr.* 58, 1998–2012. doi: 10.4319/lo.2013.58.6.1998
- Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120. doi: 10.1093/bioinformatics/btu170
- Bouchard, F., Francus, P., Pienitz, R., Laurion, I., and Feyte, S. (2014). Subarctic thermokarst ponds: Investigating recent landscape evolution and sediment dynamics in thawed permafrost of Northern Québec (Canada). *Arct. Antarct. Alp. Res.* 46, 251–271. doi: 10.1657/1938-4246-46.1.251
- Brown, C. T., Hug, L. A., Thomas, B. C., Sharon, I., Castelle, C. J., Singh, A., et al. (2015). Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* 523:208. doi: 10.1038/nature14486
- Buttigieg, P. L., and Ramette, A. (2014). A guide to statistical analysis in microbial ecology: a community-focused, living review of multivariate data analyses. *FEMS Microbiol. Ecol.* 90, 543–550. doi: 10.1111/1574-6941.12437
- Cantarel, B. L., Coutinho, P. M., Rancurel, C., Bernard, T., Lombard, V., and Henrissat, B. (2009). The carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* 37, D233–D238. doi: 10.1093/nar/gkn663
- Caspi, R., Altman, T., Billington, R., Dreher, K., Foerster, H., Fulcher, C. A., et al. (2014). The MetaCyc database of metabolic pathways and enzymes and

- the BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* 42, D459–D471. doi: 10.1093/nar/gkt1103
- CEN (2017). *Climate Station Data from Whapmagoostui-Kuujuuarapik Region in Nunavik*. Canada, CA: CEN, v. 1.4, (1987–2016).
- Comte, J., Monier, A., Crevecoeur, S., Lovejoy, C., and Vincent, W. F. (2016). Microbial biogeography of permafrost thaw ponds across the changing northern landscape. *Ecography* 39, 609–618. doi: 10.1111/ecog.01667
- Crevecoeur, S., Vincent, W. F., Comte, J., and Lovejoy, C. (2015). Bacterial community structure across environmental gradients in permafrost thaw ponds: methanotroph-rich ecosystems. *Front. Microbiol.* 6:192. doi: 10.3389/fmicb.2015.00192
- Crevecoeur, S., Vincent, W. F., Comte, J., Matveev, A., and Lovejoy, C. (2017). Diversity and potential activity of methanotrophs in high methane-emitting permafrost thaw ponds. *PLoS One* 12:e0188223. doi: 10.1371/journal.pone.0188223
- Crevecoeur, S., Vincent, W. F., and Lovejoy, C. (2016). Environmental selection of planktonic methanogens in permafrost thaw ponds. *Sci. Rep.* 6:31312. doi: 10.1038/srep31312
- Cruaud, P., Vigneron, A., Fradette, M. S., Charette, S. J., Rodriguez, M. J., Dorea Caetano, C., et al. (2017). Open the Sterivex™ casing: an easy and effective way to improve DNA extraction yields. *Limnol. Oceanogr. Methods* 15, 1015–1020. doi: 10.1002/lom3.10221
- de Bok, F. A. M., Plugge, C. M., and Stams, A. J. M. (2004). Interspecies electron transfer in methanogenic propionate degrading consortia. *Water Res.* 38, 1368–1375. doi: 10.1016/j.watres.2003.11.028
- Denfeld, B. A., Baulch, H. M., del Giorgio, P. A., Hampton, S. E., and Karlsson, J. (2018). A synthesis of carbon dioxide and methane dynamics during the ice-covered period of northern lakes. *Limnol. Oceanogr. Lett.* 3, 117–131. doi: 10.1002/lo.10079
- Deshpande, B. N., Crevecoeur, S., Matveev, A., and Vincent, W. F. (2016). Bacterial production in subarctic peatland lakes enriched by thawing permafrost. *Biogeosciences* 13, 4411–4427. doi: 10.5194/bg-13-4411-2016
- Deshpande, B. N., Maps, F., Matveev, A., and Vincent, W. F. (2017). Oxygen depletion in subarctic peatland thaw lakes. *Arct. Sci.* 3, 406–428. doi: 10.1371/journal.pone.0188223
- Farquharson, L. M., Mann, D. H., Grosse, G., Jones, B. M., and Romanovsky, V. E. (2016). Spatial distribution of thermokarst terrain in Arctic Alaska. *Geomorphology* 273, 116–133. doi: 10.1016/j.geomorph.2016.08.007
- Graham, D. E., Wallenstein, M. D., Vishnivetskaya, T. A., Waldrop, M. P., Phelps, T. J., Pfliffer, S. M., et al. (2011). Microbes in thawing permafrost: the unknown variable in the climate change equation. *ISME J.* 6:709. doi: 10.1038/ismej.2011.163
- Hammer, Ø, Harper, D., and Ryan, P. (2001). PAST—palaeontological statistics, ver. 1.89. *Palaeontol. Electron* 4:9.
- Hug, L. A., Castelle, C. J., Wrighton, K. C., Thomas, B. C., Sharon, I., Frischkorn, K. R., et al. (2013). Community genomic analyses constrain the distribution of metabolic traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling. *Microbiome* 1:22. doi: 10.1186/2049-2618-1-22
- Jansson, J. K., and Taş, N. (2014). The microbial ecology of permafrost. *Nat. Rev. Microbiol.* 12:414. doi: 10.1038/nrmicro3262
- Jetten, M. S. M., Stams, A. J. M., and Zehnder, A. J. B. (1992). Methanogenesis from acetate: a comparison of the acetate metabolism in *Methanotheroxobacter* and *Methanosarcina* spp. *FEMS Microbiol. Lett.* 88, 181–197. doi: 10.1016/0378-1097(92)90802-U
- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., et al. (2008). KEGG for linking genomes to life and the environment. *Nucleic Acids Res.* 36, D480–D484. doi: 10.1093/nar/gkm882
- Kang, D. D., Froula, J., Egan, R., and Wang, Z. (2015). MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ* 3:e1165. doi: 10.7717/peerj.1165
- Kertesz, M. A. (2000). Riding the sulfur cycle – metabolism of sulfonates and sulfate esters in Gram-negative bacteria. *FEMS Microbiol. Rev.* 24:135. doi: 10.1016/s0168-6445(99)00033-9
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., et al. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 41:e1. doi: 10.1093/nar/gks808
- Koudelakova, T., Bidmanova, S., Dvorak, P., Pavelka, A., Chaloupkova, R., Prokop, Z., et al. (2013). Haloalkane dehalogenases: Biotechnological applications. *Biotechnol. J.* 8, 32–45. doi: 10.1002/biot.201100486
- Laczny, C. C., Sternal, T., Plugaru, V., Gawron, P., Atashpendar, A., Margossian, H. H., et al. (2015). VizBin - an application for reference-independent visualization and human-augmented binning of metagenomic data. *Microbiome* 3:1. doi: 10.1186/s40168-014-0066-1
- Langer, M., Westermann, S., Walter Anthony, K., Wischniewski, K., and Boike, J. (2015). Frozen ponds: production and storage of methane during the Arctic winter in a lowland tundra landscape in northern Siberia, Lena River delta. *Biogeosciences* 12, 977–990. doi: 10.5194/bg-12-977-2015
- Laurion, I., Vincent, W. F., MacIntyre, S., Retamal, L., Dupont, C., Francus, P., et al. (2010). Variability in greenhouse gas emissions from permafrost thaw ponds. *Limnol. Oceanogr.* 55, 115–133. doi: 10.4319/lo.2010.55.1.0115
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. (2015). MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31, 1674–1676. doi: 10.1093/bioinformatics/btv033
- Liang, B., Wang, L.-Y., Mbadinga, S. M., Liu, J.-F., Yang, S.-Z., Gu, J.-D., et al. (2015). *Anaerolineaceae* and *Methanoseta* turned to be the dominant microorganisms in alkanes-dependent methanogenic culture after long-term of incubation. *AMB Express* 5:37. doi: 10.1186/s13568-015-0117-4
- Mackelprang, R., Burkert, A., Haw, M., Mahendrarajah, T., Conaway, C. H., Douglas, T. A., et al. (2017). Microbial survival strategies in ancient permafrost: insights from metagenomics. *ISME J.* 11, 2305–2318. doi: 10.1038/ismej.2017.93
- Magoč, T., and Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27, 2957–2963. doi: 10.1093/bioinformatics/btr507
- Markowitz, V. M., Mavromatis, K., Ivanova, N. N., Chen, I.-M. A., Chu, K., and Kyrpides, N. C. (2009). IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25, 2271–2278. doi: 10.1093/bioinformatics/btp393
- Matveev, A. (2018). *Dynamique du méthane et du dioxyde de carbone dans les mares de thermokarst du Québec subarctique*. PhD Thesis, Université Laval, Canada.
- Matveev, A., Laurion, I., Deshpande, B. N., Bhiry, N., and Vincent, W. F. (2016). High methane emissions from thermokarst lakes in subarctic peatlands. *Limnol. Oceanogr.* 61, S150–S164. doi: 10.1002/lno.10311
- Matveev, A., Laurion, I., Deshpande, B. N., and Vincent, W. F. (2019). *Concentrations of Dissolved Methane, Carbon Dioxide and Oxygen in Thermokarst Lakes and Ponds in Palsa Peatlands, Northern Quebec, Canada*. Quebec, QC: Nordicana D. doi: 10.5885/45588XX-5A12A84DFAAF4D36
- Meckenstock, R. U., Boll, M., Mouttaki, H., Koelschbach, J. S., Cunha Tarouco, P., Weyrauch, P., et al. (2016). Anaerobic degradation of benzene and polycyclic aromatic hydrocarbons. *J. Mol. Microbiol. Biotechnol.* 26, 92–118. doi: 10.1159/000441358
- Neuenschwander, S. M., Ghai, R., Pernthaler, J., and Salcher, M. M. (2017). Microdiversification in genome-streamlined ubiquitous freshwater Actinobacteria. *ISME J.* 12:185. doi: 10.1038/ismej.2017.156
- Newton, R. J., Jones, S. E., Helmus, M. R., and McMahon, K. D. (2007). Phylogenetic ecology of the freshwater Actinobacteria acI lineage. *Appl. Environ. Microbiol.* 73, 7169–7176. doi: 10.1128/aem.00794-07
- Newton, R. J., Kent, A. D., Triplett, E. W., and McMahon, K. D. (2006). Microbial community dynamics in a humic lake: differential persistence of common freshwater phylotypes. *Environ. Microbiol.* 8, 956–970. doi: 10.1111/j.1462-2920.2005.00979.x
- Oliva, M., and Fritz, M. (2018). Permafrost degradation on a warmer earth: challenges and perspectives. *Sustain. Soil Manag. Land Restor.* 5, 14–18. doi: 10.1016/j.coesh.2018.03.007
- Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., and Tyson, G. W. (2015). CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 25, 1043–1055. doi: 10.1101/gr.186072.114
- Parmentier, F.-J. W., Christensen, T. R., Rysgaard, S., Bendtsen, J., Glud, R. N., Else, B., et al. (2017). A synthesis of the arctic terrestrial and marine carbon cycles under pressure from a dwindling cryosphere. *Ambio* 46, 53–69. doi: 10.1007/s13280-016-0872-8

- Payette, S., Delwaide, A., Caccianiga, M., and Beauchemin, M. (2004). Accelerated thawing of subarctic peatland permafrost over the last 50 years. *Geophys. Res. Lett.* 31:L18208. doi: 10.1029/2004GL020358
- Peng, Y., Leung, H. C. M., Yiu, S. M., and Chin, F. Y. L. (2012). IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28, 1420–1428. doi: 10.1093/bioinformatics/bts174
- Pokrovsky, O. S., Shirokova, L. S., Kirpotin, S. N., Audry, S., Viers, J., and Dupré, B. (2011). Effect of permafrost thawing on organic carbon and trace element colloidal speciation in the thermokarst lakes of western Siberia. *Biogeosciences* 8, 565–583. doi: 10.5194/bg-8-565-2011
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., et al. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* 35, 7188–7196. doi: 10.1093/nar/gkm864
- Ricão Canelhas, M., Denfeld, B. A., Weyhenmeyer, G. A., Bastviken, D., and Bertilsson, S. (2016). Methane oxidation at the water-ice interface of an ice-covered lake. *Limnol. Oceanogr.* 61, S78–S90. doi: 10.1002/lno.10288
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541. doi: 10.1128/AEM.01541-09
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., et al. (2015). Climate change and the permafrost carbon feedback. *Nature* 520:171. doi: 10.1038/nature14338
- Teske, A., and Sorensen, K. B. (2007). Uncultured archaea in deep marine subsurface sediments: have we caught them all? *ISME J.* 2, 3–18. doi: 10.1038/ismej.2007.90
- Tran, P., Ramachandran, A., Khawasek, O., Beisner, B. E., Rautio, M., Huot, Y., et al. (2018). Microbial life under ice: metagenome diversity and *in situ* activity of Verrucomicrobia in seasonally ice-covered lakes: Sub-ice Verrucomicrobia genomes in Quebec lakes. *Environ. Microbiol.* 20, 2568–2584. doi: 10.1111/1462-2920.14283
- Tveit, A. T., Urich, T., Frenzel, P., and Svenning, M. M. (2015). Metabolic and trophic interactions modulate methane production by Arctic peat microbiota in response to warming. *Proc. Natl. Acad. Sci. U.S.A.* 112:E2507. doi: 10.1073/pnas.1420797112
- Vigneron, A., Alsop, E. B., Lomans, B. P., Kyrpides, N. C., Head, I. M., and Tsesmetzis, N. (2017). Succession in the petroleum reservoir microbiome through an oil field production lifecycle. *ISME J.* 11:2141. doi: 10.1038/ismej.2017.78
- Vigneron, A., Cruaud, P., Pignet, P., Caprais, J.-C., Cambon-Bonavita, M.-A., Godfroy, A., et al. (2013). Archaeal and anaerobic methane oxidizer communities in the Sonora Margin cold seeps. Guaymas Basin (Gulf of California). *ISME J.* 7, 1595–1608. doi: 10.1038/ismej.2013.18
- Vincent, W. F., Lemay, M., and Allard, M. (2017). Arctic permafrost landscapes in transition: towards an integrated Earth system approach. *Arct. Sci.* 3, 39–64. doi: 10.1139/as-2016-0027
- Wagner, M., and Horn, M. (2006). The Planctomycetes, Verrucomicrobia, Chlamydiae and sister phyla comprise a superphylum with biotechnological and medical relevance. *Environ. Biotechnol. Biotechnol.* 17, 241–249. doi: 10.1016/j.copbio.2006.05.005
- Walter Anthony, K., Schneider von Deimling, T., Nitze, I., Frolking, S., Emond, A., Daanen, R., et al. (2018). 21st-century modeled permafrost carbon emissions accelerated by abrupt thaw beneath lakes. *Nat. Commun.* 9:3262. doi: 10.1038/s41467-018-05738-9
- Wauthy, M., Rautio, M., Christoffersen, K. S., Forsström, L., Laurion, I., Mariash, H. L., et al. (2018). Increasing dominance of terrigenous organic matter in circumpolar freshwaters due to permafrost thaw. *Limnol. Oceanogr. Lett.* 3, 186–198. doi: 10.1002/lo2.10063
- Woodcroft, B. J., Singleton, C. M., Boyd, J. A., Evans, P. N., Emerson, J. B., Zayed, A. A. F., et al. (2018). Genome-centric view of carbon processing in thawing permafrost. *Nature* 560, 49–54. doi: 10.1038/s41586-018-0338-1
- Wu, Y.-W., Simmons, B. A., and Singer, S. W. (2016). MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32, 605–607. doi: 10.1093/bioinformatics/btv638
- Wurzbacher, C., Nilsson, R. H., Rautio, M., and Peura, S. (2017). Poorly known microbial taxa dominate the microbiome of permafrost thaw ponds. *ISME J.* 11:1938. doi: 10.38/ismej.2017.54
- Yamada, T., Sekiguchi, Y., Imachi, H., Kamagata, Y., Ohashi, A., and Harada, H. (2005). Diversity, localization, and physiological properties of filamentous microbes belonging to Chloroflexi Subphylum I in mesophilic and thermophilic methanogenic sludge granules. *Appl. Environ. Microbiol.* 71, 7493–7503. doi: 10.1128/AEM.71.11.7493-7503.2005
- Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., and Xu, Y. (2012). dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res.* 40, W445–W451. doi: 10.1093/nar/gks479
- Yuan, C., Lei, J., Cole, J., and Sun, Y. (2015). Reconstructing 16S rRNA genes in metagenomic data. *Bioinformatics* 31, i35–i43. doi: 10.1093/bioinformatics/btv231

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Vigneron, Lovejoy, Cruaud, Kalenitchenko, Culley and Vincent. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.