



The Independent Effects of Psychosocial Stressors on Subclinical Psychosis: Findings From the Multinational EU-GEI Study

Baptiste Pignon, Mohamed Lajnef, James B Kirkbride, Hugo Peyre, Aziz Ferchiou, Jean-Romain Richard, Grégoire Baudin, Sarah Tosato, Hannah Jongsma, Lieuwe de Haan, et al.

► To cite this version:

Baptiste Pignon, Mohamed Lajnef, James B Kirkbride, Hugo Peyre, Aziz Ferchiou, et al.. The Independent Effects of Psychosocial Stressors on Subclinical Psychosis: Findings From the Multinational EU-GEI Study. *Schizophrenia Bulletin*, 2021, 10.1093/schbul/sbab060 . hal-03337986

HAL Id: hal-03337986

<https://hal.science/hal-03337986v1>

Submitted on 14 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Draft Manuscript for Review. Please review online at <http://mc.manuscriptcentral.com/oup/szbltn>

**The independent effects of psychosocial stressors on
subclinical psychosis: findings from the multinational EU-
GEI study**

Journal:	<i>Schizophrenia Bulletin</i>
Manuscript ID	SZBLTN-ART-20-0835.R2
Manuscript Type:	Regular Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Pignon, Baptiste; Albert Chenevier Hospital, ; Lajnef, Mohamed Kirkbride, James; University College London, Division of Psychiatry; Peyre, Hugo Ferchiou, Aziz Richard, Jean-Romain; Univ Paris Est Créteil, INSERM, IMRB, Translational Neuropsychiatry Baudin, Grégoire Tosato, Sarah; University of Verona, Department of Neurosciences, Biomedicine and Movement Sciences, Section of Psychiatry and Clinical Psychology Jongsma, Hannah de Haan, Lieuwe; Amsterdam UMC -AMC Campus, Psychiatry Tarricone, Ilaria; University of Bologna, Department of Medical and Surgical Science, Psychiatry Unit, Alma Mater Studiorum Bernardo, Miquel; Hospital Clinic of Barcelona, Psychiatry Velthorst, Eva; Icahn School of Medicine at Mount Sinai, Department of Psychiatry, Early Psychosis; Icahn School of Medicine at Mount Sinai, departments of psychiatry and preventive medicine Braca, Mauro Arango, Celso; Child and Adolescent Psychiatry Department, Instituto de Investigación Sanitaria Gregorio Marañón, IISGM. Hospital General Universitario Gregorio Marañón, CIBERSAM Arrojo, Manuel; Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS) Servizo Galego de Saúde (SERGAS), Hospital Clínico Universitario; Complejo Hospitalario Universitario de Santiago de Compostela, Servizo de Psiquiatría Bobes, Julio; Universidad de Oviedo, Facultad de Medicina Del-Ben, Cristina; Universidade de Sao Paulo Campus de Ribeirao Preto, Neuroscience and Behavior, Division of Psychiatry Di Forti, Marta; Institute of Psychiatry, Psychosis Studies Gayer-Anderson, Charlotte; King's College London, Health Services and Population Research Jones, Peter; University of Cambridge, Psychiatry La Cascia, caterina; Palermo University, Dept of Experimental Biomedicine and Clinical Neuroscience -section of Psychiatry Lasalvia, Antonio; University of Verona, Public Health and Community</p>

	Medicine, Section of Psychiatry Menezes, Paulo; Universidade de Sao Paulo Faculdade de Medicina, Preventive Medicine Quattrone, Diego; Institute of Psychiatry Psychology and Neuroscience, Sanjuan, Julio; Faculty of Medicine, Psychiatry; Selten, Jean-Paul ; Rivierduinen Psychiatric Institute; Maastricht University MHeNS School for Mental Health and Neuroscience, Dept. of Psychiatry & Neuropsychology Tortelli, Andrea ; Etablissement Public de Sante Maison Blanche Ilorca, Pierre; CHU Clermont-Ferrand, CMP B; Universite Clermont Auvergne, van Os, Jim; Maastricht University, Rutten, Bart; Maastricht University, School for Mental Health and Neuroscience Murray, Robin; King's College London, psychosis Morgan, Craig; Institute of Psychiatry, Social Psychiatry Leboyer, Marion; AP-HP, Departmen of Psychiatry ; INSERM, IMRB, Dept of neuroscience szöke, andrei; Assistance Publique - Hopitaux de Paris, Hôpitaux "H. Mondor", Pole Psychiatrie Schurhoff, Franck; INSERM U955, AP-HP
Keywords:	Subclinical psychosis, Schizotypy, Psychotic symptoms, Psychosocial stress, Childhood trauma, Stressful life events, Social capital, Discrimination, Environment*environment interaction

SCHOLARONE™
Manuscripts

The independent effects of psychosocial stressors on subclinical psychosis: findings from the multinational EU-GEI study

Baptiste Pignon^{1,+}, Mohamed Lajnef¹, James B. Kirkbride², Hugo Peyre³, Aziz Ferchiou¹, Jean-Romain Richard¹, Grégoire Baudin⁴, Sarah Tosato⁵, Hannah Jongsma⁶, Lieuwe de Haan⁷, Ilaria Tarricone⁸, Miguel Bernardo⁹, Eva Velthorst¹⁰, Mauro Braca¹¹, Celso Arango¹², Manuel Arrojo¹³, Julio Bobes¹⁴, Cristina Marta Del-Ben¹⁵, Marta Di Forti¹⁶, Charlotte Gayer-Anderson¹⁷, Peter B. Jones¹⁸, Caterina La Cascia¹⁹, Antonio Lasalvia²⁰, Paulo Rossi Menezes²¹, Diego Quattrone²², Julio Sanjuán²³, Jean-Paul Selten²⁴, Andrea Tortelli²⁵, Pierre-Michel Llorca²⁶, Jim van Os²⁷, Bart P. F. Rutten²⁸, Robin M. Murray²⁹, Craig Morgan¹⁷, Marion Leboyer¹, Andrei Szöke¹, Franck Schürhoff¹

¹Univ Paris Est Creteil, INSERM, IMRB, AP-HP, Hôpitaux Universitaires « H. Mondor », DMU IMPACT, Fondation FondaMental, F-94010 Créteil, France.

²PsyLife group, Division of Psychiatry, UCL, London, W1T 7NF, UK.

³AP-HP, Hôpital universitaire Robert Debré, Service de pédopsychiatrie, 75019 Paris, France.

⁴Université de Paris, Laboratoire de Psychopathologie et Processus de Santé, F-92100 Boulogne Billancourt, France.

⁵Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement Sciences

, Verona, Italy.

⁶Centre for Transcultural Psychiatry 'Veldzicht'. Balkbrug, the Netherlands; VR Mental Health Group, University Center for Psychiatry, University Medical Centre Groningen, Groningen, the Netherlands; Centre for Longitudinal Studies, UCL, London, UK.

⁷Amsterdam UMC, Amsterdam, The Netherlands; Arkin, Amsterdam, The Netherlands

⁸Department of Medical and Surgical Sciences, Bologna University.

⁹Barcelona Clínic Schizophrenia Unit, Hospital Clínic of Barcelona, Institute of Neuroscience, University of Barcelona; Institut d'investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona; Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Spain.

¹⁰Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, US.

¹¹Department of Mental Health and Pathological Addictions, Local Health Authority, Bologna, 40100, Italy.

¹²Department of Child and Adolescent Psychiatry, Institute of Psychiatry and Mental Health, Hospital General Universitario Gregorio Marañón, Gregorio Marañón, (IiGSM), School of Medicine, Universidad Complutense de Madrid; CIBERSAM, Madrid, Spain.

¹³Department of Psychiatry, Psychiatric Genetic Group, Instituto de Investigación Sanitaria de Santiago de Compostela, Complejo Hospitalario Universitario de Santiago de Compostela, Spain.

¹⁴Faculty of Medicine and Health Sciences - Psychiatry, Universidad de Oviedo; ISPA, INEUROPA CIBERSAM. Oviedo, Spain.

1

2

338¹⁵Department of Neuroscience and Behaviour, Ribeirão Preto Medical School, University of São Paulo, São Paulo, Brazil.

439¹⁶Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King’s

540College London, London SE5 8AF, UK; South London and Maudsley NHS Mental Health Foundation Trust, London, UK

641¹⁷Department of Health Service and Population Research, Institute of Psychiatry, King's College London, De Crespigny Park,

742Denmark Hill, SE5 8AF, London, UK.

8

943¹⁸Department of Psychiatry, University of Cambridge, CB2 0SZ; CAMEO, Cambridgeshire & Peterborough NHS

1044Foundation Trust, UK.

11

1245¹⁹Dept. Biomedicine, Neuroscience and Advanced Diagnostics. School of Medicine, University of Palermo,Palermo, Italy.

1346²⁰Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement Sciences; University of Verona; Verona;

1447Italy.

15

1648²¹Department of Preventive Medicine, Faculdade de Medicina, Universidade of São Paulo, São Paulo, Brazil.

1749²²Social, Genetic, and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King’s

1850College London.

19

2051²³Biomedical Research Networking Centre in Mental Health (CIBERSAM), Madrid, Spain; Department of Psychiatry,

2152Hospital Clínico Universitario de Valencia, School of Medicine, Universidad de Valencia, Valencia, Spain.

2253Biomedical Research Institute INCLIVA, Valencia, Spain.

23

2454²⁴Rivierduinen Institute for mental Health, Leiden, The Netherlands; Maastricht University Medical Center, Dept. of

2555Psychiatry & Neuropsychology, School for Mental Health and Neuroscience, Maastricht, The Netherlands.

2656²⁵French National Institute of Health and Medical Research (INSERM), U955 - 15, Créteil, France and EPS Maison Blanche,

2757Paris, France.

28

2958²⁶CHU Clermont-Ferrand, Psychiatrie B, Clermont-Ferrand, F-63003, France ; Université Clermont Auvergne, EA7280, F-

305963000 Clermont-Ferrand, France.

31

3260²⁷Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University

3361Medical Centre, Maastricht, The Netherlands; Department of Psychiatry, UMC Utrecht Brain Centre, University Medical

3462Centre Utrecht, Utrecht University, Utrecht, The Netherlands; Department of Psychosis Studies, Institute of Psychiatry,

3563Psychology & Neuroscience, King's College London, London, UK.

36

3764²⁸Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University

3865Medical Centre, Maastricht, The Netherlands.

3966²⁹The Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK.

4067

41

4268⁺Dr. Baptiste Pignon

43

4469Hôpital Albert Chenevier, Groupe hospitalier Henri-Mondor, CHU de Créteil, Assistance

4570Publique-Hôpitaux de Paris (AP-HP), 40 rue de Mesly, 94 000, Créteil, France

46

4771**baptistepignon@yahoo.fr**

48

4972☎ : + 33 1 49 81 31 31 ; **Fax** : +33 1 49 81 30 59

50

51

52

53

54

55

56

57

58

59

60

ABSTRACT

The influence of psychosocial stressors on psychosis risk has usually been studied in isolation and after the onset of the disorder, potentially ignoring important confounding relationships or the fact that some stressors that may be the consequence of the disorder rather than preexisting. The study of subclinical psychosis could help to address some of these issues. In this study, we investigated whether there was (i) an association between dimensions of subclinical psychosis and several psychosocial stressors including: childhood trauma, self-reported discrimination experiences, low social capital, and stressful life experiences, and (ii) any evidence of environment-environment (ExE) interactions between these factors.

Data were drawn from the EUGEI study, in which healthy controls (N=1497) and siblings of subjects with a psychotic disorder (N=265) were included in six countries. The association between psychosocial stressors and subclinical psychosis dimensions (positive, negative and depressive dimension as measured by the Community Assessment of Psychic Experiences (CAPE) scale) and possible ExE interactions were assessed using linear regression models.

After adjusting for sex, age, ethnicity, country, and control/sibling status, childhood trauma (β for positive dimension: 0.13, negative: 0.49, depressive: 0.26) and stressful life events (positive: 0.08, negative: 0.16, depressive: 0.17) were associated with the three dimensions. Lower social capital was associated with the negative and depression dimensions (negative: 0.26, depressive: 0.13), and self-reported discrimination experiences with the positive dimension (0.06).

Our findings are in favor of independent, cumulative and non-specific influences of social adversities in subclinical psychosis in non-clinical populations, without arguments for ExE interactions.

Keywords

Subclinical psychosis, Schizotypy, Psychotic symptoms, Positive subclinical symptoms, Negative subclinical symptoms, Depressive subclinical symptoms, Psychosocial stress, Childhood trauma, Stressful life events, Social capital, Discrimination, Community Assessment of Psychic Experiences (CAPE)

104 **INTRODUCTION**

105 The stress-vulnerability theory of psychotic disorders posits that genetic factors interact
106 with environmental stressors in the development of a disorder¹⁻³. In this theoretical model,
107 increased sensitivity to stress plays an important role in both onset and relapse of psychotic
108 disorders, including schizophrenia. Extending this model, the stress sensitization hypothesis
109 proposes that repeated exposure to environmental stressors sensitizes key neurobiological
110 pathways to psychosis^{4,5}. Early, severe or prolonged exposure to stress would lead to a
111 dysregulated stress response and potentially explain both the role of early and current stress in
112 the etiology of psychotic disorders.

113 Several environmental factors that could be conceptualized as psychosocial stressors,
114 such as childhood trauma, stressful life events, discrimination experiences or a low level of
115 social capital have been found to increase the risk of psychotic disorders⁶⁻⁹. Furthermore,
116 several studies support a stress sensitization model in psychosis, showing for example that
117 exposure to an early stressor (childhood trauma) increases later sensitivity to other stressors
118 (e.g., social stress, population density, etc.) in patients (environment-environment ExE
119 interaction)^{10,11}. However, the fact that these studies have taken place after the onset of
120 psychotic disorder raises causality concerns. Indeed, the assessment of recent stress could be
121 confounded by several stressors associated with the disease itself, such as hospitalizations,
122 stigma, substance use disorders or social drift after onset¹²⁻¹⁴.

123 Psychotic symptoms may be present, to varying degrees, in non-clinical, general
124 population samples who do not meet criteria for a clinical disorder¹⁵⁻¹⁷. The continuum theory
125 of psychosis posits that subclinical experiences have a similar origin/etiology to full psychotic
126 disorders¹⁸⁻²². Thus, studying risk factors associated with subclinical psychosis may provide
127 insights into the etiology of psychosis, while reducing the potential interference of reverse
128 causation, i.e., stress caused by the clinical disorders themselves. Furthermore, in line with the
129 continuum theory, subclinical psychosis can be characterized by quantitative (continuous)
130 variables, improving statistical power and the capacity to control for more potential
131 confounders.

132 Several studies have previously reported associations between psychosocial stressors
133 and subclinical psychosis. For instance, childhood trauma has been associated with increased
134 rates of isolated psychotic symptoms in several studies²³⁻²⁶. Likewise, negative and/or stressful
135 life events^{27,28} or self-reported discrimination experiences²⁹ have also been associated with
136 subclinical psychosis.

One major limitation of studies published to date is that psychosocial stressors have usually been studied in isolation, which might lead to spurious or incomplete conclusions given the likely presence of confounding/interaction with other stressors. Further work is needed to determine whether different psychosocial stressors have independent effects on subclinical psychosis and underlying dimensions (i.e., positive, negative, depressive), while controlling for other relevant stressors. Moreover, the role of population-level (e.g., social capital) factors that might interact with psychosocial stressors has rarely been explored. For example, social capital has been related to the incidence and outcome of psychotic disorders^{30,31} and low levels of social capital could be a stressful condition *per se*³² (i.e., an independent stressor), or exacerbate the effects of other psychosocial factors (discrimination, trauma, stressful life events) on subsequent psychosis risk, but evidence is lacking. Relationships between social capital and subclinical psychosis have never been investigated. Moreover, to the best of our knowledge, only two studies have analyzed the role of multiple psychosocial stressors on subclinical psychosis. The first found arguments in favor of the sensitization hypothesis³³. The second, studying the role of childhood trauma alongside other environmental and genetic risk factors, found additive effects of these factors on subclinical psychosis scores²¹.

Furthermore, recent studies have reported that exposure to psychosocial stressors was in part dependent on genetic vulnerability to psychosis^{34,35}, and no study has compared associations between psychosocial stressors and subclinical psychosis in both control participants and (healthy) siblings of people with a psychotic disorder. Including both groups may help tease out the genetic and environmental etiology of psychosis.

To address some of the limitations of previous work, we used a large, cross-national sample of population-based controls and siblings of subjects with a psychotic disorder, to investigate whether there was (i) an association between subclinical psychosis dimensions (positive, negative and depressive) and psychosocial stressors, and (ii) any evidence of ExE interactions between different psychosocial stressors in line with the stress sensitization hypothesis. The psychosocial stressors we explored can be conceptualized as either “early” (childhood trauma) or “prolonged” (discrimination, low level of social capital, stressful life experiences). We hypothesized that exposure to “early” stressors would enhance the effect of adversity later in life.

METHODS

EU-GEI study

Data were collected in the “European network of national schizophrenia networks studying gene-environment interactions” (EU-GEI) study, a multicentre case-sibling-control study of genetic and environmental determinants of the occurrence, severity and outcome of psychotic disorders. For the second work-package of the study (WP2: Functional Enviromics), three categories of participants were recruited between 2010 and 2015: (i) subjects presenting with a first-episode of psychotic disorder (FEP), (ii) population-based healthy controls, and; (iii) siblings of participants with FEP³⁶. Participants were recruited across 6 countries: Brazil, France, Italy, the Netherlands, Spain, and the United Kingdom^{37,38}. In the present analyses, only controls and siblings were included.

Subjects

Population-based controls and siblings had no personal history of psychosis, and controls were recruited from the same catchment areas as the cases. In each centre, controls were recruited using a mixture of random and quota sampling to ensure control participants were broadly representative of the population at-risk from which cases could present in each catchment area on predefined variables (age, sex, and migration)³⁶.

Subclinical psychosis measure

The Community Assessment of Psychic Experiences (CAPE) is a 42-item, self-report questionnaire that has been developed to measure lifetime subclinical psychotic dimensions in the general population³⁹. For each item, 4 answers were possible according to the frequency of their occurrences (from never to nearly always). A meta-analysis found that the CAPE displays a three-dimensional symptom structure: positive, negative and depressive dimensions⁴⁰. We therefore used the sum of endorsed items to quantify each of the three psychotic dimensions consistent with previous studies^{22,41,42}. To construct these dimension scores, we dichotomized answers on each CAPE item (never vs. sometimes or more) and summed the positive answers. This led to theoretical positive dimension scores between 0-20, negative dimension scores between 0-14, and depressive dimension scores between 0-8.

We have previously demonstrated the cross-national invariance of the CAPE assessment in the EUGEI WP2 samples: equivalent factorial structures, factor loadings and thresholds

across the six countries⁴³. Thus, CAPE results can be reliably used across the different EUGEI countries.

Psychosocial stressors measure

Childhood trauma was assessed with the Childhood Trauma Questionnaire (CTQ), a 25-item questionnaire, that measures five different domains (emotional and physical neglect; emotional, physical and sexual abuse). All items are scored on a five-point Likert-scale (1: never, to 5: very often). We used the CTQ total score as the sum of all items, thus theoretically ranging from 25-125⁴⁴.

Lifetime self-reported discrimination experiences were assessed with a modified version of the Williams' major experiences of discrimination measure (available in **Supplementary Material 1**), a 12-item scale assessing several experiences of discrimination (unfairly fired or not hired because of your ethnicity/sex/weight/etc., unfairly stopped/questioned/physically threatened or abused by the police, etc.)^{45,46}. This version of the Williams' scale has already been used in a paper studying the prevalence of discrimination in South London, and its relationships with psychiatric disorders⁴⁷. We used a total discrimination score by adding all endorsed items, ranging theoretically from 0-12.

Perceived social capital in each participant's immediate neighborhood was assessed using the Social Environment Assessment Tool (SEAT), a 23-item questionnaire. This tool was designed to capture four dimensions of social capital: civic disorder (CD), impact of civic disorder (ICD), informal social control (ISC), and social cohesion and trust (SCT)⁴⁸⁻⁵¹. Respondents answer according to a five-point Likert-scale (1: unusual, to 5: very common). Sum scores for 4 subscales were derived then standardized to Z-scores (i.e., to a mean equal to 0, and a standard-deviation equal to 1). The total social capital score was obtained by adding the weighted scale scores ($SEAT\ score = zCD + 0.51 * zICD + 1.6 * zISC + zSCT$) based on the factorial structure of the instrument. This scale has shown excellent goodness-of-fit statistics (data available on request forthcoming). Our analyses were restricted to the total social capital score, which we inverted for analyses so that higher scores were associated with *lower* social capital.

Finally, stressful life events were assessed using the List of Threatening Experiences (LTE) which comprises 20 binary items^{52,53}. This scale assesses 20 events usually associated with major stress over the course of the previous 6 months including: serious injury or illness in oneself or a close relative, death of parent/child/partner, death of a family member, death of

a friend, separation from a partner, loss of job or financial difficulties. The total score ranges theoretically from 0-20.

Other adjustment variables

We also collected information on age, sex, country, and ethnicity as possible confounding factors. Age was measured at the time of the interview. Ethnicity was self-defined according to one of 6 categories: Asian, Black, North African, White, Mixed, Other.

Ethical procedures

Ethical approval was obtained from local research ethics committees in each country. The EU-GEI Project was funded by the European Community’s Seventh Framework Program under grant agreement no. HEALTH-F2-2010-241909.

Statistical methods

First, we assessed the associations between psychosocial stressors using Spearman correlation tests. Then we analyzed the relationships between these stressors and the three CAPE dimensions scores also using Spearman correlation tests. We used Mann-Whitney U-tests to assess the differences in exposure to psychosocial stressors between controls and siblings.

Second, we fitted linear regression models to analyze independent and specific effects of each psychosocial stressor on each of our three subclinical psychosis outcomes (i.e., each of the three CAPE dimensions). We fitted multivariable models, controlling for other relevant psychosocial stressors as well as age, sex, ethnicity, country, and control-sibling status (i.e., *a priori* confounders)^{18,54,55}. As CAPE scores did not follow a normal distribution (as shown by Shapiro tests with a p-value < 0.05, and graphical methods), a Box-Cox transformation of all continuous variables (CAPE scores) was performed to fulfil the normality assumption required by the parametric procedure. Complete case analysis was used.

Third, we tested for evidence of ExE interactions between our four psychosocial stressors on each outcome in our multivariable models. Each interaction (i.e., CTQxLTE, CTQxSEAT, LTExSEAT, etc.) was tested separately, by introducing interaction terms in the 4 multivariable models. In a final model, all the interaction terms were pooled into a single model for each of the outcomes.

265 To facilitate comparisons of effect sizes of the different psychosocial stressor measures,
266 Z-scores of these variables were calculated and used in the multivariable models.

267 The analyses were repeated among sibling and control subsamples, as siblings may
268 experience different exposure and different response to the same exposure.

269 Significance was based on a two-sided p-value of 0.05 throughout. R software version
270 3.6.0, with “stats”, “car” and “lattice” packages, was used.

1
2
3
4 271 **RESULTS**

5
6 272
7
8 273 ***Sample characteristics***

9
10 274 The sample was composed of 1,762 subjects, including 1,497 controls (85.0%) and 265
11 275 siblings (15.0%), of those 972 were men (55.2%) and 790 were women (44.8%). The median
12 276 age for the controls was 33 years (IQR [26-47]), and 30 years (IQR [23-38]) for the siblings.
13 277 The proportion of women was higher among controls, and controls were older. The proportion
14 278 of people from non-white ethnic backgrounds differed between controls (21.3%) and siblings
15 279 (24.1%), with a higher proportion of subjects of Black ethnicity (8.1% vs. 4.9%) and a lower
16 280 proportion of subjects of Mixed ethnicity in controls (7.7% vs. 15.5%). Regarding CAPE
17 281 scores, positive dimensions were higher among controls in comparison with siblings. Scores on
18 282 the CTQ were higher among siblings while siblings reported fewer discrimination experiences.
19 283 More details are available in the **Table 1**.
20
21
22
23
24
25
26
27 284

28
29 285 **- TABLE 1 -**
30
31 286

32
33 287 ***Correlations between psychosocial stressors***

34 288 The correlation matrix (**Table 2**) revealed small but statistically robust ($p<0.001$)
35 289 correlations between all stressors. We observed positive correlations between childhood
36 290 trauma, self-reported discrimination experiences, and stressful life events (ρ between 0.14
37 291 and 0.20), and negative correlations between high level of social capital and the 3 psychosocial
38 292 stressors (ρ between -0.15 and -0.08). Correlations between psychosocial stressors showed
39 293 similar patterns of magnitude and direction within both the control and sibling groups (except
40 294 for social capital and stressful life events and social capital among siblings which were not
41 295 associated, see **Supplementary Table 2** for analyses within controls and siblings).
42
43
44
45
46
47
48
49 296

50
51 297 **- TABLE 2 -**
52
53 298

54 299 ***Correlations between subclinical psychosis scores and psychosocial***
55 300 ***stressors***

56 301 All psychosocial stressors were positively correlated with the different CAPE
57 302 dimensions (**Table 3**). Higher perceived social capital scores were associated, as expected, with

lower positive, negative and depressive dimension scores. Correlations ranged from -0.13 (higher social capital associated with lower depressive score) to 0.29 (childhood trauma and negative dimension). Correlations showed similar patterns of direction within both the control and sibling groups, with higher levels of correlation between CAPE scores and self-reported discrimination experiences among siblings vs. controls (see **Supplementary Table 3** for analyses within controls and siblings).

- TABLE 3 -

Multivariable modelling

After adjusting for sex, age, ethnicity, country, and control/sibling status, the different CAPE scores were independently associated with childhood trauma (β with Z-score of childhood trauma between 0.13 and 0.49) and stressful life events (β between 0.08 and 0.17). Lower social capital was associated with negative and depressive dimensions (β between 0.05 and 0.24), while self-reported discrimination experiences were only associated with the positive dimension ($\beta=0.06$). None of the interaction terms of the psychosocial stressor measures (both when introduced one by one in the multivariable analyses, and in the models with all the interactions terms) were associated with any of the 3 dimensions (see **Supplementary Table 4** for the results of the interactions). Of note, sibling status was associated with lower scores on the positive dimension ($\beta=-0.16$). The detailed results of the multivariable analyses with Z-scores of the psychosocial measures are available in the **Table 4**.

The same multivariable analyses were repeated separately for siblings and controls, and revealed globally similar results. Of note, among siblings, fewer associations reached statistical significance, and the interaction between childhood trauma and discrimination was significant (negative interaction: $\beta=-0.15$, more details of the multivariable analyses within controls and siblings in **Supplementary Table 5**).

- TABLE 4 -

DISCUSSION

In the cross-national and non-clinical EUGEI sample we assessed the effects of several psychosocial stressors (childhood trauma, stressful life-events, self-reported discrimination experiences and low social capital) on different subclinical psychosis dimensions for the first time. Subclinical psychosis was assessed with the CAPE and all analyses were adjusted for relevant sociodemographic factors (age, sex, country and ethnicity). This revealed that childhood trauma and stressful life events were associated with higher scores on the positive, negative and depressive dimensions. Lower social capital was associated with higher scores on negative and depressive dimensions, while self-reported discrimination experiences were associated with the positive dimension.

Overall, as all evaluated psychosocial stressors were associated with subclinical psychosis, and as no interaction between these stressors was significant, these findings are consistent with an independent effect of the different psychosocial stressors. Moreover, as the different psychosocial stressors were (with the exception of discrimination) similarly associated with the different dimensions, our findings are consistent with a common etiology for the 3 dimensions.

This study contributes to our understanding of the relationships between psychosis and early, recent, and prolonged psychosocial stressors. The major strength of this study is the concomitant analysis of several psychosocial stressors. In concordance with a recent study in the American general population⁵⁶, we confirm the existence of significant correlations between different stressors and we show that these stressors have independent effects on the 3 dimensions of subclinical psychosis. In addition, we found that a low level of social capital is associated with higher levels of negative and depressive dimensions. This ecological neighborhood-level factor had never been studied in relation with subclinical psychosis outcomes in adults before (of note, Solmi et al. found an association between maternal neighborhood stress and the rate of psychotic symptoms among 13 year old adolescents³²), while previous studies found associations with the incidence of psychotic disorders³⁰. This result is consistent with other studies regarding the influence of neighborhood characteristics such as deprivation or social fragmentation on psychosis outcomes⁶, including subclinical psychosis^{57,58}.

Surprisingly, the absence of any strong evidence of interactions between the psychosocial stressors and especially between early and recent stress was not consistent with our hypothesis and the sensitization hypothesis. This result also differs from the results of

Lataster et al.³³. In this 10-year follow-up of adolescents and young adults from the general population in Germany (N=1,722), authors found non-additive interactions between early and recent psychosocial risk factors on the risk for positive psychotic symptoms (i.e., delusional symptoms and/or hallucinations): unadjusted analyses showed associations between early and recent adversity and further psychotic symptoms, but these associations disappeared after statistical adjustment. Additive interactions between early and recent adversity were significant only for the fourth and highest level of recent adversity (adjusted risk ratio for the combined early and higher level of recent adversity = 4.08, 95% CI [2.02-8.24], p-value=0.032). Differences in results of this and our study might be explained by important methodological differences. For example, Lataster et al. study was longitudinal (three follow-up surveys covering a mean period of 8.4 years) and analyzed only two psychosocial stressors categorical variables (childhood trauma, dichotomized; and recent trauma and negative life events, four levels).

Analyzing the influence of environmental (including childhood trauma, cannabis use and urbanicity), sensory (hearing impairment), and familial risk factors on the occurrence of subclinical psychosis in 6,646 subjects from the general population, Pries et al. found additive effects of the risk factors: the greater the number of risk factors, the greater the odds of symptoms²¹. Consistent with this finding, in our study, the effect of recent adverse experiences (measured by the LTE) was significant for each of the 3 dimensions, independently of early adverse experience (measured by childhood trauma), and we did not find any significant interactions between the different psychosocial stressors. This may be due to a lack of statistical power to detect an ExE interaction⁵⁹, however the β values of the interactions, which are close to 0 make this unlikely.

Childhood trauma, i.e., the earliest psychosocial stressor, had the strongest associations with each of the three dimensions measured. Childhood adverse experiences have been linked with long-term changes in the hypothalamic-pituitary axis (HPA) axis⁶⁰ which may be involved in early stressor and emotional dysregulation, leading to later aberrant salience, involved in the positive dimension of psychosis⁶¹. Of note, childhood trauma is known to be associated with psychopathology as a whole^{62,63}, and consistent with prior research, childhood trauma in our study was also associated with the negative dimension⁶⁴ as well as with depressive symptoms⁶⁵ in both controls and siblings.

Self-reported discrimination experiences were associated with the positive dimension of psychosis in the global sample. Several studies, including longitudinal studies⁶⁶, have shown associations, particularly with psychotic-like experiences. They have been confirmed in recent

1
2
3 399 meta-analyses^{9,29}. Consistently with the present study, some studies suggest that among the
4 400 different dimensions of subclinical psychosis discrimination is specifically associated with
5 401 positive psychotic symptoms^{9,67}. However in contrast with the present study, discrimination
6 402 experiences have also been linked to depression⁶⁸.

7
8
9
10 403 The associations between CAPE scores and stressful life events support the role of
11 404 recent adverse experiences in the development of psychosis^{8,61}. Psychosis has been theorized
12 405 to combine cognitive (external attribution) bias and emotional dysregulation⁶⁹. Stressful life
13 406 events could play a trigger role for a disruption in cognitive processes in subjects at risk, leading
14 407 to delusional ideas and hallucinatory experiences, the content of which may be influenced by
15 408 the emotional change induced by the stressful events⁷⁰.

16
17
18
19 409 One point that also deserves discussion is the lower levels of the positive dimensions of
20 410 subclinical psychosis among siblings in comparison to controls. This result is surprising, as
21 411 siblings of people with psychotic disorder face both a greater genetic predisposition to psychotic
22 412 disorders^{71,72}, and higher levels of psychosocial stressors, because they likely share some of the
23 413 environmental characteristics of their sibling (e.g., discrimination)^{73,74}, and the stress associated
24 414 with the psychiatric disorder of their sibling⁷⁵. Indeed, consistent with the stress-vulnerability
25 415 hypothesis, several studies found higher levels of subclinical psychosis among siblings, in
26 416 comparison to controls^{64,74,76}. One hypothesis that might explain the negative association in our
27 417 study is that siblings might minimize their symptoms, either because they compare them to the
28 418 symptoms of their sibling with a FEP, or because they fear having the same disorder, and thus
29 419 deny presenting the same symptoms. Moreover, we cannot exclude a selection bias, which may
30 420 have occurred if siblings with potentially higher subclinical psychosis scores refused
31 421 participation. Nevertheless, in our study the stronger association between CAPE scores and
32 422 self-reported discrimination in siblings is consistent with the stress-vulnerability hypothesis.
33 423 Indeed, in siblings who share the genetic liability to psychosis (i.e., a vulnerability) levels of
34 424 self-reported discrimination experiences are lower than in controls, but their effect is stronger.
35 425 A similar result has been shown concerning urbanicity, which has been associated with a higher
36 426 risk for psychotic disorders among subjects with familial liability as compared to subjects
37 427 without⁷⁷.

38
39
40
41 428 Several limitations should be acknowledged. First, we performed cross-sectional
42 429 analyses of both subclinical psychosis and psychosocial stressors with retrospective assessment
43 430 for some of the variables. Due to the retrospective assessment, recall bias and reverse causality
44 431 cannot be excluded entirely. Indeed, several studies showed discrepancies between prospective
45 432 and retrospective measures⁷⁸. Retrospective assessment (e.g., for childhood trauma) may be

biased by clinical outcomes (i.e., reverse causation issue)⁷⁹ for instance. Similarly, the perception of discrimination could be distorted by the presence of positive symptoms. Likewise, the perception of a low level of social capital could be influenced by depressive symptoms. With regard to the measure of stressful life events specifically, several issues have been mentioned, including intra-category variability (i.e., the fact that subjects have differing views of what comprises a “*major*” or “*serious*” event or disease), and that this variability could also be related to psychiatric symptoms^{53,80}. Moreover, the mean discrimination score (median of discrimination measure: 0, IQR=1) was quite low in comparison with other studies (which unlike our study, were conducted among ethnic minorities facing higher levels of discrimination^{29,81,82}). This low score has been occasionally found in previous studies^{66,83}, but should be interpreted taking into consideration that the version of the Williams’ Major experiences of discrimination measure^{46,47} has not been validated, and its cross-national invariance has not yet been studied. Thus, the findings may not be generalizable to other countries (e.g., USA). Moreover, the view of discrimination as a prolonged stressor might be misconstrued, as a major part of the experiences measured occur during adulthood. Certain experiences may however take place during childhood, adolescence or youth (unfair treatment when seeking medical care, discouragement from continuing education, etc.). Furthermore, except for ethnicity, the analyses were not adjusted for other important risk factors of psychosis, such as urbanicity and cannabis use^{84,85}. However, adjustment for all the known risk factors of psychosis (economic deprivation, obstetrical complications, paternal age, etc.) was not possible, and such adjustment could cause statistical overadjustment and affect genuine relationships between subclinical psychosis and psychosocial stressors. Finally, as the sampling was not fully at random, but a mixture of random and quota sampling and thus non-probabilistic, we cannot assume that our sample was representative of the general population. However, the quota sampling method warrant same socio-demographic characteristics of the general population (age, sex, and migration) in the different countries. Moreover, we cannot exclude the possibility of selective refusal of study inclusion according to subclinical psychosis and/or psychosocial factors.

Overall, this international and multicentre study assessed positive, negative and depressive dimensions of subclinical psychosis among controls and siblings from the general population and simultaneously analyzed the role of different psychosocial stressors. Childhood trauma, and stressful life events were significantly associated with the three subclinical psychosis dimensions, while lower social capital was associated with the negative and depressive dimensions, and self-reported discrimination experiences associated with the

1
2
3 467 positive dimension, consistent with independent effects of these different psychosocial
4
5 468 stressors.
6
7 469

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

EU-GEI WP2 Group: Kathryn Hubbard, Stephanie Beards, Simona A. Stilo, Mara Parellada, Pedro Cuadrado, José Juan Rodríguez Solano, Angel Carracedo, David Fraguas, Álvaro Andreu-Bernabeu, Gonzalo López, Bibiana Cabrera, Esther Lorente-Rovira, Paz Garcia-Portilla, Javier Costas, Estela Jiménez-López, Mario Matteis, Marta Rapado-Castro, Emiliano González, Covadonga M. Díaz-Caneja, Emilio Sánchez, Manuel Durán-Cutilla, Nathalie Franke, Fabian Termorshuizen, Daniella van Dam, Elsje van der Ven, Elles Messchaart, Stéphane Jamain, Thomas Charpeaud, Anne-Marie Tronche, Flora Frijda, Giovanna Marrazzo, Lucia Sideli, Crocettarachele Sartorio, Fabio Seminerio, Camila Marcelino Loureiro, Rosana Shuhama, Mirella Ruggeri, Chiara Bonetto, Doriana Cristofalo, Domnico Berardi, Marco Seri, Elena Bonora, Giuseppe D'Andrea, Laura Ferraro, Giada Tripoli, Silvia Amoretti, Gisela Mezquida

Conflicts of interest

The funder had no involvement in study design, data collection, analysis, interpretation of findings, manuscript preparation or the decision to submit the paper for publication.

Dr. Kirkbride is supported by the National Institute for Health Research (NIHR) University College London Hospital (UCLH) Biomedical Research Centre (BRC).

Dr. Bernardo has been a consultant for, received grant/research support and honoraria from, and been on the speakers/advisory board of ABBiotics, Adamed, Angelini, Casen Recordati, Janssen-Cilag, Menarini, Rovi and Takeda.

Dr. Arango has received support by the Spanish Ministry of Science and Innovation. Instituto de Salud Carlos III (SAM16PE07CP1, PI16/02012, PI19/024), co-financed by ERDF Funds from the European Commission, "A way of making Europe", CIBERSAM. Madrid Regional Government (B2017/BMD-3740 AGES-CM-2), European Union Structural Funds. European Union Seventh Framework Program under grant agreements FP7-4-HEALTH-2009-2.2.1-2-241909 (Project EU-GEI) and FP7-HEALTH-2013-2.2.1-2-603196 (Project PSYSCAN); and European Union H2020 Program under the Innovative Medicines Initiative 2 Joint Undertaking (grant agreement No 115916, Project PRISM, and grant agreement No 777394, Project AIMS-2-TRIALS), Fundación Familia Alonso and Fundación Alicia Koplowitz.

Dr. Di Forti reports personal fees from Janssen outside the submitted work, without conflict of interest with the present study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

503

REFERENCES

1. Corcoran C, Walker E, Huot R, et al. The Stress Cascade and Schizophrenia: Etiology and Onset. *Schizophr Bull.* 2003;29(4):671-692.
2. van Os J, Kenis G, Rutten BPF. The environment and schizophrenia. *Nature.* 2010;468(7321):203-212.
3. Davis J, Eyre H, Jacka FN, et al. A review of vulnerability and risks for schizophrenia: Beyond the two hit hypothesis. *Neurosci Biobehav Rev.* 2016;65:185-194.
4. Howes OD, Murray RM. Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet.* 2014;383(9929):1677-1687.
5. Weidenauer A, Bauer M, Sauerzopf U, et al. Making Sense of: Sensitization in Schizophrenia. *Int J Neuropsychopharmacol.* 2017;20(1):1-10.
6. March D, Hatch SL, Morgan C, et al. Psychosis and place. *Epidemiol Rev.* 2008;30(1):84-100.
7. Varese F, Smeets F, Drukker M, et al. Childhood Adversities Increase the Risk of Psychosis: A Meta-analysis of Patient-Control, Prospective- and Cross-sectional Cohort Studies. *Schizophr Bull.* 2012;38(4):661-671.
8. Beards S, Gayer-Anderson C, Borges S, Dewey ME, Fisher HL, Morgan C. Life events and psychosis: a review and meta-analysis. *Schizophr Bull.* 2013;39(4):740-747.
9. Pearce J, Rafiq S, Simpson J, Varese F. Perceived discrimination and psychosis: a systematic review of the literature. *Soc Psychiatry Psychiatr Epidemiol.* 2019;54(9):1023-1044.
10. Lardinois M, Lataster T, Mengelers R, Os JV, Myin-Germeys I. Childhood trauma and increased stress sensitivity in psychosis. *Acta Psychiatr Scand.* 2011;123(1):28-35.
11. Veling W, Pot-Kolder R, Counotte J, van Os J, van der Gaag M. Environmental Social Stress, Paranoia and Psychosis Liability: A Virtual Reality Study. *Schizophr Bull.* 2016;42(6):1363-1371.
12. Zipursky RB. Why are the outcomes in patients with schizophrenia so poor? *J Clin Psychiatry.* 2014;75 Suppl 2:20-24.
13. Sariaslan A, Fazel S, D'Onofrio BM, et al. Schizophrenia and subsequent neighborhood deprivation: revisiting the social drift hypothesis using population, twin and molecular genetic data. *Transl Psychiatry.* 2016;6(5):e796.
14. Pignon B, Eaton S, Schürhoff F, Szöke A, McGorry P, O'Donoghue B. Residential social drift in the two years following a first episode of psychosis. *Schizophr Res.*

537 2019;210:323-325.

538 15. Verdoux H, van Os J. Psychotic symptoms in non-clinical populations and the
539 continuum of psychosis. *Schizophr Res.* 2002;54(1-2):59-65.

540 16. McGrath JJ, Saha S, Al-Hamzawi A, et al. Psychotic Experiences in the General
541 Population: A Cross-National Analysis Based on 31,261 Respondents From 18 Countries.
542 *JAMA Psychiatry.* 2015;72(7):697-705.

543 17. Pignon B, Schürhoff F, Szöke A, et al. Sociodemographic and clinical correlates
544 of psychotic symptoms in the general population: findings from the MHGP survey. *Schizophr*
545 *Res.* 2018;193:336-342.

546 18. Linscott RJ, van Os J. An updated and conservative systematic review and meta-
547 analysis of epidemiological evidence on psychotic experiences in children and adults: on the
548 pathway from proneness to persistence to dimensional expression across mental disorders.
549 *Psychol Med.* 2013;43(06):1133-1149.

550 19. Szöke A, Kirkbride JB, Schürhoff F. Universal prevention of schizophrenia and
551 surrogate endpoints at population level. *Soc Psychiatry Psychiatr Epidemiol.* 2014;49(9):1347-
552 1351.

553 20. van Os J. The many continua of psychosis. *JAMA Psychiatry.* 2014;71(9):985-
554 986.

555 21. Pries L-K, Gülöksüz S, Ten Have M, et al. Evidence That Environmental and
556 Familial Risks for Psychosis Additively Impact a Multidimensional Subthreshold Psychosis
557 Syndrome. *Schizophr Bull.* 2018;44(4):710-719.

558 22. Schürhoff F, Pignon B, Lajnef M, et al. Psychotic experiences are associated
559 with paternal age but not with delayed fatherhood in a large, multinational, community sample.
560 *Schizophr Bull.* 2020;46(5):1327-1334.

561 23. Cougnard A, Marcelis M, Myin-Germeys I, et al. Does normal developmental
562 expression of psychosis combine with environmental risk to cause persistence of psychosis? A
563 psychosis proneness-persistence model. *Psychol Med.* 2007;37(4):513-527.

564 24. Morgan C, Reininghaus U, Reichenberg A, et al. Adversity, cannabis use and
565 psychotic experiences: evidence of cumulative and synergistic effects. *Br J Psychiatry J Ment*
566 *Sci.* 2014;204:346-353.

567 25. McGrath JJ, Saha S, Lim CCW, et al. Trauma and psychotic experiences:
568 transnational data from the World Mental Health Survey. *Br J Psychiatry J Ment Sci.*
569 2017;211(6):373-380.

570 26. Pignon B, Peyre H, Szöke A, et al. A latent class analysis of psychotic symptoms

in the general population. *Aust N Z J Psychiatry*. 2018;52(6):573-584.

27. Johns LC, Cannon M, Singleton N, et al. Prevalence and correlates of self-reported psychotic symptoms in the British population. *Br J Psychiatry J Ment Sci*. 2004;185:298-305.

28. Kocsis-Bogár K, Miklósi M, Forintos DP. Impact of adverse life events on individuals with low and high schizotypy in a nonpatient sample. *J Nerv Ment Dis*. 2013;201(3):208-215.

29. Bardol O, Grot S, Oh H, et al. Perceived ethnic discrimination as a risk factor for psychotic symptoms: a systematic review and meta-analysis. *Psychol Med*. 2020;50(7):1077-1089.

30. Kirkbride JB, Boydell J, Ploubidis GB, et al. Testing the association between the incidence of schizophrenia and social capital in an urban area. *Psychol Med*. 2008;38(8):1083-1094.

31. O'Donoghue B, Lyne JP, Renwick L, et al. Neighbourhood characteristics and the incidence of first-episode psychosis and duration of untreated psychosis. *Psychol Med*. 2016;46(7):1367-1378.

32. Solmi F, Colman I, Weeks M, Lewis G, Kirkbride JB. Trajectories of Neighborhood Cohesion in Childhood, and Psychotic and Depressive Symptoms at Age 13 and 18 Years. *J Am Acad Child Adolesc Psychiatry*. 2017;56(7):570-577.

33. Lataster J, Myin-Germeys I, Lieb R, Wittchen H-U, van Os J. Adversity and psychosis: a 10-year prospective study investigating synergism between early and recent adversity in psychosis. *Acta Psychiatr Scand*. 2012;125(5):388-399.

34. Gage SH, Jones HJ, Burgess S, et al. Assessing causality in associations between cannabis use and schizophrenia risk: a two-sample Mendelian randomization study. *Psychol Med*. 2017;47(5):971-980.

35. Colodro-Conde L, Couvy-Duchesne B, Whitfield JB, et al. Association Between Population Density and Genetic Risk for Schizophrenia. *JAMA Psychiatry*. 2018;75(9):901-910.

36. Gaye-Anderson C, Jongsma HE, Di Forti M, et al. The EUropean Network of National Schizophrenia Networks Studying Gene-Environment Interactions (EU-GEI): Incidence and First-Episode Case-Control Programme. *Soc Psychiatry Psychiatr Epidemiol*. 2020;55(5):645-657.

37. European Network of National Networks studying Gene-Environment Interactions in Schizophrenia (EU-GEI). Identifying Gene-Environment Interactions in

- Schizophrenia: Contemporary Challenges for Integrated, Large-scale Investigations. *Schizophr Bull.* 2014;40(4):729-736.
38. Jongsma HE, Gayer-Anderson C, Lasalvia A, et al. Treated incidence of psychotic disorders in the multinational EU-GEI study. *JAMA Psychiatry.* 2018;75(1):36-46.
39. Stefanis NC, Hanssen M, Smirnis NK, et al. Evidence that three dimensions of psychosis have a distribution in the general population. *Psychol Med.* 2002;32(2):347-358.
40. Mark W, Touloupoulou T. Psychometric Properties of “Community Assessment of Psychic Experiences”: Review and Meta-analyses. *Schizophr Bull.* 2016;42(1):34-44.
41. Yung AR, Nelson B, Baker K, Buckby JA, Baksheev G, Cosgrave EM. Psychotic-like experiences in a community sample of adolescents: implications for the continuum model of psychosis and prediction of schizophrenia. *Aust N Z J Psychiatry.* 2009;43(2):118-128.
42. Wigman JTW, Vollebergh WAM, Raaijmakers QAW, et al. The structure of the extended psychosis phenotype in early adolescence: a cross-sample replication. *Schizophr Bull.* 2011;37(4):850-860.
43. Pignon B, Peyre H, Ferchiou A, et al. Assessing cross-national invariance of the Community Assessment of Psychic Experiences (CAPE). *Psychol Med.* 2019;49(16):2600-2607.
44. Bernstein DP, Stein JA, Newcomb MD, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl.* 2003;27(2):169-190.
45. Williams DR, Yu Y, Jackson JS, Anderson NB. Racial Differences in Physical and Mental Health Socio-economic Status, Stress and Discrimination. *J Health Psychol.* 1997;2(3):335-351.
46. Jongsma HE, Gayer-Anderson C, Tarricone I, et al. Social disadvantage, linguistic distance, ethnic minority status and first-episode psychosis: results from the EU-GEI case-control study. *Psychol Med.* 2020;in press:1-13.
doi:doi.org/10.1017/S003329172000029X
47. Hatch SL, Gizard B, Williams DR, Frissa S, Goodwin L, Hotopf M. Discrimination and common mental disorder among migrant and ethnic groups: findings from a South East London Community sample. *Soc Psychiatry Psychiatr Epidemiol.* 2016;51:689-701.
48. Lochner K, Kawachi I, Kennedy BP. Social capital: a guide to its measurement. *Health Place.* 1999;5(4):259-270.

49. Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and Violent Crime: A Multilevel Study of Collective Efficacy. *Science*. 1997;277(5328):918-924.
50. Drukker M, Krabbendam L, Driessen G, van Os J. Social disadvantage and schizophrenia. A combined neighbourhood and individual-level analysis. *Soc Psychiatry Psychiatr Epidemiol*. 2006;41(8):595-604.
51. McCulloch A. An examination of social capital and social disorganisation in neighbourhoods in the British household panel study. *Soc Sci Med* 1982. 2003;56(7):1425-1438.
52. Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med*. 1985;15(1):189-194.
53. Motrico E, Moreno-Küstner B, de Dios Luna J, et al. Psychometric properties of the List of Threatening Experiences—LTE and its association with psychosocial factors and mental disorders according to different scoring methods. *J Affect Disord*. 2013;150(3):931-940.
54. Tortelli A, Nakamura A, Suprani F, et al. Subclinical psychosis in adult migrants and ethnic minorities: systematic review and meta-analysis. *BJPsych Open*. 2018;4(6):510-518.
55. Leane E, Dealberto M-J, Luck D, et al. Ethnic minority position and migrant status as risk factors for psychotic symptoms in the general population: a meta-analysis. *Psychol Med*. 2019;49(4):545-558.
56. Murphy D, Vallières F, Murphy J, McElroy E, Hyland P. Risk factors associated with general and specific dimensions of psychosis in a nationally representative sample of adults from the United States. *Psychosis*. 2020;12(4):303-3013. doi:10.1080/17522439.2020.1791238
57. Kirkbride JB, Stochl J, Zimbrón J, et al. Social and spatial heterogeneity in psychosis proneness in a multilevel case–prodrome–control study. *Acta Psychiatr Scand*. 2015;132(4):283-292.
58. O'Donoghue B, Yung AR, Wood S, et al. Neighbourhood characteristics and the rate of identification of young people at ultra-high risk for psychosis. *Schizophr Res*. 2015;169(1):214-216.
59. Hunter DJ. Gene–environment interactions in human diseases. *Nat Rev Genet*. 2005;6(4):287-298.
60. Aas M, Pizzagalli DA, Laskemoen JF, et al. Elevated hair cortisol is associated with childhood maltreatment and cognitive impairment in schizophrenia and in bipolar disorders. *Schizophr Res*. 2019;213:65-71.

- 673 61. Reininghaus U, Kempton MJ, Valmaggia L, et al. Stress Sensitivity, Aberrant
674 Salience, and Threat Anticipation in Early Psychosis: An Experience Sampling Study.
675 *Schizophr Bull.* 2016;42(3):712-722.
- 676 62. Green JG, McLaughlin KA, Berglund PA, et al. Childhood adversities and adult
677 psychopathology in the National Comorbidity Survey Replication (NCS-R) I: Associations
678 with first onset of DSM-IV disorders. *Arch Gen Psychiatry.* 2010;67(2):113.
- 679 63. McLaughlin K, Green J, Gruber M, Sampson NA, Zaslavsky A, Kessler R.
680 Childhood adversities and adult psychiatric disorders in the national comorbidity survey
681 replication II: Associations with persistence of DSM-IV disorders. *Arch Gen Psychiatry.*
682 2010;67(2):124-132.
- 683 64. Schürhoff F, Laguerre A, Fisher H, et al. Self-reported childhood trauma
684 correlates with schizotypal measures in schizophrenia but not bipolar pedigrees. *Psychol Med.*
685 2009;39(3):365-370.
- 686 65. van Dam DS, van Nierop M, Viechtbauer W, et al. Childhood abuse and neglect
687 in relation to the presence and persistence of psychotic and depressive symptomatology.
688 *Psychol Med.* 2015;45(7):1363-1377.
- 689 66. Janssen I, Hanssen M, Bak M, et al. Discrimination and delusional ideation. *Br*
690 *J Psychiatry.* 2003;182:71-76.
- 691 67. Kong DT. Ethnic minorities' paranoia and self-preservative work behaviors in
692 response to perceived ethnic discrimination, with collective self-esteem as a buffer. *J Occup*
693 *Health Psychol.* 2016;21(3):334-351.
- 694 68. Schmitt MT, Branscombe NR, Postmes T, Garcia A. The consequences of
695 perceived discrimination for psychological well-being: A meta-analytic review. *Psychol Bull.*
696 2014;140(4):921.
- 697 69. Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE. A cognitive model
698 of the positive symptoms of psychosis. *Psychol Med.* 2001;31(2):189-195.
- 699 70. Myin-Germeys I, Delespaul P, Os JV. Behavioural sensitization to daily life
700 stress in psychosis. *Psychol Med.* 2005;35(5):733-741.
- 701 71. Mortensen PB, Pedersen C, Westergaard T, et al. Effects of family history and
702 place and season of birth on the risk of schizophrenia. *N Engl J Med.* 1999;340(8):603-608.
- 703 72. Trotta A, Murray RM, Fisher HL. The impact of childhood adversity on the
704 persistence of psychotic symptoms: a systematic review and meta-analysis. *Psychol Med.*
705 2015;45(12):2481-2498.
- 706 73. Polanczyk G, Moffitt TE, Arseneault L, et al. Etiological and clinical features of

- childhood psychotic symptoms: results from a birth cohort. *Arch Gen Psychiatry*. 2010;67(4):328-338.
74. Binbay T, Drukker M, Elbi H, et al. Testing the psychosis continuum: differential impact of genetic and nongenetic risk factors and comorbid psychopathology across the entire spectrum of psychosis. *Schizophr Bull*. 2012;38(5):992-1002.
75. Sin J, Murrells T, Spain D, Norman I, Henderson C. Wellbeing, mental health knowledge and caregiving experiences of siblings of people with psychosis, compared to their peers and parents: an exploratory study. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51(9):1247-1255.
76. van Dam DS, Korver-Nieberg N, Velthorst E, Meijer CJ, de Haan L, For Genetic Risk and Outcome in Psychosis (GROUP). Childhood maltreatment, adult attachment and psychotic symptomatology: a study in patients, siblings and controls. *Soc Psychiatry Psychiatr Epidemiol*. 2014;49(11):1759-1767.
77. van Os J, Hanssen M, Bak M, Bijl RV, Vollebergh W. Do urbanicity and familial liability coparticipate in causing psychosis? *Am J Psychiatry*. 2003;160(3):477-482.
78. Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement Between Prospective and Retrospective Measures of Childhood Maltreatment: A Systematic Review and Meta-analysis. *JAMA Psychiatry*. 2019;76(6):584-593.
79. MacDonald K, Thomas ML, MacDonald TM, Sciolla AF. A Perfect Childhood? Clinical Correlates of Minimization and Denial on the Childhood Trauma Questionnaire. *J Interpers Violence*. 2015;30(6):988-1009.
80. Dohrenwend BP. Inventorying stressful life events as risk factors for psychopathology: Toward resolution of the problem of intracategory variability. *Psychol Bull*. 2006;132(3):477-495.
81. Oh H, Yang LH, Anglin DM, DeVlyder JE. Perceived discrimination and psychotic experiences across multiple ethnic groups in the United States. *Schizophr Res*. 2014;157(1):259-265.
82. Oh H, Cogburn CD, Anglin D, Lukens E, DeVlyder J. Major discriminatory events and risk for psychotic experiences among Black Americans. *Am J Orthopsychiatry*. 2016;86(3):277-285.
83. Rakhshan Rouhakhtar PJ, Pitts SC, Schiffman J. Associations between Race, Discrimination, Community Violence, Traumatic Life Events, and Psychosis-Like Experiences in a Sample of College Students. *J Clin Med*. 2019;8(10).
84. Di Forti M, Quattrone D, Freeman TP, et al. The contribution of cannabis use to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. *Lancet Psychiatry*. 2019;6(5):427-436.

85. Vassos E, Pedersen CB, Murray RM, Collier DA, Lewis CM. Meta-analysis of the association of urbanicity with schizophrenia. *Schizophr Bull*. 2012;38(6):1118-1123.

Table 1. Description and comparisons of the CAPE and of the psychosocial stress measures according to control/sibling status			
	Controls (N = 1497)	Siblings (N = 265)	Comparisons
	Median (IQR), mean (SD) or N (%)	Median (IQR), mean (SD) or N (%)	
Age	33 (21), 36.1 (12.9)	30 (15), 31.3 (9.4), 1.9%	< 0.01 ²
Sex			
Women	791 (47.2%)	181 (31.7%)	< 0.01 ¹
Men	706 (52.8%)	84 (68.3%)	
Ethnicity			
Asian	33 (2.2%)	3 (1.1%)	< 0.01 ¹
Black	121 (8.1%)	13 (4.9%)	
North African	24 (1.6%)	6 (2.3%)	
White	1178 (78.7%)	201 (75.9%)	
Mixed	116 (7.7%)	41 (15.5%)	
Other	24 (1.7%)	1 (3.8%)	
CAPE scores			
Positive	4 (4), 4.9 (2.9)	4 (4), 4.5 (3.0)	0.02 ²
Negative	6 (6), 6.2 (3.6)	6 (6), 5.8 (3.7)	0.18 ²
Depressive	4 (3), 4.4 (2.0)	4 (3), 4.2 (2.1)	0.13 ²
Psychosocial stressors			
Childhood trauma	31 (11), 34.6 (10.9)	33 (12), 37.3 (6.9)	< 0.01 ²
Self-reported discrimination experiences	0.0 (1.0), 0.6 (1.0)	0.0 (1.0), 0.4 (0.9)	0.02 ²
Stressful life events	1 (2), 1.5 (1.4)	1 (1), 1.5 (1.5)	0.29 ²
Social capital	0.0 (3.4), 0 (2.5)	0.1 (3.4), 0 (2.5)	0.56 ²
Abbreviations: IQR = interquartile range, NA: unavailable data, SD = standard-deviation. Legend: ¹ p-value of chi-square tests; ² p-value of Mann-Whitney.			

Table 2. Spearman correlation matrix of the psychosocial stressors				
	Childhood trauma	Self-reported discrimination experiences	Stressful life events	Social capital
Childhood trauma	1.00			
Self-reported discrimination experiences	0.17***	1.00		
Stressful life events	0.14***	0.20***	1.00	
Social capital	-0.15***	-0.08***	-0.11***	1.00
Legend: p-values: *: < 0.05, **: < 0.01, ***: < 0.001.				

Table 3. Spearman correlations between CAPE scores and psychosocial stressors			
	Positive dimension ¹	Negative dimension ¹	Depressive dimension ¹
Childhood trauma	0.26	0.29	0.27
Self-reported discrimination experiences	0.15	0.13	0.12
Stressful life events	0.18	0.13	0.20
Social capital	-0.13	-0.13	-0.14
¹ all p-values < 0.001			

Table 4. Multivariable ¹ analyses of the relationships between the Z-scores of the different psychosocial stressors and the 3-dimension scores						
	Positive dimension		Negative dimension		Depressive dimension	
	β	95 % CI	β	95 % CI	β	95 % CI
Childhood trauma	0.13	[0.10; 0.17]***	0.49	[0.37; 0.61]***	0.26	[0.19; 0.33]***
Self-reported discrimination experiences	0.06	[0.02; 0.10]**	0.05	[-0.09; 0.19]	0.05	[-0.03; 0.13]
Stressful life events	0.08	[0.04; 0.12]***	0.16	[0.04; 0.28]*	0.17	[0.11; 0.24]***
Lower social capital	0.03	[0.00; 0.07]	0.26	[0.14; 0.38]***	0.13	[0.06; 0.20]***
Siblings (vs. controls)	-0.16	[-0.26; -0.05]**	-0.28	[-0.62; -0.06]	-0.19	[-0.19; 0.01]
¹ Adjusted for age, sex, country, and ethnicity. Legend: *: < 0.05, **: < 0.01, ***: < 0.001.						

DISCRIMINATION

Supplementary material 1: Discrimination measure in the EUGEI study

In the following questions we are interested in the way other people have treated you or your beliefs about how other people have treated you. Can you tell me if any of the following has ever happened to you? Please indicate number of times, age at first occurrence and note the main reason for this. **Then:**

For any reason, have you ever been unfairly...

	Yes	No	N of times	Age (first occurred)
1. Fired	<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

	Yes	No	N of times	Age (first occurred)
2. Not hired for a job	<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

	Yes	No	N of times	Age
3. Denied promotion	<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

	Yes	No	N of times	Age
4. Stopped, questioned threatened by police	<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

DISCRIMINATION

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

5. Treated by court system

Yes	No	N of times	Age
<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

For any reason, have you ever been unfairly...

6. Discouraged from continuing education

Yes	No	N of times	Age
<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

7. Prevented from buying, renting flat or house

Yes	No	N of times	Age
<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason O1 Gender O2 Race, ethnicity O3 Religion O4 Mental Illness O5 Sexuality O6 Age O7 Other (specify):

8. Treated by neighbours or your family

Yes	No	N of times	Age
<input type="text"/>	<input type="text"/>	<input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/>

Reason

DISCRIMINATION

O1 Gender	O2 Race, ethnicity	O3 Religion	O4 Mental Illness	O5 Sexuality	O6 Age	O7 Other (specify):
--------------	-----------------------	----------------	----------------------	-----------------	-----------	------------------------

Yes

No

N of
times

Age

9. Denied a loan or preferable mortgage rate

Reason	O1 Gender	O2 Race, ethnicity	O3 Religion	O4 Mental Illness	O5 Sexuality	O6 Age	O7 Other (specify):
---------------	--------------	-----------------------	----------------	----------------------	-----------------	-----------	------------------------

Yes

No

N of
times

Age

10. Received worse service than other people

Reason	O1 Gender	O2 Race, ethnicity	O3 Religion	O4 Mental Illness	O5 Sexuality	O6 Age	O7 Other (specify):
---------------	--------------	-----------------------	----------------	----------------------	-----------------	-----------	------------------------

Yes

No

N of
times

Age

11. Treated when getting medical care

Reason	O1 Gender	O2 Race, ethnicity	O3 Religion	O4 Mental Illness	O5 Sexuality	O6 Age	O7 Other (specify):
---------------	--------------	-----------------------	----------------	----------------------	-----------------	-----------	------------------------

Yes

No

N of
times

Age

12. Treated when using public transport

Reason	O1 Gender	O2 Race, ethnicity	O3 Religion	O4 Mental Illness	O5 Sexuality	O6 Age	O7 Other (specify):
---------------	--------------	-----------------------	----------------	----------------------	-----------------	-----------	------------------------

Supplementary Table 1. Proportions of unavailable data (%)		
	Controls (N = 1497)	Siblings (N = 265)
Age	0.2%	1.9%
Sex	0.0%	0.0%
Ethnicity	0.1%	0.0%
CAPE scores		
Positive	5.3%	8.3%
Negative	4.5%	6.8%
Depressive	4.8%	6.8%
Psychosocial stressors		
Childhood trauma	0.8%	3.8%
Self-reported discrimination experiences	4.4%	4.5%
Stressful life events	16.5%	17.4%
Social capital	10.4%	7.5%

Supplementary Table 2. Spearman correlation matrix of the psychosocial stressors among controls and siblings				
<i>CONTROLS</i>				
	Childhood trauma	Self-reported discrimination experiences	Stressful life events	Social capital
Childhood trauma	1.00			
Self-reported discrimination experiences	0.17***	1.00		
Stressful life events	0.13***	0.20***	1.00	
Social capital	-0.14***	-0.09***	-0.13***	1.00
<i>SIBLINGS</i>				
	Childhood trauma	Self-reported discrimination experiences	Stressful life events	Social capital
Childhood trauma	1.00			
Self-reported discrimination experiences	0.24***	1.00		
Stressful life events	0.21**	0.26***	1.00	
Social capital	-0.19***	-0.08	0.03	1.00
Legend: p-values: *: < 0.05, **: < 0.01, ***: < 0.001.				

Supplementary Table 3. Spearman correlations between CAPE scores and psychosocial stressors among controls and siblings			
<i>CONTROLS</i>			
	Positive dimension	Negative dimension	Depressive dimension
Childhood trauma	0.26***	0.30***	0.26***
Self-reported discrimination experiences	0.14***	0.12***	0.10***
Stressful life events	0.19***	0.13***	0.19***
Social capital	-0.13***	-0.12***	-0.13***
<i>SIBLINGS</i>			
	Positive dimension	Negative dimension	Depressive dimension
Childhood trauma	0.26***	0.27***	0.35***
Self-reported discrimination experiences	0.24***	0.21***	0.19**
Stressful life events	0.19**	0.18*	0.27***
Social capital	-0.12	-0.20**	-0.22***
Legend: p-values: *: < 0.05, **: < 0.01, ***: < 0.001.			

Supplementary Table 4. Multivariable analyses of the relationships between the Z-scores of the different psychosocial stressors and the CAPE scores among pooled siblings and controls						
	Positive dimension		Negative dimension		Depressive dimension	
	β	95 % CI	β	95 % CI	β	95 % CI
<i>One model per interaction¹</i>						
Childhood trauma x self-reported discrimination experiences	-0.03	[-0.06; 0.01]	0.00	[-0.09; 0.10]	0.00	[-0.01; 0.00]
Self-reported discrimination experiences x stressful life events	-0.01	[-0.03; 0.01]	-0.01	[-0.08; 0.06]	-0.03	[-0.06; 0.06]
Self-reported discrimination experiences x low social capital	0.00	[-0.03; 0.03]	-0.06	[-0.17; 0.05]	-0.03	[-0.07; 0.01]
Childhood trauma x stressful life events	-0.01	[-0.03; 0.02]	-0.02	[-0.09; 0.05]	-0.04	[-0.02; 0.06]
Childhood trauma x low social capital	-0.03	[-0.06; 0.00]	-0.04	[-0.15; 0.07]	0.02	[-0.10; 0.02]
Low social capital x stressful life events	0.02	[-0.01; 0.04]	0.04	[-0.04; 0.12]	0.01	[-0.03; 0.06]
<i>All interaction terms in one single model²</i>						
Childhood trauma x self-reported discrimination experiences	-0.10	[-0.23; 0.02]	-0.10	[-0.61; 0.20]	-0.01	[-0.26; 0.24]
Self-reported discrimination experiences x stressful life events	-0.13	[-0.32; 0.05]	-0.13	[-0.59; 0.57]	0.01	[-0.34; 0.37]
Self-reported discrimination experiences x low social capital	-0.08	[-0.31; 0.15]	0.02	[-0.76; 0.73]	-0.14	[-0.58; 0.30]
Childhood trauma x stressful life events	0.00	[-0.07; 0.08]	0.00	[-0.26; 0.24]	0.05	[-0.09; 0.20]
Childhood trauma x low social capital	-0.04	[-0.14; 0.06]	-0.04	[-0.14; 0.10]	-0.12	[-0.31; 0.07]
Low social capital x stressful life events	0.02	[-0.07; 0.10]	0.02	[-0.04; 0.08]	0.09	[-0.07; 0.26]
¹ The interaction terms were introduced one by one in the multivariable analyses with the following factors: psychosocial stressors (childhood trauma, self-reported discrimination experiences, stressful life events, low social capital) and the adjustment factors (age, sex, ethnicity, country and sibling/control status); ² All interaction terms in the same model, with the same adjustment factors. Legend: *: < 0.05, **: < 0.01, ***: < 0.001.						

Supplementary Table 5. Multivariable ¹ analyses of the associations between the CAPE and the Z-scores of the psychosocial stressors among controls and siblings						
CONTROLS						
	Positive dimension		Negative dimension		Depressive dimension	
	β	95 % CI	β	95 % CI	β	95 % CI
Childhood trauma	0.14	[0.10; 0.18]***	0.44	[0.33; 0.60]***	0.24	[0.17; 0.32]***
Self-reported discrimination experiences	0.05	[0.01; 0.10]*	0.05	[-0.09; 0.20]	0.05	[-0.03; 0.13]
Stressful life events	0.08	[0.04; 0.12]***	0.16	[0.03; 0.29]*	0.18	[0.11; 0.25]***
Lower social capital	0.03	[-0.01; 0.07]	0.22	[0.09; 0.35]**	0.10	[0.03; 0.18]**
Interactions between psychosocial stressors						
Childhood trauma x self-reported discrimination experiences	-0.03	[-0.06; 0.01]	0.03	[-0.08; 0.13]	0.00	[-0.06; 0.06]
Self-reported discrimination experiences x stressful life events	-0.01	[-0.03; 0.01]	0.00	[-0.07; 0.07]	-0.03	[-0.07; 0.01]
Self-reported discrimination experiences x lower social capital	0.01	[-0.03; 0.04]	-0.04	[-0.15; 0.07]	-0.02	[-0.09; 0.04]
Childhood trauma x stressful life events	-0.01	[-0.03; 0.02]	-0.02	[-0.10; 0.06]	0.02	[-0.03; 0.06]
Childhood trauma x lower social capital	-0.03	[-0.07; 0.01]	-0.03	[-0.15; 0.10]	-0.04	[-0.11; 0.03]
Lower social capital x stressful life events	0.02	[-0.01; 0.05]	0.03	[-0.05; 0.12]	0.01	[-0.04; 0.06]
SIBLINGS						
	Positive dimension		Negative dimension		Depressive dimension	
	β	95 % CI	β	95 % CI	β	95 % CI
Childhood trauma	0.09	[-0.01; 0.20]	0.58	[0.26; 0.90]***	0.32	[0.12; 0.52]**
Self-reported discrimination experiences	0.21	[0.03; 0.39]*	0.27	[-0.29; 0.82]	0.17	[-0.18; 0.51]
Stressful life events	0.12	[0.00; 0.24]	0.18	[-0.19; 0.55]	0.11	[-0.11; 0.34]
Lower social capital	0.08	[-0.03; 0.19]	0.50	[0.16; 0.85]**	0.30	[0.09; 0.51]**
Interactions between psychosocial stressors						
Childhood trauma x self-reported discrimination experiences	-0.15	[-0.27; -0.03]*	-0.25	[-0.61; 0.12]	-0.04	[-0.26; 0.19]
Self-reported discrimination experiences x stressful life events	-0.15	[-0.31; 0.01]	-0.11	[-0.60; 0.38]	0.05	[-0.25; 0.35]
Self-reported discrimination experiences x lower social capital	-0.09	[-0.29; 0.10]	-0.17	[-0.79; 0.44]	-0.21	[-0.58; 0.15]
Childhood trauma x stressful life events	-0.01	[-0.08; 0.06]	-0.03	[-0.25; 0.19]	0.06	[-0.07; 0.20]
Childhood trauma x lower social capital	-0.01	[-0.04; 0.06]	-0.18	[-0.44; 0.09]	-0.10	[-0.27; 0.06]
Lower social capital x stressful life events	0.01	[-0.06; 0.09]	0.08	[-0.15; 0.32]	0.06	[-0.09; 0.20]

¹Adjusted for age, sex, ethnicity, and country.
Legend: *: < 0.05, **: < 0.01, ***: < 0.001.