



HAL
open science

The MPA Guide: A framework to achieve global goals for the ocean

Kirsten Grorud-Colvert, Jenna Sullivan-Stack, Callum Roberts, Vanessa Constant, Barbara Horta E Costa, Elizabeth P Pike, Naomi Kingston, Dan Laffoley, Enric Sala, Joachim Claudet, et al.

► To cite this version:

Kirsten Grorud-Colvert, Jenna Sullivan-Stack, Callum Roberts, Vanessa Constant, Barbara Horta E Costa, et al.. The MPA Guide: A framework to achieve global goals for the ocean. *Science*, 2021, 373 (6560), 10.1126/science.abf0861 . hal-03340433

HAL Id: hal-03340433

<https://hal.science/hal-03340433v1>

Submitted on 10 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title: The MPA Guide: A Framework to Achieve Global Goals for the Ocean

Authors: Kirsten Grorud-Colvert^{1*}, Jenna Sullivan-Stack¹, Callum Roberts², Vanessa Constant³, Barbara Horta e Costa⁴, Elizabeth P. Pike⁵, Naomi Kingston⁶, Dan Laffoley⁷, Enric Sala⁸, Joachim Claudet⁹, Alan M. Friedlander^{10,11}, David A. Gill¹², Sarah E. Lester¹³, Jon C. Day¹⁴, Emanuel J. Gonçalves^{15,16}, Gabby N. Ahmadi¹⁷,
5 Matt Rand¹⁸, Angelo Villagomez¹⁸, Natalie C. Ban¹⁹, Georgina G. Gurney²⁰, Ana K. Spalding^{21,22}, Nathan J. Bennett²³, Johnny Briggs¹⁸, Lance E. Morgan²⁴, Russell Moffitt⁵, Marine Deguignet⁶, Ellen K. Pikitch²⁵, Emily S. Darling²⁶, Sabine Jessen²⁷, Sarah O. Hameed²⁸, Giuseppe Di Carlo²⁹, Paolo Guidetti^{30,31}, Jean M. Harris³², Jorge Torre³³, Zafer Kizilkaya³⁴, Tundi Agardy³⁵, Philippe Cury³⁶, Nirmal J. Shah³⁷, Karen Sack³⁸, Ling Cao³⁹, Miriam Fernandez⁴⁰, Jane Lubchenco¹

Affiliations:

¹ Department of Integrative Biology, Oregon State University, 3029 Cordley Hall, Corvallis, OR, USA.

² Department of Environment and Geography, University of York, York YO10 5DD, UK.

³ Department of Integrative Biology, Oregon State University, 3029 Cordley Hall, Corvallis, OR, USA (Current affiliation: The National Academies of Sciences, Engineering, and Medicine, 500 5th St., NW, Washington, DC 20001).

⁴ Center of Marine Sciences, CCMAR, University of Algarve, Campus de Gambelas, Faro, 8005-139, Portugal.

⁵ Marine Protection Atlas, Marine Conservation Institute, 1914 N 24th Street Ste 400, Seattle, WA, 98103-9090, USA.

⁶ UN Environment Programme World Conservation Monitoring Centre, Cambridge, UK.

⁷ IUCN World Commission on Protected Areas, IUCN (International Union for Conservation of Nature), 28 rue Mauverney, CH-1196 Gland, Switzerland.

⁸ National Geographic Society, Washington, DC, USA.

⁹ National Center for Scientific Research, PSL Université Paris, CRIOBE, USR 3278 CNRS-EPHE-UPVD, Maison des Océans, 195 rue Saint-Jacques 75005 Paris, France.

¹⁰ Hawai'i Institute of Marine Biology, University of Hawai'i, Kāne'ohe, HI, 96744, USA.

¹¹ Pristine Seas, National Geographic Society, Washington, DC, 20036, USA.

¹² Duke University Marine Laboratory, Nicholas School of the Environment, Duke University, Beaufort, North Carolina 28516, USA.

¹³ Department of Geography, Florida State University, 113 Collegiate Loop, Tallahassee FL 32306-2190, USA.

¹⁴ ARC Centre of Excellence in Coral Reef Studies, James Cook University, Townsville QLD 4811, Australia.

¹⁵ MARE - Marine and Environmental Sciences Centre, ISPA – Instituto Universitário, Rua Jardim do Tabaco 34, 1149-041 Lisbon, Portugal.

¹⁶ Oceano Azul Foundation, Oceanário de Lisboa, Esplanada D. Carlos I, 1990-005 Lisbon, Portugal.

- 17 Ocean Conservation, World Wildlife Fund, Washington, D.C. 20037, USA.
- 18 Pew Bertarelli Ocean Legacy Project, The Pew Charitable Trusts, 901 E Street NW, Washington, DC 20004-2008, USA.
- 19 School of Environmental Studies, University of Victoria, Victoria, BC V8W 2Y2, Canada.
- 5 20 Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland 4811, Australia.
- 21 School of Public Policy, Oregon State University, Corvallis, OR 97330, USA.
- 22 Smithsonian Tropical Research Institute, Panama City, Panama; Coiba Scientific Station (Coiba AIP), Panama City, Panama.
- 10 23 The Peopled Seas Initiative, Vancouver, BC, Canada.
- 24 Marine Conservation Institute, Seattle, WA 98103, USA.
- 25 School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794, USA.
- 26 Wildlife Conservation Society, 2300 Southern Blvd, Bronx, NY 10460, USA.
- 15 27 National Ocean Program, Canadian Parks and Wilderness Society, Ottawa, ON, K2P 0A4, Canada.
- 28 Blue Parks Program, Marine Conservation Institute, Seattle, WA 98103, USA.
- 29 World Wide Fund For Nature (WWF) - Mediterranean, Rome 00198, Italy.
- 30 30 Department of Integrative Marine Ecology (EMI), Stazione Zoologica A. Dohrn - National Institute of Marine Biology, Ecology and Biotechnology, Villa Comunale, 80121, Naples, Italy.
- 31 National Research Council, Institute for the Study of Anthropic Impact and Sustainability in the Marine Environment (CNR-IAS), Via de Marini 6, 16149, Genoa, Italy.
- 32 Wildlands Conservation Trust, Hilton, South Africa.
- 33 Comunidad y Biodiversidad, A.C. Isla del Peruano 215, Col. Lomas de Miramar, Guaymas, Sonora, 85454, Mexico.
- 25 34 Mediterranean Conservation Society, Folkart Time 1. Block 807, Bornova, Izmir 35100 Turkey.
- 35 Sound Seas, 26 Van Nuys Road, Colrain, MA 01340, USA.
- 36 MARBEC, Montpellier University, CNRS, IRD, IFREMER, Sète, France.
- 30 37 Nature Seychelles, Centre for Environment and Education, Sanctuary at Roche Caiman, Mahe, Seychelles.
- 38 Ocean Unite, 2336 Wisconsin Ave, NW #32043 Washington, DC, 20007, USA.
- 39 School of Oceanography, Shanghai Jiao Tong University, Shanghai 230000, China.
- 35 40 Estación Costera de Investigaciones Marinas de Las Cruces and Departamento de Ecología, Pontificia Universidad Católica de Chile, Santiago, Chile.

*Correspondence to: grorudck@oregonstate.edu.

Abstract: Marine Protected Areas (MPAs) are conservation tools intended to protect biodiversity, promote healthy and resilient marine ecosystems, and provide societal benefits. Despite codification of MPAs in international agreements, MPA effectiveness is currently undermined by confusion about the many MPA types and consequent wildly differing outcomes. We present a clarifying science-driven framework – The MPA Guide – to aid design and evaluation. The guide categorizes MPAs by stage of establishment and level of protection, specifies the resulting direct and indirect outcomes for biodiversity and human well-being, and describes the key conditions necessary for positive outcomes. Use of this MPA guide by scientists, managers, policy makers, and communities can improve effective design, implementation, assessment, and tracking of existing and future MPAs to achieve conservation goals using scientifically grounded practices.

One Sentence Summary: A new framework facilitates design and evaluation of marine protected areas to protect biodiversity and benefit people.

Structured Abstract:

Background

Marine Protected Areas (MPAs) are places in the ocean that receive protection to safeguard biodiversity from abatable threats. Despite international agreements and decades of research, confusion exists about the definition of ‘protection’, the conditions under which an MPA is effective, and potential MPA outcomes. Not all MPAs are the same; they range from full

protection in “no-take” areas to minimal protection with many extractive activities. Some exist only on paper, not in practice. The resulting divergent outcomes and confusion fuels controversies about efficacy, undermines confidence in MPAs, and jeopardizes conservation goals, including those prescribed by the Convention on Biological Diversity and UN Sustainable Development Agenda. Clarity is needed about types of MPAs, outcomes, and conditions for success.

Advances

We propose a science-based, policy-relevant framework – The MPA Guide – to categorize, track, evaluate, and plan MPAs. The guide specifies when protection begins, which activities are allowed, what conditions are required for success, and what outcomes can be expected. It emphasizes effectiveness of protection by considering MPA quality and not just quantity. It complements the well-known IUCN Protected Areas Categories, which are based on management objectives and governance types, not level of protection. Together, the IUCN categories and this MPA guide enable a comprehensive picture of any MPA.

This guide consists of four elements. The first describes the stage of establishment (STAGE) of an MPA from initial proposal through to completion. This information is important because an MPA does not provide protection until it is activated in the water. The four STAGES are: 1) Proposed or committed by a governing or other organizing body; 2) Designated by law or other authoritative rulemaking; 3) Implemented, with activated changes in management; and 4) Actively managed, with ongoing monitoring and adaptive management.

The second element describes the level of protection (LEVEL) from abatable extractive and destructive activities within an MPA (or MPA zone). This is important because MPA outcomes

depend on the level of protection. LEVEL is based on allowed activities: 1) Fully protected – no extractive or destructive activities; 2) Highly protected – minimal extractive or destructive activities; 3) Lightly protected – moderate extractive or destructive activities; and 4) Minimally protected – activities with high total impact, although still an MPA by IUCN criteria.

5 The final two elements focus on the processes used to create and maintain an MPA and the results from that MPA. To be effective, an MPA should be established and sustained through the enabling conditions (CONDITIONS) that describe the processes, principles, and considerations for effective MPA planning, design, governance, and management.


10 Finally, this guide integrates ecological and social sciences to identify the likely ecological and social outcomes (OUTCOMES) of an MPA (e.g., for biodiversity, fisheries yield, higher incomes, or climate resilience). OUTCOMES depend directly upon STAGE, LEVEL, and CONDITIONS.

Outlook

15 This guide minimizes confusion and enables smart planning, design, and evaluation of new or existing MPAs by informing decisions about scientific, societal, and policy priorities. For example, if the MPA goal is to restore and protect biodiverse and healthy ecosystems, this guide points to fully or highly protected areas as having the greatest likelihood of success once the MPA is implemented or actively managed, as long as enabling CONDITIONS are met. The guide also provides a blueprint to assess and monitor progress on international conservation

20 targets. Different types of MPAs make it confusing to interpret a single percent area that is “protected”, necessitating the concurrent assessment of MPA quality enabled by this guide. Our synthesis of existing evidence also identifies research priorities, including improved

understanding of MPAs' effectiveness across LEVEL of protection for climate mitigation and adaptation, social change, and comprehensive marine spatial planning.



5

Summary Figure. The level of protection, and therefore the effectiveness of marine protected areas (MPAs), will greatly influence the future state of the ocean. Past ocean ecosystems were abundant and diverse in species and habitats. Over time, expanded and intensified human activities depleted and disrupted ocean ecosystems and reduced their services. MPAs, in conjunction with climate mitigation strategies and more sustainable uses of the ocean, can conserve and restore biodiversity and the resilient ecosystems needed for human well-being. Different levels of protection will result in different outcomes, as shown, if enabling conditions are satisfied.

10

15

Main Text:

Protected Areas Play a Vital Role in the Ocean

Marine Protected Areas (MPAs) are one of many tools that policy-makers, managers, and communities use to stem the loss of biodiversity, disruption of ocean ecosystems, and the decline of the many benefits provided to people by healthy ocean ecosystems (1, 2). Although most of the ocean used to be a de facto MPA due to limited access, technology has enabled exploitation of almost all of the ocean (3). In addition, although there are numerous examples of successful traditional resource management, customary marine governance including the use of closed areas has been eroded in many countries as a result of processes such as colonization and market expansion (4). Because degradation, pollution, and exploitation have significantly impacted the open ocean, the coast, and adjoining lands (1), integrated efforts are urgently needed to make extractive uses sustainable, minimize impact of destructive activities, and expand effective protection of species, habitats, and ecosystem functioning (5, 6).

MPAs by definition prioritize the conservation of nature (IUCN; 7) and are the primary area-based tool for marine biodiversity conservation. In this review we focus only on MPAs, due to their prevalence and extensive scientific underpinnings based on decades of tracking and evaluation (8, 9). Other area-based management tools where biodiversity conservation is not the primary goal are not MPAs, although they may confer some conservation benefit. For example, Locally Managed Marine Areas (e.g., 10) or Fisheries Management Areas (11) have different management priorities. If the protection they provide effectively conserves biodiversity, they may qualify as Other Effective area-based Conservation Measures (OECMs) (12, 13).

International governing bodies have set global targets for MPAs and OECMs, e.g., to protect 10% of the ocean by 2020 (Convention on Biological Diversity's Aichi Target 11, 14, United

Nations Sustainable Development Goal 14.5, 15). Calls are increasing for the more ambitious target of effectively protecting at least 30% of the ocean by 2030 (16, 17).

5 However, confusion and disagreements pervade many discussions of MPAs and detract from conservation efforts. Quantifying how much of the ocean's biodiversity is effectively protected is challenging. Significant discrepancies exist over what 'protected' means, when to 'count' an area as protected, and which types of MPAs achieve the intended conservation goals (8, 9, 18, 19). At present, global databases document that a relatively small proportion of the ocean is protected in MPAs. Specifically, at the time of this writing 7.7% of the ocean is self-reported by countries as
10 existing in some type of designated MPA (20), but only 6.4% is in MPAs that have been implemented, with likely much less actively managed (21). Importantly, not all of the tallied areas in those percentages meet the IUCN definition of an MPA (7). The race to simply protect a certain percentage of the ocean could detract from the importance of MPA quality, leading to
15 perverse outcomes from establishing MPAs that are insufficiently protected or not adequately designed to achieve conservation goals (22).

Removing Confusion around Marine Protected Areas

We posit that much of this confusion can be resolved by addressing three critical questions.

1. What does 'protected' mean for biodiversity conservation? Even under the IUCN definition,
20 the 'protected' in 'marine protected area' encompasses numerous levels of protection with an almost endless variety and combination of activities that are allowed or not allowed, and consequently, lead to a wide range of impacts on biodiversity (8, 23). As a result, the ecological and social outcomes expected from MPAs, or a zone within an MPA, vary widely (e.g., 18).

Clarifying why MPAs differ from one another and which types will deliver specific desired outcomes is essential to help evaluate whether any given MPA is set up with the appropriate protection to achieve its aims.

2. When should an MPA ‘count’ as effectively protected? There are many steps in the process to create an MPA. Global tallies differ from one another in part because they use different criteria to ‘count’ MPAs (e.g., 20, 21), e.g., when it is proclaimed in law vs. when it is implemented in the water. This disparity becomes problematic when some MPAs are counted as achievements towards global targets but no real protection is in place in the water (24). There is a need to track all stages of MPAs and clarify that biodiversity protection is not expected to begin until the MPA rules and regulations are in place and active.

3. What is needed to achieve effective ocean protection? To prevent overestimation of how much ocean is actually protected (9, 19, 25, 26), knowledge of the total MPA coverage across different levels of protection is needed at the global scale. This requires assessment of the number, area, and impact of MPAs to ensure these are sufficient to achieve local, national, or international goals for healthy, productive, and resilient ocean ecosystems that support biodiversity and sustainable use (27).

A New Framework to Understand Protected Areas in the Ocean

With input from diverse global collaborators, we reviewed MPA science and its implications for global biodiversity conservation targets to develop a multidisciplinary, collaborative scientific synthesis that addresses the above three questions. We present our findings as a new framework called The MPA Guide. This guide organizes MPAs according to stage of establishment (STAGE) and level of protection (LEVEL), defined below. We then link these MPA types to

measured outcomes (OUTCOMES), based on the enabling social and ecological conditions (CONDITIONS) that research shows are key to an MPA successfully achieving its goals.

This guide strategically complements the IUCN Protected Area Categories, an existing
5 framework that categorizes areas by their management objectives and governance types (IUCN; 28), not STAGE or LEVEL of protection. Together, the MPA Guide and the IUCN Categories provide a comprehensive picture of an MPA. This guide helps consolidate and advance the reporting framework of UNEP-World Conservation Monitoring Centre's (WCMC) World Database on Protected Areas (WDPA), and the IUCN MPA Standards (7), which summarize and
10 distill approved motions by the global conservation community in past World Conservation Congresses. As long as an MPA (or zone within a multi-zone MPA) meets the IUCN definition of an MPA (7), it will fit into one STAGE of establishment and one LEVEL of protection at any point in time.

15 **Stage of establishment (STAGE) and when to 'count' an MPA**

MPA establishment generally occurs as a series of steps by governing or other authorities based on their local and national context. This guide specifies minimum criteria for an MPA to achieve each STAGE and provides guidelines for best practices (STAGES Expanded Guidance; Figure S1). In some cases, it may take several years between an announcement of intent to create an
20 MPA to the time when in situ protection and management occurs. In other situations, an MPA may be designated and implemented simultaneously if the announcement has legal authority and a management plan. Below we describe each STAGE and provide examples.

Proposed/Committed: The intent to create an MPA is made public. An MPA must be announced in some formal (though non-binding) manner via a statement by a government, community, conservation organization, or other organizing group; for example via an international meeting, a press release, or online. The MPA site must be identified, ideally with clear goals and informed by stakeholder and rights-holder participation, and Indigenous or other local and scientific knowledge of the social-ecological context. At the time of this writing, two examples of proposed/committed MPAs are in the East Antarctic (29) and in the Weddell Sea (30), where potential MPAs are currently under consideration by the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR).

Designated: The MPA is established or recognized through legal means or other authoritative rulemaking. A designated MPA must satisfy three minimum criteria: (1) defined boundaries, (2) legal gazetting or equivalent Indigenous/traditional authorization or customary recognition, and (3) clearly stated goals and process to define allowed uses and associated regulations or rules to control impact. MPA boundaries (including zones within the MPA) are ideally published, unambiguous, and known to local users. A designated MPA should have a database ID number in the WDPA signifying official recognition of the MPA. The MPA should be long-term; e.g., it should not have a sunset clause or review process that allows for rescinding protection in fewer than 25 years (7). As an example, Seychelles recently legally designated 30% of its ocean territory as an MPA network, which is currently in the process of being implemented (31).

It is important to note that MPAs that are proposed/committed or designated are not yet implemented with changes in activities and thus will not accrue biodiversity conservation

benefits. Protection does not begin until implementation. MPAs that are designated for an extended period of time without being implemented are often referred to as ‘paper parks’. These situations may reflect a lack of capacity and support (24).

5 Implemented: The MPA has transitioned from existence ‘on paper’ to being operational ‘in the water’ with management plans activated. Biodiversity conservation benefits begin to accrue at this stage, not before. Resource users are aware of the rules, and mechanisms to promote compliance and enforcement exist. Plans for regulating MPA activities are in place. Stakeholders are engaged, users are aware of regulations, financial and human resource
10 management systems are established, and performance measures are part of a plan to evaluate and monitor the MPA. Ideally, governance and administrative structures for management, implementation, and sustainable financing are specified (e.g., in management plans). Zones and their goals should be described, if applicable (e.g., 32). A management body should exist to implement and review plans. For example, Niue’s Moana Mahu MPA is implemented (33) and
15 includes 40% of the EEZ as fully protected with enforcement activities underway, partnerships in place, and ongoing stakeholder engagement.

Actively Managed: MPA management is ongoing, including monitoring, periodic review, and adjustments made as needed to achieve biodiversity conservation and other ecological and social
20 goals. All necessary MPA management activities for sustained functioning and achievement of goals continue. The MPA management authority documents, monitors, and evaluates MPA outcomes. Adaptive management will lead to adjustments in plans and activities as needed to ensure good compliance, stakeholder and rights-holder collaboration, and achievement of MPA

goals. Comprehensive systems exist to evaluate actively managed MPAs, such as the IUCN Green List (34) and the Marine Conservation Institute's Blue Parks Program (35). Periodic reviews of actively managed areas are based on evaluations of MPA management function such as sustainable financing, staffing, and outreach as well as data collected frequently inside and outside the MPA. These should involve a social-ecological systems approach (36), and ideally the participation of local communities and stakeholders (37). For example, the California network of MPAs established by the Marine Life Protection Act is actively managed and undergoes a systematic and comprehensive periodic five-year review using monitoring data to evaluate current management and inform future decisions (38).

Level of protection (LEVEL) for biodiversity conservation

By IUCN's definition, an MPA's primary goal is the conservation of nature (7). Thus, this guide focuses on evaluating protection based on the biodiversity outcomes that different activities at different scales are expected to produce.

Extensive peer-reviewed research shows that MPAs, or specific MPA zones, effectively protect biodiversity if they adequately prohibit extractive and destructive uses, (e.g., 39–41, 42; for a list of others see LEVELS Expanded Guidance, Fig. S1) and if key factors for positive desired outcomes are in place (i.e., CONDITIONS, see below). It is possible to conserve biodiversity while also balancing sustainable uses (43) – assuming full compliance with rules, some extractive and destructive activities may be allowed in an MPA, albeit with conservation outcomes that are likely more limited (e.g., 18, 42, 44).

This MPA Guide describes four LEVELS of protection based on the impact of allowed activities. It incorporates guidance from the Regulations-Based Classification System for MPAs (23) and IUCN’s guidelines (7). Impact is determined via activity type, intensity, scale, duration, and frequency relative to biodiversity conservation goals, and is described as ‘none’, ‘low’, ‘moderate’, ‘high/large’, or ‘incompatible with biodiversity conservation’ (Fig. 1; Figure S2). Using this impact scale, a LEVEL of protection can be assigned for any given MPA or zone regardless of location, species, or circumstances. Impacts of certain activities may scale differently considering specific features of an MPA or zone, such as size; for example, distribution of an activity across areas of different sizes may render it high impact in a smaller MPA but moderate impact in a larger MPA. Incompatible activities include industrial extraction such as industrial fishing (e.g., vessels > 12m using towed/dragged gears), oil and gas exploration, mining, or other extremely impactful activities such as fishing with dynamite or poison (LEVELS Expanded Guidance, Figure S1). Any activity that may be conducted for scientific research purposes in an MPA or zone is subject to the review and approval of the MPA management authority based on its impact.

Fully Protected: No extractive or destructive activities are allowed; all abatable impacts are minimized. Minimizing impacts requires attention to the scale of the protected area and the scale of the activity. MPAs cannot abate or prevent some impacts (e.g., climate change, coastal urbanization, pollution), although in certain circumstances they can enhance ecosystem resistance and resilience (i.e., both the ability of the ecosystem to resist impacts of disturbance and to return to a healthy state following disturbance) to some of these threats (45). The meaning of other, similar terms such as ‘strong’ or ‘strict’ protection, ‘marine reserves’, or ‘no-take’ areas

varies considerably from user to user (46). We use and clearly define the term ‘fully protected’ because it encompasses more than just extractive activities and emphasizes the positive intent of the action (compared to ‘no-take’, which emphasizes what is prohibited). Non-extractive low-impact tourism or low-impact cultural activities may be compatible with fully protected areas, provided collective impact is low (Fig. 1; Fig. S2). Potentially impactful activities such as aquaculture are only allowed for restoration purposes and not extraction. Examples include small-scale, decades-old community co-managed MPAs in the Philippines (47), large-scale MPAs such as the Palau National Marine Sanctuary (which covers 80% of the country’s EEZ, 48), or zones within multi-zone MPAs (19).

Highly Protected: Only light extractive activities with low total impact are allowed, with all other abatable impacts minimized. Some allow a small amount of subsistence or small-scale fishing with minimal impact, depending on the number of fishers and gear types (e.g., use by few fishers of highly selective gear such as hand lines or collection by freedivers may be compatible with highly protected status; five or fewer gears, 23). Allowed activities include low-impact tourism and low-density, unfed aquaculture. Highly protected areas may allow low-impact cultural and traditional activities such as sustainable fishing by Indigenous communities (e.g., 49), which are supported by clear property rights affording local stakeholders and rights-holders the authority to govern areas, including restricting exploitation by non-local actors (50). The 2016 expansion zone of the US’s Papahānaumokuākea Marine National Monument, which allows only low-frequency and low-impact activities, is highly protected (51).

Lightly Protected: Some protection of biodiversity exists but moderate to significant extraction and other impacts are allowed. These MPAs can achieve some protection of biodiversity for certain species or habitats, but the number and impacts of activities allowed are greater than for highly protected areas. A larger number of fishing gears might be used (10 or fewer, 23), or fishing occurs with less selective gear types (e.g., gill, trammel, or small-scale drift nets).

Tourism could have moderate impacts on habitats and species, such as damage caused by high intensity recreational diving. Aquaculture may occur via semi-intensive, unfed methods or small-scale and low-density fed methods. The vast majority of MPAs worldwide are lightly protected or minimally protected (9, 19, 19, 21) and often attempt to balance biodiversity conservation goals with resource use and development goals. For example, Habitat Protection Zones in Australia's Great Barrier Reef Marine Park are lightly protected since they allow multiple types of fishing (52).

Minimally Protected: Extensive extraction and other impacts are allowed, but the site still provides some conservation benefit in the area. Extensive extraction and other impacts occur in a minimally protected area, but the area still achieves sufficient biodiversity conservation to satisfy the IUCN definition of an MPA. For example, the area must not allow industrial fishing (53).

Nonetheless, minimally protected areas are unlikely to deliver substantial biodiversity conservation benefits for nature and people. A recent analysis showed that more than 10 fishing gear types used in an MPA either recreationally or commercially likely leads to large-scale impacts (18, 23). Minimally protected MPAs often allow many or high-impact gear types for extraction, and may include medium- to high-density aquaculture, and/or large-impact anchoring

or infrastructure. For example, the US Hawaiian Islands Humpback Whale National Marine Sanctuary is minimally protected since it allows extensive fishing and anchoring (54).

LEVELS of protection are designed to harmonize with and build upon, but not replicate, the information provided by the IUCN Protected Area Categories. For this reason, LEVEL in The MPA Guide does not map directly to an IUCN Category. The zones in the Great Barrier Reef (GBR) Marine Park provide useful examples. Some fully protected zones correspond with IUCN Category Ia; for example, the GBR Preservation Zones are “no-go” areas with all extractive and destructive activities prohibited. Other Category 1a areas, such as the GBR Scientific Research Zones, are highly protected with low-impact extractive research and traditional resource use allowed. The GBR Conservation Park and Buffer zones are both IUCN Category IV MPAs, but Buffer Zones are highly protected whereas Conservation Park zones are lightly protected due to the range of fishing gears allowed.

Within The MPA Guide, LEVEL of protection for any particular MPA, or zone within a multi-zone MPA, depends on activity types that are explicitly permitted or prohibited by the MPA rules, or based on overlapping regulations for the surrounding area (Fig. 1). Some activity types or impact levels are not explicitly stated in MPA rules and regulations, often because they are not within the management jurisdiction of the MPA authority. In these circumstances, knowledge of whether or not that activity occurs may be used. Since it is the current activities that influence the degree to which an MPA is protecting biodiversity at a given point in time, the assessment of MPA LEVEL should reflect activities actually occurring in the site at the time of reporting, whether or not they are explicitly stated in the management plans.

Seven main types of activities determine LEVEL: (1) Mining/Oil and Gas Extraction, (2) Dredging and Dumping, (3) Anchoring, (4) Infrastructure, (5) Aquaculture, (6) Fishing, whether it is subsistence, professional, or recreational fishing; this activity encompasses extraction of wild fish and other marine species and includes gleaning, and (7) Non-extractive activities, including recreational, traditional, and cultural (LEVELS Expanded Guidance, Figure S1). The compatibility of each activity with conservation goals was evaluated through multiple, iterative workshops using peer-reviewed literature, scientific judgment, expert opinion, and IUCN resolutions and protected area guidance (e.g., 23).

This guide does not include every possible activity but provides best practices wherever possible. For example, shipping is not explicitly addressed, because the right of innocent passage is mandated under international law and regulated by International Maritime Organization treaties. As a result, it is challenging for an MPA managing authority to restrict shipping movement. Nonetheless, it is recommended that ships with dangerous goods or toxic anti-fouling chemicals not transit MPAs, and that shipping activity be restricted to shipping lanes to minimize noise pollution and other negative impacts such as collisions with marine life (55) (LEVELS Expanded Guidance, Figure S1). Guidance is intended to evolve with new knowledge, activities, and technology. Emerging threats due to electromagnetic fields, excessive or persistent noise, high energy active sonar, or other technologies not explicitly addressed here are subject to the burden of proof (e.g. 55, 56), meaning management bodies should receive evidence of their expected impacts before allowing their use, and they should monitor to assess and actively manage their actual impacts. Impacts should not exceed those associated with a given LEVEL.

Enabling conditions (CONDITIONS) for Effective MPAs

MPAs cannot achieve their goals unless key CONDITIONS are in place. These are the
5 conditions by which an MPA is effectively planned, designed, implemented, governed, and
managed to achieve desired ecological outcomes and the direct and indirect human well-being
outcomes that result. These CONDITIONS may vary in their importance during the process of
achieving each of the four STAGES (e.g., 57, 58, 59; Table 1), but aspects of each apply when
moving from proposed/committed to designated (e.g., 60–62) to implemented (e.g., 34, 63) and
10 actively managed (e.g., 34, 63, 64). They will also vary based on local challenges, opportunities,
and resources, requiring engagement in a prioritization process that is specific to each context.

The beneficial governance practices that these CONDITIONS span—such as inclusivity,
transparency, and accountability—increase legitimacy, ownership, support, and overall
15 effectiveness of conservation (65, 66). These practices give voice to those who often
disproportionately bear the costs of degradation or conservation and identify livelihood support
or other strategies to help mitigate impacts and increase benefits. For example, MPAs in the
Mediterranean received greater support from community members with transparent decision-
making that recognized and strengthened the rights of local resource users (66). MPAs that
20 exclude resource users from decision-making and ignore their rights and livelihood dependencies
can erode their well-being and undermine compliance. In Mnazi-Bay, Tanzania, exclusion of
resource users from the MPA process led to negative social outcomes, including increased food
insecurity, violent conflict, and lower educational outcomes (67).

Linking MPA goals to measurable outcomes: Achieving ocean protection

We integrate peer-reviewed scientific literature and expert working group products to link STAGE and LEVEL with the ecological and social OUTCOMES expected from different types of MPAs. If biodiversity is conserved, an MPA would be considered successful at meeting its primary goal. However, this does not preclude other outcomes from also occurring and producing benefits, including those for human well-being. Once an MPA is implemented with CONDITIONS in place, it can lead to interrelated ecological and social outcomes based on LEVEL of protection.

Ecological Outcomes of MPAs

Thousands of MPA studies document the ecological effects of MPAs across almost all ocean regions and seas, demonstrating that MPAs are an effective tool to conserve biodiversity and improve ecosystem functioning (Table 2; expanded references in Table S1). Outside their borders they can also enhance fish stocks through egg and larval export and spillover of juveniles and adults to areas outside the MPA boundaries (68). Interconnected networks of MPAs are expected to deliver scaled benefits (69). Highly mobile species and those with very large home ranges may receive lower benefit levels from MPA protection than more sedentary species, unless MPAs are larger or dynamic with mobile boundaries (70, 71) or they protect critical life stages [e.g., spawning aggregations, nursery or feeding grounds or migration bottlenecks (72, 73)]. Long-ranging species require well-designed MPA networks and effective management outside MPAs (74).

Research is often biased towards ecological and fisheries responses to MPA protection (e.g., 75) since these are related to the biodiversity conservation goals of MPAs (7) and the main impact

that MPAs abate (fishing). However, other benefits are possible (Table 2; Table S1). Water quality can improve if MPAs restore and recover vegetated habitats and filter-feeding bottom communities (76). While direct evidence is limited for MPAs, climate change mitigation, adaptation, and resilience could help offset climate-change related impacts (45, 77, 78).

5 Protecting ‘blue carbon’ habitats can preserve their ability to provide carbon sequestration and coastal protection, particularly if supplemented by restoration (79).

In well-managed MPAs, ecological benefits relative to surrounding unprotected areas are more prominent where species have previously been depleted, particularly by factors that can be managed or excluded. Significant prior habitat damage or human impacts outside MPAs can
10 slow recovery (65). In more intact areas, protection can guard against future losses (80). Threats that cannot be abated by protection may reduce benefits, especially in the short term. However, protection may partially mitigate some of these impacts by protecting functioning ecosystems, boosting resilience, and hastening recovery (45, 81, 82). Extractive activities displaced by protection may lead to impacts outside MPAs, underscoring the need for integrating MPAs into
15 comprehensive marine spatial planning to ensure damaging activities are not displaced onto more sensitive habitats or the ranges of more vulnerable species.

While some benefits occur quickly following protection, others can take decades. Species respond to protection at different rates depending on factors such as life history characteristics, behavior, depletion at the time of protection (e.g., for fished species), and other human impacts
20 (40). Early results often include increases in species already common within the MPA, but as time passes, such benefits also include increases in rare and vulnerable species, re-establishment of natural population age structure (especially for long-lived species), and recovery of degraded

structural ecosystem elements and habitats (83). The Outcomes in Table 2 assume that adequate protection has been in place long enough for effects to develop.

Recovery is more likely, faster, and more complete at the higher LEVELS of protection: positive ecological outcomes are more substantial and less variable in fully and highly than lightly and minimally protected areas (Table 2) (e.g., 18, 23, 84) with greater potential for ecosystem restoration when areas are fully protected (85). In protection levels with more activities occurring, management often addresses competing or conflicting uses of an area and may advantage certain groups of users (e.g., small-scale or recreational over larger commercial fishers). Decisions about the appropriate protection LEVEL will depend on conservation and management goals, social context, and CONDITIONS, which enable OUTCOMES (see Table 1). For example, poorly designed, managed, and resourced MPAs, with low compliance and staff, will deliver fewer benefits (57, 86), and a highly protected area could produce better outcomes than a fully protected area if it has stronger enabling conditions.

Social Outcomes of MPAs

MPAs can directly and indirectly affect all aspects of human well-being (i.e., social, health, culture, economic, and governance, 87) for different rights-holders and stakeholders (e.g., Indigenous peoples, fishers, tourism operators, coastal residents). When key CONDITIONS are in place, positive benefits of MPAs can be enhanced and negative impacts minimized. A recent comprehensive review found that about half of all documented human well-being outcomes of MPAs were positive and about one-third were negative, with the remaining showing no change or change that was not attributable as positive or negative (88). Common positive outcomes were

community involvement, increased catch per unit of fishing effort (CPUE), and higher income, whereas negative outcomes commonly manifested through increasing costs of activities (i.e., fishing) and conflict (89). Both positive and negative impacts can occur at the same time (e.g., 90, 91). Four MPAs in Indonesia had positive effects on material wealth and scientific
5 environmental knowledge but negative effects on perceived well-being, fish catch, and marine resource control (92).

Direct effects of MPAs on human well-being can be immediate due to changes in access or decision-making (93). For example, discussions about whether to have MPAs, where to place them, and what management measures to include can directly affect levels of conflict,
10 perceptions of procedural fairness, access to resources and incomes, and sense of agency in resource management, either negatively or positively (94, 95). Indirect effects also occur via subsequent management actions and ecosystem changes, including altered catches, CPUE, and income from resource extraction or non-extractive activities (65, 96, 97). Such effects are the most common positive MPA outcomes for human well-being and they may take time based on
15 ecological recovery rates. Both direct and indirect effects may shift over time (e.g., 92); negative impacts on the fish catch for certain commercial fishers increased over four years in two MPAs in the Gulf of Mexico (98).

The effects of MPAs can vary significantly across and within stakeholder groups depending on previous rights, dependence, and uses (99, 100) and across the broader social-ecological context
20 (100, 101). Differential MPA effects have been examined most commonly for fishers, particularly by fishing method (e.g., commercial, artisanal, using different gear types) (88, 90). This variability can depend on level of resource dependency (e.g., dietary dependency, livelihood

diversity, 102, 103), ability to adapt to changes (e.g., fishing areas, jobs, 94, 103, 104), involvement in MPA establishment processes (e.g., 65), and other socio-cultural characteristics that structure society (e.g. age, gender, ethnicity, 92).

The direction and strength of MPA impacts on different societal groups can also change temporally (e.g., 100) and can affect power dynamics within coastal communities as some members of the community benefit and others are excluded (e.g., 99). Individuals from marginalized groups with high resource dependency and low adaptive capacity often bear disproportionate costs (67, 105), particularly when excluded from decision-making processes (104). Alternatively, if protection strengthens local community property rights and excludes outside users, and/or provides economic benefits (e.g., from tourism), an MPA may benefit local communities. Achieving more positive outcomes requires attention to the MPA goals and the CONDITIONS during all STAGES to support stakeholders and rights-holders, the contribution of marine ecosystems to their wellbeing including livelihoods, and long-term MPA functioning (Table 1).

Protection LEVEL influences all indirect social impacts but only some direct impacts. A higher LEVEL of protection can generate greater recovery of socially, culturally, and economically important species or habitats, especially over the longer term (an indirect impact). Such protection could also increase the likelihood of conflict resulting from fishers being displaced, but may not impact other direct effects (e.g., empowerment in decision-making). In some cases, lightly or minimally protected areas may meet the needs of the local community, at least in the short term. Overall, when key CONDITIONS are met (e.g., long-term protection; high levels of

compliance, Table 1), fully and highly protected areas are associated with more positive outcomes (88), aligning with the positive outcomes found in ecological studies (e.g., 106).

Moving forward with clarity and transparency

5 Existing international targets highlight the key role of MPAs in conserving biodiversity and supporting a sustainable ocean economy—the blue economy. Achieving these goals has become even more important due to escalating threats to ocean biodiversity and ecosystem functioning (1, 5) and the disproportionately large impacts of the COVID-19 pandemic on near-shore communities (107). We argue that three key actions using this framework would strengthen
10 MPA understanding and use at local, national, and global scales.

1. Incorporate STAGE of establishment and LEVEL of protection into global reporting on progress towards international targets. The WDPA (20) reports on progress towards the 10% Aichi Target 11 and will report on the subsequent Post-2020 Target yet to be adopted by the Convention on Biological Diversity (CBD) at the time of this writing. The WDPA is mandated to
15 report designated MPAs (along with implemented and actively managed areas) in their tally of total protected area worldwide and does not track LEVEL of protection. Tracking STAGE and LEVEL, for example by using them as indicators of progress towards the future CBD Target, provides a full matrix evaluation (Fig. 2) of MPA quality and thereby moves global assessment of protection beyond a single percentage metric. This matrix approach may be similarly useful
20 for OECMs and terrestrial protected areas.

2. Use this framework to identify immediate opportunities to strengthen existing or to create new MPAs. An urgent need to recover ocean health and concomitant benefits to people means that high-priority and ample pay-off opportunities exist to create new MPAs and strengthen the level of protection and compliance for existing MPAs. Doing so was one of the Ocean Panel's five immediate action opportunities for COVID-19 recovery (107). The MPA Guide can help local, regional, and national bodies develop, implement, and manage new and existing MPAs.

Recognizing the distinct STAGE of an MPA can help MPA agencies and those working in civil society to progress an MPA to the next step, for example by thinking through and addressing capacity constraints such as lack of financial, social, and scientific capital. When developing a new MPA, decision makers and managers can also assess different protection LEVELS and their expected outcomes when deciding which activities to allow. A review of the rules and regulations in existing MPAs and how they map to protection LEVELS can help to determine whether these activities are consistent with desired ecological and social outcomes. Indonesia recently underwent an evaluation of their MPAs by STAGE and LEVEL, highlighting the impressive resource and capacity the country has invested toward active management, while also identifying MPAs that may require increased protection to achieve their goals (108). At a regional level, we can track how much of an ecosystem or habitat type is in each LEVEL of protection and identify sites in need of increased protection for biodiversity (19). Identifying how much ocean is still in proposed/committed or designated MPAs shows what has been promised for protection but is still in need of further action to implement (e.g., 109).

3. Develop research agendas to link MPA protection LEVEL, CONDITIONS, and OUTCOMES.

Although some types of MPAs have been studied for decades, two paths forward are required in a new era of MPA research. First, datasets should be organized around the protection provided by different MPAs in different LEVELS. Most existing ecological research lumps MPAs into fully protected areas and ‘partially protected’ areas, the latter of which combines highly, lightly, and minimally protected (e.g., 42, 44, 84). Combining these levels limits our ability to understand and predict OUTCOMES (e.g., Table 2) and to assess trade-offs to biodiversity conservation and trade-offs among different stakeholder groups. Explicit research across these three levels of protection is now possible using this framework.

More research is also needed to better understand MPA effects on specific social outcomes, across different societal groups (e.g., gender, age, ethnic groups), and over time (88, 90).

Research should expand geographically to assess how MPAs affect the multiple dimensions of human well-being in diverse contexts (36, 88, 90) and should employ an impact-evaluation lens, including rigorous counterfactual study designs (see 37, 92) in qualitative as well as quantitative studies (88). Further research is also needed to better understand the CONDITIONS as they relate to an MPA’s STAGE and LEVEL of protection, and the specific aspects of MPA planning, governance, and management that produce positive or negative outcomes for equity (110) and other dimensions of human well-being.

Conclusion

The stakes have never been higher for connecting MPA science to policy and action.

Development of the new CBD and other MPA goals and targets requires improved clarity and harmonization to be effective from local to global scales. Use of The MPA Guide would shift the

conversation from arguments about what MPAs can deliver, to answering questions such as ‘What level of protection is needed for an MPA to produce the desired outcomes for biodiversity and human well-being?’ and ‘What is the global tally of MPAs by stage of establishment and level of protection, and what does this tell us about progress towards ocean conservation goals?’

5 This scientific synthesis and guide offers a novel framework, language, and detailed guidance towards doing so.

10 **References and Notes:**

1. IPBES, “Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services” (E. S. Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors), IPBES secretariat, Bonn, Germany, 2019).
- 15 2. United Nations, Ed., *The First Global Integrated Marine Assessment: World Ocean Assessment I* (Cambridge University Press, Cambridge, 2017; <https://www.cambridge.org/core/books/first-global-integrated-marine-assessment/574E6BC58D440BD95B9DECFBC42070FB>).
3. J. Lubchenco, S. D. Gaines, A new narrative for the ocean. *Science*. **364**, 911–911 (2019).
- 20 4. A. M. Friedlander, Marine conservation in Oceania: Past, present, and future. *Mar. Poll. Bull.* **135**, 139–149 (2018).
5. IPCC, “IPCC Special Report on the Ocean and Cryosphere in a Changing Climate” ([H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (editors)]., Geneva, Switzerland, 2019).
- 25 6. E. Sala, J. Mayorga, D. Bradley, R. B. Cabral, T. B. Atwood, A. Auber, W. Cheung, C. Costello, F. Ferretti, A. M. Friedlander, S. D. Gaines, C. Garilao, W. Goodell, B. S. Halpern, A. Hinson, K. Kaschner, K. Kesner-Reyes, F. Leprieur, J. McGowan, L. E. Morgan, D. Mouillot, J. Palacios-Abrantes, H. P. Possingham, K. D. Rechberger, B. Worm, J. Lubchenco, Protecting the global ocean for biodiversity, food and climate. *Nature*, 1–6 (2021).
- 30

7. IUCN and WCPA, “Applying IUCN’s Global Conservation Standards to Marine Protected Areas (MPAs). Delivering effective conservation action through MPAs, to secure ocean health & sustainable development. Version 1.0.” (Gland, Switzerland, 2018).
- 5 8. J. Lubchenco, K. Grorud-Colvert, Making waves: The science and politics of ocean protection. *Science*. **350**, 382–383 (2015).
9. E. Sala, J. Lubchenco, K. Grorud-Colvert, C. Novelli, C. Roberts, U. R. Sumaila, Assessing real progress towards effective ocean protection. *Mar. Pol.* **91**, 11–13 (2018).
- 10 10. J. A. Kawaka, M. A. Samoilys, M. Murunga, J. Church, C. Abunge, G. W. Maina, Developing locally managed marine areas: Lessons learnt from Kenya. *Ocean Coastal Manage.* **135**, 1–10 (2017).
11. D. Vilas, M. Coll, X. Corrales, J. Steenbeek, C. Piroddi, D. Macias, A. Ligas, P. Sartor, J. Claudet, Current and potential contributions of the Gulf of Lion Fisheries Restricted Area to fisheries sustainability in the NW Mediterranean Sea. *Mar Pol.* **123**, 104296 (2020).
- 15 12. D. Laffoley, N. Dudley, H. Jonas, D. MacKinnon, K. MacKinnon, M. Hockings, S. Woodley, An introduction to ‘other effective area-based conservation measures’ under Aichi Target 11 of the Convention on Biological Diversity: Origin, interpretation and emerging ocean issues. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **27**, 130–137 (2017).
13. J. M. Reimer, R. Devillers, J. Claudet, Benefits and gaps in area-based management tools for the ocean Sustainable Development Goal. *Nat. Sustain.*, 1–9 (2020).
- 20 14. Woodley S, Bertsky B, Crawhall N, Dudley N, Londono JM, MacKinnon K, Redford K, Sandwith T, Meeting Aichi Target 11: what does success look like for Protected Area systems? *Parks*, 1 (2012).
15. E. S. Nocito, C. M. Brooks, A. L. Strong, Gazing at the crystal ball: predicting the future of marine protected areas through voluntary commitments. *Front. Mar. Sci.* **6** (2020), doi:10.3389/fmars.2019.00835.
- 25 16. B. C. O’Leary, M. Winther-Janson, J. M. Bainbridge, J. Aitken, J. P. Hawkins, C. M. Roberts, Effective coverage targets for ocean protection. *Cons. Lett.* **9**, 398–404 (2016).
17. K. R. Jones, C. J. Klein, H. S. Grantham, H. P. Possingham, B. S. Halpern, N. D. Burgess, S. H. M. Butchart, J. G. Robinson, N. Kingston, N. Bholá, J. E. M. Watson, Area requirements to safeguard Earth’s marine species. *One Earth.* **2**, 188–196 (2020).
- 30 18. M. Zupan, E. Fragkopoulou, J. Claudet, K. Erzini, B. H. e Costa, E. J. Gonçalves, Marine partially protected areas: drivers of ecological effectiveness. *Frontiers in Ecology and the Environment.* **16**, 381–387 (2018).
19. J. Claudet, C. Loiseau, M. Sostres, M. Zupan, Underprotected marine protected areas in a global biodiversity hotspot. *One Earth.* **2**, 380–384 (2020).
- 35

20. UNEP-WCMC, World Database on Protected Areas. *Protected Planet* (2020), (available at <https://www.protectedplanet.net/marine>).
21. Marine Conservation Institute, The Atlas of Marine Protection (2020), (available at <http://mpatlas.org/>).
- 5 22. M. D. Barnes, L. Glew, C. Wyborn, I. D. Craigie, Prevent perverse outcomes from global protected area policy. *Nat. Ecol. Evol.* **2**, 759–762 (2018).
23. B. Horta e Costa, J. Claudet, G. Franco, K. Erzini, A. Caro, E. J. Gonçalves, A regulation-based classification system for Marine Protected Areas (MPAs). *Marine Policy.* **72**, 192–198 (2016).
- 10 24. M. Pieraccini, S. Coppa, G. A. D. Lucia, Beyond marine paper parks? Regulation theory to assess and address environmental non-compliance. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **27**, 177–196 (2017).
25. B. Horta e Costa, J. M. dos S. Gonçalves, G. Franco, K. Erzini, R. Furtado, C. Mateus, E. Cadeireiro, E. J. Gonçalves, Categorizing ocean conservation targets to avoid a potential false sense of protection to society: Portugal as a case-study. *Mar. Pol.* **108**, 103553 (2019).
26. J. Claudet, C. Loiseau, A. Pebayle, Critical gaps in the protection of the second largest exclusive economic zone in the world. *Mar. Pol.* **124**, 104379 (2021).
27. CBD, “Decisions Adopted by the Conference of the Parties to the Convention on Biological Diversity at its Eighth Meeting (Decision VIII/15, Annex IV).,” *Convention on Biological Diversity* (Curitiba, Brazil, 2006).
28. “Guidelines for applying the IUCN protected area management categories to marine protected areas. Second edition.” (IUCN, Gland, Switzerland, 2019), (available at <https://www.iucn.org/content/guidelines-applying-iucn-protected-area-management-categories-marine-protected-areas-0>).
- 25 29. CCAMLR, “Proposal to establish an East Antarctic Marine Protected Area,” *CCAMLR-38/21*, (available at <https://www.ccamlr.org/en/ccamlr-38/21>).
30. CCAMLR, “Proposal to establish a Marine Protected Area across the Weddell Sea region (Phase 1),” *CCAMLR-38/23*, (available at <https://www.ccamlr.org/en/ccamlr-38/23>).
- 30 31. Speech by President Danny Faure on the occasion of 30% of Seychelles’ EEZ Designated as Marine Protected Area, (available at <https://www.statehouse.gov.sc/speeches/4786/speech-by-president-danny-faure-on-the-occasion-of-30-of-seychelles-eez-designated-as-marine-protected-area>).
- 35 32. J. C. Day, R. A. Kenchington, J. M. Tanzer, D. S. Cameron, Marine zoning revisited: How decades of zoning the Great Barrier Reef has evolved as an effective spatial planning

approach for marine ecosystem-based management. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **29**, 9–32 (2019).

33. Department of Agriculture, Forestry, and Fisheries, “Niue Moana Mahu Marine Protected Area Regulations 2020,” *No.2020/04*, (available at https://old.mpatlas.org/media/filer_public/bc/95/bc959065-13b7-42d7-97dd-507503fc4b01/reg_2020-04_niue_moana_mahu_marine_protected_area_regulations_1.pdf).
34. S. Wells, P. F. E. Addison, P. A. Bueno, M. Costantini, A. Fontaine, L. Germain, T. Lefebvre, L. Morgan, F. Staub, B. Wang, A. White, M. X. Zorrilla, Using the IUCN Green List of Protected and Conserved Areas to promote conservation impact through marine protected areas. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **26**, 24–44 (2016).
35. Marine Conservation Institute, BlueParks. *BlueParks* (2020), (available at <https://blueparks.org/>).
36. G. G. Gurney, E. S. Darling, S. D. Jupiter, S. Mangubhai, T. R. McClanahan, P. Lestari, S. Pardede, S. J. Campbell, M. Fox, W. Naisilisili, N. A. Muthiga, S. D’agata, K. E. Holmes, N. A. Rossi, Implementing a social-ecological systems framework for conservation monitoring: lessons from a multi-country coral reef program. *Biol. Cons.* **240**, 108298 (2019).
37. G. N. Ahmadia, L. Glew, M. Provost, D. Gill, N. I. Hidayat, S. Mangubhai, null Purwanto, H. E. Fox, Integrating impact evaluation in the design and implementation of monitoring marine protected areas. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370** (2015), doi:10.1098/rstb.2014.0275.
38. S. Murray, T. T. Hee, A rising tide: California’s ongoing commitment to monitoring, managing and enforcing its marine protected areas. *Ocean. Coast. Manag.* **182**, 104920 (2019).
39. S. E. Lester, B. S. Halpern, K. Grorud-Colvert, J. Lubchenco, B. I. Ruttenberg, S. D. Gaines, S. Airame, R. R. Warner, Biological effects within no-take marine reserves: a global synthesis. *Mar Ecol Prog Ser.* **384**, 33–46 (2009).
40. J. Claudet, C. W. Osenberg, L. Benedetti-Cecchi, P. Domenici, J.-A. García-Charton, Á. Pérez-Ruzafa, F. Badalamenti, J. Bayle-Sempere, A. Brito, F. Bulleri, J.-M. Culioli, M. Dimech, J. M. Falcón, I. Guala, M. Milazzo, J. Sánchez-Meca, P. J. Somerfield, B. Stobart, F. Vandeperre, C. Valle, S. Planes, Marine reserves: size and age do matter. *Ecol. Lett.* **11**, 481–489 (2008).
41. P. P. Molloy, I. B. McLean, I. M. Côté, Effects of marine reserve age on fish populations: a global meta-analysis. *J Appl Ecol.* **46**, 743–751 (2009).
42. S. Giakoumi, C. Scianna, J. Plass-Johnson, F. Micheli, K. Grorud-Colvert, P. Thiriet, J. Claudet, G. Di Carlo, A. Di Franco, S. D. Gaines, J. A. c García-Charton, J. Lubchenco, J.

Reimer, E. Sala, P. Guidetti, Ecological effects of full and partial protection in the crowded Mediterranean Sea: a regional meta-analysis. *Sci. Rep.* **7**, 8940 (2017).

43. M. Zupan, F. Bulleri, J. Evans, S. Frascchetti, P. Guidetti, A. Garcia-Rubies, M. Sostres, V. Asnaghi, A. Caro, S. Deudero, R. Goñi, G. Guarnieri, F. Guilhaumon, D. Kersting, A. Kokkali, C. Kruschel, V. Macic, L. Mangialajo, S. Mallol, E. Macpherson, A. Panucci, M. Radolovic, M. Ramdani, P. J. Schembri, A. Terlizzi, E. Villa, J. Claudet, How good is your marine protected area at curbing threats? *Biol. Cons.* **221**, 237–245 (2018).
44. M. Sciberras, S. R. Jenkins, R. Mant, M. J. Kaiser, S. J. Hawkins, A. S. Pullin, Evaluating the relative conservation value of fully and partially protected marine areas. *Fish. Fish.* **16**, 58–77 (2015).
45. C. M. Roberts, B. C. O’Leary, D. J. McCauley, P. M. Cury, C. M. Duarte, J. Lubchenco, D. Pauly, A. Sáenz-Arroyo, U. R. Sumaila, R. W. Wilson, B. Worm, J. C. Castilla, Marine reserves can mitigate and promote adaptation to climate change. *PNAS*, 201701262 (2017).
46. J. Claudet, France must impose strict levels of marine protection. *Nature.* **570**, 36–36 (2019).
47. A. C. Alcala, G. R. Russ, No-take marine reserves and reef fisheries management in the Philippines: a new people power revolution. *Ambio.* **35**, 245–254 (2006).
48. Palau International Coral Reef Center-PICRC, “Palau National Marine Sanctuary Act Overview,” (available at <https://picrc.org/picrcpage/wp-content/uploads/2019/12/PNMS-Act-Overview-2019AUG20.pdf>).
49. K. Kikiloi, A. M. Friedlander, ’Aulani Wilhelm, N. Lewis, K. Quiocho, W. ’Āila, S. Kaho’ohalahala, Papahānaumokuākea: Integrating culture in the design and management of one of the world’s largest marine protected areas. *Coast. Manage.* (2017) (available at <https://agris.fao.org/agris-search/search.do?recordID=US201800255948>).
50. M. B. Mascia, C. A. Claus, A property rights approach to understanding human displacement from protected areas: the case of marine protected areas. *Conserv. Biol.* **23**, 16–23 (2009).
51. National Oceanic and Atmospheric Administration, United States Fish and Wildlife Service, Hawai’i Department of Land and Natural Resources, “Papahānaumokuākea Marine National Monument Management Plan” (2008), (available at https://nmspapahanaumokuakea.blob.core.windows.net/papahanaumokuakea-prod/media/archive/management/mp/vol1_mmp08.pdf).
52. Great Barrier Reef Marine Park Authority, Interpreting zones, (available at <https://www.gbrmpa.gov.au/access-and-use/zoning/interpreting-zones>).

53. International Union for the Conservation of Nature, “Motion 066 - Guidance to identify industrial fishing incompatible with protected areas,” (available at <https://www.iucncongress2020.org/motion/066>).
54. Office of National Marine Sanctuaries, National Oceanic and Atmospheric Administration, United States Department of Commerce, State of Hawai‘i, Hawai‘i Department of Land and Natural Resources, “Hawai‘ian Islands Humpback Whale National Marine Sanctuary Management Plan” (2020), (available at <https://nmshawaiihumpbackwhale.blob.core.windows.net/hawaiihumpbackwhale-prod/media/docs/2020-hihwnms-management-plan.pdf>).
55. L. M. Petruny, A. J. Wright, C. E. Smith, Getting it right for the North Atlantic right whale (*Eubalaena glacialis*): A last opportunity for effective marine spatial planning? *Mar. Poll. Bull.* **85**, 24–32 (2014).
56. C. Bracciali, D. Campobello, C. Giacoma, G. Sarà, Effects of nautical traffic and noise on foraging patterns of Mediterranean damselfish (*Chromis chromis*). *PLOS ONE.* **7**, e40582 (2012).
57. D. A. Gill, M. B. Mascia, G. N. Ahmadi, L. Glew, S. E. Lester, M. Barnes, I. Craigie, E. S. Darling, C. M. Free, J. Geldmann, S. Holst, O. P. Jensen, A. T. White, X. Basurto, L. Coad, R. D. Gates, G. Guannel, P. J. Mumby, H. Thomas, S. Whitmee, S. Woodley, H. E. Fox, Capacity shortfalls hinder the performance of marine protected areas globally. *Nature.* **543**, 665–669 (2017).
58. M. Lockwood, Good governance for terrestrial protected areas: A framework, principles and performance outcomes. *Journal of Environmental Management.* **91**, 754–766 (2010).
59. N. J. Bennett, T. Satterfield, Environmental governance: A practical framework to guide design, evaluation, and analysis. *Conservation Letters.* **11**, e12600 (2018).
60. IUCN-WCPA, “Establishing marine protected area networks--Making it happen.” (IUCN World Commission on Protected Areas, National Oceanic and Atmospheric Administration, and The Nature Conservancy, Washington, D.C., 2008), p. 118.
61. R. L. Pressey, M. C. Bottrill, Approaches to landscape- and seascape-scale conservation planning: convergence, contrasts and challenges. *Oryx.* **43**, 464–475 (2009).
62. H. E. Fox, M. B. Mascia, X. Basurto, A. Costa, L. Glew, D. Heinemann, L. B. Karrer, S. E. Lester, A. V. Lombana, R. S. Pomeroy, C. A. Recchia, C. M. Roberts, J. N. Sanchirico, L. Pet-Soede, A. T. White, Reexamining the science of marine protected areas: linking knowledge to action. *Conserv Lett.* **5**, 1–10 (2012).
63. M. Hockings, S. Stolton, F. Leverington, N. Dudley, J. Courrau, *Evaluating effectiveness: a framework for assessing management effectiveness of protected areas* (IUCN, Gland, Switzerland, 2006; <https://portals.iucn.org/library/node/8932>), *Best Practice Protected Area Guidelines Series.*

64. B. Spergel, M. Moye, *Financing marine conservation a menu of options* (World Wildlife Fund, Center for Conservation Finance, Washington, D.C., 2004).
65. P. Guidetti, J. Claudet, Comanagement practices enhance fisheries in marine protected areas. *Conserv Biol* **24**, 312–318 (2010).
- 5 66. N. J. Bennett, A. D. Franco, A. Calò, E. Nethery, F. Niccolini, M. Milazzo, P. Guidetti, Local support for conservation is associated with perceptions of good governance, social impacts, and ecological effectiveness. *Conservation Letters*. **12**, e12640 (2019).
67. V. R. Kamat, “The ocean is our farm”: Marine conservation, food insecurity, and social suffering in southeastern Tanzania. *Hum. Organ.* **73**, 289 (2014).
- 10 68. H. B. Harrison, D. H. Williamson, R. D. Evans, G. R. Almany, S. R. Thorrold, G. R. Russ, K. A. Feldheim, L. van Herwerden, S. Planes, M. Srinivasan, M. L. Berumen, G. P. Jones, Larval export from marine reserves and the recruitment benefit for fish and fisheries. *Curr. Biol.* **22**, 1023–1028 (2012).
- 15 69. K. Grorud-Colvert, J. Claudet, B. N. Tissot, J. E. Caselle, M. H. Carr, J. C. Day, A. M. Friedlander, S. E. Lester, T. L. de Loma, D. Malone, W. J. Walsh, Marine protected area networks: Assessing whether the whole is greater than the sum of its parts. *PLOS ONE*. **9**, e102298 (2014).
- 20 70. A. Di Franco, J. G. Plass-Johnson, M. Di Lorenzo, B. Meola, J. Claudet, S. D. Gaines, J. A. García-Charton, S. Giakoumi, K. Grorud-Colvert, C. W. Hackradt, F. Micheli, P. Guidetti, Linking home ranges to protected area size: The case study of the Mediterranean Sea. *Biological Conservation*. **221**, 175–181 (2018).
71. S. M. Maxwell, K. M. Gjerde, M. G. Conners, L. B. Crowder, Mobile protected areas for biodiversity on the high seas. *Science*. **367**, 252–254 (2020).
- 25 72. B. Erisman, W. Heyman, S. Kobara, T. Ezer, S. Pittman, O. Aburto-Oropeza, R. S. Nemeth, Fish spawning aggregations: where well-placed management actions can yield big benefits for fisheries and conservation. *Fish and Fisheries*. **18**, 128–144 (2017).
73. C. M. Hernández, J. Witting, C. Willis, S. R. Thorrold, J. K. Llopiz, R. D. Rotjan, Evidence and patterns of tuna spawning inside a large no-take Marine Protected Area. *Sci. Rep.* **9**, 10772 (2019).
- 30 74. M. H. Carr, S. P. Robinson, C. Wahle, G. Davis, S. Kroll, S. Murray, E. J. Schumacker, M. Williams, The central importance of ecological spatial connectivity to effective coastal marine protected areas and to meeting the challenges of climate change in the marine environment. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **27**, 6–29 (2017).
- 35 75. M. Di Lorenzo, J. Claudet, P. Guidetti, Spillover from marine protected areas to adjacent fisheries has an ecological and a fishery component. *J. Nat. Cons.* **32**, 62–66 (2016).

76. “Reef 2050 Water Quality Improvement Plan 2017-2020” (State of Queensland, 2018), p. 63.
77. J. E. Duffy, J. S. Lefcheck, R. D. Stuart-Smith, S. A. Navarrete, G. J. Edgar, Biodiversity enhances reef fish biomass and resistance to climate change. *PNAS*. **113**, 6230–6235 (2016).
78. S. D. Gaines, C. Costello, B. Owashi, T. Mangin, J. Bone, J. G. Molinos, M. Burden, H. Dennis, B. S. Halpern, C. V. Kappel, K. M. Kleisner, D. Ovando, Improved fisheries management could offset many negative effects of climate change. *Sci. Adv.* **4**, eaao1378 (2018).
79. L. Sheehan, E. T. Sherwood, R. P. Moyer, K. R. Radabaugh, S. Simpson, Blue carbon: an additional driver for restoring and preserving ecological services of coastal wetlands in Tampa Bay (Florida, USA). *Wetlands*. **39**, 1317–1328 (2019).
80. G. W. Allison, S. D. Gaines, J. Lubchenco, H. P. Possingham, Ensuring persistence of marine reserves: Catastrophes require adopting an insurance factor. *Ecol. Appl.* **13**, 8–24 (2003).
81. C. Mellin, M. A. MacNeil, A. J. Cheal, M. J. Emslie, M. J. Caley, Marine protected areas increase resilience among coral reef communities. *Ecology Letters*. **19**, 629–637 (2016).
82. M. J. Emslie, M. Logan, D. H. Williamson, A. M. Ayling, M. A. MacNeil, D. Ceccarelli, A. J. Cheal, R. D. Evans, K. A. Johns, M. J. Jonker, I. R. Miller, K. Osborne, G. R. Russ, H. P. A. Sweatman, Expectations and outcomes of reserve network performance following re-zoning of the Great Barrier Reef Marine Park. *Curr. Biol.* **25**, 983–992 (2015).
83. R. C. Babcock, A. C. Alcala, K. D. Lafferty, T. McClanahan, G. R. Russ, N. T. Shears, N. S. Barrett, G. J. Edgar, Conservation or restoration: decadal trends in marine reserves. *Proceedings of the National Academy of Sciences of the United States of America*. **107**, 18256–18261 (2010).
84. A. M. Friedlander, M. K. Donovan, H. Koike, P. Murakawa, W. Goodell, Characteristics of effective marine protected areas in Hawai‘i. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **29**, 103–117 (2019).
85. O. Aburto-Oropeza, B. Erisman, G. R. Galland, I. Mascareñas-Osorio, E. Sala, E. Ezcurra, Large recovery of fish biomass in a no-take marine reserve. *PLOS ONE*. **6**, e23601 (2011).
86. P. Guidetti, M. Milazzo, S. Bussotti, A. Molinari, M. Murenu, A. Pais, N. Spano, R. Balzano, T. Agardy, F. Boero, G. Carrada, R. Cattaneo-Vietti, A. Cau, R. Chemello, S. Greco, A. Manganaro, G. N. di Sciara, G. F. Russo, L. Tunesi, Italian marine reserve effectiveness: Does enforcement matter? *Biol Conserv Biol Conserv*. **141**, 699–709 (2008).

87. M. Kaplan-Hallam, N. J. Bennett, Adaptive social impact management for conservation and environmental management. *Conservation Biology*. **32**, 304–314 (2018).
88. N. C. Ban, G. G. Gurney, N. A. Marshall, C. K. Whitney, M. Mills, S. Gelcich, N. J. Bennett, M. C. Meehan, C. Butler, S. Ban, T. C. Tran, M. E. Cox, S. J. Breslow, Well-being outcomes of marine protected areas. *Nat Sustain.* **2**, 524–532 (2019).
89. D. A. Gill, S. H. Cheng, L. Glew, E. Aigner, N. J. Bennett, M. B. Mascia, Social synergies, tradeoffs, and equity in marine conservation impacts. *Annu. Rev. Environ. Resour.* **44**, 347–372 (2019).
90. D. A. Gill, H. A. Oxenford, P. W. Schuhmann, in *Viability and Sustainability of Small-Scale Fisheries in Latin America and The Caribbean*, S. Salas, M. J. Barragán-Paladines, R. Chuenpagdee, Eds. (Springer International Publishing, Cham, 2019; https://doi.org/10.1007/978-3-319-76078-0_13), *MARE Publication Series*, pp. 295–328.
91. M. Schratzberger, S. Neville, S. Painting, K. Weston, L. Paltriguera, Ecological and socio-economic effects of Highly Protected Marine Areas (HPMAs) in temperate waters. *Front. Mar. Sci.* **6** (2019), doi:10.3389/fmars.2019.00749.
92. G. G. Gurney, J. Cinner, N. C. Ban, R. L. Pressey, R. Pollnac, S. J. Campbell, S. Tasidjawa, F. Setiawan, Poverty and protected areas: An evaluation of a marine integrated conservation and development project in Indonesia. *Glob. Environ. Change-Human Policy Dimens.* **26**, 98–107 (2014).
93. C. E. Hattam, S. C. Mangi, S. C. Gall, L. D. Rodwell, Social impacts of a temperate fisheries closure: Understanding stakeholders’ views. *Mar. Pol.* **45**, 269–278 (2014).
94. R. Chuenpagdee, J. J. Pascual-Fernández, E. Szeliánszky, J. Luis Alegret, J. Fraga, S. Jentoft, Marine protected areas: Re-thinking their inception. *Marine Policy*. **39**, 234–240 (2013).
95. H. S. Macedo, R. P. Medeiros, P. McConney, Are multiple-use marine protected areas meeting fishers’ proposals? Strengths and constraints in fisheries’ management in Brazil. *Mar. Pol.* **99**, 351–358 (2019).
96. C. Leisher, P. van Beukering, L. Scherl, “Nature’s investment bank: how marine protected areas contribute to poverty reduction.” (The Nature Conservancy, Arlington, Virginia, 2007).
97. E. Sala, C. Costello, D. Dougherty, G. Heal, K. Kelleher, J. H. Murray, A. A. Rosenberg, R. Sumaila, A general business model for marine reserves. *PLoS ONE*. **8**, e58799 (2013).
98. M. D. Smith, J. Zhang, F. Coleman, Effectiveness of marine reserves for large-scale fisheries management. *Can. J. Fish. Aquat. Sci.* **63** (153AD), doi:10.1139/f05-205.

99. K. Hogg, T. Gray, P. Noguera-Méndez, M. Semitiel-García, S. Young, Interpretations of MPA winners and losers: a case study of the Cabo De Palos- Islas Hormigas Fisheries Reserve. *Maritime Studies*. **18**, 159–171 (2019).
- 5 100. G. G. Gurney, R. L. Pressey, J. E. Cinner, R. Pollnac, S. J. Campbell, Integrated conservation and development: Evaluating a community-based marine protected area project for equality of socioeconomic impacts. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20140277 (2015).
- 10 101. S. M. Alexander, D. Armitage, P. J. Carrington, O. Bodin, Examining horizontal and vertical social ties to achieve social-ecological fit in an emerging marine reserve network. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **27**, 1209–1223 (2017).
102. C. G. McNally, E. Uchida, A. J. Gold, The effect of a protected area on the tradeoffs between short-run and long-run benefits from mangrove ecosystems. *PNAS*. **108**, 13945–13950 (2011).
- 15 103. M. Gustavsson, L. Lindström, N. S. Jiddawi, M. de la Torre-Castro, Procedural and distributive justice in a community-based managed Marine Protected Area in Zanzibar, Tanzania. *Marine Policy*. **46**, 91–100 (2014).
104. E. G. Oracion, M. L. Miller, P. Christie, Marine protected areas for whom? Fisheries, tourism, and solidarity in a Philippine community. *Ocean Coast Manage Ocean Coast Manage*. **48**, 393–410 (2005).
- 20 105. J. E. Cinner, T. Daw, C. Huchery, P. Thoya, A. Wamukota, M. Cedras, C. Abunge, Winners and losers in marine conservation: Fishers’ displacement and livelihood benefits from marine reserves. *Soc. Nat. Resour.* **27**, 994–1005 (2014).
- 25 106. G. J. Edgar, R. D. Stuart-Smith, T. J. Willis, S. Kininmonth, S. C. Baker, S. Banks, N. S. Barrett, M. A. Becerro, A. T. F. Bernard, J. Berkhout, C. D. Buxton, S. J. Campbell, A. T. Cooper, M. Davey, S. C. Edgar, G. Foerster, D. E. Galvan, A. J. Irigoyen, D. J. Kushner, R. Moura, P. E. Parnell, N. T. Shears, G. Soler, E. M. A. Strain, R. J. Thomson, Global conservation outcomes depend on marine protected areas with five key features. *Nature*. **506**, 216–220 (2014).
- 30 107. E. Northrop, Manaswita Konar, Nicola Frost, Elizabeth Hollaway, “A Sustainable and Equitable Blue Recovery to the COVID-19 Crisis” (World Resources Institute, Washington, D.C., 2020), (available at <http://www.oceanpanel.org/bluerecovery>).
108. Kementerian Kelautan dan Perikanan, *Management of Marine Protected Areas in Indonesia: Status and Challenges* (Ministry of Marine Affairs and Fisheries (MMAF) and Yayasan WWF Indonesia, 2021; <https://doi.org/10.6084/m9.figshare.13341476.v1>).
- 35 109. K. Grorud-Colvert, V. Constant, J. Sullivan-Stack, K. Dziedzic, S. L. Hamilton, Z. Randell, H. Fulton-Bennett, Z. D. Meunier, S. Bachhuber, A. J. Rickborn, B. Spiecker, J.

Lubchenco, High-profile international commitments for ocean protection: Empty promises or meaningful progress? *Mar. Pol.* **105**, 52–66 (2019).

- 5 110. H. Österblom, C. Wabnitz, D. Tladi, E. Allison, S. Arnaud-Haond, J. Bebbington, N. Bennett, R. Blasiak, W. Boonstra, A. Choudhury, A. Cisneros-Montemayor, T. Daw, M. Fabinyi, N. Franz, H. Harden-Davies, D. Kleiber, P. Lopes, C. McDougall, B. Resosudarmo, S. Selim, “Towards ocean equity” (High Level Panel for a Sustainable Ocean Economy, 2020).
- 10 111. I. M. Côté, I. Mosqueira, J. D. Reynolds, Effects of marine reserve characteristics on the protection of fish populations: a meta-analysis. *J. Fish Biol.* **59**, 178–189 (2001).
- 10 112. S. Lester, B. Halpern, Biological responses in marine no-take reserves versus partially protected areas. *Mar. Ecol. Prog. Ser.* **367**, 49–56 (2008).
113. C. M. Roberts, J. A. Bohnsack, F. Gell, J. P. Hawkins, R. Goodridge, Effects of marine reserves on adjacent fisheries. *Science*. **294**, 1920–1923 (2001).
- 15 114. J. Claudet, D. Pelletier, J.-Y. Jouvenel, F. Bachet, R. Galzin, Assessing the effects of marine protected area (MPA) on a reef fish assemblage in a northwestern Mediterranean marine reserve: Identifying community-based indicators. *Biol. Cons.* **130**, 349–369 (2006).
115. B. I. Ruttenberg, S. L. Hamilton, S. M. Walsh, M. K. Donovan, A. Friedlander, E. DeMartini, E. Sala, S. A. Sandin, Predator-Induced Demographic Shifts in Coral Reef Fish Assemblages. *PLOS ONE*. **6**, e21062 (2011).
- 20 116. A. García-Rubies, B. Hereu, M. Zabala, Long-term recovery patterns and limited spillover of large predatory fish in a Mediterranean MPA. *PLOS ONE*. **8**, e73922 (2013).
117. R. A. Abesamis, A. L. Green, G. R. Russ, C. R. L. Jadloc, The intrinsic vulnerability to fishing of coral reef fishes and their differential recovery in fishery closures. *Rev. Fish. Biol. Fisheries*. **24**, 1033–1063 (2014).
- 25 118. H. A. Malcolm, A. L. Schultz, P. Sachs, N. Johnstone, A. Jordan, Decadal changes in the abundance and length of snapper (*Chrysophrys auratus*) in subtropical marine sanctuaries. *PLOS ONE*. **10**, e0127616 (2015).
119. D. Harasti, J. Williams, E. Mitchell, S. Lindfield, A. Jordan, Increase in relative abundance and size of snapper *Chrysophrys auratus* within partially-protected and no-take areas in a temperate marine protected area. *Front. Mar. Sci.* **5** (2018), doi:10.3389/fmars.2018.00208.
- 30 120. E. Sala, E. Ballesteros, P. Dendrinis, A. D. Franco, F. Ferretti, D. Foley, S. Fraschetti, A. Friedlander, J. Garrabou, H. Güçlüsoy, P. Guidetti, B. S. Halpern, B. Hereu, A. A. Karamanlidis, Z. Kizilkaya, E. Macpherson, L. Mangialajo, S. Mariani, F. Micheli, A. Pais, K. Riser, A. A. Rosenberg, M. Sales, K. A. Selkoe, R. Starr, F. Tomas, M. Zabala,
- 35

The structure of Mediterranean rocky reef ecosystems across environmental and human gradients, and conservation implications. *PLOS ONE*. **7**, e32742 (2012).

- 5 121. P. Guidetti, P. Baiata, E. Ballesteros, A. Di Franco, B. Hereu, E. Macpherson, F. Micheli, A. Pais, P. Panzalis, A. A. Rosenberg, M. Zabala, E. Sala, Large-scale assessment of Mediterranean marine protected areas effects on fish assemblages. *PLOS ONE*. **9**, e91841 (2014).
122. E. Sala, S. Giakoumi, No-take marine reserves are the most effective protected areas in the ocean. *ICES J Mar Sci*. **75**, 1166–1168 (2018).
- 10 123. D. Agnetta, F. Badalamenti, F. Colloca, G. D’Anna, M. Di Lorenzo, F. Fiorentino, G. Garofalo, M. Gristina, L. Labanchi, B. Patti, C. Pipitone, C. Solidoro, S. Libralato, Benthic-pelagic coupling mediates interactions in Mediterranean mixed fisheries: An ecosystem modeling approach. *PLOS ONE*. **14** (2019), doi:10.1371/journal.pone.0210659.
124. G. R. Russ, A. C. Alcala, Enhanced biodiversity beyond marine reserve boundaries: The cup spillith over. *Ecol. Appl.* **21**, 241–250 (2011).
- 15 125. K. L. Nash, N. A. J. Graham, Ecological indicators for coral reef fisheries management. *Fish. Fish.* **17**, 1029–1054 (2016).
126. R. S. Nemeth, Population characteristics of a recovering US Virgin Islands red hind spawning aggregation following protection. *Mar. Ecol. Prog. Ser.* **286**, 81–97 (2005).
- 20 127. M. J. Kaiser, R. E. Blyth-Skyrme, P. J. Hart, G. Edwards-Jones, D. Palmer, Evidence for greater reproductive output per unit area in areas protected from fishing. *Can. J. Fish. Aquat. Sci.* **64**, 1284–1289 (2007).
128. R. Crec’hriou, F. Alemany, E. Roussel, A. Chassanite, J. Y. Marinaro, J. Mader, E. Rochel, S. Planes, Fisheries replenishment of early life taxa: potential export of fish eggs and larvae from a temperate marine protected area. *Fish. Oceanog.* **19**, 135–150 (2010).
- 25 129. B. M. Taylor, J. L. McIlwain, Beyond abundance and biomass: effects of marine protected areas on the demography of a highly exploited reef fish. *Mar. Ecol. Prog. Ser.* **411**, 243–258 (2010).
130. D. Díaz, S. Mallol, A. M. Parma, R. Goñi, Decadal trend in lobster reproductive output from a temperate marine protected area. *Mar. Ecol. Prog. Ser.* **433**, 149–157 (2011).
- 30 131. M. A. Hixon, D. W. Johnson, S. M. Sogard, BOFFFFs: on the importance of conserving old-growth age structure in fishery populations. *ICES J. Mar. Sci.* **71**, 2171–2185 (2014).
132. D. R. Barneche, D. R. Robertson, C. R. White, D. J. Marshall, Fish reproductive-energy output increases disproportionately with body size. *Science*. **360**, 642–645 (2018).

133. D. J. Marshall, S. Gaines, R. Warner, D. R. Barneche, M. Bode, Underestimating the benefits of marine protected areas for the replenishment of fished populations. *Front. Ecol. Environ.* **17**, 407–413 (2019).
- 5 134. R. A. Pelc, R. R. Warner, S. D. Gaines, C. B. Paris, Detecting larval export from marine reserves. *PNAS.* **107**, 18266–18271 (2010).
135. M. R. Christie, B. N. Tissot, M. A. Albins, J. P. Beets, Y. Jia, D. M. Ortiz, S. E. Thompson, M. A. Hixon, Larval connectivity in an effective network of marine protected areas. *PLOS ONE.* **5**, e15715 (2010).
- 10 136. A. D. Franco, B. M. Gillanders, G. D. Benedetto, A. Pennetta, G. A. D. Leo, P. Guidetti, Dispersal Patterns of Coastal Fish: Implications for Designing Networks of Marine Protected Areas. *PLOS ONE.* **7**, e31681 (2012).
137. C. M. Roberts, J. P. Hawkins, “Establishment of fish stock recovery areas” (European Parliament, 2012), p. 70.
- 15 138. M. Andrello, F. Guilhaumon, C. Albouy, V. Parravicini, J. Scholtens, P. Verley, M. Barange, U. R. Sumaila, S. Manel, D. Mouillot, Global mismatch between fishing dependency and larval supply from marine reserves. *Nat. Commun.* **8**, 1–9 (2017).
- 20 139. S. Manel, N. Loiseau, M. Andrello, K. Fietz, R. Goñi, A. Forcada, P. Lenfant, S. Kininmonth, C. Marcos, V. Marques, S. Mallol, A. Pérez-Ruzafa, C. Breusing, O. Puebla, D. Mouillot, Long-distance benefits of marine reserves: Myth or reality? *Trends Ecol. Evol.* **34**, 342–354 (2019).
140. J. Assis, E. Fragkopoulou, E. A. Serrão, B. Horta e Costa, M. Gandra, D. Abecasis, Weak biodiversity connectivity in the European network of no-take marine protected areas. *Sci. Total Env.* **773**, 145664 (2021).
- 25 141. D. Mouillot, J. M. Culioli, D. Pelletier, J. A. Tomasini, Do we protect biological originality in protected areas? A new index and an application to the Bonifacio Strait Natural Reserve. *Biol. Cons.* **141**, 1569–1580 (2008).
142. L. Pichegru, D. Grémillet, R. J. M. Crawford, P. G. Ryan, Marine no-take zone rapidly benefits endangered penguin. *Biol. Lett.* **6**, 498–501 (2010).
- 30 143. A. M. Gormley, E. Sooten, S. Dawson, R. J. Barker, W. Rayment, S. du Fresne, S. Bräger, First evidence that marine protected areas can work for marine mammals. *J. Appl. Ecol.* **49**, 474–480 (2012).
144. J. S. Goetze, S. D. Jupiter, T. J. Langlois, S. K. Wilson, E. S. Harvey, T. Bond, W. Naisilisili, Diver operated video most accurately detects the impacts of fishing within periodically harvested closures. *J. Exp. Mar. Biol. Ecol.* **462**, 74–82 (2015).

145. B. W. McLaren, T. J. Langlois, E. S. Harvey, H. Shortland-Jones, R. Stevens, A small no-take marine sanctuary provides consistent protection for small-bodied by-catch species, but not for large-bodied, high-risk species. *J. Exp. Mar. Biol. Ecol.* **471**, 153–163 (2015).
- 5 146. R. G. Dwyer, N. C. Krueck, V. Udyawer, M. R. Heupel, D. Chapman, H. L. Pratt, R. Garla, C. A. Simpfendorfer, Individual and population benefits of marine reserves for reef sharks. *Curr. Biol.* **30**, 480-489.e5 (2020).
147. T. Miete, C. Dytham, U. Dieckmann, J. W. Pitchford, Marine reserves and the evolutionary effects of fishing on size at maturation. *ICES J. Mar. Sci.* **67**, 412–425 (2010).
- 10 148. R. Y. Fidler, J. Carroll, K. W. Rynerson, D. F. Matthews, R. G. Turingan, Coral reef fishes exhibit beneficial phenotypes inside marine protected areas. *PLOS ONE*. **13**, e0193426 (2018).
149. K. R. Jones, C. J. Klein, B. S. Halpern, O. Venter, H. Grantham, C. D. Kuempel, N. Shumway, A. M. Friedlander, H. P. Possingham, J. E. M. Watson, The location and protection status of Earth’s diminishing marine wilderness. *Curr. Biol.* **28**, 2506-2512.e3 (2018).
- 15 150. T. K. Sørvalen, K. T. Halvorsen, H. B. Harrison, C. D. Ellis, L. A. Vøllestad, H. Knutsen, E. Moland, E. M. Olsen, Harvesting changes mating behaviour in European lobster. *Evol. Appl.* **11**, 963–977 (2018).
- 20 151. P. Guidetti, Potential of marine reserves to cause community-wide changes beyond their boundaries. *Conserv. Biol.* **21**, 540–545 (2007).
152. M. J. Costello, Long live Marine Reserves: A review of experiences and benefits. *Biol. Cons.* **176**, 289–296 (2014).
153. D. H. Williamson, D. M. Ceccarelli, R. D. Evans, G. P. Jones, G. R. Russ, Habitat dynamics, marine reserve status, and the decline and recovery of coral reef fish communities. *Ecol. Evol.* **4**, 337–354 (2014).
- 25 154. J. W. Turnbull, Y. Shah Esmaeili, G. F. Clark, W. F. Figueira, E. L. Johnston, R. Ferrari, Key drivers of effectiveness in small marine protected areas. *Biodivers. Conserv.* **27**, 2217–2242 (2018).
- 30 155. P. Guidetti, Marine reserves reestablish lost predatory interactions and cause community changes in rocky reefs. *Ecol. Appl.* **16**, 963–976 (2006).
156. J. Claudet, C. W. Osenberg, P. Domenici, F. Badalamenti, M. Milazzo, J. M. Falcón, I. Bertocci, L. Benedetti-Cecchi, J.-A. García-Charton, R. Goñi, J. A. Borg, A. Forcada, G. A. de Lucia, Á. Pérez-Ruzafa, P. Afonso, A. Brito, I. Guala, L. L. Diréach, P. Sanchez-Jerez, P. J. Somerfield, S. Planes, Marine reserves: Fish life history and ecological traits matter. *Ecol. Appl.* **20**, 830–839 (2010).
- 35

157. T. R. McClanahan, N. a. J. Graham, Marine reserve recovery rates towards a baseline are slower for reef fish community life histories than biomass. *Proc. R. Soc. B.* **282**, 20151938 (2015).
- 5 158. G. R. Russ, K. I. Miller, J. R. Rizzari, A. C. Alcala, Long-term no-take marine reserve and benthic habitat effects on coral reef fishes. *Mar. Ecol. Prog. Ser.* **529**, 233–248 (2015).
159. D. Acuña-Marrero, A. N. H. Smith, N. Hammerschlag, A. Hearn, M. J. Anderson, H. Calich, M. D. M. Pawley, C. Fischer, P. Salinas-de-León, Residency and movement patterns of an apex predatory shark (*Galeocerdo cuvier*) at the Galapagos Marine Reserve. *PLOS ONE.* **12**, e0183669 (2017).
- 10 160. R. L. Selden, S. D. Gaines, S. L. Hamilton, R. R. Warner, Protection of large predators in a marine reserve alters size-dependent prey mortality. *Proc. R. Soc. B.* **284**, 20161936 (2017).
161. E. McLeod, R. Salm, A. Green, J. Almany, Designing marine protected area networks to address the impacts of climate change. *Front. Ecol. Environ.* **7**, 362–370 (2009).
- 15 162. S. D. Ling, C. R. Johnson, S. D. Frusher, K. R. Ridgway, Overfishing reduces resilience of kelp beds to climate-driven catastrophic phase shift. *PNAS.* **106**, 22341–22345 (2009).
163. F. Micheli, A. Saenz-Arroyo, A. Greenley, L. Vazquez, J. A. E. Montes, M. Rossetto, G. A. D. Leo, Evidence that marine reserves enhance resilience to climatic impacts. *PLOS ONE.* **7**, e40832 (2012).
- 20 164. L. A. K. Barnett, M. L. Baskett, Marine reserves can enhance ecological resilience. *Ecol. Lett.* **18**, 1301–1310 (2015).
165. K. L. Wilson, D. P. Tittensor, B. Worm, K. L. Heike, Incorporating climate change adaptation into marine protected area planning. *Global Change Biol.*, 3251–3267 (2020).
- 25 166. R. A. Abesamis, G. R. Russ, Density-dependent spillover from a marine reserve: Long-term evidence. *Ecol. Appl.* **15**, 1798–1812 (2005).
167. B. S. Halpern, S. E. Lester, J. B. Kellner, Spillover from marine reserves and the replenishment of fished stocks. *Env. Cons.* **36**, 268–276 (2009).
168. M. D. Lorenzo, P. Guidetti, A. D. Franco, A. Calò, J. Claudet, Assessing spillover from marine protected areas and its drivers: A meta-analytical approach. *Fish. Fish.* **21**, 906–915 (2020).
- 30 169. P. H. Manríquez, J. C. Castilla, Significance of marine protected areas in central Chile as seeding grounds for the gastropod *Concholepas concholepas*. *Mar. Ecol. Prog. Ser.* **215**, 201–211 (2001).

170. S. Planes, G. Jones, S. Thorrold, Larval dispersal connects fish populations in a network of marine protected areas. *PNAS* (2009), doi:10.1073/pnas.0808007106.
- 5 171. A. Di Franco, A. Calò, A. Pennetta, G. De Benedetto, S. Planes, P. Guidetti, Dispersal of larval and juvenile seabream: Implications for Mediterranean marine protected areas. *Biol. Cons.* **192**, 361–368 (2015).
172. T. Lauck, C. W. Clark, M. Mangel, G. R. Munro, Implementing the Precautionary Principle in Fisheries Management Through Marine Reserves. *Ecol. Appl.* **8**, S72–S78 (1998).
- 10 173. C. M. Roberts, J. P. Hawkins, F. R. Gell, The role of marine reserves in achieving sustainable fisheries. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **360**, 123–132 (2005).
174. N. C. Krueck, G. N. Ahmadi, H. P. Possingham, C. Riginos, E. A. Treml, P. J. Mumby, Marine Reserve Targets to Sustain and Rebuild Unregulated Fisheries. *PLOS Biology*. **15**, e2000537 (2017).
- 15 175. J. Beets, A. Friedlander, Evaluation of a conservation strategy: a spawning aggregation closure for red hind, *Epinephelus guttatus*, in the U.S. Virgin Islands. *Env. Biol. Fish.* **55**, 91–98 (1999).
176. L. Rogers-Bennett, J. S. Pearse, Indirect benefits of marine protected areas for juvenile abalone. *Cons. Biol.* **15**, 642–647 (2001).
- 20 177. E. Sala, E. Ballesteros, R. M. Starr, Rapid decline of Nassau Grouper spawning aggregations in Belize: Fishery management and conservation needs. *Fisheries*. **26**, 23–30 (2001).
178. R. C. Garla, D. D. Chapman, B. M. Wetherbee, M. Shivji, Movement patterns of young Caribbean reef sharks, *Carcharhinus perezi*, at Fernando de Noronha Archipelago, Brazil: the potential of marine protected areas for conservation of a nursery ground. *Mar. Biol.* **149**, 189–199 (2006).
- 25 179. P. R. Armsworth, B. A. Block, J. Eagle, J. E. Roughgarden, The economic efficiency of a time–area closure to protect spawning bluefin tuna. *J. Appl. Ecol.* **47**, 36–46 (2010).
180. A. Grüss, D. M. Kaplan, J. Robinson, Evaluation of the effectiveness of marine reserves for transient spawning aggregations in data-limited situations. *ICES J. Mar. Sci.* **71**, 435–449 (2014).
- 30 181. N. A. Farmer, W. D. Heyman, M. Karnauskas, S. Kobara, T. I. Smart, J. C. Ballenger, M. J. M. Reichert, D. M. Wyanski, M. S. Tishler, K. C. Lindeman, S. K. Lowerre-Barbieri, T. S. Switzer, J. J. Solomon, K. McCain, M. Marhefka, G. R. Sedberry, Timing and locations of reef fish spawning off the southeastern United States. *PLOS ONE*. **12**, e0172968 (2017).
- 35

182. Y. Sadovy de Mitcheson, P. L. Colin, S. J. Lindfield, A. Bukurrou, A decade of monitoring an Indo-Pacific grouper spawning aggregation: Benefits of protection and importance of survey design. *Front. Mar. Sci.* **7** (2020), doi:10.3389/fmars.2020.571878.
- 5 183. A. D. Olds, K. A. Pitt, P. S. Maxwell, R. C. Babcock, D. Rissik, R. M. Connolly, Marine reserves help coastal ecosystems cope with extreme weather. *Globe Change Biol.* **20**, 3050–3058 (2014).
184. D. M. Alongi, N. L. Patten, D. McKinnon, N. Köstner, D. G. Bourne, R. Brinkman, Phytoplankton, bacterioplankton and virioplankton structure and function across the southern Great Barrier Reef shelf. *J. Mar. Sys.* **142**, 25–39 (2015).
- 10 185. A. D. McKinnon, S. Duggan, M. Logan, C. Lønborg, Plankton Respiration, Production, and Trophic State in Tropical Coastal and Shelf Waters Adjacent to Northern Australia. *Front. Mar. Sci.* **4** (2017), doi:10.3389/fmars.2017.00346.
- 15 186. L. Bergström, M. Karlsson, U. Bergström, L. Pihl, P. Kraufvelin, Relative impacts of fishing and eutrophication on coastal fish assessed by comparing a no-take area with an environmental gradient. *Ambio.* **48**, 565–579 (2019).
187. E. M. A. Strain, G. J. Edgar, D. Ceccarelli, R. D. Stuart-Smith, G. R. Hosack, R. J. Thomson, A global assessment of the direct and indirect benefits of marine protected areas for coral reef conservation. *Divers. Distr.* **25**, 9–20 (2019).
- 20 188. E. Cotou, A. Gremare, F. Charles, I. Hatzianestis, E. Sklivagou, Potential toxicity of resuspended particulate matter and sediments: Environmental samples from the Bay of Banyuls-sur-Mer and Thermaikos Gulf. *Cont. Shelf Res.* **25**, 2521–2532 (2005).
- 25 189. X. Durrieu de Madron, B. Ferré, G. Le Corre, C. Grenz, P. Conan, M. Pujo-Pay, R. Buscail, O. Bodiou, Trawling-induced resuspension and dispersal of muddy sediments and dissolved elements in the Gulf of Lion (NW Mediterranean). *Cont. Shelf Res.* **25**, 2387–2409 (2005).
190. J. B. Lamb, J. A. J. M. van de Water, D. G. Bourne, C. Altier, M. Y. Hein, E. A. Fiorenza, N. Abu, J. Jompa, C. D. Harvell, Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes, and invertebrates. *Science.* **355**, 731–733 (2017).
- 30 191. F. J. Pollock, J. B. Lamb, S. N. Field, S. F. Heron, B. Schaffelke, G. Shedrawi, D. G. Bourne, B. L. Willis, Sediment and turbidity associated with offshore dredging increase coral disease prevalence on nearby reefs. *PLOS ONE.* **9** (2014), doi:10.1371/journal.pone.0102498.
192. State of Queensland, “Reef 2050 Water Quality Improvement Plan 2017-2022” (State of Queensland, 2018), p. 56.

193. E. J. Powell, M. C. Tyrrell, A. Milliken, J. M. Tirpak, M. D. Staudinger, A review of coastal management approaches to support the integration of ecological and human community planning for climate change. *J. Coast Conserv.* **23**, 1–18 (2019).
- 5 194. L. Pendleton, D. C. Donato, B. C. Murray, S. Crooks, W. A. Jenkins, S. Sifleet, C. Craft, J. W. Fourqurean, J. B. Kauffman, N. Marbà, P. Megonigal, E. Pidgeon, D. Herr, D. Gordon, A. Baldera, Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems. *PLOS ONE*. **7**, e43542 (2012).
- 10 195. T. B. Atwood, R. M. Connolly, E. G. Ritchie, C. E. Lovelock, M. R. Heithaus, G. C. Hays, J. W. Fourqurean, P. I. Macreadie, Predators help protect carbon stocks in blue carbon ecosystems. *Nat. Clim. Change*. **5**, 1038–1045 (2015).
- 15 196. F. Mineur, F. Arenas, J. Assis, A. J. Davies, A. H. Engelen, F. Fernandes, E. Malta, T. Thibaut, T. Van Nguyen, F. Vaz-Pinto, S. Vranken, E. A. Serrão, O. De Clerck, European seaweeds under pressure: Consequences for communities and ecosystem functioning. *J. Sea Res.* **98**, 91–108 (2015).
- 20 197. T. G. Zarate-Barrera, J. H. Maldonado, Valuing Blue Carbon: Carbon Sequestration Benefits Provided by the Marine Protected Areas in Colombia. *PLOS ONE*. **10**, e0126627 (2015).
- 25 198. D. Krause-Jensen, C. M. Duarte, Substantial role of macroalgae in marine carbon sequestration. *Nature Geosci.* **9**, 737–742 (2016).
- 30 199. J. Howard, E. McLeod, S. Thomas, E. Eastwood, M. Fox, L. Wenzel, E. Pidgeon, The potential to integrate blue carbon into MPA design and management. *Aquat. Conserv.-Mar. Freshw. Ecosyst.* **27**, 100–115 (2017).
- 35 200. C. M. Duarte, S. Agusti, E. Barbier, G. L. Britten, J. C. Castilla, J.-P. Gattuso, R. W. Fulweiler, T. P. Hughes, N. Knowlton, C. E. Lovelock, H. K. Lotze, M. Predragovic, E. Poloczanska, C. Roberts, B. Worm, Rebuilding marine life. *Nature*. **580**, 39–51 (2020).
201. G. Mariani, W. W. L. Cheung, A. Lyet, E. Sala, J. Mayorga, L. Velez, S. D. Gaines, T. Dejean, M. Troussellier, D. Mouillot, Let more big fish sink: Fisheries prevent blue carbon sequestration—half in unprofitable areas. *Sci. Adv.* **6**, eabb4848 (2020).
202. G. K. Saba, A. B. Burd, J. P. Dunne, S. Hernández-León, A. H. Martin, K. A. Rose, J. Salisbury, D. K. Steinberg, C. N. Trueman, R. W. Wilson, S. E. Wilson, Toward a better understanding of fish-based contribution to ocean carbon flux. *Limnol. Oceanog.* **n/a**, doi:10.1002/lno.11709.
203. R. K. F. Unsworth, C. J. Collier, G. M. Henderson, L. J. McKenzie, Tropical seagrass meadows modify seawater carbon chemistry: implications for coral reefs impacted by ocean acidification. *Environ. Res. Lett.* **7**, 024026 (2012).

204. C. M. Duarte, J. Wu, X. Xiao, A. Bruhn, D. Krause-Jensen, Can seaweed farming play a role in climate change mitigation and adaptation? *Front. Mar. Sci.* **4** (2017), doi:10.3389/fmars.2017.00100.
- 5 205. D. A. Koweek, R. C. Zimmerman, K. M. Hewett, B. Gaylord, S. N. Giddings, K. J. Nickols, J. L. Ruesink, J. J. Stachowicz, Y. Takeshita, K. Caldeira, Expected limits on the ocean acidification buffering potential of a temperate seagrass meadow. *Ecol. Appl.* **28**, 1694–1714 (2018).
206. D. Grémillet, T. Boulinier, Spatial ecology and conservation of seabirds facing global climate change: a review. *Mar. Ecol. Prog. Ser.* **391**, 121–137 (2009).
- 10 207. D. Reed, L. Washburn, A. Rassweiler, R. Miller, T. Bell, S. Harrer, Extreme warming challenges sentinel status of kelp forests as indicators of climate change. *Nat. Commun.* **7** (2016), doi:10.1038/ncomms13757.
- 15 208. E. L. A. Kelly, Y. Eynaud, I. D. Williams, R. T. Sparks, M. L. Dailer, S. A. Sandin, J. E. Smith, A budget of algal production and consumption by herbivorous fish in an herbivore fisheries management area, Maui, Hawaii. *Ecosphere.* **8**, e01899 (2017).
209. L. Rogers-Bennett, C. A. Catton, Marine heat wave and multiple stressors tip bull kelp forest to sea urchin barrens. *Sci. Rep.* **9**, 15050 (2019).
210. S. Luo, F. Cai, H. Liu, G. Lei, H. Qi, X. Su, Adaptive measures adopted for risk reduction of coastal erosion in the People’s Republic of China. *Ocean Coast. Manag.* **103**, 134–145 (2015).
- 20 211. D. A. Miteva, B. C. Murray, S. K. Pattanayak, Do protected areas reduce blue carbon emissions? A quasi-experimental evaluation of mangroves in Indonesia. *Ecol. Econ.* **119**, 127–135 (2015).
212. S. Narayan, M. W. Beck, B. G. Reguero, I. J. Losada, B. van Wesenbeeck, N. Pontee, J. N. Sanchirico, J. C. Ingram, G.-M. Lange, K. A. Burks-Copes, The effectiveness, costs and coastal protection benefits of natural and nature-based defences. *PLOS ONE.* **11**, e0154735 (2016).
- 25 213. D. L. Harris, A. Rovere, E. Casella, H. Power, R. Canavesio, A. Collin, A. Pomeroy, J. M. Webster, V. Parravicini, Coral reef structural complexity provides important coastal protection from waves under rising sea levels. *Sci. Adv.* **4**, eaao4350 (2018).
- 30

Acknowledgments: The authors thank the many ocean experts from over 40 countries who generously shared their knowledge about MPAs during this process. The MPA Guide is

facilitated by the founding partners: The MPA Project at Oregon State University, IUCN-Marine, UNEP-WCMC, the World Database on Protected Areas and Protected Planet, National Geographic Society Pristine Seas, and Marine Conservation Institute's Marine Protection Atlas.

We thank the following partner organizations for in-kind support: Pew Bertarelli Ocean Legacy,

5 Centro de Ciências do Mar (CCMAR), Fundação para a Ciência e Tecnologia (FCT), Centre National de la Recherche Scientifique (CNRS), Prince Albert I of Monaco Foundation's Institut Océanographique de Paris, the Prince Albert II of Monaco Foundation, the Ocean Sanctuary

Alliance, and the Oceano Azul Foundation. Graphics were developed in collaboration with

Communications, INC. **Funding:** This project was funded by Oceans 5, the Kingfisher

10 Foundation, the David and Lucile Packard Foundation, and JL's Ocean Science Innovation Fund

at Oregon State University. J.C was supported by BiodivERsA. E.G. was supported by FCT -

Foundation for Science and Technology through UID/MAR/04292/2018/2019,

MARE/UIDB/MAR/04292/2020 and MARE/UIDP/MAR/04292/2020, and funds from the

Oceano Azul Foundation. **Author contributions:** K.G-C., J.L., and J.S-S. conceived the

15 research; all authors contributed to the analysis of the literature, and the concepts and writing of

the manuscript. All authors approved the submission. **Competing interests:** The authors declare

no competing interests. **Data and materials availability:** All data are available in the

manuscript or the supplementary material.

20 **Supplementary Materials:**

Fig. S1. The MPA Guide: Expanded Resources

Fig. S2. The MPA Guide Protection Level Decision Tree

Table S1. Expanded Ecological Outcomes

References (101-203)



Fig. 1. Level of protection based on maximum allowed impact of seven potential activities in

MPAs. An MPA or MPA zone can be categorized into one of four LEVELS of protection: Fully,

5 Highly, Lightly, or Minimally, based on seven types of activities and their impacts (for a decision tree approach, see Fig. S2). Dials indicate the scale of impact that may be occurring at a given protection level: none, minimal, low, moderate, or high/large. If impacts are high/large, the site must still provide some conservation benefit in order to meet the definition of an MPA. If the impact of any of these activities is greater than “high”, the MPA is incompatible with the

conservation of nature (See Fig. S2). For example, some activities such as mining and mineral and oil prospecting have such a high impact that they are incompatible with biodiversity conservation and should not occur in any MPA; here, the allowed impact of mining is scored as “none” across all four LEVELS.




Fig. 2. Matrix based on LEVEL of protection and STAGE of establishment of MPAs. Any MPA or MPA zone sits in one of the sixteen cells in this matrix, based on its LEVEL and STAGE, and global area of ocean protected in MPAs can also be tallied by each matrix cell.

5 Hooks symbolize extractive use; divers indicate recreational, traditional, and cultural use; and fish indicate biodiversity outcomes. As long as CONDITIONS are in place, the OUTCOMES of an MPA will depend primarily on its protection LEVEL and STAGE, as depicted (noting that other factors such as state of ecosystem degradation prior to establishment of the MPA may also enhance or reduce outcomes). Protection does not begin until an MPA is implemented or actively

managed. The most effective biodiversity conservation OUTCOMES from an MPA are likely in the top right quadrant of this matrix, where MPAs are fully or highly protected and implemented or actively managed. In considering the global area protected, a larger percentage in the upper right quadrant would signal more effective protection than a larger percentage in the lower left quadrant.

<p>Enabling conditions across all stages of establishment</p>	<ul style="list-style-type: none"> • Clearly defined vision and objectives • Long-term political will and commitment • Sustainable financing • Public participation with contextual and procedural fairness • Evidence-based decision-making • Knowledge integration, e.g., across academic disciplines, local, Indigenous, practitioner domains • Coordination with related governance institutions • Collaboration across jurisdictions 	<ul style="list-style-type: none"> • Transparency and communication • Upward and downward accountability to legal mandate and to stakeholders • Recognition and support of existing governance by Indigenous peoples and local rights-holders, including sovereignty, self-determination, and rights of access, use, and management • Conflict resolution mechanisms
<p>Enabling conditions from Proposed/ Committed to Designated</p>	<p><i>All the Enabling Conditions above, plus:</i></p> <p>Ecological design principles:</p> <ul style="list-style-type: none"> • Viability based on MPA location, size, spacing, shape, and permanence • Representativeness and replication of habitats • Incorporation of habitats and species of unique conservation value • Design for connectivity and resilience • Precautionary approach considering current and emerging threats • Consideration of existing threats and mitigation 	<p>Social design principles:</p> <ul style="list-style-type: none"> • Inclusion of social objectives for multi-dimensional human well-being • Recognition of pre-existing rights, tenure, uses: extractive and non-extractive • Consideration of pre-existing resource use and socio-economic status • Accounting for unequal costs and benefits to different social groups • Impact- and benefit-sharing with distributional fairness
<p>Enabling conditions from Designated to Implemented</p>	<p><i>All the Enabling Conditions above, plus:</i></p> <ul style="list-style-type: none"> • Sufficient and properly organized staffing and funding • Appropriate and adequate administrative structures and processes • Stakeholder engagement plan 	<ul style="list-style-type: none"> • Compliance and enforcement (including graduated sanctioning) • Education and outreach initiatives • Clarity of rules, rights, and boundaries
<p>Enabling conditions from Implemented to Actively Managed</p>	<p><i>All the Enabling Conditions above, plus:</i></p> <ul style="list-style-type: none"> • Ongoing monitoring, evaluation, and knowledge sharing • Adaptive management • Support for livelihoods, e.g. development programs, capacity building, hiring • Effective management of broader seascape and external pressures 	<ul style="list-style-type: none"> • Ongoing efforts to build trust, strong local leadership, partnerships with local users • Local collaboration in monitoring, enforcement, and management • Ongoing consideration of cultural values, traditions, and activities in site management

Table 1. Enabling conditions for effective MPAs. These CONDITIONS may vary in their importance during the process of achieving each of the four STAGES.

Outcome		Level of Protection			
		Fully	Highly	Lightly	Minimally
Biodiversity conservation	Abundance	●	●	●	●
	Population age structure	●	●	●	●
	Biomass	●	●	●	●
	Species richness (no. of species)	●	●	●	●
	Reproductive output and replenishment	●	●	●	●
	Connectivity of populations	●	●	●	●
	Rare and endangered species protected	●	●	●	●
	Genetic diversity	●	●	●	●
	Habitats	●	●	●	●
	Ecosystem functioning	●	●	●	●
	Ecosystem resilience (ability to recover after disturbance)	●	●	●	●
	Effects on exploited species	Spillover	●	●	●
Larval export		●	●	●	●
Insurance against management failure or stock collapse		●	●	●	●
Protection of vulnerable life stages		●	●	●	●
Water quality	Eutrophication	●	●	●	●
	Pathogens and pollutants	●	●	●	●
	Suspended sediment	●	●	●	●
Climate resilience	Carbon	●	●	●	●
	Acidification	●	●	●	●
	Productivity	●	●	●	●
	Coastal protection	●	●	●	●

Table 2. Ecological OUTCOMES of MPAs as a result of LEVEL of protection. The

outcomes discussed here assume that best practices in CONDITIONS have been met and the system has had time to progress from a degraded state to one with relatively few fluctuations.

5 Not all OUTCOMES can be expected from all MPAs, as they vary by habitat type, oceanographic conditions, and previous state of degradation. Levels of confidence in Table 2 are denoted by shaded circles; the darker the circle, the higher the confidence, either high, moderate,

or low confidence. Confidence level represents expert judgments based on the quantity and quality of research available. Citations available (Table S1).