



HAL
open science

Investigating the CO trimer geometry using Coulomb imaging technique

X Fléchar, A Méry, B Gervais, S Guillous, V Kumar, J Matsumoto, J Rangamma, F Ropars, H Shiromaru, A Cassimi

► **To cite this version:**

X Fléchar, A Méry, B Gervais, S Guillous, V Kumar, et al.. Investigating the CO trimer geometry using Coulomb imaging technique. *Journal of Physics: Conference Series*, 2020, 1412, pp.132047. 10.1088/1742-6596/1412/13/132047 . hal-03345630

HAL Id: hal-03345630

<https://hal.science/hal-03345630v1>

Submitted on 2 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

PAPER • OPEN ACCESS

Investigating the CO trimer geometry using Coulomb imaging technique

To cite this article: X Flécharde *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 132047

View the [article online](#) for updates and enhancements.

You may also like

- [A method of false coincidence removal from measurements of quadruple coincidences between two photoelectrons and two photoions generated in molecular double photoionization](#)

Andrew E Slattery, Thomas A Field, Muthanna Ahmad *et al.*

- [Statistical analysis of fragmentation channels of small multicharged molecular ions](#)

C Cornaggia

- [Polarisation-based coincidence event discrimination: an *in silico* study towards a feasible scheme for Compton-PET](#)

M Toghiani, J E Gillam, A L McNamara *et al.*



ECS Membership = Connection

ECS membership connects you to the electrochemical community:

- Facilitate your research and discovery through ECS meetings which convene scientists from around the world;
- Access professional support through your lifetime career;
- Open up mentorship opportunities across the stages of your career;
- Build relationships that nurture partnership, teamwork—and success!

Join ECS!

Visit electrochem.org/join



Investigating the CO trimer geometry using Coulomb imaging technique

X Fléchar¹*, A Méry², B Gervais², S Guillous², V Kumar², J Matsumoto³, J Rangamma², F Ropars²,
H Shiromaru³ and A Cassimi²

¹Normandie Univ, ENSICAEN, UNICAEN, CNRS/IN2P3, LPC Caen, 14000 Caen, France

²CIMAP, CEA-CNRS-ENSICAEN-UNICAEN, Normandie Université, BP5133, F-14050 Caen Cedex 04, France

³Department of Chemistry, Tokyo Metropolitan University, 1-1 Minamiosawa, Hachioji-shi, Tokyo 192-0397, Japan

Synopsis The Coulombic explosion of multi-charged $(\text{CO})_3^{4+}$ molecular ions produced in low energy collisions with Ar^{9+} projectiles was investigated using recoil ion momentum spectrometry (RIMS). A preliminary analysis of the data clearly shows a dominant triangular cyclic structure. Events that may correspond to a linear chain are also observed, but the latter could as well originate from false coincidences, associated to the fragmentation of dimers. A careful analysis of this source of background is proposed to disentangle trimer fragmentation events from false coincidences.

Direct observation of weakly bound structures, such as small molecular clusters, remains an experimental challenge. Infrared (IR) spectroscopy, combined with calculations of intermolecular potential surfaces, has for long been the only way to access such structures. In the last decade, a new technique providing more direct measurements has been developed: by Coulomb exploding the cluster, all momentum vectors of the charged ions or molecular ions can be measured in coincidence, giving access to the initial 3D relative position prior dissociation. A nice illustration of this technique, called “Coulomb imaging”, can be found in [1], where such measurements were performed to determine the geometry of argon and neon dimer, trimer, and tetramer.

Our study focuses on the structure of molecular CO trimers. Few experimental results from IR spectroscopy were made available for this structure, as the $(\text{CO})_3$ transitions were found partly overlapped and obscured by those of $(\text{CO})_2$. The work from Rezaei and collaborators [2] has previously shown a strong evidence for a planar, cyclic and C-bonded structure of the trimer. But satellite peaks and other spectroscopic features remained unexplained. This shortcoming called for a new investigation of the structure of $(\text{CO})_3$ using a completely different and independent technique.

In the present experiment, Coulomb explosion is induced by collisions between a supersonic gas jet, providing the $(\text{CO})_3$ molecular trimers, and Ar^{9+} ions delivered by the low energy beam line of the GANIL/ARIBE facility.

The fragments momenta are measured using a COLTRIMS setup. A clean data selection is first achieved by imposing momentum conservation laws to the fragments resulting from Coulomb explosion. The momentum vectors of the fragments are then further analyzed to infer the geometry of the trimer.




Figure 1. Dalitz plot obtained for the fragmentation of $(\text{CO})_3^{3+}$ in three singly charged $(\text{CO})^+$ fragments.

The plot of Fig. 1 clearly shows two distinct contributions. The dominant one, in the center, can be associated to a regular triangular structure. The weaker contribution, visible on the right edge of the Dalitz plot, could either indicate a linear structure or originate from false coincidences between a dissociating dimer and a monomer. A careful analysis of the background has been performed to quantify this possible contribution.

References

- [1] Ulrich B *et al* 2011 *J. Phys. Chem. A* **115** 6936
- [2] Rezaei M *et al* 2013 *J. Chem. Phys.* **138** 071102

* E-mail: flechar@lpccaen.in2p3.fr

