



HAL
open science

From gut to brain: Microbiota depletion in mice as a tool to explore causality

Sofia Cussotto, Bruno Aouizerate, Lucile Capuron

► **To cite this version:**

Sofia Cussotto, Bruno Aouizerate, Lucile Capuron. From gut to brain: Microbiota depletion in mice as a tool to explore causality. *Brain, Behavior, and Immunity*, 2021, 94, pp.4-5. <10.1016/j.bbi.2021.02.029>. <hal-03347451>

HAL Id: hal-03347451

<https://hal.science/hal-03347451v1>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

From Gut to Brain: Microbiota Depletion in Mice as a Tool to Explore Causality

Sofia Cussotto¹, Bruno Aouizerate^{1,2}, Lucile Capuron^{1*}

¹ University of Bordeaux, INRAE, Bordeaux INP, NutriNeuro, UMR 1286, Bordeaux, France

² CH Charles Perrens, Pôle de Psychiatrie Générale et Universitaire, Centre de référence régional des pathologies anxieuses et dépressives, Bordeaux, France

Invited Commentary Viewpoint for the 2020 BBI Impact Award Winner

(Fröhlich et al. 2016, “Cognitive impairment by antibiotic-induced gut dysbiosis: Analysis of gut microbiota-brain communication”)

Words: 849

Keywords: Microbiota; Gut-Brain Communication ; Antibiotic Treatment ; Cognitive Function ; Mice

*** Corresponding author:**

Dr Lucile Capuron

INRAE 1286, Laboratory of Nutrition and Integrative Neurobiology (NutriNeuro)

University of Bordeaux, 146 rue Léo Saignat

F-33076 Bordeaux, France

Tel. (33) 557571233; Fax. (33) 557571227

Email: lucile.capuron@inrae.fr

The gut microbiota is receiving increasing attention for its involvement in brain function, spanning from cognition to stress response known to occupy a pivotal position in mental health. With the field rapidly expanding, more mechanistic insights are required in order to move beyond associations and examine causal relationships between gut microbiota composition and brain function (Cryan et al. 2019). The paper by Fröhlich and colleagues, that has won the 2020 BBI Impact Award (assigned to the paper published in 2016/2017 that has received most citations in 2018), reports the effects of a short-term antibiotic-induced microbiota depletion on cognitive function in mice (Fröhlich et al. 2016). Following an intragastric administration of an antibiotic mixture, the authors performed a systematic, comprehensive and multi-level assessment of: i) gut microbiota composition, ii) metabolomics patterns in both the colon and plasma, iii) expression of neuronal signaling molecules in different brain areas, and iv) cognitive performance. The authors described a strong antibiotic-mediated effect on microbial composition and cognitive behaviors coupled with abnormal levels of peripheral metabolites and functional changes in neural signaling pathways in the brain.

Germ-free (GF) mice, which are born and raised in sterile conditions, have been instrumental in studying the causal involvement of the gut microbiota in cognitive function. However, they represent a highly artificial model with intrinsic alterations in behavior and neuronal circuits (Heijtz et al. 2011). Short-term antibiotic administration is therefore an appealing alternative, less intrusive and closer to the human setting, nevertheless carrying possible limitations. A recurrent methodological concern that accompanies antibiotic administration involves the pharmacokinetic properties of the antibiotics chosen, as those with high oral bioavailability can enter the circulation and potentially cross the blood-brain barrier (BBB) causing, in turn, a direct central effect (Tochitani et al. 2016). Among the antibiotics used by Fröhlich and colleagues (ampicillin, bacitracin, meropenem, neomycin, vancomycin), only ampicillin is

absorbed to some extent from the human gut. Thus, to exclude the risk of an ampicillin-mediated direct central effects, the authors quantified its levels in the brain. They found them to be below the detection limit, suggesting that the cognitive impairment (disruption of novel object recognition memory) observed in antibiotic-treated mice more likely resulted from gut dysbiosis rather than from a systemic antibiotic response.

Antibiotic administration caused a task-dependent effect, inducing a reduction in memory index (time spent exploring the novel object *vs* known object) in the novel object recognition test (NORT), whilst not affecting spatial learning and memory in the Barnes maze test. These results suggest that gut dysbiosis does not have a generalized effect on cognitive function, but that certain aspects of cognition may be more susceptible to antibiotic-induced alterations in gut microbiota than others. The question arises regarding the possibility to modify the behavioral outcome by changing the composition of the antibiotic mixture and/or route of administration, thus targeting a different range of gut bacteria (or depleting selectively some specific taxa). This important issue opens up an interesting research avenue for further investigation into the influence of specific gut bacteria on distinct components of cognition.

The authors moved on to explore the possible mechanisms involved in the gut-to-brain communication and likely to mediate cognitive alterations. To this end, several cognition-relevant signaling pathways were explored into specific brain areas. Interestingly, the hippocampal and hypothalamic expression of brain-derived neurotrophic factor (BDNF) was decreased and correlated with reduced memory index in the NORT in antibiotic-treated mice. In addition, NPY showed an increased expression in the amygdala and hypothalamus of antibiotic-treated mice, an observation that parallels earlier findings in GF mice (Schéle et al. 2013). Similarly, antibiotic-treated mice exhibited increased expression of the serotonin transporter SLC6A4 and the ionotropic glutamate NMDA receptor N2B subunit in the amygdala. Lastly, circulating levels of corticosterone were elevated in antibiotic-treated mice.

Given that that prolonged exposure to corticosterone was found to dramatically impact brain connectivity, neurotransmitters systems, and cognitive function (McEwen 2002, Lupien et al. 2009), the authors hypothesized that increased corticosterone could play a major role in mediating cognitive deficits in antibiotic-treated mice. Increase in circulating corticosterone could, in turn, contribute to the downregulation of BDNF in the hippocampus and hypothalamus (Naert et al. 2015). Noteworthy, the antibiotic administration did not affect concentrations of central and peripheral cytokines, hence the authors excluded inflammation as potential mechanism responsible for gut-to-brain related cognitive-alterations.

What emerges from the paper here discussed is that antibiotic administration represents a useful pharmacological tool to investigate the causal relationships between changes in gut microbiota, brain function and behavior. Not only the administration of an antibiotic mixture offers chronological flexibility (i.e. the microbiota depletion can be performed acutely or chronically at any stage of the animal's lifespan), but it also permits to titrate the dose of antibiotics and to mimic as closely as possible the clinical setting in humans. Once the experimental model is refined and verified, understanding how gut microbes influence brain function requires a better mechanistic effort into exploring all the possible mediators involved in gut-to-brain communication. The intriguing findings from Fröhlich and colleagues should be considered as a starting point for a translational investigation into the role of gut microbiota in human cognition.

References

- Cryan, J. F., O'Riordan, K. J., Cowan, C. S. M., Sandhu, K. V., Bastiaanssen, T. F. S., Boehme, M., et al. (2019). "The Microbiota-Gut-Brain Axis." Physiol Rev **99**(4): 1877-2013.
- Fröhlich, E. E., Farzi, A., Mayerhofer, R., Reichmann, F., Jačan, A., Wagner, B., et al. (2016). "Cognitive impairment by antibiotic-induced gut dysbiosis: Analysis of gut microbiota-brain communication." Brain Behav Immun **56**: 140-155.
- Heijtz, R. D., Wang, S., Anuar, F., Qian, Y., Björkholm, B., Samuelsson, A., et al. (2011). "Normal gut microbiota modulates brain development and behavior." Proceedings of the National Academy of Sciences **108**(7): 3047-3052.
- Lupien, S. J., McEwen, B. S., Gunnar, M. R. and Heim, C. (2009). "Effects of stress throughout the lifespan on the brain, behaviour and cognition." Nat Rev Neurosci **10**(6): 434-445.
- McEwen, B. S. (2002). "Sex, stress and the hippocampus: allostasis, allostatic load and the aging process." Neurobiol Aging **23**(5): 921-939.
- Naert, G., Zussy, C., Tran Van Ba, C., Chevallier, N., Tang, Y. P., Maurice, T., et al. (2015). "Involvement of Endogenous Brain-Derived Neurotrophic Factor in Hypothalamic-Pituitary-Adrenal Axis Activity." J Neuroendocrinol **27**(11): 850-860.
- Schéle, E., Grahemo, L., Anesten, F., Hallén, A., Bäckhed, F. and Jansson, J. O. (2013). "The gut microbiota reduces leptin sensitivity and the expression of the obesity-suppressing neuropeptides proglucagon (Gcg) and brain-derived neurotrophic factor (Bdnf) in the central nervous system." Endocrinology **154**(10): 3643-3651.
- Tochitani, S., Ikeno, T., Ito, T., Sakurai, A., Yamauchi, T. and Matsuzaki, H. (2016). "Administration of Non-Absorbable Antibiotics to Pregnant Mice to Perturb the Maternal Gut Microbiota Is Associated with Alterations in Offspring Behavior." PLoS One **11**(1): e0138293.