



HAL
open science

Generalized principal eigenvalues of space-time periodic, weakly coupled, cooperative, parabolic systems

Léo Girardin, Idriss Mazari

► **To cite this version:**

Léo Girardin, Idriss Mazari. Generalized principal eigenvalues of space-time periodic, weakly coupled, cooperative, parabolic systems. 2021. hal-03349049v6

HAL Id: hal-03349049

<https://hal.science/hal-03349049v6>

Preprint submitted on 26 Feb 2024 (v6), last revised 7 Jun 2024 (v7)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GENERALIZED PRINCIPAL EIGENVALUES OF SPACE-TIME PERIODIC, WEAKLY COUPLED, COOPERATIVE, PARABOLIC SYSTEMS

LÉO GIRARDIN AND IDRIS MAZARI-FOUQUER

ABSTRACT. This paper is concerned with generalizations of the notion of principal eigenvalue in the context of space-time periodic cooperative systems. When the spatial domain is the whole space, the Krein–Rutman theorem cannot be applied and this leads to more sophisticated constructions and to the notion of generalized principal eigenvalues. These are not unique in general and we focus on a one-parameter family corresponding to principal eigenfunctions that are space-time periodic multiplicative perturbations of exponentials of the space variable. Besides existence and uniqueness properties of such principal eigenpairs, we also prove various dependence and optimization results illustrating how known results in the scalar setting can, or cannot, be extended to the vector setting. We especially prove an optimization property on minimizers and maximizers among mutation operators valued in the set of bistochastic matrices that is, to the best of our knowledge, new.

CONTENTS

1. Introduction	2
1.1. Organization of the paper	2
1.2. Notations	3
1.3. Motivation	4
1.4. Generalized principal eigenvalues in space-time periodic media	5
1.5. Results	6
1.6. Extension to systems with a coupling default	21
1.7. Relation with KPP-type semilinear systems	22
2. Preliminaries	22
2.1. Strong maximum principle	23
2.2. Harnack inequality	23
3. Proofs	27
3.1. Existence, characterization and concavity: proof of Theorems 1.1–1.3	27
3.2. Asymptotic dependence: proof of Theorems 1.5–1.9	40
3.3. Formulas and estimates in special cases: proof of Theorems 1.12–1.16	59
3.4. Optimization: proof of Theorems 1.18–1.20	71
4. Acknowledgements	84

(L. G.) CNRS, ICJ UMR5208, ÉCOLE CENTRALE DE LYON, INSA LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, UNIVERSITÉ JEAN MONNET, 69622 VILLEURBANNE, FRANCE

(I. M.-F.) CEREMADE, UMR CNRS 7534, UNIVERSITÉ PARIS-DAUPHINE, UNIVERSITÉ PSL, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY, 75775 PARIS CEDEX 16, FRANCE

E-mail addresses: leo.girardin@math.cnrs.fr, mazari@ceremade.dauphine.fr.

2010 *Mathematics Subject Classification.* 35K40, 35K57, 47A13, 47A75, 49J20, 92D25.

Key words and phrases. principal eigenvalues, space-time periodicity, cooperative systems.

1. INTRODUCTION

In recent years, the study of principal eigenvalues has proved very fruitful, especially (but not exclusively) for the study of several biological phenomena. Indeed, these eigenvalues encode several informations that are crucial in the understanding of population dynamics; we refer to Section 1.3 below for an overview of the motivations. Although the scalar case is now rather well understood, several problems remain open in the case of systems. In this paper, we propose a systematic approach for the case of space-time periodic cooperative systems, and we offer several contributions to their spectral analysis and optimization.

Formally, this paper is concerned with eigenvalues of linear operators of the form

$$\mathcal{Q} : \mathbf{u} \mapsto \text{diag}(\mathcal{P})\mathbf{u} - \mathbf{L}\mathbf{u},$$

where $\mathbf{u} : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^N$ is a vector-valued function of size $N \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}$, with a time variable $t \in \mathbb{R}$ and a space variable $x \in \mathbb{R}^n$, $n \in \mathbb{N}^*$ being the dimension of the underlying space, $\mathbf{L} : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^{N \times N}$ is a square matrix-valued function of t and x , and each operator of the family $\mathcal{P} = (\mathcal{P}_i)_{i \in [N]}$, where $[N] = \mathbb{N} \cap [1, N]$, has the form

$$\mathcal{P}_i : u \mapsto \partial_t u - \nabla \cdot (A_i \nabla u) + q_i \cdot \nabla u,$$

with $A_i : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times n}$ and $q_i : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ functions of t and x , respectively square matrix-valued and vector-valued. The functions A_i , q_i and \mathbf{L} are smooth and periodic (in a sense made precise later on).

The standing assumptions on \mathcal{P} and \mathbf{L} are the following.

(A₁) The family $(A_i)_{i \in [N]}$ is *uniformly elliptic*:

$$0 < \min_{i \in [N]} \min_{y \in \mathbb{S}^{n-1}} \min_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} (y \cdot A_i(t,x)y).$$

(A₂) The matrix $\underline{\mathbf{L}} \in \mathbb{R}^{N \times N}$, whose entries are

$$\underline{l}_{i,j} = \min_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} l_{i,j}(t,x) \quad \text{for all } (i,j) \in [N]^2,$$

is *essentially nonnegative*: its off-diagonal entries are nonnegative.

(A₃) The matrix $\bar{\mathbf{L}} \in \mathbb{R}^{N \times N}$, whose entries are

$$\bar{l}_{i,j} = \max_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} l_{i,j}(t,x) \quad \text{for all } (i,j) \in [N]^2,$$

is *irreducible*: it does not have a stable subspace of the form $\text{span}(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k})$, where $k \in [N-1]$, $i_1, \dots, i_k \in [N]$ and $\mathbf{e}_i = (\delta_{ij})_{j \in [N]}$. By convention, $[0] = \emptyset$ and 1×1 matrices are irreducible, even if zero.

Under these assumptions, the linear system $\mathcal{Q}\mathbf{u} = \mathbf{0}$ is uniformly parabolic, space-time periodic, weakly coupled (namely, coupled only in the zeroth order term [64]), cooperative and fully coupled (namely, coupled in such a way that there is no independent subsystem [11, 66]).

1.1. Organization of the paper. The remainder of Section 1 is devoted to a detailed introduction (notations, motivation, definitions, main results and applications to semilinear systems). Section 2 is devoted to technical preliminaries. Section 3 contains the proofs.

1.2. **Notations.** We fix once and for all $n+1$ positive numbers $T, L_1, \dots, L_n \in \mathbb{R}_+^*$. For the sake of brevity, we use the notations $L = (L_1, \dots, L_n)$, $(0, L) = (0, L_1) \times \dots \times (0, L_n)$ and $[[0, L]] = \prod_{\alpha=1}^n L_\alpha$. Unless otherwise specified, temporal and spatial periodicities refer to, respectively, T -periodicity with respect to t and L_α -periodicity with respect to x_α for each $\alpha \in [n]$ (or L -periodicity with respect to x for short). The space-time periodicity cell $(0, T) \times (0, L)$ is denoted Ω_{per} and its volume is $T[[0, L]]$.

Vectors in \mathbb{R}^N and matrices in $\mathbb{R}^{N \times N}$ are denoted in bold font. Functional operators are denoted in calligraphic typeface (bold if they act on functions valued in \mathbb{R}^N). Functional spaces, e.g. $\mathcal{W}^{1, \infty}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^N)$, are also denoted in calligraphic typeface. A functional space \mathcal{X} denoted with a subscript \mathcal{X}_{per} , $\mathcal{X}_{t\text{-per}}$ or $\mathcal{X}_{x\text{-per}}$ is restricted to functions that are space-time periodic, time periodic or space periodic respectively.

For clarity, Hölder spaces of functions with $k \in \mathbb{N} \cup \{0\}$ derivatives that are all Hölder-continuous with exponent $\alpha \in (0, 1)$ are denoted $\mathcal{C}^{k+\alpha}$; when the domain is $\mathbb{R} \times \mathbb{R}^n$, it should be unambiguously understood that $\mathcal{C}^{k+\alpha, k'+\alpha'}$ denotes the set of functions that have k α -Hölder-continuous derivatives in time and k' α' -Hölder-continuous derivatives in space.

For any two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^N$, $\mathbf{u} \leq \mathbf{v}$ means $u_i \leq v_i$ for all $i \in [N]$, $\mathbf{u} < \mathbf{v}$ means $\mathbf{u} \leq \mathbf{v}$ together with $\mathbf{u} \neq \mathbf{v}$ and $\mathbf{u} \ll \mathbf{v}$ means $u_i < v_i$ for all $i \in [N]$. If $\mathbf{u} \geq \mathbf{0}$, we refer to \mathbf{u} as *nonnegative*; if $\mathbf{u} > \mathbf{0}$, as *nonnegative nonzero*; if $\mathbf{u} \gg \mathbf{0}$, as *positive*. The sets of all nonnegative, nonnegative nonzero, positive vectors are respectively denoted $[\mathbf{0}, \infty)$, $[\mathbf{0}, \infty) \setminus \{\mathbf{0}\}$ and $(\mathbf{0}, \infty)$. The vector whose entries are all equal to 1 is denoted by $\mathbf{1}$ and this never refers to an indicator function. Similar notations and terminologies might be used in other dimensions and for matrices. The identity matrix is denoted \mathbf{I} .

Similarly, a function can be nonnegative, nonnegative nonzero, positive. For clarity, a positive function is a function with only positive values.

To avoid confusion between operations in the state space \mathbb{R}^N and operations in the spatial domain \mathbb{R}^n , Latin indexes i, j, k are assigned to vectors and matrices of size N whereas Greek indexes α, β, γ are assigned to vectors and matrices of size n . We use mostly subscripts to avoid confusion with algebraic powers, but when both Latin and Greek indexes are involved, we move the Latin ones to a superscript position, e.g. $A_{\alpha, \beta}^i(t, x)$. We denote scalar products in \mathbb{R}^N with the transpose operator, $\mathbf{u}^T \mathbf{v} = \sum_{i=1}^N u_i v_i$, and scalar products in \mathbb{R}^n with a dot, $x \cdot y = \sum_{\alpha=1}^n x_\alpha y_\alpha$.

For any vector $\mathbf{u} \in \mathbb{R}^N$, $\text{diag}(\mathbf{u})$, $\text{diag}(u_i)_{i \in [N]}$ or $\text{diag}(u_i)$ for short refer to the diagonal matrix in $\mathbb{R}^{N \times N}$ whose i -th diagonal entry is u_i . These notations can also be used if \mathbf{u} is a function valued in \mathbb{R}^N .

Finite dimensional Euclidean norms are denoted $|\cdot|$ whereas the notation $\|\cdot\|$ is reserved for norms in functional spaces.

The notation \circ is reserved in the paper for the Hadamard product (component-wise product of vectors or matrices) and never refers to the composition of functions.

Finally, when the focus of the paper is on the dependence of an eigenvalue on (a parameter of) the underlying operator, and when the context is unambiguous, we write with a slight abuse of notation the eigenvalue as a function of the varying parameter (e.g., an eigenvalue λ of the operator \mathcal{Q} might be denoted $\lambda(\mathcal{Q})$, $\lambda(A_1)$, $\lambda(q_1, \dots, q_n)$, $\lambda(\mathbf{L})$, and so on).

1.3. Motivation. The linear parabolic system $\mathcal{Q}\mathbf{u} = \mathbf{0}$ can be understood as the linearization at the homogeneous steady state $\mathbf{0}$ of a semilinear, non-cooperative, space-time periodic, reaction–diffusion system $\text{diag}(\mathcal{P})\mathbf{u}(t, x) = \mathbf{f}(t, x, \mathbf{u}(t, x))$.

From a modeling viewpoint, such systems appear for instance in population dynamics, where each component of \mathbf{u} models a population density for an age class or a phenotypical trait class [22, 23, 30, 35, 42]. In this context, the sign of the principal eigenvalue of the linearization at $\mathbf{0}$ indicates, at least in simple spatio-temporal settings, whether small populations survive and persist or, on the contrary, go extinct.

The population dynamics models we have in mind use a simple growth term of logistic type, also referred to as a Fisher–KPP, or simply KPP, reaction term [31, 49], that satisfies $\mathbf{L}(t, x)\mathbf{u}(t, x) = D_{\mathbf{u}}\mathbf{f}(t, x, \mathbf{0})\mathbf{u} \geq \mathbf{f}(t, x, \mathbf{u})$. Non-cooperative Fisher–KPP systems whose linearization around 0 is cooperative have been the object of a growing literature in the past few years, especially in the case of two components $N = 2$ (see, *e.g.*, [1, 19, 21, 23, 24, 34–37, 39, 41, 44, 45, 56, 58]). It turns out that for such models population persistence is often equivalent to small population persistence, and this makes the study of the principal eigenvalue even more crucial.

When the underlying model is a population structured with respect to a phenotypical trait, then \mathbf{L} typically takes the form $\mathbf{L} = \text{diag}(r_i) + \mathbf{M}$, where each $r_i > 0$ is an intrinsic growth rate and the matrix \mathbf{M} is a mutation matrix; in the simplest case \mathbf{M} is a discrete Laplacian with Neumann boundary conditions:

$$(1) \quad \mathbf{M} = \mu \begin{pmatrix} -1 & 1 & 0 & \dots & 0 \\ 1 & -2 & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & -2 & 1 \\ 0 & \dots & 0 & 1 & -1 \end{pmatrix}$$

where $\mu > 0$ is a mutation rate.

When the underlying model is a population structured with respect to age, then \mathbf{L} is a diagonally perturbed Leslie matrix:

$$(2) \quad \mathbf{L} = -\text{diag}(d_i + a_i) + \begin{pmatrix} b_1 & b_2 & b_3 & \dots & b_N \\ a_1 & 0 & 0 & \dots & 0 \\ 0 & a_2 & 0 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & a_{N-1} & 0 \end{pmatrix}$$

where each $d_i \geq 0$ is a death rate, each $a_i > 0$ an aging rate and each $b_i \geq 0$ a birth rate with $b_N > 0$.

Each one of these models can be understood as a discretized version of some nonlocal equation [35].

The second example (2) above explains in particular why we do not make any *a priori* assumption on the symmetry of \mathbf{L} .

Let us also point out that the periodic cooperative systems we consider find applications in the chemistry of nuclear reactor cores [2, 25]. Due to our long-term goals (see Section 1.7), in this paper, we favor a population dynamics viewpoint.

1.4. Generalized principal eigenvalues in space-time periodic media. In [59], Nadin analyzed the scalar case $N = 1$. Following previous efforts [13, 15, 16, 46], he introduced and studied the following quantities:

$$\lambda_1 = \sup \left\{ \lambda \in \mathbb{R} \mid \exists u \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (0, \infty)) \mathcal{Q}u \geq \lambda u \right\},$$

$$\lambda'_1 = \inf \left\{ \lambda \in \mathbb{R} \mid \exists u \in \mathcal{W}^{1,\infty} \cap \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (0, \infty)) \mathcal{Q}u \leq \lambda u \right\}.$$

These two quantities turn out to be eigenvalues of \mathcal{Q} (in the sense that associated eigenfunctions exist), and are referred to as *generalized principal eigenvalues* (their eigenfunctions are referred to as *generalized principal eigenfunctions*). Due to the lack of compactness in the spatial variable, the existence of these eigenvalues cannot be directly deduced from the Krein–Rutman theorem. However, they can be related with classical Krein–Rutman principal eigenvalues: the first one, λ_1 , is the limit of the principal eigenvalues associated with the time periodic problem with Dirichlet boundary conditions in a sequence of growing balls; the second one, λ'_1 , coincides with the principal eigenvalue of the space-time periodic problem. Actually, both eigenvalues are related to the family $(\lambda_{1,z})_{z \in \mathbb{R}^n}$ of principal eigenvalues of the space-time periodic problems associated with the operators

$$\mathcal{Q}_z : u \mapsto e_{-z} \mathcal{Q}(e_z u) \quad \text{where } e_{\pm z} : x \mapsto e^{\pm z \cdot x},$$

which can be expanded as

$$\mathcal{Q}_z u = \mathcal{Q}u - (A + A^T)z \cdot \nabla u - (z \cdot Az + \nabla \cdot (Az) - q \cdot z)u.$$

Since $\mathcal{Q}(e_z u) = \lambda_{1,z} e_z u$, $\lambda_{1,z}$ can be understood as the principal eigenvalue of \mathcal{Q} acting on the set $e_z \cdot \mathcal{C}_{\text{per}}^{1,2}$ of space-time periodic multiplicative perturbations of the planar exponential e_z . Nadin showed that $\lambda'_1 = \lambda_{1,0} \leq \lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}$ and subsequently exhibited sufficient conditions for the equality $\lambda_1 = \lambda'_1$ to hold; his study is completed by several dependence and optimization results.

Our aim in this paper is twofold. First, we want to generalize the results of Nadin; second, we want to illustrate the originality of systems compared to scalar equations by means of new results and counter-examples without scalar counterpart. Let us point out that most generalizations of scalar results we consider here require work indeed. On one hand, many proofs of [59] rely on algebraic operations that are at least ambiguous, at worst unavailable, in the vector setting, like powers or quotients, and this often leads to counter-examples. On the other hand, the full coupling assumption that we use to emulate the scalar strong comparison principle, (A_3) , is not a pointwise property but rather a global property, and this makes some adaptations quite technical.

Replacing scalar operators and test functions ($N = 1$) by vector ones ($N \in \mathbb{N}^*$), we will therefore study the following quantities:

$$(3) \quad \lambda_1 = \sup \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)) \mathcal{Q}\mathbf{u} \geq \lambda \mathbf{u} \right\},$$

$$(4) \quad \lambda'_1 = \inf \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{W}^{1,\infty} \cap \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)) \mathcal{Q}\mathbf{u} \leq \lambda \mathbf{u} \right\},$$

as well as the family $(\lambda_{1,z})_{z \in \mathbb{R}^n}$, where:

$$(5) \quad \lambda_{1,z} = \lambda_{1,\text{per}}(\mathcal{Q}_z),$$

$$(6) \quad \mathcal{Q}_z = e_{-z} \mathcal{Q}e_z = \mathcal{Q} - \text{diag} \left((A_i + A_i^T)z \cdot \nabla + z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z \right).$$

We admit that the Krein–Rutman theorem can be successfully applied to the operator \mathcal{Q}_z under additional spatial compactness assumptions, so that the *periodic principal eigenvalue* $\lambda_{1,z} = \lambda_{1,\text{per}}(\mathcal{Q}_z)$ is well-defined; similarly, for any nonempty smooth bounded connected open set $\Omega \subset \mathbb{R}^n$, the *Dirichlet principal eigenvalue* $\lambda_{1,\text{Dir}}(\mathcal{Q}_z, \Omega)$ is well-defined. The first one corresponds to the operator \mathcal{Q}_z acting on $\mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n)$, and hereafter we denote \mathbf{u}_z such a positive principal eigenfunction. The second one corresponds to the operator acting on $\mathcal{C}_{t-\text{per}}^{1,2}(\mathbb{R} \times \Omega) \cap \mathcal{C}_0^1(\mathbb{R} \times \bar{\Omega})$, where the subscript 0 denotes functions that vanish on $\partial\Omega$. Eigenfunctions for these principal eigenvalues are unique up to multiplication by a constant. For detailed applications of the Krein–Rutman theory in the Dirichlet case, we refer to Bai–He [11] or Antón–López-Gómez [8].

Let us emphasize once more that the Krein–Rutman theorem cannot be used for λ_1 and λ'_1 . This is why the generalized principal eigenproblem is mathematically challenging and has richer outcomes.

Definition 1.1. A *generalized principal eigenfunction associated with λ_1* is a function $\mathbf{u} \in \mathcal{C}_{t-\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ such that $\mathcal{Q}\mathbf{u} = \lambda_1 \mathbf{u}$.

A *generalized principal eigenfunction associated with λ'_1* is a function $\mathbf{u} \in \mathcal{W}^{1,\infty} \cap \mathcal{C}_{t-\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ such that $\mathcal{Q}\mathbf{u} = \lambda'_1 \mathbf{u}$.

1.5. Results. Although the theorems and definitions in Subsection 1.5.1 are completely analogous to the scalar setting [59], the ones in Subsections 1.5.2–1.5.6, will require new restrictions specific to the parabolic vector setting and will show how the time structure, the spatial structure and the multidimensional state space interact intricately.

Before stating the results, we make the standing assumptions on the smoothness and periodicity of the coefficients of \mathcal{Q} precise.

(A₄) There exists $\delta \in (0, 1)$ such that $\mathbf{L} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ and, for any $i \in [N]$, $A_i \in \mathcal{C}_{\text{per}}^{\delta/2, 1+\delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{n \times n})$ and $q_i \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$. Moreover, $A_i = A_i^T$ for each $i \in [N]$.

We point out that, in such a smooth and generic framework, the symmetry of the diffusion matrices is assumed without loss of generality¹.

1.5.1. *Existence and characterization of generalized principal eigenpairs.*

Theorem 1.1. *The generalized principal eigenvalues λ_1 and λ'_1 are well-defined real numbers related to the family $(\lambda_{1,z})_{z \in \mathbb{R}^n}$:*

$$\lambda'_1 = \lambda_{1,0}, \quad \lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}.$$

The maximum is uniquely achieved.

Consequently, $\lambda'_1 \leq \lambda_1$, \mathbf{u}_0 is a generalized principal eigenfunction associated with λ'_1 and there exists a unique $z^ \in \mathbb{R}^n$ such that $\mathbf{e}_{z^*} \mathbf{u}_{z^*}$ is a generalized principal eigenfunction associated with λ_1 .*

¹Indeed, if A_i is not symmetric, then we can write it as the sum of its symmetric part $A_i^{\text{sym}} = \frac{1}{2}(A_i + A_i^T)$ and its skew-symmetric part $A_i^{\text{skew}} = \frac{1}{2}(A_i - A_i^T)$. The operator $\nabla \cdot (A_i^{\text{skew}} \nabla)$ acting on the space of functions of class \mathcal{C}^2 can be rewritten as an advection operator $a_i \cdot \nabla$, so that $-\nabla \cdot (A_i \nabla) + q_i \cdot \nabla = -\nabla \cdot (A_i^{\text{sym}} \nabla) + (q_i - a_i) \cdot \nabla$. The operator on the right-hand side has the same structure and has “gained” the symmetry of its diffusion matrix.

Furthermore, the following max–min and min–max characterizations hold:

$$\lambda_{1,z} = \max_{\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \min_{i \in [N]} \min_{\Omega_{\text{per}}} \left(\frac{(\mathcal{Q}_z \mathbf{u})_i}{u_i} \right) \quad \text{for all } z \in \mathbb{R}^n,$$

$$\lambda_{1,z} = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \max_{i \in [N]} \max_{\Omega_{\text{per}}} \left(\frac{(\mathcal{Q}_z \mathbf{u})_i}{u_i} \right) \quad \text{for all } z \in \mathbb{R}^n,$$

$$\lambda_1 = \max_{\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \min_{i \in [N]} \inf_{\mathbb{R} \times \mathbb{R}^n} \left(\frac{(\mathcal{Q} \mathbf{u})_i}{u_i} \right).$$

By simplicity of the periodic principal eigenvalue, the only non-negative periodic eigenfunctions are periodic principal eigenfunctions. Under assumptions (A₁)–(A₄), if we further impose standard normalisation conditions on the eigenfunction (*e.g.*, $|\mathbf{u}_z(0, 0)| = 1$), classical compactness estimates on the family $(\lambda_{1,z}, \mathbf{u}_z)$ imply that the spectral elements $(\lambda_{1,z}, \mathbf{u}_z)$ are continuous with respect to the coefficients of \mathcal{Q}_z . In particular, this shows the continuity of λ_1 and λ'_1 as functions of the coefficients of \mathcal{Q} .

Since generalized principal eigenfunctions associated with λ'_1 are globally bounded, a simple comparison argument with the uniformly positive \mathbf{u}_0 shows that it is, up to a multiplicative constant, the unique generalized principal eigenfunction associated with λ'_1 . On the contrary, generalized principal eigenfunctions associated with λ_1 cannot, in general, be compared. The possible existence of generalized principal eigenfunctions for λ_1 that are not of the form $e_{z^*} \mathbf{u}_{z^*}$ remains an open question.

It is well-known that the equality $\lambda'_1 = \lambda_1$ can be false: in the scalar case, the differential operator $u \mapsto -u'' + u'$ is a classical counter-example. The key to this counter-example is the nonzero advection term that moves the maximum of $\lambda_{1,z}$ away from $z = 0$; a similar counter-example of a fully coupled cooperative parabolic system that does not reduce trivially to an elliptic scalar equation is, in spatial dimension $n = 1$, $\mathcal{Q} = \partial_t - \partial_{xx} + \partial_x - (1/8)\mathbf{I} - \mathbf{M}$, where \mathbf{I} is the identity matrix in $\mathbb{R}^{N \times N}$ and \mathbf{M} is the discrete Laplacian defined in (1). By uniqueness of the periodic principal eigenpair and the fact that the coefficients depend neither on time nor space,

$$\lambda_{1,z} = -\lambda_{\text{PF}} \left(-z(1-z)\mathbf{I} + \frac{1}{8}\mathbf{I} + \mathbf{M} \right) = z(1-z) - \frac{1}{8} - \lambda_{\text{PF}}(\mathbf{M}) = z(1-z) - \frac{1}{8},$$

where λ_{PF} denotes the *Perron–Frobenius eigenvalue* of an essentially nonnegative irreducible matrix in $\mathbb{R}^{N \times N}$. Therefore $\lambda'_1 = -1/8 < \lambda_1 = 1/8$, and this also confirms that, as in the scalar case, λ_1 and λ'_1 need not have the same sign.

In the elliptic scalar setting, the absence of advection implies that $z \mapsto \lambda_{1,z}$ is even, whence the equality $\lambda_1 = \lambda'_1$ follows [59, Proposition 3.2]. In the elliptic vector setting, Griette and Matano have very recently proved with a counter-example that this is not the case [40, Proposition 4.1]: the mere asymmetry of $\mathbf{L}(x)$ can induce the strict inequality $\lambda'_1 < \lambda_1$. For the sake of completeness, we recall their counter-example in Remark 3.5.

As in [59], our method of proof actually establishes a few results on λ_1 in arbitrary domains². For any nonempty open connected set $\Omega \subset \mathbb{R}^n$, we define:

$$(7) \quad \lambda_1(\Omega) = \sup \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega}) \quad \mathcal{Q}\mathbf{u} \geq \lambda \mathbf{u} \right\}.$$

Since $\partial\Omega$ is not necessarily smooth, the set $\mathcal{C}^1(\mathbb{R} \times \bar{\Omega})$ is understood here as the set of functions $\mathbf{u} \in \mathcal{C}^1(\mathbb{R} \times \Omega)$ such that both \mathbf{u} and $\nabla \mathbf{u}$ can be continuously extended at any boundary point admitting a strong barrier (see Berestycki–Nirenberg–Varadhan [15]). The subset $\mathcal{C}_0^1(\mathbb{R} \times \bar{\Omega})$ is the set of functions in $\mathcal{C}^1(\mathbb{R} \times \bar{\Omega})$ vanishing continuously at such boundary points.

Definition 1.2. Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set. A *generalized principal eigenfunction associated with $\lambda_1(\Omega)$* is a function $\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}_0^1(\mathbb{R} \times \bar{\Omega})$ such that $\mathcal{Q}\mathbf{u} = \lambda_1 \mathbf{u}$.

Theorem 1.2. Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$. Then the generalized principal eigenvalue $\lambda_1(\Omega)$ is a well-defined real number and there exists an associated generalized principal eigenfunction.

If $\Omega = \mathbb{R}^n$, $\lambda_1(\Omega) = \lambda_1$. If Ω is bounded and smooth, $\lambda_1(\Omega) = \lambda_{1,\text{Dir}}(\Omega)$.

Furthermore, the following max–min characterization holds true:

$$\lambda_1(\Omega) = \max_{\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega})} \min_{i \in [N]} \inf_{\mathbb{R} \times \Omega} \left(\frac{(\mathcal{Q}\mathbf{u})_i}{u_i} \right).$$

1.5.2. *Monotonic or convex dependence with respect to the coefficients.* As an immediate corollary of the max–min characterization of Theorem 1.1, we already know that the eigenvalues $\lambda_{1,z}$, as functions of the matrix entries $l_{i,j}$, are decreasing: if $l_{i,j} < \tilde{l}_{i,j}$ (i.e., $(t, x) \mapsto l_{i,j}(t, x) - \tilde{l}_{i,j}(t, x)$ is a nonnegative nonzero function), then $\lambda_{1,z}(l_{i,j}) > \lambda_{1,z}(\tilde{l}_{i,j})$. This applies in particular to λ_1 and λ'_1 , by virtue of the identifications $\lambda_1 = \max \lambda_{1,z}$ and $\lambda'_1 = \lambda_{1,0}$.

Our first theorem on coefficient dependence is concerned with the concavity of the eigenvalues $\lambda_{1,z}$ as functions of the entries $l_{i,j}$. It generalizes a well-known result by Nussbaum [63] on matrices in $\mathbb{R}^{N \times N}$ as well as a result by Nadin [59] on the scalar parabolic case.

Theorem 1.3. Let $z \in \mathbb{R}^n$ and let

$$(\mathbf{L}[s])_{s \in [0,1]} \in \left(\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N}) \right)^{[0,1]}$$

be a family of matrices satisfying the same assumptions as \mathbf{L} (i.e., (\mathbf{A}_2) , (\mathbf{A}_3)) and such that, for all $(t, x) \in \mathbb{R} \times \mathbb{R}^n$ and $i \in [N]$,

- (1) $s \mapsto l_{i,i}[s](t, x)$ is convex;
- (2) for all $j \in [N] \setminus \{i\}$, $s \mapsto l_{i,j}[s](t, x)$ is either identically zero or log-convex.

Then the map

$$s \in [0, 1] \mapsto \lambda_{1,\text{per}}(\mathcal{Q}_z[s]),$$

where $\mathcal{Q}_z[s]$ is the operator \mathcal{Q}_z with \mathbf{L} replaced by $\mathbf{L}[s]$, is affine or strictly concave.

²In the spirit of Berestycki–Rossi [17], λ'_1 can also be defined in an arbitrary domain Ω and further results on $\lambda_1(\Omega)$ and $\lambda'_1(\Omega)$ are likely achievable. As the focus of this paper is on the influence of space–time periodicity, we do not pursue this direction here.

It is affine if and only if there exist a constant vector $\mathbf{b} \gg \mathbf{0}$, a function $\mathbf{c} \in \mathcal{C}_{\text{per}}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ and a function $\mathbf{f} \in \mathcal{C}_{\text{per}}(\mathbb{R}, \mathbb{R}^N)$ satisfying $\int_0^T \mathbf{f} \in \text{span}(\mathbf{1})$ such that the entries of $\mathbf{L}[s]$ have the form:

$$l_{i,j}[s] : (t, x) \mapsto \begin{cases} l_{i,i}[0](t, x) - s f_i(t) & \text{if } i = j, \\ l_{i,j}[0](t, x) \left(\frac{b_j}{c_i(t, x)} \right)^s e^{s \left(\int_0^t f_j - \frac{t}{T} \int_0^T f_j \right)} & \text{if } i \neq j, \end{cases}$$

and such that the function \mathbf{c} satisfies, at all $(t, x) \in \overline{\Omega_{\text{per}}}$ and for each $i \in [N]$,

$$c_i(t, x) = b_i e^{\int_0^t f_i - \frac{t}{T} \int_0^T f_i} \quad \text{or} \quad \forall j \in [N] \setminus \{i\}, l_{i,j}[0](t, x) = 0.$$

As explained in Remark 3.2, the function \mathbf{c} in the above statement is in general not uniquely determined, but it is so if, for instance, $\mathbf{L}[0]$ is pointwise irreducible.

Although Theorem 1.3 directly applies to $\lambda'_1 = \lambda_{1,0}$, we are only able to prove a weaker concavity property on the generalized principal eigenvalue λ_1 in arbitrary domains – in bounded and smooth domains, a result exactly analogous to Theorem 1.3 applies, see Proposition 3.7. Similarly, in the elliptic case with general spatial heterogeneities in \mathbb{R}^n , Arapostathis–Biswas–Pradhan [9, Lemma 2.3] proved the concavity of λ_1 with respect to the diagonal entries of \mathbf{L} – they did not consider the off-diagonal entries but, their arguments being the same as ours, their result can be extended accordingly.

Theorem 1.4. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$.*

Let

$$(\mathbf{L}[s])_{s \in [0,1]} \in \left(\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \Omega, \mathbb{R}^{N \times N}) \right)^{[0,1]}$$

be a family of matrices satisfying the same assumptions as \mathbf{L} (i.e., (A_2) , (A_3)) and such that, for all $(t, x) \in \mathbb{R} \times \Omega$ and $i \in [N]$,

- (1) $s \mapsto l_{i,i}[s](t, x)$ is convex;
- (2) for all $j \in [N] \setminus \{i\}$, $s \mapsto l_{i,j}[s](t, x)$ is either identically zero or log-convex.

Then the mapping $s \in [0, 1] \mapsto \lambda_1(\Omega, \mathcal{Q}[s])$, where $\mathcal{Q}[s]$ is the operator \mathcal{Q} with \mathbf{L} replaced $\mathbf{L}[s]$, is concave.

Monotonicity or convexity results on the dependence on the diffusion matrices A_i or the advection vectors q_i are in full generality false (in the scalar setting, cases of non-monotonic and non-concave dependence with respect to the diffusion rate are exhibited in Hutson–Mischaikow–Polacik [47]).

1.5.3. Asymptotic dependence with respect to the coefficients. The next theorem shows how the generalized principal eigenvalues $\lambda_{1,z}$ and λ_1 behave close to the boundary where (A_1) , (A_2) and (A_4) remain satisfied but the full coupling assumption (A_3) does not³. We recall that a nonnegative square matrix can be conjugated into a block upper triangular Frobenius normal form by a permutation matrix, with each diagonal block an irreducible nonnegative square matrix (recall that 1×1 matrices are by convention referred to as irreducible even if zero). For a space-time periodic cooperative parabolic operator of the form $\text{diag}(\mathcal{P}) - \mathbf{L}$ but where \mathbf{L} does

³If \mathcal{Q} is spatio-temporally homogeneous, then the theorem reduces to the well-known continuity of the dominant eigenvalue in the set of essentially nonnegative square matrices.

not satisfy (A_3) , conjugating with the permutation matrix associated with the aforementioned Frobenius normal form of the matrix $\bar{\mathbf{L}}$ brings similarly the system into block upper triangular form with each block satisfying (A_3) . Therefore we can assume without loss of generality that the limiting matrix \mathbf{L} is already in block upper triangular form with each block satisfying (A_3) .

Theorem 1.5. *Let $\mathbf{L}^\Delta \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ be a block upper triangular essentially nonnegative matrix. Let $N' \in [N]$ and $(N_k)_{k \in [N'-1]}$ such that*

$$N_0 = 0 < 1 \leq N_1 \leq N_2 \leq \dots \leq N_{N'-1} \leq N_{N'} = N$$

and such that

$$(l_{i,j}^\Delta)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2}$$

is the k -th diagonal block of \mathbf{L}^Δ (with the convention $[0] = \emptyset$). Assume

$$\left(\max_{(t,x) \in \Omega_{\text{per}}} l_{i,j}^\Delta(t,x) \right)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2} \quad \text{is irreducible for all } k \in [N'].$$

Let

$$\mathcal{Q}_k = \text{diag}(\mathcal{P}_i)_{i \in [N_k] \setminus [N_{k-1}]} - (l_{i,j}^\Delta)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2} \quad \text{for all } k \in [N'].$$

Then, as $\mathbf{L} \rightarrow \mathbf{L}^\Delta$ in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$,

$$\lambda_{1,z}(\mathcal{Q}) \rightarrow \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \quad \text{for all } z \in \mathbb{R}^n,$$

$$\lambda_1(\mathcal{Q}) \rightarrow \max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathcal{Q}_k).$$

We comment specifically on this important result in Section 1.6.

The next theorem is concerned with concurrently vanishing diffusion and advection rates – the question of vanishing diffusion rates when the advection rates remain nonnegligible is much more difficult, even in the scalar case [54], and is beyond the scope of this paper; for now, it remains open.

In the statement below, we use the generalized principal eigenvalues of a time periodic weakly coupled linear degenerate parabolic operator, that combines uniformly parabolic equations and ordinary differential equations through cooperative coupling. Such operators satisfy a form of strong comparison principle and, consequently, admit generalized principal eigenvalues $\lambda'_1, \lambda_1, (\lambda_{1,z})_{z \in \mathbb{R}^n}$ that could be studied quite similarly. For the case of bounded domains with homogeneous Dirichlet boundary conditions, we refer to Liang, Zhang and Zhao [51]; the adaptation to space-time periodic settings should be straightforward, following ideas developed in the present paper.

Theorem 1.6. *Let $\mathbf{f} \in \mathcal{C}^1([0, +\infty), [\mathbf{0}, \infty))$ such that $\mathbf{f}^{-1}(\{\mathbf{0}\}) = \{0\}$ and $\mathbf{f}'(0) \neq \mathbf{0}$.*

For all $\varepsilon \geq 0$, let $(q_i^\varepsilon)_{i \in [N]} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$. Assume $(q_i^\varepsilon)_{i \in [N]} \rightarrow (q_i^0)_{i \in [N]}$ in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$ as $\varepsilon \rightarrow 0$.

Let \mathcal{Q}_ε be the operator \mathcal{Q} with $(A_i)_{i \in [N]}$ and $(q_i)_{i \in [N]}$ replaced respectively by $(f_i(\varepsilon)^2 A_i)_{i \in [N]}$ and $(f_i(\varepsilon) q_i^\varepsilon)_{i \in [N]}$.

Denote (with a slight abuse of notation), for every $x \in [0, L]$,

$$(\mathbf{L}(x), (A_i(x))_{i \in [N]}, (q_i(x))_{i \in [N]}) : t \mapsto (\mathbf{L}(t, x), (A_i(t, x))_{i \in [N]}, (q_i(t, x))_{i \in [N]}).$$

Then, for all $z \in \mathbb{R}^n$,

$$\liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda_{1,z}(\mathcal{Q}_\varepsilon) \geq \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right).$$

Furthermore, the equality

$$\lim_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda_{1,z}(\mathcal{Q}_\varepsilon) = \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right).$$

holds true for all $z \in \mathbb{R}^n$ if at least one of the following conditions is satisfied:

- (1) all coefficients of \mathcal{Q}_ε do not depend on x ;
- (2) there exists $\underline{x} \in [0, L]$ such that

$$\lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(\underline{x}) \right) = \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right)$$

and such that the operator

$$\tilde{\mathcal{Q}} = \partial_t - \text{diag}(f'_i(0)^2 \nabla \cdot (A_i(\underline{x}) \nabla) - f'_i(0) q_i^0(\underline{x}) \cdot \nabla) - \mathbf{L}(\underline{x})$$

satisfies

$$\lambda_1(\tilde{\mathcal{Q}}) = \lambda'_1(\tilde{\mathcal{Q}}).$$

The assumptions on \mathbf{f} could be relaxed with a marginal impact on the proof; they are mostly used to simplify the statement of the Theorem and are in any case sufficient for our purposes.

We conjecture that, under reasonable regularity assumptions, the inferior and superior limits of $\lambda_{1,z}(\mathcal{Q}_\varepsilon)$ should always coincide; at this stage, it remains an open question.

The above two conditions that are sufficient to guarantee that

$$\limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda_{1,z}(\mathcal{Q}_\varepsilon) \leq \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right)$$

cover very different cases. However, as explained in Remark 3.10, there also exist cases where the limit of $\lambda_{1,z}(\mathcal{Q}_\varepsilon)$ is well-defined and differs from $\min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right)$. More generally, the problem of characterizing the limit is subtle and, in our opinion, worthy of future attention. It requires a better understanding of the impact of non-vanishing advection rates; indeed, the condition $\lambda_1(\tilde{\mathcal{Q}}) = \lambda'_1(\tilde{\mathcal{Q}})$ is, by virtue of Corollary 1.11, satisfied as soon as $\tilde{\mathcal{Q}}$ contains no advection term, *i.e.* $f'_1(0)q_1^0 = f'_2(0)q_2^0 = \dots = f'_N(0)q_N^0 = 0$.

Beyond pointing out cases where $z \mapsto \lambda_{1,z}$ converges pointwise to a constant in a correctly scaled vanishing diffusion–advection limit⁴, the possible dependence on ε of the advection rates q_i^ε implies that the following two distinct limits are true:

$$\liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1 \left(\partial_t - \text{diag}(\varepsilon^2 \nabla \cdot (A_i \nabla) - \varepsilon q_i \cdot \nabla) - \mathbf{L} \right) \geq \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right),$$

⁴Actually, since $z \mapsto \lambda_{1,z}$ is concave, this convergence is locally uniform in z , but there is really no hope for uniform convergence since $\lambda_{1,z} \rightarrow -\infty$ as $|z| \rightarrow +\infty$. Also, in general, $\lambda_1 = \max \lambda_{1,z}$ does not converge to the same limit: indeed, even for the spatio-temporally homogeneous one-dimensional operator $\partial_t - \varepsilon^2 \partial_{xx} + \varepsilon \partial_x - \mathbf{M}$, with \mathbf{M} the discrete Laplacian defined in (1), the maximum of $z \mapsto \lambda_{1,z}$ is $1/4$, independently of ε , whereas the pointwise limit as $\varepsilon \rightarrow 0$ is 0.

$$\lim_{\substack{\mathbf{d} \rightarrow \mathbf{0} \\ \mathbf{d} \gg \mathbf{0}}} \lambda_1'(\partial_t - \text{diag}(\mathbf{d}) \text{diag}(\nabla \cdot (A_i \nabla) - q_i \cdot \nabla) - \mathbf{L}) = \min_{x \in [0, L]} \lambda_{1, \text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right).$$

Although the two limits look similar, they do not refer to the same underlying questions.

The first one is related to a slowly varying medium. Indeed, assume, for the sake of simplicity, that all L_α coincide and denote $\varepsilon = L_1^{-1}$. Then the change of variable $x \rightarrow \varepsilon x$ changes the $[0, T] \times [0, \varepsilon^{-1}]^n$ -periodic operator \mathcal{Q} into the $[0, T] \times [0, 1]^n$ -periodic operator $\partial_t - \varepsilon^2 \text{diag}(\nabla \cdot (A_i^\varepsilon \nabla) + \varepsilon \text{diag}(q_i^\varepsilon \cdot \nabla) - \mathbf{L}^\varepsilon$, where

$$((A_i^\varepsilon)_{i \in [n]}, (q_i^\varepsilon)_{i \in [N]}, \mathbf{L}^\varepsilon) : (t, x) \mapsto ((A_i)_{i \in [N]}, (q_i)_{i \in [N]}, \mathbf{L}) \left(t, \frac{x}{\varepsilon} \right).$$

In the scalar case, the limit $\varepsilon \rightarrow 0$ has been studied by Hamel, Nadin and Roques [43] in the elliptic case and by Nadin [59] in the parabolic case. In the vector case with temporally homogeneous coefficients and an extra time scaling, it has been studied by Allaire and Hutridurga [2] (parabolic scaling) and by Mirrahimi and Souganidis [57] (hyperbolic scaling).

The second one corresponds for instance to the early stages $t \rightarrow t/\varepsilon$ of a fast-reaction system $\mathbf{L} \rightarrow \frac{1}{\varepsilon} \mathbf{L}$, when spatial processes (dispersal, transport) are still negligible. In bounded domains with Dirichlet boundary conditions, the singular limit $\varepsilon \rightarrow 0$ has been studied by Bai and He [11]. As explained by Lam and Lou in their paper on the Neumann elliptic case [50], the fact that the vanishing parameter is the vector $\mathbf{d} \in \mathbb{R}^N$ and not a scalar amplitude parameter $\varepsilon \in \mathbb{R}$ is meaningful: the spatial processes for one species may be much faster than for the others (*e.g.*, $d_N = \varepsilon \max_{i \in [N-1]} d_i$), as long as they are all slow compared to the parameter ε measuring the time scale and the speed of the reaction.

By considering a uniform limit and advection rates q_i^ε that might vanish as $\varepsilon \rightarrow 0$, we bring together these two frameworks and prove both limits concurrently. We believe this approach is new.

The next theorem is, on the contrary, concerned with how very large diffusion rates impact the periodic principal eigenvalue $\lambda_{1, \text{per}}$. The large diffusivity limit for the whole family $(\lambda_{1, z}(\mathcal{Q}_{\mathbf{d}}))_{z \in \mathbb{R}^n}$ is an entirely different problem, since the large parameter \mathbf{d} appears also in the zeroth order term which will therefore blow-up as soon as z is nonzero⁵. This problem is beyond our scope and is left open.

The question of very large advection rates, already much more delicate in the scalar case [55], is also beyond our scope.

Theorem 1.7. *Let*

$$((\langle A_i \rangle, \langle q_i \rangle)_{i \in [N]}, \langle \mathbf{L} \rangle) : t \mapsto \frac{1}{|[0, L]|} \int_{[0, L]} ((A_i, q_i)_{i \in [N]}, \mathbf{L})(t, x) dx$$

and, for all $\mathbf{d} \in (\mathbf{0}, \infty)$, let $\mathcal{Q}_{\mathbf{d}}$ be the operator \mathcal{Q} with $(A_i)_{i \in [N]}$ replaced by $(d_i A_i)_{i \in [N]}$.

Then, as $\min_{i \in [N]} d_i \rightarrow +\infty$,

$$\lambda_{1, \text{per}}(\mathcal{Q}_{\mathbf{d}}) \rightarrow \lambda_{1, \text{per}} \left(\frac{d}{dt} - \langle \mathbf{L} \rangle \right).$$

⁵More precisely, as will be shown in Corollary 3.13, $-|P_1(z)| \max_{i \in [N]} d_i \leq \lambda_{1, z}(\mathcal{Q}_{\mathbf{d}}) \leq -|P_2(z)| \min_{i \in [N]} d_i$ for some second-order polynomials P_1, P_2 . What would then be relevant would be to figure out an asymptotic expansion of $\lambda_{1, z}(\mathcal{Q}_{\mathbf{d}})$.

This homogenization theorem basically means that, for the periodic principal eigenvalue $\lambda_{1,\text{per}}$, very large diffusion rates tend to replace spatially heterogeneous coefficients by their averages in space. Again, the fact that the multiplicative coefficients d_i can go to $+\infty$ at different rates is meaningful.

In the scalar case [59], the inequality

$$\lambda_{1,\text{per}} \left(\frac{d}{dt} - \langle l_{1,1} \rangle \right) \geq \min_{x \in [0,L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - l_{1,1}(x) \right)$$

holds, and implies a comparison between the large diffusion asymptotic and the vanishing diffusion asymptotic. In the vector case, this inequality is still true if for instance the periodic principal eigenfunction associated with $\partial_t - \mathbf{L}(x)$ depends neither on t nor on x . Indeed, by integrating the equality it satisfies over $[0, L]$,

$$\lambda_{1,\text{per}} \left(\frac{d}{dt} - \langle \mathbf{L} \rangle \right) = \frac{1}{|[0, L]|} \int_{[0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(\cdot) \right) \geq \min_{x \in [0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right).$$

However it is not true in full generality, as shown by the counter-example of Remark 3.12.

1.5.4. *Dependence on the space or time frequencies.* As explained before, assuming that all spatial periods L_α coincide and changing appropriately the variables t and x , the $\overline{\Omega_{\text{per}}}$ -periodic operator \mathcal{Q} becomes the following $[0, 1] \times [0, 1]^n$ -periodic operator:

$$\mathcal{Q}_{T, L_1} = \frac{1}{T} \partial_t - \frac{1}{L_1^2} \text{diag}(\nabla \cdot (A_i^\diamond \nabla)) + \frac{1}{L_1} \text{diag}(q_i^\diamond \cdot \nabla) - \mathbf{L}^\diamond,$$

where

$$\left((A_i^\diamond, q_i^\diamond)_{i \in [N]}, \mathbf{L}^\diamond \right) : (t, x) \mapsto ((A_i, q_i)_{i \in [N]}, \mathbf{L})(Tt, L_1 x).$$

Theorems 1.6 and 1.7 and Remark 3.12 have immediate interpretations in this context, summarized in the following corollary.

However, before stating the corollary, we draw attention on the fact that after such changes of variables, the family of generalized principal eigenvalues $(\lambda_{1,z})_{z \in \mathbb{R}^n}$ is dilated. Indeed, $\lambda_{1,z}(\mathcal{Q}) = \lambda'_1(\mathcal{Q}_z)$ coincides with

$$\lambda_{1, L_1 z}(\mathcal{Q}_{T, L_1}) = \lambda'_1 \left(\mathcal{Q}_{T, L_1} - \text{diag} \left(2 \frac{1}{L_1} A_i^\diamond z \cdot \nabla + z \cdot A_i^\diamond z + \frac{1}{L_1} \nabla \cdot (A_i^\diamond z) - q_i^\diamond \cdot z \right) \right)$$

and not with

$$\lambda_{1,z}(\mathcal{Q}_{T, L_1}) = \lambda'_1 \left(\mathcal{Q}_{T, L_1} - \text{diag} \left(2 \frac{1}{L_1^2} A_i^\diamond z \cdot \nabla + \frac{1}{L_1^2} z \cdot A_i^\diamond z + \frac{1}{L_1^2} \nabla \cdot (A_i^\diamond z) - \frac{1}{L_1} q_i^\diamond \cdot z \right) \right).$$

It is in some sense disappointing that the limit $L_1 \rightarrow +\infty$ in the corollary below is concerned with the wrong family. In fact it is known that the right family requires a different asymptotic analysis. It is outside the scope of the present paper and instead we refer for instance to [43, Proposition 3.1] where the particular case of scalar space periodic elliptic equations is investigated.

Corollary 1.8. *If $q_i^\diamond = 0$ for each $i \in [N]$, then the generalized principal eigenvalues of \mathcal{Q}_{T, L_1} satisfy the following properties.*

(1) *If $L_1 \rightarrow +\infty$, then*

$$\lambda_{1,z}(\mathcal{Q}_{T, L_1}) \rightarrow \min_{x \in [0, 1]^n} \lambda_{1,\text{per}} \left(\frac{1}{T} \frac{d}{dt} - \mathbf{L}^\diamond(x) \right) \quad \text{for all } z \in \mathbb{R}^n.$$

(2) If $L_1 \rightarrow 0$, then

$$\lambda_{1,\text{per}}(\mathcal{Q}_{T,L_1}) \rightarrow \lambda_{1,\text{per}}\left(\frac{1}{T} \frac{d}{dt} - \langle \mathbf{L}^\diamond \rangle\right).$$

(3) There exist a choice of $(A_i^\diamond)_{i \in [N]}$ and \mathbf{L}^\diamond such that $L_1 \mapsto \lambda_{1,\text{per}}(\mathcal{Q}_{T,L_1})$ is decreasing, a choice such that it is constant and a choice such that it is neither.

It remains to investigate the effect of the time frequency $1/T$. In the case of a scalar equation in a bounded domain with Dirichlet boundary conditions, this problem was recently studied thoroughly by Liu, Lou, Peng and Zhou [53]. They identified cases where $T \mapsto \lambda_{1,\text{Dir}}(\mathcal{Q}_{T,L_1})$ is constant, cases where it is decreasing and cases where it is neither; additionally, they studied the asymptotics $T \rightarrow 0$ and $T \rightarrow +\infty$ – reusing the proof of Nadin [59] for the limit $T \rightarrow 0$. Recently, similar results were obtained for the space homogeneous, time periodic system [12]. We will adapt the scalar arguments [53, 59] to prove the following result.

Theorem 1.9. *For all $\omega \in (0, +\infty)$, let \mathcal{Q}_ω be the operator \mathcal{Q} with ∂_t replaced by $\omega \partial_t$. Then:*

(1) if $\omega \rightarrow 0$, then for all $z \in \mathbb{R}^n$,

$$\lambda_{1,z}(\mathcal{Q}_\omega) \rightarrow \frac{1}{T} \int_0^T \lambda_{1,z}(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)) dt,$$

$$\lambda_1(\mathcal{Q}_\omega) \rightarrow \frac{1}{T} \int_0^T \lambda_1(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)) dt,$$

where we denote (with a slight abuse of notation) $((A_i(t), q_i(t))_{i \in [N]}, \mathbf{L}(t)) : x \mapsto ((A_i(t, x), q_i(t, x))_{i \in [N]}, \mathbf{L}(t, x))$;

(2) if $\omega \rightarrow +\infty$, then for all $z \in \mathbb{R}^n$,

$$\lambda_{1,z}(\mathcal{Q}_\omega) \rightarrow \lambda_{1,z}\left(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}}\right),$$

$$\lambda_1(\mathcal{Q}_\omega) \rightarrow \lambda_1\left(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}}\right),$$

where

$$\left((\hat{A}_i, \hat{q}_i)_{i \in [N]}, \hat{\mathbf{L}}\right) : x \mapsto \frac{1}{T} \int_0^T ((A_i, q_i)_{i \in [N]}, \mathbf{L})(t, x) dt.$$

Corollary 1.10. *The generalized principal eigenvalues of \mathcal{Q}_{T,L_1} satisfy the following properties.*

(1) If $T \rightarrow +\infty$, then for all $z \in \mathbb{R}^n$,

$$\lambda_{1,z}(\mathcal{Q}_{T,L_1}) \rightarrow \int_0^1 \lambda_{1,z}\left(-\frac{1}{L_1^2} \text{diag}(\nabla \cdot (A_i^\diamond(t)\nabla)) + \frac{1}{L_1} \text{diag}(q_i^\diamond(t) \cdot \nabla) - \mathbf{L}^\diamond(t)\right) dt.$$

$$\lambda_1(\mathcal{Q}_{T,L_1}) \rightarrow \int_0^1 \lambda_1\left(-\frac{1}{L_1^2} \text{diag}(\nabla \cdot (A_i^\diamond(t)\nabla)) + \frac{1}{L_1} \text{diag}(q_i^\diamond(t) \cdot \nabla) - \mathbf{L}^\diamond(t)\right) dt.$$

(2) If $T \rightarrow 0$, then for all $z \in \mathbb{R}^n$,

$$\lambda_{1,z}(\mathcal{Q}_{T,L_1}) \rightarrow \lambda_{1,z}\left(-\frac{1}{L_1^2} \text{diag}(\nabla \cdot (\hat{A}_i^\diamond \nabla)) + \frac{1}{L_1} \text{diag}(\hat{q}_i^\diamond \cdot \nabla) - \hat{\mathbf{L}}^\diamond\right).$$

$$\lambda_1(\mathcal{Q}_{T,L_1}) \rightarrow \lambda_1 \left(-\frac{1}{L_1^2} \text{diag}(\nabla \cdot (\hat{A}_i^\diamond \nabla)) + \frac{1}{L_1} \text{diag}(\hat{q}_i^\diamond \cdot \nabla) - \hat{\mathbf{L}}^\diamond \right).$$

Theorem 1.9 shows that large time frequencies tend to replace time heterogenous coefficients by their averages in time, whereas small time frequencies tend to replace the parabolic operator by the elliptic operator parametrized by t before averaging the eigenvalue.

1.5.5. *Formulas and estimates in special cases.* Recall the notations \hat{A}_i , \hat{q}_i , $\hat{\mathbf{L}}$ for the averages in time and $\langle A_i \rangle$, $\langle q_i \rangle$, $\langle \mathbf{L} \rangle$ for the averages in space. In this section, for the sake of brevity, we use these notations repeatedly. The combined notation, e.g. $\langle \hat{\mathbf{L}} \rangle$, denotes naturally a space-time average.

As a preliminary, we note that, by simplicity of the periodic principal eigenspace, the generalized principal eigenvalue $\lambda_{1,z}$ can be simplified when coefficients do not depend on space:

$$(8) \quad \lambda_{1,z} = \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L} - \text{diag}(z \cdot A_i z - q_i \cdot z) \right)$$

or when they do not depend on time:

$$(9) \quad \lambda_{1,z} = \lambda_{1,\text{per}} \left(-\text{diag}(\nabla \cdot (A_i \nabla) - (q_i - 2A_i z) \cdot \nabla + (z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z)) - \mathbf{L} \right).$$

When all coefficients are space-time constant, the expression of generalized principal eigenvalue can be simplified even further:

$$(10) \quad \lambda_{1,z} = -\lambda_{\text{PF}}(\mathbf{L} + \text{diag}(z \cdot A_i z - q_i \cdot z)).$$

These reductions to ordinary differential, elliptic partial differential or matrix operators are valid for any $z \in \mathbb{R}^n$. Moreover, when considering only the specific case $z = 0$, the condition of space, time or space-time homogeneity needs to be verified only by \mathbf{L} , and not by $(A_i)_{i \in [N]}$ or $(q_i)_{i \in [N]}$. These reductions will be used repeatedly in the proofs and in the constructions of counter-examples.

In the statements below, the Perron–Frobenius eigenvalue of a reducible matrix is defined by continuous extension of the dominant eigenvalue on the set of essentially nonnegative matrices; for the sake of simplicity, its nonnegative eigenvectors are still referred to as Perron–Frobenius eigenvectors, even though they might not be positive and the eigenvalue might not be simple (algebraically and/or geometrically).

As an immediate consequence of (8) and of the monotonicity of $\lambda_{1,\text{per}}$ with respect to the diagonal entries of \mathbf{L} , we have the following corollary, which shows that in the absence of advection and of spatial heterogeneities, there is no gap between λ_1 and λ'_1 .

Corollary 1.11. *Assume:*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on x ,
- (2) for each $i \in [N]$, $q_i = 0$.

Then $\lambda_1 = \lambda'_1$.

Our next two theorems are concerned with operators whose coefficients depend only on time or only on space, and generalize [59, Propositions 3.1 and 3.2].

Theorem 1.12. *Assume:*

- (1) $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ and \mathbf{L} do not depend on x ,

- (2) there exists a constant positive vector $\mathbf{u} \in (\mathbf{0}, \infty)$ such that \mathbf{u} is a Perron–Frobenius eigenvector of $\mathbf{L}(t)$ for all $t \in \mathbb{R}$.

Let $z \in \mathbb{R}^n$. The equality

$$\lambda_{1,z} = -z \cdot \hat{A}_1 z + \hat{q}_1 \cdot z - \lambda_{\text{PF}}(\hat{\mathbf{L}})$$

is true if $z = 0$ or $(A_1, q_1) = (A_2, q_2) = \dots = (A_N, q_N)$.

Consequently, if:

- (1) $(A_1, q_1) = (A_2, q_2) = \dots = (A_N, q_N)$,
(2) $\hat{q}_1 = 0$,

then

$$\lambda_1 = \lambda'_1 = -\lambda_{\text{PF}}(\hat{\mathbf{L}}).$$

We explain in Remark 3.13 that if the assumption on the existence of a constant positive eigenvector is not satisfied, then the claimed equality on $\lambda_{1,z}$ is false in general. This is striking, since in the scalar setting, the existence of a constant positive eigenvector is not required.

Theorem 1.13. *Assume:*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
(2) $\mathbf{L}(x)$ is symmetric for all $x \in \mathbb{R}^n$,
(3) there exists $z \in \mathbb{R}^n$ and $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ such that $\int_{[0,L]} \nabla Q = 0$ and

$$A_1^{-1} q_1 = A_2^{-1} q_2 = \dots = A_N^{-1} q_N = 2z + \nabla Q.$$

Then

$$\lambda_1 = \lambda_{1,z} = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L}_{Q,z} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2},$$

where

$$\mathbf{L}_{Q,z} = \mathbf{L} + \text{diag} \left(\frac{1}{2} \nabla \cdot (A_i \nabla Q) - \frac{1}{4} \nabla Q \cdot A_i \nabla Q + \nabla \cdot (A_i z) - z \cdot A_i (z + \nabla Q) \right).$$

Furthermore, if there exists a constant positive vector $\mathbf{u} \in (\mathbf{0}, \infty)$ such that \mathbf{u} is a Perron–Frobenius eigenvector of $\mathbf{L}_{Q,z}(x)$ for all $x \in \mathbb{R}^n$, then

$$\lambda_1 = \lambda'_1 \leq -\lambda_{\text{PF}}(\langle \mathbf{L}_{Q,z} \rangle).$$

We will explain in Remark 3.15 that if \mathbf{L} is not symmetric, then, even in the simple case $z = q_1 = q_2 = \dots = 0$, there are counter-examples where

$$\lambda'_1 > \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2}.$$

As noted before, we will also recall in Remark 3.5 the counter-example of Griette–Matano [40] where the mere asymmetry of \mathbf{L} breaks the equality $\lambda_1 = \lambda'_1$.

We will also explain in Remark 3.19 that if the assumption on the existence of a constant positive eigenvector is not satisfied, then the inequality between $-\lambda'_1$ and

the Perron–Frobenius eigenvalue of $\langle \mathbf{L}_{Q,z} \rangle$ can fail. Again, in the scalar case, this assumption is not required [13].

The following theorem is similar in spirit and requires a line-sum-symmetry assumption ($\mathbf{L}\mathbf{1} = \mathbf{L}^T\mathbf{1}$). Examples of line-sum-symmetric essentially nonnegative matrices are doubly stochastic matrices, essentially nonnegative symmetric matrices and essentially nonnegative circulant matrices. For more details on line-sum-symmetric matrices, we refer for instance to Eaves–Hoffman–Rothblum–Schneider [29].

Theorem 1.14. *Assume $\mathbf{L}(t, x)$ is line-sum-symmetric at all $(t, x) \in \overline{\Omega_{\text{per}}}$.*

Let $z \in \mathbb{R}^n$. If, for all $i \in [N]$, $q_i \in C^1_{\text{per}}(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) = 0$, then

$$\lambda_{1,z} \leq -\frac{1}{N} \left(\sum_{i,j=1}^N \langle \hat{l}_{i,j} \rangle + z \cdot \sum_{i=1}^N (\langle \hat{A}_i \rangle z - \langle \hat{q}_i \rangle) \right)$$

and this inequality is an equality if $\mathbf{L} + \text{diag}(\nabla \cdot (A_i z) + z \cdot (A_i z - q_i))$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$ with Perron–Frobenius eigenvector $\mathbf{1}$ and constant Perron–Frobenius eigenvalue.

This theorem has several interesting consequences, which we detail in Corollaries 3.27 and 3.28.

Two similar results without line-sum-symmetry follow.

Theorem 1.15. *Let $z \in \mathbb{R}^n$. If, for all $i \in [N]$, $q_i \in C^1_{\text{per}}(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) = 0$, then*

$$\lambda_{1,z} \leq \lambda_{1,z} \left(\partial_t - \text{diag}(\nabla \cdot (\langle A_i \rangle \nabla) - \langle q_i \rangle) - \mathbf{L}^\# \right),$$

where the entries of the matrix $\mathbf{L}^\# = (l_{i,j}^\#)_{(i,j) \in [N]^2}$ are defined by:

$$l_{i,j}^\# : t \mapsto \begin{cases} \frac{1}{|[0,L]|} \int_{[0,L]} l_{i,i}(t, x) dx & \text{if } i = j, \\ \exp \left(\frac{1}{|[0,L]|} \int_{[0,L]} \ln l_{i,j}(t, x) dx \right) & \text{if } i \neq j \text{ and } \min_{(t,x) \in \overline{\Omega_{\text{per}}}} l_{i,j}(t, x) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 1.16. *Let $z \in \mathbb{R}^n$. If $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ and \mathbf{L} do not depend on x , then*

$$\lambda_{1,z} \leq -\lambda_{\text{PF}} \left(\mathbf{L}^b + \text{diag} \left(z \cdot \hat{A}_i z - \hat{q}_i \cdot z \right) \right),$$

where the entries of the matrix $\mathbf{L}^b = (l_{i,j}^b)_{(i,j) \in [N]^2}$ are defined by:

$$l_{i,j}^b = \begin{cases} \frac{1}{T} \int_0^T l_{i,i} & \text{if } i = j, \\ \exp \left(\frac{1}{T} \int_0^T \ln l_{i,j} \right) & \text{if } i \neq j \text{ and } \min_{t \in [0,T]} l_{i,j}(t) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

The operator introduced in Theorem 1.15 is spatially homogeneous, so that

$$\lambda_{1,z} \left(\partial_t - \text{diag}(\nabla \cdot (\langle A_i \rangle \nabla) - \langle q_i \rangle \cdot \nabla) - \mathbf{L}^\# \right) = \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}^\# - \text{diag}(z \cdot \langle A_i \rangle z - \langle q_i \rangle \cdot z) \right).$$

Therefore the last two theorems can be applied consecutively to find the following corollary.

Corollary 1.17. *Let $z \in \mathbb{R}^n$. If, for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) = 0$, then*

$$\lambda_{1,z} \leq -\lambda_{\text{PF}} \left(\mathbf{L}^{\#b} + \text{diag} \left(z \cdot \langle \hat{A}_i \rangle z - \langle \hat{q}_i \rangle \cdot z \right) \right)$$

where

$$l_{i,j}^{\#b} = \begin{cases} \frac{1}{T|[0,L]|} \int_0^T \int_{[0,L]} l_{i,i} & \text{if } i = j, \\ \exp \left(\frac{1}{T|[0,L]|} \int_0^T \int_{[0,L]} \ln l_{i,j} \right) & \text{if } i \neq j \text{ and } \min_{(t,x) \in \overline{\Omega}_{\text{per}}} l_{i,j}(t,x) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

We emphasize that this upper estimate accounts for off-diagonal entries of \mathbf{L} and is therefore better than the one that could be obtained by writing $\mathbf{L} \geq \text{diag}(l_{i,i})$ and then using the well-known scalar estimate $\lambda_{1,z}(\mathcal{P}_i - l_{i,i}) \leq -\frac{1}{T|[0,L]|} \int_0^T \int_{[0,L]} (l_{i,i} + z \cdot A_i z - z \cdot q_i)$ under the assumption $\nabla \cdot (q_i - 2A_i z) = 0$.

Theorems 1.16 and 1.15 show that when comparing heterogeneous environments with averaged environments, heterogeneities tend to decrease the generalized principal eigenvalues, provided the geometric average is used for the off-diagonal entries of \mathbf{L} . This is of course related to the convexity property of Theorem 1.3. This is also related to the asymptotic results of Theorems 1.7 and 1.9, although in these asymptotics the off-diagonal entries are averaged with the arithmetic mean instead of the geometric mean. By comparing the arithmetic and geometric averages and using the monotonicity of $\lambda_{1,z}$ with respect to \mathbf{L} , we can try to compare these results; however, inequalities are in the wrong sense. For instance, in the simple case $z = 0$ with each q_i divergence-free, what we get is:

$$\begin{aligned} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}^{\#} \right) &\geq \max \left[\lambda'_1(\mathcal{Q}), \lambda_{1,\text{per}} \left(\frac{d}{dt} - \langle \mathbf{L} \rangle \right) \right] \\ &= \max \left[\lambda'_1(\mathcal{Q}), \lim_{\min_{i \in [N]} d_i \rightarrow +\infty} \lambda'_1(\mathcal{Q}_d) \right]. \end{aligned}$$

1.5.6. Optimization. Our first optimization result is a highly nontrivial generalization of a result on matrices of Neumann–Sze [62]. To the best of our knowledge, in the context of cooperative partial differential operators, it is the first time such a result is stated and proved.

Recall that a doubly stochastic matrix $\mathbf{S} \in \mathbb{R}^{N \times N}$ is a nonnegative matrix such that $\mathbf{S}\mathbf{1} = \mathbf{S}^T \mathbf{1} = \mathbf{1}$. Denote $\mathcal{S} \subset \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ the set of all periodic functions whose values are doubly stochastic matrices almost everywhere and $\mathcal{S}_{\{0,1\}}$ the restriction to functions valued in the set of permutation matrices almost everywhere.

A decomposition $\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ of a given essentially nonnegative matrix \mathbf{L} with \mathbf{S} doubly stochastic and $\boldsymbol{\mu}$ nonnegative exists in many cases (see Lemma 3.31). Such a decomposition is not unique: replacing $(\mathbf{S}, \boldsymbol{\mu})$ by $(\mathbf{I} + \gamma(\mathbf{S} - \mathbf{I}), \gamma^{-1} \boldsymbol{\mu})$ with a small $\gamma > 0$ gives another decomposition. The main property of this decomposition is that the so-called mutation part $(\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ admits $\mathbf{1}$ as left Perron–Frobenius eigenvector, with eigenvalue 0. In other words, summing the lines of the system makes the mutations disappear: if the phenotypes do not differ in intrinsic growth rate (all r_i coincide), then the phenotype distribution has no effect on the growth of the meta-population $\sum_{i=1}^N u_i$. This is indeed under this form

that \mathbf{L} appears in several papers on reaction–diffusion models for phenotypically structured populations [23, 41, 58].

Theorem 1.18. *Assume \mathbf{L} has the form*

$$\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$$

with $\mathbf{S} \in \mathcal{S}$, $\mathbf{r} \in \mathcal{L}_{\text{per}}^{\infty}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^N)$ and $\boldsymbol{\mu} \in \mathcal{L}_{\text{per}}^{\infty}(\mathbb{R} \times \mathbb{R}^n, [\mathbf{0}, \infty))$.

Then, for all $z \in \mathbb{R}^n$,

$$\min_{\mathbf{S} \in \mathcal{S}_{\{0,1\}}} \lambda_{1,z}(\mathbf{S}) = \min_{\mathbf{S} \in \mathcal{S}} \lambda_{1,z}(\mathbf{S}) \leq \max_{\mathbf{S} \in \mathcal{S}} \lambda_{1,z}(\mathbf{S}) = \max_{\mathbf{S} \in \mathcal{S}_{\{0,1\}}} \lambda_{1,z}(\mathbf{S}).$$

This theorem does not require the assumption (A_3) (which is not satisfied for some choices of \mathbf{S} ; in such cases, the generalized principal eigenvalues $\lambda_{1,z}$ are defined by continuous extension, see Theorem 1.5). In particular, the set of optimal permutation matrices might *a priori* be reduced to the singleton $\{\mathbf{I}\}$. Also, in this theorem, and as usual in optimization problems, we consider \mathcal{L}^{∞} constraints on \mathbf{S} instead of Hölder-continuity constraints; the optimizers might be for instance “bang-bang” discontinuous piecewise-constant functions. Let us also point out that, as explained in Remark 3.24, the result remains true with any more general decomposition $\mathbf{L} = \mathbf{B} + \mathbf{S}\mathbf{A}$ with \mathbf{A} nonnegative and \mathbf{B} essentially nonnegative.

The modeling viewpoint on this result is natural and enlightening. Say we want to optimize the chances of, for instance, survival of a population, and, for simplicity, that the environment is homogeneous; the phenotypes are labelled as follows: u_1 is the best phenotype when there are no mutations, u_2 is the second best phenotype, and so forth. Intuitively we should select a (reducible) mutation strategy such that the type u_1 is 100% heritable. Thus the first column of \mathbf{S} should be \mathbf{e}_1 . Since \mathbf{S} is doubly stochastic, its first line is then \mathbf{e}_1^T , whence the first phenotype is in fact completely isolated from the others. Subsequently, whatever the mutation strategy for the phenotypes u_2, u_3, \dots , is, the periodic principal eigenvalue is optimal and equal to the periodic principal eigenvalue of the scalar equation satisfied by u_1 . If u_2 is just as good as u_1 , then similarly the pair $\{u_1, u_2\}$ has to be isolated, but apart from this restriction the two blocks of \mathbf{S} can be chosen freely, and in particular they can have the form of permutation matrices. The extension of this intuition to spatio-temporally heterogeneous environments explains why the optimal \mathbf{S} is not in general constant; it has to “switch” as soon as the optimal family of phenotypes changes.

Let us stress that although the set of doubly stochastic matrices is the convex hull of the set of permutation matrices (a classical result known as the Birkhoff–von Neumann theorem), $\mathbf{S} \in \mathcal{S} \mapsto \lambda_{1,z}(\mathbf{S})$ is not concave (see Theorem 1.3), so that Theorem 1.18 does not follow from mere convexity considerations. Let us also stress that as soon as all (\mathcal{P}_i, r_i) coincide with constant r_i , $\mathbf{S} \mapsto \lambda_{1,z}(\mathbf{S})$ is constant: maximizers and minimizers need not be in $\mathcal{S}_{\{0,1\}}$ and can coincide.

The proof of Theorem 1.18 is in fact quite involved and requires the construction of an explicit rank-one perturbation of \mathbf{S} .

Our second optimization result, closely related to Theorem 1.18, generalizes a theorem due to Karlin and later generalized by Altenberg [3, 48] which states that, for any irreducible stochastic matrix \mathbf{S} and any diagonal matrix \mathbf{D} with positive diagonal entries, the mapping $\tau \in [0, 1] \mapsto \lambda_{\text{PF}}(((1 - \tau)\mathbf{I} + \tau\mathbf{S})\mathbf{D})$ is nonincreasing. The Karlin theorem has been interpreted as “greater mixing yields slower growth”

and shows how, in a space-time homogeneous setting, mutations reduce the chances of survival.

Theorem 1.19. *Assume $(A_i)_{i \in [N]}$ is independent of t , $(q_i)_{i \in [N]} = 0$, and \mathbf{L} has the form $\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ with $\mathbf{r} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R}^n, \mathbb{R}^N)$, $\boldsymbol{\mu} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R}^n, (\mathbf{0}, \infty))$ and $\mathbf{S} \in \mathcal{S}$ all independent of t .*

For any $\rho > 0$, let \mathcal{Q}_ρ be the operator with $(A_i)_{i \in [N]}$ and \mathbf{L} replaced by $(\rho A_i)_{i \in [N]}$ and $\text{diag}(\mathbf{r}) + \rho(\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ respectively.

Then $\rho \in [0, 1] \mapsto \lambda'_1(\mathcal{Q}_\rho)$ is concave and nondecreasing. Furthermore, if \mathbf{r} depends on x and $s > 0$, then it is strictly concave and increasing.

Consequently, λ'_1 is maximized at $\rho = 1$ and minimized at $\rho = 0$: “greater mutation+diffusion yields slower growth”.

We emphasize that the main interest of Theorem 1.19 is that it does not require the symmetry of the mutation matrix $(\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$. When it is symmetric, the variational formula of Theorem 1.13 can be used to deduce a stronger result, the concavity and monotonicity with respect to the diffusion rate on one hand and to the mutation rate on the other hand, with no need to couple the two rates.

Our last optimization result deals with the spatial distribution in the matrix \mathbf{L} in one dimension of space. In this context, the spatial periodicity cell is then the interval $(0, L_1)$. Our result is a generalization of a result by Nadin [59] and makes use of the periodic rearrangement. We recall that for any scalar L_1 -periodic function u there exists a unique L_1 -periodic function u^\dagger whose restriction to $[0, L_1]$ is symmetric (with respect to the midpoint $L_1/2$) and non-increasing in $[L_1/2, L_1]$ and that has the same distribution function as u . The distribution function of u is:

$$\mu_u : t \mapsto |\{u \geq t\} \cap [0, L_1]|.$$

For a time dependent scalar function u , u^\dagger stands for the function rearranged, at every t , with respect to x . For a time-dependent, vector (respectively matrix) valued function \mathbf{u} , the notation \mathbf{u}^\dagger is understood as the vector-valued function with i -th (resp. (i, j) -th) component u_i^\dagger (resp. $u_{i,j}^\dagger$).

Theorem 1.20. *Assume $n = 1$ and $\text{diag}(\mathcal{P}) = \partial_t - \mathbf{D}\Delta$ for some diagonal matrix \mathbf{D} with constant, positive diagonal entries.*

Then

$$\lambda_{1,\text{per}}(\mathcal{Q}) \geq \lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - \mathbf{L}^\dagger)$$

where \mathbf{L}^\dagger is the entry-wise periodic rearrangement of \mathbf{L} .

Note that this theorem optimizes the distribution of each $l_{i,j}$ but does not optimize the distribution of mass in the matrix \mathbf{L} . Brenier [20] showed that the rearrangement of a function of x and the polar decomposition of an invertible matrix in $\mathbb{R}^{N \times N}$ are related notions, via the relations $\mu(x) = \mu^\#(u(x))$ ($\mu^\#$ is the spatial rearrangement, u is unitary) and $\mathbf{M} = \mathbf{R}\mathbf{U}$ ($\mathbf{R} = (\mathbf{M}\mathbf{M}^\text{T})^{\frac{1}{2}}$ is symmetric positive definite, \mathbf{U} is orthogonal). In particular, it is well-known that, similarly to $\lambda_{1,\text{per}}(-\Delta - \mu^\#) \leq \lambda_{1,\text{per}}(-\Delta - \mu)$, any essentially nonnegative matrix $\mathbf{M} \in \mathbb{R}^{N \times N}$ satisfies $\lambda_{\text{PF}}(\mathbf{M}) \leq \lambda_{\text{max}}((\mathbf{M}\mathbf{M}^\text{T})^{\frac{1}{2}})$, where λ_{max} denotes the maximal eigenvalue of a real symmetric matrix. In other words, if \mathcal{Q} has only constant coefficients,

$$\lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - (\mathbf{L}^\text{T}\mathbf{L})^{\frac{1}{2}}) \leq \lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - \mathbf{L}),$$

where the periodic principal eigenvalue $\lambda_{1,\text{per}}$ on the left-hand side is defined via the spectral theorem for self-adjoint compact operators instead of via the Krein–Rutman theorem – the matrix $(\mathbf{L}^T \mathbf{L})^{\frac{1}{2}}$ is not, in general, essentially nonnegative. However the proofs of $\lambda_{1,\text{per}}(-\Delta - \mu^\#) \leq \lambda_{1,\text{per}}(-\Delta - \mu)$ and of $\lambda_{\text{PF}}(\mathbf{M}) \leq \lambda_{\max}((\mathbf{M}\mathbf{M}^T)^{\frac{1}{2}})$ differ strongly. The first one typically uses the Hardy–Littlewood inequality, which is false for matrices as showed by Brenier [20]. Therefore it seems that optimizing \mathbf{L} in the spatial sense and in the matrix sense simultaneously is much more difficult and we leave it as a very interesting open problem.

1.6. Extension to systems with a coupling default. Theorem 1.5 shows how results on fully coupled cooperative systems (and especially the results of Subsections 1.5.1–1.5.6) can be applied to more general cooperative systems, that need not satisfy (\mathbf{A}_3) , by understanding them as networks of fully coupled subsystems. It also shows that such a perspective is limited regarding λ_1 , as we are now going to explain.

Recall that the Perron–Frobenius eigenvalue λ_{PF} can be understood as the restriction to the set of irreducible essentially nonnegative matrices of the dominant eigenvalue, which is a well-defined continuous mapping from the set of essentially nonnegative matrices to \mathbb{R} . Therefore it is natural to suggest the following extension of the generalized principal eigenvalues $\lambda_{1,z}$ and $\lambda_1(\Omega)$:

$$\begin{aligned}\lambda_{1,z}(\mathcal{Q}) &= \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) = \min_{k \in [N']} \lambda_{1,\text{per}}(e_{-z} \mathcal{Q}_k e_z), \\ \lambda_1(\mathcal{Q}, \Omega) &= \min_{k \in [N']} \lambda_1(\mathcal{Q}_k, \Omega),\end{aligned}$$

where \mathcal{Q}_k denotes as in the statement of Theorem 1.5 the k -th fully coupled block of $\mathcal{Q} = \text{diag}(\mathcal{P}) - \mathbf{L}^\Delta$ in block upper triangular form. With these definitions, Theorem 1.5 shows that the extension of each $\lambda_{1,z}$, and in particular that of λ_1' , is continuous. However, as explained in Remark 3.9, the inequality

$$\lim_{\mathbf{L} \rightarrow \mathbf{L}^\Delta} \lambda_1(\mathcal{Q}) = \max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathcal{Q}_k)$$

is in some cases strict: the extension of λ_1 suggested above is not lower semi-continuous, and *a fortiori* not continuous.

It might be tempting to think that this discontinuity is caused by a wrong choice of generalized definition, and that the correct choice should be continuous. For instance, defining λ_1 as $\max_{z \in \mathbb{R}^n} \lambda_{1,z}$ would give a continuous extension to systems with a coupling default. In view of the literature [9, 15, 16, 59], it is also natural to consider the original definition (3) of λ_1 , and since the coupling default induces a weaker maximum principle, it is also natural to consider a relaxed definition with nonnegative nonzero super-solutions instead of positive super-solutions. In order to compare these quantities, let us denote them as follows:

$$\begin{aligned}\lambda_1^0 &= \min_{k \in [N']} \lambda_1(\mathcal{Q}_k), \\ \lambda_1^1 &= \max_{z \in \mathbb{R}^n} \lambda_{1,z}, \\ \lambda_1^2 &= \sup \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)) \ \mathcal{Q}\mathbf{u} \geq \lambda\mathbf{u} \right\}, \\ \lambda_1^3 &= \sup \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n), [\mathbf{0}, \infty), \ \mathbf{u} \neq \mathbf{0}, \ \mathcal{Q}\mathbf{u} \geq \lambda\mathbf{u} \right\}.\end{aligned}$$

Then we can show⁶ that

$$\lambda_1^1 \leq \lambda_1^0 \leq \max_{k \in [N']} \lambda_1(\mathcal{Q}_k) = \lambda_1^3, \quad \lambda_1^2 \leq \lambda_1^0.$$

The inequality $\lambda_1^0 \leq \lambda_1^3$ is strict as soon as two $\lambda_1(\mathcal{Q}_k)$ differ. The equality $\lambda_1^0 = \lambda_1^2$ can be verified if \mathcal{Q} is block diagonal; although the proof seems to require some work, we believe that it remains true even if \mathcal{Q} is not block diagonal. In any case, since the counter-example of Remark 3.9 is block diagonal, there are block diagonal operators \mathcal{Q} such that $\lambda_1^1 < \lambda_1^0 = \lambda_1^2 < \lambda_1^3$. This shows that reasonable definitions of λ_1 other than λ_1^1 cannot be continuous as (A_3) ceases to be true.

Let us point out that λ_1^1 is indisputably the least natural definition. In particular, having in mind that $\lambda_1 < 0$ should be a criterion for population growth (see Section 1.7 below), then the natural definitions would be either λ_1^0 (growth of at least one population) or λ_1^3 (growth of all populations). In both cases, the default of lower semi-continuity means that populations with vanishingly small couplings might have much stronger chances than decoupled populations. This has strong implications for modeling, as simplifying a vanishingly coupled model into a decoupled one is often tempting. It has been related to the emergence in eco-evolutionary models of unexpectedly large spreading speeds in the vanishing mutation limit. We refer to Elliott–Cornell [30] for the first formal calculations and to Morris–Börger–Crooks [58] for the rigorous analysis.

1.7. Relation with KPP-type semilinear systems. In the scalar framework of KPP-type reaction–diffusion equations, $\lambda_1 < 0$ implies the locally uniform convergence of all solutions to the unique periodic and uniformly positive entire solution, whereas $\lambda_1 \geq 0$ implies the uniform convergence of all solutions to 0, as proved by Nadin [61]. The study of entire solutions is much more delicate in the multidimensional setting, simply due to topological freedom [34,37,39,58], and their uniqueness and stability properties cannot in general be inferred from the linearization at $\mathbf{0}$. However, we will show in a sequel [38] that in the multidimensional case, the results of Nadin [61] can be generalized in the following weak form: $\lambda_1 < 0$ implies the locally uniform persistence of all solutions and the existence of a periodic and uniformly positive entire solution, whereas $\lambda_1 \geq 0$ implies the uniform convergence of all solutions to $\mathbf{0}$.

Going toward these results is one of our main motivations for the present work, the other one being the future construction of pulsating traveling waves [60].

2. PRELIMINARIES

Many of our proofs will use a strong maximum principle and a Harnack inequality for parabolic cooperative systems. These already exist in the literature under slightly different forms (we refer for instance to [11,32,64] or to [7,10,18,26–28,66] for the elliptic case). For the sake of self-containment and because the parabolic Harnack inequality in [32] is insufficient for our purposes, in this section, we state or prove what we need afterward.

⁶The proof is voluntarily not detailed, for the sake of brevity.

2.1. Strong maximum principle. The strong maximum principle for time periodic nonnegative solutions of $\mathcal{Q}\mathbf{u} + K\mathbf{u} = \mathbf{0}$ with large $K > 0$ (actually, $K > \lambda_1$) is established as a side result of the preparation of the application of the Krein–Rutman theorem, just as in Bai–He [11]. In fact, we can repeat the argument of [11, p. 9882] to obtain the strong maximum principle for all values of $K \in \mathbb{R}$, including $K = 0$ (large values of K are required only for the inversion of the operator), and for super-solutions that might not be time periodic but are well-defined in a sufficiently distant past. For clarity, we state this version of the strong maximum principle below.

Proposition 2.1 (Strong maximum principle). *Let $\mathbf{u} \in \mathcal{C}^{1,2}((0, +\infty) \times \mathbb{R}^n, [\mathbf{0}, \infty)) \cap \mathcal{C}([0, +\infty) \times \mathbb{R}^N)$ such that $\mathcal{Q}\mathbf{u} \geq \mathbf{0}$ in $(0, +\infty) \times \mathbb{R}^N$.*

If there exist $t^ > T$, $x^* \in \mathbb{R}^n$ and $i^* \in [N]$ such that $u_{i^*}(t^*, x^*) = 0$, then $\mathbf{u} = \mathbf{0}$ in $[0, +\infty) \times \mathbb{R}^N$.*

A similar property is satisfied in bounded domains. For the sake of simplicity, we only consider smooth boundaries.

Proposition 2.2 (Strong maximum principle in bounded domains). *Let $\Omega \subset \mathbb{R}^n$ be a nonempty smooth bounded open connected set and $\mathbf{u} \in \mathcal{C}^{1,2}((0, +\infty) \times \Omega, [\mathbf{0}, \infty)) \cap \mathcal{C}^{0,1}([0, +\infty) \times \bar{\Omega})$ such that $\mathcal{Q}\mathbf{u} \geq \mathbf{0}$ in $(0, +\infty) \times \Omega$.*

Assume that there exists $x_0 \in \Omega$ such that $[x_0, x_0 + L] \subset \Omega$.

If there exist $t^ > T$, $x^* \in \Omega$ and $i^* \in [N]$ such that $u_{i^*}(t^*, x^*) = 0$, then $\mathbf{u} = \mathbf{0}$ in $[0, +\infty) \times \bar{\Omega}$.*

If there exist $t^ > T$, $x^* \in \partial\Omega$ and $i^* \in [N]$ such that $u_{i^*}(t^*, x^*) = \nu(x^*) \cdot \nabla u_{i^*}(t^*, x^*) = 0$, where $\nu(x^*) \in \mathbb{R}^n$ is the outward pointing unit normal vector, then $\mathbf{u} = \mathbf{0}$ in $[0, +\infty) \times \bar{\Omega}$.*

These versions of the strong maximum principle exploit the full coupling assumption (A_3) : if one component of \mathbf{u} is zero, then so are the others. Nonnegative super-solutions are either zero or positive. Without (A_3) , this alternative is false in general; we refer, for weaker statements applicable to general cooperative systems, to the celebrated book by Protter and Weinberger [64, Chapter 3, Section 8].

2.2. Harnack inequality. In this section, we denote by $\sigma > 0$ the smallest positive entry of $\bar{\mathbf{L}}$ and by $K \geq 1$ the smallest positive number such that

$$\begin{aligned} K^{-1} &\leq \min_{i \in [N]} \min_{y \in \mathbb{S}^{n-1}} \min_{(t,x) \in \bar{\Omega}_{\text{per}}} (y \cdot A_i(t, x)y), \\ \max_{i \in [N]} \max_{y \in \mathbb{S}^{n-1}} \max_{(t,x) \in \bar{\Omega}_{\text{per}}} (y \cdot A_i(t, x)y) &\leq K, \\ \max_{i \in [N]} \max_{\alpha \in [n]} \max_{(t,x) \in \bar{\Omega}_{\text{per}}} |q_{i,\alpha}(t, x)| &\leq K, \\ \max_{i,j \in [N]} \sup_{(t,x) \in \bar{\Omega}_{\text{per}}} |l_{i,j}(t, x)| &\leq K. \end{aligned}$$

Applying Földes–Poláčik’s Harnack inequality [32, Theorem 3.9] to the operator \mathcal{Q} , we obtain the following property.

Proposition 2.3. *Let $\theta > 0$. Assume the irreducibility of the matrix*

$$\underline{\mathbf{L}} = \left(\min_{(t,x) \in \bar{\Omega}_{\text{per}}} l_{i,j}(t, x) \right)_{(i,j) \in [N]^2}$$

and denote $\eta > 0$ its smallest positive entry.

There exists a constant $\bar{\kappa}_{\theta, \eta} > 0$, determined only by n, N, η, K and the parameter θ such that, if $\mathbf{u} \in \mathcal{C}([-2\theta, 6\theta] \times [-\frac{3\theta}{2}, \frac{3\theta}{2}]^n, [\mathbf{0}, \infty))$ is a solution of $\mathbf{Q}\mathbf{u} = \mathbf{0}$, then

$$\min_{i \in [N]} \min_{(t, x) \in [5\theta, 6\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t, x) \geq \bar{\kappa}_{\theta, \eta} \max_{i \in [N]} \max_{(t, x) \in [0, 2\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t, x).$$

However, our irreducibility assumption (A_3) is concerned with the matrix

$$\bar{\mathbf{L}} = \left(\max_{(t, x) \in \Omega_{\text{per}}} l_{i, j}(t, x) \right)_{(i, j) \in [N]^2}$$

and not with $\underline{\mathbf{L}}$. By continuity and essential nonnegativity, $\bar{\mathbf{L}}$ is irreducible if and only if

$$(T|[0, L]|)^{-1} \int_{\Omega_{\text{per}}} \mathbf{L}(t, x) dt dx$$

is itself irreducible. Hence we can understand the assumption (A_3) as “ $\mathbf{L}(t, x)$ is irreducible on average”. It is known that such an assumption is sufficient, and in some sense necessary, for full coupling of the parabolic or elliptic operator; refer, for instance, to [10, 11, 18, 66].

Since Földes–Poláčik’s Harnack inequality requires the pointwise irreducibility of \mathbf{L} , which is a much stronger assumption than the irreducibility on average (there are simple examples of matrices that are irreducible on average but reducible pointwise at all (t, x) , see for instance Remark 3.13), it is not satisfying for our purposes. Actually, going through the proof of [32, Theorem 3.9], it appears that its adaptation to our setting is not straightforward, as Földes and Poláčik overcome the key obstacle by constructing a nonnegative nonzero sub-solution smaller than η multiplied by some positive constant. Nevertheless, since (A_3) is known to be the optimal assumption for full coupling, it is natural to expect a similar Harnack inequality to hold, provided the parabolic cylinder under consideration is sufficiently larger than the periodicity cell Ω_{per} . This is what we prove below, drawing inspiration from the elliptic case studied in Araposthathis–Ghosh–Marcus [10].

By convenience for future use, we state the result for a zeroth order, diagonal, non-necessarily periodic perturbation of \mathbf{Q} . The diffusion and advection terms can be perturbed similarly if needed.

Proposition 2.4 (Fully coupled Harnack inequality). *Let $\theta \geq \max(T, L_1, \dots, L_n)$ and $\mathbf{f} \in \mathcal{L}^\infty \cap \mathcal{C}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^N)$ with $\delta \in (0, 1)$. Let $F > 0$ such that*

$$\max_{i \in [N]} \sup_{(t, x) \in \mathbb{R} \times \mathbb{R}^n} |f_i(t, x)| \leq F.$$

There exists a constant $\bar{\kappa}_{\theta, F} > 0$, determined only by n, N, σ, K and the parameters θ and F such that, if $\mathbf{u} \in \mathcal{C}([-2\theta, 6\theta] \times [-\frac{3\theta}{2}, \frac{3\theta}{2}]^n, [\mathbf{0}, \infty))$ is a solution of $\mathbf{Q}\mathbf{u} = \text{diag}(\mathbf{f})\mathbf{u}$, then

$$\min_{i \in [N]} \min_{(t, x) \in [5\theta, 6\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t, x) \geq \bar{\kappa}_{\theta, F} \max_{i \in [N]} \max_{(t, x) \in [0, 2\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t, x).$$

Proof. Define, for all $i \in [N]$, the $n + 1$ -dimensional hypercube

$$Q_i = \left(5\theta - \frac{\theta}{2^{i-1}}, 6\theta \right) \times \left(-\frac{\theta}{2} - \frac{\theta}{2^i}, \frac{\theta}{2} + \frac{\theta}{2^i} \right)^n \subset \mathbb{R} \times \mathbb{R}^n.$$

Note the series of compact inclusions

$$Q_1 = (4\theta, 6\theta) \times (-\theta, \theta)^n \supset Q_2 \supset \cdots \supset Q_N \supset (5\theta, 6\theta) \times \left(-\frac{\theta}{2}, \frac{\theta}{2}\right)^n.$$

Following carefully the proof of Földes–Poláčik’s Harnack inequality [32], we observe that we only have to prove the following claim.

Claim 1: let $k \in [N - 1]$. If there exists $I \subset [N]$ of cardinal k and a positive constant κ_k determined only by $k, n, N, \sigma, K, \theta$ and F , such that, for all $j \in I$,

$$\min_{(t,x) \in \overline{Q_k}} u_j(t,x) \geq \kappa_k \max_{(t,x) \in [0,2\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_1(t,x),$$

then there exists $i \in [N] \setminus I$ and a positive constant $\kappa_{k+1} \leq \kappa_k$ determined only by $k, n, N, \sigma, K, \theta$ and F , such that

$$\min_{(t,x) \in \overline{Q_{k+1}}} u_i(t,x) \geq \kappa_{k+1} \max_{(t,x) \in [0,2\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_1(t,x).$$

We prove first the following simpler claim, inspired by [10, Lemma 3.6].

Claim 2: Let $k \in [N - 1]$, $i \in [N]$ and $g \in \mathcal{C}(\overline{Q_k}, [0, +\infty))$. There exists a positive constant C_k determined only by k, n, K, θ and F , such that, if u is a solution of $\mathcal{P}_i u - l_{i,i} u - f_i u = g$ in Q_k with $u = 0$ on the parabolic boundary $\partial_P Q_k$, then

$$\min_{(t,x) \in \overline{Q_{k+1}}} u(t,x) \geq C_k \max_{(t,x) \in \overline{Q_k}} g(t,x).$$

Proof of Claim 2. When $g = 0$, $u = 0$ as well and the result is obvious (with, say, $C_k = 1$). Therefore we assume without loss of generality that $g > 0$.

Up to dividing u by $\max_{\overline{Q_k}} g$, we assume without loss of generality $\max_{\overline{Q_k}} g = 1$. Since the solution u of the Cauchy–Dirichlet problem with zero data on the parabolic boundary is unique, we only have to prove that this solution has a positive minimum in $\overline{Q_{k+1}}$, and that the infimum of these minima, when \mathcal{Q}, f and g vary in the correct class, is still positive.

The nonnegativity of u is a direct consequence of the (weak) maximum principle. The positivity of its minimum in $\overline{Q_{k+1}}$ is a consequence of the strong maximum principle and the fact that 0 cannot be the solution.

Now, define \mathcal{U}_k as the set of all $U = (A, q, l, f, g)$ such that

$$A \in \mathcal{C}^{\delta/2, 1+\delta}(\overline{Q_k}, \mathbb{R}^{n \times n}),$$

$$(q, l, f, g) \in \mathcal{C}^{\delta/2, \delta}(\overline{Q_k}, [-K, K]^n \times [-K, K] \times [-F, F] \times [0, 1]),$$

such that $A = A^T$, $\max_{\overline{Q_k}} g = 1$ and

$$K^{-1} \leq \min_{y \in \mathbb{S}^{n-1}} \min_{(t,x) \in \overline{Q_k}} (y \cdot A(t,x)y) \leq \max_{y \in \mathbb{S}^{n-1}} \max_{(t,x) \in \overline{Q_k}} (y \cdot A(t,x)y) \leq K.$$

For all $U \in \mathcal{U}_k$, denote u_U the solution of

$$\begin{cases} \partial_t u - \nabla \cdot (A \nabla u) + q \cdot \nabla u - l u - f u = g & \text{in } Q_k, \\ u = 0 & \text{on } \partial_P Q_k, \end{cases}$$

and denote $m(U) = \min_{\overline{Q_{k+1}}} u_U > 0$. Let us verify that $\inf_{U \in \mathcal{U}_k} m(U) > 0$.

Assume by contradiction $\inf_{U \in \mathcal{U}_k} m(U) = 0$. Then there exists a minimizing sequence $(U_p)_{p \in \mathbb{N}}$ such that $m(U_p) \rightarrow 0$ as $p \rightarrow +\infty$. By classical compactness

and regularity estimates [52], up to extraction, (U_p) converges uniformly to a limit $U_\infty = (A_\infty, q_\infty, l_\infty, f_\infty, g_\infty)$ such that

$$A_\infty \in \mathcal{C}^{0,1}(\overline{Q_k}, \mathbb{R}^{n \times n}),$$

$$(q_\infty, l_\infty, f_\infty, g_\infty) \in \mathcal{C}(\overline{Q_k}, \mathbb{R}^{n \times n} \times [-K, K]^n \times [-K, K] \times [-F, F] \times [0, 1]),$$

such that $A_\infty = A_\infty^T$, $\max_{\overline{Q_k}} g_\infty = 1$ and

$$K^{-1} \leq \min_{y \in \mathbb{S}^{n-1}} \min_{(t,x) \in \overline{Q_k}} (y \cdot A_\infty(t,x)y) \leq \max_{y \in \mathbb{S}^{n-1}} \max_{(t,x) \in \overline{Q_k}} (y \cdot A_\infty(t,x)y) \leq K$$

and $(u_p)_{p \in \mathbb{N}} = (u_{U_p})_{p \in \mathbb{N}}$ converges uniformly to the solution u_∞ of

$$\begin{cases} \partial_t u - \nabla \cdot (A_\infty \nabla u) + q_\infty \cdot \nabla u - l_\infty u - f_\infty u = g_\infty & \text{in } Q_k, \\ u = 0 & \text{on } \partial_P Q_k. \end{cases}$$

Moreover, by definition of (U_p) , $m(U_\infty) = \lim_{p \rightarrow +\infty} m(U_p) = 0$. But then the strong maximum principle yields $u_\infty = 0$, and this contradicts $g_\infty > 0$. Hence $\inf_{U \in \mathcal{U}_k} m(U) > 0$ and Claim 2 is proved with $C_k = \inf_{U \in \mathcal{U}_k} m(U) > 0$. \square

Proof of Claim 1. Let $k \in [N-1]$, $I \subset [N]$ of cardinal k ,

$$M = \max_{(t,x) \in [0, 2\theta] \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_1(t,x),$$

and assume that for all $j \in I$,

$$\min_{(t,x) \in \overline{Q_k}} u_j(t,x) \geq \kappa_k M.$$

By (A₃), there exists $i \in [N] \setminus I$ and $j \in I$ such that $\max_{\overline{\Omega_{\text{per}}}} l_{i,j} > 0$.

Let \underline{u} be the solution of $\mathcal{P}_i \underline{u} - l_{i,i} \underline{u} - f_i \underline{u} = \kappa_k M l_{i,j}$ in Q_k , $\underline{u} = 0$ on $\partial_P Q_k$. Since Q_k contains a translation of $\overline{\Omega_{\text{per}}}$ and $l_{i,j}$ is periodic, applying Claim 2, we get:

$$\min_{\overline{Q_{k+1}}} \underline{u} \geq C_k \kappa_k M \max_{\overline{\Omega_{\text{per}}}} l_{i,j}.$$

Moreover, in Q_k ,

$$\mathcal{P}_i u_i - l_{i,i} u_i - f_i u_i = \sum_{k \in [N] \setminus \{i\}} l_{i,k} u_k \geq l_{i,j} u_j \geq l_{i,j} \min_{(t,x) \in \overline{Q_k}} u_j \geq \kappa_k M l_{i,j}.$$

Also, on the parabolic boundary $\partial_P Q_k$, $u_i \gg 0 = \underline{u}$. Therefore, by virtue of the comparison principle, $u_i \geq \underline{u}$ in Q_k , and subsequently, using the definition of σ ,

$$\min_{\overline{Q_{k+1}}} u_i \geq \min_{\overline{Q_{k+1}}} \underline{u} \geq C_k \kappa_k M \max_{\overline{\Omega_{\text{per}}}} l_{i,j} \geq C_k \kappa_k M \sigma.$$

Setting $\kappa_{k+1} = C_k \kappa_k \sigma$, we have proved Claim 1. \square

This ends the proof. \square

Remark 2.1. As an immediate corollary, if \mathbf{u} is time periodic, then $\overline{\kappa}_{\theta, F} < 1$ and

$$\min_{i \in [N]} \min_{(t,x) \in \mathbb{R} \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t,x) \geq \overline{\kappa}_{\theta, F} \max_{i \in [N]} \max_{(t,x) \in \mathbb{R} \times [-\frac{\theta}{2}, \frac{\theta}{2}]^n} u_i(t,x).$$

If \mathbf{u} is space-time periodic, then an even stronger estimate holds:

$$\min_{i \in [N]} \min_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} u_i(t,x) \geq \overline{\kappa}_{\theta, F} \max_{i \in [N]} \max_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} u_i(t,x).$$

3. PROOFS

3.1. Existence, characterization and concavity: proof of Theorems 1.1–1.3. The main result of this subsection is Theorem 1.1. It is actually a consequence of Theorems 1.2 and of a concavity result on $z \mapsto \lambda_{1,z}$, Corollary 3.6, that will follow from a more general concavity result, Proposition 3.5, that also contains 1.3.

Most proofs in this subsection are direct adaptations to the vector case of the proofs by Nadin [59], written here for the paper to be self-contained. The only proofs whose adaptations truly require some care are those of Propositions 3.5 and 3.9.

3.1.1. *The generalized principal eigenvalue λ_1 in arbitrary domains: proof of Theorem 1.2.*

Proposition 3.1. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$. Then the generalized principal eigenvalue $\lambda_1(\Omega) \in \mathbb{R}$ is well-defined.*

Furthermore, if $\partial\Omega$ is bounded and smooth, then $\lambda_1(\Omega) = \lambda_{1,\text{Dir}}(\Omega)$.

Proof. We begin with the case of bounded smooth domains. The inequality $\lambda_{1,\text{Dir}}(\Omega) \leq \lambda_1(\Omega)$ follows by using the Dirichlet principal eigenfunction as test function in the definition of $\lambda_1(\Omega)$. The converse inequality is proved by contradiction: assume that $\lambda_{1,\text{Dir}}(\Omega) < \lambda_1(\Omega)$. Then there exists $\mu \in (\lambda_{1,\text{Dir}}(\Omega), \lambda_1(\Omega))$ and $\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega})$ such that $\mathcal{Q}\mathbf{u} \geq \mu\mathbf{u}$. By boundedness of the Dirichlet principal eigenfunction \mathbf{v} , the quantity

$$\kappa^* = \inf \{ \kappa > 0 \mid \kappa\mathbf{u} - \mathbf{v} \gg \mathbf{0} \}$$

is well-defined in \mathbb{R} . The function $\mathbf{w} = \kappa^*\mathbf{u} - \mathbf{v}$ satisfies

$$\mathcal{Q}\mathbf{w} = \kappa^*\mu\mathbf{u} - \lambda_{1,\text{Dir}}(\Omega)\mathbf{v} \gg \lambda_{1,\text{Dir}}(\Omega)\mathbf{w} \quad \text{in } \mathbb{R} \times \Omega,$$

$$\mathbf{w} \geq \mathbf{0} \quad \text{in } \mathbb{R} \times \bar{\Omega},$$

$$\mathbf{w} \geq \mathbf{0} \quad \text{on } \mathbb{R} \times \partial\Omega,$$

and there exists $(i^*, t^*, x^*) \in [N] \times [0, T] \times \bar{\Omega}$ such that $w_{i^*}(t^*, x^*) = 0$. If $x^* \in \Omega$, then by virtue of the strong maximum principle (see Proposition 2.2), \mathbf{w} is the zero function, which contradicts $\mu > \lambda_{1,\text{Dir}}(\Omega)$. Hence $\mathbf{w} \gg \mathbf{0}$ in $\mathbb{R} \times \Omega$. Since $\mathbf{u} \in \mathcal{C}^1(\mathbb{R} \times \bar{\Omega})$, the normal derivative of \mathbf{w} at any point $(t, x) \in \mathbb{R} \times \partial\Omega$ is well-defined. The optimality of κ^* implies the existence of $(i', t', x') \in [N] \times [0, T] \times \partial\Omega$ such that both $w_{i'}(t', x')$ and the normal derivative of $w_{i'}$ at (t', x') are zero, which contradicts the boundary version of the strong maximum principle. Hence $\lambda_1(\Omega) \leq \lambda_{1,\text{Dir}}(\Omega)$. This ends the proof in the case of bounded smooth domains.

Then we turn to general, not necessarily bounded and smooth, domains. Let $\nu = -\lambda_{\text{PF}}(\bar{\mathbf{L}}) \in \mathbb{R}$, where the square matrix $\bar{\mathbf{L}}$ is defined in (A₃), and let $\mathbf{u} \in \mathbb{R}^N$ be a positive Perron–Frobenius eigenvector for $\bar{\mathbf{L}}$, namely $\bar{\mathbf{L}}\mathbf{u} = -\nu\mathbf{u}$. Then clearly $\mathcal{Q}\mathbf{u} \geq \nu\mathbf{u}$, which proves that the set

$$\left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega}) \quad \mathcal{Q}\mathbf{u} \geq \lambda\mathbf{u} \right\}$$

is nonempty. Hence its supremum, $\lambda_1(\Omega)$, is well-defined in $\mathbb{R} \times \{\infty\}$.

Next, it follows directly from the definition that $\lambda_1(\Omega) \leq \lambda_1(\Omega')$ for any open set $\Omega' \subset \Omega$. Since Ω is open and contains a periodicity cell $[x_0, x_0 + L]$, it contains

a bounded smooth connected open set Ω' satisfying $[x_0, x_0 + L] \subset \Omega' \subset \overline{\Omega'} \subset \Omega$. Therefore

$$\lambda_1(\Omega) \leq \lambda_1(\Omega') = \lambda_{1,\text{Dir}}(\Omega') < +\infty.$$

This ends the proof. \square

Proposition 3.2. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set and let $(\Omega_k)_{k \in \mathbb{N}}$ be a sequence of nonempty open connected sets such that, for some $x_0 \in \Omega$,*

$$[x_0, x_0 + L] \subset \Omega_1, \quad \Omega_k \subset \Omega_{k+1}, \quad \bigcup_{k \in \mathbb{N}} \Omega_k = \Omega.$$

Then $\lambda_1(\Omega_k) \rightarrow \lambda_1(\Omega)$ as $k \rightarrow +\infty$.

Furthermore, there exists a generalized principal eigenfunction associated with $\lambda_1(\Omega)$.

Proof. In order to work with bounded and smooth domains, we consider a family $(\tilde{\Omega}_k)_{k \in \mathbb{N}}$, nondecreasing and convergent to Ω in the inclusion sense, and such that $\tilde{\Omega}_k \subset \Omega_k$ for all $k \in \mathbb{N}$ (with $[x_0, x_0 + L] \subset \tilde{\Omega}_1$, which is always possible since $[x_0, x_0 + L]$ is closed and Ω_1 is open). Denote $(\mu_k)_{k \in \mathbb{N}} = (\lambda_{1,\text{Dir}}(\Omega_k))_{k \in \mathbb{N}}$, $(\nu_k)_{k \in \mathbb{N}} = (\lambda_{1,\text{Dir}}(\tilde{\Omega}_k))_{k \in \mathbb{N}}$, and note that both sequences converge, with limits satisfying

$$\lambda_1(\Omega) \leq \lim_{k \rightarrow +\infty} \mu_k \leq \lim_{k \rightarrow +\infty} \nu_k.$$

Let $\nu = \lim \nu_k$. We now aim to prove that $\nu \leq \lambda_1(\Omega)$ by constructing an eigenfunction for the eigenvalue ν of the operator \mathcal{Q} acting on $\mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}_0^1(\mathbb{R} \times \overline{\Omega})$. Since such an eigenfunction will in fact be a generalized principal eigenfunction for the generalized principal eigenvalue $\lambda_1(\Omega)$, this will complete the proof.

Fix $y \in \tilde{\Omega}_1 = \bigcap_{k \in \mathbb{N}} \tilde{\Omega}_k$ and consider the sequence $(\mathbf{u}_k)_{k \in \mathbb{N}}$ of positive principal eigenfunctions associated with ν_k and normalized by $\max_{i \in [N]} u_{i,k}(0, y) = 1$. Extend these eigenfunctions as functions defined in $\mathbb{R} \times \Omega$ by setting $\mathbf{u}_k = \mathbf{0}$ in $\mathbb{R} \times \Omega \setminus \tilde{\Omega}_k$.

By virtue of the time periodicity of \mathbf{u}_k , of the normalization at time $t = 0$ and of the Harnack inequality of Proposition 2.4, the sequence $\left(\|\mathbf{u}_k\|_{\mathcal{L}^\infty([0, T] \times \tilde{\Omega}_{k_0})} \right)_{k \in \mathbb{N}, k > k_0}$ is bounded for any $k_0 \in \mathbb{N}$. By standard regularity estimates [52], $(\mathbf{u}_k)_{k \in \mathbb{N}}$ converges up to a diagonal extraction to a function $\mathbf{u}_\infty \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega)$ satisfying

$$\mathcal{Q}\mathbf{u}_\infty = \nu\mathbf{u}_\infty \quad \text{in } \mathbb{R} \times \Omega.$$

Moreover, \mathbf{u}_∞ is nonnegative, nonzero at $(t, x) = (0, y)$, and by the maximum principle it is therefore positive in $\mathbb{R} \times \Omega$.

In order to establish $\nu \leq \lambda_1(\Omega)$, it only remains to verify that $\mathbf{u}_\infty \in \mathcal{C}_0^1(\mathbb{R} \times \overline{\Omega})$.

Let

$$C = |\lambda_1(\tilde{\Omega}_1)| \sup_{k \in [N]} \|\mathbf{u}_k\|_{\mathcal{L}^\infty([0, T] \times \tilde{\Omega}_k)}$$

and define $\hat{\mathbf{u}} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}_0^1(\mathbb{R} \times \overline{\Omega}, [\mathbf{0}, \infty))$ as the time periodic solution of the following (decoupled) system:

$$\begin{cases} \text{diag}(\mathcal{P})\hat{\mathbf{u}} = \mathbf{1} & \text{in } \mathbb{R} \times \Omega, \\ \hat{\mathbf{u}} = \mathbf{0} & \text{on } \mathbb{R} \times \partial\Omega. \end{cases}$$

Then, for any $k \in \mathbb{N}$,

$$\text{diag}(\mathcal{P})((C\hat{\mathbf{u}} - \mathbf{u}_k) \geq C\mathbf{1} - \sup_{k \in [N]} \left(\lambda_1(\tilde{\Omega}_k) \right) \mathbf{u}_k = C\mathbf{1} - \lambda_1(\tilde{\Omega}_1) \mathbf{u}_k \geq \mathbf{0}.$$

This leads to $\mathbf{u}_k \leq C\hat{\mathbf{u}}$ for all $k \in [N]$, and then, passing to the limit, $\mathbf{u}_\infty \leq C\hat{\mathbf{u}}$ in $\mathbb{R} \times \Omega$. Hence $\mathbf{u}_\infty \in \mathcal{C}_0(\mathbb{R} \times \bar{\Omega})$. The continuity of its gradient $\nabla \mathbf{u}_\infty$ on the regular boundary points follows from classical regularity estimates up to the boundary [52]. This ends the proof. \square

Remark 3.1. The proof uses the interior Harnack inequality of Proposition 2.4, which, as stated, requires that the domain of definition contains a translation of $[0, 3\theta]^n$, with $\theta \geq \max(T, L_1, \dots, L_n)$. This is not optimal and just for convenience of notation; what truly matters for the interior Harnack inequality is that the domain of definition is strictly larger than a closed periodicity cell, as expressed in the preceding statement. We leave the necessary correction of the proof of Proposition 2.4 as an exercise for interested readers.

Proposition 3.3. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$. Then the generalized principal eigenvalue $\lambda_1(\Omega)$ can be characterized as:*

$$\lambda_1(\Omega) = \max_{\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega})} \min_{i \in [N]} \inf_{\mathbb{R} \times \Omega} \left(\frac{(\mathcal{Q}\mathbf{u})_i}{u_i} \right).$$

Proof. Testing \mathcal{Q} against a generalized principal eigenfunction (whose existence is guaranteed by Proposition 3.2), we directly find

$$\lambda_1(\Omega) \leq \sup_{\mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty)) \cap \mathcal{C}^1(\mathbb{R} \times \bar{\Omega})} \min_{i \in [N]} \inf_{\mathbb{R} \times \Omega} \left(\frac{(\mathcal{Q}\mathbf{u})_i}{u_i} \right).$$

Next we assume by contradiction that the above inequality is actually strict. Then there exists $\mu > \lambda_1(\Omega)$ and a test function \mathbf{u} such that $\mathcal{Q}\mathbf{u} \geq \mu\mathbf{u}$. This contradicts the definition of $\lambda_1(\Omega)$.

Finally, the existence of a generalized principal eigenfunction shows that the supremum is in fact a maximum, as in the statement. \square

3.1.2. Characterizations of the periodic principal eigenvalues $\lambda_{1,z}$. For any $z \in \mathbb{R}^n$, the existence and uniqueness of the eigenpair $(\lambda_{1,z}, \mathbf{u}_z)$, up to multiplication of the eigenfunction by a constant, follows from the Krein–Rutman theorem. We do not detail the proof of this claim. Below, we prove a generalization of the classical Collatz–Wielandt formula for Perron–Frobenius eigenvalues.

Proposition 3.4. *Let $z \in \mathbb{R}^n$. Then the periodic principal eigenvalue $\lambda_{1,z}$ can be characterized as:*

$$(11) \quad \lambda_{1,z} = \max_{\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \min_{i \in [N]} \min_{\bar{\Omega}_{\text{per}}} \left(\frac{(\mathcal{Q}_z \mathbf{u})_i}{u_i} \right),$$

$$(12) \quad \lambda_{1,z} = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \max_{i \in [N]} \max_{\bar{\Omega}_{\text{per}}} \left(\frac{(\mathcal{Q}_z \mathbf{u})_i}{u_i} \right).$$

Proof. We prove only the max–min characterization, the min–max one being proved quite similarly.

Using the existence of the periodic principal eigenfunction \mathbf{u}_z , we immediately obtain

$$\lambda_{1,z} \leq \sup_{\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))} \min_{i \in [N]} \min_{\overline{\Omega_{\text{per}}}} \left(\frac{(\mathcal{Q}_z \mathbf{u})_i}{u_i} \right).$$

Next we assume by contradiction that the above inequality is actually strict. Then there exists a test function $\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ and a real number $\mu > \lambda_{1,z}$ such that $\mathcal{Q}_z \mathbf{u} \geq \mu \mathbf{u}$. Let

$$\kappa^* = \inf \{ \kappa > 0 \mid \kappa \mathbf{u} - \mathbf{u}_z \gg \mathbf{0} \}.$$

Applying the strong maximum principle to $\kappa^* \mathbf{u} - \mathbf{u}_z$, just as in the proof of Proposition 3.1, we find a contradiction.

Finally, the existence of \mathbf{u}_z shows that the supremum is in fact a maximum, as in the statement. \square

3.1.3. Concave dependence on z and \mathbf{L} . In order to show later on that $\lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_z$, we need to establish first the strict concavity of $z \mapsto \lambda_z$. This is stated below in Corollary 3.6. Since the proof of Theorem 1.3 on the concavity of $\mathbf{L} \mapsto \lambda_z(\mathbf{L})$ is quite similar, we prove the two results directly together.

Proposition 3.5. *Let $z_1, z_2 \in \mathbb{R}^n$.*

Let

$$(\mathbf{L}[s])_{s \in [0,1]} \in \left(\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N}) \right)^{[0,1]}$$

a family of matrices satisfying the same assumptions as \mathbf{L} (i.e., (A_2) , (A_3)) and such that, for all $(t, x) \in \mathbb{R} \times \mathbb{R}^n$ and $i \in [N]$,

- (1) $s \mapsto l_{i,i}[s](t, x)$ is convex;*
- (2) for all $j \in [N] \setminus \{i\}$, $s \mapsto l_{i,j}[s](t, x)$ is either identically zero or log-convex.*

For all $s \in [0, 1]$, denote

$$\mathcal{Q}[s] = e_{-(1-s)z_1 - s z_2} (\text{diag}(\mathcal{P}_i) - \mathbf{L}[s]) e_{(1-s)z_1 + s z_2}$$

and $\lambda[s] = \lambda_{1, \text{per}}(\mathcal{Q}[s])$ the associated periodic principal eigenvalue.

Then $s \in [0, 1] \mapsto \lambda[s]$ is affine or strictly concave and it is affine if and only if the following conditions are both satisfied:

- (Cond. 1) $z_1 = z_2$;*
- (Cond. 2) there exist a constant vector $\mathbf{b} \gg \mathbf{0}$, a function $\mathbf{c} \in \mathcal{C}_{\text{per}}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ and a function $\mathbf{f} \in \mathcal{C}_{\text{per}}(\mathbb{R}, \mathbb{R}^N)$ satisfying $\int_0^T \mathbf{f} \in \text{span}(\mathbf{1})$ such that the entries of \mathbf{L} have the form:*

$$l_{i,j}[s] : (t, x) \mapsto \begin{cases} l_{i,i}[0](t, x) - s f_i(t) & \text{if } i = j, \\ l_{i,j}[0](t, x) \left(\frac{b_j}{c_i(t, x)} \right)^s e^{s \left(\int_0^t f_j - \frac{t}{T} \int_0^T f_j \right)} & \text{if } i \neq j \end{cases}$$

and such that the function \mathbf{c} satisfies, at all $(t, x) \in \overline{\Omega_{\text{per}}}$ and for each $i \in [N]$,

$$c_i(t, x) = b_i e^{\int_0^t f_i - \frac{t}{T} \int_0^T f_i} \quad \text{or} \quad \forall j \in [N] \setminus \{i\}, l_{i,j}[0](t, x) = 0.$$

Proof. We divide the proof into three steps: the concavity of $s \mapsto \lambda[s]$, the alternative between affinity or strict concavity, the characterization of the affinity case.

Step 1: concavity. Fix $s \in [0, 1]$, set $z = (1-s)z_1 + sz_2$ and, for all $(t, x) \in \mathbb{R} \times \mathbb{R}^n$, define the auxiliary matrix $\tilde{\mathbf{L}}[s](t, x)$ whose entries are:

$$\tilde{l}_{i,j}[s](t, x) = \begin{cases} (1-s)l_{i,i}[0](t, x) + sl_{i,i}[1](t, x) & \text{if } i = j \\ (l_{i,j}[0](t, x))^{1-s} (l_{i,j}[1](t, x))^s & \text{if } i \neq j \end{cases}$$

(with $0^0 = 0$ by convention). By construction, and by our convexity assumptions, $\mathbf{L}[s] \leq \tilde{\mathbf{L}}[s]$ in $\mathbb{R} \times \mathbb{R}^n$. Hence, as a direct consequence of the min-max/max-min characterizations of the periodic principal eigenvalue of Proposition 3.4, we get:

$$(13) \quad \lambda[s] \geq \lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[s]),$$

where $\tilde{\mathbf{Q}}[s] = \mathbf{Q}[s] + \mathbf{L}[s] - \tilde{\mathbf{L}}[s]$.

Recall the notation $e_{z'} : x \mapsto e^{z' \cdot x}$ and note that, by definition of $\tilde{\mathbf{Q}}[s]$,

$$\frac{\left(\tilde{\mathbf{Q}}[s](e_{-z}\mathbf{u})\right)_i}{e_{-z}u_i} = \frac{\mathcal{P}_i u_i - \left(\tilde{\mathbf{L}}[s]\mathbf{u}\right)_i}{u_i} \quad \text{for all } \mathbf{u} \in \mathcal{C}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)).$$

Hence there is a bijection between space-time periodic eigenfunctions of $\tilde{\mathbf{Q}}[s]$ and time periodic eigenfunctions of $\text{diag}(\mathcal{P}) - \tilde{\mathbf{L}}[s]$ whose product with e_z is space periodic.

Let $\mu = \lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[0])$, $\nu = \lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[1])$ and $e_{-z_1}\mathbf{u}$, $e_{-z_2}\mathbf{v}$ two respectively associated space-time periodic positive eigenfunctions:

$$\tilde{\mathbf{Q}}[0](e_{-z_1}\mathbf{u}) = \mu e_{-z_1}\mathbf{u}, \quad \tilde{\mathbf{Q}}[1](e_{-z_2}\mathbf{v}) = \nu e_{-z_2}\mathbf{v},$$

i.e.

$$\text{diag}(\mathcal{P})\mathbf{u} - \tilde{\mathbf{L}}[0]\mathbf{u} = \mu\mathbf{u}, \quad \text{diag}(\mathcal{P})\mathbf{v} - \tilde{\mathbf{L}}[1]\mathbf{v} = \nu\mathbf{v}.$$

Define $\mathbf{w} = (u_i^{1-s}v_i^s)_{i \in [N]}$. Since $e_{-z}w_i = (e_{-z_1}u_i)^{1-s}(e_{-z_2}v_i)^s$ for all $i \in [N]$, $e_{-z}\mathbf{w}$ is space-time periodic and therefore we can use it as test function for $\tilde{\mathbf{Q}}[s]$. Following Nadin [59] for the expansion of the \mathcal{P}_i part and using the uniform ellipticity assumption (A_1) , we find:

$$\begin{aligned} \frac{\mathcal{P}_i w_i - \left(\tilde{\mathbf{L}}[s]\mathbf{w}\right)_i}{w_i} &= (1-s)\frac{\mathcal{P}_i u_i}{u_i} + s\frac{\mathcal{P}_i v_i}{v_i} + s(1-s)\left(\frac{\nabla u_i}{u_i} - \frac{\nabla v_i}{v_i}\right) \cdot A_i\left(\frac{\nabla u_i}{u_i} - \frac{\nabla v_i}{v_i}\right) \\ &\quad - (1-s)\tilde{l}_{i,i}[0] - \tilde{s}l_{i,i}[1] - \frac{1}{w_i} \sum_{j \in [N] \setminus \{i\}} (\tilde{l}_{i,j}[0]u_j)^{1-s} (\tilde{l}_{i,j}[1]v_j)^s \\ &\geq (1-s)\frac{\mathcal{P}_i u_i}{u_i} + s\frac{\mathcal{P}_i v_i}{v_i} \\ &\quad - (1-s)\tilde{l}_{i,i}[0] - \tilde{s}l_{i,i}[1] - \frac{1}{w_i} \sum_{j \in [N] \setminus \{i\}} (\tilde{l}_{i,j}[0]u_j)^{1-s} (\tilde{l}_{i,j}[1]v_j)^s. \end{aligned}$$

Following Nussbaum [63] and using the Hölder inequality, the equalities satisfied by \mathbf{u} and \mathbf{v} and the inequality between arithmetic and geometric means, we get

$$(14) \quad \frac{\left(\tilde{\mathbf{Q}}[s](e_{-z}\mathbf{w})\right)_i}{e_{-z}w_i} \geq (1-s)\mu + s\nu \quad \text{for all } i \in [N],$$

and, eventually, the max-min characterization yields:

$$(15) \quad \lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[s]) \geq (1-s)\mu + s\nu = (1-s)\lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[0]) + s\lambda_{1,\text{per}}(\tilde{\mathbf{Q}}[1]).$$

Combining (13) and (15) and using the fact that $\mathcal{Q}[s]$ and $\tilde{\mathcal{Q}}[s]$ coincide at $s = 0$ and $s = 1$, we find indeed the claimed concavity:

$$(16) \quad \lambda_{1,\text{per}}(\mathcal{Q}[s]) \geq (1-s)\lambda_{1,\text{per}}(\mathcal{Q}[0]) + s\lambda_{1,\text{per}}(\mathcal{Q}[1]).$$

□

Step 2: affinity or strict concavity. Assume that $s \mapsto \lambda[s]$ is not strictly concave. This means that there exists $s_0 \in [0, 1]$ such that (16) is an equality at $s = s_0$.

The equality in (16) at $s = s_0$ implies the equality in (13) at $s = s_0$, which in turn implies the equality $\mathbf{L}[s_0] = \tilde{\mathbf{L}}[s_0]$ in $\mathbb{R} \times \mathbb{R}^n$. Since all $s \mapsto \tilde{l}_{i,i}[s](t, x)$ are linear and all $s \mapsto l_{i,i}[s](t, x)$ are convex, $l_{i,i}[s](t, x) \leq \tilde{l}_{i,i}[s](t, x)$ together with the equality at $s = 0$, $s = s_0$, $s = 1$ imply $l_{i,i} = \tilde{l}_{i,i}$ identically for all $i \in [N]$. Similarly, $l_{i,j} = \tilde{l}_{i,j}$ identically for all $i, j \in [N]$. Hence, as functions of (s, t, x) , $\mathbf{L} = \tilde{\mathbf{L}}$ identically in $[0, 1] \times \mathbb{R} \times \mathbb{R}^n$.

Similarly, the equality in (16) at $s = s_0$ implies the equality in (15) at $s = s_0$, and then the max-min characterization (Proposition 3.4) implies equality in (14) at $s = s_0$ for all $i \in [N]$ in $\mathbb{R} \times \mathbb{R}^n$. Then, this implies, for all $i \in [N]$:

- $\nabla u_i/u_i = \nabla v_i/v_i$, that is there exists a function a_i of the variable t only such that $u_i(t, x) = a_i(t)v_i(t, x)$;
- for all $j \in [N] \setminus \{i\}$, there exists a positive function c_i of t and x such that $\tilde{l}_{i,j}[0]u_j = c_i \tilde{l}_{i,j}[1]v_j$ (equality in the Hölder inequality);
- $\frac{\mathcal{P}_i u_i}{u_i} - \tilde{l}_{i,i}[0] - \mu = \frac{\mathcal{P}_i v_i}{v_i} - \tilde{l}_{i,i}[1] - \nu$ (equality in the inequality between geometric and arithmetic averages).

Putting the two together, the equality in (16) at $s = s_0$ implies:

- (Cond. 1') $\mathbf{L} = \tilde{\mathbf{L}}$ identically in $[0, 1] \times \mathbb{R} \times \mathbb{R}^n$;
- (Cond. 2') there exists a function a_i of the variable t only such that $u_i(t, x) = a_i(t)v_i(t, x)$;
- (Cond. 3') for all $j \in [N] \setminus \{i\}$, there exists a positive function c_i of t and x such that $\tilde{l}_{i,j}[0]a_j = c_i \tilde{l}_{i,j}[1]$;
- (Cond. 4') $\frac{\mathcal{P}_i u_i}{u_i} - \tilde{l}_{i,i}[0] - \mu = \frac{\mathcal{P}_i v_i}{v_i} - \tilde{l}_{i,i}[1] - \nu$.

These four conditions do not depend on s_0 . Going back through Step 1, it appears that under these conditions, all inequalities are equalities. Hence (16) is an equality at all $s \in [0, 1]$, or in other words $s \mapsto \lambda[s]$ is affine. It will be useful in the next step to note that this argument precisely shows that (Cond. 1')–(Cond. 4') are equivalent to the affinity of $s \mapsto \lambda[s]$.

□

Step 3: necessary and sufficient conditions for affinity. From Step 2, we know that $s \mapsto \lambda[s]$ is affine if and only if (Cond. 1')–(Cond. 4'). Let us prove that this group of conditions is equivalent to the group (Cond. 1)–(Cond. 2).

Note first that without loss of generality, we can assume that \mathbf{u} and \mathbf{v} are uniquely identified by the following normalizations:

$$\|u_1(0, \cdot)\|_{\mathcal{L}^\infty(\mathbb{R}^n, \mathbb{R})} = 1, \quad \|v_1(0, \cdot)\|_{\mathcal{L}^\infty(\mathbb{R}^n, \mathbb{R})} = 1.$$

First, we prove that (Cond. 1')–(Cond. 4') imply (Cond. 1)–(Cond. 2). From $u_i(t, x) = a_i(t)v_i(t, x)$, we deduce $z_1 = z_2$ (recall that $e_{-z_1}\mathbf{u}$ and $e_{-z_2}\mathbf{v}$ are both space-time periodic) and $\frac{\mathcal{P}_i u_i}{u_i} = \frac{\mathcal{P}_i v_i}{v_i} + \frac{a_i'}{a_i}$. The equality $\frac{\mathcal{P}_i u_i}{u_i} - \tilde{l}_{i,i}[0] - \mu =$

$\frac{\mathcal{P}_i v_i}{v_i} - \tilde{l}_{i,i}[1] - \nu$ reads $\frac{a'_i}{a_i} = \tilde{l}_{i,i}[0] - \tilde{l}_{i,i}[1] + \mu - \nu$, or in other words there exists $\mathbf{b} \in \mathbb{R}^N$ such that

$$a_i : t \mapsto b_i \exp \left(\int_0^t \left(\tilde{l}_{i,i}[0](t', x) - \tilde{l}_{i,i}[1](t', x) \right) dt' + (\mu - \nu)t \right).$$

This directly implies that $f_i = \tilde{l}_{i,i}[0] - \tilde{l}_{i,i}[1]$ does not depend on x . Moreover, the positivity of both u_i and v_i implies $b_i > 0$, the normalizations imply $b_1 = 1$, and the time periodicity implies that f_i is periodic with average $\nu - \mu$, independent of i . To characterize \mathbf{c} , we sum the $N - 1$ equalities $l_{i,j}[0]a_j = c_i l_{i,j}[1]$ for $j \in [N] \setminus \{i\}$ coming from (Cond. 3') and rearrange terms as follows:

$$\begin{aligned} 0 &= \left(\sum_{j \in [N] \setminus \{i\}} l_{i,j}[1]v_j \right) c_i - \sum_{j \in [N] \setminus \{i\}} l_{i,j}[0]u_j \\ &= ((\mathbf{L}[1]\mathbf{v})_i - l_{i,i}[1]v_i) c_i - ((\mathbf{L}[0]\mathbf{u})_i - l_{i,i}[0]u_i) \\ &= \left(\frac{\mathcal{P}_i v_i}{v_i} - l_{i,i}[1] - \nu \right) v_i c_i - \left(\frac{\mathcal{P}_i u_i}{u_i} - l_{i,i}[0] - \mu \right) u_i \end{aligned}$$

Now, using (Cond. 2') and (Cond. 4'), we get

$$0 = \left(\frac{\mathcal{P}_i v_i}{v_i} - l_{i,i}[1] - \nu \right) v_i (c_i - a_i) = \left(\sum_{j \in [N] \setminus \{i\}} l_{i,j}[1]v_j \right) (c_i - a_i).$$

By nonnegativity of each term in the sum, we deduce that, at each $(t, x) \in \overline{\Omega_{\text{per}}}$,

$$c_i(t, x) = a_i(t) \quad \text{or} \quad \forall j \in [N] \setminus \{i\}, l_{i,j}[0](t, x) = 0.$$

Second, to verify that (Cond. 1)–(Cond. 2) imply (Cond. 1')–(Cond. 4'), it suffices to set

$$a_i : t \mapsto \frac{b_i}{b_1} \exp \left(\int_0^t f_i - \frac{t}{T} \int_0^T f_i \right),$$

and to check $\frac{1}{T} \int_0^T f_i = \mu - \nu$ and $\mathbf{u} = \mathbf{a} \circ \mathbf{v}$. Actually, $\tilde{\mathbf{u}} = \mathbf{a} \circ \mathbf{v}$ satisfies

$$\begin{aligned} \mathcal{P}_i \tilde{u}_i - (\mathbf{L}[0]\tilde{\mathbf{u}})_i &= a'_i v_i + a_i \mathcal{P}_i v_i - (\mathbf{L}[0](\mathbf{a} \circ \mathbf{v}))_i \\ &= \left(f_i - \frac{1}{T} \int_0^T f_i \right) a_i v_i + a_i (\mathbf{L}[1]\mathbf{v})_i + a_i \lambda_{1,\text{per}}(\mathcal{Q}[1])v_i - (\mathbf{L}[0](\mathbf{a} \circ \mathbf{v}))_i \\ &= \left(\lambda_{1,\text{per}}(\mathcal{Q}[1]) - \frac{1}{T} \int_0^T f_i \right) \tilde{u}_i + \sum_{j \in [N] \setminus \{i\}} ((l_{i,j}[1]a_i - l_{i,j}[0]a_j) v_j) \\ &= \left(\lambda_{1,\text{per}}(\mathcal{Q}[1]) - \frac{1}{T} \int_0^T f_i \right) \tilde{u}_i + (a_i - c_i) \sum_{j \in [N] \setminus \{i\}} l_{i,j}[1]v_j \\ &= \left(\lambda_{1,\text{per}}(\mathcal{Q}[1]) - \frac{1}{T} \int_0^T f_i \right) \tilde{u}_i. \end{aligned}$$

Since $e_{-z_2} \mathbf{v}$ is space-time periodic, $e_{-z_2} \tilde{\mathbf{u}} = e_{-z_2} \mathbf{a} \circ \mathbf{v}$ is also space-time periodic, whence by uniqueness $\lambda_{1,\text{per}}(\mathcal{Q}[1]) - T^{-1} \int_0^T f_i = \lambda_{1,\text{per}}(\mathcal{Q}[0])$ and $e_{-z_1} \tilde{\mathbf{u}} \in \text{span}(\mathbf{u}_{z_1})$. This exactly proves the existence of $C > 0$ such that $C\tilde{\mathbf{u}} = \mathbf{u}$, and, in view of the chosen normalizations on \mathbf{u} and \mathbf{v} , $C = 1$, *i.e.* $\tilde{\mathbf{u}} = \mathbf{u}$. \square

The proof of the theorem is complete. \square

Remark 3.2. When $\mathbf{L}[0](t, x)$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$, the characterization of the function \mathbf{c} in (Cond. 2) is immediately strengthened as: c_i only depends on t and $c_i : t \mapsto b_i e^{\int_0^t f_i - \frac{t}{T} \int_0^T f_i}$.

On the contrary, \mathbf{c} cannot be uniquely determined when a line of $\mathbf{L}[0]$ vanishes somewhere in $\overline{\Omega_{\text{per}}}$, and from (Cond. 2) it is actually clear that if the i -th line of $\mathbf{L}[0]$ vanishes in an open space-time ball $B \subset \overline{\Omega_{\text{per}}}$, then basically any nonnegative scalar function can be added to a_i in B and the resulting sum still forms an admissible \mathbf{c} . This might come as a surprise, especially since pointwise irreducible matrices are dense⁷ in the set of admissible matrices (namely, matrices in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ satisfying (\mathbf{A}_2) and (\mathbf{A}_3)).

Corollary 3.6. *With the notations of Proposition 3.5, if $z_1 \neq z_2$, then $s \in [0, 1] \mapsto \lambda[s]$ is strictly concave. In particular, $z \mapsto \lambda_{1,z}$ is strictly concave.*

Very minor adaptations of the proof of Proposition 3.5, not detailed here, lead to the following analogous result in the Dirichlet case⁸.

Proposition 3.7. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty, bounded, smooth, open, connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$.*

Let

$$(\mathbf{L}[s])_{s \in [0,1]} \in \left(\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \Omega, \mathbb{R}^{N \times N}) \right)^{[0,1]}$$

a family of matrices satisfying the same assumptions as \mathbf{L} (i.e., (\mathbf{A}_2) , (\mathbf{A}_3)) and such that, for all $(t, x) \in \mathbb{R} \times \Omega$ and $i \in [N]$,

(1) $s \mapsto l_{i,i}[s](t, x)$ is convex;

(2) for all $j \in [N] \setminus \{i\}$, $s \mapsto l_{i,j}[s](t, x)$ is either identically zero or log-convex.

Then $s \in [0, 1] \mapsto \lambda_{1, \text{Dir}}(\Omega, \mathcal{Q}[s])$, where $\mathcal{Q}[s]$ is the operator \mathcal{Q} with \mathbf{L} replaced $\mathbf{L}[s]$, is affine or strictly concave and it is affine if and only if there exist a constant vector $\mathbf{b} \gg \mathbf{0}$, a function $\mathbf{c} \in \mathcal{C}_{t\text{-per}}(\mathbb{R} \times \Omega, (\mathbf{0}, \infty))$ and a function $\mathbf{f} \in \mathcal{C}_{t\text{-per}}(\mathbb{R}, \mathbb{R}^N)$ satisfying $\int_0^T \mathbf{f} \in \text{span}(\mathbf{1})$ such that the entries of \mathbf{L} have the form:

$$l_{i,j}[s] : (t, x) \mapsto \begin{cases} l_{i,i}[0](t, x) - s f_i(t) & \text{if } i = j, \\ l_{i,j}[0](t, x) \left(\frac{b_j}{c_i(t, x)} \right)^s e^{s \left(\int_0^t f_j - \frac{t}{T} \int_0^T f_j \right)} & \text{if } i \neq j, \end{cases}$$

and such that the function \mathbf{c} satisfies, at all $(t, x) \in \mathbb{R} \times \Omega$ and for each $i \in [N]$,

$$c_i(t, x) = b_i e^{\int_0^t f_i - \frac{t}{T} \int_0^T f_i} \quad \text{or} \quad \forall j \in [N] \setminus \{i\}, l_{i,j}[0](t, x) = 0.$$

As a corollary, we obtain the concavity of λ_1 in arbitrary domains, namely Theorem 1.4.

Corollary 3.8. *Let $\Omega \subset \mathbb{R}^n$ be a nonempty open connected set such that there exists $x_0 \in \Omega$ satisfying $[x_0, x_0 + L] \subset \Omega$.*

Let

$$(\mathbf{L}[s])_{s \in [0,1]} \in \left(\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \Omega, \mathbb{R}^{N \times N}) \right)^{[0,1]}$$

⁷Just change \mathbf{L} into $\mathbf{L} + \varepsilon \mathbf{1}_{N \times N}$.

⁸The absence of z actually makes the proof shorter.

a family of matrices satisfying the same assumptions as \mathbf{L} (i.e., (A_2) , (A_3)) and such that, for all $(t, x) \in \mathbb{R} \times \Omega$ and $i \in [N]$,

- (1) $s \mapsto l_{i,i}[s](t, x)$ is convex;
- (2) for all $j \in [N] \setminus \{i\}$, $s \mapsto l_{i,j}[s](t, x)$ is either identically zero or log-convex.

Then the mapping $s \in [0, 1] \mapsto \lambda_1(\Omega, \mathbf{Q}[s])$, where $\mathbf{Q}[s]$ is the operator \mathbf{Q} with \mathbf{L} replaced $\mathbf{L}[s]$, is concave.

Proof. Just as in the proof of Proposition 3.2, we work with a sequence $(\Omega_k)_{k \in \mathbb{N}}$ of smooth, bounded, nonempty, open, connected subsets of Ω such that

$$[x_0, x_0 + L] \subset \Omega_1, \quad \Omega_k \subset \Omega_{k+1}, \quad \bigcup_{k \in \mathbb{N}} \Omega_k = \Omega.$$

By virtue of Proposition 3.7, all $s \in [0, 1] \mapsto \lambda_{1, \text{Dir}}(\Omega_k, \mathbf{Q}[s])$ are concave. By virtue of Proposition 3.2, $\lambda_{1, \text{Dir}}(\Omega_k, \mathbf{Q}[s]) \rightarrow \lambda_1(\Omega, \mathbf{Q}[s])$ as $k \rightarrow +\infty$, for all $s \in [0, 1]$.

The pointwise convergence of a sequence of concave functions on the compact set $[0, 1]$ is automatically improved as uniform convergence in $[0, 1]$, and the limit is concave on $[0, 1]$ as well. This ends the proof. \square

Remark 3.3. We will establish in the next section that $\lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_z$. However, the maximum of a family of concave functions is in general not a concave function itself, so that this identity cannot be used to prove the concavity of λ_1 .

3.1.4. Relations between λ_1 , λ'_1 and $\lambda_{1,z}$.

Proposition 3.9. *There exists $z \in \mathbb{R}^n$ such that $e_z \mathbf{u}_z$ is a generalized principal eigenfunction of \mathbf{Q} associated with λ_1 and $\lambda_1 = \lambda_{1,z}$.*

Proof. From Proposition 3.2, there exists a generalized principal eigenfunction $\mathbf{u} \in C^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ associated with λ_1 .

We first prove that there exists $z_1 \in \mathbb{R}$ and a new generalized principal eigenfunction \mathbf{u}^1 such that $(t, x) \mapsto e^{-z_1 x_1} \mathbf{u}^1(t, x)$ is L_1 -periodic with respect to x_1 .

Define the translation $\tau : x \in \mathbb{R}^n \mapsto x + L_1 e_1$, where $e_1 = (\delta_{1\alpha})_{\alpha \in [n]}$, and denote $\mathbf{u}^\tau : (t, x) \mapsto \mathbf{u}(t, \tau(x))$ and $\mathbf{v} = (u_i^\tau / u_i)_{i \in [N]}$. By virtue of the fully coupled Harnack inequality of Proposition 2.4 and periodicity of the coefficients of \mathbf{Q} , \mathbf{v} is globally bounded. Let

$$z_1 = L_1^{-1} \ln \left(\max_{i \in [N]} \sup_{(t,x) \in \mathbb{R} \times \mathbb{R}^n} v_i(t, x) \right).$$

Recalling that \mathbf{u} and consequently \mathbf{v} are time periodic, there exists $\bar{i} \in [N]$ and $(t_k, x_k)_{k \in \mathbb{N}} \in ([0, T] \times \mathbb{R}^n)^{\mathbb{N}}$ such that $v_{\bar{i}}(t_k, x_k) \rightarrow e^{z_1 L_1}$ as $k \rightarrow +\infty$. Moreover, there exists $(y_k)_{k \in \mathbb{N}}$ such that, for all $k \in \mathbb{N}$, $x_k - y_k \in L_1 \mathbb{Z} \times \cdots \times L_n \mathbb{Z}$. Up to extraction, we assume that $(t_k, y_k) \rightarrow (t_\infty, y_\infty) \in \overline{\Omega_{\text{per}}}$.

Now, define, for all $k \in \mathbb{N}$,

$$\hat{\mathbf{u}}_k : (t, x) \mapsto \frac{1}{u_{\bar{i}}(t_k, x_k)} \mathbf{u}(t + t_k, x + x_k),$$

$$\hat{\mathbf{u}}_k^\tau : (t, x) \mapsto \hat{\mathbf{u}}_k(t, \tau(x)),$$

$$\mathbf{w}_k : (t, x) \mapsto e^{z_1 L_1} \hat{\mathbf{u}}_k - \hat{\mathbf{u}}_k^\tau.$$

Once more by virtue of the Harnack inequality and the periodicity of the coefficients of \mathcal{Q} , $(\hat{\mathbf{u}}_k)_{k \in \mathbb{N}}$ is globally bounded. By periodicity of the coefficients of \mathcal{Q} , it satisfies:

$$\mathcal{Q}(t + t_k, x + y_k) \hat{\mathbf{u}}_k(t, x) = \lambda_1 \hat{\mathbf{u}}_k(t, x) \quad \text{for all } (t, x) \in \mathbb{R} \times \mathbb{R}^n, \quad k \in \mathbb{N}.$$

Therefore, by classical regularity estimates [52], $(\hat{\mathbf{u}}_k)_{k \in \mathbb{N}}$ converges up to a diagonal extraction to $\hat{\mathbf{u}}_\infty \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, [\mathbf{0}, \infty))$ which satisfies:

$$\mathcal{Q}(t + t_\infty, x + y_\infty) \hat{\mathbf{u}}_\infty(t, x) = \lambda_1 \hat{\mathbf{u}}_\infty(t, x) \quad \text{for all } (t, x) \in \mathbb{R} \times \mathbb{R}^n,$$

i.e.

$$\mathcal{Q}(t, x) \hat{\mathbf{u}}_\infty(t - t_\infty, x - y_\infty) = \lambda_1 \hat{\mathbf{u}}_\infty(t - t_\infty, x - y_\infty) \quad \text{for all } (t, x) \in \mathbb{R} \times \mathbb{R}^n.$$

Moreover, $\hat{u}_{i,\infty}(0, 0) = 1$, whence $\hat{\mathbf{u}}_\infty$ is nonzero. By the strong maximum principle (see Proposition 2.1), it is in fact positive.

We can now extend the family (\mathbf{w}_k) in $\mathbb{N} \cup \{\infty\}$ with $\mathbf{w}_\infty = e^{z_1 L_1} \hat{\mathbf{u}}_\infty - \hat{\mathbf{u}}_\infty^\tau$. Since, for all $k \in \bar{\mathbb{N}}$,

$$\mathbf{w}_k = \hat{\mathbf{u}}_k \circ (e^{z_1 L_1} \mathbf{1} - \mathbf{v}_k), \quad \text{where } \mathbf{v}_k : (t, x) \mapsto \mathbf{v}(t + t_k, x + x_k),$$

we deduce by definition of z_1 that $\mathbf{w}_\infty \geq \mathbf{0}$ with $w_{i,\infty}(0, 0) = 0$. Moreover, \mathbf{w}_∞ satisfies the same equation than $\hat{\mathbf{u}}_\infty$. Therefore, by virtue of the strong maximum principle, \mathbf{w}_∞ is the zero function. This exactly means that $e^{z_1 L_1} \hat{\mathbf{u}}_\infty = \hat{\mathbf{u}}_\infty^\tau$.

It is now clear that $\mathbf{u}^1 : (t, x) \mapsto \hat{\mathbf{u}}_\infty(t - t_\infty, x - y_\infty)$ is positive, time periodic, a solution of $\mathcal{Q}\mathbf{u}^1 = \lambda_1 \mathbf{u}^1$, and that the function $(t, x) \mapsto e^{-z_1 x_1} \mathbf{u}^1(t, x)$ is L_1 -periodic with respect to x_1 . Indeed, for any $(t, x) \in \mathbb{R} \times \mathbb{R}^n$,

$$\begin{aligned} e^{-z_1(x_1 + L_1)} \mathbf{u}^1(t, x + L_1 e_1) &= e^{-z_1 x_1} e^{-z_1 L_1} \mathbf{u}^1(t, \tau(x)) \\ &= e^{-z_1 x_1} e^{-z_1 L_1} \hat{\mathbf{u}}_\infty^\tau(t - t_\infty, x - y_\infty) \\ &= e^{-z_1 x_1} \hat{\mathbf{u}}_\infty(t - t_\infty, x - y_\infty) \\ &= e^{-z_1 x_1} \mathbf{u}^1(t, x). \end{aligned}$$

The first part of the proof is done.

Next, we iterate this construction, replacing \mathbf{u} by \mathbf{u}^1 , in order to obtain a new generalized principal eigenfunction \mathbf{u}^2 such that $(t, x) \mapsto e^{-z_1 x_1} e^{-z_2 x_2} \mathbf{u}^2(t, x)$ is L_1 -periodic with respect to x_1 and L_2 -periodic with respect to x_2 . Iterating again, we finally obtain $z \in \mathbb{R}^n$ and $\mathbf{u}^n \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ such that \mathbf{u}^n is a generalized principal eigenfunction associated with λ_1 and such that $e_{-z} \mathbf{u}^n$ is space periodic. The uniqueness of the eigenpair $(\lambda_{1,z}, \mathbf{u}_z)$, up to multiplication of \mathbf{u}_z by a constant, yields finally $\lambda_1 = \lambda_{1,z}$ and $e_{-z} \mathbf{u}^n \in \text{span}(\mathbf{u}_z)$. \square

Corollary 3.10. *The generalized principal eigenvalue λ_1 satisfies:*

$$(17) \quad \lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}$$

and there exists a unique $z \in \mathbb{R}^n$ such that $\lambda_1 = \lambda_{1,z}$.

Proof. Proposition 3.9 already shows that λ_1 is in the image of $z \mapsto \lambda_{1,z}$ and Corollary 3.6 already shows that $z \mapsto \lambda_{1,z}$ is strictly concave. Thus it only remains to show $\lambda_1 \geq \sup_{z \in \mathbb{R}^n} \lambda_{1,z}$. This is actually obvious, since the equality $\mathcal{Q}(e_z \mathbf{u}_z) = \lambda_{1,z} e_z \mathbf{u}_z$ (which is just the definition of the eigenpair $(\lambda_{1,z}, \mathbf{u}_z)$) directly implies, in view of the definition of λ_1 , the inequality $\lambda_1 \geq \lambda_{1,z}$. \square

Remark 3.4. Let

$$E = \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)) \quad \mathcal{Q}\mathbf{u} = \lambda \mathbf{u} \right\}$$

and denote $\Lambda \subset \mathbb{R}$ the image of $z \in \mathbb{R}^n \mapsto \lambda_{1,z}$. From the equality $\mathcal{Q}e_z \mathbf{u}_z = \lambda_{1,z} e_z \mathbf{u}_z$, the following set inclusions hold true:

$$\Lambda \subset E \subset \left\{ \lambda \in \mathbb{R} \mid \exists \mathbf{u} \in \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty)) \quad \mathcal{Q}\mathbf{u} \geq \lambda \mathbf{u} \right\}.$$

By strict concavity, $\Lambda = (-\infty, \max \lambda_{1,z}]$, and since $\lambda_1 = \max \lambda_{1,z}$ is by definition the supremum of the larger set above, all inclusions above are actually set equalities.

This shows in particular that the set E of eigenvalues of \mathcal{Q} acting on the set $\mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ is $(-\infty, \lambda_1]$. This is of course in striking contrast with the case of smooth bounded domains, where the Krein–Rutman theorem can be applied and the principal eigenvalue is unique. For the same result in the elliptic case with general spatial heterogeneities, refer to Berestycki–Rossi [16, Theorem 1.4] (scalar setting) and Arapostathis–Biswas–Pradhan [9, Theorem 1.2] (cooperative vector setting).

Proposition 3.11. *The generalized principal eigenvalue λ'_1 satisfies:*

$$(18) \quad \lambda'_1 = \lambda_{1,0}.$$

Proof. Since $\mathbf{u}_0 = e_0 \mathbf{u}_0$ is globally bounded, we can use it as test function in the definition of λ'_1 and obtain $\lambda'_1 \leq \lambda_{1,0}$.

Now, we assume by contradiction that this inequality is actually strict, so that by definition of λ'_1 , there exists $\mu \in (\lambda'_1, \lambda_{1,0})$ and $\mathbf{u} \in \mathcal{W}^{1,\infty} \cap \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ such that $\mathcal{Q}\mathbf{u} \leq \mu \mathbf{u}$.

We can now define

$$\kappa^* = \inf \{ \kappa > 0 \mid \kappa \mathbf{u}_0 - \mathbf{u} \gg \mathbf{0} \}$$

and study the sign of $\mathbf{v} = \kappa^* \mathbf{u}_0 - \mathbf{u}$. This function satisfies

$$\mathcal{Q}\mathbf{v} \geq (\lambda_{1,0} - \mu) \kappa^* \mathbf{u}_0 + \mu \mathbf{v},$$

is time periodic and nonnegative, and by optimality there exists $((t_k, x_k))_{k \in \mathbb{N}} \in ([0, T] \times \mathbb{R}^n)^{\mathbb{N}}$ and $i \in [N]$ such that

$$v_i(t_k, x_k) \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Moreover, there exists $(y_k)_{k \in \mathbb{N}}$ such that, for all $k \in \mathbb{N}$, $x_k - y_k \in L_1 \mathbb{Z} \times \cdots \times L_n \mathbb{Z}$. Up to extraction, we assume that $(t_k, y_k) \rightarrow (t_\infty, y_\infty) \in \overline{\Omega_{\text{per}}}$.

Define

$$\mathbf{v}_k : (t, x) \mapsto \mathbf{v}(t + t_k, x + x_k) \quad \text{for all } k \in [N].$$

By standard regularity estimates [52], $(\mathbf{v}_k)_{k \in \mathbb{N}}$ converges up to a diagonal extraction to a function $\mathbf{v}_\infty \in \mathcal{L}^\infty \cap \mathcal{C}_{t\text{-per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, [\mathbf{0}, \infty))$ which satisfies $v_{i,\infty}(0, 0) = 0$ and, for all $(t, x) \in \mathbb{R} \times \mathbb{R}^n$,

$$(\mathcal{Q} - \mu)(t + t_\infty, x + y_\infty) \mathbf{v}_\infty(t, x) \geq (\lambda_{1,0} - \mu) \kappa^* \min_{i \in [N]} \min_{\overline{\Omega_{\text{per}}}} (u_{0,i}) \mathbf{1} \gg \mathbf{0}.$$

By virtue of the strong maximum principle (see Proposition 2.1), $\mathbf{v}_\infty = \mathbf{0}$, but then this contradicts the preceding inequality. This ends the proof. \square

Remark 3.5. It is natural to investigate the equality between $\lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}$ and $\lambda'_1 = \lambda_{1,0}$. The scalar counter-example with constant coefficients $\mathcal{Q} = \partial_t - \partial_{xx} + q\partial_x - l$ shows that both outcomes are possible, since $\lambda_{1,z} = z(q - z) - l$ is maximal at $z = 0$ if and only if $q = 0$. Identifying precise conditions for the maximality at $z = 0$ becomes then one of our main goals. A very recent contribution by Griette and Matano [40, Proposition 4.1] shows that in the vector setting, the absence of advection is not enough.

Their two-dimensional counter-example in one-dimensional space is:

$$\mathcal{Q} = \partial_t - \partial_x \left(\text{diag} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \partial_x \right) - \begin{pmatrix} r_1 - \frac{1}{\varepsilon} p & \frac{1}{\varepsilon} (1-p) \\ \frac{1}{\varepsilon} p & r_2 - \frac{1}{\varepsilon} (1-p) \end{pmatrix}$$

with a_1, a_2, r_1, r_2 and p periodic functions of x . As $\varepsilon \rightarrow 0$, locally uniformly with respect to z ,

$$\lambda_{1,z}(\mathcal{Q}) \rightarrow \lambda_{1,z}(-\partial_x(a\partial_x) + q\partial_x - (r - q'))$$

with

$$a = (1-p)a_1 + pa_2, \quad r = (1-p)r_1 + pr_2, \quad q = (a_1 - a_2)p'.$$

Under the condition $\int_0^{L_1} q/a \neq 0$, the limit is not maximal at $z = 0$ [40, Appendix A], whence $\lambda_{1,z}(\mathcal{Q})$ is also not maximal at $z = 0$ when ε is sufficiently small. For more details, we refer to [40].

3.1.5. Rough estimates. Here we state rough upper and lower estimates that will be used later on in the proofs. The more precise estimates of Subsection 1.5.5, that use special assumptions on the coefficients of \mathcal{Q} , will be proved later.

Using the cooperativity assumption (A₂), the min–max characterization of Proposition 3.4, the equalities $\lambda_1 = \max \lambda_{1,z}$ and $\lambda'_1 = \lambda_{1,0}$ of Corollary 3.10 and Proposition 3.11 respectively, and the corresponding scalar results [59], we deduce the following corollary which relates the generalized principal eigenvalues of the operator \mathcal{Q} to the generalized principal eigenvalues of the scalar operators $\mathcal{P}_i - l_{i,i}$.

Corollary 3.12. *For all $z \in \mathbb{R}^n$,*

$$\lambda_{1,z}(\mathcal{Q}) \leq \min_{i \in [N]} \lambda_{1,z}(\mathcal{P}_i - l_{i,i}).$$

Consequently,

$$\lambda_1(\mathcal{Q}) \leq \min_{i \in [N]} \lambda_1(\mathcal{P}_i - l_{i,i}) \quad \text{and} \quad \lambda'_1(\mathcal{Q}) \leq \min_{i \in [N]} \lambda'_1(\mathcal{P}_i - l_{i,i}).$$

Remark 3.6. Rougher but more explicit estimates can subsequently be derived by considering constant test functions in the min–max characterization of $\lambda_{1,z}(\mathcal{P}_i - l_{i,i})$

and the discrete Cauchy–Schwarz inequality:

$$\begin{aligned}
 \lambda_{1,z}(\mathcal{P}_i - l_{i,i}) &\leq \frac{\max}{\Omega_{\text{per}}}(-z \cdot A_i z - \nabla \cdot (A_i z) + q_i \cdot z - l_{i,i}) \\
 &\leq -\frac{\min}{\Omega_{\text{per}}}(z \cdot A_i z) + \frac{\max}{\Omega_{\text{per}}}(-\nabla \cdot (A_i z)) + \frac{\max}{\Omega_{\text{per}}}(q_i \cdot z) - \frac{\min}{\Omega_{\text{per}}}l_{i,i} \\
 &\leq -l_{i,i} - \frac{\min}{\Omega_{\text{per}}} \min_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y) |z|^2 + \frac{\max}{\Omega_{\text{per}}} \sum_{\alpha=1}^n \partial_{\alpha} \left(\sum_{\beta=1}^n -A_{\alpha,\beta}^i z_{\beta} \right) + \frac{\max}{\Omega_{\text{per}}} (|q_i|) |z| \\
 &\leq -l_{i,i} - \frac{\min}{\Omega_{\text{per}}} \min_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y) |z|^2 + \sum_{\alpha=1}^n \frac{\max}{\Omega_{\text{per}}} \sum_{\beta=1}^n \partial_{\alpha} (-A_{\alpha,\beta}^i) z_{\beta} + \| |q_i| \| |z| \\
 &\leq -l_{i,i} - \frac{\min}{\Omega_{\text{per}}} \min_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y) |z|^2 + \sum_{\alpha=1}^n \left\| \left(\sum_{\beta=1}^n |\partial_{\alpha} A_{\alpha,\beta}^i|^2 \right)^{1/2} \right\| |z| + \| |q_i| \| |z|,
 \end{aligned}$$

where \mathbf{L} is defined in (A₂) and the notation $\| \cdot \|$ refers to the norm in the space $\mathcal{L}^{\infty}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R})$.

Another way to obtain rough upper and lower estimates consists in using

$$\mathbf{L} + \left(\frac{\min}{\Omega_{\text{per}}} \min_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y) |z|^2 - K |z| \right) \mathbf{I} \leq \mathbf{L} + \text{diag} (z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z)$$

and

$$\mathbf{L} + \text{diag} (z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z) \leq \bar{\mathbf{L}} + \left(\frac{\max}{\Omega_{\text{per}}} \max_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y) |z|^2 + K |z| \right) \mathbf{I},$$

where

$$K = \sum_{\alpha=1}^n \left\| \left(\sum_{\beta=1}^n |\partial_{\alpha} A_{\alpha,\beta}^i|^2 \right)^{1/2} \right\| + \| |q_i| \|.$$

Although \mathbf{L} might not be irreducible, its Perron–Frobenius eigenvalue is still well-defined by continuous extension; it admits nonnegative nonzero eigenvectors that can be used as sub-solutions. Using Perron–Frobenius eigenvectors as test functions in the min–max and max–min formulas of Proposition 3.4, we deduce the following corollary.

Corollary 3.13. *Let*

$$\begin{aligned}
 A &= \max_{i \in [N]} \left(\max \left(\frac{\max}{\Omega_{\text{per}}} \max_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y), \frac{1}{\min_{\Omega_{\text{per}}} \min_{y \in \mathbb{S}^{n-1}} (y \cdot A_i y)} \right) \right) > 0, \\
 B &= \max_{i \in [N]} \left(\sum_{\alpha=1}^n \left(\sum_{\beta=1}^n \frac{\max}{\Omega_{\text{per}}} |\partial_{\alpha} A_{\alpha,\beta}^i|^2 \right)^{1/2} + \frac{\max}{\Omega_{\text{per}}} |q_i| \right) \geq 0.
 \end{aligned}$$

Then, for all $z \in \mathbb{R}^n$,

$$-\lambda_{\text{PF}}(\bar{\mathbf{L}}) - A |z|^2 - B |z| \leq \lambda_{1,z} \leq -\lambda_{\text{PF}}(\mathbf{L}) - \frac{1}{A} |z|^2 + B |z|.$$

3.2. Asymptotic dependence: proof of Theorems 1.5–1.9. Proposition 3.5 already proves Theorem 1.3. Below, we prove the remaining theorems on coefficient dependence.

3.2.1. Continuity: proof of Theorem 1.5. We begin with the proof of Theorem 1.5, whose statement is recalled below.

Proposition 3.14. *Let $\mathbf{L}^\Delta \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ be a block upper triangular essentially nonnegative matrix. Let $N' \in [N]$ and $(N_k)_{k \in [N'-1]}$ such that*

$$N_0 = 0 < 1 \leq N_1 \leq N_2 \leq \dots \leq N_{N'-1} \leq N_{N'} = N$$

and such that

$$(l_{i,j}^\Delta)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2}$$

is the k -th diagonal block of \mathbf{L}^Δ (with the convention $[0] = \emptyset$). Assume

$$\left(\max_{(t,x) \in \Omega_{\text{per}}} l_{i,j}^\Delta(t,x) \right)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2} \quad \text{is irreducible for all } k \in [N'].$$

Let

$$\mathbf{Q}_k = \text{diag}(\mathcal{P}_i)_{i \in [N_k] \setminus [N_{k-1}]} - (l_{i,j}^\Delta)_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2} \quad \text{for all } k \in [N'].$$

Then, as $\mathbf{L} \rightarrow \mathbf{L}^\Delta$ in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$,

$$\lambda_{1,z}(\mathbf{Q}) \rightarrow \min_{k \in [N']} \lambda_{1,z}(\mathbf{Q}_k) \quad \text{for all } z \in \mathbb{R}^n,$$

$$\lambda_1(\mathbf{Q}) \rightarrow \max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathbf{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathbf{Q}_k).$$

Proof. Step 1: the special case $z=0$. Let $(\mathbf{L}_p)_{p \in \mathbb{N}}$ be a sequence of matrices satisfying (A_2) , (A_3) and that converges to \mathbf{L}^Δ in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$. Denote $\mathbf{Q}_p = \text{diag}(\mathcal{P}_i) - \mathbf{L}_p$ and $\mathbf{Q}_{k,p} = \text{diag}(\mathcal{P}_i)_{i \in [N_k] \setminus [N_{k-1}]} - (l_{p,i,j})_{(i,j) \in ([N_k] \setminus [N_{k-1}])^2}$.

Since

$$0 \leq \mathbf{L}_p \leq \left(\sup_{p \in \mathbb{N}} \max_{(t,x) \in \Omega_{\text{per}}} l_{p,i,j}(t,x) \right),$$

we can derive from the max–min and min–max characterizations of $\lambda_1(\mathbf{L}_p)$ (see Proposition 3.4) uniform bounds on $(\lambda_1(\mathbf{L}_p))_{p \in \mathbb{N}}$. Therefore up to extraction this sequence converges to a limit $\lambda \in \mathbb{R}$. Similarly, up to extraction, the associated generalized principal eigenfunction $\mathbf{u}_p \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ with normalization $|\mathbf{u}_p(0,0)| = 1$ converges to a nonnegative nonzero limit $\mathbf{u} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, [\mathbf{0}, \infty))$ satisfying the same normalization and satisfying

$$\text{diag}(\mathcal{P}_i)\mathbf{u} - \mathbf{L}^\Delta \mathbf{u} = \lambda \mathbf{u}.$$

Note that for each $k \in [N']$,

$$\begin{aligned} \mathbf{Q}_k(u_i)_{i \in [N_k] \setminus [N_{k-1}]} &= \lambda(u_i)_{i \in [N_k] \setminus [N_{k-1}]} + \left(\sum_{j \in [N_{k-1}] \cup [N] \setminus [N_k]} l_{i,j}^\Delta u_j \right)_{i \in [N_k] \setminus [N_{k-1}]} \\ &\geq \lambda(u_i)_{i \in [N_k] \setminus [N_{k-1}]} \end{aligned}$$

Therefore, from the strong maximum principle of Proposition 2.1, either $(u_i)_{i \in [N_k] \setminus [N_{k-1}]} = \mathbf{0}$ or $(u_i)_{i \in [N_k] \setminus [N_{k-1}]} \gg \mathbf{0}$. For all $k \in [N']$ such that $(u_i)_{i \in [N_k] \setminus [N_{k-1}]} \gg \mathbf{0}$, it follows from the characterization of $\lambda_{1,\text{per}}(\mathbf{Q}_k)$ (see Proposition 3.4) that $\lambda \leq \lambda_{1,\text{per}}(\mathbf{Q}_k)$.

Since \mathbf{u} is nonzero, there exists at least one such k . Let $I \subset [N']$ be the set of all such k and let $J = [N'] \setminus I$.

If $N' \in I$, then from the special block upper triangular form of \mathcal{Q} , $\lambda = \lambda'_1(\mathcal{Q}_{N'})$. Otherwise, there exists $k \in [N' - 1] \cap I$. It follows then from a classical inductive argument that there exists indeed $k \in [N']$ such that $\lambda = \lambda'_1(\mathcal{Q}_k)$.

Now, assume by contradiction that there exists $k' \in [N']$ such that $\lambda > \lambda'_1(\mathcal{Q}_{k'})$. Let $\eta = \lambda - \lambda'_1(\mathcal{Q}_{k'}) > 0$. Let $\mathbf{u}_{k'}$ be a periodic principal eigenfunction associated with $\lambda_{1,\text{per}}(\mathcal{Q}_{k'})$. Let $\underline{\mathbf{u}}$ be defined as

$$\underline{u}_i = \begin{cases} u_{k',i-N_{k'-1}} & \text{if } i \in [N_{k'}] \setminus [N_{k'-1}], \\ 0 & \text{otherwise.} \end{cases}$$

Then, for all $i \in [N]$,

$$(\mathcal{Q}_p \underline{\mathbf{u}})_i = \begin{cases} ((\mathcal{Q}'_{k',p} - \mathcal{Q}'_{k'}) \underline{\mathbf{u}})_i + \lambda'_1(\mathcal{Q}_{k'}) \underline{u}_i & \text{if } i \in [N_{k'}] \setminus [N_{k'-1}], \\ - \sum_{j \in [N_{k'}] \setminus [N_{k'-1}]} l_{p,i,j} \underline{u}_j & \text{otherwise.} \end{cases}$$

On one hand,

$$- \sum_{j \in [N_{k'}] \setminus [N_{k'-1}]} l_{p,i,j} \underline{u}_j \leq 0 = \underline{u}_i \quad \text{for all } i \notin [N_{k'}] \setminus [N_{k'-1}].$$

On the other hand, by convergence of \mathbf{L}_p and the Harnack inequality of Proposition 2.4 applied to the fully coupled operator $\mathcal{Q}'_{k'}$, we can assume that $p \in \mathbb{N}$ is so large that, for all $i \in [N_{k'}] \setminus [N_{k'-1}]$,

$$((\mathcal{Q}'_{k',p} - \mathcal{Q}'_{k'}) \underline{\mathbf{u}})_i = ((\mathbf{L}'_{k',p} - \mathbf{L}'_{k'} \Delta) \underline{\mathbf{u}})_i \leq \frac{\eta}{2} \underline{u}_i,$$

where $\mathbf{L}'_{k',p} = (l_{p,i,j})_{(i,j) \in ([N_{k'}] \setminus [N_{k'-1}])^2}$ and $\mathbf{L}'_{k'} = (l'_{i,j})_{(i,j) \in ([N_{k'}] \setminus [N_{k'-1}])^2}$. Hence

$$\mathcal{Q}_p \underline{\mathbf{u}} \leq \left(\lambda'_1(\mathcal{Q}_{k'}) + \frac{\eta}{2} \right) \underline{\mathbf{u}} = \left(\lambda - \frac{\eta}{2} \right) \underline{\mathbf{u}}.$$

If $\lambda'_1(\mathcal{Q}_p) > \lambda - \frac{\eta}{2}$, then we can study $\kappa^* \mathbf{u}_p - \underline{\mathbf{u}}$ with

$$\kappa^* = \frac{\max_{i \in [N]} \max_{(t,x) \in \Omega_{\text{per}}} u_i(t,x)}{\min_{i \in [N]} \min_{(t,x) \in \Omega_{\text{per}}} u_{p,i}(t,x)} > 0$$

and, by full coupling of \mathcal{Q}_p , deduce a contradiction from the strong maximum principle. Hence $\lambda'_1(\mathcal{Q}_p) \leq \lambda - \frac{\eta}{2}$. But now, assuming in addition that p is so large that $\lambda'_1(\mathcal{Q}_p) > \lambda - \frac{\eta}{3}$, we find a contradiction. Therefore, for all $k' \in [N']$, $\lambda \leq \lambda'_1(\mathcal{Q}_{k'})$, or in other words:

$$\lambda \leq \min_{k' \in [N']} \lambda'_1(\mathcal{Q}_{k'}).$$

Combining this with $\lambda = \lambda'_1(\mathcal{Q}_k)$, we deduce that the preceding inequality is an equality.

This argument shows that any convergent subsequence of the sequence $(\lambda'_1(\mathbf{L}_p))_{p \in \mathbb{N}}$ converges to $\min_{k \in [N']} \lambda'_1(\mathcal{Q}_k)$. The conclusion follows. \square

Step 2: the general case $z \in \mathbb{R}^n$. In view of

$$\lambda_{1,z}(\mathcal{Q}) = \lambda'_1(\mathcal{Q}_z) = \lambda'_1(\mathcal{Q} - \text{diag}((A_i + A_i^T)z \cdot \nabla + z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z)),$$

in order to prove the convergence of $\lambda_{1,z}$ for any $z \in \mathbb{R}^n$, we only have to apply the preceding step to the operator \mathcal{Q}_z . \square

Step 3: convergence of $\lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}$. Since all $z \mapsto \lambda_{1,z}(\mathcal{Q}_p)$, $p \in \mathbb{N}$, are concave, the pointwise convergence is automatically improved to locally uniform convergence.

On one hand, recall from Corollary 3.13 and the ellipticity assumption (A_1) that there exists $A > 0$, $B \geq 0$ and $C \in \mathbb{R}$ such that, for all $p \in \mathbb{N}$,

$$\lambda_{1,z}(\mathcal{Q}_p) \leq -A|z|^2 + B|z| + C \quad \text{for all } p \in \mathbb{N}.$$

On the other hand, for all $p \in \mathbb{N}$, $\lambda_1(\mathcal{Q}_p) \geq \lambda'_1(\mathcal{Q}_p)$. In particular, $\lambda_1(\mathcal{Q}_p) \geq \inf_{p \in \mathbb{N}} \lambda'_1(\mathcal{Q}_p)$ and this lower bound is finite by virtue of Step 1 above.

Consequently, for all $p \in \mathbb{N}$, the point z_p where the maximum is achieved (which is indeed uniquely defined, see Corollary 3.10) is necessarily in the set Z defined as:

$$Z = \left\{ z \in \mathbb{R}^n \mid \inf_{p \in \mathbb{N}} \lambda'_1(\mathcal{Q}_p) \leq -A|z|^2 + B|z| + C \right\}.$$

This set is compact.

To conclude, from the already established equality:

$$\lim_{p \rightarrow +\infty} \lambda_{1,z}(\mathcal{Q}_p) = \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k),$$

and from the uniform convergence in Z and the concavity in \mathbb{R}^n , we deduce

$$\begin{aligned} \lim_{p \rightarrow +\infty} \lambda_1(\mathcal{Q}_p) &= \lim_{p \rightarrow +\infty} \max_{z \in Z} \lambda_{1,z}(\mathcal{Q}_p) \\ &= \max_{z \in Z} \lim_{p \rightarrow +\infty} \lambda_{1,z}(\mathcal{Q}_p) \\ &= \max_{z \in \mathbb{R}^n} \lim_{p \rightarrow +\infty} \lambda_{1,z}(\mathcal{Q}_p) \\ &= \max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k). \end{aligned}$$

Finally, from the inequality $\lambda_{1,z}(\mathcal{Q}_k) \leq \lambda_1(\mathcal{Q}_k)$ for all k and z , it follows that

$$\min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathcal{Q}_k),$$

whence

$$\max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathcal{Q}_k).$$

\square

This ends the proof. \square

Remark 3.7. It should be noted here that the only part of this proof that is seemingly specific to the case of Hölder-continuous coefficients is the Harnack inequality provided by Proposition 2.4. However, this is merely used to show that $\kappa^* \in (0, \infty)$. In the case of coefficients in \mathcal{L}^∞ , a simple reasoning by contradiction shows that $\kappa^* \in (0, \infty)$ is still true. A minor adaptation of the arguments then shows the continuity of λ'_1 for the weak- \star \mathcal{L}^∞ convergence of coefficients.

Remark 3.8. We will use repeatedly the arguments of Step 2 and Step 3 above in what follows, in order to deduce the convergence of $\lambda_{1,z}$ and λ_1 when the convergence of λ'_1 has been established.

Note however that the estimate $\lambda_{1,z}(\mathcal{Q}) \leq -A|z|^2 + B|z| + C$ with $A > 0$ and $B \geq 0$ of Corollary 3.13 becomes useless when the diffusion matrices A_i vanish. This is consistent with the fact that, in Theorem 1.6, the convergence of λ_1 is in general false.

Remark 3.9. The inequality

$$\max_{z \in \mathbb{R}^n} \min_{k \in [N']} \lambda_{1,z}(\mathcal{Q}_k) \leq \min_{k \in [N']} \lambda_1(\mathcal{Q}_k)$$

is strict in some cases. Consider for instance the following space-time homogeneous, one-dimensional, two-component counter-example:

$$\mathcal{Q} = \mathcal{Q}_\varepsilon = \text{diag} \left(\begin{pmatrix} \partial_t u_1 - \partial_{xx} u_1 \\ \partial_t u_2 - \partial_{xx} u_2 + 2\partial_x u_2 - 1 \end{pmatrix} \right) - \varepsilon \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix},$$

where $\varepsilon > 0$. The operator is diagonal if $\varepsilon = 0$. By standard reduction (cf. (10)), the two scalar operators on the diagonal, $\mathcal{Q}_1 = \partial_t - \partial_{xx}$ and $\mathcal{Q}_2 = \partial_t - \partial_{xx} + 2\partial_x - 1$, satisfy $\lambda_{1,z}(\mathcal{Q}_1) = -z^2$ and $\lambda_{1,z}(\mathcal{Q}_2) = -(z-1)^2$. In particular, $\lambda_1(\mathcal{Q}_1) = \lambda_1(\mathcal{Q}_2) = 0$. However, the function $z \mapsto \min(-z^2, -(z-1)^2)$ coincides with

$$z \mapsto \begin{cases} -(z-1)^2 & \text{if } z < 1/2, \\ -z^2 & \text{if } z \geq 1/2, \end{cases}$$

whose maximal value is $-1/4 < 0$, which is attained at $1/2$.

3.2.2. Vanishing diffusion and advection: proof of Theorem 1.6. Below, we prove Theorem 1.6 on vanishing diffusion and advection rates.

Following the statement of the theorem, we fix a function $\mathbf{f} \in \mathcal{C}^1([0, +\infty), [\mathbf{0}, \infty))$ such that $\mathbf{f}^{-1}(\{\mathbf{0}\}) = \{0\}$ and $\mathbf{f}'(0) \neq \mathbf{0}$, as well as a family $((q_i^\varepsilon)_{i \in [N]})_{\varepsilon \geq 0}$ such that, for all $\varepsilon \geq 0$, $(q_i^\varepsilon)_{i \in [N]} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$ and such that $(q_i^\varepsilon)_{i \in [N]} \rightarrow (q_i^0)_{i \in [N]}$ in $\mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^n)$ as $\varepsilon \rightarrow 0$. Note that, by assumption, $\mathbf{f}'(0) > \mathbf{0}$.

We denote \mathcal{Q}_ε the operator \mathcal{Q} where $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ and \mathbf{L} are replaced respectively by $(f_i(\varepsilon)^2 A_i)_{i \in [N]}$, $(f_i(\varepsilon) q_i^\varepsilon)_{i \in [N]}$ and a parameterized matrix \mathbf{L}^ε that still satisfies (A_2) – (A_4) and that converges uniformly to $\mathbf{L}^0 = \mathbf{L}$ as $\varepsilon \rightarrow 0$. We use the (slightly abusing) notations $A_i(x) : t \mapsto A_i(t, x)$, $q_i^\varepsilon(x) : t \mapsto q_i^\varepsilon(t, x)$ and $\mathbf{L}^\varepsilon(x) : t \mapsto \mathbf{L}^\varepsilon(t, x)$.

Recall that, for any $z \in \mathbb{R}^n$,

$$\begin{aligned} e_{-z} \mathcal{Q}_\varepsilon e_z &= \partial_t - \text{diag} (f_i(\varepsilon)^2 \nabla \cdot (A_i \nabla)) \\ &\quad - \text{diag} (f_i(\varepsilon) (2f_i(\varepsilon) A_i z - q_i^\varepsilon) \cdot \nabla) \\ &\quad - \text{diag} ((f_i(\varepsilon)^2 z \cdot A_i z + f_i(\varepsilon)^2 \nabla \cdot (A_i z) - f_i(\varepsilon) q_i^\varepsilon \cdot z)) - \mathbf{L}^\varepsilon \end{aligned}$$

Therefore the convergence in the case where the coefficients of \mathcal{Q}_ε do not depend on space is a straightforward consequence of (8) and of the continuity of the periodic principal eigenvalue (see Theorem 1.5).

The above expansion of $e_{-z} \mathcal{Q}_\varepsilon e_z$ also shows that, thanks to the replacement of \mathbf{L} by \mathbf{L}^ε , this operator has again the form of \mathcal{Q}_ε . Therefore, by virtue of the above

expansion and of the equality, for all $x \in [0, L]$,

$$\lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right) = \lambda'_1 \left(\partial_t - \text{diag}(f'_i(0)^2 \nabla \cdot (A_i(x) \nabla) - f'_i(0) q_i^0(x) \cdot \nabla) - \mathbf{L}(x) \right),$$

the following result is actually sufficient to prove the remaining part of Theorem 1.6.

Proposition 3.15. *For any $x \in [0, L]$, let*

$$\tilde{\mathcal{Q}}(x) = \partial_t - \text{diag}(f'_i(0)^2 \nabla \cdot (A_i(x) \nabla) - f'_i(0) q_i^0(x) \cdot \nabla) - \mathbf{L}(x).$$

Then the generalized principal eigenvalue $\lambda'_1(\mathcal{Q}_\varepsilon)$ satisfies:

$$\min_{x \in [0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right) \leq \liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_\varepsilon) \leq \limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_\varepsilon) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)).$$

Proof. The proof is done in three steps.

Step 1: the pointwise irreducibility of every \mathbf{L}^ε can be assumed without loss of generality.

Assume the result has been proved provided every $\mathbf{L}^\varepsilon(t, x)$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$.

Define

$$\mathbf{L}^\varepsilon : s \in [0, +\infty) \mapsto \mathbf{L}^\varepsilon + (e^s - 1) \mathbf{1}_{N \times N} - (e^s - 1) \mathbf{I}.$$

Obviously, $\mathbf{L}^\varepsilon(0) = \mathbf{L}^\varepsilon$ and, for all $s \in (0, +\infty)$, $\mathbf{L}^\varepsilon(s, t, x)$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$. Moreover, by virtue of Propositions 3.5, 3.4 and 3.14, the periodic principal eigenvalue $\lambda'_1(\varepsilon, s)$ associated with the operator

$$\mathcal{Q}_{\varepsilon, s} = \partial_t - \text{diag}(f_i(\varepsilon)^2 \nabla \cdot (A_i \nabla) - f_i(\varepsilon) q_i^\varepsilon \cdot \nabla) - \mathbf{L}^\varepsilon(s)$$

is, as a function of s , continuous in $[0, +\infty)$, decreasing in $[0, +\infty)$, strictly concave in $[0, +\infty)$.

In particular, by decreasing monotonicity, for any $s > 0$,

$$\liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, 0}) \geq \liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, s}) \geq \min_{x \in [0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(s, x) \right).$$

Passing to the limit $s \rightarrow 0$ in the right-hand side, which is continuous indeed with respect to s , shows the first inequality:

$$\min_{x \in [0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right) \leq \liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_\varepsilon).$$

Similarly, defining for all $s \in (0, +\infty)$ and $x \in [0, L]$ the operator

$$\tilde{\mathcal{Q}}_s(x) = \partial_t - \text{diag}(f'_i(0)^2 \nabla \cdot (A_i(x) \nabla) - f'_i(0) q_i^0(x) \cdot \nabla) - \mathbf{L}(s, x),$$

we find by monotonicity

$$\limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, s}) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}_s(x)) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)).$$

To pass to the limit $s \rightarrow 0$ in the left-hand side, we use the concavity as follows. Let $\gamma > 0$ and $s \in (0, 1)$. There exists $\varepsilon_{s, \gamma} > 0$ such that

$$\lambda'_1(\mathcal{Q}_{\varepsilon, s}) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)) + \gamma \quad \text{for all } \varepsilon \in (0, \varepsilon_{s, \gamma}).$$

Let $\varepsilon \in (0, \varepsilon_{s,\gamma})$. By concavity, using $s < 1$,

$$\lambda'_1(\varepsilon, 2) - \lambda'_1(\varepsilon, 1) \leq \lim_{\substack{s'' < s \\ s'' \rightarrow s}} \frac{\lambda'_1(\varepsilon, s'') - \lambda'_1(\varepsilon, s)}{s'' - s} \leq 0$$

and for all $s' \in (0, s)$,

$$\lambda'_1(\mathcal{Q}_{\varepsilon, s'}) \leq \lambda'_1(\mathcal{Q}_{\varepsilon, s}) + \left(\lim_{\substack{s'' < s \\ s'' \rightarrow s}} \frac{\lambda'_1(\varepsilon, s'') - \lambda'_1(\varepsilon, s)}{s'' - s} \right) (s' - s)$$

i.e.

$$\lambda'_1(\mathcal{Q}_{\varepsilon, s'}) + \left(\lim_{\substack{s'' < s \\ s'' \rightarrow s}} \frac{\lambda'_1(\varepsilon, s'') - \lambda'_1(\varepsilon, s)}{s'' - s} \right) (s - s') \leq \lambda'_1(\mathcal{Q}_{\varepsilon, s}).$$

Hence

$$\lambda'_1(\mathcal{Q}_{\varepsilon, s'}) + \left(\lim_{\substack{s'' < s \\ s'' \rightarrow s}} \frac{\lambda'_1(\varepsilon, s'') - \lambda'_1(\varepsilon, s)}{s'' - s} \right) (s - s') \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)) + \gamma$$

and consequently

$$\lambda'_1(\mathcal{Q}_{\varepsilon, s'}) + (\lambda'_1(\varepsilon, 2) - \lambda'_1(\varepsilon, 1))(s - s') \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)) + \gamma.$$

Passing to the limit $s' \rightarrow 0$, we get:

$$\lambda'_1(\mathcal{Q}_{\varepsilon, 0}) + (\lambda'_1(\varepsilon, 2) - \lambda'_1(\varepsilon, 1))s \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)) + \gamma$$

and this inequality holds for any $\gamma > 0$, $s \in (0, 1)$ and $\varepsilon \in (0, \varepsilon_{s,\gamma})$.

There exists a sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ such that, as $k \rightarrow +\infty$, $\varepsilon_k \rightarrow 0$ and

$$\lambda'_1(\mathcal{Q}_{\varepsilon_k, 0}) \rightarrow \limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, 0}).$$

Passing to the limit along this sequence, we obtain:

$$\limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, 0}) + \liminf_{k \rightarrow +\infty} (\lambda'_1(\varepsilon_k, 2) - \lambda'_1(\varepsilon_k, 1))s \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)) + \gamma.$$

Finally, passing to the limit $s \rightarrow 0$ and $\gamma \rightarrow 0$, we obtain the second inequality:

$$\limsup_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \lambda'_1(\mathcal{Q}_{\varepsilon, 0}) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x)).$$

This ends the proof of this step. \square

Consequently, in the following steps, we assume without loss of generality that every $\mathbf{L}^\varepsilon(t, x)$ is indeed irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$.

In order to ease the reading, we assume without loss of generality (up to a spatial translation) that the minimum of $x \mapsto \lambda_{1, \text{per}}\left(\frac{d}{dt} - \mathbf{L}(x)\right)$ is attained at $x = 0$.

Step 2: $\liminf_{\varepsilon \rightarrow 0} \lambda'_1(\mathcal{Q}_\varepsilon) \geq \lambda_{1, \text{per}}\left(\frac{d}{dt} - \mathbf{L}(0)\right)$. For each $\varepsilon > 0$, let \mathbf{u}_ε be the space-time periodic generalized principal eigenfunction associated with $\lambda'_1(\mathcal{Q}_\varepsilon)$ and normalized by

$$\max_{i \in [N]} \max_{\overline{\Omega_{\text{per}}}} u_{i, \varepsilon} = 1.$$

Let $(t_\varepsilon, x_\varepsilon) \in \overline{\Omega_{\text{per}}}$ such that $\max_{i \in [N]} u_{i,\varepsilon}(t_\varepsilon, x_\varepsilon) = 1$. By compactness, there exists a sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ that converges to 0 as $k \rightarrow \infty$ and such that:

- (1) $(t_k, x_k) = (t_{\varepsilon_k}, x_{\varepsilon_k})$ converges in $\overline{\Omega_{\text{per}}}$ to a limit (t_∞, x_∞) ;
- (2) $\lambda_k = \lambda_1'(\mathcal{Q}_{\varepsilon_k})$ converges in $[-\lambda_{\text{PF}}(\overline{\mathbf{L}}), -\lambda_{\text{PF}}(\underline{\mathbf{L}})]$ to a limit λ_∞ (see Corollary 3.13).

Let $g_i : \varepsilon \in (0, +\infty) \mapsto f_i(\varepsilon)/\varepsilon$. For each $k \in \mathbb{N}$, define

$$\begin{aligned} \mathbf{v}_k &: (t, x) \mapsto \mathbf{u}_{\varepsilon_k}(t, \varepsilon_k x + x_k), \\ (A_{i,k})_{i \in [N]} &: (t, x) \mapsto (g_i(\varepsilon_k))^2 A_i(t, \varepsilon_k x + x_k)_{i \in [N]}, \\ (q_{i,k})_{i \in [N]} &: (t, x) \mapsto (g_i(\varepsilon_k)) q_i^{\varepsilon_k}(t, \varepsilon_k x + x_k)_{i \in [N]}, \\ \mathbf{L}_k &: (t, x) \mapsto \mathbf{L}^{\varepsilon_k}(t, \varepsilon_k x + x_k), \\ \mathcal{Q}_k &= \partial_t - \text{diag}(\nabla \cdot (A_{i,k} \nabla) - q_{i,k} \cdot \nabla) - \mathbf{L}_k. \end{aligned}$$

The function \mathbf{v}_k is space-time periodic with periodicity cell $[0, T] \times [0, L/\varepsilon_k]$. It satisfies

$$\mathcal{Q}_k \mathbf{v}_k = \lambda_k \mathbf{v}_k \quad \text{in } \mathbb{R} \times \mathbb{R}^n.$$

Denote, for each $i \in [N]$,

$$\sigma_i = f_i'(0) = \lim_{\varepsilon \rightarrow 0} g_i(\varepsilon) \geq 0.$$

Let $I \subset [N]$ such that $\sigma_i > 0$ if and only if $i \in I$. By assumption on \mathbf{f} , I is nonempty.

The coefficients of the above system, $(A_{i,k})$, $(q_{i,k})$ and \mathbf{L}_k , converge locally uniformly to their respective space homogeneous $[0, T]$ -time periodic limits $(\sigma_i^2 A_i(x_\infty))_{i \in [N]}$, $(\sigma_i q_i^0(x_\infty))_{i \in [N]}$, $\mathbf{L}(x_\infty)$.

Note that the limiting operator,

$$\mathcal{Q}_\infty = \partial_t - \text{diag}(\nabla \cdot (\sigma_i^2 A_i(x_\infty) \nabla) - \sigma_i q_i^0(x_\infty) \cdot \nabla) - \mathbf{L}(x_\infty),$$

is a non-degenerate parabolic operator only in the special case $I = [N]$ and is in general a parabolic–ordinary operator. Hence the standard parabolic estimates cannot be used to pass directly to the limit in the solution \mathbf{v}_k .

Since $\mathbf{v}_k \leq \mathbf{1}$ for all $k \in \mathbb{N}$, by virtue of the Banach–Alaoglu theorem, the sequence $(\mathbf{v}_k)_{k \in \mathbb{N}}$ converges up to extraction to a weak- \star limit $\mathbf{v}_\infty \in \mathcal{L}_{\text{per}}^\infty([0, T], \mathcal{L}^\infty(\mathbb{R}^n))$. Up to other extractions, by standard parabolic estimates [52], all components with index $i \in I$ (there is at least one) converge in a Sobolev stronger sense that preserves the fact that they are weak solutions. We introduce, for every $k \leq +\infty$, the adjoint operator:

$$\mathcal{Q}_k^* = -\partial_t - \text{diag}(\nabla \cdot (A_{i,k} \nabla) + q_{i,k} \cdot \nabla + \nabla \cdot q_{i,k}) - \mathbf{L}_k^T.$$

By passing to the limit in

$$\lambda_k \int_{[0, T] \times \mathbb{R}^n} \varphi^T \mathbf{v}_k = \int_{[0, T] \times \mathbb{R}^n} \mathbf{v}_k^T (\mathcal{Q}_k^* \varphi),$$

for any test function $\varphi \in \mathcal{C}_{\text{per}}^1(\mathbb{R}, \mathcal{C}_c^2(\mathbb{R}^n, [\mathbf{0}, \infty)))$, and then integrating by parts in space-time, we find that:

- if $i \in I$, then $v_{i,\infty}$ is a bounded weak solution of a parabolic equation with bounded continuous coefficients and bounded forcing term, whence it benefits from bootstrapped parabolic regularity, and in particular it is continuous;

- if $i \in [N] \setminus I$, then $v_{i,\infty}$ is a bounded weak solution of an ordinary differential equation with bounded continuous coefficients and bounded forcing term, whence it is actually Lipschitz-continuous.

Consequently, \mathbf{v}_∞ is continuous. Iterating, we find that it is actually a classical solution, nonnegative, time periodic and globally bounded by $\mathbf{1}$, of

$$\mathcal{Q}_\infty \mathbf{v}_\infty = \lambda_\infty \mathbf{v}_\infty \quad \text{in } \mathbb{R} \times \mathbb{R}^n.$$

Let us show now that it is a nonzero solution. Actually, let us show that it is positive. Up to permutations, we can assume that there exists $N_0 \in [N]$ such that $I = [N_0]$. Since

$$\mathbf{L}_\infty \geq \begin{pmatrix} (l_{i,j,\infty})_{i,j \in [N_0]} & 0 \\ 0 & (l_{i,j,\infty})_{i,j \in [N] \setminus [N_0]} \end{pmatrix}$$

and both diagonal blocks in the right-hand side are fully coupled (*cf.* Step 1), we can apply the comparison principle for cooperative systems of parabolic partial differential equations of Proposition 2.1 on $(v_{i,\infty})_{i \in [N_0]}$ and a form of comparison principle for cooperative systems of ordinary differential equations [65] on $(v_{i,\infty})_{i \in [N] \setminus [N_0]}$ to deduce that they are both either positive or zero. Subsequently, the full coupling of \mathbf{L}_∞ (*cf.* Step 1) implies that \mathbf{v}_∞ is either positive or zero. The standard parabolic estimates [52] on $(v_{i,k})_{i \in [N_0]}$ imply that if

$$\inf_{k \in \mathbb{N}} \max_{i \in [N_0]} v_{i,k}(t_\infty, 0) > 0,$$

then $\max_{i \in [N_0]} v_{i,\infty}(t_\infty, 0) > 0$ and subsequently \mathbf{v}_∞ is positive. If, on the contrary,

$$\inf_{k \in \mathbb{N}} \max_{i \in [N_0]} v_{i,k}(t_\infty, 0) = 0,$$

then by the parabolic strong comparison principle applied to each uniformly parabolic operator \mathcal{Q}_k , it actually means that, up to extraction, $(v_{i,k})_{i \in [N_0]}$ converges locally uniformly to $\mathbf{0}$ as $k \rightarrow +\infty$. Due to the normalization $\max_{j \in [N]} v_{j,k}(t_k, 0) = 1$, there exists $\bar{j} \in [N] \setminus [N_0]$ such that, up to extraction,

$$v_{\bar{j},k}(t_\infty, 0) \rightarrow 1 \quad \text{as } k \rightarrow +\infty.$$

Moreover, due to the full coupling of $\mathbf{L}(x_\infty)$, there exists $\bar{i} \in [N_0]$ such that

$$l_{\bar{i},\bar{j}}(t_\infty, x_\infty) > 0.$$

Evaluating at $(t, x) = (t_\infty, 0)$ and passing to the limit $k \rightarrow +\infty$ in the equality

$$\partial_t v_{\bar{i},k} - \nabla \cdot (A_{\bar{i},k} \nabla v_{\bar{i},k}) + q_{\bar{i},k} \cdot \nabla v_{\bar{i},k} - \sum_{j \in [N_0]} l_{\bar{i},j,k} v_{j,k} - \lambda_k v_{\bar{i},k} = \sum_{j \in [N] \setminus [N_0]} l_{\bar{i},j,k} v_{j,k},$$

the left-hand side converges to 0 (by standard parabolic regularity estimates) whereas the right-hand side admits a limit inferior bounded below by $l_{\bar{i},\bar{j}}(t_\infty, x_\infty) > 0$. This contradiction means that

$$\inf_{k \in \mathbb{N}} \max_{i \in [N_0]} v_{i,k}(t_\infty, 0) > 0,$$

whence \mathbf{v}_∞ is positive.

Now, consider $\mathbf{v} \in \mathcal{C}_{\text{per}}^1(\mathbb{R}, \mathbb{R}^N)$ the time periodic principal eigenfunction satisfying

$$\mathbf{v}' = \mathbf{L}(x_\infty) \mathbf{v} + \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x_\infty) \right) \mathbf{v}, \quad \min_{i \in [N]} \min_{[0,T]} v_i = 1.$$

Since it is spatially homogeneous, it is also a solution of

$$\mathcal{Q}_\infty \mathbf{v} = \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x_\infty) \right) \mathbf{v}.$$

A comparison argument (distinguishing, as before, the parabolic part and the ordinary part of the operator \mathcal{Q}_∞) shows that if $\lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x_\infty) \right) \geq \lambda_\infty$, then \mathbf{v} and \mathbf{v}_∞ are proportional, and then the preceding inequality is an equality. In other words,

$$\lambda_\infty \geq \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x_\infty) \right).$$

Since this applies to any accumulation point of $(x_\varepsilon, \lambda'_1(\mathcal{Q}_\varepsilon))_{\varepsilon>0}$, the conclusion of this step follows:

$$\liminf_{\varepsilon \rightarrow 0} \lambda'_1(\mathcal{Q}_\varepsilon) \geq \min_{x \in [0, L]} \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right) = \lambda_{1,\text{per}} \left(\frac{d}{dt} - \mathbf{L}(0) \right).$$

□

Step 3: $\limsup_{\varepsilon \rightarrow 0} \lambda'_1(\mathcal{Q}_\varepsilon) \leq \min_{x \in [0, L]} \lambda_1(\tilde{\mathcal{Q}}(x))$. Let us fix arbitrarily $x_0 \in [0, L]$ and prove that

$$\limsup_{\varepsilon \rightarrow 0} \lambda'_1(\mathcal{Q}_\varepsilon) \leq \lambda_1(\tilde{\mathcal{Q}}(x_0)).$$

We repeat the construction of Step 2 with the (important) modification $(t_\varepsilon, x_\varepsilon) = (0, x_0)$ for each $\varepsilon > 0$. We use, for each $k \in \mathbb{N}$ and up to $k = +\infty$ when it makes sense, the same notations $\lambda_k, t_k, x_k, \mathbf{v}_k, \sigma_i, I, A_{i,k}, q_{i,k}, \mathbf{L}_k, \mathcal{Q}_k$. With the new spatial shift, $\mathcal{Q}_\infty = \tilde{\mathcal{Q}}(x_0)$.

Since $(0, x_0) \in \Omega_{\text{per}}$ is *a priori* not in $\text{argmax}(\max_{i \in [N]} v_{i,k})$, it is now unclear if \mathbf{v}_k remains positive at the limit $k \rightarrow +\infty$. Thus, instead of directly passing to the limit, we are going to construct a sub-solution appropriate for every finite k .

Let $R > 0$ and, for each $k \in \mathbb{N}$, define $\mu_{R,k} = \lambda_{1,\text{Dir}}(\mathcal{Q}_k, B(x_0, R))$. The solution $\mathbf{v}_k \in \mathcal{C}_{t-\text{per}}^{1,2}(\mathbb{R} \times B(x_0, R), \mathbb{R}^N) \cap \mathcal{C}_{t-\text{per}}^{0,1}(\mathbb{R} \times \overline{B(x_0, R)}, \mathbb{R}^N)$ of the Dirichlet periodic-parabolic problem:

$$\begin{cases} \mathcal{Q}_k \mathbf{v}_k = \mu_{R,k} \mathbf{v}_k & \text{in } \mathbb{R} \times B(x_0, R) \\ \mathbf{v}_k = \mathbf{0} & \text{on } \mathbb{R} \times \partial B(x_0, R) \\ \max_{i \in [N]} \max_{(t,x) \in [0, T] \times B(x_0, R)} v_{i,k}(t, x) = 1 \end{cases}$$

can be multiplied by an appropriately small positive constant and extended in $\mathbb{R} \times \mathbb{R}^n$ by setting $\mathbf{v}_k = \mathbf{0}$ in $\mathbb{R} \times (\mathbb{R}^n \setminus B(x_0, R))$ in order to show, thanks to a standard comparison argument, that

$$(19) \quad \lambda'_1(\mathcal{Q}_k) \leq \mu_{R,k}.$$

Now, let us verify that $\mu_{R,k} \rightarrow \lambda_{1,\text{Dir}}(\tilde{\mathcal{Q}}(x_0), B(x_0, R))$ as $k \rightarrow +\infty$. We point out that when $\min_{i \in [N]} \sigma_i > 0$, this convergence is obvious, by locally uniform convergence of the coefficients $(A_{i,k})_{i \in [N]}$, $(q_{i,k})_{i \in [N]}$ and \mathbf{L}_k and by continuity of the Dirichlet principal eigenvalue. Hence we only have to verify that this continuity remains true when some, but not all, equations of the system become ordinary differential equations.

Repeating the compactness and bootstrap argument of Step 2, we obtain after passing to the limit a classical solution \mathbf{v}_∞ of the Dirichlet periodic-parabolic problem:

$$\begin{cases} \tilde{\mathcal{Q}}(x_0)\mathbf{v}_\infty = \mu_{R,\infty}\mathbf{v}_\infty & \text{in } \mathbb{R} \times B(x_0, R) \\ \mathbf{v}_\infty = \mathbf{0} & \text{on } \mathbb{R} \times \partial B(x_0, R) \\ \max_{i \in [N]} \max_{(t,x) \in [0,T] \times \overline{B(x_0,R)}} v_{i,\infty}(t,x) = 1, \end{cases}$$

where $\mu_{R,\infty}$ is an accumulation point of $(\mu_{R,k})_{k \in \mathbb{N}}$. To prove by a comparison argument the uniqueness of the Dirichlet periodic-parabolic principal eigenpair, so that $\mu_{R,\infty} = \lambda_{1,\text{Dir}}(\tilde{\mathcal{Q}}(x_0), B(x_0, R))$, we need to control the behavior of the solution close to $\partial B(x_0, R)$. More precisely, it suffices to prove that for every Dirichlet principal eigenfunction \mathbf{v} , there exists $C > 0$ such that, for each $i \in [N]$,

$$(20) \quad 0 < \frac{1}{C} \leq \frac{v_i(t,x)}{||x - x_0| - R|} \leq C \quad \text{for all } (t,x) \in [0,T] \times B(x_0, R).$$

For each $i \in I$, (20) follows directly from the parabolic regularity up to the boundary and the Hopf lemma. It remains to prove (20) for each $i \in [N] \setminus I$.

As in Step 2, assume up to permutations that $I = [N_0]$ with $N_0 \in [N]$. If $N_0 = N$, then (20) is true for all $i \in [N]$. If $N_0 < N$, then $1 \leq N_0 \leq N - 1$ and we can write $\mathbf{L}(x_0)$ in block form:

$$\mathbf{L}(t, x_0) = \begin{pmatrix} \mathbf{A}(t) & \mathbf{B}(t) \\ \mathbf{C}(t) & \mathbf{D}(t) \end{pmatrix} \quad \text{for all } t \in [0, T]$$

with

$$\mathbf{A}(t) \in \mathbb{R}^{N_0 \times N_0}, \mathbf{B}(t) \in \mathbb{R}^{N_0 \times (N - N_0)}, \mathbf{C}(t) \in \mathbb{R}^{(N - N_0) \times N_0}, \mathbf{D}(t) \in \mathbb{R}^{(N - N_0) \times (N - N_0)}.$$

By Step 1, we can assume without loss of generality that, for all $t \in [0, T]$, $\mathbf{A}(t)$ is essentially positive, $\mathbf{B}(t)$ is positive, $\mathbf{C}(t)$ is positive and $\mathbf{D}(t)$ is essentially positive. Hence, decomposing the principal eigenfunction \mathbf{v} as

$$\mathbf{v} = \begin{pmatrix} \mathbf{w} \\ \tilde{\mathbf{w}} \end{pmatrix} \quad \text{with } \mathbf{w} \in \mathbb{R}^{N_0}, \tilde{\mathbf{w}} \in \mathbb{R}^{N - N_0},$$

we find

$$\partial_t \tilde{\mathbf{w}} = \mathbf{D} \tilde{\mathbf{w}} + \mathbf{C} \mathbf{w} \quad \text{in } [0, T] \times B(x_0, R)$$

On one hand, this immediately leads to the Lipschitz-continuity of $\tilde{\mathbf{w}}$ at the boundary of the spatial domain, namely the upper estimate of (20) for each $i \in [N] \setminus [N_0]$. On the other hand, using the inequality

$$\partial_t \tilde{\mathbf{w}} = \mathbf{D} \tilde{\mathbf{w}} + \mathbf{C} \mathbf{w} \geq \text{diag}(d_{i,i})_{i \in [N - N_0]} \tilde{\mathbf{w}} + \mathbf{C} \mathbf{w} \quad \text{in } [0, T] \times B(x_0, R)$$

multiplied by

$$\mathbf{E} : t \mapsto \text{diag} \left(e^{-\int_0^t d_{i,i}(\tau) d\tau} \right)_{i \in [N - N_0]}$$

and integrated in $[0, t]$ for an arbitrary $t \in (0, T]$, we obtain for all $x \in B(x_0, R)$

$$\begin{aligned} \tilde{\mathbf{w}}(t, x) &\geq \mathbf{E}(t)^{-1} \tilde{\mathbf{w}}(0, x) + \mathbf{E}(t)^{-1} \int_0^t \mathbf{E}(t') \mathbf{C}(t') \mathbf{w}(t', x) dt' \\ &\geq \mathbf{E}(t)^{-1} \int_0^t \mathbf{E}(t') \mathbf{C}(t') \mathbf{w}(t', x) dt' \end{aligned}$$

Since \mathbf{w} satisfies (20) component-wise, it follows that, up to increasing C , for each $i \in [N] \setminus [N_0]$,

$$0 < \frac{1}{C} \leq \frac{\tilde{w}_i(t, x)}{\|x - x_0\| - R} \quad \text{for all } (t, x) \in [0, T] \times B(x_0, R).$$

Therefore for each $i \in [N] \setminus [N_0]$ the lower estimate of (20) is also satisfied.

Hence (20) is satisfied for each $i \in [N]$.

Now, take two principal eigenpairs (λ, \mathbf{v}) and $(\tilde{\lambda}, \tilde{\mathbf{v}})$, assume for instance $\lambda \geq \tilde{\lambda}$, and let us prove by comparison that $\lambda = \tilde{\lambda}$ and $\mathbf{v} = \kappa \tilde{\mathbf{v}}$ for some $\kappa > 0$. By (20), there exists $\kappa > 0$ such that $\kappa \tilde{\mathbf{v}} \leq \mathbf{v}$. Hence $\mathbf{w} = \mathbf{v} - \kappa \tilde{\mathbf{v}}$ is nonnegative and satisfies:

$$\tilde{\mathcal{Q}}(x_0) \mathbf{w} = \tilde{\lambda} \mathbf{w} + (\lambda - \tilde{\lambda}) \mathbf{v} \geq \tilde{\lambda} \mathbf{w} \quad \text{in } \mathbb{R} \times B(x_0, R).$$

Verifying that the Hopf-type lower estimate of (20) remains true for super-solutions (the argument is the same), we find that \mathbf{w} is either positive or zero. Increasing continuously κ , we deduce from the comparison principle an optimal κ^* such that $\mathbf{v} = \kappa^* \tilde{\mathbf{v}}$. It follows subsequently, from the system satisfied by \mathbf{w} , that $\lambda = \tilde{\lambda}$.

Hence the Dirichlet principal eigenvalue is unique indeed and $\mu_{R,k} \rightarrow \lambda_{1, \text{Dir}}(\tilde{\mathcal{Q}}(x_0))$. Passing to the limit $k \rightarrow +\infty$ in (19), we deduce:

$$\limsup_{\varepsilon \rightarrow 0} \lambda'_1(\mathcal{Q}_\varepsilon) \leq \lambda_{1, \text{Dir}}(\tilde{\mathcal{Q}}(x_0), B(x_0, R)).$$

Since this is true for all $R > 0$, it only remains to verify that

$$\lambda_{1, \text{Dir}}(\tilde{\mathcal{Q}}(x_0), B(x_0, R)) \rightarrow \lambda_1(\tilde{\mathcal{Q}}(x_0)) \quad \text{as } R \rightarrow +\infty.$$

Again, this is classical if $N_0 = N$ and only requires work if $N_0 < N$.

In fact, it can be verified that, in order to adapt the proof of Proposition 3.2 to an operator which is degenerate parabolic but still satisfies the strong comparison principle, we only need to be able to extract a convergent subsequence of Dirichlet principal eigenfunctions \mathbf{v}_R as $R \rightarrow +\infty$ with nonzero limit. In view of the compactness procedure detailed in Step 2, we actually only need to verify that the pointwise normalization $\max_{i \in [N]} v_{i,R}(0, x_0) = 1$ implies that, for any $R_0 > 0$, there exists $C > 0$ such that, for any $R > R_0$,

$$\mathbf{v}_R \leq C \mathbf{1} \quad \text{in } [0, T] \times B(x_0, R_0).$$

As a matter of fact, such local bounds can be derived from parabolic interior regularity estimates [52] applied to components with index $i \in [N_0]$ combined with standard regularity results for parameterized ordinary differential equations applied to components with index $i \in [N] \setminus [N_0]$. This is standard and not detailed. \square

The proof is now ended. \square

Remark 3.10. Contrarily to what was claimed by Nadin in his work on the scalar case [59, Theorem 3.6], the convergence to $\min_{x \in [0, L]} \lambda_{1, \text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right)$ cannot be true in general, even if $N = 1$. Indeed, if:

- $f_i(\varepsilon) = \varepsilon$ for each $i \in [N]$;
- the coefficients of \mathcal{Q}_ε do not depend on time;
- \mathbf{L} is symmetric;
- and there exists $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ such that $\int_{[0, L]} \nabla Q = 0$ and

$$(\varepsilon^2 A_i)^{-1} (\varepsilon q_i^\varepsilon) = \nabla Q \quad \text{for each } i \in [N];$$

then by the variational formula of Theorem 1.13,

$$\lambda_1(\mathcal{Q}_\varepsilon) = \lambda'_1(\mathcal{Q}_\varepsilon) = \lambda'_1(-\varepsilon^2 \operatorname{diag}(\nabla \cdot (A_i \nabla)) - \mathbf{L}_Q)$$

with

$$\begin{aligned} \mathbf{L}_Q &= \mathbf{L} + \operatorname{diag} \left(\frac{1}{2} \nabla \cdot (\varepsilon^2 A_i \nabla Q) - \frac{1}{4} \nabla Q \cdot \varepsilon^2 A_i \nabla Q \right) \\ &= \mathbf{L} + \operatorname{diag} \left(\frac{1}{2} \varepsilon \nabla \cdot (q_i) - \frac{1}{4} A_i^{-1} q_i \cdot q_i \right). \end{aligned}$$

If the convergence of $\lambda'_1(\mathcal{Q}_\varepsilon)$ to $\min_{x \in [0, L]} \lambda_{1, \text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right)$ was true, then by passing to the limit in the equality $\lambda'_1(\mathcal{Q}_\varepsilon) = \lambda'_1(-\varepsilon^2 \operatorname{diag}(\nabla \cdot (A_i \nabla)) - \mathbf{L}_Q)$, we would obtain

$$\min_{x \in [0, L]} \lambda_{1, \text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) \right) = \min_{x \in [0, L]} \lambda_{1, \text{per}} \left(\frac{d}{dt} - \mathbf{L}(x) + \frac{1}{4} \operatorname{diag}(|A_i^{-1/2} q_i|^2)(x) \right),$$

which yields an obvious contradiction after a careful choice of \mathbf{L} , $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ – for instance, in spatial dimension $n = 1$,

$$A_i = 1, \quad q_i : x_1 \mapsto \cos \left(\frac{2\pi}{L_1} x_1 \right), \quad \mathbf{L} : x_1 \mapsto -\frac{1}{4} \sin \left(\frac{2\pi}{L_1} x_1 \right)^2 \mathbf{I} + \mathbf{M}$$

with \mathbf{M} the discrete Laplacian defined in (1).

The mistake in Nadin's proof [59] can be corrected with the additional assumption $\lambda_1(\tilde{\mathcal{Q}}) = \lambda'_1(\tilde{\mathcal{Q}})$. In this sense, our result and its proof provide as a by-product a correction of the scalar counterpart in [59].

3.2.3. *Large diffusion: proof of Theorem 1.7.* We now prove Theorem 1.7.

Proposition 3.16. *Let*

$$((A_i), (q_i))_{i \in [N]}, (\mathbf{L}) : t \mapsto \frac{1}{|[0, L]|} \int_{[0, L]} ((A_i, q_i)_{i \in [N]}, \mathbf{L})(t, x) dx$$

and, for all $\mathbf{d} \in (\mathbf{0}, \infty)$, let $\mathcal{Q}_{\mathbf{d}}$ be the operator \mathcal{Q} with $(A_i)_{i \in [N]}$ replaced by $(d_i A_i)_{i \in [N]}$.

Then

$$\lim_{\min_{i \in [N]} d_i \rightarrow +\infty} \lambda_{1, \text{per}}(\mathcal{Q}_{\mathbf{d}}) = \lambda_{1, \text{per}}(\partial_t - \langle \mathbf{L} \rangle).$$

Proof. Let $\mathbf{d} \gg \mathbf{0}$ and $\mathbf{u}_{\mathbf{d}}$ be the periodic principal eigenfunction associated with $\lambda'_1(\mathcal{Q}_{\mathbf{d}})$ and normalized by

$$\frac{1}{T|[0, L]|} \int_{\Omega_{\text{per}}} |\mathbf{u}_{\mathbf{d}}|^2 = 1.$$

Below, we assume \mathcal{C}^1 space regularity of the coefficients $(q_i)_{i \in [N]}$. The proof in the general case with mere Hölder-continuity is not detailed – it can be deduced by a standard regularization procedure and the continuity of periodic principal eigenvalues.

Multiplying $(\mathcal{Q}_{\mathbf{d}} \mathbf{u}_{\mathbf{d}})_i - \lambda'_1(\mathcal{Q}_{\mathbf{d}}) u_{\mathbf{d}, i}$ by $u_{\mathbf{d}, i}$ and then integrating over $\overline{\Omega_{\text{per}}}$, we find for each $i \in [N]$:

$$d_i \int_{\overline{\Omega_{\text{per}}}} \nabla u_{\mathbf{d}, i} \cdot A_i \nabla u_{\mathbf{d}, i} = \int_{\overline{\Omega_{\text{per}}}} \left(\left(\frac{\nabla \cdot q_i}{2} + \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \right) u_{\mathbf{d}, i}^2 + \sum_{j=1}^N l_{i, j} u_{\mathbf{d}, i} u_{\mathbf{d}, j} \right).$$

Recall from Corollary 3.13 the estimate $-\lambda_{\text{PF}}(\bar{\mathbf{L}}) \leq \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \leq -\lambda_{\text{PF}}(\mathbf{L})$, whence, by the Young inequality $|u_{\mathbf{d},i}u_{\mathbf{d},j}| \leq \frac{1}{2}(|u_{\mathbf{d},i}|^2 + |u_{\mathbf{d},j}|^2)$, there exists a constant $K > 0$ independent of \mathbf{d} such that

$$0 \leq \sum_{i=1}^N \int_{\Omega_{\text{per}}} \nabla u_{\mathbf{d},i} \cdot A_i \nabla u_{\mathbf{d},i} \leq \frac{K}{\min_{i \in [N]} d_i}.$$

Consequently,

$$\frac{1}{T} \int_0^T \left(\sum_{i=1}^N \int_{[0,L]} \nabla u_{\mathbf{d},i} \cdot A_i \nabla u_{\mathbf{d},i} \right) \rightarrow 0 \quad \text{as } \min_{i \in [N]} d_i \rightarrow +\infty.$$

Let $\langle \mathbf{u}_{\mathbf{d}} \rangle : t \mapsto \frac{1}{|[0,L]|} \int_{[0,L]} \mathbf{u}_{\mathbf{d}}(t,x) dx$ and $\mathbf{v}_{\mathbf{d}} = \mathbf{u}_{\mathbf{d}} - \langle \mathbf{u}_{\mathbf{d}} \rangle$. By the Poincaré inequality, there exists another constant $K' > 0$ such that

$$\sum_{i=1}^N \int_{[0,L]} \nabla u_{\mathbf{d},i} \cdot A_i \nabla u_{\mathbf{d},i} = \sum_{i=1}^N \int_{[0,L]} \nabla v_{\mathbf{d},i} \cdot A_i \nabla v_{\mathbf{d},i} \geq K' \int_{[0,L]} |\mathbf{v}_{\mathbf{d}}|^2.$$

Since the average in $[0, T]$ of the nonnegative function on the left-hand side converges to 0 as $\min_{i \in [N]} d_i \rightarrow +\infty$, so does the average in $[0, T]$ of $\int_{[0,L]} |\mathbf{v}_{\mathbf{d}}|^2$, whence $\mathbf{v}_{\mathbf{d}}$ itself converges to $\mathbf{0}$ almost everywhere (up to extraction of a subsequence).

Since, for each $i \in [N]$,

$$\begin{aligned} \int_0^T \langle \mathbf{u}_{\mathbf{d}} \rangle_i &= |[0, L]|^{-1} \int_{\Omega_{\text{per}}} u_{\mathbf{d},i} \\ &\leq \left(\frac{T}{|[0, L]|} \right)^{1/2} \left(\int_{\Omega_{\text{per}}} u_{\mathbf{d},i}^2 \right)^{1/2} \\ &\leq \left(\frac{T}{|[0, L]|} \right)^{1/2} \left(\int_{\Omega_{\text{per}}} \sum_{i=1}^N u_{\mathbf{d},i}^2 \right)^{1/2} = T, \end{aligned}$$

$\langle \mathbf{u}_{\mathbf{d}} \rangle_i$ is bounded in $\mathcal{L}^1([0, T])$ uniformly with respect to \mathbf{d} .

Integrating $\mathcal{Q}_{\mathbf{d}} \mathbf{u}_{\mathbf{d}} = \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \mathbf{u}_{\mathbf{d}}$ over $[0, L]$ and dividing by $|[0, L]|$, we find:

$$\begin{aligned} \partial_t \langle \mathbf{u}_{\mathbf{d}} \rangle &= |[0, L]|^{-1} \int_{[0,L]} (\text{diag}(\nabla \cdot q_i) \mathbf{u}_{\mathbf{d}}) + |[0, L]|^{-1} \int_{[0,L]} (\mathbf{L} \mathbf{u}_{\mathbf{d}}) + \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \langle \mathbf{u}_{\mathbf{d}} \rangle \\ &= |[0, L]|^{-1} \int_{[0,L]} \text{diag}(\nabla \cdot q_i) \langle \mathbf{u}_{\mathbf{d}} \rangle + |[0, L]|^{-1} \int_{[0,L]} \mathbf{L} \langle \mathbf{u}_{\mathbf{d}} \rangle + \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \langle \mathbf{u}_{\mathbf{d}} \rangle \\ &\quad + |[0, L]|^{-1} \int_{[0,L]} ((\text{diag}(\nabla \cdot q_i) + \mathbf{L}) \mathbf{v}_{\mathbf{d}}) \\ &= \langle \mathbf{L} \rangle \langle \mathbf{u}_{\mathbf{d}} \rangle + \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \langle \mathbf{u}_{\mathbf{d}} \rangle + |[0, L]|^{-1} \int_{[0,L]} ((\text{diag}(\nabla \cdot q_i) + \mathbf{L}) \mathbf{v}_{\mathbf{d}}). \end{aligned}$$

Now, by the discrete and continuous Cauchy–Schwarz inequalities, denoting $\|\cdot\|$ the norm in $\mathcal{L}^\infty(\overline{\Omega_{\text{per}}}, \mathbb{R})$,

$$\begin{aligned} \left| \int_{[0,L]} \left((\nabla \cdot q_i) v_{\mathbf{d},i} + \sum_{j=1}^N l_{i,j} v_{\mathbf{d},j} \right) \right| &\leq \left(\|\nabla \cdot q_i\| + \max_{j \in [N]} \|l_{i,j}\| \right) \int_{[0,L]} \sum_{j=1}^N |v_{\mathbf{d},j}| \\ &\leq \left(\|\nabla \cdot q_i\| + \max_{j \in [N]} \|l_{i,j}\| \right) \int_{[0,L]} \sqrt{N} \left(\sum_{j=1}^N |v_{\mathbf{d},j}|^2 \right)^{\frac{1}{2}} \\ &\leq \left(\|\nabla \cdot q_i\| + \max_{j \in [N]} \|l_{i,j}\| \right) \sqrt{N|[0,L]|} \int_{[0,L]} |\mathbf{v}_{\mathbf{d}}|^2. \end{aligned}$$

Since $\int_{[0,L]} |\mathbf{v}_{\mathbf{d}}|^2$ term converges to 0 as $\min_{i \in [N]} d_i \rightarrow +\infty$, each component of $\partial_t \langle \mathbf{u}_{\mathbf{d}} \rangle$ is bounded in $\mathcal{L}^1([0, T])$ uniformly with respect to \mathbf{d} . Hence each component of $\langle \mathbf{u}_{\mathbf{d}} \rangle$ is bounded uniformly in $\mathcal{W}^{1,1}([0, T])$, and then via the fundamental theorem of calculus it is bounded uniformly in $\mathcal{L}^\infty([0, T])$, whence it is bounded uniformly in the space of functions of bounded variation $\mathcal{BV}([0, T])$. By compactness of the embedding $\mathcal{W}^{1,1} \hookrightarrow \mathcal{L}^1$, each component of $\langle \mathbf{u}_{\mathbf{d}} \rangle$ converges up to extraction in $\mathcal{L}^1([0, T])$. Using the equation

$$\partial_t \langle \mathbf{u}_{\mathbf{d}} \rangle = \langle \mathbf{L} \rangle \langle \mathbf{u}_{\mathbf{d}} \rangle + \lambda'_1(\mathcal{Q}_{\mathbf{d}}) \langle \mathbf{u}_{\mathbf{d}} \rangle + |[0, L]|^{-1} \int_{[0,L]} ((\text{diag}(\nabla \cdot q_i) + \mathbf{L}) \mathbf{v}_{\mathbf{d}})$$

and assuming up to another extraction that $\lambda'_1(\mathcal{Q}_{\mathbf{d}}) \rightarrow \lambda \in \mathbb{R}$, so does each component of $\partial_t \langle \mathbf{u}_{\mathbf{d}} \rangle$. Denoting by \mathbf{u}_∞ the limit of $\langle \mathbf{u}_{\mathbf{d}} \rangle$, we deduce that the limit of $\partial_t \langle \mathbf{u}_{\mathbf{d}} \rangle$ is, in distributional sense, the derivative of \mathbf{u}_∞ , so that \mathbf{u}_∞ satisfies

$$\partial_t \mathbf{u}_\infty = \langle \mathbf{L} \rangle \mathbf{u}_\infty + \lambda \mathbf{u}_\infty \quad \text{in } (\mathcal{L}^\infty)'(\mathbb{R}).$$

Again by virtue of the fundamental theorem of calculus, each component of \mathbf{u}_∞ is actually in $\mathcal{L}^\infty([0, T])$, and now from the equation it appears that so does each component of $\partial_t \mathbf{u}_\infty$. Therefore \mathbf{u}_∞ is in fact Lipschitz-continuous, and using again the equation it is \mathcal{C}^1 . Since it is periodic, nonnegative (by almost everywhere convergence, up to another extraction) and nonzero (if on the contrary it was zero, then $\mathbf{u}_{\mathbf{d}}$ would converge to 0 almost everywhere and this would contradict the normalization on $\mathbf{u}_{\mathbf{d}}$) and since the operator $\partial_t - \langle \mathbf{L} \rangle$ is fully coupled in $[0, T]$ by (A_3) , we deduce by uniqueness of the classical solution that

$$\lambda = \lambda_{1,\text{per}}(\partial_t - \langle \mathbf{L} \rangle).$$

By uniqueness, the family $(\lambda'_1(\mathcal{Q}_{\mathbf{d}}))_{\mathbf{d} \gg \mathbf{0}}$ has a unique accumulation point and thus:

$$\lim_{\min_{i \in [N]} d_i \rightarrow +\infty} \lambda'_1(\mathcal{Q}_{\mathbf{d}}) = \lambda_{1,\text{per}}(\partial_t - \langle \mathbf{L} \rangle).$$

□

Remark 3.11. Contrarily to what was claimed by Nadin in [59, Theorem 3.6], the large diffusion limit of the family $(\lambda_{1,z})_{z \in \mathbb{R}^n}$ and of the generalized principal eigenvalue λ_1 cannot be directly deduced from the above proof. Indeed, the large parameter \mathbf{d} appears in the zeroth order term of the operator \mathcal{Q}_z and makes the eigenvalue $\lambda_{1,z}$ blow-up to $-\infty$ as $\min d_i \rightarrow +\infty$, see Corollary 3.13.

Remark 3.12. Contrarily to the scalar setting [59] or the special cases where $\partial_t - \mathbf{L}(x)$ admits a space-time homogeneous periodic principal eigenfunction, it is in general false that for systems without diffusion and advection, spatial average and periodic principal eigenvalue commute, namely

$$\lambda_{1,\text{per}}(\partial_t - \langle \mathbf{L} \rangle) \neq \langle x \mapsto \lambda_{1,\text{per}}(\partial_t - \mathbf{L}(x)) \rangle.$$

In fact, even the inequality

$$\lambda_{1,\text{per}}(\partial_t - \langle \mathbf{L} \rangle) \geq \min_{x \in [0, L]} \lambda_{1,\text{per}}(\partial_t - \mathbf{L}(x))$$

is false in general, as shown by the following very simple time homogeneous one-dimensional counter-example

$$\mathbf{L} : (t, x) \mapsto \begin{cases} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{if } x \in [0, L_1/2] + L_1\mathbb{Z}, \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{if } x \in [L_1/2, L_1] + L_1\mathbb{Z}. \end{cases}$$

In a time homogeneous setting,

$$\lambda_{1,\text{per}}(\partial_t - \langle \mathbf{L} \rangle) = -\lambda_{\text{PF}}(\langle \mathbf{L} \rangle), \quad \min_{x \in [0, L]} \lambda_{1,\text{per}}(\partial_t - \mathbf{L}(x)) = -\max_{x \in [0, L]} \lambda_{\text{PF}}(\mathbf{L}(x)).$$

With the counter-example above, these two quantities turn out to be respectively $-\frac{3}{2}$ and -1 : the averaged matrix has a larger Perron–Frobenius eigenvalue than the matrix at any point in space. In other words, considering for instance the operator $\partial_t - d\Delta - \mathbf{L}$, the limit $d \rightarrow 0$ of the periodic principal eigenvalue is larger than the limit $d \rightarrow +\infty$. This is in sharp contrast with the variational formula of Theorem 1.13, which indicates a nondecreasing dependence on d but does not apply here due to the asymmetry of \mathbf{L} . Of course \mathbf{L} in this counter-example is not continuous and therefore does not satisfy (A_4) ; however, any smooth sufficiently precise approximation of \mathbf{L} will give the same conclusion, by continuity of the Perron–Frobenius eigenvalue. As a side result, this counter-example also shows that the variational formula of Theorem 1.13 does not hold if only $\langle \mathbf{L} \rangle$ is symmetric, namely if the pointwise symmetry assumption is replaced by an assumption of symmetry on average.

Let us also point out that this counter-example is used to prove Corollary 1.8. More precisely, the map $L_1 \mapsto \lambda_{1,\text{per}}(\mathcal{Q}_{T, L_1})$ is

- constant if for instance the coefficients of \mathcal{Q}_{T, L_1} do not depend on space;
- decreasing if for instance the coefficients of \mathcal{Q}_{T, L_1} do not depend on time and the operator is self-adjoint and spatially heterogeneous, so that the variational formula of Theorem 1.13 applies and periodic principal eigenfunctions are spatially heterogeneous and therefore have nonzero gradients;
- neither if for instance $\mathcal{Q}_{T, L_1} = \frac{1}{T}\partial_t - \frac{1}{L_1^2}\Delta - \mathbf{L}$ with \mathbf{L} the above counter-example, so that the limit $L_1 \rightarrow +\infty$ of the periodic principal eigenvalue is larger than the limit $L_1 \rightarrow 0$.

3.2.4. Small and large time frequency: proof of Theorem 1.9. Now we turn to the proof of Theorem 1.9. Denoting by \mathcal{Q}_ω the operator \mathcal{Q} with ∂_t replaced by $\omega\partial_t$, we first prove the small frequency limit $\omega \rightarrow 0$ in Proposition 3.17, then the high frequency one $\omega \rightarrow +\infty$ in Proposition 3.18.

Proposition 3.17. *For all $z \in \mathbb{R}^n$,*

$$\lim_{\substack{\omega > 0 \\ \omega \rightarrow 0}} \lambda_{1,z}(\mathcal{Q}_\omega) = \frac{1}{T} \int_0^T \lambda_{1,z}(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)) dt,$$

$$\lim_{\substack{\omega > 0 \\ \omega \rightarrow 0}} \lambda_1(\mathcal{Q}_\omega) = \frac{1}{T} \int_0^T \lambda_1(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)) dt,$$

where, with a slight abuse of notation, for all $t \in [0, T]$,

$$((A_i(t), q_i(t))_{i \in [N]}, \mathbf{L}(t)) : x \mapsto ((A_i(t, x), q_i(t, x))_{i \in [N]}, \mathbf{L}(t, x)).$$

Proof. It is sufficient to prove only the case $z = 0$, since we can deduce the general case for $\lambda_{1,z}$ by applying the result to the operator \mathcal{Q}_z , and then we can deduce the result for λ_1 by applying the same argument as in the proof of Proposition 3.14, using Corollary 3.12 and the strict concavity of $z \mapsto \lambda_{1,z}$.

The proof requires two steps.

Step 1: the pointwise irreducibility of \mathbf{L} can be assumed without loss of generality. Assume the limit has been proved provided $\mathbf{L}(t, x)$ is irreducible at all $(t, x) \in \overline{\Omega}_{\text{per}}$.

Define

$$\mathbf{L} : s \in [0, +\infty) \mapsto \mathbf{L} + (e^s - 1)\mathbf{1}_{N \times N} - (e^s - 1)\mathbf{L}.$$

Obviously, $\mathbf{L}(0) = \mathbf{L}$ and, for all $s \in (0, +\infty)$, $\mathbf{L}(s, t, x)$ is irreducible at all $(t, x) \in \overline{\Omega}_{\text{per}}$. Moreover, by virtue of Propositions 3.5, 3.4 and 3.14, the periodic principal eigenvalue $\lambda'_1(\omega, s)$ associated with the operator

$$\mathcal{Q}_{\omega, s} = \omega \partial_t - \text{diag}(\nabla \cdot (A_i \nabla) - q_i \cdot \nabla) - \mathbf{L}(s)$$

is, as a function of s , continuous in $[0, +\infty)$, decreasing in $[0, +\infty)$, strictly concave in $[0, +\infty)$. By concavity, for all $\omega \in (0, 1]$, the one-sided derivatives of $s \mapsto \lambda'_1(\omega, s)$ are well-defined and satisfy:

$$\begin{aligned} \lambda'_1(\omega, 2) - \lambda'_1(\omega, 1) &< \lim_{\substack{s' < 1 \\ s' \rightarrow 1}} \frac{\lambda'_1(\omega, s') - \lambda'_1(\omega, 1)}{s' - 1} \\ &\leq \lim_{\substack{s' < s \\ s' \rightarrow s}} \frac{\lambda'_1(\omega, s') - \lambda'_1(\omega, s)}{s' - s} \\ &\leq \lim_{\substack{s' > s \\ s' \rightarrow s}} \frac{\lambda'_1(\omega, s') - \lambda'_1(\omega, s)}{s' - s} \\ &\leq \lim_{\substack{s' > 0 \\ s' \rightarrow 0}} \frac{\lambda'_1(\omega, s') - \lambda'_1(\omega, 0)}{s'} \\ &\leq 0. \end{aligned}$$

By assumption, $\mathbf{L}(s)$ being irreducible when $s > 0$, $\lambda'_1(\omega, 1)$ and $\lambda'_1(\omega, 2)$ both converge as $\omega \rightarrow 0$, whence

$$\inf_{\omega \in (0, 1]} \lambda'_1(\omega, 2) - \lambda'_1(\omega, 1) > -\infty.$$

Therefore the family $(s \in [0, 1] \mapsto \lambda'_1(\omega, s))_{\omega \in (0, 1]}$ is uniformly Lipschitz-continuous, and *a fortiori* equicontinuous. By virtue of the Arzelà–Ascoli theorem, it is relatively compact in $\mathcal{C}([0, 1])$.

Let $\lambda \in \mathcal{C}([0, 1])$ be any accumulation point of the family as $\omega \rightarrow 0$. Since we assumed the pointwise convergence

$$\lim_{\substack{\omega > 0 \\ \omega \rightarrow 0}} \lambda'_1(\mathcal{Q}_{\omega, s}) = \frac{1}{T} \int_0^T \lambda'_1(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(s, t)) dt,$$

when $s > 0$, it follows that λ coincides in $(0, 1]$ with

$$s \mapsto \frac{1}{T} \int_0^T \lambda_{1, z}(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(s, t)) dt.$$

By continuity of λ and of the above function (due to Proposition 3.14), they also coincide at $s = 0$. Hence there is a unique accumulation point for the sequence, whence the whole family $(s \mapsto \lambda'_1(\omega, s))_{\omega \in (0, 1]}$ converges uniformly to the above function as $\omega \rightarrow 0$. This implies the pointwise convergence at $s = 0$, and this ends the proof of this step. \square

In the following step we assume, without loss of generality, that $\mathbf{L}(t, x)$ is indeed irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$.

Step 2: the proof in the pointwise irreducible case. Let $\omega > 0$ and $\varepsilon > 0$.

Let

$$\lambda : t \mapsto \lambda'_1(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)).$$

By virtue of the pointwise irreducibility of \mathbf{L} , which implies the full coupling of all operators in the family

$$(-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t))_{t \in [0, T]},$$

the function λ is continuous and periodic. Let $\mathbf{v} : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^N$ be the function such that, for any $t \in [0, T]$, $x \in [0, L] \mapsto \mathbf{v}(t, x)$ is the periodic principal eigenfunction of $-\text{diag}(\nabla \cdot (A_i(t)\nabla) - q_i(t) \cdot \nabla) - \mathbf{L}(t)$, with an appropriate normalization ensuring the continuity of \mathbf{v} in time, *e.g.* $\max_{i \in [N]} v_i(t, 0) = 1$.

Assuming sufficient time regularity of the coefficients $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$, \mathbf{L} , we deduce from classical regularity estimates [52] that $\mathbf{v} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$. The proof in the general case with mere Hölder-continuity in time is not detailed – it can be deduced by a standard regularization procedure and the continuity of periodic principal eigenvalues, *cf.* Step 1.

Since \mathbf{v} is \mathcal{C}^1 with respect to time, so is $(\ln v_i)_{i \in [N]}$, and therefore $(\partial_t v_i / v_i)_{i \in [N]}$ is globally bounded in $\overline{\Omega_{\text{per}}}$. Hence there exists $K > 0$, independent of ω , such that $-K\mathbf{v} \leq \partial_t \mathbf{v} \leq K\mathbf{v}$. Provided $\omega \leq \frac{\varepsilon}{K}$,

$$(-\varepsilon + \lambda)\mathbf{v} \leq \mathcal{Q}_{\omega} \mathbf{v} \leq (\varepsilon + \lambda)\mathbf{v} \quad \text{in } \overline{\Omega_{\text{per}}}.$$

Let

$$v : t \mapsto \exp\left(\frac{1}{\omega} \left(\frac{t}{T} \int_0^T \lambda(t') dt' - \int_0^t \lambda(t') dt'\right)\right)$$

which is positive, periodic and satisfies $\omega v' = \left(\frac{1}{T} \int_0^T \lambda - \lambda\right)v$. Then

$$\left(-\varepsilon + \frac{1}{T} \int_0^T \lambda\right)v\mathbf{v} \leq \omega v' \mathbf{v} + v \mathcal{Q}_{\omega} \mathbf{v} \leq \left(\varepsilon + \frac{1}{T} \int_0^T \lambda\right)v\mathbf{v}.$$

Since $\omega v' \mathbf{v} + v \mathcal{Q}_\omega \mathbf{v} = \mathcal{Q}_\omega(v \mathbf{v})$ and $v \mathbf{v} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$, this shows that $v \mathbf{v}$ can be used both as a super-solution and as a sub-solution to derive from Proposition 3.4 the following inequalities:

$$\frac{1}{T} \int_0^T \lambda(t) dt - \varepsilon \leq \lambda'_1(\mathcal{Q}_\omega) \leq \frac{1}{T} \int_0^T \lambda(t) dt + \varepsilon.$$

Passing to the limit $\varepsilon \rightarrow 0$ ends the proof of this step. \square

Putting the two steps together, the claim is proved. \square

Next we prove the limit $\omega \rightarrow +\infty$.

Proposition 3.18. *For all $z \in \mathbb{R}^n$,*

$$\lim_{\omega \rightarrow +\infty} \lambda_{1,z}(\mathcal{Q}_\omega) = \lambda_{1,z} \left(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}} \right),$$

$$\lim_{\omega \rightarrow +\infty} \lambda_1(\mathcal{Q}_\omega) = \lambda_1 \left(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}} \right),$$

where

$$\left((\hat{A}_i, \hat{q}_i)_{i \in [N]}, \hat{\mathbf{L}} \right) : x \mapsto \frac{1}{T} \int_0^T ((A_i, q_i)_{i \in [N]}, \mathbf{L})(t, x) dt.$$

Proof. Similarly to the proof of Proposition 3.17, it is sufficient to prove only the case $z = 0$.

By virtue of Corollary 3.13, $-\lambda_{\text{PF}}(\bar{\mathbf{L}}) \leq \lambda'_1(\mathcal{Q}_\omega) \leq -\lambda_{\text{PF}}(\underline{\mathbf{L}})$, whence there exists a sequence $(\omega_k)_{k \in \mathbb{N}}$ and $\lambda_\infty \in \mathbb{R}$ such that, as $k \rightarrow +\infty$, $\omega_k \rightarrow +\infty$ and $\lambda_k = \lambda'_1(\mathcal{Q}_{\omega_k}) \rightarrow \lambda_\infty$.

Let $\mathbf{u}_k \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ be the unique generalized principal eigenfunction associated with λ_k satisfying the normalization $\int_{\overline{\Omega_{\text{per}}}} |\mathbf{u}_k|^2 = 1$.

Multiplying $(\mathcal{Q}_{\omega_k} \mathbf{u}_k)_i - \lambda_k u_{k,i}$ by $u_{k,i}$, integrating by parts over $\overline{\Omega_{\text{per}}}$, and using the Cauchy–Schwarz inequality $\int_{\overline{\Omega_{\text{per}}}} u_{k,j} u_{k,i} \leq \|u_{k,j}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})} \|u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})}$, we obtain the uniform boundedness of $(\nabla u_{k,i})_{k \in \mathbb{N}}$ in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$ for each $i \in [N]$, just as in Nadin [59, Proof of Theorem 3.10].

From now on, we assume \mathcal{C}^1 regularity in time of the coefficients $(A_i)_{i \in [N]}$. The proof in the general case with mere Hölder-continuity in time is not detailed – it can be deduced from the \mathcal{C}^1 case by a standard regularization procedure and the continuity of periodic principal eigenvalues.

By integration by parts in time, for each $i \in [N]$ and $k \in \mathbb{N}$,

$$\int_{\overline{\Omega_{\text{per}}}} A_i \nabla u_{k,i} \cdot \nabla \partial_t u_{k,i} = - \int_{\overline{\Omega_{\text{per}}}} \partial_t A_i \nabla u_{k,i} \cdot \nabla u_{k,i} - \int_{\overline{\Omega_{\text{per}}}} A_i \nabla \partial_t u_{k,i} \cdot \nabla u_{k,i},$$

whence, by virtue of the symmetry of A_i , the following identity holds:

$$\int_{\overline{\Omega_{\text{per}}}} A_i \nabla u_{k,i} \cdot \nabla \partial_t u_{k,i} = -\frac{1}{2} \int_{\overline{\Omega_{\text{per}}}} \partial_t A_i \nabla u_{k,i} \cdot \nabla u_{k,i}.$$

From this identity and a space-time integration by parts of $((\mathcal{Q}_{\omega_k} \mathbf{u}_k)_i - \lambda_k u_{k,i}) \partial_t u_{k,i}$, we deduce

$$\begin{aligned} \omega_k \int_{\overline{\Omega_{\text{per}}}} (\partial_t u_{k,i})^2 &= \frac{1}{2} \int_{\overline{\Omega_{\text{per}}}} \partial_t A_i \nabla u_{k,i} \cdot \nabla u_{k,i} \\ &\quad - \int_{\overline{\Omega_{\text{per}}}} (q_i \cdot \nabla u_{k,i}) \partial_t u_{k,i} \\ &\quad + \sum_{j=1}^N \int_{\overline{\Omega_{\text{per}}}} l_{i,j} u_{k,j} \partial_t u_{k,i}. \end{aligned}$$

By the Cauchy–Schwarz inequality and the Young inequality, there exist $A > 0$ and $B > 0$, that only depend on \mathcal{L}^∞ bounds on $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ and \mathbf{L} , such that, for each $i \in [N]$,

$$(\omega_k - A) \|\partial_t u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})}^2 \leq B \|\nabla u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})}^2.$$

Therefore, by finiteness of $\sup_{k \in \mathbb{N}} \|\nabla u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})}$,

$$\|\partial_t u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})} \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Hence, for each $i \in [N]$, $(u_{k,i})_k$, $(\partial_t u_{k,i})_k$ and $(\nabla u_{k,i})_k$ are all uniformly bounded in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$, with $\|\partial_t u_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})} \rightarrow 0$ as well. Therefore, up to extraction of a subsequence, $u_{k,i}$ converges in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$ to a limit $u_{\infty,i}$ and $\nabla u_{k,i}$ and $\partial_t u_{k,i}$ converge weakly in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$ to limits $\nabla u_{\infty,i}$ and $\partial_t u_{\infty,i}$ respectively. By weak lower-semicontinuity of the norm in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$, the convergence $\partial_t u_{k,i} \rightarrow 0$ occurs in fact in the sense of the strong convergence in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$.

Let $\hat{\mathbf{u}}_k : x \mapsto \frac{1}{T} \int_0^T \mathbf{u}_k(t, x) dt$ and $\mathbf{v}_k = \mathbf{u}_k - \hat{\mathbf{u}}_k$. By the Poincaré inequality, there exists a constant $K > 0$ such that, for each $i \in [N]$,

$$\int_0^T (\partial_t u_{k,i})^2 = \int_0^T (\partial_t v_{k,i})^2 \geq K \int_0^T v_{k,i}^2,$$

whence

$$\|v_{k,i}\|_{\mathcal{L}^2(\overline{\Omega_{\text{per}}})} \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Also, since

$$\begin{aligned} \int_{[0,L]} |\hat{u}_{k,i} - u_{\infty,i}| &= \int_{[0,L]} \left| \frac{1}{T} \int_0^T u_{k,i} - \frac{1}{T} \int_0^T u_{\infty,i} \right| \\ &\leq \frac{1}{T} \int_{\overline{\Omega_{\text{per}}}} |u_{k,i} - u_{\infty,i}| \\ &\leq \sqrt{\frac{|[0,L]|}{T}} \left(\int_{\overline{\Omega_{\text{per}}}} (u_{k,i} - u_{\infty,i})^2 \right)^{1/2} \end{aligned}$$

for each $i \in [N]$, $\hat{\mathbf{u}}_k$ converges to \mathbf{u}_∞ in $\mathcal{L}^1([0, L])$. Similarly, for any test function $\varphi \in \mathcal{L}^2_{\text{per}}(\mathbb{R}^n)$,

$$\left| \int_{[0,L]} (\nabla \hat{u}_{k,i} - \nabla u_{\infty,i}) \varphi \right| \leq \frac{1}{T} \left| \int_{\overline{\Omega_{\text{per}}}} (\nabla u_{k,i} - \nabla u_{\infty,i}) \varphi \right|,$$

so that $\nabla \hat{\mathbf{u}}_k \rightharpoonup \nabla \mathbf{u}_\infty$ in $\mathcal{L}^2_{\text{per}}(\mathbb{R}^n)$.

Integrating for any $k \in \mathbb{N}$ the quantity $(\mathcal{Q}_{\omega_k} \mathbf{u}_k)_i - \lambda_k u_{k,i}$ in $[0, T]$ and dividing by T , we deduce

$$\begin{aligned} 0 &= \frac{1}{T} \nabla \cdot \left(\int_0^T (A_i \nabla u_{k,i}) \right) - \frac{1}{T} \int_0^T (q_i \cdot \nabla u_{k,i}) + \frac{1}{T} \sum_{j=1}^N \int_0^T l_{i,j} u_{k,j} + \lambda_k \hat{u}_{k,i} \\ &= \nabla \cdot (\hat{A}_i \nabla \hat{u}_{k,i}) - \hat{q}_i \cdot \nabla \hat{u}_{k,i} + \sum_{j=1}^N \hat{l}_{i,j} \hat{u}_{k,j} + \lambda_k \hat{u}_{k,i} \\ &\quad + \frac{1}{T} \nabla \cdot \left(\int_0^T (A_i \nabla v_{k,i}) \right) - \frac{1}{T} \int_0^T (q_i \cdot \nabla v_{k,i}) + \frac{1}{T} \sum_{j=1}^N \int_0^T l_{i,j} v_{k,j}. \end{aligned}$$

Testing this identity against a test function in $\mathcal{C}_{\text{per}}^2(\mathbb{R}^n)$ and using the convergence of $(\mathbf{v}_k)_k$ to $\mathbf{0}$ in $\mathcal{L}^2(\overline{\Omega_{\text{per}}})$ as well as the convergence of $(\hat{\mathbf{u}}_k)_k$ to \mathbf{u}_{∞} in $\mathcal{L}^1(\overline{\Omega_{\text{per}}})$, we deduce that $x \mapsto \mathbf{u}_{\infty}(x)$ is a weak solution in the dual of $\mathcal{C}_{\text{per}}^2(\mathbb{R}^n)$ of

$$\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) \mathbf{u}_{\infty} + \hat{\mathbf{L}} \mathbf{u}_{\infty} + \lambda \mathbf{u}_{\infty} = \mathbf{0}.$$

By density, this remains true with test functions in $\mathcal{H}_{\text{per}}^1(\mathbb{R}^n)$, or in other words \mathbf{u}_{∞} is a weak solution on $\mathcal{H}_{\text{per}}^{-1}(\mathbb{R}^n)$. By elliptic regularity [33], $\mathbf{u}_{\infty} \in \mathcal{H}_{\text{per}}^1(\mathbb{R}^n)$ is in fact a classical solution, in $\mathcal{C}_{\text{per}}^2(\mathbb{R}^n)$. Since \mathbf{u}_{∞} is nonnegative and satisfies the normalization $\int_{\overline{\Omega_{\text{per}}}} |\mathbf{u}_{\infty}|^2 = |[0, L]| \int_0^T |\mathbf{u}_{\infty}|^2 = 1$, it is nonnegative nonzero, and then positive by the maximum principle (the elliptic operator under consideration is fully coupled in $[0, L]$ by (A_3)), whence it is a generalized principal eigenfunction associated with $\lambda'_1(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}})$. Thus $\lambda = \lambda'_1(-\text{diag}(\nabla \cdot (\hat{A}_i \nabla) - \hat{q}_i \cdot \nabla) - \hat{\mathbf{L}})$.

As a conclusion, the accumulation point of $(\lambda_k)_{k \in \mathbb{N}}$ is unique and therefore the whole sequence converges. Subsequently, the whole family $(\lambda'_1(\mathcal{Q}_{\omega}))_{\omega > 0}$ converges. \square

3.3. Formulas and estimates in special cases: proof of Theorems 1.12–1.16. We begin this subsection by recalling that space, time or space-time homogeneous coefficients in \mathcal{Q} lead to the reduced formulas (8), (9), (10), respectively.

3.3.1. Formulas for operators with space homogeneity: proof of Theorem 1.12. Recall the notations $\hat{A}_i, \hat{q}_i, \hat{\mathbf{L}}$ for the averages in time, $\langle A_i \rangle, \langle q_i \rangle, \langle \mathbf{L} \rangle$ for the averages in space and $\langle \hat{A}_i \rangle, \langle \hat{q}_i \rangle, \langle \hat{\mathbf{L}} \rangle$ for the averages in space-time.

Proposition 3.19. *Let $z \in \mathbb{R}^n$. If*

- (1) A_1, q_1 and \mathbf{L} do not depend on x ,
- (2) there exists a constant positive vector $\mathbf{u} \in (\mathbf{0}, \infty)$ such that \mathbf{u} is a Perron–Frobenius eigenvector of $\mathbf{L}(t)$ for all $t \in \mathbb{R}$,
- (3) either $z = 0$ or $(A_1, q_1) = (A_2, q_2) = \dots = (A_N, q_N)$,

then

$$\lambda_{1,z} = -z \cdot \hat{A}_1 z + \hat{q}_1 \cdot z - \lambda_{\text{PF}}(\hat{\mathbf{L}}).$$

Proof. First, writing the equality satisfied by \mathbf{u} and taking the average in time, we obtain $\frac{1}{T} \int_0^T \lambda_{\text{PF}}(\mathbf{L}(t)) dt = \lambda_{\text{PF}}(\hat{\mathbf{L}})$. Note that $\hat{\mathbf{L}}$ is irreducible.

Next, let $f : t \mapsto -z \cdot A_1(t)z + q_1(t) \cdot z - \lambda_{\text{PF}}(\mathbf{L}(t))$. By uniqueness of the periodic principal eigenpair of \mathcal{Q} , it suffices to verify that the space-independent function

$$(t, x) \mapsto \exp \left(- \int_0^t f(t') dt' + \frac{t}{T} \int_0^T f \right) \mathbf{u}$$

is a $\mathcal{C}^{1,2}$, periodic, positive eigenfunction of \mathcal{Q}_z associated with the eigenvalue $\frac{1}{T} \int_0^T f(t) dt$. The continuity of $t \mapsto \lambda_{\text{PF}}(\mathbf{L}(t))$ follows from (A₄) and the continuity of the Perron–Frobenius eigenvalue as function of the entries of the matrix. \square

Corollary 3.20. *If*

- (1) A_1, q_1 and \mathbf{L} do not depend on x ,
- (2) there exists a constant positive vector $\mathbf{u} \in (\mathbf{0}, \infty)$ such that \mathbf{u} is a Perron–Frobenius eigenvector of $\mathbf{L}(t)$ for all $t \in \mathbb{R}$,

then

$$\lambda'_1 = -\lambda_{\text{PF}}(\hat{\mathbf{L}}).$$

Furthermore, if $(A_1, q_1) = (A_2, q_2) = \dots = (A_N, q_N)$, then $\lambda_1 = \lambda'_1$ if and only if $\hat{q}_1 = 0$.

Remark 3.13. Although we do not know if the second condition in the statement is truly optimal, we know that the first condition alone cannot be sufficient. Indeed, simple counter-examples exist.

For instance, consider in dimension $N = 2$ the matrix

$$\mathbf{L} : t \mapsto \begin{pmatrix} 0 & \eta(t) \\ \eta(t - T/2) & 0 \end{pmatrix}$$

where η is the continuous T -periodic function that coincides on $[0, T]$ with $t \mapsto \max(\sin(\frac{2\pi}{T}t), 0)$.

Even though $\mathbf{L}(t)$ is actually always reducible, its Perron–Frobenius eigenvalue, understood as the continuous extension of the Perron–Frobenius eigenvalue to essentially nonnegative matrices, is 0 for all $t \in [0, T]$, it is always a geometrically simple eigenvalue and its unit Perron–Frobenius eigenvector is $(1, 0)^T$ in $(0, T/2)$ and $(0, 1)^T$ in $(T/2, T)$. The matrix $\hat{\mathbf{L}}$ is symmetric and admits $\mathbf{1}$ as Perron–Frobenius eigenvector and $1/\pi$ as Perron–Frobenius eigenvalue.

Due to the uniqueness of the periodic principal eigenfunction and the symmetries of \mathbf{L} , the periodic principal eigenfunction of $\partial_t - \Delta - \mathbf{L}$, namely that of $\frac{d}{dt} - \mathbf{L}$, necessarily has the form $\mathbf{u} : t \mapsto (u(t), u(t - T/2))^T$. Moreover, we can choose to normalize it with $u(0) = 1$. The scalar function u satisfies $u'(t) = \eta(t)u(t - T/2) + \lambda'_1 u(t)$ for all $t \in [0, T]$, i.e. $u'(t) = \sin(\frac{2\pi}{T}t)u(t - T/2) + \lambda'_1 u(t)$ for all $t \in [0, T/2]$ and $u'(t) = \lambda'_1 u(t)$ for all $t \in [T/2, T]$. It follows that u satisfies:

$$u(t)e^{-\lambda'_1 t} = 1 + \int_0^t e^{-\lambda'_1 t'} \sin\left(\frac{2\pi}{T}t'\right) u(t' - T/2) dt' \quad \text{for all } t \in [0, T/2],$$

$$u(t)e^{-\lambda'_1(t-T/2)} = u(T/2) \quad \text{for all } t \in [T/2, T].$$

Since $u(T/2) = u(T)e^{-\lambda'_1(T-T/2)} = u(0)e^{-\lambda'_1 T/2} = e^{-\lambda'_1 T/2}$ and since, for all $t' \in [0, T/2]$, $u(t' - T/2) = u(t' + T/2) = u(T/2)e^{\lambda'_1(t'+T/2-T/2)} = e^{\lambda'_1(t'-T/2)}$, the first

equality, for $t \in [0, T/2]$, is simplified as

$$\begin{aligned} u(t) &= e^{\lambda'_1 t} + e^{\lambda'_1(t-T/2)} \int_0^t \sin\left(\frac{2\pi}{T}t'\right) dt' \\ &= e^{\lambda'_1 t} + e^{\lambda'_1(t-T/2)} \left(-\frac{T}{2\pi}\right) \left(\cos\left(\frac{2\pi}{T}t\right) - 1\right) \end{aligned}$$

whence, evaluating at $t = T/2$,

$$e^{-\lambda'_1 T/2} = u(T/2) = e^{\lambda'_1 T/2} + \frac{T}{\pi}$$

i.e. $\sinh\left(\frac{\lambda'_1 T}{2}\right) = -\frac{T}{2\pi}$, i.e.

$$\lambda'_1 = \frac{2}{T} \sinh^{-1}\left(-\frac{T}{2\pi}\right) = -\frac{2}{T} \ln\left(\frac{T}{2\pi} + \sqrt{\frac{T^2}{4\pi^2} + 1}\right).$$

On one hand, the above equality shows that $\lambda'_1 < 0$, independently of the value of T . On the other hand, it is easily verified that 1 is not in the image of $\tau \in (0, +\infty) \mapsto \frac{\ln(\tau + \sqrt{\tau^2 + 1})}{\tau}$, whence $\lambda'_1 \neq -\frac{1}{\pi}$, also independently of the value of T .

Therefore this counter-example shows that in general, $\lambda_{1,\text{per}}\left(\frac{d}{dt} - \mathbf{L}\right)$ coincides neither with $-\frac{1}{T} \int_0^T \lambda_{\text{PF}}(\mathbf{L}) = 0$ nor with $-\lambda_{\text{PF}}(\hat{\mathbf{L}}) = -\frac{1}{\pi}$.

Note also that this counter-example is consistent with the formulas for asymptotics $T \rightarrow 0$ and $T \rightarrow +\infty$ of Theorem 1.9, that predict $\lambda'_1(T) \rightarrow \frac{1}{T} \int_0^T -\lambda_{\text{PF}}(\mathbf{L})$ as $T \rightarrow \infty$ and $\lambda'_1(T) \rightarrow -\lambda_{\text{PF}}(\hat{\mathbf{L}})$ as $T \rightarrow 0$.

Remark 3.14. Consider a diagonal perturbation of \mathbf{L} of the form $\mathbf{L}_\nu = \mathbf{L} - \nu \mathbf{I}$ with $\nu > 0$. At some arbitrary time $t_0 \in [0, T]$, consider the “frozen in time” system of ordinary differential equations $\mathbf{u}'(t) = \mathbf{L}_\nu(t_0)\mathbf{u}(t)$: its periodic principal eigenvalue is $-\lambda_{\text{PF}}(\mathbf{L}_\nu(t_0)) = \nu > 0$. However, the periodic principal eigenvalue of the “unfrozen” nonautonomous system $\mathbf{u}'(t) = \mathbf{L}_\nu(t)\mathbf{u}(t)$ is $\lambda'_1 + \nu$, that remains negative provided $\nu > 0$ is small enough. Therefore, although the trajectories of the “frozen” system converge exponentially fast to $\mathbf{0}$, the nonnegative nonzero solutions of the “unfrozen” system diverge from $\mathbf{0}$ exponentially fast. In this sense, the stability properties of the two systems are unrelated.

This fact should not surprise readers familiar with nonautonomous dynamical systems, since the existence of such counter-examples, relying strongly upon the non-symmetry, is classical. We highlight it here for other readers.

3.3.2. Formulas for operators with time homogeneity: proof of Theorem 1.13. Now we turn to the proof of Theorem 1.13. We will use the following well-known property concerning variational formulas in the self-adjoint elliptic case.

Proposition 3.21. *If*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
- (2) $\mathbf{L}(x)$ is symmetric for all $x \in \mathbb{R}^n$,
- (3) $q_1 = q_2 = \dots = q_N = 0$,

then the periodic principal eigenvalue of $\mathcal{L} = \text{diag}(\nabla \cdot (A_i \nabla)) + \mathbf{L}$ satisfies:

$$\lambda'_1(-\mathcal{L}) = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2}.$$

and, for any nonempty bounded smooth open set Ω , the Dirichlet principal eigenvalue satisfies:

$$\lambda_1(-\mathcal{L}, \Omega) = \min_{\mathbf{u} \in \mathcal{C}_0^1(\Omega, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{\Omega} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{\Omega} |\mathbf{u}|^2}.$$

These formulas being recalled, we are in a position to prove Theorem 1.13. We first focus on the case $z = 0$, and will then deduce the general case as a consequence of this one. Also, since the statement of Theorem 1.13 involves the inverses of the diffusion matrices A_i , we recall that their invertibility follows from their uniform ellipticity (\mathbf{A}_1) . Moreover, the periodicity and regularity of the inverses follows from the periodicity and regularity of the diffusion matrices (\mathbf{A}_4) .

Proposition 3.22. *If*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
- (2) $\mathbf{L}(x)$ is symmetric for all $x \in \mathbb{R}^n$,
- (3) there exists $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ such that $\int_{[0,L]} \nabla Q = 0$ and

$$A_1^{-1} q_1 = A_2^{-1} q_2 = \cdots = A_N^{-1} q_N = \nabla Q,$$

then

$$\lambda_1 = \lambda'_1 = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L}_Q \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2},$$

where

$$\mathbf{L}_Q = \mathbf{L} + \frac{1}{4} \text{diag} (2\nabla \cdot q_i - A_i^{-1} q_i \cdot q_i) = \mathbf{L} + \frac{1}{4} \text{diag} (2\nabla \cdot (A_i \nabla Q) - \nabla Q \cdot A_i \nabla Q).$$

Proof. Let $\mathcal{L} = \text{diag}(\nabla \cdot (A_i \nabla) - q_i \cdot \nabla) + \mathbf{L}$.

Step 1: the case $q_1 = q_2 = \cdots = q_N = 0$, i.e. Q constant. By (9), we only have to prove the statement for the elliptic operator \mathcal{L} . Also, we already know that $\lambda'_{1,0} = \lambda_{1,0} \leq \lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1,z}$.

Following Berestycki–Rossi [16], we consider an even function $\chi \in \mathcal{C}^\infty(\mathbb{R}, [0, 1])$ such that $\chi = 0$ in $\mathbb{R} \setminus [-1, 1]$ and $\chi(0) = 1$. Next, we construct a family of radial smooth cut-off functions $(\chi_R)_{R>1}$ such that, for each $R > 1$, $\chi_R = 1$ in B_{R-1} and $\chi_R(x) = \chi(|x| - (R-1))$ if $x \in \mathbb{R}^n \setminus B_{R-1}$, where $B_{R-1} = B(0, R-1)$. By construction, the family $(\|\chi_R\|_{\mathcal{C}^\infty(\mathbb{R}^n, \mathbb{R})})_{R>1}$ is bounded.

Let $R > 1$ and denote μ_R the Dirichlet principal eigenvalue $\lambda_{1,\text{Dir}}(-\mathcal{L}, B_R)$, where $B_R = B(0, R)$. By Proposition 3.21,

$$\mu_R = \min_{\mathbf{u} \in \mathcal{C}_0^1(B_R, \mathbb{R}^N) \setminus \{0\}} \frac{\int_{B_R} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{B_R} |\mathbf{u}|^2}.$$

Taking $\chi_R \mathbf{u}_0$ as test function (recall that \mathbf{u}_0 is a positive periodic principal eigenfunction of \mathcal{L}) and using the equality $-\mathcal{L} \mathbf{u}_0 = \lambda'_1 \mathbf{u}_0$ satisfied pointwise in B_{R-1} , we find:

$$\begin{aligned} \mu_R &\leq \frac{\int_{B_{R-1}} \chi_R \mathbf{u}_0^T (-\mathcal{L})(\chi_R \mathbf{u}_0) + \int_{B_R \setminus B_{R-1}} \chi_R \mathbf{u}_0^T (-\mathcal{L})(\chi_R \mathbf{u}_0)}{\int_{B_R} \chi_R^2 |\mathbf{u}_0|^2} \\ &= \frac{\int_{B_{R-1}} \mathbf{u}_0^T (-\mathcal{L}) \mathbf{u}_0 - \int_{B_R \setminus B_{R-1}} \chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)}{\int_{B_R} \chi_R^2 |\mathbf{u}_0|^2} \\ &= \frac{\lambda'_1 \int_{B_{R-1}} |\mathbf{u}_0|^2 - \int_{B_R \setminus B_{R-1}} \chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)}{\int_{B_R} \chi_R^2 |\mathbf{u}_0|^2} \\ &= \frac{\lambda'_1 \int_{B_{R-1}} \chi_R^2 |\mathbf{u}_0|^2 - \int_{B_R \setminus B_{R-1}} \chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)}{\int_{B_R} \chi_R^2 |\mathbf{u}_0|^2} \\ &= \lambda'_1 - \frac{\lambda'_1 \int_{B_R \setminus B_{R-1}} \chi_R^2 |\mathbf{u}_0|^2 + \int_{B_R \setminus B_{R-1}} \chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)}{\int_{B_R} \chi_R^2 |\mathbf{u}_0|^2}. \end{aligned}$$

Using now the triangle inequality, the inequality $\chi_R \geq (\chi_R)|_{B_{R-1}}$, the discrete Cauchy–Schwarz inequality and then pointwise upper and lower bounds, we find:

$$\begin{aligned} \mu_R &\leq \lambda'_1 + \frac{|\lambda'_1| \int_{B_R \setminus B_{R-1}} \chi_R^2 |\mathbf{u}_0|^2 + \int_{B_R \setminus B_{R-1}} |\chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)|}{\int_{B_{R-1}} \chi_R^2 |\mathbf{u}_0|^2} \\ &\leq \lambda'_1 + \frac{\int_{B_R \setminus B_{R-1}} (|\lambda'_1| \chi_R^2 |\mathbf{u}_0|^2 + |\chi_R \mathbf{u}_0^T \mathcal{L}(\chi_R \mathbf{u}_0)|)}{\int_{B_{R-1}} |\mathbf{u}_0|^2} \\ &\leq \lambda'_1 + \frac{\int_{B_R \setminus B_{R-1}} (|\lambda'_1| \chi_R^2 |\mathbf{u}_0|^2 + |\chi_R \mathbf{u}_0| |\mathcal{L}(\chi_R \mathbf{u}_0)|)}{\min_{x \in \Omega_{\text{per}}} |\mathbf{u}_0(x)|^2 \int_{B_{R-1}} 1} \\ &\leq \lambda'_1 + \frac{|\lambda'_1| \|\chi_R \mathbf{u}_0\|^2 + \|\chi_R \mathbf{u}_0\| \|\mathcal{L}(\chi_R \mathbf{u}_0)\| \int_{B_R \setminus B_{R-1}} 1}{\min_{x \in \Omega_{\text{per}}} |\mathbf{u}_0(x)|^2 \int_{B_{R-1}} 1} \end{aligned}$$

where the norm $\|\cdot\|$ is defined as $\|\mathbf{v}\| = \sup_{x \in \mathbb{R}^n} |\mathbf{v}(x)|$ (appropriate for $\mathcal{C}(\mathbb{R}^n, \mathbb{R}^N)$). Thus, from the boundedness of the operator $\mathcal{L} : \mathcal{C}^2(\mathbb{R}^n, \mathbb{R}^N) \rightarrow \mathcal{C}(\mathbb{R}^n, \mathbb{R}^N)$ and the boundedness in $\mathcal{C}^2(\mathbb{R}^n, \mathbb{R}^N)$ of the family $(\chi_R \mathbf{u}_0)_{R>1}$, there exists a constant $K > 0$, independent of R , such that

$$\mu_R \leq \lambda'_1 + K \frac{R^{n-1}}{(R-1)^n},$$

and, passing to the limit $R \rightarrow +\infty$, we deduce finally $\lambda_1 \leq \lambda'_1$ (the proof of the convergence of the Dirichlet principal eigenvalues in balls of increasing radius to the generalized principal eigenvalue λ_1 for the elliptic operator $-\mathcal{L}$ is done exactly as in the parabolic case, see Proposition 3.2). Hence $\lambda'_1 = \lambda_1$.

The conclusion of this step follows from Proposition 3.21. \square

Step 2: the general case. From now on, for all $i \in [N]$, $q_i = A_i \nabla Q$ with $\int_{[0,L]} \nabla Q = 0$. Following Berestycki–Hamel–Rossi [14], the idea is to change variables to reduce this case to the previous one.

Preliminarily, we check that $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ is necessarily space periodic. Fix $\alpha \in [n]$. The function $x \mapsto Q(x + L_\alpha e_\alpha) - Q(x)$, where $e_\alpha = (\delta_{\alpha\beta})_{\beta \in [n]}$, is constant, since

$$\nabla(Q(x + L_\alpha e_\alpha) - Q(x)) = A_1^{-1}(x + L_\alpha e_\alpha)q_1(x + L_\alpha e_\alpha) - A_1^{-1}(x)q_1(x) = 0.$$

Then

$$\begin{aligned} Q(L_\alpha e_\alpha) - Q(0) &= (|[0, L]|)^{-1} \int_{[0, L]} Q(x + L_\alpha e_\alpha) - Q(x) dx \\ &= (|[0, L]|)^{-1} \int_0^{L_\alpha} \int_{[0, L]} \frac{\partial Q}{\partial x_\alpha}(x + s e_\alpha) dx ds \\ &= 0 \end{aligned}$$

Hence Q is indeed periodic with respect to x_α , and then with respect to x .

Then, introducing for any $\mathbf{u} \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R}^N)$ the transformation

$$\mathbf{v} : x \mapsto \exp(Q(x)/2)\mathbf{u}(x)$$

and following [14], we get:

$$\begin{aligned} -(\mathcal{L}\mathbf{v})_i &= e^{Q/2} \left[-(\mathcal{L}\mathbf{u})_i - \frac{1}{2} \left(u_i \nabla \cdot (A_i \nabla Q) + 2 \nabla u_i \cdot (A_i \nabla Q) + \frac{1}{2} u_i \nabla Q \cdot (A_i \nabla Q) - u_i q_i \cdot \nabla Q \right) \right] \\ &= e^{Q/2} \left[-(\mathcal{L}\mathbf{u})_i - \frac{1}{2} \left(u_i \nabla \cdot q_i + 2 \nabla u_i \cdot q_i - \frac{1}{2} u_i \nabla Q \cdot q_i \right) \right] \\ &= e^{Q/2} \left[-\nabla \cdot (A_i \nabla u_i) - (\mathcal{L}\mathbf{u})_i - \frac{1}{2} \left(\nabla \cdot q_i - \frac{1}{2} \nabla Q \cdot q_i \right) u_i \right]. \end{aligned}$$

Therefore \mathbf{v} is an eigenfunction of $-\mathcal{L}$ if and only if \mathbf{u} is an eigenfunction of the new periodic elliptic operator:

$$-\mathcal{L}_Q = -\text{diag}(\nabla \cdot (A_i \nabla)) - \mathbf{L}_Q.$$

Consequently, the periodic and Dirichlet principal eigenvalues coincide: for instance, with \mathbf{u} a periodic principal eigenfunction of $-\mathcal{L}_Q$, \mathbf{v} satisfies $-\mathcal{L}\mathbf{v} = \lambda'_1(-\mathcal{L}_Q)\mathbf{v}$, and since $\mathbf{v} = e^{Q/2}\mathbf{u}$ is periodic, it is then (by uniqueness) a periodic principal eigenfunction of $-\mathcal{L}$, whence $\lambda'_1(-\mathcal{L}) = \lambda'_1(-\mathcal{L}_Q)$.

By virtue of Step 1,

$$\lim_{R \rightarrow +\infty} \lambda_1(-\mathcal{L}_Q, B_R) = \lambda'_1(-\mathcal{L}_Q)$$

and consequently

$$\lim_{R \rightarrow +\infty} \lambda_1(-\mathcal{L}, B_R) = \lambda'_1(-\mathcal{L}).$$

Therefore

$$\lambda_1(-\mathcal{L}) = \lambda'_1(-\mathcal{L}) = \lambda'_1(-\mathcal{L}_Q).$$

The conclusion of this step follows from the variational formula for the operator $-\mathcal{L}_Q$, see Proposition 3.21. \square

The proof is ended. \square

Remark 3.15. The symmetry assumption on \mathbf{L} is crucial, both for the equality between λ_1 and λ'_1 (as explained above in Remark 3.5) and for the equality between λ'_1 and the minimized integral.

Denote

$$R = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N |\nabla u_i|^2 - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2},$$

which is the quotient appearing in the variational formula in the special case $q_i = 0$ and $A_i = \text{Id}$ for each $i \in [N]$.

It is well-known that for a general non-symmetric square matrix, the maximum of the Rayleigh quotient needs not coincide with the dominant eigenvalue. More precisely, the maximum of the Rayleigh quotient of a matrix \mathbf{L} coincides with the dominant eigenvalue of the symmetric part $\frac{1}{2}(\mathbf{L} + \mathbf{L}^T)$. Similarly, R is the periodic principal eigenvalue of the symmetrized operator $-\Delta - \frac{1}{2}(\mathbf{L} + \mathbf{L}^T)$.

Therefore, using a constant irreducible non-symmetric matrix

$$\mathbf{L} = \begin{pmatrix} 1 & 1 \\ \varepsilon & 1 \end{pmatrix} \quad \text{with } \varepsilon > 0,$$

we obtain a counter-example of the equality between $\lambda'_1 = -1 - \sqrt{\varepsilon}$ and $R = -(3 + \varepsilon)/2$.

Subsequently, replacing $(q_i)_{i \in [N]}$ by $(q_i - 2A_i z)_{i \in [N]}$ and \mathbf{L} by $\mathbf{L} + \text{diag}(z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z)$, we obtain the following corollary, which is the full statement of Theorem 1.13.

Corollary 3.23. *If*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
- (2) $\mathbf{L}(x)$ is symmetric for all $x \in \mathbb{R}^n$,
- (3) there exists $z \in \mathbb{R}^n$ and $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ such that $\int_{[0,L]} \nabla Q = 0$ and

$$A_1^{-1} q_1 = A_2^{-1} q_2 = \dots = A_N^{-1} q_N = 2z + \nabla Q,$$

then

$$\lambda_1 = \lambda_{1,z} = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N \nabla u_i \cdot A_i \nabla u_i - \mathbf{u}^T \mathbf{L}_{Q,z} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2},$$

where

$$\mathbf{L}_{Q,z} = \mathbf{L}_Q + \text{diag}(\nabla \cdot (A_i z) - z \cdot A_i(z + \nabla Q))$$

and \mathbf{L}_Q is defined as in the statement of Proposition 3.22.

With no symmetry assumption on \mathbf{L} and more general advection terms, we can still compare $\lambda_{1,z}$ with the variational formula.

Corollary 3.24. *Let $z \in \mathbb{R}^n$. If*

- (1) $(A_i, q_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
- (2) for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) \leq 0$,
- (3) for all $i \in [N]$, $q_i \cdot z \geq 0$,

then

$$\lambda_{1,z} \geq \lambda_{1,z} \left(\partial_t - \text{diag}(\nabla \cdot (A_i \nabla)) - \frac{1}{2}(\mathbf{L} + \mathbf{L}^T) \right).$$

Proof. By time homogeneity of the coefficients, the periodic principal eigenfunction of the parabolic operator \mathcal{Q}_z is time homogeneous. Taking the scalar product between the periodic principal eigenfunction $\mathbf{u}_z \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, (\mathbf{0}, \infty))$ associated with $\lambda_{1,z}$ and $\mathcal{Q}_z \mathbf{u}_z$ and then integrating in $[0, L]$, we get immediately:

$$\begin{aligned} \lambda_{1,z} \int_{[0,L]} |\mathbf{u}_z|^2 &= \sum_{i=1}^N \int_{[0,L]} \nabla u_{z,i} \cdot A_i \nabla u_{z,i} - \int_{[0,L]} \mathbf{u}_z^T \mathbf{L} \mathbf{u}_z \\ &\quad - \sum_{i=1}^N \int_{[0,L]} \left(\frac{1}{2} \nabla \cdot (q_i - 2A_i z) + (z \cdot A_i z) + \nabla \cdot (A_i z) - (q_i \cdot z) \right) u_{z,i}^2. \end{aligned}$$

The conclusion follows from $\mathbf{u}_z^T \mathbf{L} \mathbf{u}_z = \mathbf{u}_z^T \frac{1}{2}(\mathbf{L} + \mathbf{L}^T) \mathbf{u}_z$, the sign assumptions on $\nabla \cdot (q_i - 2A_i z)$ and $q_i \cdot z$ and the variational formula of Theorem 1.13 applied to the operator $\partial_t - \text{diag}(\nabla \cdot (A_i \nabla)) - \frac{1}{2}(\mathbf{L} + \mathbf{L}^T)$. \square

Remark 3.16. In the nonnegative square matrix context, the inequality $\lambda_{\text{PF}}(\mathbf{L}) \leq \lambda_{\text{PF}}\left(\frac{1}{2}(\mathbf{L} + \mathbf{L}^T)\right)$ is a consequence of a theorem by Levinger which states that $t \in [0, 1] \mapsto \lambda_{\text{PF}}\left(t\mathbf{L} + (1-t)\mathbf{L}^T\right)$ is nondecreasing in $[0, 1/2]$, nonincreasing in $[1/2, 1]$, and that the function is constant if and only if the unit Perron–Frobenius eigenvectors of \mathbf{L} and \mathbf{L}^T coincide. There are many works on this theorem and on its extension to Banach spaces. We refer for instance to the recent paper of Altenberg–Cohen [4] and references therein.

Remark 3.17. The second and third assumptions are obviously satisfied if q_i is divergence-free and $z = 0$, but it is also interesting to consider for instance the case $z \neq 0$ with shear flows $q_i : x \mapsto (\alpha_i(x_2, \dots, x_n), 0, \dots, 0)^T$ with α_i of constant sign. In biological applications (climate change at constant speed towards the north, fish populations living in a river, etc.) or when studying planar spreading, such shear flows appear naturally.

Remark 3.18. We emphasize that the estimate of the above corollary on $\lambda_{1,z}$ is a lower estimate, contrarily to most estimates in this work which are upper estimates (see Subsection 1.5.5).

Taking as a test function in the variational formula of Theorem 1.13 any constant eigenvector of $\mathbf{L}_{Q,z}(x)$, we obtain that the average of the corresponding eigenvalue is smaller than or equal to $-\lambda_1 = -\lambda_{1,z}$. In particular, noting that a constant Perron–Frobenius eigenvector implies $\frac{1}{|[0,L]|} \int_{[0,L]} \lambda_{\text{PF}}(\mathbf{L}_{Q,z}(x)) dx = \lambda_{\text{PF}}(\langle \mathbf{L}_{Q,z} \rangle)$, the following corollary holds.

Corollary 3.25. *If*

- (1) $(A_i)_{i \in [N]}$ and \mathbf{L} do not depend on t ,
- (2) $\mathbf{L}(x)$ is symmetric for all $x \in \mathbb{R}^n$,
- (3) there exists $z \in \mathbb{R}^n$ and $Q \in \mathcal{C}^2(\mathbb{R}^n, \mathbb{R})$ such that $\int_{[0,L]} \nabla Q = 0$ and

$$A_1^{-1} q_1 = A_2^{-1} q_2 = \dots = A_N^{-1} q_N = 2z + \nabla Q,$$

(4) there exists a constant positive vector $\mathbf{u} \in (\mathbf{0}, \infty)$ such that \mathbf{u} is a Perron–Frobenius eigenvector of $\mathbf{L}_{Q,z}(x)$ for all $x \in \mathbb{R}^n$,

then

$$\lambda_1 = \lambda_{1,z} \leq -\lambda_{\text{PF}}(\langle \mathbf{L}_{Q,z} \rangle).$$

Remark 3.19. Again, denote

$$R = \min_{\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, \mathbb{R}^N) \setminus \{\mathbf{0}\}} \frac{\int_{[0,L]} \left(\sum_{i=1}^N |\nabla u_i|^2 - \mathbf{u}^T \mathbf{L} \mathbf{u} \right)}{\int_{[0,L]} |\mathbf{u}|^2}.$$

Let us construct a counter-example where all the conditions of the statement are satisfied but where, due to heterogeneities in $\mathbf{L}(x)$,

$$R < -\lambda_{\text{PF}}(\langle \mathbf{L} \rangle) = -\frac{1}{|[0,L]|} \int_{[0,L]} \lambda_{\text{PF}}(\mathbf{L}(x)) \, dx.$$

(The existence of such counter-examples in the scalar setting is well-known, we provide a vector counter-example just for the sake of completeness.)

In a spirit similar to that of Remark 3.13, we set

$$\mathbf{L} : x \mapsto \begin{pmatrix} 1 & \eta(x_1) \\ \eta(x_1) & 1 \end{pmatrix}$$

where η is the continuous L_1 -periodic function that coincides on $[0, L_1]$ with $x_1 \mapsto \max(L_1/4 - |x_1 - L_1/4|, 0)$.

For all $x \in [0, L]$, $\lambda_{\text{PF}}(\mathbf{L}(x)) = 1 + \eta(x_1)$ with constant eigenvector $\mathbf{1}$, whence

$$R \leq -\frac{1}{L_1} \int_0^{L_1} (1 + \eta(x_1)) \, dx_1 = -1 - \frac{L_1}{16}.$$

Considering test functions of the form $\mathbf{u}(x) = u(x_1)\mathbf{1}$, we get

$$R \leq \min_{u \in \mathcal{C}_{\text{per}}^2(\mathbb{R}), \|u\|_{\mathcal{C}^2} = 1} \int_0^{L_1} (|u'(x_1)|^2 - (1 + \eta(x_1)) u(x_1)^2) \, dx_1$$

Testing against (a \mathcal{C}^2 approximation of) $u : x_1 \mapsto \frac{8\sqrt{3}}{L_1^{3/2}} \eta(x_1)$, we find:

$$\begin{aligned} R &\leq \frac{96}{L_1^3} \int_0^{L_1} (1 - \eta(x_1)^2 - \eta(x_1)^3) \, dx_1 \\ &= -1 + \frac{96}{L_1^2} - \frac{3L_1}{16} \\ &< -1 - \frac{L_1}{16} \quad \text{if } L_1 > 768^{1/3}. \end{aligned}$$

Remark 3.20. More interestingly, in the vector setting, the inequality

$$-R \geq \frac{1}{|[0,L]|} \int_{[0,L]} \lambda_{\text{PF}}(\mathbf{L})$$

might not be satisfied if the fourth assumption of the statement, regarding the existence of a constant positive eigenvector, fails.

To verify this claim, we consider the counter-example $\mathcal{Q}\mathbf{u} = \partial_t \mathbf{u} - \Delta \mathbf{u} - \mathbf{L}\mathbf{u}$, where

$$\mathbf{L} : x \mapsto \frac{1}{1 + \eta(x_1) + \eta(x_1 - \frac{L_1}{3}) + \eta(x_1 - \frac{2L_1}{3})} \begin{pmatrix} 1 + \eta(x_1 - L_1/3) & \eta(x_1) \\ \eta(x_1) & 1 + \eta(x_1 - 2L_1/3) \end{pmatrix},$$

where, this time, η is the continuous L_1 -periodic function that coincides on $[-L_1/2, L_1/2]$ with $x_1 \mapsto \max(L_1/6 - |x_1|, 0)$. Interval by interval, \mathbf{L} satisfies:

$$\begin{aligned}\mathbf{L}(x) &= \frac{1}{1 + \eta(x_1 - \frac{2L_1}{3})} \begin{pmatrix} 1 & 0 \\ 0 & 1 + \eta(x_1 - \frac{2L_1}{3}) \end{pmatrix} \quad \text{if } x_1 \in \left[-\frac{L_1}{2}, -\frac{L_1}{6}\right] + \mathbb{Z}L_1, \\ \mathbf{L}(x) &= \frac{1}{1 + \eta(x_1)} \begin{pmatrix} 1 & \eta(x_1) \\ \eta(x_1) & 1 \end{pmatrix} \quad \text{if } x_1 \in \left[-\frac{L_1}{6}, \frac{L_1}{6}\right] + \mathbb{Z}L_1, \\ \mathbf{L}(x) &= \frac{1}{1 + \eta(x_1 - \frac{L_1}{3})} \begin{pmatrix} 1 + \eta(x_1 - \frac{L_1}{3}) & 0 \\ 0 & 1 \end{pmatrix} \quad \text{if } x_1 \in \left[\frac{L_1}{6}, \frac{L_1}{2}\right] + \mathbb{Z}L_1.\end{aligned}$$

Hence the Perron–Frobenius eigenvalue of $\mathbf{L}(x)$ is 1, for all $x \in [0, L]$, and the unique one-dimensional left-continuous unit Perron–Frobenius eigenvector is:

$$\mathbf{u}_{\text{PF}} : x \mapsto \begin{cases} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} & \text{if } x_1 \in (0, \frac{L_1}{6}] \cup (\frac{5L_1}{6}, L_1] + \mathbb{Z}L_1, \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \text{if } x_1 \in (\frac{L_1}{6}, \frac{L_1}{2}] + \mathbb{Z}L_1, \\ \begin{pmatrix} 0 \\ 1 \end{pmatrix} & \text{if } x_1 \in (\frac{L_1}{2}, \frac{5L_1}{6}] + \mathbb{Z}L_1. \end{cases}$$

All other unit Perron–Frobenius eigenvectors coincide with this one at all continuity points.

Now, let $\mathbf{u} \in \mathcal{C}_{\text{per}}^2(\mathbb{R}^n, [\mathbf{0}, \infty))$ and $\lambda \leq -1$ such that $-\Delta\mathbf{u} - \mathbf{L}\mathbf{u} \leq \lambda\mathbf{u}$. Taking the scalar product in \mathbb{R}^2 with \mathbf{u} and integrating by parts in $[0, L]$, we obtain:

$$0 \leq \int_{[0, L]} \sum_{i=1}^2 |\nabla u_i|^2 \leq \int_{[0, L]} \mathbf{u}^T (\mathbf{L} - \mathbf{I}) \mathbf{u}.$$

Since, at all $x \in [0, L]$, $\mathbf{L}(x) - \mathbf{I}$ is a symmetric matrix with nonpositive eigenvalues, $\mathbf{u}(x)^T (\mathbf{L}(x) - \mathbf{I}) \mathbf{u}(x) \leq 0$. Therefore all inequalities above are actually equalities, and in particular \mathbf{u} is a constant vector satisfying $\mathbf{u}^T (\mathbf{L}(x) - \mathbf{I}) \mathbf{u} = 0$. Since no Perron–Frobenius eigenvector is constant, necessarily $\mathbf{u} = \mathbf{0}$.

Therefore no $\lambda \leq -1$ can satisfy $-\Delta\mathbf{u} - \mathbf{L}\mathbf{u} = \lambda\mathbf{u}$ for some positive periodic \mathcal{C}^2 eigenfunction \mathbf{u} , and this fact, together with the reduction (9), directly implies that $\lambda'_1 > -1$.

Independently of this observation, Theorem 1.13 implies that $\lambda'_1 = R$. Therefore we have indeed proved that, for this counter-example,

$$-R < \frac{1}{|[0, L]|} \int_{[0, L]} \lambda_{\text{PF}}(\mathbf{L}).$$

3.3.3. Upper estimates: proof of Theorems 1.14, 1.15, 1.16. In a similar spirit, the following property requires a line-sum-symmetry assumption ($\mathbf{L}\mathbf{1} = \mathbf{L}^T\mathbf{1}$) and uses the property described in Eaves–Hoffman–Rothblum–Schneider [29, Corollary 3].

Proposition 3.26. *Let $z \in \mathbb{R}^n$. Assume:*

- (1) for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) = 0$,
- (2) $\mathbf{L}(t, x)$ is line-sum-symmetric for all $(t, x) \in \overline{\Omega}_{\text{per}}$.

Then

$$\lambda_{1,z} \leq -\frac{1}{N} \left(\sum_{i,j=1}^N \langle \hat{l}_{i,j} \rangle + z \cdot \sum_{i=1}^N \left(\langle \hat{A}_i \rangle z - \langle \hat{q}_i \rangle \right) \right)$$

and this inequality is an equality if $\mathbf{L} + \text{diag}(\nabla \cdot (A_i z) + z \cdot (A_i z - q_i))$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$ with Perron–Frobenius eigenvector $\mathbf{1}$ and constant Perron–Frobenius eigenvalue.

Proof. Denote, for all $i \in [N]$, $q_i - 2A_i z = b_i$ and recall

$$\begin{aligned} \mathcal{Q}_z &= \mathcal{Q} - \text{diag}(2A_i z \cdot \nabla + z \cdot A_i z + \nabla \cdot (A_i z) - q_i \cdot z) \\ &= \text{diag}(\partial_t - \nabla \cdot (A_i \nabla) + b_i \cdot \nabla - \nabla \cdot (A_i z) - z \cdot (A_i z - q_i)) - \mathbf{L} \end{aligned}$$

Denote $\mathbf{u} = \mathbf{u}_z$ the unit periodic principal eigenfunction associated with $\lambda_{1,z}$. Taking the scalar product in \mathbb{R}^N between $(1/u_i)_{i \in [N]}$ and $\mathcal{Q}_z \mathbf{u} - \lambda_{1,z} \mathbf{u}$, integrating by parts in $\overline{\Omega_{\text{per}}}$, using the fact that all b_i are divergence-free and using

$$(21) \quad \sum_{i=1}^N \frac{(\mathbf{L}\mathbf{u})_i}{u_i} \geq \sum_{i,j=1}^N l_{i,j} \quad \text{and} \quad \int_{\overline{\Omega_{\text{per}}}} \frac{\nabla u_i}{u_i} \cdot A_i \frac{\nabla u_i}{u_i} \geq 0,$$

we get

$$\lambda_{1,z} \leq -\frac{1}{NT|[0, L]|} \left(\sum_{i,j=1}^N \int_{\overline{\Omega_{\text{per}}}} l_{i,j} + z \cdot \sum_{i=1}^N \int_{\overline{\Omega_{\text{per}}}} (A_i z - q_i) \right).$$

From the equality case in (21), we deduce that this inequality is an equality if $\mathbf{u}_z \in \text{span}(\mathbf{1})$ and $\mathbf{L}(t, x)$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$. These conditions are satisfied if and only if $\mathbf{L} + \text{diag}(\nabla \cdot (A_i z) + z \cdot (A_i z - q_i))$ is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$ with Perron–Frobenius eigenvector $\mathbf{1}$ and Perron–Frobenius eigenvalue $\lambda_{1,z}$, both constant. Finally, by uniqueness of the periodic principal eigenvalue, the assumption that the Perron–Frobenius eigenvalue is $\lambda_{1,z}$ can be replaced without loss of generality by the assumption that the Perron–Frobenius eigenvalue is constant. \square

Remark 3.21. Circulant matrices and doubly stochastic matrices are line-sum-symmetric and always admit $\mathbf{1}$ as eigenvector. Hence all inequalities on $\lambda_{1,z}$ are equalities if:

- (1) all A_i are constant and coincide and all q_i are constant and coincide,
- (2) \mathbf{L} is, at all (t, x) , irreducible and either circulant or doubly stochastic,
- (3) its Perron–Frobenius eigenvalue $\lambda_{\text{PF}}(\mathbf{L}(t, x)) = \sum_{j=1}^N l_{1,j}(t, x)$ is constant (this condition being automatically satisfied in the doubly stochastic case).

This shows in particular that the inequalities can all be equalities even if \mathbf{L} is not spatio-temporally constant.

The following two corollaries are concerned with special cases.

Corollary 3.27. *Assume:*

- (1) for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot q_i = 0$,
- (2) $\mathbf{L}(t, x)$ is line-sum-symmetric for all $(t, x) \in \overline{\Omega_{\text{per}}}$.

Then

$$\lambda'_1 \leq -\frac{1}{N} \sum_{i,j=1}^N \langle \hat{l}_{i,j} \rangle = -\frac{1}{N} \mathbf{1}^T \langle \hat{\mathbf{L}} \rangle \mathbf{1}$$

and this inequality is an equality if \mathbf{L} is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$ with Perron–Frobenius eigenvector $\mathbf{1}$ and constant Perron–Frobenius eigenvalue.

Corollary 3.28. *Assume:*

- (1) for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and q_i and each column of A_i are divergence-free,
- (2) $\mathbf{L}(t, x)$ is line-sum-symmetric for all $(t, x) \in \overline{\Omega_{\text{per}}}$.

Denote

$$[A] = \frac{1}{N} \sum_{i=1}^N \langle \hat{A}_i \rangle, \quad [q] = \frac{1}{N} \sum_{i=1}^N \langle \hat{q}_i \rangle.$$

Then

$$\lambda_1 \leq -\frac{1}{N} \mathbf{1}^T \langle \hat{\mathbf{L}} \rangle \mathbf{1} + \frac{1}{4} [q] \cdot [A][q]$$

with equality if

$$\mathbf{L} + \frac{1}{2} \text{diag} \left(\nabla \cdot (A_i [A]^{-1} [q] + [A]^{-1} [q] \cdot \left(\frac{1}{2} A_i [A]^{-1} [q] - q_i \right)) \right)$$

is irreducible at all $(t, x) \in \overline{\Omega_{\text{per}}}$ with Perron–Frobenius eigenvector $\mathbf{1}$ and constant Perron–Frobenius eigenvalue.

Proof. The assumption that all q_i and all columns of all A_i are divergence-free implies that, for all $i \in [N]$ and $z \in \mathbb{R}^n$, $q_i - 2A_i z$ is divergence-free (and actually the converse implication is also true: consider the special cases $z = 0, e_1, \dots, e_n$, where $e_\alpha = (\delta_{\alpha\beta})_{\beta \in [n]}$).

By (A_1) , $[A]$ is invertible, so that the inequality of Proposition 3.26 reads

$$\lambda_{1, z + \frac{1}{2} [A]^{-1} [q]} \leq -\frac{1}{N} \mathbf{1}^T \langle \hat{\mathbf{L}} \rangle \mathbf{1} - z \cdot [A]z + \frac{1}{4} [q] \cdot [A][q].$$

The inequality on $\lambda_1 = \max_{z \in \mathbb{R}^n} \lambda_{1, z}$ and the associated equality case follow directly. \square

Now we turn to the proof of Theorem 1.15.

Proposition 3.29. *Let $z \in \mathbb{R}^n$. If, for all $i \in [N]$, $q_i \in \mathcal{C}_{\text{per}}^1(\mathbb{R}^n, \mathbb{R}^n)$ and $\nabla \cdot (q_i - 2A_i z) = 0$, then*

$$\lambda_{1, z} \leq \lambda_{1, \text{per}} \left(\partial_t - \mathbf{L}^\# - \text{diag} (z \cdot (\langle A_i \rangle z - \langle q_i \rangle)) \right),$$

where the entries of the matrix $\mathbf{L}^\# = (l_{i,j}^\#)_{(i,j) \in [N]^2}$ are defined by:

$$l_{i,j}^\# = \begin{cases} \frac{1}{|[0,L]|} \int_{[0,L]} l_{i,i} & \text{if } i = j, \\ \exp \left(\frac{1}{|[0,L]|} \int_{[0,L]} \ln l_{i,j} \right) & \text{if } i \neq j \text{ and } \min_{(t,x) \in \overline{\Omega_{\text{per}}}} l_{i,j}(t,x) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The proof is quite similar to that of Proposition 3.26, we only sketch it.

Multiply each line of $\mathcal{Q}_z \mathbf{u} - \lambda_{1,z} \mathbf{u}$ by $1/u_i$, integrate by parts in $[0, L]$, divide by $|[0, L]|$, define $J_i = \{j \in [N] \setminus \{i\} \mid \min_{\overline{\Omega}_{\text{per}}} l_{i,j} > 0\}$, use the Jensen inequality:

$$\begin{aligned} \frac{1}{|[0, L]|} \int_{[0, L]} \sum_{j \in [N] \setminus \{i\}} \frac{l_{i,j} u_j}{u_i} &\geq \frac{1}{|[0, L]|} \int_{[0, L]} \sum_{j \in J_i} e^{-\ln(u_i) + \ln(l_{i,j}) + \ln(u_j)} \\ &\geq \sum_{j \in J_i} e^{\frac{1}{|[0, L]|} \int_{[0, L]} (-\ln(u_i) + \ln(l_{i,j}) + \ln(u_j))} \\ &= \sum_{j \in J_i} \frac{e^{\frac{1}{|[0, L]|} \int_{[0, L]} \ln(l_{i,j})} e^{\frac{1}{|[0, L]|} \int_{[0, L]} \ln(u_j)}}{e^{\frac{1}{|[0, L]|} \int_{[0, L]} \ln(u_i)}}, \end{aligned}$$

define the positive function $\mathbf{v} : t \mapsto \left(\exp \left(\frac{1}{|[0, L]|} \int_{[0, L]} \ln(u_i(t, x)) dx \right) \right)_{i \in [N]}$, find

$$\mathbf{v}' - \left[\mathbf{L}^\# + \text{diag} \left(z \cdot \frac{1}{|[0, L]|} \int_{[0, L]} (A_i z - q_i) \right) \right] \mathbf{v} \geq \lambda_{1,z} \mathbf{v},$$

and subsequently use the min–max formula for the periodic principal eigenvalue. The result follows. \square

Repeating the exact same procedure but this time with averages in $[0, T]$, we also find the estimate of Theorem 1.16, recalled below.

Proposition 3.30. *Let $z \in \mathbb{R}^n$. If $(A_i)_{i \in [N]}$, $(q_i)_{i \in [N]}$ and \mathbf{L} do not depend on x , then*

$$\lambda_{1,z} \leq -\lambda_{\text{PF}} \left(\mathbf{L}^b + \text{diag} \left(z \cdot \hat{A}_i z - \hat{q}_i \cdot z \right) \right),$$

where the entries of the matrix $\mathbf{L}^b = (l_{i,j}^b)_{(i,j) \in [N]^2}$ are defined by:

$$l_{i,j}^b = \begin{cases} \frac{1}{T} \int_0^T l_{i,i} & \text{if } i = j, \\ \exp \left(\frac{1}{T} \int_0^T \ln l_{i,j} \right) & \text{if } i \neq j \text{ and } \min_{t \in [0, T]} l_{i,j}(t) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

The estimate of Corollary 1.17 follows directly.

3.4. Optimization: proof of Theorems 1.18–1.20.

3.4.1. *Optimization of the normalized mutation matrix: proof of Theorem 1.18.* In this section we prove Theorem 1.18.

We recall that a doubly stochastic matrix $\mathbf{S} \in \mathbb{R}^{N \times N}$ is a nonnegative matrix such that $\mathbf{S} \mathbf{1} = \mathbf{S}^T \mathbf{1} = \mathbf{1}$. Denote $\mathcal{S} \subset \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ the set of all periodic functions whose values are doubly stochastic matrices almost everywhere and $\mathcal{S}_{\{0,1\}}$ the subset of all functions valued almost everywhere in the set of permutation matrices.

Although we assumed until now that the zeroth order term \mathbf{L} of \mathcal{Q} is Hölder-continuous, it can be verified that the family $(\lambda_{1,z})_{z \in \mathbb{R}^n}$ can still be defined if \mathbf{L} has only an \mathcal{L}^∞ regularity, using for instance a standard regularization procedure not detailed here.

We begin with the following decomposition lemma.

Lemma 3.31. *Let $\mathbf{A} \in \mathbb{R}^{N \times N}$ be a non-diagonal, essentially nonnegative matrix and $\mathbf{r} = \mathbf{A}^T \mathbf{1}$.*

Assume $\mathbf{A} - \text{diag}(\mathbf{r})$ admits a positive Perron–Frobenius eigenvector $\mathbf{v} \in (\mathbf{0}, \infty)$ and let $\boldsymbol{\nu} = (1/v_i)_{i \in [N]}$.

Then there exists $m_0 > 0$ such that, for any $m \in (0, m_0]$, the matrix $\mathbf{S} = m(\mathbf{A} - \text{diag}(\mathbf{r})) \text{diag}(\boldsymbol{\nu})^{-1} + \mathbf{I}$ is doubly stochastic.

In other words, \mathbf{A} can be decomposed as $\mathbf{A} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$, with $\mathbf{r} \in \mathbb{R}^N$, $\mathbf{S} \in [0, 1]^{N \times N}$ doubly stochastic, $\boldsymbol{\mu} \in (\mathbf{0}, \infty)$.

Proof. By definition of \mathbf{r} , the matrix $\mathbf{A} - \text{diag}(\mathbf{r})$ admits $\mathbf{1}$ as left Perron–Frobenius eigenvector with Perron–Frobenius eigenvalue 0. Hence

$$\lambda_{\text{PF}}(\mathbf{A} - \text{diag}(\mathbf{r})) = \lambda_{\text{PF}}((\mathbf{A} - \text{diag}(\mathbf{r}))^T) = 0$$

and \mathbf{v} satisfies $(\mathbf{A} - \text{diag}(\mathbf{r}))\mathbf{v} = \mathbf{0}$.

For any $m > 0$, the essentially nonnegative matrix

$$\mathbf{S} = m(\mathbf{A} - \text{diag}(\mathbf{r})) \text{diag}(\boldsymbol{\nu})^{-1} + \mathbf{I}$$

satisfies:

$$\mathbf{1}^T \mathbf{S} = m \mathbf{1}^T [(\mathbf{A} - \text{diag}(\mathbf{r})) \text{diag}(\boldsymbol{\nu})^{-1} + \mathbf{I}] = \mathbf{0}^T \text{diag}(\boldsymbol{\nu})^{-1} + \mathbf{1}^T = \mathbf{1}^T,$$

$$\mathbf{S} \mathbf{1} = [m(\mathbf{A} - \text{diag}(\mathbf{r})) \text{diag}(\boldsymbol{\nu})^{-1} + \mathbf{I}] \mathbf{1} = m(\mathbf{A} - \text{diag}(\mathbf{r}))\mathbf{v} + \mathbf{1} = \mathbf{1},$$

i.e. $\mathbf{1}$ is a left and right Perron–Frobenius eigenvector with Perron–Frobenius eigenvalue 1.

Note that, by definition of \mathbf{r} , $r_i - a_{i,i} \geq 0$ for each $i \in [N]$. Since \mathbf{A} is non-diagonal, $\max_{i \in [N]} (r_i - a_{i,i}) v_i > 0$ and $\max_{i,j \in [N], i \neq j} a_{i,j} v_j > 0$. Therefore, for \mathbf{S} to be doubly stochastic, it only remains to choose $m \in (0, m_0]$ with m_0 defined as

$$m_0 = \min \left[\frac{1}{\max_{i,j \in [N], i \neq j} a_{i,j} v_j}, \frac{1}{\max_{i \in [N]} (r_i - a_{i,i}) v_i} \right].$$

□

Remark 3.22. It can be easily verified that any other decomposition $\mathbf{A} = \text{diag}(\tilde{\mathbf{r}}) + (\tilde{\mathbf{S}} - \mathbf{I}) \text{diag}(\tilde{\boldsymbol{\mu}})$ with $\tilde{\mathbf{r}} \in \mathbb{R}^N$, $\tilde{\mathbf{S}} \in [0, 1]^{N \times N}$ doubly stochastic, $\tilde{\boldsymbol{\mu}} \in (\mathbf{0}, \infty)$ satisfies, for some $m > 0$, $\mathbf{r} = \tilde{\mathbf{r}}$, $\boldsymbol{\mu} = \frac{1}{m} \tilde{\boldsymbol{\mu}}$ and $\mathbf{S} = \mathbf{I} + m(\tilde{\mathbf{S}} - \mathbf{I})$.

Remark 3.23. When such a decomposition $\mathbf{A} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ exists, with $\mathbf{r} \in \mathbb{R}^N$, $\mathbf{S} \in [0, 1]^{N \times N}$ doubly stochastic and $\boldsymbol{\mu} \in [0, \infty)$, the positivity of all μ_i is equivalent with the positivity of the Perron–Frobenius eigenvector of $\mathbf{A} - \text{diag}(\mathbf{r})$. On the contrary, when \mathbf{A} is reducible and $\mathbf{A} - \text{diag}(\mathbf{r})$ has Perron–Frobenius eigenvectors only in $\partial(\mathbf{0}, \infty)$, the existence of such a decomposition can be both a true or a false statement, as shown by the following two examples:

$$\mathbf{A} - \text{diag}(\mathbf{r}) = \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & -2 \end{pmatrix} = \left(\begin{pmatrix} 0 & 1/2 & 1/2 \\ 0 & 1/2 & 1/2 \\ 1 & 0 & 0 \end{pmatrix} - \mathbf{I} \right) \text{diag}(0, 2, 2)$$

$$\mathbf{A} - \text{diag}(\mathbf{r}) = \begin{pmatrix} -1 & 1 & 1/3 \\ 1 & -1 & 2/3 \\ 0 & 0 & -1 \end{pmatrix}.$$

Proposition 3.32. *Assume \mathbf{L} has the form*

$$\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$$

with $\mathbf{S} \in \mathcal{S}$, $\mathbf{r} \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^N)$ and $\boldsymbol{\mu} \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, [0, \infty))$.

Then, for all $z \in \mathbb{R}^n$,

$$\inf_{\mathbf{S} \in \mathcal{S}_{\{0,1\}}} \lambda_{1,z}(\mathbf{S}) = \inf_{\mathbf{S} \in \mathcal{S}} \lambda_{1,z}(\mathbf{S}) \leq \sup_{\mathbf{S} \in \mathcal{S}} \lambda_{1,z}(\mathbf{S}) = \inf_{\mathbf{S} \in \mathcal{S}_{\{0,1\}}} \lambda_{1,z}(\mathbf{S}).$$

Furthermore, all inf and max above are actually min and max respectively.

Proof. It suffices to prove that there exists an element of $\mathcal{S}_{\{0,1\}}$ that minimizes $\mathbf{S} \in \mathcal{S} \mapsto \lambda_{1,z}$, the property on the maximum being proved similarly. Also, it is sufficient to prove only the case $z = 0$.

Step 1: exhibiting a minimizer in \mathcal{S} . The closed and bounded set

$$\mathcal{S} = \left\{ \mathbf{S} \in \mathcal{L}^\infty(\Omega_{\text{per}}, \mathbb{R}^{N \times N}) \mid \mathbf{S} \geq \mathbf{0}, \mathbf{S}\mathbf{1} = \mathbf{S}^T\mathbf{1} = \mathbf{1} \text{ a.e.} \right\}$$

is, by virtue of the Banach–Alaoglu theorem, compact in the weak- \star topology of $\mathcal{L}^\infty(\Omega_{\text{per}}) = (\mathcal{L}^1(\Omega_{\text{per}}))'$. Hence a minimizing sequence $(\mathbf{S}_k)_{k \in \mathbb{N}}$ converges, up to extraction, to a weak- \star limit $\mathbf{S}_\infty \in \mathcal{L}^\infty(\Omega_{\text{per}}, \mathbb{R}^{N \times N})$. Extending periodically $\mathbf{S}_\infty \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$, it only remains to verify $\mathbf{S} \in \mathcal{S}$ and $\lambda'_1(\mathbf{S}_\infty) = \lim_{k \rightarrow +\infty} \lambda'_1(\mathbf{S}_k)$.

The nonnegativity of \mathbf{S}_∞ in the sense of linear forms is immediate, testing the convergence against arbitrary nonnegative functions in $\mathcal{L}^1(\Omega_{\text{per}})$. Subsequently, testing for all $(i, j) \in [N]^2$ against \mathbf{e}_j multiplied by the indicator of $\{s_{\infty, i, j} < 0\} \cap \overline{\Omega_{\text{per}}}$, we deduce the nonnegativity almost everywhere. Testing for any $(t_0, x_0) \in \overline{\Omega_{\text{per}}}$, $\rho > 0$, against $\mathbf{1}$ multiplied by the indicator of $B((t_0, x_0), \rho)$ and divided by $|B((t_0, x_0), \rho)|$, we deduce by virtue of the Lebesgue differentiation theorem $\mathbf{S}_\infty \mathbf{1} = \mathbf{1}$ almost everywhere. Next, testing the convergence against all $(\mathbf{e}_j)_{j \in [N]}$, we find that all entries $s_{k, i, j}$ converge in the weak- \star topology of $\mathcal{L}^\infty(\Omega_{\text{per}}, \mathbb{R})$, so that $(\mathbf{S}_k^T)_{k \in \mathbb{N}}$ also converges. Similarly, $\mathbf{S}_\infty^T \mathbf{1} = \mathbf{1}$ almost everywhere. Therefore $\mathbf{S}_\infty \in \mathcal{S}$ indeed.

By continuity of λ'_1 with respect to the weak- \star topology of $\mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ (see Theorem 1.5 and Remark 3.7), $\lim_{k \rightarrow +\infty} \lambda'_1(\mathbf{S}_k) = \lambda'_1(\mathbf{S}_\infty)$. This ends this step of the proof. \square

Define the function:

$$\begin{aligned} \Phi : \mathcal{S} \times \mathbb{R} \times \mathbb{R}^n &\rightarrow [N] \\ (\mathbf{S}, t, x) &\mapsto \#\{(i, j) \in [N]^2 \mid s_{i, j}(t, x) = 1\} \end{aligned}$$

and remark that, for any $\mathbf{S} \in \mathcal{S}$,

$$\mathbf{S} \in \mathcal{S}_{\{0,1\}} \iff |\{(t, x) \in \overline{\Omega_{\text{per}}} \mid \Phi(\mathbf{S}, t, x) = N\}| = |\overline{\Omega_{\text{per}}}| = T|[0, L]|.$$

In other words, defining $\Omega_{\text{per}}(\mathbf{S}, N_0) = \{(t, x) \in \overline{\Omega_{\text{per}}} \mid \Phi(\mathbf{S}, t, x) = N_0\}$,

$$\mathbf{S} \in \mathcal{S}_{\{0,1\}} \iff \forall N_0 \in [N-1] \quad |\Omega_{\text{per}}(\mathbf{S}, N_0)| = 0.$$

Let $\mathbf{S}^\wedge \in \mathcal{S}$ be a minimizer. Assume $\mathbf{S}^\wedge \notin \mathcal{S}_{\{0,1\}}$. This means that there exists $N_0 \in [N-1]$ such that $\Omega_{N_0} = \Omega_{\text{per}}(\mathbf{S}^\wedge, N_0)$ has a positive measure. We are now going to correct this minimizer step by step.

In what follows, we first consider the case where \mathbf{S}^\wedge is irreducible on average in Ω_{per} . Of course this is not necessarily the case, and we will generalize afterward. In order to ease the reading, we denote $\mathbf{B} = \text{diag}(\mathbf{r}) - \text{diag}(\boldsymbol{\mu})$.

Step 2: when \mathbf{S}^\wedge is irreducible on average, correcting the minimizer in a large subset of Ω_{N_0} .

Let $\mathbf{u}, \mathbf{v} \in \mathcal{C}_{\text{per}}^{1,2}(\mathbb{R} \times \mathbb{R}^n, (\mathbf{0}, \infty))$ be respectively a periodic principal eigenfunction of \mathcal{Q} and a periodic principal eigenfunction of the adjoint operator

$$\mathcal{Q}^* = -\partial_t - \text{diag}(\nabla \cdot (A_i \nabla) + q_i \cdot \nabla + \nabla \cdot q_i) - \mathbf{B}^\text{T} - \text{diag}(\boldsymbol{\mu})(\mathbf{S}^\wedge)^\text{T}.$$

By full coupling of the operator \mathcal{Q} , \mathbf{u} and \mathbf{v} are positive. With the normalizations $\int_{\Omega_{\text{per}}} |\mathbf{u}|^2 = \int_{\Omega_{\text{per}}} \mathbf{v}^\text{T} \mathbf{u} = 1$ (the second one is possible because, by positivity, \mathbf{u} and \mathbf{v} cannot be orthogonal), \mathbf{u} and \mathbf{v} are uniquely defined.

Let $(t, x) \in \Omega_{N_0}$. Following exactly the construction of Neumann–Sze [62], there exist two permutation matrices $\mathbf{P}(t, x) \in \mathcal{S}_{\{0,1\}}$ and $\mathbf{Q}(t, x) \in \mathcal{S}_{\{0,1\}}$ such that:

- (1) the matrix $\tilde{\mathbf{S}}^\wedge(t, x) = \mathbf{P}(t, x) \mathbf{S}^\wedge(t, x) \mathbf{Q}(t, x)^\text{T}$ is doubly stochastic and has a block diagonal form:

$$\begin{pmatrix} \tilde{\mathbf{S}}_{\text{top}}^\wedge(t, x) & \mathbf{0} \\ \mathbf{0} & \tilde{\mathbf{S}}_{\text{bottom}}^\wedge(t, x) \end{pmatrix}$$

with $\tilde{\mathbf{S}}_{\text{bottom}}^\wedge(t, x) \in \{0, 1\}^{N_0 \times N_0}$ empty (if $N_0 = 0$) or a permutation matrix (if $N_0 > 0$) and all entries in $\tilde{\mathbf{S}}_{\text{top}}^\wedge(t, x)$ smaller than 1;

- (2) for all $i \in [N - N_0]$,

$$\mathbf{e}_i^\text{T} \mathbf{P}(t, x) \mathbf{v}(t, x) \leq \mathbf{e}_1^\text{T} \mathbf{P}(t, x) \mathbf{v}(t, x);$$

- (3) for all $j \in [N - N_0]$,

$$\mathbf{e}_j^\text{T} \mathbf{Q}(t, x) \text{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x) \leq \mathbf{e}_1^\text{T} \mathbf{Q}(t, x) \text{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x).$$

The three properties together imply that,

$$(22) \quad (\tilde{s}_{i,1}^\wedge(t, x))_{i \in [N]}^\text{T} \mathbf{P}(t, x) \mathbf{v}(t, x) \leq \mathbf{e}_1^\text{T} \mathbf{P}(t, x) \mathbf{v}(t, x),$$

$$(23) \quad (\tilde{s}_{1,j}^\wedge(t, x))_{j \in [N]}^\text{T} \mathbf{Q}(t, x) \text{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x) \leq \mathbf{e}_1^\text{T} \mathbf{Q}(t, x) \text{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x).$$

Next, define, for the same $(t, x) \in \Omega_{N_0}$,

$$\mathbf{a}(t, x) = -\mathbf{e}_1 + (\tilde{s}_{i,1}^\wedge(t, x))_{i \in [N]} = \begin{pmatrix} -(1 - \tilde{s}_{1,1}^\wedge(t, x)) \\ \tilde{s}_{2,1}^\wedge(t, x) \\ \vdots \\ \tilde{s}_{N-N_0,1}^\wedge(t, x) \\ 0 \\ \vdots \\ 0 \end{pmatrix},$$

$$\mathbf{b}(t, x) = -\mathbf{e}_1 + (\tilde{s}_{1,j}^\wedge(t, x))_{j \in [N]} = \begin{pmatrix} -(1 - \tilde{s}_{1,1}^\wedge(t, x)) \\ \tilde{s}_{1,2}^\wedge(t, x) \\ \vdots \\ \tilde{s}_{1,N-N_0}^\wedge(t, x) \\ 0 \\ \vdots \\ 0 \end{pmatrix},$$

$$\mathbf{T}^\wedge(t, x) = \frac{1}{1 - \tilde{s}_{1,1}^\wedge(t, x)} \mathbf{a}(t, x) \mathbf{b}(t, x)^\text{T}.$$

Let us verify that $\tilde{\mathbf{S}}^\wedge + \mathbf{T}^\wedge$ is doubly stochastic at (t, x) . Since $\tilde{\mathbf{S}}^\wedge(t, x)$ is doubly stochastic, we only have to verify that $\tilde{\mathbf{S}}^\wedge(t, x) + \mathbf{T}^\wedge(t, x)$ is nonnegative and that $\mathbf{a}(t, x)\mathbf{b}(t, x)^\top \mathbf{1} = \mathbf{b}(t, x)\mathbf{a}(t, x)^\top \mathbf{1} = \mathbf{0}$. Both properties turn out to be obvious. By construction, $\Phi(\tilde{\mathbf{S}}^\wedge + \mathbf{T}^\wedge, t, x) = N_0 + 1$. Indeed, $\tilde{\mathbf{S}}^\wedge(t, x) + \mathbf{T}^\wedge(t, x)$ has N_0 entries equal to 1 in its bottom right block and its upper left entry satisfies

$$\tilde{s}_{1,1}^\wedge(t, x) + t_{1,1}^\wedge(t, x) = \tilde{s}_{1,1}^\wedge(t, x) + \frac{1}{1 - \tilde{s}_{1,1}^\wedge(t, x)}(1 - \tilde{s}_{1,1}^\wedge(t, x))^2 = 1.$$

Let $\omega \subset \Omega_{N_0}$ be a measurable subset. Setting

$$\mathbf{T}_\omega^\wedge : (t, x) \in \overline{\Omega_{\text{per}}} \mapsto \begin{cases} \mathbf{T}^\wedge(t, x) & \text{if } (t, x) \in \omega, \\ \mathbf{0} & \text{if } (t, x) \in \overline{\Omega_{\text{per}}} \setminus \omega, \end{cases}$$

extending \mathbf{T}_ω^\wedge periodically in $\mathbb{R} \times \mathbb{R}^n$ and verifying routinely that \mathbf{T}_ω^\wedge is measurable, we are now in a position to verify that this construction does not modify the periodic principal eigenvalue λ_1' , namely $\tilde{\mathbf{S}}^\wedge + \mathbf{T}_\omega^\wedge \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ is still a minimizer, provided ω is appropriately chosen.

Denote, for any $\alpha \in [0, 1]$,

$$\mathcal{Q}_\alpha = \text{diag}(\mathcal{P}_i) - \mathbf{B} - \mathbf{P}^\top \left(\alpha \mathbf{T}_\omega^\wedge + \tilde{\mathbf{S}}^\wedge \right) \mathbf{Q} \text{diag}(\boldsymbol{\mu}),$$

$\lambda(\alpha) = \lambda_1'(\mathcal{Q}_\alpha)$, and let \mathbf{u}_α and \mathbf{v}_α be two positive periodic principal eigenfunctions of respectively \mathcal{Q}_α and of the adjoint operator

$$\mathcal{Q}_\alpha^* = -\partial_t - \text{diag}(\nabla \cdot (A_i \nabla) + q_i \cdot \nabla + \nabla \cdot q_i) - \mathbf{B}^\top - \text{diag}(\boldsymbol{\mu}) \mathbf{Q}^\top \left(\alpha \mathbf{T}_\omega^\wedge + \tilde{\mathbf{S}}^\wedge \right)^\top \mathbf{P},$$

normalized so that $\int_{\overline{\Omega_{\text{per}}}} |\mathbf{u}_\alpha|^2 = \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\alpha^\top \mathbf{u}_\alpha = 1$. For any $\alpha, \beta \in [0, 1]$, $\alpha \neq \beta$,

$$\begin{aligned} \frac{\lambda(\beta) - \lambda(\alpha)}{\beta - \alpha} &= \frac{\lambda(\beta) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha - \lambda(\alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha}{(\beta - \alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \\ &= \frac{1}{(\beta - \alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \left(\int_{\overline{\Omega_{\text{per}}}} \mathbf{u}_\alpha^\top (\mathcal{Q}_\beta^* \mathbf{v}_\beta) - \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top (\mathcal{Q}_\alpha \mathbf{u}_\alpha) \right) \\ &= \frac{1}{(\beta - \alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \left(\int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top (\mathcal{Q}_\beta \mathbf{u}_\alpha) - \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top (\mathcal{Q}_\alpha \mathbf{u}_\alpha) \right) \\ &= \frac{1}{(\beta - \alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \left(\int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top ((\mathcal{Q}_\beta - \mathcal{Q}_\alpha) \mathbf{u}_\alpha) \right) \\ &= \frac{1}{(\beta - \alpha) \int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \left(\int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top (((\beta - \alpha) \mathcal{Q}_1 - (\beta - \alpha) \mathcal{Q}_0) \mathbf{u}_\alpha) \right) \\ &= -\frac{1}{\int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \left(\int_{\overline{\Omega_{\text{per}}}} (\mathbf{P} \mathbf{v}_\beta)^\top \mathbf{T}_\omega^\wedge \mathbf{Q} \text{diag}(\boldsymbol{\mu}) \mathbf{u}_\alpha \right) \\ &= -\frac{1}{\int_{\overline{\Omega_{\text{per}}}} \mathbf{v}_\beta^\top \mathbf{u}_\alpha} \int_\omega \frac{(\mathbf{a}^\top \mathbf{P} \mathbf{v}_\beta)(\mathbf{b}^\top \mathbf{Q} \text{diag}(\boldsymbol{\mu}) \mathbf{u}_\alpha)}{1 - \tilde{s}_{1,1}^\wedge}. \end{aligned}$$

Taking the limit $\beta \rightarrow \alpha$, this leads to

$$\lambda'(\alpha) = - \int_{\omega} \frac{(\mathbf{a}^T \mathbf{P} \mathbf{v}_{\alpha})(\mathbf{b}^T \mathbf{Q} \operatorname{diag}(\boldsymbol{\mu}) \mathbf{u}_{\alpha})}{1 - \widehat{s}_{1,1}^{\alpha}}.$$

In view of this equality and of (22)–(23), $\lambda'(0) \leq 0$. We claim that in fact $\lambda'(0) = 0$. Indeed, if this is not the case, then there exists a small $\alpha > 0$ such that $\lambda(\alpha) < \lambda(0)$. Then the minimality of $\lambda(0) = \lambda'_1(\mathbf{P}^T \widehat{\mathbf{S}}^{\wedge} \mathbf{Q}) = \lambda'_1(\mathbf{S}^{\wedge})$ in \mathcal{S} is contradicted⁹.

Since $\lambda'(0) = 0$ for any choice of ω , using the Lebesgue differentiation theorem, we obtain:

$$-\frac{(\mathbf{a}^T \mathbf{P} \mathbf{v}) \left(\mathbf{b}^T \mathbf{Q} \operatorname{diag}(\boldsymbol{\mu}) \mathbf{u} \right)}{1 - \widehat{s}_{1,1}^{\alpha}} = 0 \quad \text{almost everywhere in } \Omega_{N_0}.$$

Subsequently, for almost every $(t, x) \in \Omega_{N_0}$,

$$\mathbf{T}_{\Omega_{N_0}}^{\wedge}(t, x) \mathbf{Q}(t, x) \operatorname{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x) = \mathbf{0} \quad \text{or} \quad \mathbf{T}_{\Omega_{N_0}}^{\wedge}(t, x)^T \mathbf{P}(t, x) \mathbf{v}(t, x) = \mathbf{0}.$$

Let

$$\begin{aligned} \omega_{\mathbf{u}} &= \left\{ (t, x) \in \Omega_{N_0} \mid \mathbf{T}_{\Omega_{N_0}}^{\wedge}(t, x) \mathbf{Q}(t, x) \operatorname{diag}(\boldsymbol{\mu}(t, x)) \mathbf{u}(t, x) = \mathbf{0} \right\}, \\ \omega_{\mathbf{v}} &= \left\{ (t, x) \in \Omega_{N_0} \mid \mathbf{T}_{\Omega_{N_0}}^{\wedge}(t, x)^T \mathbf{P}(t, x) \mathbf{v}(t, x) = \mathbf{0} \right\}. \end{aligned}$$

The subsets $\omega_{\mathbf{u}}$ and $\omega_{\mathbf{v}} \setminus \omega_{\mathbf{u}}$ are measurable, disjoint and satisfy $|\omega_{\mathbf{u}} \cup (\omega_{\mathbf{v}} \setminus \omega_{\mathbf{u}})| = |\Omega_{N_0}|$. One of the two, denoted below ω , satisfies $|\omega| \geq \frac{1}{2} |\Omega_{N_0}|$. Choosing this ω in the definition of $\mathbf{T}_{\omega}^{\wedge}$, we deduce directly that the corresponding eigenvector at $\alpha = 0$ (\mathbf{u} if $\omega = \omega_{\mathbf{u}}$, \mathbf{v} if $\omega = \omega_{\mathbf{v}}$) remains a periodic principal eigenvector for any $\alpha \in [0, 1]$, with in addition $\lambda(\alpha) = \lambda(0)$ for any $\alpha \in [0, 1]$. In particular,

$$\lambda'_1(\mathbf{S}^{\wedge}) = \lambda(0) = \lambda(1) = \lambda'_1(\mathbf{S}^{\wedge} + \mathbf{P}^T \mathbf{T}_{\omega}^{\wedge} \mathbf{Q}).$$

□

Step 3: when \mathbf{S}^{\wedge} is irreducible on average, correcting the minimizer in Ω_{N_0} almost everywhere.

Let $\mathbf{S}_1^{\wedge} = \mathbf{S}^{\wedge} + \mathbf{P}^T \mathbf{T}_{\omega}^{\wedge} \mathbf{Q}$ and $\Omega_{N_0,1} = \Omega_{\text{per}}(\mathbf{S}_1^{\wedge}, N_0)$. Note that $\omega = \Omega_{N_0} \setminus \Omega_{N_0,1}$ up to a negligible set. The new minimizer \mathbf{S}_1^{\wedge} satisfies, by construction:

- $|\Omega_{\text{per}}(\mathbf{S}_1^{\wedge}, N_0 + 1)| = |\omega| + |\Omega_{\text{per}}(\mathbf{S}^{\wedge}, N_0 + 1)|$,
- $(\mathbf{S}_1^{\wedge})_{|\Omega_{\text{per}} \setminus \Omega_{N_0}} = (\mathbf{S}^{\wedge})_{|\Omega_{\text{per}} \setminus \Omega_{N_0}}$,
- $|\Omega_{N_0,1}| \leq \frac{1}{2} |\Omega_{N_0}|$,
- $\Omega_{N_0,1} \subset \Omega_{N_0}$.

Iterating the construction, we obtain a sequence $(\mathbf{S}_k^{\wedge})_{k \in \mathbb{N}} \in \mathcal{S}^{\mathbb{N}}$ of minimizers and a sequence of measurable sets $(\Omega_{N_0,k})_{k \in \mathbb{N}}$ such that, for each $k \geq 2$,

- (1) $\Omega_{\text{per}}(\mathbf{S}_k^{\wedge}, N_0) = \Omega_{N_0,k}$,
- (2) $|\Omega_{\text{per}}(\mathbf{S}_k^{\wedge}, N_0 + 1)| = |\Omega_{N_0,k-1} \setminus \Omega_{N_0,k}| + |\Omega_{\text{per}}(\mathbf{S}_{k-1}^{\wedge}, N_0 + 1)|$,
- (3) $(\mathbf{S}_k^{\wedge})_{|\Omega_{\text{per}} \setminus \Omega_{N_0,k-1}} = (\mathbf{S}_{k-1}^{\wedge})_{|\Omega_{\text{per}} \setminus \Omega_{N_0,k-1}}$,
- (4) $|\Omega_{N_0,k}| \leq \frac{1}{2} |\Omega_{N_0,k-1}|$,
- (5) $\Omega_{N_0,k} \subset \Omega_{N_0,k-1}$.

⁹Note that we cannot in general extend λ on the left of $\alpha = 0$, since for $\alpha < 0$, the matrix $\alpha \mathbf{T}^{\wedge} + \widehat{\mathbf{S}}^{\wedge}$ might loose the crucial property of essential nonnegativity. Thus the minimizer of $\lambda(\alpha)$, $\alpha = 0$, is not in general an interior critical point and $\lambda'(0) = 0$ cannot be deduced only from the first-order optimality condition. The role played by (22)–(23) is indeed crucial. By reversing one of the two inequalities, we obtain the proof of the complementary result on maximizers.

In particular,

$$0 \leq |\Omega_{N_0, k}| \leq \frac{1}{2k} |\Omega_{N_0}|, \quad |\Omega_{\text{per}}(\mathbf{S}_k^\wedge, N_0 + 1)| = |\Omega_{N_0}| - |\Omega_{N_0, k}| + |\Omega_{\text{per}}(\mathbf{S}^\wedge, N_0 + 1)|,$$

whence, as $k \rightarrow +\infty$,

$$|\Omega_{N_0, k}| \rightarrow 0, \quad |\Omega_{\text{per}}(\mathbf{S}_k^\wedge, N_0 + 1)| \rightarrow |\Omega_{N_0}| + |\Omega_{\text{per}}(\mathbf{S}^\wedge, N_0 + 1)|.$$

Let

$$\mathbf{S}_\infty^\wedge : (t, x) \in \overline{\Omega_{\text{per}}} \mapsto \begin{cases} \mathbf{S}^\wedge(t, x) & \text{if } (t, x) \in \overline{\Omega_{\text{per}}} \setminus \Omega_{N_0}, \\ \mathbf{S}_1^\wedge(t, x) & \text{if } (t, x) \in \Omega_{N_0} \setminus \Omega_{N_0, 1}, \\ \mathbf{S}_2^\wedge(t, x) & \text{if } (t, x) \in \Omega_{N_0, 1} \setminus \Omega_{N_0, 2}, \\ \vdots & \\ \mathbf{I} & \text{if } (t, x) \in \bigcap_{k \in \mathbb{N}} \Omega_{N_0, k}. \end{cases}$$

and extend it periodically in $\mathbb{R} \times \mathbb{R}^n$, so that $\mathbf{S}_\infty^\wedge \in \mathcal{S}$. Note that $|\bigcap_{k \in \mathbb{N}} \Omega_{N_0, k}| = 0$.

Then the sequence $(\mathbf{S}_k^\wedge)_{k \in \mathbb{N}}$ converges almost everywhere, and in any $\mathcal{L}_{\text{per}}^p(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$ with $p \in [1, +\infty)$, to \mathbf{S}_∞^\wedge . Moreover, by continuity of the mapping $\mathbf{S} \mapsto \lambda_1'(\mathbf{S})$ with respect to the topology of, say, $\mathcal{L}_{\text{per}}^2(\mathbb{R} \times \mathbb{R}^n, \mathbb{R}^{N \times N})$, \mathbf{S}_∞^\wedge is still a minimizer. Finally, it satisfies

$$|\Omega_{\text{per}}(\mathbf{S}_\infty^\wedge, N_0)| = 0 \quad \text{and} \quad |\Omega_{\text{per}}(\mathbf{S}_\infty^\wedge, N_0 + 1)| = |\Omega_{\text{per}}(\mathbf{S}^\wedge, N_0)| + |\Omega_{\text{per}}(\mathbf{S}^\wedge, N_0 + 1)|.$$

□

Step 4: when \mathbf{S}^\wedge is irreducible on average, correcting the minimizer in all possible sets Ω_{N_0} . Performing the construction of Steps 3 and 4, first for

$$N_0^\wedge = \min\{N_0 \in [N - 1] \mid |\Omega_{\text{per}}(\mathbf{S}^\wedge, N_0)| > 0\},$$

and then for $N_0^\wedge + 1$, etc., up to $N - 1$, we obtain in the end a new minimizer whose restriction to $\overline{\Omega_{\text{per}}}$ is valued in the set of permutation matrices almost everywhere, that is a new minimizer in $\mathcal{S}_{\{0, 1\}}$. □

Step 5: when \mathbf{S}^\wedge is reducible on average. The key tool for this generalization is a Frobenius normal form of the matrix with constant coefficients $\frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} \mathbf{S}^\wedge$.

This matrix is doubly stochastic. Indeed, its entries are valued in $[0, 1]$ and

$$\begin{aligned} \mathbf{1} &= \frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} \mathbf{1} = \frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} (\mathbf{S}^\wedge \mathbf{1}) = \left(\frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} \mathbf{S}^\wedge \right) \mathbf{1} \\ &= \frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} ((\mathbf{S}^\wedge)^T \mathbf{1}) = \left(\frac{1}{T|[0, L]|} \int_{\overline{\Omega_{\text{per}}}} \mathbf{S}^\wedge \right)^T \mathbf{1}. \end{aligned}$$

By nonnegativity, there exists a permutation matrix $\mathbf{F} \in \{0, 1\}^{N \times N}$ such that

$$\mathbf{S}^{\mathbf{F}} = \mathbf{F}^T \left[\frac{1}{T|[0, L]|} \left(\int_{\overline{\Omega_{\text{per}}}} \mathbf{S}^\wedge \right) \right] \mathbf{F}$$

is a block upper triangular matrix whose diagonal blocks are irreducible nonnegative square matrices (recall that 1×1 matrices are by convention referred to as irreducible even if zero) and whose off-diagonal blocks are nonnegative.

Let us verify that this Frobenius normal form $\mathbf{S}^{\mathbf{F}}$ is actually block diagonal, with doubly stochastic diagonal blocks.

Indeed, since $\frac{1}{T|[0,L]|} \int_{\Omega_{\text{per}}} \mathbf{S}^\wedge$ is doubly stochastic and since \mathbf{F} is a permutation matrix, then \mathbf{S}^F is also doubly stochastic. Moreover, the first diagonal block is a left-stochastic matrix (all columns sum to 1) and, since all off-diagonal blocks are nonnegative, its lines sum to at most 1. Let $N_1 \in [N]$ such that this first block is a $N_1 \times N_1$ matrix; then the sum of all entries of the block is exactly N_1 . Consequently, each line-sum is actually exactly 1, and the block is doubly stochastic. This, in turn, implies that all entries indexed by $(i, j) \in [N_1] \times [N] \setminus [N_1]$ are zero. Iterating on each diagonal block, we deduce that \mathbf{S}^F has indeed the claimed form.

Now, remark that each diagonal block of \mathbf{S}^F corresponds to a fully coupled subsystem in Ω_{per} .

Hence, up to permutations that are constant in space-time (for regularity reasons, this matters), we can assume without loss of generality that the operator $\text{diag}(\mathbf{P}) - \mathbf{B} - \mathbf{S}^\wedge \text{diag}(\boldsymbol{\mu})$ is in block diagonal form with each block fully coupled, and with each block of \mathbf{S}^\wedge a doubly stochastic matrix.

To conclude, it only remains to apply the correction of Steps 1–4 block by block. In the end, we obtain indeed a minimizer in $\mathcal{S}_{\{0,1\}}$. \square

This ends the proof. \square

Remark 3.24. Consistently with Neumann–Sze [62], the decomposition $\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ can be replaced by a more general decomposition $\mathbf{L} = \mathbf{B} + \mathbf{S}\mathbf{A}$ with \mathbf{A} nonnegative and \mathbf{B} essentially nonnegative. The generalization of the proof is straightforward.

Remark 3.25. From Proposition 3.32 and the fact that \max_z and $\max_{\mathbf{S}}$ commute, we can deduce a similar result on the maximizers of $\max_{z \in \mathbb{R}^n} \lambda_{1,z}(\mathbf{S})$. Yet we do not insist on it, for two reasons:

- (1) from the discussion in Section 1.6, we know that $\max_{z \in \mathbb{R}^n} \lambda_{1,z}(\mathbf{S})$ is not a satisfying generalization of $\lambda_1(\mathbf{S})$ when \mathbf{S} ceases to satisfy (A_3) , and clearly there are many $\mathbf{S} \in \mathcal{S}_{\{0,1\}}$ that do not satisfy (A_3) ;
- (2) since there is no reason why $\min_{\mathbf{S}}$ and \max_z should commute (in particular, $(\mathbf{S}, z) \mapsto \lambda_{1,z}(\mathbf{S})$ is not convex–concave), the argument does not apply to minimizers.

The problem of optimizing $\mathbf{S} \in \mathcal{S} \mapsto \lambda_1(\mathbf{S})$, that needs both a unambiguous definition of λ_1 when \mathbf{L} ceases to satisfy (A_3) and a new method of proof that applies to minimizers, remains therefore open.

3.4.2. *Optimization of the mutation rate: proof of Theorem 1.19.* Next we prove Theorem 1.19. The proof relies on a dual convexity lemma of Altenberg [3, Lemma 1] whose statement is recalled below.

Lemma 3.33 (Altenberg’s dual convexity lemma). *Let $f : (0, +\infty) \times [0, +\infty) \rightarrow \mathbb{R}$ be a function of two variables r and s , positively homogeneous of degree 1, and convex with respect to its second variable s .*

Then:

- (1) *f is convex with respect to its first variable r ; furthermore, $r \mapsto f(r, s)$ is strictly convex if $s \neq 0$ and if the convexity with respect to s is strict;*
- (2) *for all $(r, s) \in (0, +\infty) \times [0, +\infty)$, $z \mapsto f(r, s) + zf(1, 0) - f(r + z, s)$ is either identically zero or positive; furthermore it is positive if $s \neq 0$ and if f is strictly convex with respect to s ;*

(3) for all $r \in (0, +\infty)$,

$$\lim_{\substack{r' \rightarrow r \\ r' < r}} \frac{f(r', s) - f(r, s)}{r' - r} \leq \lim_{\substack{r' \rightarrow r \\ r' > r}} \frac{f(r', s) - f(r, s)}{r' - r} \leq f(1, 0) \quad \text{for all } s \in [0, +\infty),$$

and the first inequality is an equality except possibly at a countable number of values of r .

In our context, this lemma brings forth Theorem 1.19, as shown below.

Corollary 3.34. *Assume $(A_i)_{i \in [N]}$ is independent of t , $(q_i)_{i \in [N]} = 0$, and \mathbf{L} has the form $\mathbf{L} = \text{diag}(\mathbf{r}) + (\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ with $\mathbf{r} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R}^n, \mathbb{R}^N)$, $\boldsymbol{\mu} \in \mathcal{C}_{\text{per}}^{\delta/2, \delta}(\mathbb{R}^n, (\mathbf{0}, \infty))$ and $\mathbf{S} \in \mathcal{S}$ all independent of t .*

For any $\rho > 0$, let \mathcal{Q}_ρ be the operator with $(A_i)_{i \in [N]}$ and \mathbf{L} replaced by $(\rho A_i)_{i \in [N]}$ and $\text{diag}(\mathbf{r}) + \rho(\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu})$ respectively.

Then $\rho \in [0, 1] \mapsto \lambda'_1(\mathcal{Q}_\rho)$ is concave and nondecreasing. Furthermore, if \mathbf{r} depends on x and $s > 0$, then it is strictly concave and increasing.

Proof. We reduce the eigenvalue thanks to (9) and apply Lemma 3.33 to the function

$$f : (r, s) \mapsto -\lambda'_1(-r \text{diag}(\nabla \cdot (A_i \nabla)) - s \text{diag}(\mathbf{r}) - r(\mathbf{S} - \mathbf{I}) \text{diag}(\boldsymbol{\mu}))$$

which is, by virtue of Theorem 1.3, convex with respect to $s \in (0, +\infty)$, strictly if \mathbf{r} depends on x , and which is of class \mathcal{C}^1 in (r, s) away from $r = 0$. Subsequently, $f(1, 0) = 0$ yields the monotonicity of f with respect to r . \square

Note that a version of the above corollary appropriate for $\lambda_{1,z}$, with an assumption $A_i^{-1} q_i = 2z + \nabla Q$ reminiscent of Theorem 1.13, could be just as easily established. For the sake of brevity, we focus here on λ'_1 only.

3.4.3. Optimization of the spatial distribution of \mathbf{L} in dimension 1: proof of Theorem 1.20. In this section, we prove Theorem 1.20.

First, we investigate a Talenti inequality for cooperative elliptic systems, as such estimates are milestones to proving spectral comparison [59, Theorem 3.9]. Let us recall that the core idea underlying these estimates is to compare some \mathcal{L}^p norms (here, the \mathcal{L}^∞ norms) of the solution of an elliptic problem with that of a related equation, the coefficients of which have been replaced by their symmetrization. It should be noted that our results would also hold for boundary conditions of Dirichlet type in the spatial domain $B(0, R)$, $R > 0$.

Remark 3.26. In what follows, we will use a few specific notations.

It will be convenient to identify $[0, L_1]$ with $[-\frac{L_1}{2}, \frac{L_1}{2}]$; this amounts to translating the functions, and has the advantage of having 0 as a symmetry point. In this context, let us recall the fundamental ordering on the set of functions: for two functions $\mathbf{f}, \mathbf{g} \in \mathcal{L}^2([0, L_1], [\mathbf{0}, \infty))$, the notation $\mathbf{f} \prec \mathbf{g}$ stands for:

$$\int_{-\frac{r}{2}}^{\frac{r}{2}} \mathbf{f} \leq \int_{-\frac{r}{2}}^{\frac{r}{2}} \mathbf{g} \quad \text{for all } r \leq \frac{L_1}{2}.$$

In particular, these inequalities hold component wise.

For any non-negative scalar function, we may identify its rearrangement u^\dagger with a non-increasing mapping $\bar{u} : [0, L_1/2] \rightarrow \mathbb{R}$. For any non-negative \mathbf{u} , we denote by \mathbf{u}^\dagger its periodic rearrangement.

The first step in the proof of Theorem 1.20 is the following comparison result for elliptic systems.

Proposition 3.35. *Assume $\text{diag}(\mathcal{P}) = \partial_t - \mathbf{D}\Delta$ for some diagonal matrix \mathbf{D} with constant, positive diagonal entries and assume that \mathbf{L} is nonnegative and depends only on x .*

Let $c > 0$ and $\phi, \psi \in \mathcal{L}^2([-\frac{L_1}{2}, \frac{L_1}{2}], [0, \infty))$ such that $\phi \prec \psi$. Let \mathbf{u} and \mathbf{v} be the (unique) solutions of

$$\begin{cases} -\mathbf{D}\Delta\mathbf{u} + c\mathbf{u} = \mathbf{L}\phi & \text{in } [-\frac{L_1}{2}, \frac{L_1}{2}], \\ \mathbf{u} \in \mathcal{W}_{\text{per}}^{1,2} \end{cases}$$

and

$$\begin{cases} -\mathbf{D}\Delta\mathbf{v} + c\mathbf{v} = \mathbf{L}^\dagger\psi^\dagger & \text{in } [-\frac{L_1}{2}, \frac{L_1}{2}] \\ \mathbf{v} \in \mathcal{W}_{\text{per}}^{1,2} \end{cases}$$

Then $\mathbf{u} \prec \mathbf{v}$.

Proof. First, let us verify that if ϕ, ψ are nonnegative and satisfy $\phi \prec \psi$, then

$$(24) \quad \mathbf{L}^\dagger\phi^\dagger \prec \mathbf{L}^\dagger\psi^\dagger.$$

First of all, for any $s \in [0, \frac{L_1}{2}]$ and any $j \in [N]$,

$$\chi_{[-s, s]}\phi_j^\dagger \prec \chi_{[-s, s]}\psi_j^\dagger,$$

where χ denotes the characteristic function of a set. This property is stable by addition and multiplication by a nonnegative constant [6]. As a consequence, since any nonnegative nonincreasing function can be approximated from below by a nonnegative step function, (24) follows from the monotone convergence theorem.

For the sake of simplicity, assume the level sets of each u_i have zero Lebesgue measure – should this not be the case, we can argue exactly as in [67]. Let $\tau \geq 0$ be a fixed real number and let $i \in [N]$. Integrating the i -th equation on the level set $\{u_i \geq \tau\}$, we get

$$d_i \int_{\{u_i = \tau\}} |\nabla u_i| = -c \int_{\{u_i \geq \tau\}} u_i + \sum_{j=1}^N \int_{\{u_i \geq \tau\}} l_{i,j} \phi_j.$$

Since u and u^\dagger are equimeasurable by the definition of the periodic rearrangement, there holds

$$c \int_{\{u_i \geq \tau\}} u_i = c \int_{\{u_i^\dagger \geq \tau\}} u_i^\dagger.$$

By the Hardy–Littlewood inequality, since all the $l_{i,j}$ are nonnegative,

$$\sum_{j=1}^N \int_{\{u_i \geq \tau\}} l_{i,j} \phi_j \leq \sum_{j=1}^N \int_{\{u_i^\dagger \geq \tau\}} l_{i,j}^\dagger \phi_j^\dagger.$$

At this point, we have obtained

$$d_i \int_{\{u_i = \tau\}} |\nabla u_i| \leq -c \int_{\{u_i^\dagger \geq \tau\}} u_i^\dagger + \sum_{j=1}^N \int_{\{u_i^\dagger \geq \tau\}} l_{i,j}^\dagger \phi_j^\dagger.$$

By (24), we thus conclude that

$$d_i \int_{\{u_i=\tau\}} |\nabla u_i| \leq -c \int_{\{u_i^\dagger \geq \tau\}} u_i^\dagger + \sum_{j=1}^N \int_{\{u_i^\dagger \geq \tau\}} l_{i,j}^\dagger \psi_j^\dagger.$$

We introduce the distribution function μ_i of u_i ,

$$\mu_i(\tau) = |\{u_i > \tau\}|.$$

From the co-area formula,

$$-\mu_i'(\tau) = \int_{\{u_i=\tau\}} \frac{1}{|\nabla u_i|}.$$

Since the periodic rearrangement decreases the perimeter of level-sets, we have

$$\text{Per}(\{u_i^\dagger = \tau\}) \leq \text{Per}(\{u_i = \tau\}).$$

From the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} \text{Per}(\{u_i^\dagger = \tau\})^2 &\leq \text{Per}(\{u_i = \tau\})^2 \\ &\leq \int_{\{u_i=\tau\}} \frac{1}{|\nabla u_i|} \int_{\{u_i=\tau\}} |\nabla u_i| \\ &\leq -\mu_i'(\tau) \int_{\{u_i=\tau\}} |\nabla u_i| \\ &\leq -\frac{\mu_i'(\tau)}{d_i} \left(-c \int_{\{u_i^\dagger \geq \tau\}} u_i^\dagger + \sum_{j=1}^N \int_{\{u_i^\dagger \geq \tau\}} l_{i,j}^\dagger \psi_j^\dagger \right). \end{aligned}$$

Since we are working in one dimension, for any $\tau \in (\min(u_i), \max(u_i))$, there holds

$$4 \leq \text{Per}(\{u_i^\dagger = \tau\})^2.$$

Furthermore, by definition of the rearrangement,

$$\int_{\{u_i > \tau\}} u_i = \int_0^{\mu_i(\tau)} \bar{u}_i.$$

We define, for $i \in [N]$,

$$k_i^{\mathbf{u}} : \xi \in \left[0, \frac{L_1}{2}\right] \mapsto \int_0^\xi \bar{u}_i.$$

From this definition, we obtain

$$(k_i^{\mathbf{u}})''(\mu_i(\tau)) = \bar{u}_i'(\mu_i(\tau)) = \frac{1}{\mu_i'(\tau)}.$$

With these notations, we obtain the following differential inequality: for any $\xi \in [0, L_1/2]$,

$$-4(k_i^{\mathbf{u}})''(\xi) \leq -\frac{c}{d_i} k_i^{\mathbf{u}}(\xi) + \sum_{j=1}^N \int_0^\xi l_{i,j}^\dagger \psi_j^\dagger.$$

Furthermore,

$$k_i^{\mathbf{u}}(0) = 0.$$

Working with $\boldsymbol{\psi}$ instead of $\boldsymbol{\phi}$ and with \mathbf{v} instead of \mathbf{u} , all the previous inequalities are equalities: indeed, by the variational formulation of the equation on each of the

coordinates v_i , it appears that $v_i^\dagger = v_i$, so that, for any $\tau \in (\min(v_i), \max(v_i))$, $\text{Per}(\{v_i = \tau\})^2 = 4$. Thus, with transparent notations, $k_i^{\mathbf{v}}$ solves the differential equation

$$-4(k_i^{\mathbf{v}})''(\xi) = -\frac{c}{d_i}k_i^{\mathbf{v}}(\xi) + \sum_{j=1}^N \int_0^\xi l_{i,j}^\dagger \psi_j^\dagger.$$

Similarly,

$$k_i^{\mathbf{v}}(0) = 0.$$

Hence, from (24), the vector $\mathbf{k} = \mathbf{k}^{\mathbf{v}} - \mathbf{k}^{\mathbf{u}}$ satisfies

$$-4 \text{diag}(\mathbf{d})\mathbf{k}'' + \mathbf{c}\mathbf{k} \geq \mathbf{0}, \quad \mathbf{k}(0) = \mathbf{0}.$$

Finally, integrating both equations in \mathbf{u} and \mathbf{v} on the domain we obtain

$$\int_{[0, L_1/2]} \mathbf{u}_i = \frac{1}{c} \int_{[0, L_1/2]} (\mathbf{L}\phi)_i \leq \frac{1}{c} \int_{[0, L_1/2]} (\mathbf{L}\psi)_i = \int_{[0, L_1/2]} \mathbf{v}_i,$$

so that

$$\mathbf{k} \left(\frac{L_1}{2} \right) \geq \mathbf{0}.$$

From the maximum principle,

$$\mathbf{k} \geq \mathbf{0} \quad \text{in} \left(0, \frac{L_1}{2} \right).$$

However, this is exactly the desired conclusion. \square

We now apply Proposition 3.35 to derive a comparison principle.

Proposition 3.36. *Assume $\text{diag}(\mathcal{P}) = \partial_t - \mathbf{D}\Delta$ for some diagonal matrix \mathbf{D} with constant, positive diagonal entries.*

Let $\mathbf{u}_0 \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R}, [\mathbf{0}, \infty))$ and let \mathbf{u}, \mathbf{v} be the respective space periodic solutions of

$$\begin{cases} \mathcal{Q}\mathbf{u} = \mathbf{0} & \text{in } (0, T) \times \mathbb{R}, \\ \mathbf{u} = \mathbf{u}_0 & \text{on } \{0\} \times \mathbb{R} \end{cases}$$

and

$$\begin{cases} \text{diag}(\mathcal{P})\mathbf{v} - \mathbf{L}^\dagger \mathbf{v} = \mathbf{0} & \text{in } (0, T) \times \mathbb{R}, \\ \mathbf{v} = \mathbf{u}_0^\dagger & \text{on } \{0\} \times \mathbb{R}. \end{cases}$$

Then, for all $t \in [0, T]$, $\mathbf{u}(t, \cdot) \prec \mathbf{v}(t, \cdot)$.

Proof. We use a classical time discretization of the system, following [5, Proof of Theorem 3].

Let $c > 0$ so large that $l_{i,i} + c \geq 0$ for all $i \in [N]$. Let $K \in \mathbb{N}$ and $\delta = \frac{T}{K} > 0$. We define, for any $\omega \in [K]$,

$$\mathbf{L}_\omega = K \int_{(\omega-1)\delta}^{\omega\delta} (\mathbf{L}(\tau, \cdot) + c\mathbf{I})d\tau, \quad \mathbf{L}_{\omega, \dagger} = K \int_{(\omega-1)\delta}^{\omega\delta} (\mathbf{L}^\dagger(\tau, \cdot) + c\mathbf{I})d\tau.$$

Clearly $\mathbf{L}_\omega + K\mathbf{I} \prec \mathbf{L}_{\omega, \dagger} + K\mathbf{I}$ and both are nonnegative and only depend on space. We set $\mathbf{u}^0 = \mathbf{u}_0$, $\mathbf{v}^0 = \mathbf{u}_0^\dagger$ and consider, for any $\omega \in [K]$, the space periodic solutions of the elliptic systems

$$\begin{aligned} (K + c)\mathbf{u}^\omega - \mathbf{D}\Delta\mathbf{u}^\omega &= K\mathbf{u}^{\omega-1} + \mathbf{L}_\omega\mathbf{u}^{\omega-1}, \\ (K + c)\mathbf{v}^\omega - \mathbf{D}\Delta\mathbf{v}^\omega &= K\mathbf{v}^{\omega-1} + \mathbf{L}_{\omega, \dagger}\mathbf{v}^{\omega-1}. \end{aligned}$$

By an immediate recursion, and since the coefficients of the system satisfied by \mathbf{v}^ω are spatially rearranged, \mathbf{v}^ω itself is rearranged, for any $\omega \in [K]$. Indeed, this follows from the uniqueness of the solutions of the above systems and from the existence of radial solutions by using radial coordinates. Subsequently, by Proposition 3.35, for any $\omega \in [K]$, $\mathbf{u}^\omega \prec \mathbf{v}^\omega$.

Proceeding as in [5], we may pass to the limit $K \rightarrow \infty$ to conclude the proof. \square

Remark 3.27. Regarding the convergence of the sequence of elliptic problems to the parabolic one, although we do not detail it for the sake of conciseness, the easiest way to proceed here is to assume that \mathbf{L} is \mathcal{C}^1 in the time variable, which is always possible through a standard approximation argument. From this point of view, a Taylor expansion shows that, if \mathbf{u} is the solution of the parabolic equation, if $\bar{\mathbf{u}}^\omega := K \int_{(\omega-1)\delta}^{\omega\delta} \mathbf{u}$ and if \mathbf{u} is the (piecewise) affine interpolation of $(\mathbf{u}^\omega)_{\omega \in [K]}$, then we have $\|\bar{\mathbf{u}} - \mathbf{u}\|_{\mathcal{L}^2((0,T), \mathcal{W}^{1,2})} \leq C\delta$. Although the idea of such a discretization was used in a systematic way in [5], it was introduced in [68] where the discretization procedure was justified by the use of the (more abstract) Crandall–Liggett theorem.

Proposition 3.37. *Assume $\text{diag}(\mathcal{P}) = \partial_t - \mathbf{D}\Delta$ for some diagonal matrix \mathbf{D} with constant, positive diagonal entries.*

Then

$$\lambda_{1,\text{per}}(\mathcal{Q}) \geq \lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - \mathbf{L}^\dagger)$$

where \mathbf{L}^\dagger is the entry-wise periodic rearrangement of \mathbf{L} .

Proof. The proof relies on Proposition 3.36.

We proceed as in [59] and introduce, for some $c > 0$ so large that $\mathbf{L} + c\mathbf{I} \geq \mathbf{0}$, the Poincaré mapping

$$\mathcal{G}_{\mathbf{L}+c\mathbf{I}} : \begin{cases} \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N) & \rightarrow \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N) \\ \mathbf{u}_0 & \mapsto \mathbf{u}(\mathbf{u}_0, \mathbf{L} + c\mathbf{I}; T, \cdot) \end{cases}$$

where $(t, x) \mapsto \mathbf{u}(\mathbf{u}_0, \mathbf{L} + c\mathbf{I}; t, x)$ is the solution of $\text{diag}(\mathcal{P})\mathbf{u} + c\mathbf{u} = (\mathbf{L} + c\mathbf{I})\mathbf{u}$ with initial condition \mathbf{u}_0 .

We define $r(\mathbf{L} + c\mathbf{I})$ as the principal eigenvalue of the operator $\mathcal{G}_{\mathbf{L}+c\mathbf{I}}$. As is classical, this eigenvalue can be obtained as

$$r(\mathbf{L} + c\mathbf{I}) = \lim_{k \rightarrow \infty} \|\mathcal{G}_{\mathbf{L}+c\mathbf{I}}^k\|^{1/k},$$

where the notation $\|\cdot\|$ stands for the norm on the vector space of linear mappings from $\mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N)$ into itself (for the $\mathcal{L}^\infty - \mathcal{L}^\infty$ norm), and the two quantities $r(\mathbf{L} + c\mathbf{I})$ and $\lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - \mathbf{L})$ are immediately related through $r(\mathbf{L} + c\mathbf{I}) = -\frac{1}{T} \lambda_{1,\text{per}}(\text{diag}(\mathcal{P}) - \mathbf{L})$. As a consequence, in order to obtain the required comparison result, it suffices to establish that, for any $\mathbf{u}_0 \in \mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N)$,

$$\|\mathcal{G}_{\mathbf{L}+c\mathbf{I}}\mathbf{u}_0\|_{\mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N)} \leq \|\mathcal{G}_{\mathbf{L}^\dagger+c\mathbf{I}}\mathbf{u}_0\|_{\mathcal{L}_{\text{per}}^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R}^N)}.$$

Yet this is a direct consequence of Proposition 3.36. \square

Remark 3.28. The previous result established the optimality of periodically rearranged entries of \mathbf{L} when rearrangements are all centered at the same arbitrary spatial position, $x = 0$ in our construction. The spatially periodic entries of \mathbf{L} must be “in phase”. If rearrangements of different entries of \mathbf{L} are centered at different positions, the optimality fails, as the following counter-example shows.

We consider, in the one-dimensional case $n = 1$ with $L_1 = 2$, a space periodic function χ whose restriction to $[-1, 1]$ is the indicator function of $(-y, y)$, $0 < y < 1$, a real number η , and the matrix

$$\mathbf{L} : x \mapsto \begin{pmatrix} -1 + \chi(x) & 1 + \chi(x - \eta) \\ 1 + \chi(x - \eta) & -1 + \chi(x) \end{pmatrix}.$$

The vector $\mathbf{1}$ is a Perron–Frobenius eigenvector of $\mathbf{L}(x)$ with Perron–Frobenius eigenvalue $\mu_\eta(x) = \chi(x) + \chi(x - \eta)$. Let \mathbf{u} be a periodic principal eigenfunction of $\mathcal{Q} = \partial_t - \Delta - \mathbf{L}$. The function $u_\eta = \mathbf{1}^T \mathbf{u}$ is positive, time homogeneous, space periodic and solves

$$-\Delta u_\eta = \mu_\eta u_\eta + \lambda_{1,\text{per}}(\mathcal{Q})u_\eta.$$

Therefore $\lambda_{1,\text{per}}(\mathcal{Q}) = \lambda_{1,\text{per}}(-\Delta - \mu_\eta)$, where the last operator is a scalar space periodic elliptic operator. Note that χ and $\chi(\cdot - \eta)$ are invariant by periodic rearrangement centered at $x = 0$ and $x = \eta$ respectively. Nonetheless, \mathbf{L} is not optimal as soon as $\eta \neq 0$. Indeed,

$$\mu_\eta(x) = \begin{cases} 0 & \text{if } x \in (-1, -y) \cup (y + \eta, 1), \\ 1 & \text{if } x \in (-y, -y + \eta) \cup (y, y + \eta), \\ 2 & \text{if } x \in (-y + \eta, y). \end{cases}$$

Hence all $(\mu_\eta)_{\eta \in \mathbb{R}}$ are piecewise-constant, space periodic functions of total mass equal to $4y > 0$. It is well-known [59] that among these the one that minimizes $\lambda_{1,\text{per}}(-\Delta - \mu_\eta)$ is the one corresponding to $\eta = 0$. This is of course consistent with our optimization result.

4. ACKNOWLEDGEMENTS

The authors are grateful to the anonymous reviewer for a careful reading of the manuscript and insightful comments that lead to substantial clarifications and improvements. They are also grateful to G. Nadin for several inspiring discussions and useful advice.

L. G. acknowledges support from the ANR via the project Indycana under grant agreement ANR-21-CE40-0008. I. M.-F. was partially funded by a PSL Young Researcher Starting Grant 2023.

REFERENCES

- [1] Matthieu Alfaro and Quentin Griette. Pulsating fronts for Fisher-KPP systems with mutations as models in evolutionary epidemiology. *Nonlinear Anal. Real World Appl.*, 42:255–289, 2018.
- [2] Grégoire Allaire and Harsha Hutridurga. On the homogenization of multicomponent transport. *Discrete Contin. Dyn. Syst. Ser. B*, 20(8):2527–2551, 2015.
- [3] Lee Altenberg. Resolvent positive linear operators exhibit the reduction phenomenon. *Proc. Natl. Acad. Sci. USA*, 109(10):3705–3710, 2012.
- [4] Lee Altenberg and Joel E. Cohen. Nonconcavity of the spectral radius in Levinger’s theorem. *Linear Algebra Appl.*, 606:201–218, 2020.
- [5] Alvino Alvino, Guido Trombetti, and Pierre-Louis Lions. Comparison results for elliptic and parabolic equations via Schwarz symmetrization. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 7(2):37–65, 1990.
- [6] Angelo Alvino, Pierre-Louis Lions, and Guido Trombetti. Comparison results for elliptic and parabolic equations via symmetrization: a new approach. *Differential Integral Equations*, 4(1):25–50, 1991.

- [7] Bénédicte Alziary, Jacqueline Fleckinger, and Marie-Hélène Lécureux. Principal eigenvalue and maximum principle for some elliptic systems defined on general domains with refined Dirichlet boundary condition. *Commun. Math. Anal.*, 7(2):1–11, 2009.
- [8] Inmaculada Antón and Julián López-Gómez. The strong maximum principle for cooperative periodic-parabolic systems and the existence of principal eigenvalues. In *World Congress of Nonlinear Analysts '92, Vol. I–IV (Tampa, FL, 1992)*, pages 323–334. de Gruyter, Berlin, 1996.
- [9] Ari Arapostathis, Anup Biswas, and Somnath Pradhan. On the monotonicity property of the generalized eigenvalue for weakly-coupled cooperative elliptic systems. *J. Differential Equations*, 351:156–193, 2023.
- [10] Ari Arapostathis, Mrinal K. Ghosh, and Steven I. Marcus. Harnack’s inequality for cooperative weakly coupled elliptic systems. *Comm. Partial Differential Equations*, 24(9-10):1555–1571, 1999.
- [11] Xueli Bai and Xiaoqing He. Asymptotic behavior of the principal eigenvalue for cooperative periodic-parabolic systems and applications. *J. Differential Equations*, 269(11):9868–9903, 2020.
- [12] Michel Benaim, Claude Lobry, Tewfik Sari, and Edouard Strickler. When can a population spreading across sink habitats persist ? *arXiv e-prints*, page arXiv:2311.04706, November 2023.
- [13] Henri Berestycki, François Hamel, and Lionel Roques. Analysis of the periodically fragmented environment model. I. Species persistence. *J. Math. Biol.*, 51(1):75–113, 2005.
- [14] Henri Berestycki, François Hamel, and Luca Rossi. Liouville-type results for semilinear elliptic equations in unbounded domains. *Ann. Mat. Pura Appl. (4)*, 186(3):469–507, 2007.
- [15] Henri Berestycki, Louis Nirenberg, and SR Srinivasa Varadhan. The principal eigenvalue and maximum principle for second-order elliptic operators in general domains. *Communications on Pure and Applied Mathematics*, 47(1):47–92, 1994.
- [16] Henri Berestycki and Luca Rossi. On the principal eigenvalue of elliptic operators in \mathbb{R}^N and applications. *Journal of the European Mathematical Society*, 8(2):195–215, 2006.
- [17] Henri Berestycki and Luca Rossi. Generalizations and properties of the principal eigenvalue of elliptic operators in unbounded domains. *Comm. Pure Appl. Math.*, 68(6):1014–1065, 2015.
- [18] Isabeau Birindelli, Enzo Mitidieri, and Guido Sweers. Existence of the principal eigenvalue for cooperative elliptic systems in a general domain. *Differ. Uravn.*, 35(3):325–333, 1999.
- [19] Sidi Mohammed Bouguima, S. Fekih, and W. Hennaoui. Spacial structure in a juvenile-adult model. *Nonlinear Anal. Real World Appl.*, 9(3):1184–1201, 2008.
- [20] Yann Brenier. Polar factorization and monotone rearrangement of vector-valued functions. *Comm. Pure Appl. Math.*, 44(4):375–417, 1991.
- [21] K. J. Brown and Yanping Zhang. On a system of reaction-diffusion equations describing a population with two age groups. *J. Math. Anal. Appl.*, 282(2):444–452, 2003.
- [22] Robert Stephen Cantrell, Chris Cosner, and Salomé Martínez. Persistence for a two-stage reaction-diffusion system. *Mathematics*, 8(3), 2020.
- [23] Robert Stephen Cantrell, Chris Cosner, and Xiao Yu. Dynamics of populations with individual variation in dispersal on bounded domains. *Journal of Biological Dynamics*, 12(1):288–317, 2018.
- [24] Robert Stephen Cantrell, Chris Cosner, and Xiao Yu. Populations with individual variation in dispersal in heterogeneous environments: Dynamics and competition with simply diffusing populations. *Science China Mathematics*, Jan 2020.
- [25] Yves Capdeboscq. Homogenization of a neutronic critical diffusion problem with drift. *Proc. Roy. Soc. Edinburgh Sect. A*, 132(3):567–594, 2002.
- [26] Zhen-Qing Chen and Zhong X. Zhao. Harnack principle for weakly coupled elliptic systems. *J. Differential Equations*, 139(2):261–282, 1997.
- [27] Djairo G. de Figueiredo. Monotonicity and symmetry of solutions of elliptic systems in general domains. *NoDEA Nonlinear Differential Equations Appl.*, 1(2):119–123, 1994.
- [28] Djairo G. de Figueiredo and Enzo Mitidieri. Maximum principles for linear elliptic systems. *Rend. Istit. Mat. Univ. Trieste*, 22(1-2):36–66, 1990.
- [29] B. Curtis Eaves, Alan J. Hoffman, Uriel G. Rothblum, and Hans Schneider. Line-sum-symmetric scalings of square nonnegative matrices. Number 25, pages 124–141. 1985. Mathematical programming, II.

- [30] Elizabeth C. Elliott and Stephen J. Cornell. Dispersal polymorphism and the speed of biological invasions. *PLOS ONE*, 7(7):1–10, 07 2012.
- [31] Ronald Aylmer Fisher. The wave of advance of advantageous genes. *Annals of eugenics*, 7(4):355–369, 1937.
- [32] Juraj Földes and Peter Poláčik. On cooperative parabolic systems: Harnack inequalities and asymptotic symmetry. *Discrete Contin. Dyn. Syst.*, 25(1):133–157, 2009.
- [33] David Gilbarg and Neil S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Springer-Verlag, 2001.
- [34] Léo Girardin. Non-cooperative Fisher–KPP systems: Asymptotic behavior of traveling waves. *Mathematical Models and Methods in Applied Sciences*, 28(06):1067–1104, 2018.
- [35] Léo Girardin. Non-cooperative Fisher–KPP systems: traveling waves and long-time behavior. *Nonlinearity*, 31(1):108, 2018.
- [36] Léo Girardin. Addendum to ‘Non-cooperative Fisher–KPP systems: traveling waves and long-time behavior’. *Nonlinearity*, 32(1):168, 2019.
- [37] Léo Girardin. Two components is too simple: an example of oscillatory Fisher–KPP system with three components. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, pages 1–24, 2019.
- [38] Léo Girardin. Persistence, extinction and spreading properties of non-cooperative Fisher–KPP systems in space-time periodic media. *arXiv e-prints*, page arXiv:2305.18857, May 2023.
- [39] Léo Girardin and Quentin Griette. A Liouville-type result for non-cooperative Fisher–KPP systems and nonlocal equations in cylinders. *Acta Applicandae Mathematicae*, 170(1):123–139, Dec 2020.
- [40] Quentin Griette and Hiroshi Matano. Propagation dynamics of solutions to spatially periodic reaction-diffusion systems with hybrid nonlinearity. *arXiv e-prints*, page arXiv:2108.10862, August 2021.
- [41] Quentin Griette and Gaël Raoul. Existence and qualitative properties of travelling waves for an epidemiological model with mutations. *J. Differential Equations*, 260(10):7115–7151, 2016.
- [42] Quentin Griette, Gaël Raoul, and Sylvain Gandon. Virulence evolution at the front line of spreading epidemics. *Evolution*, 69(11):2810–2819, 2015.
- [43] François Hamel, Grégoire Nadin, and Lionel Roques. A viscosity solution method for the spreading speed formula in slowly varying media. *Indiana University Mathematics Journal*, 60:1229–1247, 2011.
- [44] Li Jun Hei and Jian Hua Wu. Existence and stability of positive solutions for an elliptic cooperative system. *Acta Math. Sin. (Engl. Ser.)*, 21(5):1113–1120, 2005.
- [45] Ouassila Henaoui. An elliptic system modeling two subpopulations. *Nonlinear Anal. Real World Appl.*, 13(6):2447–2458, 2012.
- [46] Peter Hess. *Periodic-parabolic boundary value problems and positivity*, volume 247 of *Pitman Research Notes in Mathematics Series*. Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1991.
- [47] Vivian Hutson, Konstantin Mischaikow, and Peter Poláčik. The evolution of dispersal rates in a heterogeneous time-periodic environment. *J. Math. Biol.*, 43(6):501–533, 2001.
- [48] Samuel Karlin. Classifications of selection-migration structures and conditions for a protected polymorphism. *Evol. Biol.*, 14(61):204, 1982.
- [49] Andrei N. Kolmogorov, I. G. Petrovsky, and N. S. Piskunov. Étude de l’équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique. *Bulletin Université d’État à Moscou*, 1:1–25, 1937.
- [50] King-Yeung Lam and Yuan Lou. Asymptotic behavior of the principal eigenvalue for cooperative elliptic systems and applications. *J. Dynam. Differential Equations*, 28(1):29–48, 2016.
- [51] Xing Liang, Lei Zhang, and Xiao-Qiang Zhao. The principal eigenvalue for degenerate periodic reaction-diffusion systems. *SIAM J. Math. Anal.*, 49(5):3603–3636, 2017.
- [52] Gary M. Lieberman. *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [53] Shuang Liu, Yuan Lou, Rui Peng, and Maolin Zhou. Monotonicity of the principal eigenvalue for a linear time-periodic parabolic operator. *Proc. Amer. Math. Soc.*, 147(12):5291–5302, 2019.

- [54] Shuang Liu, Yuan Lou, Rui Peng, and Maolin Zhou. Asymptotics of the principal eigenvalue for a linear time-periodic parabolic operator I: large advection. *SIAM J. Math. Anal.*, 53(5):5243–5277, 2021.
- [55] Shuang Liu, Yuan Lou, Rui Peng, and Maolin Zhou. Asymptotics of the principal eigenvalue for a linear time-periodic parabolic operator II: Small diffusion. *Trans. Amer. Math. Soc.*, 374(7):4895–4930, 2021.
- [56] Ruyun Ma, Ruipeng Chen, and Yanqiong Lu. Steady states of a nonlinear elliptic system arising from population dynamics. *Bound. Value Probl.*, pages 2014:185, 7, 2014.
- [57] Sepideh Mirrahimi and Panagiotis E. Souganidis. A homogenization approach for the motion of motor proteins. *NoDEA Nonlinear Differential Equations Appl.*, 20(1):129–147, 2013.
- [58] Aled Morris, Luca Börger, and Elaine C. M. Crooks. Individual variability in dispersal and invasion speed. *Mathematics*, 7(9), 2019.
- [59] Grégoire Nadin. The principal eigenvalue of a space–time periodic parabolic operator. *Annali di Matematica Pura ed Applicata*, 188(2):269–295, 2009.
- [60] Grégoire Nadin. Traveling fronts in space-time periodic media. *J. Math. Pures Appl. (9)*, 92(3):232–262, 2009.
- [61] Grégoire Nadin. Existence and uniqueness of the solution of a space-time periodic reaction-diffusion equation. *J. Differential Equations*, 249(6):1288–1304, 2010.
- [62] Michael Neumann and Nung-Sing Sze. Optimization of the spectral radius of nonnegative matrices. *Oper. Matrices*, 1(4):593–601, 2007.
- [63] Roger D. Nussbaum. Convexity and log convexity for the spectral radius. *Linear Algebra Appl.*, 73:59–122, 1986.
- [64] Murray H. Protter and Hans F. Weinberger. *Maximum Principles in Differential Equations*. Springer-Verlag, 1984.
- [65] Wojciech Słomczyński. Irreducible cooperative systems are strongly monotone. *Univ. Iagel. Acta Math.*, (30):159–163, 1993.
- [66] Guido Sweers. Strong positivity in $C(\overline{\Omega})$ for elliptic systems. *Math. Z.*, 209(2):251–271, 1992.
- [67] Giorgio Talenti. Elliptic equations and rearrangements. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 3(4):697–718, 1976.
- [68] Juan Luis Vázquez. Symétrisation pour $u_1 = \Delta\varphi(u)$ et applications. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(2):71–74, 1982.