

# Influence of ionizing radiation on molecular recognition in the gas phase: VUV and X-ray photoabsorption of antibiotic-receptor non-covalent complexes

M Abdelmouleh, M Lalande, A Espinosa Rodriguez, V Vizcaino, J Rangama, T Schlathölter, J-C Poully

### ▶ To cite this version:

M Abdelmouleh, M Lalande, A Espinosa Rodriguez, V Vizcaino, J Rangama, et al.. Influence of ionizing radiation on molecular recognition in the gas phase: VUV and X-ray photoabsorption of antibiotic-receptor non-covalent complexes. 31st International Conference on Photonic, Electronic and Atomic Collisions (ICPEAC XXXI) 23-30 July 2019, Deauville, France, 2020, Deauville, France. pp.152061, 10.1088/1742-6596/1412/15/152061. hal-03349486

HAL Id: hal-03349486

https://hal.science/hal-03349486

Submitted on 20 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### **PAPER • OPEN ACCESS**

Influence of ionizing radiation on molecular recognition in the gas phase: VUV and X-ray photoabsorption of antibiotic-receptor non-covalent complexes

To cite this article: M Abdelmouleh et al 2020 J. Phys.: Conf. Ser. 1412 152061

View the article online for updates and enhancements.



**1412** (2020) 152061 doi:10.1088/1742-6596/1412/15/152061

## Influence of ionizing radiation on molecular recognition in the gas phase: VUV and X-ray photoabsorption of antibiotic-receptor non-covalent complexes

M Abdelmouleh<sup>1\*</sup>, M Lalande<sup>1</sup>, A Espinosa Rodriguez<sup>1,2</sup>, V Vizcaino<sup>1</sup>, J Rangama<sup>1</sup>, T Schlathölter<sup>2</sup> and J-C Poullv1

<sup>1</sup>CIMAP, UMR 6252 (CEA/CNRS/ENSICAEN/Université de Caen Normandie), Caen, France <sup>2</sup>Dpto. de Física Atómica, Molecular y Nuclear. Facultad de CC Físicas. Avda. Complutense, s/n - 28040 Madrid, Spain <sup>3</sup>Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747AG Groningen, The Netherlands

Synopsis The intrinsic effects of ionizing radiation on molecular recognition between an antibiotic and its receptor have been probed by coupling a synchrotron beamline to a tandem mass spectrometer. Protonated vancomycin-receptor non-covalent complexes in the gas phase undergo inter- as well as intramolecular bond cleavage due to internal energy transfer after photoabsorption in the VUV-X energy range. The deprotonated complex undergoes electron detachment, but the native interactions underlying molecular recognition are not systematically

The structure of biological molecules plays an important role in their activity, especially in the molecular recognition between drugs and their receptors. This is illustrated in figure 1 for the glycopeptide antibiotic vancomycin bound to its peptidic receptor Ac2-Lys-DAla-DAla in condensed phase, where the two structures tightly fit each other. When isolated in the gas phase, this native structure is lost for gas-phase protonated but retained for deprotonated complexes.[1,2] Moreover, in the case of a chemioradiotherapy to cure cancer, drugs can interact with ionizing radiation. Therefore, it is crucial to know if molecular recognition is affected by irradiation. We have thus studied photoionization of isolated protonated as well as deprotonated vancomycin/receptor complexes by means of a synchrotron beamline coupled to a tandem mass spectrometer, which allowed identifying the irradiation products. After absorption of one VUV photon by doubly-protonated Vancomycin, we observe non-dissociative ionization (NDI), but also peptide backbone fragmentation as well as sugar bond cleavages due to electronic to vibrational energy transfer. The latter rises with photon energy, making NDI to vanish at X-ray energies. For the doubly protonated vancomycin-receptor complex, NDI is not observed following photoabsorption, probably because of the low binding energy of this noncovalent complex. Interestingly, when deprotonated, the intact complex is observed after electron detachment, together with fewer fragmen-

tation channels, the main ones being loss of receptor and CO<sub>2</sub>. The latter comes only from vancomycin, not from the receptor, which can be explained by the retained native structure of the deprotonated complex, where the receptor's CO<sub>2</sub> group is strongly bound to Vancomycin, even after electron detachment. Our results indicate that the geometrical structure of a drugreceptor complex has a large effect on the processes induced by photon irradiation in the VUV-X energy range. Furthermore, molecular recognition is not systematically lost after photoabsorption.




Figure 1. molecular surfaces of the native structure of vancomycin (white) and its receptor (purple) in a crystal (Protein Data Bank 1FVM).

#### References

- [1] Poully J C et al 2010 International Journal of Mass *Spectrometry* **297** 28–35
- Poully J C et al 2010 Phys. Chem. Chem. Phys. 12

<sup>\*</sup> E-mail: marwa.abdelmouleh@ganil.fr

Content from this work may be used under the terms of the . Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. 1