



HAL
open science

Lipid-mediated regulation of flowering time

Yvon Jaillais, François Parcy

► **To cite this version:**

Yvon Jaillais, François Parcy. Lipid-mediated regulation of flowering time. *Science*, 2021, 373 (6559), pp.1086-1087. 10.1126/science.abl4883. hal-03349739

HAL Id: hal-03349739

<https://hal.science/hal-03349739v1>

Submitted on 20 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lipid-mediated regulation of flowering time

An anionic phospholipid tells plants when to flower based on ambient temperature

By Yvon Jaillais¹ and François Parcy²

¹Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, CNRS, INRAE, Lyon, France. ²Laboratoire Physiologie Cellulaire et Végétale, Univ. Grenoble Alpes, CNRS, CEA, INRAE, IRIGDBSCI-LPCV, Grenoble, France.

Email: yvon.jaillais@ens-lyon.fr

FLOWERING LOCUS T (FT) is a mobile protein, known as florigen, that induces flowering in plants. In *Arabidopsis thaliana*, FT transcription is induced in the leaf vascular tissues (phloem) when days become longer (1). It then moves at long distance through the phloem to reach the shoot apical meristem, where it induces floral transitions by regulating nuclear transcription complexes (1, 2). FT has homologies with characterized lipid-binding proteins in animals and belongs to the phosphatidylethanolamine-binding protein (PEBP) family (2). Although conserved throughout flowering plants, the function of FT lipid-binding capacity has remained enigmatic. On page XXX of this issue, Susila *et al.* (3) show that FT binds to the anionic phospholipid phosphatidylglycerol, which sequesters FT in membrane compartments of phloem cells at low temperature, thereby delaying flowering in cold conditions. Thus, a temperature-dependent regulation of FT translocation by a membrane phospholipid tunes the plant response to ambient temperature.

Susila *et al.* found that FT also binds to the anionic phospholipid phosphatidylglycerol *in vitro* and in phloem cells *in vivo*. Anionic phospholipids modulate the physicochemical properties of membranes, notably their electrostatic field, and provide docking platforms for proteins (4). Each anionic phospholipid species localizes to precise compartments within plant cells, thereby regulating the membrane identity of these organelles (4). Phosphatidylglycerol mostly accumulates in chloroplast and mitochondria (5), but has also been found in various compartments of the endomembrane system (6). However, its precise localization is largely unknown, raising the question of where FT interacts with phosphatidylglycerol.

Susila *et al.* found that, in *A. thaliana* protoplasts, FT associates with various membranes that contain phosphatidylglycerol: plasma membrane, Golgi apparatus, trans-Golgi network, endoplasmic reticulum, and the chloroplast envelop. Low temperature (i.e. long day at 16°C) induced the interaction between FT and cellular membranes in a phosphatidylglycerol-dependent manner.

Differentiated phloem tissues consist of conducting cells, called the sieve elements, and specialized parenchymal cells, called the companion cells (see the figure). Sieve elements are enucleated cells responsible for transporting sugars, metabolites and some proteins throughout the plants. Companion cells are nucleated cells responsible for the metabolic functioning of sieve elements. They are closely associated with sieve elements via specialized cell-to-cell connections called plasmodesmata. FT is expressed in

1 phloem companion cells and then moves to the sieve elements via the cytoplasmic canal of the
2 plasmodesmata (1, 7). Low temperature promotes the phosphatidylglycerol-dependent retention of FT in
3 cellular membranes of companion cells, which delays flowering by reducing FT movement toward the sieve
4 elements and thus the shoot meristem. In phosphatidylglycerol biosynthetic mutants, this temperature-
5 dependent retention is reduced, leading to an early flowering phenotype and temperature insensitivity.
6 During long days at 23°C, FT is not sequestered within companion cell membranes and moves into phloem
7 sieve elements to reach the shoot apical meristem and accelerate flowering.
8

9
10
11 Once in the meristem, FT binds to another phospholipid, phosphatidylcholine, to induce flowering (5, 8).
12 FT preferentially interacts with phosphatidylcholine species with low unsaturation degree (i.e. few
13 numbers of carbon-carbon double bonds in their fatty acid chains), which are more abundant under long
14 day conditions. However, it remains unclear how ambient temperature modulates FT-
15 phosphatidylglycerol interaction and thus its mobility. Lipidomic analyses by Susila *et al.* suggest that the
16 overall amounts of phosphatidylglycerol do not vary with temperature. Yet, consistent with the intense
17 temperature-dependent remodeling of lipids in plants (6, 9, 10), the abundance of polyunsaturated
18 phosphatidylglycerol species increase at 16°C compared with 23°C. Since FT interaction with
19 phosphatidylcholine is dependent on its unsaturation degree (8), it could also be the case for FT-
20 phosphatidylglycerol interaction. As such, the membrane network, through changes in lipid unsaturation,
21 could catch or release PEBP proteins, thereby regulating plant architecture in response to environmental
22 cues, notably temperature variations. However, such a model will require further investigation.
23

24
25 A major step toward understanding how different phospholipid molecular species could regulate FT will
26 be to characterize this interaction at the atomic level. High-resolution crystal structures suggest possible
27 lipid binding sites (11), but the next step will be to analyze FT structure in complex with its lipid ligand or
28 even directly inserted into membranes. Dissecting this interaction should allow the identification of FT
29 variants specifically defective in lipid binding with intact transcriptional activity, thus allowing the
30 uncoupling of temperature sensing capacity from the flowering inducing functions of FT. It will be
31 particularly interesting to analyze whether such FT variants may already exist in nature, for example in
32 plant species with temperature-insensitive flowering transition.
33

34
35 The results of Susila *et al.* provide a likely explanation for why florigen retained its capacity to bind
36 membrane lipids when it functions as a transcriptional co-activator in the nucleus of meristem cells. Susila
37 *et al.* found that FT also localizes to TERMINAL FLOWER1 (TFL1)-containing vesicles (12). TFL1 is a
38 homolog of FT that belongs to the PEBP family (2, 7), and also binds to phosphatidylglycerol in vitro (3). It
39 acts as a flowering repressor and may be involved in vacuolar trafficking (12), as well as transcriptional
40 regulations (2). It will be interesting to analyze the importance of TFL1 lipid-binding and potential
41 temperature sensitivity in regulating meristem functions antagonistically with FT. Similarly, the potential
42 relevance of temperature-sensitive membrane interaction for members of the PEBP family involved in bud
43 release or tuber formation in other plants will be worth exploring (2, 7).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

REFERENCES AND NOTES

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

1. L. Corbesier *et al.*, *science* **316**, 1030-1033 (2007).
2. Y. Zhu, S. Klasfeld, D. Wagner, *Journal of experimental botany* **72**, 2301-2311 (2021).
3. H. Susila *et al.*, *Science* **xxx**, xxx (2021).
4. L. C. Noack, Y. Jaillais, *Annual review of plant biology* **71**, 71-102 (2020).
5. L. A. Colin, Y. Jaillais, *Current opinion in plant biology* **53**, 1-9 (2019).
6. M. Uemura, R. A. Joseph, P. L. Steponkus, *Plant physiology* **109**, 15-30 (1995).
7. S. Jin, Z. Nasim, H. Susila, J. H. Ahn, *Seminars in cell & developmental biology* **109**, 20-30 (2021).
8. Y. Nakamura *et al.*, *Nature communications* **5**, 3553 (2014).
9. D. L. Falcone, J. P. Ogas, C. R. Somerville, *BMC plant biology* **4**, 17 (2004).
10. Q. Li *et al.*, *The Plant cell* **27**, 86-103 (2015).
11. Y. Nakamura *et al.*, *iScience* **21**, 577-586 (2019).
12. E. J. Sohn *et al.*, *Proceedings of the National Academy of Sciences* **104**, 18801-18806 (2007).

AUTHORS DECLARE NO COMPETING INTEREST

FIGURE: Regulation of florigen localization and activity by lipids

Model for the regulation of FLOWERING LOCUS T (FT, also called florigen) localization and activity by the phospholipids phosphatidylglycerol and phosphatidylcholine. FT expression, localization, movement, and activity are regulated by ambient temperature and the photoperiod, which together influence flowering time.

