



A fresh trim provides a new look at the human mannose receptor

E Ficko-Blean, Gerald Hart

► To cite this version:

E Ficko-Blean, Gerald Hart. A fresh trim provides a new look at the human mannose receptor. Journal of Biological Chemistry, 2021, 297, 10.1016/j.jbc.2021.100922 . hal-03350142v1

HAL Id: hal-03350142

<https://hal.science/hal-03350142v1>

Submitted on 17 Nov 2021 (v1), last revised 21 Sep 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

A Fresh Trim Provides A New Look at The Human Mannose Receptor


Elizabeth Ficko-Blean

PII: S0021-9258(21)00722-5

DOI: <https://doi.org/10.1016/j.jbc.2021.100922>

Reference: JBC 100922

To appear in: *Journal of Biological Chemistry*



Please cite this article as: Ficko-Blean E, A Fresh Trim Provides A New Look at The Human Mannose Receptor, *Journal of Biological Chemistry* (2021), doi: <https://doi.org/10.1016/j.jbc.2021.100922>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 THE AUTHORS. Published by Elsevier Inc on behalf of American Society for Biochemistry and Molecular Biology.

A fresh trim provides a new look at the human mannose receptor

Running title: Fresh trim for the mannose receptor

Elizabeth Ficko-Blean

Station Biologique de Roscoff,
Laboratory of Integrative Biology of Marine Models (LBI2M)
CNRS and Sorbonne University
Place Georges Teissier
29688 Roscoff Cedex
France

ORCID: 0000-0002-3915-4715

Edited by Gerald Hart

Keywords: Mannose receptor, CRD4, lectin, glycan array, carbohydrate, protein-carbohydrate interactions, affinity, avidity, multivalent binding

Mini Abstract

The human mannose receptor plays an important role in scavenging a variety of glycans and glycoconjugates which contributes to both innate and adaptive immunity. However, the fine details of its ligand specificity, and specifically that of CRD4, the most functionally relevant C-type lectin domain within the receptor, are not completely understood. Feinberg *et al.* use glycan arrays, crystallography, and a newly trimmed version of CRD4 to elucidate the molecular mechanisms driving binding specificity. These data contribute to our molecular understanding of Ca^{2+} ion-mediated binding promiscuity in the human mannose receptor and the scavenging role of the receptor itself, and highlight unexpected interactions that should inspire further study.

Main Text

The type I transmembrane human mannose receptor CD206 is a C-type lectin found primarily on macrophages, immature dendritic cells, and liver sinusoidal endothelial cells. Its known role is to scavenge endogenous and foreign macromolecules through carbohydrate and protein recognition, and it is involved in both innate and adaptive immune responses. Ligands for the mannose receptor include endogenous glycoproteins, such as lysosomal enzymes, collagen, and pituitary hormones, as well as microbial surface glycan structures such as yeast mannans, mycobacterial mannose-capped lipopolysaccharides, and viral high mannose oligosaccharides (1). However, the molecular determinants driving carbohydrate recognition by the mannose receptor for its multitude of diverse carbohydrate ligands are not fully understood. New work by Feinberg *et al.* provides major advances on this front with high-resolution insight into the molecular determinants driving carbohydrate recognition across 15 crystal structures for one of the key carbohydrate-binding domains from this receptor (2).

The extracellular, ligand-binding portion of the mannose receptor is composed of three types of domains. The N-terminal domain is an R-type carbohydrate-recognition domain (CRD) that binds to non-reducing sulfo-*N*-acetyl-*D*-galactosamine moieties (3). The neighboring FnIII domain binds to collagen fragments (4). This is followed by eight C-type lectin-like domains. CRD4 and CRD5 account for most of

the binding activity and CRD4 is the only C-type lectin domain capable of independently binding glycans, suggesting that further investigations of CRD4 could shed light on mannose receptor recognition overall. C-type lectin domains like CRD4 bind sugar ligands mediated through Ca^{2+} coordination in their primary carbohydrate-binding sites. CRD4 is known to bind terminal mannose, fucose, N-acetylglucosamine, and glucose residues (5). However, the only crystal structure of CRD4, published over 20 years ago, demonstrated an altered domain structure that prevented the capture of biologically relevant ligand complexes for crystallographic analyses (6).

To circumvent this problem, Feinberg *et al.* first modified the mannose receptor domain to remove predicted flexible regions anticipated to enable domain swapping during crystallization (2). The resultant “trimmed CRD4” was purified on a mannose-Sepharose column, thus demonstrating carbohydrate binding activity. The construct was further biotinylated and incubated with either streptavidin-coated plates for solid phase competition binding assays or fluorescently-labelled streptavidin for screening on the Consortium for Functional Glycomics complex glycan array.

Glycan array screening revealed a fuller picture of the variety of glycans CRD4 is capable of binding. Nearly all the oligosaccharides giving a strong signal included a non-reducing D-mannose or L-fucose moiety (2). The identified D-mannose ligands are all substructures of a Man_9 high mannose oligosaccharide and the strongest D-mannose-related signals have terminal D-mannose- α -1,2-D-mannose motifs. The screen also revealed some unexpected results, such as binding of sulfated ligands and those that bear Lewis-a epitopes, but not those that contain Lewis-x.

A solid phase competition assay allowed the authors to delve more deeply into the glycan interactions. The data showed a Man_9 oligosaccharide binds with 40-fold higher affinity than a D-mannose monosaccharide (2). Testing of di- and tri-saccharides suggested that avidity could be responsible for the tighter binding of the branched ligands. Indeed, the experimental setup of the assay concentrates CRD4 on the surface of streptavidin coated plates, thus clustering the CRD4 binding sites together and providing a multivalent binding surface. A critical assessment to advance the understanding of the binding mechanism with branched ligands, and to distinguish between surface immobilization effects versus free CRD4, could be obtained by including a similar comparison using isothermal titration calorimetry. Future solution-based investigations of binding affinity will be important to determine whether these measurements are influenced by non-natural avidity and to provide a deeper understanding of the thermodynamic forces driving these biologically relevant protein-carbohydrate interactions.

Using X-ray crystallography, Feinberg *et al.* examined the molecular interactions driving the specificity and promiscuity of CRD4 (Figure 1) (2). High resolution crystal complexes unexpectedly demonstrate that the Ca^{2+} -coordinated glycans can interact in more than one orientation, even when binding the same monosaccharide in the primary binding site. D-mannose is coordinated in alternate orientations via its 3- and 4-OH groups (Figure 1A-C), methyl-N-acetyl- α -D-glucosamine is coordinated in alternate orientations via its 3- and 4-OH groups (Figure 1D,E), and L-fucose is coordinated via its 2- and 3-OH groups (Figure 1F,G) or in alternate orientations via its 3- and 4-OH groups (Figure 1H-J). Thus, this Ca^{2+} -mediated coordination is absolutely essential in CRD4 binding overall, as expected, but also in affording a binding site able to accommodate diverse carbohydrate ligands. How, then, does CRD4 maintain any specificity? The authors also provide insight into this question by identifying secondary interactions of CRD4 with complex carbohydrate ligands. Further investigation into more distal

secondary interactions with relevant glycans would also contribute to our understanding on complex carbohydrate binding specificity. Of particular interest for potential secondary interactions is a flexible, solvent-exposed hydrophobic residue (Tyrosine 723) on a surface loop of CRD4 near the primary binding site, which seems poised to play a role in binding.

Untangling the mechanisms and driving forces of interactions and the full biological components of mannose receptor ligands remains a priority and further structure and function studies will be helpful; for example, further screening of microbial glycans could reveal unexplored host-pathogen interactions. Moreover, structural studies of the conformations of the extracellular domain of the mannose receptor have revealed it undergoes pH-dependent conformational changes, taking on an extended conformation at basic pH and a more condensed conformation at acidic pH (7-9). A tantalizing comparison of the extracellular multidomain mannose receptor components in solution demonstrating avid binding on immobilized multivalent ligands (10) provides a suggestion for the next frontier of molecular studies. Examination of conformational changes upon ligand binding, such as has already been exploited to a certain extent with the mannose receptor (8), along with biophysical characterization of binding interactions using a solution-based method, will be valuable in appreciating how avid binding contributes to mannose receptor structure and function.

Feinberg *et al.* have successfully demonstrated that even within only the one CRD4 domain, mannose receptor interaction with its carbohydrate ligands is extremely complex. This high resolution structure/function study lends important new insight into the promiscuous power of Ca^{2+} coordination in biologically relevant protein-carbohydrate interactions. Furthermore, this contributes significantly towards our molecular understanding of C-type lectin receptors, particularly the scavenging role of the mannose receptor and its mechanism of interaction with both endogenous and foreign macromolecules.

Acknowledgments

The author would like to thank Dr. Mirjam Czjzek for scientific discussion and Dr. Catherine Goodman for scientific editing.

Conflict of interest

The author declares no conflict of interest.




Figure 1. CRD4 primary monosaccharide-binding site interactions demonstrating the promiscuous power of Ca^{2+} coordination (2). Carbons of the interacting monosaccharide hydroxyl groups are labelled adjacently for all the complexes. Ca^{2+} is shown as a dark sphere. Amino acids participating in the primary binding site are labelled in **A**. CRD4 is shown in complex with: **A**. Methyl- α -D-mannose (three complexes from pdb id 7JUB, 7JUC and 7JUD), **B**. D-mannose (white, dual occupancy, pdb id 7JUG) and D-mannose- α -1,6-D-mannose (rose, only the non-reducing dual occupancy monosaccharide could be modelled, pdb id 7JUH,), **C**. D-mannose- α -1,2-D-mannose (two complexes from pdb id 7JUE and 7JUF), **D**. Methyl-*N*-acetyl- α -D-glucosamine (pdb id 7L65), **E**. Methyl-*N*-acetyl- α -D-glucosamine (pdb id 7L66), **F**. L-fucose- α -1,3-*N*-acetyl-D-glucosamine (pdb id 7L67), **G**. L-fucose- α -1-2-D-galactose- β -1-4-D-glucose (pdb id 7L63), **H**. L-fucose- α -1-2-D-galactose- β -1-4-D-glucose (pdb id 7L62), **I**. L-fucose- α 1-2-D-galactose- β -1-4-D-glucose (pdb id 7L61) and **J**. the Lewis-A trisaccharide (pdb id 7L64).

1. Stahl, P. D., and Ezekowitz, R. A. (1998) The mannose receptor is a pattern recognition receptor involved in host defense. *Curr Opin Immunol* **10**, 50-55
2. Feinberg, H., Jegouzo, S. A. F., Lasanajak, Y., Smith, D. F., Drickamer, K., Weis, W. I., and Taylor, M. E. (2021) Structural analysis of carbohydrate binding by the macrophage mannose receptor CD206. *J Biol Chem*, 100368
3. Liu, Y., Misulovin, Z., and Bjorkman, P. J. (2001) The molecular mechanism of sulfated carbohydrate recognition by the cysteine-rich domain of mannose receptor. *Journal of molecular biology* **305**, 481-490
4. Napper, C. E., Drickamer, K., and Taylor, M. E. (2006) Collagen binding by the mannose receptor mediated through the fibronectin type II domain. *Biochem J* **395**, 579-586

5. Taylor, M. E., Bezouska, K., and Drickamer, K. (1992) Contribution to ligand binding by multiple carbohydrate-recognition domains in the macrophage mannose receptor. *J Biol Chem* **267**, 1719-1726
6. Feinberg, H., Park-Snyder, S., Kolatkar, A. R., Heise, C. T., Taylor, M. E., and Weis, W. I. (2000) Structure of a C-type carbohydrate recognition domain from the macrophage mannose receptor. *J Biol Chem* **275**, 21539-21548
7. Hu, Z., Wang, Y., Cheng, C., and He, Y. (2019) Structural basis of the pH-dependent conformational change of the N-terminal region of human mannose receptor/CD206. *J Struct Biol* **208**, 107384
8. Hu, Z., Shi, X., Yu, B., Li, N., Huang, Y., and He, Y. (2018) Structural Insights into the pH-Dependent Conformational Change and Collagen Recognition of the Human Mannose Receptor. *Structure* **26**, 60-71 e63
9. Napper, C. E., Dyson, M. H., and Taylor, M. E. (2001) An extended conformation of the macrophage mannose receptor. *J Biol Chem* **276**, 14759-14766
10. Taylor, M. E., and Drickamer, K. (1993) Structural requirements for high affinity binding of complex ligands by the macrophage mannose receptor. *J Biol Chem* **268**, 399-404

