



HAL
open science

Linking changes in the soil microbial community to C and N dynamics during crop residue decomposition

Cyrine Rezgui, Isabelle Trinsoutrot-Gattin, Marie Benoit, Karine Laval,
Wassila Riah-Anglet

► **To cite this version:**

Cyrine Rezgui, Isabelle Trinsoutrot-Gattin, Marie Benoit, Karine Laval, Wassila Riah-Anglet. Linking changes in the soil microbial community to C and N dynamics during crop residue decomposition. *Journal of Integrative Agriculture*, 2021, 20 (11), pp.3039 - 3059. 10.1016/s2095-3119(20)63567-5 . hal-03353784

HAL Id: hal-03353784

<https://hal.science/hal-03353784>

Submitted on 24 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Available online at www.sciencedirect.com

ScienceDirect



RESEARCH ARTICLE

Linking changes in the soil microbial community to C and N dynamics during crop residue decomposition



CrossMark

Cyrine REZGUI¹, Isabelle TRINSOUTROT-GATTIN¹, Marie BENOIT², Karine LAVAL¹, Wassila RIAH-ANGLET¹

¹ AGHYLE Research Unit, UniLaSalle, Rouen-Team, Mont-Saint Aignan 76134, France

² Agroecology and Environment Research Unit, Isara, AgroSchool for Life, Lyon 69007, France

Abstract

Crop residues are among the main inputs that allow the organic carbon (C) and nutrients to be maintained in agricultural soil. It is an important management strategy that can improve soil fertility and enhance agricultural productivity. This work aims to evaluate the extent of the changes that may occur in the soil heterotrophic microbial communities involved in organic matter decomposition and C and nitrogen (N) mineralization after the addition of crop residues. Soil microcosm experiments were performed at 28°C for 90 days with the addition of three crop residues with contrasting biochemical qualities: pea (*Pisum sativum* L.), rapeseed (*Brassica napus* L.), and wheat (*Triticum aestivum* L.). Enzyme activities, C and N mineralization, and bacterial and fungal biomasses were monitored, along with the bacterial and fungal community composition, by the high-throughput sequencing of 16S rRNA and ITS genes. The addition of crop residues caused decreases in β -glucosidase and arylamidase activities and simultaneous enhancement of the C mineralization and net N immobilization, which were linked to changes in the soil microbial communities. The addition of crop residues decreased the bacterial and fungal biomasses 90 days after treatment and there were shifts in bacterial and fungal diversity at the phyla, order, and genera levels. Some specific orders and genera were dependent on crop residue type. For example, Chloroflexales, Inquilinus, Rubricoccus, Clitocybe, and Verticillium were identified in soils with pea residues; whereas Thermoanaerobacterales, Thermacetogenum, and Hypoxylon were enriched in soils with rapeseed residues, and Halanaerobiales, Rubrobacter, and Volutella were only present in soils with wheat residues. The findings of this study suggest that soil C and N dynamics in the presence of the crop residues were driven by the selection of specific bacterial and fungal decomposers linked to the biochemical qualities of the crop residues. If crop residue decomposition processes showed specific bacterial and fungal operational taxonomic unit (OTU) signatures, this study also suggests a strong functional redundancy that exists among soil microbial communities.

Keywords: crop residues, C and N mineralization, enzyme activities, bacterial and fungal diversity

Received 11 June, 2020 Accepted 27 November, 2020

Cyrine REZGUI, E-mail: Cyrine.Rezgui@unilasalle.fr;
Correspondence Wassila RIAH-ANGLET, Tel: +33-2-32829198,
E-mail: Wassila.RIAH-ANGLET@unilasalle.fr

© 2021 CAAS. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
doi: 10.1016/S2095-3119(20)63567-5

1. Introduction

C and N transformations from soil organic matter (OM) are mainly (micro) biologically based. Indeed, soil organic materials are the trophic resources of heterotrophic soil organisms (Paterson *et al.* 2009). OM plays an important role in many ecosystem services provided by soils (Dominati

et al. 2010). Its degradation by soil organisms allows the turnover of nutrients (mainly C, N, phosphorus (P), and sulfur (S)), which is essential for plant growth and the durability of ecosystems. Crop residue inputs have been proposed as a sustainable and cost-efficient management tool in agricultural practices to preserve soil organic matter (Lehtinen *et al.* 2014; Arcand *et al.* 2016; Shahbaz *et al.* 2017). The long-term balance between soil C inputs through crop residues and losses *via* mineralization and oxidation determines the soil organic C content and nutrient cycling in the agrosystem (Ghimire *et al.* 2017).

Various crop residues often exhibit different physicochemical properties, thereby affecting the soil in different ways. Their biochemical compositions and physical structures affect mineralization (Prescott 2010). Some studies have demonstrated, under controlled or field conditions, that hemicellulose, lignin, and the physical size of crop residues, affect the C and N decomposition rates (Hadas *et al.* 2004; Abiven *et al.* 2005; Cayuela *et al.* 2009; Moreno-Cornejo *et al.* 2014). The soluble C content determines the initial rate of crop residue decomposition (Hadas *et al.* 2004), whereas the lignin content controls the medium- to long-term fate of added C (Trinsoutrot *et al.* 2000; Nicolardot *et al.* 2007). According to Ghimire *et al.* (2017), the addition of crop residues at a higher rate potentially increases the C mineralization. Moreover, a study conducted by Redin *et al.* (2014) showed that the application of 25 species of crop residues with a broad range of chemical compositions affected C mineralization.

In addition, some researchers have investigated the effect of incorporating the residues of different plant organs (roots, leaves, stems, etc.) on C and N mineralization (Trinsoutrot *et al.* 2000; Fang *et al.* 2007; Abera *et al.* 2012; Redin *et al.* 2014). As expected, these plant organs also affected soil N mineralization. Indeed, roots displayed significantly lower N mineralization compared to the shoots and leaves (Abbasi *et al.* 2014). The biochemical composition or quality parameters, such as total N concentration, lignin, cellulose, and hemicellulose contents, exert a significant influence on crop residue decomposition. In parallel, some ratios and indexes (such as C:N ratio, lignin:N ratio, and lignocellulosic index (LCI)) are considered useful indicators for estimating both decomposition and the N release of added residues (Vahdat *et al.* 2011; Abera *et al.* 2012; Abbasi *et al.* 2014).

The heterotrophic microbial communities control the coupling of C and N cycles in soils (Recous *et al.* 2017). Indeed, the diversity and the activity of these microbial communities, as well as the proportions of the major elements (C, N, P, S, etc.) in the crop residues, determine the partition between the C and the N that is mineralized or assimilated in soils (Chen *et al.* 2014). In this case, the relationship between the addition of crop residues and the

microbial communities has attracted increasing attention (Pascault *et al.* 2010a; Kriaučiūnienė *et al.* 2012; Lee *et al.* 2017; Fang *et al.* 2018; Sauvadet *et al.* 2018). Suleiman *et al.* (2018) indicated that the addition of wheat straw residues causes a change in the diversity and function of microbial communities. Moreover, the addition of different quantities of crop residues promotes the abundance of some bacterial groups (*Bacteroidetes* and *Proteobacteria*), while low-quality crop residues favor fungal abundance (Fang *et al.* 2018). Lian *et al.* (2019) showed that soybean residues significantly impact the bacterial community, resulting in *Bacteroidetes*, *Actinobacteria*, *Firmicutes*, and *Acidobacteria* being enriched at the initial stage of crop residue input.

Microbial communities in soils produce several enzyme activities which are involved in the decomposition of crop residues (Frasier *et al.* 2016; Lashermes *et al.* 2016; Hünninghaus *et al.* 2017; Luo *et al.* 2017; Zheng *et al.* 2018). Fang *et al.* (2019) showed a significant increase in several enzyme activities (β -glucosidase, cellobiohydrolase, xylosidase, N-acetyl-glucosaminidase, and aminopeptidase) with a marked increase at 10 days after the addition of wheat residues. These authors suggested that these increases in enzyme activities were linked to the stimulation of microbial growth and the abundance of some functional genes. In parallel, Sauvadet *et al.* (2016) showed that the relationship between the evolution of microbial communities and the production of enzymes during the decomposition of crop residues is related to residue qualities. Indeed, the study conducted by Amin *et al.* (2014) found that microorganisms exhibited an initial rapid growth in the presence of a high-quality litter and produced enzymes that are not efficient in degrading recalcitrant components. Meanwhile, in the presence of the more recalcitrant maize roots, microbial biomass grew more slowly but produced enzymes of higher efficiency. This high enzyme efficiency could be explained by the synergistic action of hydrolytic and oxidative enzymes, even in the early stages of decomposition.

This study is part of a larger program that aims to evaluate the impact of innovative crop rotations (pea–wheat–barley vs. rapeseed–wheat–barley) in the Normandy region of France. As different crops produce various residues, an understanding of residue degradation kinetics and its influence on soil health is of major interest to the evaluation of the crop systems. The objective of this paper is to evaluate whether the application of crop residues will affect the soil heterotrophic microbial community involved in OM decomposition and C and N mineralization. Although the degradation of plants of various quantities and qualities is well-known, its consequences on soil communities and their functions have received less attention. To better understand these effects, we have followed the evolution of microbial

communities in relation to C-N fluxes and enzyme activities at key stages in the decomposition of different qualities of crop residues, in the pedoclimatic context of Normandy (France). The focus was on monitoring the abundance and diversity of bacterial and fungal communities, enzyme activities, and C and N mineralization as a result of incorporating contrasting crop residues (legume and non-legume crop residues) harvested in the field.

2. Materials and methods

2.1. Soil sampling

The soil was collected from an arable field located in the Normandy region (France). This region has a temperate oceanic climate, with a mean annual rainfall ranging from 80–110 mm. The soil is sandy loam, classified as luvisol. This soil was chosen because of its characteristics, which are representative of the common pedological types of soil in Normandy. Soil samples were collected randomly from a depth of 0–20 cm from three points in each plot, using a soil auger. The soil analysis indicated that the soil texture was characterized by 6.56% clay, 68.10% silt, and 25.34% sand, with a mean pH (soil H₂O) of 7.6 in the topsoil. The average soil C and N contents were (17.14±3.82) and (1.27±0.07) g kg⁻¹ of dry soil, respectively. Before drying, 20 g of fresh sieved soil was used to determine water content by drying the soil at 105°C for 24 h.

2.2. Collection of crop residues

Three predominant plant species available on-farm were selected as part of the crop rotation. These were pea (*Pisum sativum* L.), rapeseed (*Brassica napus* L.), and wheat (*Triticum aestivum* L.). Crop residues (stem and leaf) were dried at room temperature and were then analyzed, milled, and passed through a 1-mm sieve. Triplicate samples of crop residues were taken and analyzed for their C, N, soluble (SOL) compounds, hemicellulose (HEM), cellulose (CEL), and lignin (LIG) concentrations. Total C and N contents of the residues were determined using an elemental analyzer (NA, 2000; Fisons Instruments, Milan, Italy). This technique is based on the quantitative “dynamic flash combustion” method.

To characterize the biochemical composition of decomposing residues, the SOL compounds of the dried residues were determined by hot water extraction (100°C) for 30 min, followed by extraction with a neutral detergent (100°C) for 60 min using the method described by Linères and Djakovitch (1993). The resulting non-extracted material was then successively subjected to selective extraction for

HEM and CEL, as described by Van Soest (1963). The mass of the residues was weighed after each step in the serial extraction, and the difference between the mass of the material subjected to a specific extraction and the mass of the residues accounted for that specific component. The mass of the final residual non-extractable fraction of the plant material was considered as LIG mass. From biochemical fractionation, the LCI was calculated as described by Melillo et al. (1989). It corresponds to the ratio between the content of crop residues in LIG and the sum of the contents of LIG, CEL, and HEM.

2.3. Laboratory incubations

Fresh soil, equivalent to 25 g of dried soil, was weighed and transferred into glass jars. Distilled water was added to the soil to achieve a soil moisture content of 80% of field capacity. The treatments comprised a control without the addition of crop residues and three crop-residue sources, i.e., pea (*P. sativum* L.), rapeseed (*B. napus* L.) and wheat (*T. aestivum* L.). Crop residues were weighed and added to the jars at a rate equivalent to 2‰ of organic C of the crop residue weight, which corresponded to 0.114, 0.109, and 0.106 g for pea, rapeseed, and wheat, respectively. Incubation experiments were conducted under constant C and non-limiting N conditions, according to the AFNOR Standard (AFNOR XP U44-163) (AFNOR 2009). Then, 25 g of dry soil mixed with crop residues was incubated in 2-L glass jars for CO₂ monitoring, and in 50-mL plastic containers for the determination of mineral N and microbiological parameters. These containers were placed in large plastic trays with a few small holes in the covers to allow for gas exchange.

A total of 81 jars corresponded to soils with added pea, rapeseed, and wheat residues (three treatments) and soils without crop residues (control)×9 incubation times (1, 3, 7, 14, 21, 28, 49, 70, and 90 days)×3 replicates for CO₂ monitoring. A total of 45 jars for mineral N monitoring (3 treatments+controls×5 incubation times (1, 7, 14, 49, and 90 days)×3 replicates) were used, and some of them were taken for the determination of microbiological parameters (3 treatments+controls×2 incubation times (1 and 90 days)×3 replicates). All jars were placed in a completely randomized design. The soil was then incubated under controlled conditions at 28°C. Soil moisture was checked/adjusted every two days by weighing the glass jars and adding the required amount of distilled water.

2.4. C and N mineralization measurements

The CO₂ produced by the soil was trapped in 10 mL of

0.5 mol L⁻¹ NaOH in a 40-mL flat-bottom flask. The traps were changed periodically to renew the atmosphere in the jars and prevent saturation of the NaOH. The CO₂ produced by the soil and trapped by NaOH was determined by conductimetry using a conductivity meter (Metler Toledo Seven Multi, Darmstadt, Germany)

The initial concentration of total mineral N on day 1 was determined by extracting soil samples with 100 mL of 1 mol L⁻¹ KCl added directly to the flask immediately after incorporation of the crop residues. Thereafter, mineral N was extracted by shaking for 1 h with 100 mL of 1 mol L⁻¹ KCl followed by filtration (Whatman No. 40). Extracts were stored at -20°C until analysis. The total mineral N of the extract was determined by using the Gallery Automated Photometric Analyser-Thermo (Thermo Fisher Scientific, Vantaa, Finland).

2.5. Total DNA extraction and quantification

Nucleic acids were extracted from 0.5 g of soil using a FastDNA SPIN Kit for soil (MP-Biomedicals, Santa Ana, CA, USA), according to the instructions of the manufacturer. Three replicates were performed for each treatment. DNA was quantified by a fluorimetric measurement using Hoeschst 33258 fluorochrome at 360/460 nm excitation/emission wavelengths, with a Fluorescent DNA Quantitation Kit (Biorad, Hercules, CA, USA) of dsDNA stored at -20°C (Gangneux *et al.* 2011). The DNA extracted from the soil corresponded to the total microbial biomass, and it was expressed as µg of extracted DNA per g of dry soil (µg g⁻¹ dry soil).

2.6. Real-time PCR amplification for bacterial and fungal biomass quantification

The 18S rRNA amplifications for fungal biomass estimation by 18S rDNA real-time qPCR were carried out with a total volume of 25 µL. The qPCR mix was prepared as follows: 5 ng of soil microbial DNA, 0.5 µmol L⁻¹ of each primer (FU18S1 5'-GGAAACTCACCAGGTCCAGA-3' and Nu-SSU-1536 5'-ATTGCAATGCYCTATCCCCA-3') (Borneman and Hartin 2000), 25 µL of absolute qPCR SYBR Green Mix (Roche, Basel, Switzerland), and 0.25 mg mL⁻¹ BSA (GeneOn Bioscience, Germany). Standard curves were obtained using serial dilutions of linearized plasmids containing the cloned 18S rRNA gene of *Fusarium graminearum*, and the results are expressed as 18S rRNA gene copy number per gram of dry soil. The amplification protocol (40 cycles of PCR, 20 s at 95°C, 30 s at 62°C, and 30 s at 72°C) was performed using the LightCycler 480 real-time PCR system (Roche, Basel, Switzerland). The

efficiency of the qPCR ranged from 95–99%.

The 16S rRNA amplifications for bacterial biomass estimation by 16S rRNA real-time PCR were carried out under the same conditions as the 18S rDNA PCR, except for the primers (63f5'-CAGGCCTAACACATGCAAGTC-3' (Marchesi *et al.* 1998) and BU16S4 5'-CTGCTGCCTCCCGTAGG-3') derived from 341F (Muyzer *et al.* 1993). The efficiency of the qPCR ranged from 98–100%. Standard curves were obtained using serial dilutions of linearized plasmids containing the cloned 16S rRNA genes from *Pseudomonas aeruginosa*, and the results are expressed as 16S rRNA gene copy number per gram of dry soil. Two independent qPCR assays were performed for the bacterial and fungal dsDNA estimations. Three replicates were performed for each treatment.

2.7. High-throughput sequencing of the bacterial 16S rRNA and fungal ITS genes

Sixteen samples of DNA extracted from soil (three soils with added residues and the control soil without residues) × 2 replicates × 2 sampling times: T1 and T90 were analyzed for the bacterial 16S rRNA gene and for the ITS region of fungal gene amplicons by high-throughput sequencing using the Illumina MiSeq. For each sample, DNA extracted from soils was used for amplification of the V4–V5 hypervariable region of bacterial 16S rRNA gene (Claesson *et al.* 2010) and ITS2 of fungal genes using an optimized and standardized amplicon-library preparation protocol (Metabio®[®], GenoScreen, Lille, France). An internal bacterial (artificial bacterial community comprising 17 different bacteria (ABCv2)) and fungal (artificial fungal community comprising 11 different fungi (AFC)) positive control and a negative (sterile water) control were also performed. Briefly, PCR reactions were performed using 5 ng of genomic DNA and 192 fusion barcoded primers (at 0.2 µmol L⁻¹ final concentrations) with an annealing temperature of 50°C for 30 cycles. PCR products were purified using Agencourt AMPure XP magnetic beads (Beckman Coulter, Brea, CA, United States), quantified according to GenoScreen's protocol, and mixed in an equimolar amount. Sequencing was performed using 250-bp paired-end for bacteria and 300-bp paired-end for fungi-sequencing chemistry on the Illumina MiSeq platform (Illumina, San Diego, CA, United States) at GenoScreen (Lille, France).

The raw paired-end reads (495 447 for bacteria and 1 465 138 for fungi) were demultiplexed per sample and subjected to the following steps: (1) search and removal of both forward and reverse primers using CutAdapt with no mismatches allowed in the primer sequences; (2) quality-

filtering using the PRINSEQ-lite PERL script (Schmieder and Edwards 2011) by truncating bases at the 3' end with Phred quality score < 30; (3) paired-end read assembly using FLASH (Magoč and Salzberg 2011) with a minimum overlap of 30 bases and >97% overlap identity. Taxonomic and diversity analyses were performed with the Metabiote Online ver. 2.0 pipeline (GenoScreen, Lille, France), which is based in part on QIIME Software ver. 1.9.1 (Caporaso *et al.* 2010). Following pre-processing, full-length 16S rRNA gene sequences were checked for chimera sequences (in-house method based on Usearch 6.1). For both bacteria and fungi, similar sequences with a nucleic identity-defined threshold (97% identity for an affiliation at the genus level on the V4–V5 regions of the 16S rRNA gene and the ITS regions, respectively) were clustered with Uclust ver. 1.2.22q (Edgar 2010) through an open-reference OTU-picking process and complete-linkage method, finally generating groups of sequences, or "Operational Taxonomic Units." An OTU cleaning step, involving elimination of singletons, was performed. The most abundant sequence of each OTU was considered the reference sequence of its OTU and taxonomically compared to the Greengenes database. The taxonomy of each 16S rRNA and ITS gene sequence was analysed by the RDP classifier algorithm ver. 2.2 (Cole *et al.* 2014) against the database of Greengenes (version 13_8; www.greengenes.gov) for the 16S rRNA gene and UNITE7.0 (alpha version; <https://unite.ut.ee>) for the ITS. Alpha-diversity metrics (Chao1 index) within samples were computed using QIIME ver. 1.9.1.

2.8. Enzyme activities

Soil enzymes have been reported as useful soil-state bioindicators because they provide information on the soil's ability to perform biogeochemical reactions (Sinsabaugh and Shah 2012; Almeida *et al.* 2015). We have chosen to measure β -D-glucosidase (β -GLU; EC: 3.2.1.21), N-acetyl- β -glucosaminidase (NAG; EC: 3.2.1.30), and arylamidase (ARYLN; EC: 3.5.1.5) activities, which decompose different substrates with varying complexity (Sinsabaugh and Follsat Shah 2012). The GLU enzyme is related to the C cycle, acting in the cleavage of cellobiose into glucose molecules (Almeida *et al.* 2015). The ARYLN enzyme catalyses the hydrolysis of N-terminal amino acids from arylamides (Dodor and Tabatabai 2007). NAG enzymes catalyse the hydrolysis of chitin, which is important in C and N cycling in soils. Both ARYLN and NAG activities are associated with microbial N acquisition, and they play a major role in N mineralization in soils (Ekenler and Tabatabai 2004; Jian *et al.* 2016). Enzyme activity measurements were performed in 96-well microplates in triplicate according to the AFNOR Standard (AFNOR 20130:2018). Briefly, for β -GLU and

NAG activities, 4 g soil samples were mixed for 10 min at 250 r min⁻¹ with 25 mL water. Soil suspensions were incubated respectively with 0.05 mol L⁻¹ 4-nitrophenyl β -D-glucopyranoside (Sigma-Aldrich, Merck KGaA, Darmstadt, Germany) and 0.01 mol L⁻¹ 4-N-acetyl- β -D-glucosaminide (Sigma-Aldrich, Merck KGaA, Darmstadt, Germany). The reaction was stopped (with 0.5 mol L⁻¹ CaCl₂ and 0.1 mol L⁻¹ Tris at pH 12) after 1 h, each plate was centrifuged (5 min at 1500×g), and absorbance was measured on a microplate reader (Varioskan Flash-Thermo Scientific, Vantaa, Finland). The amounts of *p*-nitrophenol were obtained by measuring the absorbance at λ =405 nm, with comparison to calibration curves.

For ARYLN activity, 4 g soil samples were mixed for 10 min at 250 r min⁻¹ with 25 mL Trizma base (50 mmol L⁻¹, pH 7.5). Soil solutions were incubated with 0.008 mol L⁻¹ L-leucine β -naphthylamide hydrochloride (Sigma-Aldrich, Merck KGaA, Darmstadt, Germany). The β -naphthylamine produced was extracted with acidified ethanol and converted to an azo compound by reacting with *p*-dimethylaminocinnamaldehyde (DMCA). The amount of β -naphthylamine was obtained by measuring the absorbance at λ =540 nm, with comparison to calibration curves.

Three replicates were performed for each soil (with and without residues) at two sampling times corresponding to one day after incubation (T1) and 90 days after incubation (T90), for a total of 24 samples. Enzyme activities are expressed as nmol min⁻¹ g⁻¹ dry soil.

2.9. Statistical analysis

Statistical tests were carried out to compare the HEM, CEL, LIG, and SOL fractions between the three crop residues. Statistical tests were also carried out to evaluate the effects of the addition of crop residues on the total microbial biomass (dsDNA), the total bacterial and fungal biomasses (16S and 18S rRNA gene copy numbers), bacterial and fungal diversity (high-throughput sequencing), enzyme activities (β -GLU, NAG, and ARYLN), and C and N dynamics. The measured data were compared by Tukey's test with significant differences at 5% ($P < 0.05$). The diversity of bacterial and fungal communities was analyzed by a Venn diagram. Principal component analysis (PCA) was performed to compare crop residue effects on microbial parameters. The calculations of the correlations between the different parameters were carried out using a Spearman's rank correlation procedure, and the significance level of these correlations was tested using a pairwise *t*-test. All analyses were performed using R (The R Development Core Team 2019).

3. Results

3.1. Composition of crop residues

The C content of the residues ranged from 438 to 467 g kg⁻¹, while the N content ranged from 6.1 to 15.5 g kg⁻¹, respectively. Pea residues exhibited the highest N content, while wheat residues exhibited the highest C content. The biochemical characterization of crop residues showed significant differences among their compositions (Table 1). Pea residues exhibited the highest SOL fraction (37%) and the lowest content of HEM (2%). The LIG content varied between 6% in wheat residues and 12% in rapeseed residues, whereas the CEL content was equivalent in the three crop residues and corresponded to the most abundant component. For the three crop residues, the C:N ratio ranged from 28.1±3.06 to 76.6±4.60 (pea residues<rapeseed residues<wheat residues), while the LIG:N ratio varied from 1.24±0.17 to 0.50±0.50. LIG:N was the highest in rapeseed residues, while the lowest values were recorded in the pea residues. Regarding the LCI ratio,

rapeseed residues were the most recalcitrant compared to pea and wheat residues (Table 1).

3.2. C and N mineralization after the addition of crop residues

Soils without residues presented the lowest C mineralization (943 mg C-CO₂ kg⁻¹ dry soil) compared to soils with added crop residues. In each treatment, net C mineralization was calculated by subtracting mineralized C in the control soil from mineralized C in the soils with residues (Fig. 1-A). During the beginning of the incubation (until day 9), soils with pea residues exhibited the highest C mineralization (Fig. 1-B). Then (until day 90), net C mineralization was similar for soils amended with pea and rapeseed residues (2 200 mg C-CO₂ kg⁻¹ of dry soil) and significantly higher for soil with wheat residues (2 400 mg C-CO₂ kg⁻¹ of dry soil; Fig. 1-A).

Regarding the N dynamic, it was expressed as net cumulative mineralized N by subtracting the amount of mineral N recovered in control soils (Fig. 2). No matter

Table 1 Main characteristics of the three crop residues

	Fraction (%)				Index and ratio ¹⁾		
	Soluble	Lignin	Hemicellulose	Cellulose	LCI	C:N	Lignin:N
Pea residues	37.34±0.86 a	7.71±0.16 b	2.36±1.03 c	52.59±0.31 ns	0.12±0.01 b	28.33±3.06 c	0.50±0.05 c
Wheat residues	13.86±0.18 c	6.09±0.42 c	27.25±3.03 a	52.80±2.83 ns	0.07±0.01 c	76.88±4.60 a	1.00±0.05 b
Rapeseed residues	19.56±0.09 b	12.07±0.44 a	16.13±1.68 b	52.24±1.72 ns	0.15±0.01 a	46.90±6.60 b	1.24±0.17 a

¹⁾ LCI, lignocellulosic index (LCI=Ratio of lignin/(Lignin+Cellulose+Hemicellulose)).

Data are mean±SD (n=3). Letters (a, b, c) indicate significance in soluble, lignin, hemicellulose and cellulose fractions, index and ratios between the three crop residues at P<0.05 according to Tukey's test; ns indicates not significant.




Fig. 1 Net C mineralization in soil after the addition of crop residues after 90 days of incubation (A) and after 21 days of incubation (B). Data are mean±SD (n=3). * indicates statistical significance between soils with different crop residues at P<0.05 according to Tukey's test.

which crop residue is added, the amount of N in soil without residues is higher than the amount found in the soil with crop residues. Therefore, all the residues drove a net N immobilization during the 90 days of the experiment. Wheat residues showed the highest N immobilization (309 mg N kg⁻¹ of dry soil) compared to rapeseed (263 mg kg⁻¹ of dry soil) and pea residues (182 mg N kg⁻¹ of dry soil).

3.3. Enzyme activities after the addition of crop residues

The three tested enzymes did not respond in a similar way to the presence of crop residues (Fig. 3). The β-GLU activity was 1.4 times significantly higher in amended soils compared to control soils at the beginning of the experiment. Whereas NAG activities were 1.3 and 1.2 times significantly higher after pea and wheat residue addition, respectively, compared to the control. For ARYLN activity, no significant effect was observed after the addition of crop residues. At the end of the incubation, β-GLU activities decreased by about 6.71 and 7.28% following the addition of pea and rapeseed residues. The most marked decrease was observed for ARYLN activity, which decreased by about 38 and 35% after the addition of wheat and rapeseed residues. NAG activity was negatively correlated with LIG and LCI; however, no significant effect was observed on its activity after the addition of crop residues. Other measured enzyme activities were not correlated with the biochemical qualities of the crop residues.

3.4. Abundance of the total bacterial and fungal biomasses after the addition of crop residues

At the beginning of the incubation, the addition of crop




Fig. 2 Net N immobilization in soils under three crop residue additions at different incubation times. Data are mean±SD (n=3). * indicates statistical significance between soil with different crop residues at P<0.05 according to Tukey's test.

residues slightly increased total microbial biomass (assessed by the quantification of DNA extracted from soil) compared to the control (about 20%; Table 2). At the end of the incubation (T90), except for soils with pea residues that showed a slight decrease in microbial biomass, no difference was observed between soils amended with rapeseed or wheat residues and the control soil. A significant effect was observed between T1 and T90 on total microbial biomass.

Regarding bacterial biomass (Table 2), no crop residue effect was observed at the end of the incubation. However, a significant increase was observed between soils with added residues and the control at the beginning of the incubation (T1). For fungal biomass, various effects were observed. A significant decrease in soils with added pea and wheat residues and an increase in soils with rapeseed residues




Fig. 3 β-Glucosidase, arylamidase, and N-acetyl-glucosaminidase enzyme activities in soils with and without crop residues at two incubation times. T1, one day after incubation; T90, 90 days after incubation. Data are mean±SD (n=3). Letters (a, b and c) indicate statistical significance between soils with crop residues and soil without crop residues (control) for each sampling time at P<0.05 according to Tukey's test.

Table 2 The total microbial, bacterial and fungal biomasses from soil without residues (control) and soils with crop residues at two incubation times

	Incubation time ¹⁾	Total microbial biomass ($\mu\text{g g}^{-1}$ dry soil) ²⁾	Bacterial biomass (16S rDNA gene copy number g^{-1} dry soil ($\times 10^9$))	Fungal biomass (18S rDNA gene copy number g^{-1} dry soil ($\times 10^7$))
Control	T1	8.20 \pm 0.51 b A	2.08 \pm 0.14 b A	3.42 \pm 0.23 c A
	T90	6.96 \pm 0.33 a B	1.79 \pm 0.52 a A	1.18 \pm 0.11 b B
Pea residues	T1	10.66 \pm 0.82 a A	2.50 \pm 0.17 a A	11.1 \pm 0.65 a A
	T90	5.74 \pm 0.56 b B	1.53 \pm 0.25 a B	0.91 \pm 0.12 c B
Rapeseed residues	T1	9.50 \pm 0.63 ab A	2.69 \pm 0.33 a A	5.38 \pm 0.87 b A
	T90	6.26 \pm 1.09 ab B	1.77 \pm 0.12 a B	2.07 \pm 0.22 a B
Wheat residues	T1	9.90 \pm 0.80 ab A	2.95 \pm 0.30 a A	5.66 \pm 0.96 b A
	T90	6.88 \pm 1.10 ab B	1.82 \pm 0.11 a B	1.00 \pm 0.13 c B

¹⁾ T1, one day after incubation; T90, 90 days after incubation.

²⁾ The total microbial biomass was measured by the quantification of the total DNA extracted from the soil.

Data are mean \pm SD ($n=3$). Letters (a, b and c) indicate significant differences between soil without residues (control) and soils with crop residues. Letters (A and B) indicate significant differences between the two incubation times (T1 and T90) at $P<0.05$ according to Tukey's test.

was observed at the end of the incubation.

3.5. High-throughput sequencing analysis of the microbial communities after the addition of crop residues

For bacterial community composition, Miseq sequencing yielded a total of 427701 single reads, and their distribution ranged from a minimum of 24622 to a maximum of 35845 reads per sample of 16S rDNA. The rarefaction curves of the soil samples reached the plateau for all samples (Appendix A). The OTUs were assigned to 27 phyla, 70 orders, and 200 genera. When the abundance exceeds 1% (>1%), we have considered the phylum as dominant in the bacterial community, as reported by several authors (López *et al.* 2017; Wang *et al.* 2018; Deng *et al.* 2019). The abundant phyla are shown in Fig. 4-A. Among them, the most dominant phylum corresponded to Proteobacteria (48%), which included α -Proteobacteria (25%), δ -Proteobacteria (9%), β -Proteobacteria (7%), and γ -Proteobacteria (6%), followed by Actinobacteria (19%), Acidobacteria (8%), Bacteroidetes (8%), Chloroflexi (4%), Nitrospirae (3%), Gemmatimonadetes (2%), and Firmicutes (2%).

At the phylum level, Proteobacteria was the most abundant group regardless of the sampling time or the considered residues. Proteobacteria increased significantly between T1 and T90 in all treatments. Proteobacteria varied significantly between the amended soil with wheat residues (48%) and soils with pea and rapeseed residues (\approx 51%). At the end of the incubation, the most significant decrease of β -Proteobacteria was observed in the soil with wheat residues, while only a slight decrease for γ -Proteobacteria was observed in soil with pea residues. No significant difference was observed for δ -Proteobacteria after the addition of residues, whereas the abundance of α -Proteobacteria was significantly higher after the addition

of rapeseed residues. Actinobacteria and Bacteroidetes decreased significantly at T90, especially after the addition of pea and rapeseed residues, respectively. The relative abundances of Chloroflexi, Gemmatimonadetes, and Firmicutes increased about 1.5, 2, and 1.6 times in soils with wheat, pea, and rapeseed residues, respectively, in comparison to the control soil at the end of incubation (T90). The less abundant phyla (<1%) are shown in Fig. 4-B, and they are represented by Armatimonadetes, Elusimicrobia, Chlorobi, Fibrobacteres, GN02, OD1, WPS-2, BHI80-139, GAL15, OP11, WS2, and WS3. Among these phyla, Armatimonadetes, Chlorobi, GN02, GAL 15, WPS-2, and OD1 were, in most cases, significantly different between the control and the amended soils at the end of the incubation (T90). Indeed, the relative abundances of WPS-2, WS3, Armatimonadetes, and GN02 increased significantly compared to the control, while the abundances of Chlorobi, OP3, GAL 15, Fibrobacteres, and OD1 decreased. Overall, the diversity of the bacterial profile based on minor phyla showed similar profiles in soil amended with rapeseed residues and the control soil, and these profiles were different from those observed in soils amended with wheat or pea residues.

At the order level, five orders (Chloroflexales, Rhodothermales, Nitrososphaerales, Rickettsiales, and Ktedonobacterales) were identified after the addition of pea residues, mainly dominated by Rickettsiales (Fig. 5). Under rapeseed residues, four orders were identified, with the highest abundance in Thermoanaerobacterales OTUs. In soils with wheat residues, Halanaerobiales, Rubrobacterales, Planctomycetales, and Chthoniobacterales were identified. The bacterial orders were represented by a relatively equal number of OTUs. The number of genera-specific OTUs was higher than those identified for the order level after the addition of the three crop residues (Fig. 5). Indeed, 12




Fig. 4 The relative abundances of major bacterial phyla exceeding 1% (A) and minor bacterial phyla at less than 1% (B) in the soils with and without crop residues (control) at two incubation times. T1, one day after incubation; T90, 90 days after incubation. Data are mean \pm SD ($n=2$).

genera were identified under pea residues, with the highest abundance in *Inquilinus*. Under rapeseed residues, eight genera were observed with an average of equal abundance of OTUs. Seven genera were specifically identified after the addition of wheat residues, mostly represented by OTUs assigned to *Nocardioides*, *Aquicella*, and *Rubrobacter*.

For fungal community composition, Miseq sequencing yielded a total of 91 571 single reads, and the rarefaction curves of the soil samples reach the plateau for all samples (Appendix A). The OTUs were assigned to 13 fungal phyla, including Ascomycota, Basidiomycota, Mortierellomycota, Chytridiomycota, etc. The most dominant fungal phyla included Basidiomycota, Ascomycota, and Mortierellomycota (Fig. 6). These phyla are differentially

impacted by the addition of crop residues. The abundance of Basidiomycota and Ascomycota increased after the addition of crop residues. Basidiomycota was greater after the addition of rapeseed residues. In contrast, the relative abundances of Ascomycota identified in the study were significantly higher after the addition of wheat ($18.08\pm 2.24\%$ in control and $32.89\pm 0.78\%$ after the addition of wheat residues) and pea residues ($18.08\pm 2.24\%$ in control and $6.98\pm 1.22\%$ after the addition of pea residues). Mortierellomycota phyla decreased after the addition of the three crop residues and was significantly more important after wheat residues. Entomophthoromycota and Mucoromycota phyla were present only after the addition of pea and wheat residues, respectively. The addition of




Fig. 5 Venn diagram for bacterial diversity showing the unique and shared OTUs between soils with and without crop residues. OTUs were defined at 97% sequence similarity. Pie charts showing the abundances of unique OTUs at genera (A1, B1, C1 and D1) and order levels (A2, B2, C2 and D2) in soils without residues (A), with rapeseed residues (B), with wheat residues (C), and with pea residues (D).




Fig. 6 The relative abundances of fungal phyla in the soils with and without crop residues (control) at two incubation times. T1, one day after incubation; T90, 90 days after incubation. Data are mean±SD (n=2).

crop residues markedly enriched soils with Cercozoa; the abundance of this fungal group increased three times after the addition of pea residues and about two times after the addition of wheat and rapeseed residues.

At the order level, two orders (Chaetosphaeriales and Basidiobolales) were identified after the addition of pea residues, mainly dominated by Basidiobolales (Fig. 7). Under rapeseed residues, three orders were selected and

were represented by a relatively equal number of OTUs. In soils with wheat residues, six orders were selected, with the highest abundance in Malasseziales OTUs.

Among the 275 identified fungal genera, the number of genera-specific OTUs was higher than those identified for the order level under the addition of the three crop residues (Fig. 8). In fact, 20 genera were identified under pea residues, with the highest abundance in Basidiobolus. Fourteen and 15 genera were specifically identified after the addition of rapeseed and wheat residues, respectively.

3.6. The effects of the biochemical qualities of crop residues on the diversity of bacterial and fungal communities

The effects of crop residue biochemical qualities on the diversity of bacterial and fungal communities have been studied using a PCA (Fig. 8). The results showed that the diversity of soil bacterial and fungal communities is significantly different according to both the addition of crop residues (control vs. amended soil) and their biochemical characteristics (pea, wheat, and rapeseed). Based on the diversity of bacterial and fungal communities, the PCA showed three clusters, including control soils, soils with rapeseed or wheat residues and soil with pea residues

(Fig. 8). For bacterial and fungal diversities, the two axes of the PCA explain 65.8% of the total variance. The PCA clearly illustrates the link between the biochemical composition of crop residues and the microbial diversity at the genera and order levels. The PCA score plot revealed that the changes in bacterial and fungal diversity were more pronounced in soils with pea residues in comparison to soils with rapeseed and wheat residues. The specific bacterial genera (*Dongia*, *Collimonas*, *Candidatus Nitrososphaera*, *Inquillinus*) and fungal orders and genera (*Basidiobolales*, *Chaetosphaeriales*, *Verticillium*, *Clitocybe*, *Neosascochyta*) identified under soils amended with pea residues were linked to the soluble fraction and total N content of crop residues. While soils amended with rapeseed and wheat residues are not discriminated based on bacterial and fungal genera, some genera seem to be dependent on plant species. The addition of rapeseed residues affected *Niabella*, *Cellvibrio*, and *Hypomicrobium* bacterial genera and *Spiculogloeales*, *Microbotryomycetes_ord_Incertae_sedis*, *Xylariales* *Udeniozyma*, *Yurkovia*, and *Sordaria* fungal orders and genera, which were strongly linked to the C:N ratio, and to a lesser extent, to lignin and cellulose contents. For soil amended with wheat residues, *Flavobacterium*, *Rubrobacter*, and *Caldilinea* bacterial genera and *Leucosporidiales*, *Mucorales*, *Leucosporidium*,




Fig. 7 Venn diagram for fungal diversity showing the unique and shared OTUs between soils with and without crop residues. OTUs are defined at 97% sequence similarity. Pie charts showing the abundances of unique OTUs at genera (A1, B1, C1 and D1) and order levels (A2, B2, C2 and D2) in soils without residues (A), with rapeseed residues (B), with wheat residues (C), and with pea residues (D).




Fig. 8 Principal component analysis (PCA) of crop residue effect on bacterial and fungal diversity at the end of incubation (90 days). The circular arrow indicates the gradual changes in the bacterial and fungal diversity in relation to the initial biochemical composition of the crop residues. Bacterial diversity is designated in red color, fungal diversity designated in green color, and biochemical composition of crop residues is designated in yellow color.

and Rhizophydium genera fungal orders and genera were closely correlated with hemicellulose content. The circular arrow in Fig. 8 represents the gradual change in bacterial and fungal diversity, which should be linked to the initial biochemical composition of the crop residues. The distribution of bacterial and fungal diversity at genus and order levels highlighted by the PCAs are consistent with those described by the Venn diagram (Figs. 5 and 7).

4. Discussion

4.1. Relationship between composition of the crop residues and C mineralization

In this study, a higher rate of C mineralization at the beginning of the incubation (until day 9) was observed after the addition of pea residues, which presented the highest content of SOL compounds (37%) (Fig. 1-B). Bertrand *et al.* (2007) reported that labile compounds (SOL fraction) determine the decomposition rate of crop residues in the early stages. These results agree with those of our study since SOL fraction and net C mineralization (day 1) were significantly correlated ($r=0.74$, $P<0.05$; Appendix B). Concentrations of total sugars as soluble compounds have been considered to exert dominant control in the earlier stages of crop residue decomposition (Johansson *et al.*

2011). Some studies show that legume residues have the highest content of soluble sugars compared to wheat and rapeseed residues, which explains the highest level of C mineralization observed after the addition of pea residues (Hashem *et al.* 2013; Stewart *et al.* 2015). At the end of the incubation, a decline in mineralization rate was observed.

This observation had been previously explained by the increase in crop residue recalcitrance over time due to the greater proportion of remaining recalcitrant compounds at the end of decomposition (Šnajdr *et al.* 2011; Wickings *et al.* 2012). Some studies reported that at late stages, CEL, HEM, and LIG compounds, which are more recalcitrant, defined the decomposition of crop residues (Machinet *et al.* 2011; Šnajdr *et al.* 2011). In our study, however, no significant correlation was found between these recalcitrant fractions and the C mineralization rate (Appendix B).

Some researchers have shown a negative correlation between LIG fraction and C mineralization, and they explain that this is due to the resistant nature of LIG to the rapid microbial decomposition that promotes the formation of a complex phenyl-propanol structure, which often covers the cellulose-hemicellulose matrix (Stewart *et al.* 2015). Due to the complexity of this matrix, microbial substrate use efficiency for LIG is low because this efficiency decreases with increasing LIG content (Bahri *et al.* 2008; Cotrufo *et al.* 2013). The results of this study showed the highest C

mineralization rate in soils with wheat residues, which have the lowest amount of LIG (3%). Crop residues vary in their structural and chemical compositions (Adair *et al.* 2008), and their decomposition rates are generally negatively related to the number of recalcitrant compounds present in their biomass, such as lignin, phenols, and tannins (Bertrand *et al.* 2007; Castellano *et al.* 2015). In this case, crop residues with the lowest LIG content (less SGH-lignin content) had a higher C mineralization rate (Stewart *et al.* 2015).

4.2. Relationship between crop residue composition and N dynamics

Crop residue incorporation in soils is subject to biological degradation (Berg and McClaugherty 2008). During biological degradation, crop residue C is used in respiration by decomposers, which releases CO₂ and provides energy (Chen *et al.* 2009). With this energy source, microbial communities absorb N from the soil to promote their propagation. Therefore, the availability of mineral N in the soil can play an important role in the decomposition of crop residues (Chen *et al.* 2014). Its influence depends mainly on the N demand of the microorganisms. In this study, the monitoring of the mineralization kinetics was carried out under non-limiting N conditions so as to avoid a confounding effect of N availability (Recous *et al.* 1995; Trinsoutrot *et al.* 2000). In this case, it is expected that the N residue content will not significantly affect the decomposition rate of crop residues. This study's results showed a decrease in the rate of mineral N in the soil after the addition of crop residues. For each residue, a net N immobilization is observed for the entire duration of the incubation. It has been previously demonstrated that crop residues with low N (or with a high C:N ratio >20) lead to N immobilization in the early stages of decomposition (Jensen 1994; Johnson *et al.* 2007).

The results showed an important N immobilization in the soils, with wheat residues presenting the highest proportion between the immobilized N and the N residue content (1.68%). Moreover, a negative correlation was also observed between the rate of mineral N in soils with added residues and the C:N ratio of the crop residues ($r=-0.92$, $P<0.05$; Appendix B). Therefore, microbial activity and N immobilization are favored for residues with a higher rate of mineralized C, such as wheat residues (Hadas *et al.* 2004; Kriauciūnienė *et al.* 2012), due to the degradation of crop residues and the subsequent consumption of mineral N by the soil microbial communities.

4.3. Effect of crop residue addition on enzyme activities

In this study, the addition of crop residues induces an

immediate increase in hydrolytic enzyme activities (β -GLU and NAG) at the beginning of the experiment. This is consistent with results shown by Amin *et al.* (2014) who demonstrated that the addition of residues increases enzyme (cellulase, xylanase, and laccase) activities after only 10 min of incubation. Other authors have shown stimulation of some enzyme activities (β -glucosidase, cellobiohydrolase, xylosidase, phosphatase, N-acetylglucosaminidase) after the addition of wheat residues (Fang *et al.* 2018). This contribution stimulates the growth of starved microbial populations, and therefore, the rate of production of enzymes (Piotrowska and Wilczewsk 2012; Amin *et al.* 2014; Sauvadet *et al.* 2017). ARYLN, which is considered as an N acquiring enzyme (Dodor and Tabatabai 2007), did not change at the beginning of the incubation because of the non-limiting N conditions in the soils. At the end of the incubation, decreases of β -GLU and ARYLN activities were noted, and these two enzymes were strongly correlated with each other ($r=0.78$, $P<0.05$; Appendix B). Hydrolase enzymes are associated with the degradation of parietal polysaccharides (Cusack *et al.* 2010; Bell *et al.* 2014) and peaked at earlier stages of degradation after the addition of crop residues (Rinkes *et al.* 2011; Šnadjr *et al.* 2011). The decrease in these enzyme activities at the end of the experiment could be explained by several factors, including the depletion of less complex organic compounds (Geisseler *et al.* 2009; Piotrowska and Wilczewsk 2012) and the contribution of oxidative enzymes in the degradation of recalcitrant compounds (Piotrowska and Wilczewsk 2012) as demonstrated by the strong negative correlation between NAG activity and LCI ($r=-0.69$, $P<0.05$; Appendix B) and NAG activity and LIG ($r=-0.83$, $P<0.05$; Appendix B). In addition, the negative correlation between NAG activity and fungal biomass ($r=-0.80$, $P<0.05$; Appendix B) reinforces the previous findings (Geisseler *et al.* 2009; Kriauciūnienė *et al.* 2012; Piotrowska and Wilczewsk 2012) showing the large contribution of fungal oxidative enzymes in the degradation of crop residues at the end of incubation.

4.4. Effect of the addition of crop residues on the abundance of microbial communities

The addition of crop residues enhances total, bacterial, and fungal biomasses only at the beginning of the incubation. Several researchers have argued that the addition of crop residues induces the stimulation of total biomass by enhancing its growth activity, resulting in a significant increase in total biomass (Geisseler *et al.* 2009). Some researchers have shown the same findings as our study using different methods to analyze microbial biomass under the addition of wheat, pea, and rapeseed residues (Kriauciūnienė *et al.* 2012; Nguyen and Marschner 2017;

Sauvadet *et al.* 2017). The late stage of the residue's decomposition is marked by the colonization of crop residues by a population that slowly degrades the most recalcitrant fraction, such as LIG (Fontaine *et al.* 2003; Sauvadet *et al.* 2017). The correlations observed between fungal biomass and both LIG and LCI ($r=0.57$, $P<0.05$ and $r=0.72$, $P<0.05$; Appendix B) confirmed the findings in the literature showing the involvement of fungi in the degradation of recalcitrant compounds by using LIG as a C source (Boer *et al.* 2005; Kriauciūnienė *et al.* 2012). As suggested by the literature and as demonstrated by the significant correlations between LIG and some bacterial strains (*Cellvibrio*, *Pedomicrobium*, *Haliangium*, *Harbinensis*, *Flavobacterium*, etc.) (Appendix C), bacteria can also degrade LIG, sometimes more effectively than the fungal population (Perestelo *et al.* 1996; Vargasc-García *et al.* 2007). The observed increase in fungal biomass at the end of incubation after the addition of rapeseed residues, could be linked, as suggested by Berg *et al.* (2015), to the higher Mn content of rapeseed residues (Isikhuemhen *et al.* 2014), compared to pea and wheat residues. Indeed, Berg *et al.* (2015) showed a relationship between the crop residue decomposition rate and Mn concentrations, suggesting the stimulation of fungal Mn peroxidase enzymes (Hatakka and Hammel 2011).

4.5. Effect of the addition of crop residues on the taxonomic composition of bacterial communities

The addition of crop residues impacted the soil bacterial community by inducing a significant change in the abundance of some bacterial groups. The rapid change in the community observed at the early stages of residue degradation is explained by the modification of simple compounds (sugars, starch, etc.). The addition of crop residues enhanced copiotrophs (R-strategists) bacterial phyla, which are stimulated in C-rich environments (Fierer *et al.* 2007; Pascault *et al.* 2010b; Sauvadet *et al.* 2017; Suleiman *et al.* 2018; Zheng *et al.* 2018) and are mostly represented by β - and γ -Proteobacteria sub-classes (Acosta-Martínez *et al.* 2008; Verzeaux *et al.* 2016; Suleiman *et al.* 2018). In this study, these phyla increased because of substrate availability and their fast growth capacity (Aneja *et al.* 2006; Bernard *et al.* 2007; Fierer *et al.* 2007; Pascault *et al.* 2013; Trivedi *et al.* 2013; Tao *et al.* 2019). After 90 days, α - and δ -Proteobacteria, and Firmicutes increased after the addition of crop residues as shown in previous studies (Ramirez *et al.* 2012; Verzeaux *et al.* 2016; Lee *et al.* 2017; Suleiman *et al.* 2018). Because of the depletion of labile C compounds, these oligotrophs (K-strategists) phyla grow slowly in nutrient-poor environments (Pascault *et al.* 2010a; Sauvadet *et al.* 2017; Suleiman *et al.* 2018) and can use a wide range of recalcitrant C compounds (Gaillard *et al.*

2003). In addition, most of the bacteria affiliated with α - and δ -Proteobacteria, and Firmicutes can create cellulosomes, which are involved in the digestion of recalcitrant compounds (Ziganshin *et al.* 2013; Sacco *et al.* 2016; Lee *et al.* 2017). Actinobacteria and Bacteroidetes abundances decreased 90 days after the crop residue input into the soil. These results are in agreement with those of Suleiman *et al.* (2018) but in discrepancy with several studies describing Actinobacteria and Bacteroidetes as a bacterial group able to degrade complex substrates (Pascault *et al.* 2010b; Lee *et al.* 2017; Lian *et al.* 2019). However, the reduction in Actinobacteria may be related to the decrease of the major order Actinomycetales, which includes *Sporichtha* and *Nocardia* genera (Tao *et al.* 2019).

Minor phyla (Armatimonadetes, GN02, WPS-2, and OD1) increased in this study after the addition of crop residues, while Chlorobi decreased. Armatimonadetes phyla are present in most soils at relatively low abundances (below 1%) as shown in this study (Deng *et al.* 2019). Based on these limited phenotypic data, one can only reasonably note that all the currently available Armatimonadetes groups are oligotrophs (Lee *et al.* 2014). Regarding GN02 and OD1 bacteria, they are not culturable; this could explain the limited available information about their trophic state. Brown and Jumpponen (2015) highlighted that many of these rare bacteria are digestion-resistant and have a symbiotic nature with their environment. WPS-2 bacteria were described as dominant and positively correlated with the C:N ratio (Hermans *et al.* 2017), which disagrees with the results of this study (WPS-2 bacteria were negatively correlated with the C:N ratio, $r=-0.62$, $P<0.05$; Appendix C). Other OTUs that belonged to *Chlorobi* were abundant in the earlier stage of decomposition, but they were always found in very low proportions (Rui *et al.* 2009; Liu *et al.* 2016). Overall, our findings highlight the changes in bacterial community composition based on the turnover between copiotrophic and oligotrophic bacterial phyla.

At the order level, correlational analyses provided insight into the crop residue characteristics responsible for the observed selection in bacterial community composition. Among the orders that were most affected and appeared after the addition of each crop residue, the Halanaerobiales, Thermoanaerobacterales, Halanaerobiales, Actinomycetales, etc. (Appendix C) exhibited significant correlations with some crop residue compositions. These orders are poorly described in the literature. Thus, there remains a large amount of unexplained variation in the observed bacterial order composition as linked to their biochemical characteristics (Li *et al.* 2018). Various bacterial genera also showed significant correlations with crop residue composition (Appendix C). *Rubrobacter* has also been shown to degrade HEM ($r=0.73$, $P<0.05$; Appendix C) and

xylan and could, therefore, play a significant role in the degradation of these components (De Mandal *et al.* 2017).

The relative abundance of oligotrophic taxa, such as *Nocardioides* and *Niabella*, increased during the residue degradation (present at the end of the incubation at T90), indicating the abilities of these microorganisms to degrade polymers with different complexities, such as HEM and CEL (Appendix C) (Reddy *et al.* 2013; Kerdraon *et al.* 2019). The positive relationship between these genera and HEM suggested their direct or indirect involvement in the degradation of the HEM fraction, which may be related to their abundance after the addition of wheat residues, which presents the highest HEM content. Soils with pea residues are rich in genera like *Inquilinus*, *Dongia*, *Conexibacter*, *Collimonas*, and *Dactylosporangium*, which are involved in nutrient recycling and the degradation of crop residues. Among them, *Conexibacter* is highly correlated to SOL, TN, and the C:N ratio (Appendix C) and is described in literature as mainly associated with the degradation of litter with low C:N (Koceja *et al.* 2019). Except for *Conexibacter* and *Collimonas*, which are highly correlated with the SOL fraction ($P < 0.01$; Appendix C), the other specific genera are weakly correlated with the SOL fraction (Appendix C), but they are highly correlated among themselves and to *Collimonas* genus ($P < 0.01$; Appendix C). This may explain their presence after the addition of pea residues, which presents the highest SOL content. Some bacterial genera have been identified as CEL and LIG consumers, and *Cellvibrio* and *Flavobacterium* are included among them. These bacteria were positively correlated with LIG ($r = 0.71$, $P < 0.03$; Appendix C) and highly abundant under rapeseed residues, which presented the highest LIG content, confirming the previous findings (Zhang and Lueders 2017; Moreno-Espíndola *et al.* 2018).

4.6. Effect of the addition of crop residues on the taxonomic composition of fungal communities

In the present study, Ascomycota, Basidiomycota, and Mortierellomycota were the dominant phyla in all treatments, accounting for >18% of the fungi, which was much greater than the other phyla. This is consistent with results shown by Srour *et al.* (2020) and Zhang *et al.* (2020). Ascomycota are the main fungal decomposers in many soil ecosystems (Ma *et al.* 2013; Zhou *et al.* 2016). Ascomycota play a major role in recycling plant residues, and they might form symbioses with mycorrhizae or endophytes (Ivancevic and Karadelev 2013; Choudhary *et al.* 2018; Zhang *et al.* 2020). Ascomycota are the most diverse group of saprotrophic fungi (Xiong *et al.* 2014) and the key decomposers in agricultural soils, which expand in response to organic inputs (Ma *et al.* 2013; Wang *et al.* 2018). Basidiomycota, rather than

Ascomycota, dominates terrestrial ecosystems, probably due to relatively less data available for agricultural soils (Tederloo *et al.* 2014). Basidiomycota plays a particularly important role in degrading plant litter with high lignin content (Blackwood *et al.* 2007; Entwistle *et al.* 2018; Zhang *et al.* 2020). Our results showed that the relative abundance of Basidiomycota after the addition of rapeseed residues was significantly higher, indicating that rapeseed residues could provide a higher lignin content for the growth of Basidiomycota.

To further detail the effects of the addition of crop residues with contrasting biochemical characteristics on specific groups of soil fungi, important orders and genera were detected and appeared only after the addition of each crop residue. However, a variety of fungal genera showed significant correlations with crop residue compositions. For instance, *Clitocybe* and *Verticillium* present only after added pea residues had been shown to degrade carbohydrates (Fernando *et al.* 2016; Geethanjali and Jayashankar 2016). These specific genera are negatively correlated ($r = -0.78$, $P < 0.05$; Appendix D) to the C:N residue ratio. *Hypoxyylon* genus, known for its ability to degrade LIG, was detected only after the addition of rapeseed (which presented the highest LIG content) (López *et al.* 2017). These specific genera are positively correlated with LIG content ($r = 0.82$, $P < 0.05$; Appendix D). *Volutella* genera play an important role in HEM degradation (Masigol *et al.* 2019). The results of this study showed a positive correlation ($r = 0.76$, $P < 0.05$) between the abundance of these genera and HEM fraction (mostly present in wheat residue). This may explain their abundance after the addition of wheat residues. Finally, this study confirms that the residue biochemical qualities are the main drivers of the shift in bacterial and fungal communities during crop residue decomposition in soil.

The decomposition of crop residues is an essential process in cropping systems as they are the main exogenous source of C. Our results showed that the residue qualities have distinct effects on soil microbial communities and C and N dynamics. The literature reported that the crop residues with the highest labile fraction affect the bacterial pathway at the beginning of decomposition, and crop residues with the highest recalcitrant fraction affect the fungal pathway at the end of decomposition, which suggests that a specific function is linked to specific taxa. Our findings highlighted the concomitant presence of some bacterial and fungal groups throughout the decomposition of crop residues. Indeed, many bacterial and fungal genera and orders were co-correlated among them and with the same biochemical fraction of the crop residues. This finding highlights the strong functional redundancy of soil microbial communities and underlines the complex relationships between structure and function of soil microbial communities. Finally, the

correlations observed among fungal groups are more abundant and much stronger than those among bacterial groups, supporting the idea that potential interactions in fungal communities are stronger than those of bacterial communities in soils during crop residue decomposition.

5. Conclusion

This study complements previous knowledge about the effects of crop residue biochemical composition on microbial communities, C and N dynamics, and the soil microbiological state. It is well-known that crop residue inputs increase C mineralization and induce N immobilization in a range that is influenced by the crop residue's initial biochemical composition. The main hypothesis, which stated that these changes might be related to changes in the composition and biomass of soil heterotrophic microbial communities and enzyme activities, was confirmed. The findings of this study showed that the soil C and N dynamics in the presence of the crop residues were driven by the selection of specific bacterial and fungal decomposers linked to the initial biochemical qualities of crop residues.

These findings are of major concerns from the perspective of agroecology development. Indeed, crop residue management is a recognized practice that may help to maintain OM in soils and contribute to N management during the intercrop period. Moreover, it seems that crop residues drive the microbial community composition in soils. This is of great interest when agricultural practices need a decrease in chemical inputs and an improvement in soil biodiversity and its related functions, such as nutrient cycling.

Acknowledgements

The authors would like to thank François D'Hubert of the Chambre d'Agriculture of Normandy and the farmers of the association of Economic and Environmental Interest Group (EEIG). This research was funded by the Normandy region and was supported by the Vivepois Project (D16-12746), France.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Appendices associated with this paper are available on <http://www.ChinaAgriSci.com/V2/En/appendix.htm>

References

Abbasi K, Tahir M, Sabir N, Khurshid M. 2014. Impact of the addition of different plant residues on carbon-nitrogen

content and nitrogen mineralization-immobilization turnover in a soil incubated under laboratory conditions. *Solid Earth Discussions*, **6**, 197–205.

Abera G, Wolde-meskel E, Bakken L R. 2012. Carbon and nitrogen mineralization dynamics in different soils of the tropics amended with legume residues and contrasting soil moisture contents. *Biology and Fertility of Soils*, **48**, 51–66.

Abiven S, Recous S, Reyes-Gómez V, Oliver R. 2005. Mineralisation of C and N from root, stem and leaf residue in soil and role of their biochemical quality. *Biology and Fertility of Soils*, **42**, 119–128.

Acosta-Martínez V, Dowd S, Sun Y, Allen V. 2008. Tag-encoded pyrosequencing analysis of bacterial diversity in a single soil type as affected by management and land use. *Soil Biology and Biochemistry*, **40**, 2762–2770.

Adair E C, Parton W J, Grosso S J D, Silver W L, Harmon M E, Hall S A, Burke I C, Hart S C. 2008. Simple three-pool model accurately describes patterns of long-term litter decomposition in diverse climates. *Global Change Biology*, **14**, 2636–2660.

AFNOR (the French Standardisation Association). 2009. Norme XP U 44-163. Characterization of organic matter by potential mineralization of carbon and nitrogen. AFNOR, Official Distributor of Standards according to the decree of 16 June 2009. France. (in French)

Almeida F, Wolf J M, Casadevall A. 2015. Virulence-associated enzymes of *cryptococcus neoformans*. *Eukaryotic Cell*, **14**, 1173–1185.

Amin B A Z, Chabbert B, Moorhead D, Bertrand I. 2014. Impact of fine litter chemistry on lignocellulolytic enzyme efficiency during decomposition of maize leaf and root in soil. *Biogeochemistry*, **117**, 169–183.

Aneja M K, Sharma S, Fleischmann F, Stich S, Heller W, Bahnweg G, Munch J C, Schloter M. 2006. Microbial colonization of beech and spruce Litter — Influence of decomposition site and plant litter species on the diversity of microbial community. *Microbial Ecology*, **52**, 127–135.

Arcand M M, Helgason B L, Lemke R L. 2016. Microbial crop residue decomposition dynamics in organic and conventionally managed soils. *Applied Soil Ecology*, **107**, 347–359.

Bahri H, Rasse D P, Rumpel C, Dignac M F, Bardoux G, Mariotti A. 2008. Lignin degradation during a laboratory incubation followed by ¹³C isotope analysis. *Soil Biology and Biochemistry*, **40**, 1916–1922.

Bell C, Carrillo Y, Boot C M, Rocca J D, Pendall E, Wallenstein M D. 2014. Rhizosphere stoichiometry: Are C:N:P ratios of plants, soils, and enzymes conserved at the plant species-level? *New Phytologist*, **201**, 505–517.

Berg B, Kjønaas O J, Johansson M B, Erhagen B, Åkerblom S. 2015. Late stage pine litter decomposition: Relationship to litter N, Mn, and acid unhydrolyzable residue (AUR) concentrations and climatic factors. *Forest Ecology and Management*, **358**, 41–47.

Berg B, McClaugherty C. 2008. Initial litter chemical composition. In: *Plant Litter: Decomposition, Humus Formation, Carbon*

- Sequestration*. Springer, Berlin, Heidelberg. pp. 53–83.
- Bertrand I, Delfosse O, Mary B. 2007. Carbon and nitrogen mineralization in acidic, limed, and calcareous agricultural soils: Apparent and actual effects. *Soil Biology and Biochemistry*, **39**, 276–288.
- Bernard L, Mougél C, Maron P A, Nowak V, Lévêque J, Henault C, Haichar F Z, Marol C, Balesdent J, Gibiat F, Lemanceau P, Ranjard L. 2007. Dynamics and identification of soil microbial populations actively assimilating carbon from ¹³C-labelled wheat residue as estimated by DNA- and RNA-SIP techniques. *Environmental Microbiology*, **9**, 752–764.
- Blackwood C B, Waldrop M P, Zak D R, Sinsabaugh R L. 2007. Molecular analysis of fungal communities and laccase genes in decomposing litter reveals differences among forest types but no impact of nitrogen deposition. *Environmental Microbiology*, **9**, 1306–1316.
- Boer W, Folman L B, Summerbell R C, Boddy L. 2005. Living in a fungal world: Impact of fungi on soil bacterial niche development. *FEMS Microbiology Reviews*, **29**, 795–811.
- Borneman J, Hartin R J. 2000. PCR primers that amplify fungal rRNA genes from environmental samples. *Applied and Environmental Microbiology*, **66**, 4356–4360.
- Brown S P, Jumpponen A. 2015. Phylogenetic diversity analyses reveal disparity between fungal and bacterial communities during microbial primary succession. *Soil Biology and Biochemistry*, **89**, 52–60.
- Caporaso J G, Kuczynski J, Stombaugh J, Bittinger K, Bushman F D, Costello E K, Fierer N, Peña A G, Goodrich J K, Gordon J I, Huttley G A, Kelley S T, Knights D, Koenig J E, Ley R E, Lozupone C A, McDonald D, Muegge B D, Pirrung M, Reeder J, et al. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, **7**, 335–336.
- Castellano M J, Mueller K E, Oik D C, Sawyer J E, Six J. 2015. Integrating plant litter quality, soil organic matter stabilization, and the carbon saturation concept. *Global Change Biology*, **21**, 3200–3209.
- Cayuela M L, Sinicco T, Mondini C. 2009. Mineralization dynamics and biochemical properties during initial decomposition of plant and animal residue in soil. *Applied Soil Ecology*, **41**, 118–127.
- Chen B, Liu E, Tian Q, Yan C, Zhang Y. 2014. Soil nitrogen dynamics and crop residues. A review. *Agronomy for Sustainable Development*, **34**, 429–442.
- Chen H, Fan M, Billen N, Stahr K, Kuzyakov Y. 2009. Effect of land use types on decomposition of ¹⁴C-labelled maize residue (*Zea mays* L.). *European Journal of Soil Biology*, **45**, 123–130.
- Choudhary M, Sharma P C, Jat H S, McDonald A, Jat M L, Choudhary S, Garg N, Choudhary M, Sharma P C, Jat H S, McDonald A, Jat M L, Choudhary S, Garg N. 2018. Soil biological properties and fungal diversity under conservation agriculture in Indo-Gangetic Plains of India. *Journal of Soil Science and Plant Nutrition*, **18**, 1142–1156.
- Claesson M J, Wang Q, O'Sullivan O, Greene-Diniz R, Cole J R, Ross R P, O'Toole P W. 2010. Comparison of two next-generation sequencing technologies for resolving highly complex microbiota composition using tandem variable 16S rRNA gene regions. *Nucleic Acids Research*, **38**, e200.
- Cole J R, Wang Q, Fish J A, Chai B, McGarrell D M, Sun Y, Brown C T, Porras-Alfaro A, Kuske C R, Tiedje J M. 2014. Ribosomal Database Project: Data and tools for high throughput rRNA analysis. *Nucleic Acids Research*, **42**, D633–D642.
- Cotrufo M F, Wallenstein M D, Boot C M, Deneff K, Paul E. 2013. The microbial efficiency-matrix stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter? *Global Change Biology*, **19**, 988–995.
- Cusack D F, Torn M S, McDowell W H, Silver W L. 2010. The response of heterotrophic activity and carbon cycling to nitrogen additions and warming in two tropical soils. *Global Change Biology*, **16**, 2555–2572.
- Deng Y, Wang Y, Xia Y, Zhang A N, Zhao Y, Zhang T. 2019. Genomic resolution of bacterial populations in saccharin and cyclamate degradation. *Science of the Total Environment*, **658**, 357–366.
- Dodor D E, Tabatabai D M A. 2007. Arylamidase activity as an index of nitrogen mineralization in soils. *Communications in Soil Science and Plant Analysis*, **38**, 2197–2207.
- Dominati E, Patterson M, Mackay A. 2010. A framework for classifying and quantifying the natural capital and ecosystem services of soils. *Ecological Economics*, **69**, 1858–1868.
- Edgar R C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, **26**, 2460–2461.
- Ekenler M, Tabatabai M A. 2004. β-Glucosaminidase activity as an index of nitrogen mineralization in soils. *Communications in Soil Science and Plant Analysis*, **35**, 1081–1094.
- Entwistle E M, Zak D R, Argiroff W A. 2018. Anthropogenic N deposition increases soil C storage by reducing the relative abundance of lignolytic fungi. *Ecological Monographs*, **88**, 225–244.
- Fang M, Motavalli P P, Kremer R J, Nelson K A. 2007. Assessing changes in soil microbial communities and carbon mineralization in Bt and non-Bt corn residue-amended soils. *Applied Soil Ecology*, **37**, 150–160.
- Fang Y, Nazaries L, Singh B K, Singh B P. 2018. Microbial mechanisms of carbon priming effects revealed during the interaction of crop residues and nutrient inputs in contrasting soils. *Global Change Biology*, **24**, 2775–2790.
- Fang Y, Singh B P, Cowie A, Wang W, Arachchi M H, Wang H, Tavakkoli E. 2019. Balancing nutrient stoichiometry facilitates the fate of wheat residue carbon in physically defined soil organic matter fractions. *Geoderma*, **354**, 113–883.
- Fernando W G D, Zhang X, Amarasinghe C C. 2016. Detection of *Leptosphaeria maculans* and *Leptosphaeria biglobosa* causing blackleg disease in canola from Canadian canola seed lots and dockage. *Plants* (Basel), **5**, 12.
- Fierer N, Bradford M A, Jackson R B. 2007. Toward an

- ecological classification of soil bacteria. *Ecology*, **88**, 1354–1364.
- Fontaine S, Mariotti A, Abbadie L. 2003. The priming effect of organic matter: A question of microbial competition? *Soil Biology and Biochemistry*, **35**, 837–843.
- Frasier I, Noellemeyer E, Figuerola E, Erijman L, Permingeat H, Quiroga A. 2016. High quality residue from cover crops favor changes in microbial community and enhance C and N sequestration. *Global Ecology and Conservation*, **6**, 242–256.
- Gaillard V, Chenu C, Recous S. 2003. Carbon mineralization in soil adjacent to plant residue of contrasting biochemical quality. *Soil Biology and Biochemistry*, **35**, 93–99.
- Gangneux C, Akpa-Vinceslas M, Sauvage E, Desaire S, Houot S, Laval K. 2011. Fungal, bacterial and plant dsDNA contributions to soil total DNA extracted from silty soils under different farming practices: Relationships with chloroform-labile carbon. *Soil Biology and Biochemistry*, **43**, 431–437.
- Geethanjali P M, Jayashankar P M. 2016. A review on litter decomposition by soil fungal community. *IOSR Journal of Pharmacy and Biological Sciences*, **11**, 01–03.
- Geisseler D, Horwath W R, Doane T A. 2009. Significance of organic nitrogen uptake from plant residue by soil microorganisms as affected by carbon and nitrogen availability. *Soil Biology and Biochemistry*, **41**, 1281–1288.
- Ghimire B, Ghimire R, VanLeeuwen D, Mesbah A. 2017. Cover crop residues amount and quality effects on soil organic carbon mineralization. *Sustainability*, **9**, 23–16.
- Hadas A, Kautsky L, Goek M, Kara E E. 2004. Rates of decomposition of plant residue and available nitrogen in soil, related to residue composition through simulation of carbon and nitrogen turnover. *Soil Biology and Biochemistry*, **36**, 255–266.
- Hashem M, Ali E H, Abdel-Basset R. 2013. Recycling rice straw into biofuel. *Journal of Agriculture, Science and Technology*, **15**, 709–721.
- Hatakka A, Hammel K E. 2011. Fungal biodegradation of lignocelluloses. In: Hofrichter M, ed., *Industrial Applications, the Mycota*. Springer, Berlin, Heidelberg. pp. 319–340.
- Hermans S M, Buckley H L, Case B S, Curran-Cournane F, Taylor M, Lear G. 2017. Bacteria as emerging indicators of soil condition. *Applied Environmental Microbiology*, **83**, e02826-16.
- Hünninghaus M, Koller R, Kramer S, Marhan S, Kandeler E, Bonkowski M. 2017. Changes in bacterial community composition and soil respiration indicate rapid successions of protist grazers during mineralization of maize crop residues. *Pedobiologia*, **62**, 1–8.
- Isikhuemhen O S, Mikiashvili N A, Senwo Z N, Ohimain E I. 2014. Biodegradation and sugar release from canola plant biomass by selected white rot fungi. *Advances in Biological Chemistry*, **04**, 395.
- Ivancevic B, Karadelev M. 2013. Overview of fungi species in Prespa National Park (Albania). *International Journal of Ecosystems and Ecology Science*, **3**, 679–686.
- Jensen E S. 1994. Mineralization-immobilization of nitrogen in soil amended with low C/N ratio plant residue with different particle size. *Soil Biology and Biochemistry*, **26**, 519–521.
- Jian S, Li J, Chen J, Wang G, Mayes M A, Dzantor K E, Hui D, Luo Y. 2016. Soil extracellular enzyme activities, soil carbon and nitrogen storage under nitrogen fertilization: A meta-analysis. *Soil Biology and Biochemistry*, **101**, 32–43.
- Johansson M B, Berg B, Meentemeyer V. 2011. Litter mass-loss rates in late stages of decomposition in a climatic transect of pine forests. Long-term decomposition in a Scots pine forest. IX. *Canadian Journal of Botany*, **73**, 10.
- Johnson J M F, Barbour N W, Weyers S L. 2007. Chemical composition of crop biomass impacts its decomposition. *Soil Science Society of America Journal*, **71**, 155.
- Kerdraon L, Balesdent M H, Barret M, Laval V, Suffert F. 2019. Crop residues in wheat–oilseed rape rotation system: a pivotal, shifting platform for microbial meetings. *Microbial Ecology*, **77**, 931–945.
- Koceja M E, Bledsoe R B, Goodwillie C, Peralta A L. 2019. Nutrient enrichment increases soil bacterial diversity and decomposition rates of different litter types in a coastal plain wetland. *bioRxiv*, doi: 10.1101/732883.
- Kriauciūnienė Z, Velička R, Raudonius S. 2012. The influence of crop residues type on their decomposition rate in the soil: A litterbag study. *Zemdirbyste (Agriculture)*, **99**, 227–236.
- Lashermes G, Gainvors-Claissse A, Recous S, Bertrand I. 2016. Enzymatic strategies and carbon use efficiency of a litter-decomposing fungus grown on maize leaves, stems, and roots. *Frontiers in Microbiology*, **7**, 13–15.
- Lee J, Kim J R, Jeong S, Cho J, Kim J Y. 2017. Long-term performance of anaerobic digestion for crop residues containing heavy metals and response of microbial communities. *Waste Management*, **59**, 498–507.
- Lee K C, Dunfield P F, Stott M B. 2014. The phylum *Armatimonadete*. In: Rosenberg E, ed., *The Prokaryotes*. 4th ed. *Other Major Lineages of Bacteria and The Archaea* Springer-Verlag, Berlin. pp. 447–458.
- Lehtinen T, Schlatter N, Baumgarten A, Bechini L, Krüger J, Grignani C, Zavattaro L, Costamagna C, Spiegel H. 2014. Effect of crop residues incorporation on soil organic carbon and greenhouse gas emissions in European agricultural soils. *Soil Use and Management*, **30**, 524–538.
- Li Q, Chai S, Li Y, Huang J, Luo Y, Xiao L, Liu Z. 2018. biochemical components associated with microbial community shift during the pile-fermentation of primary dark tea. *Frontiers in Microbiology*, **9**, 1509.
- Lian T, Yu Z, Li Y, Jin J, Wang G, Liu X, Tang C, Franks A, Liu J, Liu J. 2019. The shift of bacterial community composition magnifies over time in response to different sources of soybean residue. *Applied Soil Ecology*, **136**, 163–167.
- Linères M, Djackovitch J L. 1993. Caractérisation de la stabilité biologique des apports organiques par l'analyse biochimique. In: Decroux J, Ignazi J C, eds., *Matières Organiques et Agricultures*. Quatrieme Journées de L'Analyse de Terre et Cinquieme Forum de la Fertilisation Raisonnée, GEMAS-COMIFER, Paris. pp. 159–168. (in French)

- Liu C, Li H, Zhang Y, Si D, Chen Q. 2016. Evolution of microbial community along with increasing solid concentration during high-solids anaerobic digestion of sewage sludge. *Bioresource Technology*, **216**, 87–94.
- López A M Q, Silva A L, Dos Santos E C L. 2017. The fungal ability for biobleaching/biopulping/bioremediation of lignin-like compounds of agro-industrial raw material. *Química Nova*, **40**, 916–931.
- Luo G, Ling N, Nannipieri P, Chen H, Raza W, Wang M, Guo S, Shen Q. 2017. Long-term fertilisation regimes affect the composition of the alkaline phosphomonoesterase encoding microbial community of a vertisol and its derivative soil fractions. *Biology and Fertility of Soils*, **53**, 375–388.
- Ma A, Zhuang X, Wu J, Cui M, Lv D, Liu C, Zhuang G. 2013. Ascomycota members dominate fungal communities during straw residue decomposition in Arable soil. *PLoS ONE*, **8**, e66146.
- Machinet G E, Bertrand I, Barrière Y, Chabbert B, Recous S. 2011. Impact of plant cell wall network on biodegradation in soil: Role of lignin composition and phenolic acids in roots from 16 maize genotypes. *Soil Biology and Biochemistry*, **43**, 1544–1552.
- Magoč T, Salzberg S L. 2011. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*, **27**, 2957–2963.
- De Mandal S, Chatterjee R, Kumar N S. 2017. Dominant bacterial phyla in caves and their predicted functional roles in C and N cycle. *BMC Microbiology*, **17**, 90.
- Marchesi J R, Sato T, Weightman A J, Martin T A, Fry J C, Hiom S J, Dymock D, Wade W G. 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied and Environmental Microbiology*, **64**, 795–799.
- Masigol H, Khodaparast S A, Woodhouse J N, Rojas-Jimenez K, Fonvielle J, Rezakhani F, Mostowfizadeh-Ghahamfarsa R, Neubauer D, Goldhammer T, Grossart H P. 2019. The contrasting roles of aquatic fungi and oomycetes in the degradation and transformation of polymeric organic matter. *Limnology and Oceanography*, **64**, 2662–2678.
- Melillo J M, Aber J D, Linkins A E, Ricca A, Fry B, Nadelhoffer K J. 1989. Carbon and nitrogen dynamics along the decay continuum: plant litter to soil organic matter. *Plant and Soil*, **115**, 189–198.
- Moreno-Cornejo J, Zornoza R, Faz A. 2014. Carbon and nitrogen mineralization during decomposition of crop residues in a calcareous soil. *Geoderma*, **230–231**, 58–63.
- Moreno-Espíndola I P, Ferrara-Guerrero M J, Luna-Guido M L, Ramírez-Villanueva D A, De León-Lorenzana A S, Gómez-Acata S, González-Terreros E, Ramírez-Barajas B, Navarro-Noya Y E, Sánchez-Rodríguez L M, Fuentes-Ponce M, Macedas-Jimenez J U, Dendooven L. 2018. The bacterial community structure and microbial activity in a traditional organic milpa farming system under different soil moisture conditions. *Frontiers in Microbiology*, **9**, 2737.
- Muyzer G, De Waal E C, Uitterlinden A C. 1993. Profiling of complex microbial populations by denaturing gradient gel-electrophoresis analysis of polymerase chain reaction-amplified genes-coding for 16S ribosomal-RNA. *Applied and Environmental Microbiology*, **59**, 695–700.
- Nguyen T T, Marschner P. 2017. Soil respiration, microbial biomass and nutrient availability in soil after addition of residue with adjusted N and P concentrations. *Pedosphere*, **27**, 76–85.
- Nicolardot B, Bouziri L, Bastian F, Ranjard L. 2007. A microcosm experiment to evaluate the influence of location and quality of plant residue on residue decomposition and genetic structure of soil microbial communities. *Soil Biology and Biochemistry*, **39**, 1631–1644.
- Pascual N, Cécillon L, Mathieu O, Hénault C, Sarr A, Lévêque J, Farcy P, Ranjard L, Maron P A. 2010a. *In situ* dynamics of microbial communities during decomposition of wheat, rape, and alfalfa residue. *Microbial Ecology*, **60**, 816–828.
- Pascual N, Nicolardot B, Bastian F, Thiébeau P, Ranjard L, Maron P A. 2010b. *In situ* dynamics and spatial heterogeneity of soil bacterial communities under different crop residues management. *Microbial Ecology*, **60**, 291–303.
- Pascual N, Ranjard L, Kaisermann A, Bachar D, Christen R, Terrat S, Mathieu O, Lévêque J, Mougél C, Hénault C, Lemanceau P, Péan M, Boiry S, Fontaine S, Maron P A. 2013. Stimulation of different functional groups of bacteria by various plant residue as a driver of soil priming effect. *Ecosystems*, **16**, 810–822.
- Paterson E, Midwood A J, Millard P. 2009. Through the eye of the needle: A review of isotope approaches to quantify microbial processes mediating soil carbon balance. *New Phytologist*, **184**, 19–33.
- Perestelo F, Rodríguez A, Pérez R, Carnicero A, Fuente G D I, Falcón M A. 1996. Short communication: Isolation of a bacterium capable of limited degradation of industrial and labelled, natural and synthetic lignins. *World Journal of Microbiology and Biotechnology*, **12**, 111–112.
- Piotrowska A, Wilczewski E. 2012. Effect of catch crop cultivated for green manure and mineral nitrogen fertilization on soil enzymes activities and chemical properties. *Geoderma*, **189–190**, 72–80.
- Prescott C E. 2010. Litter decomposition: What controls it and how can we alter it to sequester more carbon in forest soils? *Biogeochemistry*, **101**, 133–149.
- Ramirez K S, Craine J M, Fierer N. 2012. Consistent effects of nitrogen amendments on soil microbial communities and processes across biomes. *Global Change Biology*, **18**, 1918–1927.
- Recous S, Lashermes G, Bertrand I. 2017. Couplages et contrôles des cycles du carbone et de l'azote par les communautés microbiennes dans les sols cultivés. In: Quae S, ed., *Les Sols et la Vie Souterraine: Des Enjeux Majeurs en AgroEcologie*. Quae, Paris. p. 328. (in French)
- Recous S, Robin D, Darwis D, Mary B. 1995. Soil inorganic N availability: Effect on maize residue decomposition. *Soil Biology and Biochemistry*, **27**, 1529–1538.
- Reddy A P, Simmons C W, D'haeseleer P, Khudiyakov J, Burd H, Hadi M, Simmons B A, Singer S W, Thelen M P,

- Vandergheynst J S. 2013. Discovery of microorganisms and enzymes involved in high-solids decomposition of rice straw using metagenomic analyses. *PLoS ONE*, **8**, e77985.
- Redin M, Recous S, Aita C, Dietrich G, Skolaude A C, Ludke W H, Schmatz R, Giacomini S J. 2014. How the chemical composition and heterogeneity of crop residues mixtures decomposing at the soil surface affects C and N mineralization. *Soil Biology and Biochemistry*, **78**, 65–75.
- Rinkes Z L, Weintraub M N, DeForest J L, Moorhead D L. 2011. Microbial substrate preference and community dynamics during decomposition of *Acer saccharum*. *Fungal Ecology*, **4**, 396–407.
- Rui J, Peng J, Lu Y. 2009. Succession of bacterial populations during plant residue decomposition in rice field soil. *Applied and Environmental Microbiology*, **75**, 4879–4886.
- Sacco L P, Castellane T C L, Lopes E M, de Macedo Lemos E G, Alves L M C. 2016. Properties of polyhydroxyalkanoate granules and bioemulsifiers from *Pseudomonas* sp. and *Burkholderia* sp. Isolates growing on glucose. *Applied Biochemistry and Biotechnology*, **178**, 990–1001.
- Sauvadet M, Chauvat M, Brunet N, Bertrand I. 2017. Can changes in litter quality drive soil fauna structure and functions? *Soil Biology and Biochemistry*, **107**, 94–103.
- Sauvadet M, Chauvat M, Fanin N, Coulibaly S, Bertrand I. 2016. Comparing the effects of litter quantity and quality on soil biota structure and functioning: Application to a cultivated soil in Northern France. *Applied Soil Ecology*, **107**, 261–271.
- Sauvadet M, Lashermes G, Alavoine G, Recous S, Chauvat M, Maron P A, Bertrand I. 2018. High carbon use efficiency and low priming effect promote soil C stabilization under reduced tillage. *Soil Biology and Biochemistry*, **123**, 64–73.
- Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics*, **27**, 863–864.
- Shahbaz M, Kuzyakov Y, Sanaullah M, Heitkamp F, Zelenev V, Kumar A, Blagodatskaya E. 2017. Microbial decomposition of soil organic matter is mediated by quality and quantity of crop residues: Mechanisms and thresholds. *Biology and Fertility of Soils*, **53**, 287–301.
- Sinsabaugh R L, Follstad Shah J J. 2012. Ecoenzymatic stoichiometry and ecological theory. *Annual Review of Ecology, Evolution, and Systematics*, **43**, 313–343.
- Šnajdr J, Cajthaml T, Valášková V, Merhautová V, Petránková M, Spetz P, Leppänen K, Baldrian P. 2011. Transformation of *Quercus petraea* litter: successive changes in litter chemistry are reflected in differential enzyme activity and changes in the microbial community composition. *FEMS Microbiology Ecology*, **75**, 291–303.
- Van Soest P J. 1963. The use of detergents in the analysis of fibrous feeds. II. A rapid method for the determination of fiber and lignin. *Journal of the Association of Official Analytical Chemists*, **46**, 829–835.
- Srouf A Y, Ammar H A, Subedi A, Pimentel M, Cook R L, Bond J, Fakhoury A M. 2020. Microbial communities associated with long-term tillage and fertility treatments in a corn–soybean cropping system. *Frontiers in Microbiology*, **11**, 1363.
- Stewart C E, Moturi P, Follett R F, Halvorson A D. 2015. Lignin biochemistry and soil N determine crop residue decomposition and soil priming. *Biogeochemistry*, **124**, 335–351.
- Suleiman A K A, Lourenço K S, Pitombo L M, Mendes L W, Roesch L F W, Pijl A, Carmo J B, Cantarella H, Kuramae E E. 2018. Recycling organic residue in agriculture impacts soil-borne microbial community structure, function and N₂O emissions. *Science of the Total Environment*, **631–632**, 1089–1099.
- Tao K, Zhang X, Chen X, Liu X, Hu X, Yuan X. 2019. Response of soil bacterial community to bioaugmentation with a plant residue-immobilized bacterial consortium for crude oil removal. *Chemosphere*, **222**, 831–838.
- Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou N S, Wijesundera R, Ruiz L V, Vasco-Palacios A M, Thu P Q, Suija A, Smith M E, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Põldmaa K, Piepenbring M, et al. 2014. Global diversity and geography of soil fungi. *Science*, **346**, 1256688.
- Trinsoutrot I, Recous S, Bentz B, Linères M, Cheneby D, Nicolardot B. 2000. Biochemical quality of crop residues and carbon and nitrogen mineralization kinetics under nonlimiting nitrogen conditions. *Soil Science Society of America Journal*, **64**, 918–926.
- Trivedi P, Anderson I, Singh B. 2013. Microbial modulators of soil carbon storage: Integrating genomic and metabolic knowledge for global prediction. *Trends in Microbiology*, **21**, 641–651.
- Vahdat E, Nourbakhsh F, Basiri M. 2011. Lignin content of range plant residue controls N mineralization in soil. *European Journal of Soil Biology*, **47**, 243–246.
- Vargas-García M C, Suárez-Estrella F, López M J, Moreno J. 2007. *In vitro* studies on lignocellulose degradation by microbial strains isolated from composting processes. *International Biodeterioration and Biodegradation*, **59**, 322–328.
- Verzeaux J, Alahmad A, Habbib H, Nivelles E, Roger D, Lacoux J, Decocq G, Hirel B, Catterou M, Spicher F, Dubois F, Duclercq J, Tetu T. 2016. Cover crops prevent the deleterious effect of nitrogen fertilization on bacterial diversity by maintaining the carbon content of ploughed soil. *Geoderma*, **281**, 49–57.
- Wang H, Zeng Y, Guo C, Bao Y, Lu G, Reinfelder J R, Dang Z. 2018. Bacterial, archaeal, and fungal community responses to acid mine drainage-laden pollution in a rice paddy soil ecosystem. *Science of the Total Environment*, **616–617**, 107–116.
- Wickings K, Grandy A S, Reed S C, Cleveland C C. 2012. The origin of litter chemical complexity during decomposition. *Ecology Letters*, **15**, 1180–1188.
- Xiong J, Peng F, Sun H, Xue X, Chu H. 2014. Divergent responses of soil fungi functional groups to short-term warming. *Microbial Ecology*, **68**, 708–715.
- Zhang L, Lueders T. 2017. Micropredator niche differentiation between bulk soil and rhizosphere of an agricultural soil

- depends on bacterial prey. *FEMS Microbiology Ecology*, **93**, fix103.
- Zhang S, Wang Y, Sun L, Qiu C, Ding Y, Gu H, Wang L, Wang Z, Ding Z. 2020. Organic mulching positively regulates the soil microbial communities and ecosystem functions in tea plantation. *BMC Microbiology*, **20**, 103.
- Zheng W, Zhao Z, Gong Q, Zhai B, Li Z. 2018. Effects of cover crop in an apple orchard on microbial community composition, networks, and potential genes involved with degradation of crop residues in soil. *Biology and Fertility of Soils*, **54**, 743–759.
- Zhou J, Jiang X, Zhou B, Zhao B, Ma M, Guan D, Li J, Chen S, Cao F, Shen D, Qin J. 2016. Thirty-four years of nitrogen fertilization decreases fungal diversity and alters fungal community composition in black soil in Northeast China. *Soil Biology and Biochemistry*, **95**, 135–143.
- Ziganshin A M, Liebetrau J, Pröter J, Kleinstüber S. 2013. Microbial community structure and dynamics during anaerobic digestion of various agricultural waste materials. *Applied Microbiology and Biotechnology*, **97**, 5161–5174.

Executive Editor-in-Chief ZHANG Wei-li
Managing Editor SUN Lu-juan