

ESR/U-series dating Eemian human occupations of Northern France

Jean-Jacques Bahain, Samer Farkh, Christophe Falguères, Qingfeng Shao, Pierre Voinchet, Bassam Ghaleb, David Hérisson, Jean-Luc Locht, Nicole Limondin-Lozouet, Patrick Auguste, et al.

▶ To cite this version:

Jean-Jacques Bahain, Samer Farkh, Christophe Falguères, Qingfeng Shao, Pierre Voinchet, et al.. ESR/U-series dating Eemian human occupations of Northern France. 16th International Luminescence and Electron Spin Resonance Dating conference (LED2021), Sep 2021, Burgos, Spain. , 2021. hal-03354044

HAL Id: hal-03354044 https://hal.science/hal-03354044v1

Submitted on 24 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Introduction

Palaeolithic sites associated with the Eemian Interglacial (MIS 5e) are very rare in Northern France and their preservation is restricted to very specific geological contexts, in association with carbonated tufa (Caours) or peat deposits (Waziers) (Figure 1).

ESR/U-series dating method was hence applied on teeth recovered from different levels of the two sites, in the aim to evaluate the quality of the results obtained in very clearly chronologically constrained stratigraphic sequences. The results obtained will be displayed and discussed in this poster.

Material and method

ESR/U-series dating method was applied using the protocol of preparation and analyses described in Bahain et al. (2010).

Equivalent doses were determined using single exponential function. The dose rate was estimated from measurements by gamma-ray spectrometry, both in laboratory with a low background, high purity germanium detector and in situ using Canberra Inspector1000 spectrometer. Rn loss from each tissue was determined by combining alpha-ray and gamma-ray data (Bahain et al., 1992). The cosmic dose rate was estimated using the formula of Prescott and Hutton (1994).

The following parameters were also used in the age determination:

- a k-value of 0.13 ± 0.02 (Grün & Katzenberger-Apel, 1994);
- water content of sediments was measured for each dated level by drying it in oven at 40°C during one week; - water contents of 0 wt% in the enamel and 7 wt% in the dentine ;
- conversion factors of Adamiec and Aitken (1998);
- beta attenuations values in dental tissues were calculated using a Monte Carlo approach (Brennan et al., 1997).

Results

The ESR/U-series results are shown in Table 1. Several interesting points can be highlighted in these data.

							Та	ble 1. E	SR/U-s	series resu	lts obtaine	ed on tee	th recovered	d from tw	vo Eem	ian sites	of North	hern France
Site	Unit	Sample	Tissue	U (ppm)	234U/238U	²³⁰ Th/ ²³² Th	²³⁰ Th/ ²³⁴ U	²²² Rn/ ²³⁰ Th	Initial thickness (µm)	Rem. thickness internal side (μm)	Rem. thickness external side (µm)	^S Equivalent doses (Gy)	U-uptake parameter (p or n)	Internal D _a (μGy/an)	D _a ß (µGy/an)	D _a (γ+cosm) (μGy/an)	D _a (µGy/an)	ESR/U-series ages (US or <i>AU</i>) (ka)
		G 4 2005 01	dentine	8.235 ± 0.220	1.281 ± 0.032	318	1.342 ± 0.052	0.17	0(7 + 100	27 + 2	4 + 1	70.21 + 2.07	-0.0139 ± 0.0098	105 - 142	105 . 117	250 - 21	(20×10)	10(+ 27
Caours	N1	CA2005-01	enamel	0.332 ± 0.020	$_1.376\pm0.094$	12	0.927 ± 0.086	1.00	867 ± 108	27 ± 3	4±1	79.31 ± 3.07	-0.0123 ± 0.0037	185 ± 143	$185 \pm 11/$	239 ± 21	629 ± 186	120 ± 3 /
	N1	CA2005-02	dentine	6.560 ± 0.219	1.232 ± 0.044	90	1.419 ± 0.080	0.32	993 ± 124	50 ± 6	20 ± 2	67.53 ± 2.66	$+0.0390 \pm 0.1431$	(1)7	216 . 106	210 - 20	5 (7 . 201	110 + 42
			enamel	0.301 ± 0.017	1.247 ± 0.075	16	0.813 ± 0.078	0.55					-0.0128 ± 0.0051	41 ± 37	316 ± 196	210 ± 20	$56/\pm 201$	<i>119</i> ± 42
	N1	CA2005-03	dentine	10.270 ± 0.242	1.167 ± 0.026	133	1.150 ± 0.042	0.16	893 ± 119	59 ± 7	4 ± 1	53.39 ± 5.30	-0.0105 ± 0.0005		150 . 10			
			enamel	0.393 ± 0.029	1.063 ± 0.094	23	0.953 ± 0.102	1.00					-0.0088 ± 0.0005	121 ± 3	139 ± 10	169 ± 22	449 ± 24	119 ± 15
	N3	CA2005-04	dentine	13.771 ± 0.316	1.225 ± 0.021	67	1.135 ± 0.048	0.09	1014 ± 127	43 ± 5	17 ± 2	73.16 ± 2.90	-0.0104 ± 0.0002	116 ± 1	74 ± 3	395 ± 20	585 ± 20	<i>125</i> ± 5
			enamel	0.193 ± 0.012	1.222 ± 0.090	13	0.884 ± 0.087	1.00					-0.0077 ± 0.0001					
	N1	CA2007-01	dentine	5.281 ± 0.165	1.229 ± 0.014	113	1.008 ± 0.052	0.14	$\begin{array}{c} 14\\92\end{array} 957 \pm 120 \end{array}$	2 ± 0	29 ± 4	71.00 ± 10.62	-0.0102 ± 0.0008	59 ± 2	279 ± 21	210 ± 20	548 ± 29	<i>129</i> ± <i>23</i>
			enamel	0.338 ± 0.016	1.274 ± 0.066	8	0.923 ± 0.055	0.92					-0.0312 ± 0.0057					
Waziers	2	WBT-19	dentine	0.097 ± 0.001	1.609 ± 0.004	128	0.547 ± 0.003	1.00	1250 ± 156	38 ± 5	135 ± 17	58.08 ± 1.70	-0.0127 ± 0.0017	9 ± 22	48 ± 9	408 ± 20	465 ± 33	125 ± 8
			enamel	0.036 ± 0.001	1.467 ± 0.017	46	0.600 ± 0.009	1.00					-0.7551 ± 0.0608					
	2	WBT-22-1	dentine	0.028 ± 0.001	1.348 ± 0.005	16	0.655 ± 0.004	1.00					-0.8125 ± 0.0502					
			enamel	0.071 ± 0.001	$1.246 \pm 0.005.$	213	0.775 ± 0.004	1.00	1209 ± 151	70±9	84 ± 11	59.71 ± 1.03	-0.0106 ± 0.0010	21 ± 3	4 ± 42	408 ± 20	<i>433</i> ± <i>26</i>	<i>138</i> ± <i>8</i>
	2	WBT-22-2	dentine	0.015 ± 0.001	1.508 ± 0.007	145	0.555 ± 0.004	1.00	1113 ± 139	55 ± 7	39 ± 5	58.95 ± 1.66	-0.6443 ± 0.0608		60 40			
			enamel	0.009 ± 0.001	1.386 ± 0.011	30	0.672 ± 0.008	1.00					-0.9384 ± 0.0438	2 ± 20	69 ± 10	408 ± 20	479 ± 30	123 ± 7
	6	WBT-318	dentine	0.068 ± 0.001	1.276 ± 0.005	13	0.679 ± 0.004	1.00	$ \begin{array}{c} 00 \\ 00 \\ 00 \end{array} $ 883 ± 110	39 ± 5	75 ± 9	76.47 ± 2.72	-0.9016 ± 0.0479	20 ± 32	89 ± 12	470 ± 23	579 ± 41	132 ± 8
			enamel	0.070 ± 0.001	1.276 ± 0.005	344	0.687 ± 0.003	1.00					-0.9386 ± 0.0461					

For the Caours samples, the US-ESR model cannot be applied due to high 230Th/234U ratios higher to unity and relatively low equivalent doses, between 50 and 80 Gy. As a with very low uranium contents and often higher in the enamels than in the dentines. consequence, only the AU model was used to calculate ESR/U-series ages, indicating an initial U-uptake systematically followed by a U-leaching for quite all tissues of the five samples. This work demonstrates also the importance of the in situ dosimetry in the age calculation

Despite all these quite negative points, the obtained ages for N1 teeth are relatively US age of 132 ± 8 ka Given these very homogeneous ages for the two dated levels, a homogeneous, leading to a mean age for level 1 of 124 ± 10 ka ($\pm 2\sigma$), similar to the age weighted mean age of 129 ± 11 ka can be calculated from these four samples, confirming obtained previously by Bahain et al. (2010). The tooth analysed from level N3 provides the attribution of the deposit to the Eemian interglacial and even, in agreement with the an AU age of 129 ± 23 ka. Given these very homogeneous ages for the two dated levels, the attribution of the deposit to the Eemian interglacial is well confirmed with a mean age of 125 ± 9 ka.



16^h International Conference on Luminescence and Electron Spin Resonance Dating - 13 to17 September 2021

ESR/U-series dating Eemian human occupations of Northern France

Jean-Jacques BAHAIN^{1*}, Samer FARKH^{1, 2}, Christophe FALGUÈRES¹, Qingfeng SHAO³, Pierre VOINCHET¹, Bassam GHALEB⁴, David HERISSON⁵, Jean-Luc LOCHT^{6, 7}, Nicole LIMONDIN-LOZOUET⁷, Patrick AUGUSTE⁸, Julie DABKOWSKI⁷, Laurent DESCHODT^{7,9} & Pierre ANTOINE⁷

- ¹ UMR7194 HNHP, Muséum national d'histoire naturelle, Paris, France
- ² Lebanese University, Beirut, Lebanon
- ³ College of Geography Science, Nanjing Normal University, Nanjing, China
- ⁵ UMR7041 ArScAn, CNRS, Nanterre, France
- ⁶ INRAP, Glisy, France

Site presentation and sampling





The Caours site is located close to Abbeville in the Scardon Valley. Coarse fluvial deposits were here recovered by silts and a mm thick tufa sequence who fossilized several archeological levels containing abundant Middle Paleolithic industry and faunal remains associated with a forested temperate environment. U-series, TL and OSL dates constrain the tufa formation and Caours archeological level to ca 123 ka, allowing its clear attribution to Eemian interglacial stage (MIS 5e). Malacofauna study permits to place the climatic optimum of this stage in the lower part of the tufa formation.

Five teeth, four from archaeological level N1 and one from N3, were analyzed in the present work by ESR/U-series. Another tooth was previously dated by the same method of 124 ± 15 ka (Bahain et al., 2010).

In the Waziers teeth, we observe a very specific geochemical behavior for all the samples, Consequently the internal dose has a restricted impact in the age determination (<10%) compared to the external dose.

The equivalent doses are low but very homogeneous for the N2 teeth, , leading to a mean age for this layer of 128 ± 20 ka ($\pm 2\sigma$), while the DE value obtained for the N6 leads to a

Conclusions

The combined ESR/U-series analyses realized on teeth carried out from two Eemian sites of Northern France, Ca and Wasiers, demonstrate the reliability of the method to date Middle Paleolithic sites and, despite partic geochemical characteristics of part of the analyzed tissues, the validity of the used analytical protocols. This demonstrates in particular the importance of the in situ dosimetry in the age calculation.

At Caours, the obtained ESR/U-series ages are in excellent agreement with the other available geochronological derived from U-series and luminescence studies. They permit to calculate new mean age for the lower part o sequence $(124 \pm 8 \text{ ka for unit 5 (N1, n=7)} \text{ and } 127 \pm 15 \text{ ka for units 6-11 (including N3, n=6)}$. Associated to avai paleo-environmental and archaeological data, they make of this site one of the best known Middle Palaeol sequence of France.

For Waziers, the new established ESR/U-series chronology, with a mean age of 129 ± 11 ka, confirms the record Saalian Late Glacial deposits at the bottom of the sequence and of the subsequent Eemian interglacial stage, evider by malacological and palynological studies.

These results show the interest to integrate ESR/U-series dating analyses during the chronostratigraphical stud Middle Paleolithic sites when possible and demonstrate the necessity to involve geochronologists to the fieldworks excavations, since its earliest stages.

Acknowledgements

We would like to thank the Caours and Waziers excavation teams for their disponibility, patience and helpful discussions.

The present study was financially supported by the UMR7194 HNHP which allowed the funding of the field missions and analyses. the help of the PRC PaléHauts



⁴ GEOTOP, Université du Québec à Montréal, Montréal, Canada

Waziers site is located close to Douai, in the Scarpe river valley. Discovered during archeological survey operation in 2011 et 2013, it was excavated in 2014-2015. The stratigraphic sequence encloses coarse fluvial deposits recovered by an alternation of silts and peat levels including archaeological levels with Middle Palaeolithic lithic industries, then by sandy loess deposits. Malacofauna study indicates at the bottom of the sequence both open landscape and cold environement, then an interglacial temperate environment with first peat bog then swampy forest. The associated mammal fauna is typical from forested environment also (cervids, aurochs, beavers,...). Four teeth from Waziers were analyzed during the present work, three from N2 and one from N6.

The ESR and mobile gamma-ray spectrometers of the French National Museum of Natural History were bought with the financial su of the 'Sesame Île-de-France' program and the 'Région Centre' respectively.

	References
ours	Adamiec, G, Aitken, M, 1998. Dose-rate conversion factors: update. Ancient TL, 16, 37-50.
cular work	Bahain, JJ., Yokoyama, Y., Falguères, C., Sarcia, M.N., 1992. ESR dating of tooth enamel: a comparison with K-Ar dating. Quaternary Science reviews, 11, 245-250.
data f the lable lithic	Bahain JJ., Falguères C., Dolo JM., Antoine P., Auguste P., Limondin-Lozouet, N., Locht JL., Tuffreau A., 2010. ESR/U-series dating of teeth recovered from well-stratigraphically age-controlled sequences from Northern France. Quaternary Geochronology 5, 371-375.
rd of nced	Brennan, B.J., Rink, W.J., McGuirl, E.L., Schwarcz, H.P., Prestwich, W.V., 1997. Beta doses in tooth enamel by "One Group" theory and the Rosy ESR dating software. Radiation Measurements, 27, 307–314.
ly of and	Grün, R., Katzenberger-Apel, O., 1994. An alpha irradiator for ESR dating. Ancient TL 12, 35–38.
	Grün, R., Schwarcz, H.P., Chadam, J.M., 1988. ESR dating of tooth enamel: coupled correction for U-uptake and U-series disequilibrium. Nuclear Tracks and Radiation Measurements, 14, 237-241.
	Prescott, J.R., Hutton, J.T., 1994. Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. Radiation Measurements, 23, 497–500.
, with	Shao, Q., Bahain, J.J., Dolo, J.M., Falguères, C., 2014. Monte Carlo approach to calculate US-ESR ages and their uncertainties. Quaternary Geochronology 22, 99-106.
ıpport	Shao, Q., Bahain, JJ., Falguères, C., Dolo, JM., Garcia, T., 2012. A new U-uptake model for combined ESR/U-series dating of tooth enamel. Quaternary Geochronology, 10, 406-411.