



HAL
open science

ESR/U-series dating Eemian human occupations of Northern France

Jean-Jacques Bahain, Samer Farkh, Christophe Falguères, Qingfeng Shao, Pierre Voinchet, Bassam Ghaleb, David Hérison, Jean-Luc Locht, Nicole Limondin-Lozouet, Patrick Auguste, et al.

► **To cite this version:**

Jean-Jacques Bahain, Samer Farkh, Christophe Falguères, Qingfeng Shao, Pierre Voinchet, et al.. ESR/U-series dating Eemian human occupations of Northern France. 16th International Luminescence and Electron Spin Resonance Dating conference (LED2021), Sep 2021, Burgos, Spain. , 2021. hal-03354044

HAL Id: hal-03354044

<https://hal.science/hal-03354044>

Submitted on 24 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ESR/U-series dating Eemian human occupations of Northern France

Jean-Jacques BAHAIN^{1*}, Samer FARKH^{1, 2}, Christophe FALGUÈRES¹, Qingfeng SHAO³, Pierre VOINCHET¹, Bassam GHALEB⁴, David HÉRISSEON⁵, Jean-Luc LOCHT^{6, 7}, Nicole LIMONDIN-LOZOUET⁷, Patrick AUGUSTE⁸, Julie DABKOWSKI⁷, Laurent DESCHODT^{7, 9} & Pierre ANTOINE⁷

¹ UMR7194 HNHP, Muséum national d'histoire naturelle, Paris, France
² Lebanese University, Beirut, Lebanon
³ College of Geography Science, Nanjing Normal University, Nanjing, China

⁴ GEOTOP, Université du Québec à Montréal, Montréal, Canada
⁵ UMR7041 ArScAn, CNRS, Nanterre, France
⁶ INRAP, Glisy, France

⁷ UMR8591 LGP, CNRS, Meudon, France
⁸ UMR 8198 EEP, CNRS-USTL, Villeneuve d'Ascq, France
⁹ INRAP, Villeneuve d'Ascq, France

Introduction

Palaeolithic sites associated with the Eemian Interglacial (MIS 5e) are very rare in Northern France and their preservation is restricted to very specific geological contexts, in association with carbonated tufa (Caours) or peat deposits (Waziers) (Figure 1).

ESR/U-series dating method was hence applied on teeth recovered from different levels of the two sites, in the aim to evaluate the quality of the results obtained in very clearly chronologically constrained stratigraphic sequences. The results obtained will be displayed and discussed in this poster.


Figure 1 - Location of the studied sites in Northern France



Site presentation and sampling

Since 2003 for Caours and 2011 for Waziers, these two sites are studied by multidisciplinary teams including geologists, archaeologists, paleontologists, malacologists and geochronologists. The main results of these studies are summarized in Figure 2.

Figure 2 - Presentation of the Caours and Waziers Eemian sites in Northern France.



The Caours site is located close to Abbeville in the Scardon Valley. Coarse fluvial deposits were here recovered by silts and a mm thick tufa sequence who fossilized several archeological levels containing abundant Middle Paleolithic industry and faunal remains associated with a forested temperate environment. U-series, TL and OSL dates constrain the tufa formation and Caours archeological level to ca 123 ka, allowing its clear attribution to Eemian interglacial stage (MIS 5e). Malacofauna study permits to place the climatic optimum of this stage in the lower part of the tufa formation. Five teeth, four from archaeological level N1 and one from N3, were analyzed in the present work by ESR/U-series. Another tooth was previously dated by the same method of 124 ± 15 ka (Bahain et al., 2010).

Waziers site is located close to Douai, in the Scarpe river valley. Discovered during archeological survey operation in 2011 et 2013, it was excavated in 2014-2015. The stratigraphic sequence encloses coarse fluvial deposits recovered by an alternation of silts and peat levels including archaeological levels with Middle Paleolithic lithic industries, then by sandy loess deposits. Malacofauna study indicates at the bottom of the sequence both open landscape and cold environment, then an interglacial temperate environment with first peat bog then swampy forest. The associated mammal fauna is typical from forested environment also (cervids, aurochs, beavers,...). Four teeth from Waziers were analyzed during the present work, three from N2 and one from N6.

Material and method

ESR/U-series dating method was applied using the protocol of preparation and analyses described in Bahain et al. (2010).

Equivalent doses were determined using single exponential function. The dose rate was estimated from measurements by gamma-ray spectrometry, both in laboratory with a low background, high purity germanium detector and in situ using Canberra Inspector1000 spectrometer. Rn loss from each tissue was determined by combining alpha-ray and gamma-ray data (Bahain et al., 1992). The cosmic dose rate was estimated using the formula of Prescott and Hutton (1994).

The following parameters were also used in the age determination:

- a k-value of 0.13 ± 0.02 (Grün & Katzenberger-Apel, 1994);
- water content of sediments was measured for each dated level by drying it in oven at 40°C during one week;
- water contents of 0 wt% in the enamel and 7 wt% in the dentine;
- conversion factors of Adamiec and Aitken (1998);
- beta attenuations values in dental tissues were calculated using a Monte Carlo approach (Brennan et al., 1997).

Results

The ESR/U-series results are shown in Table 1. Several interesting points can be highlighted in these data.

Table 1. ESR/U-series results obtained on teeth recovered from two Eemian sites of Northern France

Site	Unit	Sample	Tissue	U (ppm)	²³⁴ Th/ ²³⁸ U	²³⁰ Th/ ²³² Th	²³⁰ Th/ ²³⁴ U	²²⁸ Rn/ ²³⁰ Th	Initial thickness (µm)	Rem. thickness internal side (µm)	Rem. thickness external side (µm)	Equivalent doses (Gy)	U-uptake parameter (p or n)	Internal D ₀ (µGy/ann)	D _{0β} (µGy/ann)	D _{0γ} (γ-cosm) (µGy/ann)	D ₀ (µGy/ann)	ESR/U-series ages (US or AU) (ka)	
Caours	N1	CA2005-01	dentine	8.235 ± 0.220	1.281 ± 0.032	318	1.342 ± 0.052	0.17	867 ± 108	27 ± 3	4 ± 1	79.31 ± 3.07	-0.0139 ± 0.0098	185 ± 143	185 ± 117	259 ± 21	629 ± 186	126 ± 37	
			enamel	0.332 ± 0.020	1.376 ± 0.094	12	0.927 ± 0.086	1.00	0.927 ± 0.086	1.00	90	1.419 ± 0.080	0.32	0.0123 ± 0.0037	-0.0123 ± 0.0037	41 ± 37	316 ± 196	210 ± 20	567 ± 201
	N1	CA2005-02	dentine	6.560 ± 0.219	1.232 ± 0.044	90	1.419 ± 0.080	0.32	993 ± 124	50 ± 6	20 ± 2	67.53 ± 2.66	+0.0390 ± 0.1431	-0.0128 ± 0.0051	121 ± 3	159 ± 10	169 ± 22	449 ± 24	119 ± 15
			enamel	0.301 ± 0.017	1.247 ± 0.075	16	0.813 ± 0.078	0.55	1.150 ± 0.042	0.16	133	1.020 ± 0.026	0.16	-0.0105 ± 0.0005	-0.0088 ± 0.0005	116 ± 1	74 ± 3	395 ± 20	585 ± 20
	N1	CA2005-03	dentine	10.270 ± 0.242	1.167 ± 0.026	133	1.150 ± 0.042	0.16	893 ± 119	59 ± 7	4 ± 1	53.39 ± 5.30	-0.0088 ± 0.0005	-0.0102 ± 0.0008	59 ± 2	279 ± 21	210 ± 20	548 ± 29	129 ± 23
			enamel	0.393 ± 0.029	1.063 ± 0.094	23	0.953 ± 0.102	1.00	1.135 ± 0.048	0.09	67	1.135 ± 0.048	0.09	-0.0104 ± 0.0002	-0.0077 ± 0.0001	9 ± 22	48 ± 9	408 ± 20	465 ± 33
	N3	CA2005-04	dentine	13.771 ± 0.316	1.225 ± 0.021	67	1.135 ± 0.048	0.09	1014 ± 127	43 ± 5	17 ± 2	73.16 ± 2.90	-0.0077 ± 0.0001	-0.0102 ± 0.0008	21 ± 3	4 ± 42	408 ± 20	433 ± 26	138 ± 8
			enamel	0.193 ± 0.012	1.222 ± 0.090	13	0.884 ± 0.087	1.00	1.008 ± 0.052	0.14	113	1.008 ± 0.052	0.14	-0.0102 ± 0.0008	-0.0102 ± 0.0008	2 ± 20	69 ± 10	408 ± 20	479 ± 30
	N1	CA2007-01	dentine	5.281 ± 0.165	1.229 ± 0.014	113	1.008 ± 0.052	0.14	957 ± 120	2 ± 0	29 ± 4	71.00 ± 10.62	-0.0312 ± 0.0057	-0.0127 ± 0.0017	20 ± 32	89 ± 12	470 ± 23	579 ± 41	132 ± 8
			enamel	0.338 ± 0.016	1.274 ± 0.066	8	0.923 ± 0.055	0.92	0.923 ± 0.055	0.92	8	0.923 ± 0.055	0.92	-0.0127 ± 0.0017	-0.0127 ± 0.0017	9 ± 22	48 ± 9	408 ± 20	465 ± 33
Waziers	2	WBT-19	dentine	0.097 ± 0.001	1.609 ± 0.004	128	0.547 ± 0.003	1.00	1250 ± 156	38 ± 5	135 ± 17	58.08 ± 1.70	-0.0127 ± 0.0017	9 ± 22	48 ± 9	408 ± 20	465 ± 33	125 ± 8	
			enamel	0.036 ± 0.001	1.467 ± 0.017	46	0.600 ± 0.009	1.00	0.600 ± 0.009	1.00	46	0.600 ± 0.009	1.00	-0.7551 ± 0.0608	-0.7551 ± 0.0608	2 ± 20	69 ± 10	408 ± 20	479 ± 30
	2	WBT-22-1	dentine	0.028 ± 0.001	1.348 ± 0.005	16	0.655 ± 0.004	1.00	1209 ± 151	70 ± 9	84 ± 11	59.71 ± 1.03	-0.8125 ± 0.0502	-0.8125 ± 0.0502	21 ± 3	4 ± 42	408 ± 20	433 ± 26	138 ± 8
			enamel	0.071 ± 0.001	1.246 ± 0.005	213	0.775 ± 0.004	1.00	0.775 ± 0.004	1.00	213	0.775 ± 0.004	1.00	-0.0106 ± 0.0010	-0.0106 ± 0.0010	21 ± 3	4 ± 42	408 ± 20	433 ± 26
	2	WBT-22-2	dentine	0.015 ± 0.001	1.508 ± 0.007	145	0.555 ± 0.004	1.00	1113 ± 139	55 ± 7	39 ± 5	58.95 ± 1.66	-0.6443 ± 0.0608	-0.6443 ± 0.0608	2 ± 20	69 ± 10	408 ± 20	479 ± 30	123 ± 7
			enamel	0.009 ± 0.001	1.386 ± 0.011	30	0.672 ± 0.008	1.00	0.672 ± 0.008	1.00	30	0.672 ± 0.008	1.00	-0.9384 ± 0.0438	-0.9384 ± 0.0438	20 ± 32	89 ± 12	470 ± 23	579 ± 41
6	WBT-318	dentine	0.068 ± 0.001	1.276 ± 0.005	13	0.679 ± 0.004	1.00	883 ± 110	39 ± 5	75 ± 9	76.47 ± 2.72	-0.9016 ± 0.0479	-0.9016 ± 0.0479	20 ± 32	89 ± 12	470 ± 23	579 ± 41	132 ± 8	
		enamel	0.070 ± 0.001	1.276 ± 0.005	344	0.687 ± 0.003	1.00	0.687 ± 0.003	1.00	344	0.687 ± 0.003	1.00	-0.9386 ± 0.0461	-0.9386 ± 0.0461	20 ± 32	89 ± 12	470 ± 23	579 ± 41	132 ± 8

For the Caours samples, the US-ESR model cannot be applied due to high ²³⁰Th/²³⁴U ratios higher to unity and relatively low equivalent doses, between 50 and 80 Gy. As a consequence, only the AU model was used to calculate ESR/U-series ages, indicating an initial U-uptake systematically followed by a U-leaching for quite all tissues of the five samples. This work demonstrates also the importance of the in situ dosimetry in the age calculation.

Despite all these quite negative points, the obtained ages for N1 teeth are relatively homogeneous, leading to a mean age for level 1 of 124 ± 10 ka (±2σ), similar to the age obtained previously by Bahain et al. (2010). The tooth analysed from level N3 provides an AU age of 129 ± 23 ka. Given these very homogeneous ages for the two dated levels, the attribution of the deposit to the Eemian interglacial is well confirmed with a mean age of 125 ± 9 ka.

In the Waziers teeth, we observe a very specific geochemical behavior for all the samples, with very low uranium contents and often higher in the enamels than in the dentines. Consequently the internal dose has a restricted impact in the age determination (<10%) compared to the external dose.

The equivalent doses are low but very homogeneous for the N2 teeth, leading to a mean age for this layer of 128 ± 20 ka (±2σ), while the DE value obtained for the N6 leads to a US age of 132 ± 8 ka. Given these very homogeneous ages for the two dated levels, a weighted mean age of 129 ± 11 ka can be calculated from these four samples, confirming the attribution of the deposit to the Eemian interglacial and even, in agreement with the

Conclusions

The combined ESR/U-series analyses realized on teeth carried out from two Eemian sites of Northern France, Caours and Waziers, demonstrate the reliability of the method to date Middle Paleolithic sites and, despite particular geochemical characteristics of part of the analyzed tissues, the validity of the used analytical protocols. This work demonstrates in particular the importance of the in situ dosimetry in the age calculation.

At Caours, the obtained ESR/U-series ages are in excellent agreement with the other available geochronological data derived from U-series and luminescence studies. They permit to calculate new mean age for the lower part of the sequence (124 ± 8 ka for unit 5 (N1, n=7) and 127 ± 15 ka for units 6-11 (including N3, n=6)). Associated to available paleo-environmental and archaeological data, they make of this site one of the best known Middle Palaeolithic sequence of France.

For Waziers, the new established ESR/U-series chronology, with a mean age of 129 ± 11 ka, confirms the record of Saalian Late Glacial deposits at the bottom of the sequence and of the subsequent Eemian interglacial stage, evidenced by malacological and palynological studies.

These results show the interest to integrate ESR/U-series dating analyses during the chronostratigraphical study of Middle Paleolithic sites when possible and demonstrate the necessity to involve geochronologists to the fieldworks and excavations, since its earliest stages.

Acknowledgements

We would like to thank the Caours and Waziers excavation teams for their disponibility, patience and helpful discussions.

The present study was financially supported by the UMR7194 HNHP which allowed the funding of the field missions and analyses., with the help of the PRC PaléHauts

The ESR and mobile gamma-ray spectrometers of the French National Museum of Natural History were bought with the financial support of the 'Sesame Ile-de-France' program and the 'Région Centre' respectively.

References

- Adamiec, G., Aitken, M., 1998. Dose-rate conversion factors: update. *Ancient TL*, 16, 37-50.
- Bahain, J.-J., Yokoyama, Y., Falguères, C., Sarcia, M.N., 1992. ESR dating of tooth enamel: a comparison with K-Ar dating. *Quaternary Science reviews*, 11, 245-250.
- Bahain J.-J., Falguères C., Dolo J.-M., Antoine P., Auguste P., Limondin-Lozouet, N., Loch J.-L., Tuffreau A., 2010. ESR/U-series dating of teeth recovered from well-stratigraphically age-controlled sequences from Northern France. *Quaternary Geochronology* 5, 371-375.
- Brennan, B.J., Rink, W.J., McGuirl, E.L., Schwarcz, H.P., Prestwich, W.V., 1997. Beta doses in tooth enamel by "One Group" theory and the Rosy ESR dating software. *Radiation Measurements*, 27, 307-314.
- Grün, R., Katzenberger-Apel, O., 1994. An alpha irradiator for ESR dating. *Ancient TL* 12, 35-38.
- Grün, R., Schwarcz, H.P., Chadam, J.M., 1988. ESR dating of tooth enamel: coupled correction for U-uptake and U-series disequilibrium. *Nuclear Tracks and Radiation Measurements*, 14, 237-241.
- Prescott, J.R., Hutton, J.T., 1994. Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. *Radiation Measurements*, 23, 497-500.
- Shao, Q., Bahain, J.J., Dolo, J.M., Falguères, C., 2014. Monte Carlo approach to calculate US-ESR ages and their uncertainties. *Quaternary Geochronology* 22, 99-106.
- Shao, Q., Bahain, J.-J., Falguères, C., Dolo, J.-M., Garcia, T., 2012. A new U-uptake model for combined ESR/U-series dating of tooth enamel. *Quaternary Geochronology*, 10, 406-411.