



HAL
open science

Measuring the quality of learning in a human-robot collaboration: a study of laparoscopic surgery

Eléonore Ferrier-Barbut, Philippe Gauthier, Vanda Luengo, Geoffroy Canlorbe, Marie-Aude Vitrani

► To cite this version:

Eléonore Ferrier-Barbut, Philippe Gauthier, Vanda Luengo, Geoffroy Canlorbe, Marie-Aude Vitrani. Measuring the quality of learning in a human-robot collaboration: a study of laparoscopic surgery. ACM Transactions on Human-Robot Interaction, In press, 10.1145/3476414 . hal-03355055

HAL Id: hal-03355055

<https://hal.science/hal-03355055>

Submitted on 8 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Measuring the quality of learning in a human-robot collaboration: a study of laparoscopic surgery.

ELÉONORE FERRIER-BARBUT, Sorbonne Université, France

PHILIPPE GAUTHIER, Sorbonne Université, France

VANDA LUENGO, Sorbonne Université, France

GEOFFROY CANLORBE, Sorbonne Université, France

MARIE-AUDE VITRANI, Sorbonne Université, France

Robot-Assisted Laparoscopic Surgery (RALS) is now prevalent in Operating Rooms (ORs). This situation requires future surgeons to learn Classic Laparoscopic Surgery (CLS) and RALS simultaneously. Therefore, along with the investigation of the differences in performance between the two techniques, it is essential to study the impact of training in RALS on the skills mastered in CLS. In this article, we study comanipulated RALS (Co-RALS), one of the two designs for RALS, where the human and the robot share the execution of the task. We use a rarely used in Human-Robot Interaction measuring tool: gaze tracking, and time recording to measure for the acquisition of skills in CLS either when training in Co-RALS or in CLS, and time recording to compare the learning curves between Co-RALS and CLS. These metrics allow us to observe differences in Co-RALS and CLS. Training in Co-RALS develops slightly better but not significantly better hand-eye coordination skills and significantly better time-wise performance compared with training in CLS alone. Co-RALS enhances time-wise performance in laparoscopic surgery on specific type of task that requires precision rather than depth perception skills, compared with CLS. The results obtained enable to further define the Human Robot Interaction quality in Co-RALS.

CCS Concepts: • **Human-centered computing** → **Collaborative interaction**; *Empirical studies in interaction design*.

ACM Reference Format:

Eléonore Ferrier-Barbut, Philippe Gauthier, Vanda Luengo, Geoffroy Canlorbe, and Marie-Aude Vitrani. 2018. Measuring the quality of learning in a human-robot collaboration: a study of laparoscopic surgery.. In *Woodstock '18: ACM Symposium on Neural Gaze Detection, June 03–05, 2018, Woodstock, NY*. ACM, New York, NY, USA, 19 pages. <https://doi.org/10.1145/1122445.1122456>

1 INTRODUCTION

Robotic surgery is increasingly used in Operating Rooms (ORs), especially the *Da Vinci* (cf. **figure 1-j**) with over 1 million procedures performed in 2018 [36]. This robot is essentially used to facilitate the practice of laparoscopic surgery. Without robot, e.g. Classic Laparoscopic Surgery (CLS) (cf. **figure 1-e**) is surgery done through small holes in the patient's belly, in which are inserted long instruments and a camera *i.e.* endoscope. The image of the endoscope is displayed on a 2D screen, showing the inside of the patient's body. CLS is advantageous for patients (less pain, shorter recovery, aesthetic benefits) but disadvantageous for surgeons who then have more visual as well as physical difficulties to deal with than in open surgery. Visual difficulties are partly generated by the absence of direct vision. The working space is visualized on a 2-dimensional screen while gestures are performed in a 3-dimensional space,

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2018 Association for Computing Machinery.

Manuscript submitted to ACM

53 making hand-eye coordination and depth perception complicated [7]. Physical difficulties are due to the length of
54 the instruments and to the fact that their insertion point (the small incisions in the patient body) induces kinematic
55 restrictions and the appearance of the fulcrum or lever arm effect [25]. Robot-Assisted Laparoscopic Surgery (RALS)
56 aims to alleviate varying difficulties depending on the robot's interface and features. There are two interaction designs
57 for Robot Assisted Laparoscopic Surgery (RALS): telemanipulated (Tele-RALS) (cf. figure 1-f) such as with the *Da Vinci*
58 and comanipulated (Co-RALS) (cf. figure 1-k). In Tele-RALS, as indicated by the semantics, a distance exists between
59 the surgeon (who sits at the master console) and the patient (above who stand the robotic arms holding the instruments,
60 the slave console). This distance removes the hand-eye coordination and the physical difficulties as instruments situated
61 at the slave console are controlled with joysticks situated at the master's console thanks to a communication channel
62 between the two. However, it introduces new difficulties. To use the robot surgeons and future surgeons have to learn,
63 on top of traditional skills linked with Classic Laparoscopic Surgery (CLS) more technology-oriented skills. Here, we
64 study the learning period when surgeons simultaneously learn to perform classic (without robot) and robot-assisted
65 laparoscopic surgery. During this period, the discrepancy between psycho-motor skills required to perform the two
66 techniques is problematic. Most importantly since learning laparoscopic surgery with a telemanipulated assistance
67 does not result in a transfer of skills to the classic technique [4], a situation that requires re-learning for each switch
68 between techniques.
69

73 A comanipulated robotic assistance, such as the Acrobot [17], aims to improve security and dexterity when performing
74 the surgical gesture while keeping the surgeon close to her/his patient. Comanipulation means the shared control of
75 the instruments by the robotic assistance and the surgeon [24] (cf. figure 1-l): both the robotic arms and the user
76 hold the instruments needed to perform the surgery. Same as for laparoscopic surgery, while technically possible to
77 perform viewing the image of the endoscope in 3D, Co-RALS is most frequently performed viewing the image of the
78 endoscope on a 2D screen. This design is at the fourth degree of human-robot collaboration scale [14]: Supportive. At
79 this level, the human and the robot work together at the same time and with the same workpiece to complete a common
80 task. Arguably, a successful *Supportive* human-robot collaboration is defined by its ability to augment performance
81 compared to the same task performed without a robot, but also to confer to the human a cognitive and physical load
82 that is neither too important nor too little, to diminish the difficulty of the task while keeping the human involved
83 and active [27]. For a task as complex as surgery simultaneously performed by a human operator and a robot, where
84 robots are far from being able to replace the human operators, it can be suggested that keeping the human involved
85 and active is mandatory, for him/her to maintain his/her acquired skills level. Indeed, in our case, the robot assists
86 the human in a task s/he already masters. Comanipulated robots for surgery are destined to work very closely with
87 surgeons and future surgeons without disturbing their dexterity skills and while increasing their performance. The
88 robot only augments already existing technical skills. Still, the quality of the human-robot interaction in Co-RALS
89 has not been investigated yet to our knowledge. We propose a method to measure with quantitative metrics, how a
90 collaborative robotic assistance not only can help to increase performance and learning in very complex tasks but also
91 the impact it has on the technical skills mastered by the human. We study it through two research questions, situated in
92 the specific context of learning when the acquisition of skills is of crucial importance.
93

94 The **main research question** is the following: does training with a comanipulated robot in what we call a *target*
95 *task*, in our case exercises of laparoscopic surgery, develop hand-eye coordination and time-wise skills in this target
96 -and simultaneously learned in a surgical curriculum- task? We focus on the hand-eye coordination skills as these are
97 especially complex to learn and perform in laparoscopic surgery for the previously mentioned reasons. To measure the
98 hand-eye coordination skills, we use a tool that to our knowledge has rarely been used in Human-Robot Interaction:
99
100
101
102
103
104




Fig. 1. CLS (Top); Tele-RALS (Middle); Co-RALS (Bottom). A CLS OR diagram (a) and an image of an OR in CLS (b). Mechanical tools (c), activated by pulling a trigger and a rotation knob (d). Surgeons insert instrument through small holes (e). Tele-RALS OR diagram (f) and an image of an OR in Tele-RALS (g) where a surgeon works at the master console and two assistants at the bed side. Moving joysticks (i) control tools (h) on the robotic arms at the slave console (j). Co-RALS OR diagram (k) and image of an OR in Co-RALS (l). Same tools as in CLS are used (m) manipulated the same way (n), while gestures are secured and improved by the robotic arms (o). Extended from "Impacts of Telem Manipulation in Robotic Assisted Surgery." by Avellino, I., et al. [2]. Extended with permission.

gaze tracking. The number of fixations (the event when the gaze remains on a point for 50 to 600 ms [18]) on the aimed target, the fixation rate per second and the duration of fixations, when performing in CLS are measured either after training in Co-RALS or in CLS. Previous works have shown that these metrics enable to measure hand-eye coordination skills developed when training [13], in laparoscopic surgery especially [19, 42, 43]. To measure time-wise skills, time elapsed is recorded when performing exercises in CLS, either after training in Co-RALS or in CLS. This shows the operator's performance efficiency [22] in the *target task* after training in Co-RALS. The **secondary research question** is the following: does using Co-RALS improve time-wise performance on exercises of laparoscopic surgery compared with the same exercises performed in CLS? Time elapsed is also recorded when performing repeated exercises with each technique. The repetition of the exercises highlights the shapes of the learning curves. Other metrics are used as exploratory measures: the NASA TLX [12] that serves to compare the workload between the robot-assisted technique and the classic technique, and a performance score given for the exercise performed in the classic technique either after training with the robotic-assistance or without.

157 In section 2 the existing connections between motor learning, robotic systems and their interfaces are discussed and
158 comanipulation is presented. In section 3 the experiment's methods are detailed. In section 4 results are submitted.
159 The results presented below, as measured by gaze tracking and time recording suggest that Co-RALS, contrarily to
160 what was shown for Tele-RALS, does not negatively impact skills in CLS, developing slightly but not significantly
161 better hand-eye coordination skills. Additionally, during the training session, the exercise of laparoscopic surgery that
162 develops precision skills presents a significantly shorter learning curve when performed in Co-RALS compared with
163 CLS, while an exercise of laparoscopic surgery that develops depth perception skills presents a learning curve that is
164 not statistically different from that of CLS. The results, although reporting on a relatively small number of participants,
165 demonstrate the ability of the metrics chosen to show differences in development of hand-eye coordination skills
166 between the classic and the robot-assisted techniques for laparoscopic surgery, and differences in the learning curve
167 between the two.
168
169
170

171 2 STATE OF THE ART 172

173 In this section, the consequences of interaction design for RALS on skills mastered by the user are presented and
174 comanipulation is described.
175

176 2.1 Performance, human skills, and interfaces for RALS 177

178 Numerous articles in the field of Human Computer Interfaces (HCI) have focused on the impact of interfaces for RALS
179 on non-technical skills [45] related to laparoscopic surgery: workflow, communication, situation awareness, teamwork
180 [2, 6, 29, 32–34]. These works are exclusively on Tele-RALS, and the metrics used are essentially subjective. These
181 robotic systems enhance the surgeon's experience by making them more independent to perform surgery and increasing
182 the number of instruments they can control. But the distance imposed with patients keeps them away from their
183 team. This physical distance is compounded by visual, auditory and mental distances. It disrupts access to information,
184 changes power distribution and decreases the surgeon's situation awareness. It also has consequences for students who
185 then have fewer tasks, decisions and actions to make [41] than when the surgical intervention is performed without
186 the robot. Their learning process is made more difficult: they can no longer learn by observing, hearing, and doing
187 according to the well known wording *See one, do one, teach one* [30] as they are not standing next to the surgeon any
188 more [23].
189

190 Although less studied in HCI, the impact of interaction design for robotic assistance on technical skills [26] related
191 to laparoscopic surgery is just as strong. The interaction design for robotic surgery defines the nature and the difficulty
192 of motor skills either learned or mastered by the user, and how they transfer to without robotic assistance. Robotic
193 assistance can easily improve performance when being used. However, as stressed in a review of literature on robotic
194 assistance of motor learning by Heuer et. al's from 2015 from the field of neurosciences, it is complicated to ensure
195 that after the robot is turned off, the user continues to perform as well as if s/he had never used it [15]. A risk exists
196 that the use of robotic assistance becomes "normal" and the motor learning highly dependent on the robot specific
197 dynamic environment. This situation would lead to a negative impact of training with a robotic assistance on skills
198 mastered in the classic technique, as it seems to be the case for Tele-RALS [3, 4]. Both these studies by Blavier et al.
199 from 2007 were conducted with medical students without any prior surgical experience and results suggest that training
200 in Tele-RALS has negative consequences on mastery of skills in CLS, compared with training in CLS alone. In the case
201 of a conversion from Tele-RALS to CLS during a surgical intervention, a scenario that regularly happens for different
202 reasons such as mentioned in Blavier's et al. article from 2007 [3], this negative impact can have major consequences.
203
204
205
206
207
208

209 Here, we hypothesize that training in Co-RALS results in equivalent hand-eye coordination and time-wise abilities
210 compared with training in CLS. Hence we take a different approach than previous work: with objective metrics, we
211 seek to observe the quality of the human-robot interaction through the analysis of the skills developed by the human in
212 conjunction with the performances achieved by the human-robot team.
213

214 2.2 Comanipulated Interfaces in Robotics

215 The definition of comanipulation can be found in Morel's article from 2013: *A comanipulator is thus any robotic system*
216 *performing a task, most often in contact with the environment, that can be controlled through direct contact by an operator.*
217 *It aims to increase the manipulation performance of the operator [24].*
218

219 Comanipulation is an interaction paradigm involving a robot and a user simultaneously manipulating a load or a
220 tool (cf. figure 1-m). The robot is employed as a comanipulated device, in the sense that the gesture control of the
221 instrument is shared by the robot and the surgeon. Of the difficulties of CLS mentioned earlier, comanipulation aims to
222 facilitate the physical ones in particular. The currently existing commercialized comanipulated robotic systems are
223 basically designed for specific types of surgical tasks [10, 16, 17, 28]. Research institutes have also exploited the idea for
224 precise surgical tasks [5]. Comanipulation can be applied to tasks that require both precise manipulation and human
225 judgment so as to enhance gesture quality [37]. This interaction design for robotics has the ability to at least compensate
226 for gravity and filter tremors [31] while performing the surgical gesture. Some other physical difficulties can also be
227 dealt with including the reduction of the fulcrum effect for example with the help of an active force feedback [35].
228 To improve haptic feedback, virtual fixtures can be applied [11, 40]. In this study, two comanipulated robotic arms
229 are used. One for each instrument manipulated when performing exercises in laparoscopic surgery. Only the basic
230 functions of the robots are studied. Gravity compensation for instruments is ensured, as well as tremor filtration. The
231 algorithm used for tremor filtration is a viscous field, *i.e.* a damping algorithm proportional to the velocity of movement
232 [8]. The robots exhibit high viscosity at low velocities and no viscosity at high velocities, at the instrument tip. It
233 helps the surgeons during precise surgical tasks (performed at small speed), by filtering unintentional movements and
234 augmenting precision of the gesture. This viscous field also increases the rate of adaptation of the user, forced to remain
235 active when performing her/his task. A similar algorithm, implemented in a comanipulated robotic assistance where
236 the human controls the direction and the speed while the robot ensures the precision and smoothness of motions by
237 suppressing sudden and abrupt gestures, has been shown to significantly improve performance for tasks of manual
238 welding [9]. In an other research article [39], a human robot cooperative calligraphic task is performed, in which
239 the human and the robot grasp a writing brush at the same time. The robot is controlled against the human force
240 to prevent the vibration and to enhance the accuracy, and results show the advantage of the control method. The
241 benefits of variable damping coefficient have been demonstrated. Thus, referring to the second question of research, it
242 is hypothesized that in Co-RALS, the adaptive damping algorithm implemented enables better time-wise performance
243 for exercises of laparoscopic surgery compared to the same exercises performed without the robotic assistance.
244
245
246
247
248
249
250
251

252 3 METHODS

253 To investigate 1. whether training with a comanipulated device in laparoscopic surgery results in an equivalent
254 development of skills in this *target task*, exercises of laparoscopic surgery, compared with training in the *target task* and
255 2. the comparison of the learning curve between the robot-assisted task and the classic task, we present an experimental
256
257
258
259
260

Table 1. Experimental Protocol: exercises in bold are those during which data is recorded. The others are familiarization exercises. The order of exercises performed during the Learning Step is randomized.

Group	Pre-Learning Step	Learning Step	Post-Learning Step
Classic	Task CLS Peg Transf. Domin. (4min)	Task CLS Pea on a Peg (30min)	Task CLS Peg Transf. Domin. (4min)
	Peg Transf. Non Domin. (4min)	Loops and Wire (15min)	Peg Transf. Non Domin. (4min)
Robot	Task CLS Peg Transf. Domin. (4min)	Task Co-RALS Pea on a Peg (30min)	Task CLS Peg Transf. Domin. (4min)
	Peg Transf. Non Domin. (4min)	Loops and Wire (15min)	Peg Transf. Non Domin. (4min)

protocol involving two conditions. The participants in *Classic* condition are pre-tested in CLS during the *Pre-Learning Step*, then trained in CLS, and then post-tested during the *Post-Learning Step* in CLS. The participants in *Robot* condition are also pre-tested in CLS, then trained in RALS, then post-tested in CLS (cf. table 1). Both groups are Pre-trained and Post-trained in the same task, CLS, to compare the differences in the process of learning for each group, *Classic* and *Robot*, in CLS. The main research question is studied by comparing the skills in CLS of the two groups, *Classic* and *Robot* with gaze-tracking and time recording, and the secondary research question by comparing the learning curves of the two groups *Classic* and *Robot* during their learning session with time recording. The exercises of laparoscopic surgery performed at each step were chosen among basic training exercises for this discipline, depending on the skills they enable to train and measure. During the Pre and Post-Learning steps, the exercise chosen was the Peg Transfer as it enables to observe a large panel of participant's skills in laparoscopic surgery: bimanual coordination, precision, and depth perception. It requires the participant to lift six objects place on the left side of a board with a grasper first using non-dominant (in this case, left) hand and transfer the object midair to the dominant hand. Then, the participant has to place each object on a peg on the right side of the board. It is also performed from dominant to non-dominant hand. This exercise is further described later on. Pea on a Peg and Loops and Wire were chosen because they trained precision and depth perception respectively. Exercise Pea on a Peg consists of placing 14 beads on pegs of different heights placed on a board, and exercise Loops and Wire consists of passing a wire through 4 different loops placed on a board. Both these exercises are also further described later on. The fact that each exercise trains different skills enables to better define the differences between CLS and Co-RALS in terms of learning. This experimental protocol is tested both with Resident participants and, more numerous, Non-Resident participants. The similarity of the exercises in laparoscopic surgery performed by each group is controlled. Level equivalence between group Robot and Classic is statistically verified for Non-Resident participants during the *Pre-Learning Step*. There were not enough Resident participants involved in the protocol to perform this verification, hence the Resident participants' results are presented as indication of tendency and insights for future research rather than results that can be generalized. The experiment is thus divided into three steps: 1. A pre-learning step to control the participants' level, 2. A learning step to train the participants of each group respectively in each task and (CLS and Co-RALS) and compare their performance and 3. A post-learning step to compare the mean level of each group in the target activity, *i.e.* CLS.

3.1 Participants

Fourteen Non-Resident participants and six Residents in medicine are recruited. Among them, seven Non-Resident participants and three Resident participants (group Classic) performed their training session in in CLS (task CLS). Seven other Non-Resident participants and three other Resident participants (group Robot) performed their training

313 session in RALS on a comanipulated robot (task Co-RALS). Resident participants have a small and contrasted experience
 314 in laparoscopic surgery (6 ± 4.5 hours of practice). The number of Resident participants involved in the study does
 315 not allow to perform statistics. Still, their results enable to verify whether the trend observed among novices may
 316 be applied to intermediate level users. Because of their tight agenda, Residents in medicine could only perform one
 317 exercise, *Pea on a Peg*, during the training session and, due to a lack of technical and time resources, no gaze data could
 318 be recorded during *Peg Transfer* exercise before and after the Learning session. Their results are presented separately.
 319 Five of them are first year residents and one is third year resident. Non-Resident participants all declared to be novices
 320 in laparoscopic surgery, and are all university students. All procedures are in accordance with the ethical standards of
 321 the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments
 322 or comparable ethical standards.
 323
 324
 325

3.2 Material



345 Fig. 2. (a) CLS Set-Up (b) Co-RALS Set Up

346
347
348
349 3.2.1 *Apparatus*. Material includes two set-ups: one in CLS, and one in Co-RALS.

350 **Task CLS** *The CLS set-up*: It includes two surgical graspers. Exercises are performed in a pelvi-trainer, above which is a
 351 2D screen transcribing the 3D working space inside it (cf. figure 2a). The endoscope does not move during the entire
 352 learning session.
 353

354 **Task Co-RALS** *The Co-RALS set-up*: It consists of two robotic arms, each holding the same two graspers as those used
 355 in the CLS set-up. (cf. figure 2b). The two robotic arms are modified Haption Virtuose 3D robots, characterized by six
 356 rotational joints. The robots used are 3D robots as other degrees of freedom are constrained by the entree point in the
 357 patients' belly. The first three joints are fully actuated and the other three form a free wrist that allows full motion
 358 across the surgical workspace [21]. Our research team designed the software implemented in these robotics arms:
 359 (i) tool weight compensation so that holding the tools is transparent to surgeons [38] and (ii) a damping algorithm
 360 proportional to the velocity of movement, experimentally tuned such as in [8, 20]. The same pelvi-trainer as for the
 361 CLS set-up is used (cf. figure 2a). The endoscope is not moved during the entire learning session.
 362
 363
 364

365 3.2.2 *Measuring tools and metrics.* Different measuring tools and metrics are used at each step of the experiment to
366 assess hand-eye coordination abilities, time-wise performance and, workload and scores. They were chosen for their
367 ability to show not only the observed performance at the end of a task but also the learning process itself. **Gaze-tracking**
368 is used during the Pre-Learning Step to verify the absence of heterogeneity in average hand-eye coordination skills
369 in laparoscopic surgery of groups Classic and Robot before the Learning session. It is also used during Post-Learning
370 step to quantify for differences in the same skills between these groups after the Learning session. **Time** is recorded
371 at each step of the experiment. Before and after the Learning session this metric enables to answer the first research
372 question, together with gaze-tracking. During the learning session, Time recording serves to answer the secondary
373 research question. The results obtained at the **NASA Task Load Index** during the learning session and the scores at
374 the *Peg Transfer* exercise are explored, respectively as self-perceived feeling of comfort with the task and observed
375 performance.
376
377
378

379 Pre-Learning Step

- 380 • Gaze-Tracking: Observing gaze pattern has been proven to be relevant in distinguishing novices from experts
381 [13, 19, 42, 44]. Experts have a greater ability to anticipate their gestures with their gaze than novices. This
382 results, in tasks of target reaching (such as the *Peg Transfer*), in a greater ability to look at the aimed target rather
383 than the instruments used to reach the target. In other words, experts anticipate their movement by taking
384 their eyes off it and projecting them to its goal, while novices are focused on achieving the movement itself
385 and keep their eyes on it. This translates into a struggle to detach their eyes off their hands or the instruments
386 they hold to perform the movement and to project their gaze to its goal. It can be measured by the number
387 of fixations (the maintenance of the gaze in a single location) on the aimed target before reaching it with the
388 instruments, a greater number meaning a better ability to anticipate, and greater expertise [1]. Experts also
389 tend to do less back and forth movements with the eyes -movements used by novices to make calculations of
390 distances, lengths, velocities- and a smaller number of fixations. The duration of their fixations also tends to
391 be larger than for novices. Hence, **the number of fixations on the aimed target** before it has been reached
392 with the instruments, **fixation rate per second** and **duration of fixations** are analyzed. A higher **number**
393 **of fixations on the aimed target** at the specific moment of reaching for the target in the *Peg Transfer* means
394 a greater level of comfort with the task. At the opposite, less **fixations per second** and greater **duration of**
395 **fixations** while performing the entire exercise means better hand-eye coordination skills. Gaze data analysis is
396 performed separately for each phase of the *Peg Transfer* exercise described later on in the article. It is cut into
397 3 phases: *Grab*, *Transfer* and *Drop*. *Grab* corresponds to the moment where participants grab the bead with
398 their dominant hand, *Transfer* to the moment they pass it to their non-dominant hand and *Drop* to the moment
399 they put it on the peg. The *Transfer* and *Drop* phases are the most interesting. They represent the moment
400 where the participants are specifically doing a target reaching task. Recording gaze data enables observation of
401 hand-eye coordination abilities. The Eye Tracker used to record gaze data is a Tobii X3-120 screen-based eye
402 tracker with a sampling rate of 120Hz. It is mounted at the bottom of the screen at the eye level of participants.
403 Data analysis is performed with Tobii Pro Lab and the filtering algorithm used is the IV-T Classifier. To identify
404 **the number of fixations on the aimed target**, we manually defined, using Tobii Pro Lab, dynamic zones of
405 interest. They were the zones, on the 2D screen, aimed by the participants when performing the *Peg Transfer*
406 exercise described later on. Statistics of the number of fixations in these zones can then be extracted.
407
408
409
410
411 • Time: A timer is set during performance of exercises.
412
413
414
415
416




Fig. 3. (a) Peg Transfer (b) Pea on a Peg (c) Loops and Wire

- Score: Videos of the camera used to perform the exercises is recorded to allow for scoring of the *Peg Transfer* exercise. 10 penalty points are counted per dropped sleeve. When a sleeve falls from the pegboard 20 penalty points are counted. Total score is time in seconds + penalty points. This score shows temporal as well as qualitative performance.

Learning Step

- Time: A timer is set during each trial of the two exercises performed during the learning session, described latter on in the article.
- NASA Task Load Index (TLX): It is used as an exploratory measure of comfort level with the executed tasks during the learning session. This index is a subjective, multidimensional assessment tool that rates perceived workload in order to assess a task, system, or team's effectiveness or other aspects of performance. It was developed by the Human Performance Group at NASA's Arms Research Center. The higher the score is, the more important is the perceived workload. In this study, the score is NASA weighted. Participants were required to fill the NASA TLX after they had performed the learning session.

Post-Learning Step

The measuring tools and metrics are the same as during the *Pre-Learning Step*.

3.3 Procedures

Experiment is divided into three steps (cf. **table 1**):

3.3.1 Pre-Learning Step. This step consists in the measurement of participants' base level. The data recorded enables to ensure that the two groups of participants: Classic and Robot do not show, initially, significant differences in level in the target activity. Means of each group before the learning session are compared to check for statistical differences.

The exercises performed are the following:

- *Peg Transfer Non Dominant*: familiarization exercise, data is not recorded during this task because it is the familiarization task, during which participants discover instruments and set-up for four minutes. There are 6 sleeves and 12 pegs on the board. The 6 sleeves are positioned on 6 pegs, on one side of the board. Subjects have to transfer the sleeves from one side of the board to the other. To pick up a sleeve, they use the instrument held in the non dominant hand. Then they transfer the sleeve to the instrument held in the dominant hand. Maximum time is 4 minutes.

- *Peg Transfer Dominant*: recorded exercise. Subjects have to transfer sleeves resting on pegs on the other side of the board. They pick the sleeves up with the instrument held in the dominant hand, transfer them in the instrument held in the non dominant hand, drop them on the other side of the board. They have to do as many as possible within four minutes. Data is recorded during this exercise only (cf. figure 3a).

3.3.2 *Learning Step*. Participants are randomly assigned either to group Classic which will perform one hour training session in CLS or group Robot which will perform 1h training in Co-RALS. During each learning session, participants have to perform two exercises: *Pea on a Peg*, *Loops and Wire* (cf. figure 3). These two exercises are performed in a random order. They are each performed repeatedly: participants had a maximum number of five trials for each exercise. To avoid the influence of fatigue on performance, a maximum total time per exercise and a maximum time per trial is set. The exercises are the same when performed either in Co-RALS or in CLS. Exercises performed are the following:

- *Pea on a Peg*: Participants have to place 14 beads on pegs of different heights. The beads are positioned on a peg board with the cup containing them in front. A maximum time per trial of 10 minutes is set as well as a maximum number of trials of 5 and a maximum time spent on exercise of 30 minutes. Once all the beads are placed on the pegs, the trial is finished. The psychomotor skills developed are similar to those involved during *Peg Transfer*: fine motor skills, coordination, precision and depth perception (cf. figure 3b).
- *Loops and wire*: The exercise contains a peg board with 4 loops on which is positioned a flexible wire. Participants have to insert the wire in the 4 loops in a specific order indicated on the board. A maximum time per trial of 4 minutes is set as well as a maximum number of trials of 5 and a maximum of time spent on exercise of 15 minutes. Once the wire is inserted in all the loops, the trial is finished. The psychomotor skills developed are related to depth perception and manipulation of the instruments (cf. figure 3c).

3.3.3 *Post-Learning Step*. Participants have to perform the same exercises as the one performed during the pre-learning step:

- *Peg Transfer Non Dominant*: familiarization exercise.
- *Peg Transfer Dominant*: recorded exercise.

4 RESULTS

We present the results given by each of the measuring tools and metrics used during the experiment. As mentioned before, gaze-tracking could only be recorded for Non-Resident participants while time and score were recorded for all participants. Statistical tests were performed as a comparison between the means of each group: *Classic* and *Robot* for every metric used during *Pre-Learning*, *Learning* and *Post-Learning* steps. All sets of data were normally distributed, variances were homogeneous and samples were independent, hence Student's t tests were performed for every pair of data: number of fixations, duration of fixations, time to perform *Peg Transfer*.

4.1 Main Research Question: Does learning laparoscopic surgery with a comanipulated robotic assistance result in development of skills in CLS?

4.1.1 *Pre-Learning Step*. Gaze data and time data during Pre-Learning is analyzed to compare between group Classic and group Robot and make sure that there are no statistical differences between these two groups previous to the learning session. The means in terms of time taken to perform *Peg Transfer* in the Pre-Learning step are different between group Classic (3.8 ± 0.4) and group Robot (2.7 ± 0.9) but not significantly different (*Student's t Test*, $t(11.2)=2.09$,

$p = 0.06$). Still, because of the low p -value, the difference in terms of time taken to perform *Peg Transfer* after learning between group Classic and Robot should be considered with caution. All other data measured before learning shows clearly no statistical difference between group Classic and group Robot in terms of gaze data *i.e.* duration of fixations during *Grab* phase (*Student's t Test*, $t(8.7459) = 0.62994$, $p = 0.5$), *Transfer* phase (*Student's t Test*, $t(9.4918) = -0.5219$, $p = 0.6$) and *Drop* phase (*Student's t Test*, $t(11.334) = -0.86636$, $p = 0.4$) and number of fixations during *Grab* phase (*Student's t Test*, $t(8.1897) = 0.58616$, $p = 0.5$), *Transfer* phase (*Student's t Test*, $t(8.2418) = 0.42915$, $p = 0.6$) and *Drop* phase (*Student's t Test*, $t(9.8688) = 0.54009$, $p = 0.6011$).

These tests have been performed for Non-Resident participants, Resident participants were not numerous enough to perform statistical comparisons.

4.1.2 Post-Learning Step.

- Mean number of fixations on aimed target: **for Non-Resident participants**, before learning, on average participants perform 0.6 ± 0.1 fixations on aimed target during *Transfer* phase. In the same phase, after learning, group Classic performs 0.55 ± 0.2 fixations on average on aimed targets, and group Robot 0.59 ± 0.2 fixations on average on aimed target. Before learning, on average participants perform 0.54 ± 0.19 fixations on aimed target during *Drop* phase. In the same phase, after learning, group Classic performs 0.55 ± 0.22 fixations and Group Robot performs 0.45 ± 0.13 fixations on average on aimed target (**cf. figure 4**). These differences are not significant (*Student's t Test*, $t(0.12) = 20.3$, $p = 0.9$).




Fig. 4. Number of fixations on target before reaching them with the instrument during *Transfer* and *Drop* phases: Non-Residents. **Ta1-3** (Mean number of fixations on Targets n°1, 2 and 3 before reaching them with the instrument), **Ta4-5** (Mean number of fixations on Targets n°4 and 5 before reaching them with the instrument), **Ta6** (Mean number of fixations on Target n°6 before reaching it with the instrument). Targets are separated in accordance with the % of participants who managed to reach them, this number being proportionally lower for the targets with the higher number. On average, group Robot does more fixations on targets 1 to 5 in *Transfer* phase, on targets 1 to 3 in *Drop* phase, than group Classic. Group Classic does more fixations than group Robot on target n°6, but variability is also twice as important: ± 0.28 for group Classic, ± 0.14 for group Robot.

- Mean fixation rate per second: **for Non-Resident participants**, the fixation rate per second decreases for both groups before and after learning all phases, except in *Grab* phase where it slightly increases for group Classic. After learning, in *Grab* phase, group Classic increases on average the fixation rate per second of 4% compared to before learning, group Robot stagnates. In *Transfer* phase, group Classic does on average as many fixations per second before and after learning, while group Robot does 5% less. Finally in *Drop* phase, group Classic decreases by 6% on average the fixation rate per second, group Robot decreases by 10% (cf. figure 5). Difference is not statistically significant between the groups (*Student's t Test*, $t(1.9)=9.7$, $p = 0.08$).




Fig. 5. Mean fixation rate per second and Duration of fixations during Peg Transfer Post-Learning Session: Non-Residents. Participants in group Robot reduce the **mean fixation rate per second** in phases *Transfer* and *Drop* before and after learning while group Classic stagnates in *Transfer* phase. Standard Deviation reduces strongly for the two groups before and after learning. Participants in groups Classic and Robot augment the **mean duration of fixations** before and after learning. Standard deviation also reduces strongly for the two groups.

- Duration of fixations: **for Non-Resident participants**, the duration of fixations increases for both groups before and after learning in all phases of the *Peg Transfer* exercise, except for *Grab* phase in which it slightly decreases for group Classic. After learning, in *Grab* phase, group Classic decreases on average the duration of fixations by 9% compared to before learning, group Robot increases by 6%. In *Transfer* phase, group Classic increases the duration of fixations of 3% before and after learning, while group Robot increases by 5%. Finally in *Drop* phase, group Classic increases by 2% on average the duration of fixations, group Robot increases by 7% (cf. figure 5). Difference is not statistically significant between the groups (*Student's t Test*, $t(-0.7)=11.9$, $p = 0.4$).
- Time to perform *Peg Transfer*: **for Non-Resident participants**, both groups decrease time to perform the *Peg Transfer* exercise after the learning session. Mean time to perform the exercise before the learning session is 3.1 ± 0.9 minutes, while mean time to perform the exercise after the learning session is, for group Classic of 2.6 ± 0.6 minutes and for group Robot of 1.9 ± 0.4 minutes (cf figure 6). Difference is significant between group Classic ($M=2.6$, $SD=0.23$) and group Robot ($M=1.9$, $SD=0.18$), *Student t. test*, $t(11.2)=2.09$, $p=0.05$.
- Time to perform *Peg Transfer*: **for Resident participants**, both groups decrease time to perform the *Peg Transfer* exercise after the learning session. Mean time to perform the exercise before the learning session is of




Fig. 6. Mean time to perform Peg Transfer Pre and Post-Learning Session

3.3 ± 0.9 minutes, while mean time to perform the exercise after the learning session is, for group Residents Classic of 1.8 ± 0.6 minutes and for Residents Robot of 1.8 ± 0.4 minutes (cf figure 6).

4.2 Secondary Research Question: Does using a comanipulated interface for RALS improve performance compared with CLS?

- Time to perform training exercises: **for Non-Resident participants**, during the learning session, for exercise *Pea on a Peg*, group Classic starts for Trial n°1 with a mean time of 8.6 ± 1.9 minutes and ended for Trial n°5 with a mean time of 4 ± 2.4 minutes. Group Robot starts for Trial n°1 with a mean time of 4.4 ± 2.8 minutes and ends for Trial n°5 with a mean time of 3.1 ± 1.3 minutes. Difference of time taken across conditions is significantly different for trial n°1 at the $p < 0.002$ level [$F(1,13) = 17.11, p = 0.001$]. For exercise *Loops and Wire* no significant difference is observed between group Classic and group Robot. Group Classic starts for Trial n°1 with a mean time of 2.7 ± 0.7 minutes and ended for Trial n°5 with a mean time of 1.6 ± 1.4 minutes. Group Robot starts for Trial n°1 with a mean time of 3.2 ± 1.2 minutes and ends for Trial n°5 with a mean time of 1.5 ± 0.7 minutes (cf. figure 8). After verifying the normality and sphericity of data, and the independence of samples, a repeated measures analysis of variance was performed to test for effect of learning. This effect is significant for Group Classic for exercise *Pea on a Peg* (*Anova*, $F(1,6) = 63.25, p = 0.0002$), but not for group Robot (*Anova*, $F(1,7) = 2.215, p = 0.18$). For exercise *Loops and Wire*, the effect of learning is not significant for group Classic (*Anova*, $F(1,6) = 3.68, p = 0.1$), but it is for group Robot (*Anova*, $F(2,5)=8.4, p = 0.025$).
- Time to perform training exercises: **for Resident participants**, exercise *Pea on a peg* alone is performed. Group Residents Classic starts for Trial n°1 with a mean time of 4.7 ± 1 minutes and ends for Trial n°5 with a mean time of 2.9 ± 1.4 minutes. Group Robot starts for Trial n°1 with a mean time of 6.9 ± 2.8 minutes and ends for Trial n°5 with a mean time of 1.4 ± 1.6 minutes (cf. figure 8).

4.3 Exploratory Results: Self perceived comfort level with the task and observed performance

- Score for NASA TLX: **for Non-Resident participants**, they show an almost equal score for the two groups. Group Classic rates a mean score of 61 ± 6 and group Robot of 64 ± 12 (cf. figure 9).





Fig. 7. Learning Session Non-Residents




Fig. 8. Learning Session Residents

- 721
722
723
724
725
726
727
728
- Score for NASA TLX: **for Resident participants**, both score very closely the task: 58 ± 18 for group Classic and 60 ± 19 for group Robot (cf. figure 9).
 - Score at *Peg Transfer* exercise: **for Non-Resident participants**, mean score before the learning session for two groups is of 230 ± 81 , after learning for group Classic of 163 ± 56 , and after learning for group Robot of 158 ± 56 . The two groups decrease their score in a similar way, showing equivalent improvement of skills.




Fig. 9. NASA TLX

- Score at *Peg Transfer* exercise: **for Resident participants**, mean score before the learning session for the two groups is of 222 ± 86 , after learning for group Classic of 111 ± 59 , and after learning for group Robot of 147 ± 38 . Group Classic decrease their score a little bit more than group Robot.

5 DISCUSSION

The measuring tools' and metrics' results confirm their ability to show the differences in terms of skills and performance between the two experimental conditions (Co-RALS and CLS). They also confirm the absence of negative impact of training in Co-RALS on skills in CLS. Gaze-tracking enables to observe the subtle disparities in hand-eye coordination skills in CLS between training in Co-RALS and in CLS, although showing both train the human to perform laparoscopic surgery. Time recording showed a slightly significant difference in terms of the ability for speed in CLS for group trained in Co-RALS compared with group trained in CLS, but this result should be analyzed with care as before learning, group Robot showed better time-wise skills than group Classic. The scores obtained in CLS by each group show no difference in achieved performance. Time recording when performing similar repeated exercises in Co-RALS and in CLS identifies that differences in the number of repetition needed before mastering the task, and the time performance that can be achieved for each condition depends on the type of task performed. The NASA TLX, finally, confirms that no supplementary workload was felt when performing in Co-RALS compared with CLS.

5.1 Main Research Question: *Does learning laparoscopic surgery with a comanipulated robotic assistance result in development of skills in CLS?*

5.1.1 Non-Resident participants. Both group Classic and group Robot decrease time taken to perform the *Peg Transfer* exercise. The difference between the groups is statistically significant, with group Robot taking on average 0.7 minutes less than group Classic after the learning session. Thus, group Robot seems to have developed better time-wise skills in laparoscopic surgery compared with group Classic. Gaze patterns between the two groups are very similar. Still, group Robot makes on average, after the learning session, slightly less and longer fixations in the three phases of the exercise: *Grab*, *Transfer* and *Drop*. These differences are not significant, but one can postulate that longer and repetitive learning sessions would expand them. Also, group Robot augmented, on average, the number of fixations on target before reaching it with the instrument in phases *Transfer* and *Drop* for targets n°1 to 5 while group Classic, on average,

781 slightly decreased it, showing little improvement for the first group, and consistency for the second. We interpret these
782 results as a consequence of having to deal with the algorithm implemented in the robot, the viscous field forcing the
783 participants in group Robot to remain active when performing the gesture. This may have increased their attention
784 state, despite the fact that their score is almost equivalent to that of the group Classic. In other words, the results suggest
785 that group Robot has developed skills in laparoscopic in a way that is not statistically different to those developed by
786 group Classic in the same task. Without it being statistically significant, group Robot even has better results in terms of
787 number of fixations on target, fixation rate per second and mean duration of fixations after learning than group Classic.
788
789

790
791 **5.1.2 Resident Participants.** Groups Classic and Robot, for Resident participants, took on average the same time to
792 perform the *Peg Transfer* exercise after the learning session, suggesting they have equivalently improved their skills.
793 Time taken to perform the *Peg Transfer* decreases equivalently for both group Classic and group Robot. The results
794 from the small cohort of residents show a similar trend to that observed in the non-residents' results. This suggests that
795 the same results could be obtained on this experiment with larger groups of residents in medicine.
796

797 **5.2 Secondary Research Question: Does using a comanipulated interface for RALS improve** 798 **performance compared with CLS?** 799

800
801 **5.2.1 Non-Resident Participants.** For Non-Resident participants, it appears that training with robotic assistance de-
802 creases the time taken to perform exercises that require precision and fine motor skills such as *Pea on a Peg*. The group
803 of participants trained in Co-RALS took significantly less time to perform the first trial, and visibly less time to perform
804 every other trial. Contrary to what was expected, for exercise *Loops and Wire* in which proprioceptive and instrument
805 manipulating skills are required, comanipulated robot-assistance seems to hinder learning compared to no robotic
806 assistance. The damping algorithm may, in this case, have disturbed the participants as it forces them to adapt to a
807 new eye-hand coordination which has no advantage over the goal of this exercise. The implementation of an other
808 algorithm such as virtual fixtures may help to improve performance for such an exercise but could have a negative
809 impact on skill transfer to without robot mode. Further investigation is needed. Scores obtained at the NASA Task Load
810 Index are very close for the two groups, not significantly different, showing an equivalent cognitive and physical load
811 perceived when training either in CLS or in Co-RALS, suggesting it was as difficult to perform the learning session in
812 CLS as it was in Co-RALS.
813
814
815

816
817 **5.2.2 Resident Participants.** For Resident participants, who had a better base level at laparoscopic surgery, more trials
818 were required in RALS for *Pea on a Peg* exercise than for Non-Resident participants to attain the same level as in CLS,
819 but better performance was attained with the same number of trials. This may be due to the fact that, as they were used
820 to performing in CLS, they had to re-adapt to this new setting before they could achieve their best performance on it.
821 No significant difference was observed regarding scores obtained at the NASA Task Load Index when training either in
822 CLS or in Co-RALS. Same as for the main question of research, a similar trend than that of the novices is observed. This
823 paves the way for more research on effect of Co-RALS on subjects with intermediate level in CLS, to assess whether
824 their performance can be improved significantly compared with CLS.
825
826
827

828 **5.3 Exploratory Results: Self perceived comfort level with the task and observed performance**

829
830 **5.3.1 Non-Resident Participants.** Group Classic and Robot have on average equivalently improved their score at the *Peg*
831 *Transfer* exercise before and after learning, with group Robot slightly better than group Classic. The mean score given
832

833 at the NASA TLX is almost equivalent for the two groups, showing no superior workload for group Robot compared
834 with group Classic.
835

836 5.3.2 *Resident Participants*. Resident participants started with a score equivalent to the Non-Resident participants -222
837 compared with 230- and ended with a slightly better score than the Non-Resident participants -111 compared with
838 153 for group Classic and 147 compared with 158 for group Robot-. No difference is observed in terms of perceived
839 workload between group Robot and group Classic: the mean score at the NASA TLX is very close for the two groups.
840
841

842 6 CONCLUSION

843

844 The experiment conducted and its results, first, demonstrate that the measuring tools and metrics used and the exercises
845 performed succeed in showing the learning process and performance of participants on tasks of robotic and classic
846 laparoscopic surgery. The different metrics' results also indicate the advantages and limits of Co-RALS for learning.
847 A comanipulated interface for RALS seems to succeed in maintaining active learning of the motor skills of the user
848 *i.e.* in the *target task* (CLS) while performing the robot-assisted task (Co-RALS), contrarily to what was observed by
849 Blavier et al. with the da Vinci [3, 4]. However, these results confirm the motivations that lead to the development of
850 comanipulation [24]: to support and assist the gesture to make it easier to perform without changing its characteristics.
851 Gaze-tracking and time recording permit observation of two different aspects of the learning process: the psycho-motor
852 skills' acquisition and the immediately visible performance. These two measuring tools taken together provide a
853 comprehensive overview of what has been learned when training with robotic assistance. However, they go along
854 with other measurements, whose purpose is to study the performance of the human-robot team. Thus, to evaluate
855 the learning curve on exercises of laparoscopy either when training in Co-RALS or in CLS, we used time-recording
856 also. We observed the learning curve of two exercises: one is learned faster and performed more rapidly in Co-RALS
857 than in CLS, and the other is learned and performed more slowly with the robot compared to without. Hence, the
858 superiority of the human-robot team's performance in terms of the learning curve, seems to depend greatly on the
859 type of exercise performed. Co-RALS enables to increase performance for exercises of fine motor skills and precision
860 but for a task of proprioceptive and manipulating skills, *Loops and Wire*, it does not improve the learning process. The
861 exploratory measures show other aspects of the human-robot interaction. The results obtained with the NASA TLX
862 suggest that interacting with a robotic assistance to laparoscopic surgery results in an equivalent workload compared
863 with performing CLS. The comparison of the peg transfer scores in CLS either after training in Co-RALS or in CLS
864 demonstrate and equivalent performance between the two groups, confirming the previous results.
865
866
867
868
869

870 This study encourages to pursue research in human-robot interaction using quantitative metrics to qualify the
871 conditions for interactions to be virtuous both for the human, and for the human-robot team's performance. The
872 results are also in favor of other interaction designs for RALS than the dominant one, Tele-RALS. Still, these findings
873 present some limitations. First, the number of exercises performed are limited. Future research on comanipulated
874 interfaces and RALS with longer, more complex and realistic tasks may show more clearly how it can benefit to
875 students in surgery. Second, the number of Resident participants is small. A study with a greater number of Resident
876 participants, with groups of different levels would show more precisely how the comanipulated robotic system impacts
877 performance and develops psychomotor skills depending on the level in CLS. Also, our study only focuses on one
878 aspect of laparoscopic surgery which is dexterity. One could imagine future research that would involve knowledge
879 also: of anatomical structures, procedures, risks etc. Despite these limitations, our results pledge for more research on
880 human-robot collaboration, which could lead to more adapted and hence more easily adopted technologies.
881
882
883
884

7 ACKNOWLEDGMENTS

The authors would like to thank Xavier Deffieux and Sofiane Bendifallah, hospital practitioners who helped recruiting Resident participants and Gilles Bailly for his precious contributions to experimental design and article redaction. The authors would also like to thank all members of the Agathe team at Institut des Systèmes Intelligents et de Robotique (ISIR) who helped in editing the paper: especially Rémi Chalard, Ignacio Avellino, Angéline Bellicha, Etienne Moullet, Jesus Mago, Felix Péchereau, Fabien Vérité and Mégane Millan. The authors declare that they have no conflict of interest. This work has been supported by the funds managed by the ANR within the Investissements d'Avenir program (Labex CAMI) under reference ANR-11-LABX-0004.

REFERENCES

- [1] Hajra Ashraf, Mikael H Sodergren, Nabeel Merali, George Mylonas, Harsimrat Singh, and Ara Darzi. 2018. Eye-tracking technology in medical education: A systematic review. *Medical teacher* 40, 1 (2018), 62–69.
- [2] Ignacio Avellino, Gilles Bailly, Geoffroy Canlorbe, Jérémie Belghiti, Guillaume Morel, and Marie-Aude Vitrani. 2019. Impacts of Telemanipulation in Robotic Assisted Surgery. In *CHI Conference on Human Factors in Computing Systems Proceedings (CHI 2019)*.
- [3] Adélaïde Blavier, Quentin Gaudissart, Guy-Bernard Cadière, and Anne-Sophie Nyssen. 2007. Comparison of learning curves and skill transfer between classical and robotic laparoscopy according to the viewing conditions: implications for training. *The American journal of surgery* 194, 1 (2007), 115–121.
- [4] Adélaïde Blavier, Quentin Gaudissart, Guy-Bernard Cadiere, and Anne-Sophie Nyssen. 2007. Perceptual and instrumental impacts of robotic laparoscopy on surgical performance. *Surgical endoscopy* 21, 10 (2007), 1875–1882.
- [5] Rémi Chalard, David Reversat, Guillaume Morel, Pierre Mozer, and Marie-Aude Vitrani. 2018. Precisely positioning the tip of an instrument inserted through an orifice with a free wrist robot: application to prostate biopsies. *International journal of computer assisted radiology and surgery* 13, 5 (2018), 611–618.
- [6] Amy Cheatle, Hannah Pelikan, Malte Jung, and Steven Jackson. 2019. Sensing (Co) operations: Articulation and Compensation in the Robotic Operating Room. *Proceedings of the ACM on Human-Computer Interaction* 3, CSCW (2019), 1–26.
- [7] S Dharia and T Falcone. 2005. Robotics in reproductive medicine. Fertility and Sterility. (2005).
- [8] Lin Dong. 2017. *Assistance to laparoscopic surgery through comanipulation*. Ph.D. Dissertation. Paris 6.
- [9] Mustafa Suphi Erden and Bobby Marić. 2011. Assisting manual welding with robot. *Robotics and Computer-Integrated Manufacturing* 27, 4 (2011), 818–828.
- [10] Sarah Ferrand-Sorbets, D Taussig, Martine Fohlen, Christine Bulteau, Georg Dorfmueller, and Olivier Delalande. 2010. Frameless stereotactic robot-guided placement of depth electrodes for stereo-electroencephalography in the presurgical evaluation of children with drug-resistant focal epilepsy. In *CNS Annual Meeting*. CNS Annual Meeting.
- [11] Caspar Gruijthuisen, Gianni Borghesan, Dominiek Reynaerts, and Emmanuel Vander Poorten. 2019. A Hybrid Active/Passive Wrist Approach for Increasing Virtual Fixture Stiffness in Comanipulated Robotic Minimally Invasive Surgery. *IEEE Robotics and Automation Letters* 4, 3 (2019), 3029–3036.
- [12] Sandra G Hart and Lowell E Staveland. 1988. Development of NASA-TLX (Task Load Index): Results of empirical and theoretical research. In *Advances in psychology*. Vol. 52. Elsevier, 139–183.
- [13] Adrian Harvey, Joan N Vickers, Ryan Snelgrove, Matthew F Scott, and Sheila Morrison. 2014. Expert surgeon's quiet eye and slowing down: expertise differences in performance and quiet eye duration during identification and dissection of the recurrent laryngeal nerve. *The American Journal of Surgery* 207, 2 (2014), 187–193.
- [14] Evert Helms, Rolf Dieter Schraft, and M Hagele. 2002. rob@ work: Robot assistant in industrial environments. In *Proceedings. 11th IEEE International Workshop on Robot and Human Interactive Communication*. IEEE, 399–404.
- [15] Herbert Heuer and Jenna Luettgen. 2015. Robot assistance of motor learning: A neuro-cognitive perspective. *Neuroscience & Biobehavioral Reviews* 56 (2015), 222–240.
- [16] Gwyneth Hughes, Sumeet Vadera, Juan Bulacio, and Jorge Gonzalez-Martinez. 2013. Robotic placement of intracranial depth electrodes for long-term monitoring: Utility and efficacy. *Cleveland Clinic* (2013).
- [17] Matjaz Jakopc, F Rodriguez y Baena, Simon J Harris, Paula Gomes, Justin Cobb, and Brian L Davies. 2003. The hands-on orthopaedic robot" Acrobat": Early clinical trials of total knee replacement surgery. *IEEE Transactions on Robotics and Automation* 19, 5 (2003), 902–911.
- [18] Michael Land and Benjamin Tatler. 2009. *Looking and acting: vision and eye movements in natural behaviour*. Oxford University Press.
- [19] Benjamin Law, M Stella Atkins, Arthur E Kirkpatrick, and Alan J Lomax. 2004. Eye gaze patterns differentiate novice and experts in a virtual laparoscopic surgery training environment. In *Proceedings of the 2004 symposium on Eye tracking research & applications*. ACM, 41–48.
- [20] Guillaume Morel Lin Dong. 2021. Control Strategy at Different Instrument Points Using Lever Model in Laparoscopic Surgery. *Accepted:IEEE ICARM 2021* (2021).

- 937 [21] Jesus Mago, Mario Aricò, Jimmy Da Silva, and Guillaume Morel. 2019. Safe teleoperation of a laparoscope holder with dynamic precision but low
938 stiffness. In *2019 International Conference on Robotics and Automation (ICRA)*. IEEE, 2693–2699.
- 939 [22] Jeremy A Marvel, Shelly Bagchi, Megan Zimmerman, and Brian Antonishek. 2020. Towards Effective Interface Designs for Collaborative HRI in
940 Manufacturing: Metrics and Measures. *ACM Transactions on Human-Robot Interaction (THRI)* 9, 4 (2020), 1–55.
- 941 [23] L Mondada. 2014. Instructions in the operating room: How the surgeon directs their assistant’s hands. *Discourse Studies* 2 (2014), 131–161.
- 942 [24] Guillaume Morel, Jérôme Szewczyk, and Marie-Aude Vitrani. 2013. Comanipulation. *Medical Robotics* (2013), 303–350.
- 943 [25] Ilana Nisky, Felix Huang, Amit Milstein, Carla M Pugh, Ferdinando A Mussa-Ivaldi, and Amir Karniel. 2012. Perception of stiffness in laparoscopy—the
944 fulcrum effect. *Studies in health technology and informatics* 173 (2012), 313.
- 945 [26] Society of American Gastrointestinal and Endoscopic Surgeons. 2010. Fundamentals of laparoscopic surgery (FLS). <http://www.flsprogram.org>.
- 946 [27] Raja Parasuraman and Victor Riley. 1997. Humans and automation: Use, misuse, disuse, abuse. *Human factors* 39, 2 (1997), 230–253.
- 947 [28] Andrew D Pearle, Daniel Kendoff, Volker Stueber, Volker Musahl, and John A Repicci. 2009. Perioperative management of unicompartmental knee
arthroplasty using the MAKO robotic arm system (MAKOplasty). *American Journal of Orthopedics* 38, 2 (2009), 16–19.
- 948 [29] Hannah RM Pelikan, Amy Cheatle, Malte F Jung, and Steven J Jackson. 2018. Operating at a Distance—How a Teleoperated Surgical Robot
949 Reconfigures Teamwork in the Operating Room. *Proceedings of the ACM on Human-Computer Interaction* 2, CSCW (2018), 1–28.
- 950 [30] Harsha V Polavarapu, Afif N Kulaylat, Susie Sun, and OH Hamed. 2013. 100 years of surgical education: the past, present, and future. *Bull Am Coll
951 Surg* 98, 7 (2013), 22–27.
- 952 [31] Cécile Poquet, Pierre Mozer, Marie-Aude Vitrani, and Guillaume Morel. 2014. An endorectal ultrasound probe comanipulator with hybrid actuation
953 combining brakes and motors. *IEEE/ASME Transactions on Mechatronics* 20, 1 (2014), 186–196.
- 954 [32] Rebecca Randell, Natasha Alvarado, Stephanie Honey, Joanne Greenhalgh, Peter Gardner, Arron Gill, David Jayne, Alwyn Kotze, Alan Pearman,
955 and Dawn Dowding. 2015. Impact of robotic surgery on decision making: perspectives of surgical teams. In *AMIA Annual Symposium Proceedings*,
Vol. 2015. American Medical Informatics Association, 1057.
- 956 [33] Rebecca Randell, Stephanie Honey, Natasha Alvarado, Alan Pearman, Joanne Greenhalgh, Andrew Long, Peter Gardner, Arron Gill, David Jayne,
957 and Dawn Dowding. 2016. Embedding robotic surgery into routine practice and impacts on communication and decision making: a review of the
958 experience of surgical teams. *Cognition, Technology & Work* 18, 2 (2016), 423–437.
- 959 [34] Rebecca Randell, SA Honey, Jon Hindmarsh, Natasha Alvarado, Joanne Greenhalgh, Alan Pearman, Andrew Long, Alexandra Cope, Arron Gill,
960 Peter Gardner, et al. 2017. A realist process evaluation of robot-assisted surgery: integration into routine practice and impacts on communication,
961 collaboration and decision-making. *Health Services and Delivery Research* 5, 20 (2017).
- 962 [35] François Schmitt, Josue Sulub, Ignacio Avellino, Jimmy Da Silva, Laurent Barbé, Olivier Piccin, Bernard Bayle, and Guillaume Morel. 2019. Using
963 manipulation with active force feedback to undistort stiffness perception in laparoscopy. In *2019 International Conference on Robotics and
964 Automation (ICRA)*. IEEE, 3902–3908.
- 965 [36] Intuitive Surgical. 2019. Intuitive Surgical Announces Preliminary Fourth Quarter and Full Year 2018 Results. [https://isrg.intuitive.com/news-
966 releases/news-release-details/intuitive-surgical-announces-preliminary-fourth-quarter-and-4](https://isrg.intuitive.com/news-releases/news-release-details/intuitive-surgical-announces-preliminary-fourth-quarter-and-4).
- 967 [37] Russell Taylor, Pat Jensen, Louis Whitcomb, Aaron Barnes, Rajesh Kumar, Dan Stoianovici, Puneet Gupta, ZhengXian Wang, Eugene Dejuan, and
968 Louis Kavoussi. 1999. A steady-hand robotic system for microsurgical augmentation. *The International Journal of Robotics Research* 18, 12 (1999),
1201–1210.
- 969 [38] C Poquet Torterot, MA Vitrani, and G Morel. 2014. Proximal comanipulation of a minimally invasive surgical instrument to emulate distal forces.
970 In *Proceedings of the 4th Joint Workshop on New Technologies for Computer/Robot Assisted Surgery (CRAS2014)*. 48–51.
- 971 [39] T. Tsumugiwa, R. Yokogawa, and K. Hara. 2002. Variable impedance control based on estimation of human arm stiffness for human-robot cooperative
972 calligraphic task. In *Proceedings 2002 IEEE International Conference on Robotics and Automation (Cat. No.02CH37292)*, Vol. 1. 644–650 vol.1.
- 973 [40] Marie-Aude Vitrani, Cécile Poquet, and Guillaume Morel. 2016. Applying virtual fixtures to the distal end of a minimally invasive surgery instrument.
974 *IEEE Transactions on Robotics* 33, 1 (2016), 114–123.
- 975 [41] Kristian Wasen. 2010. Replacement of highly educated surgical assistants by robot technology in working life: Paradigm shift in the service sector.
International Journal of Social Robotics 2, 4 (2010), 431–438.
- 976 [42] Mark Wilson, John McGrath, Samuel Vine, James Brewer, David Defriend, and Richard Masters. 2010. Psychomotor control in a virtual laparoscopic
977 surgery training environment: gaze control parameters differentiate novices from experts. *Surgical endoscopy* 24, 10 (2010), 2458–2464.
- 978 [43] Mark R Wilson, Jamie M Poolton, Neha Malhotra, Karen Ngo, Elizabeth Bright, and Rich SW Masters. 2011. Development and validation of a
979 surgical workload measure: the surgery task load index (SURG-TLX). *World journal of surgery* 35, 9 (2011), 1961.
- 980 [44] Mark R Wilson, Samuel J Vine, Elizabeth Bright, Rich SW Masters, David Defriend, and John S McGrath. 2011. Gaze training enhances laparoscopic
981 technical skill acquisition and multi-tasking performance: a randomized, controlled study. *Surgical endoscopy* 25, 12 (2011), 3731–3739.
- 982 [45] S Yule, Rhona Flin, S Paterson-Brown, and N Maran. 2006. Non-technical skills for surgeons in the operating room: a review of the literature.
983 *Surgery* 139, 2 (2006), 140–149.
- 984
985
986
987
988