



HAL
open science

A FORMAL MODEL OF COLEMAN FAMILIES AND APPLICATIONS TO IWASAWA INVARIANTS

Filippo Alberto Edoardo Nuccio Mortarino Majno Di Capriglio, Tadashi Ochiai, Jishnu Ray

► **To cite this version:**

Filippo Alberto Edoardo Nuccio Mortarino Majno Di Capriglio, Tadashi Ochiai, Jishnu Ray. A FORMAL MODEL OF COLEMAN FAMILIES AND APPLICATIONS TO IWASAWA INVARIANTS. 2023. hal-03355637v2

HAL Id: hal-03355637

<https://hal.science/hal-03355637v2>

Preprint submitted on 17 Mar 2023 (v2), last revised 26 Mar 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A FORMAL MODEL OF COLEMAN FAMILIES AND APPLICATIONS TO IWASAWA INVARIANTS

FILIPPO A. E. NUCCIO MORTARINO MAJNO DI CAPRIGLIO, TADASHI OCHIAI,
AND JISHNU RAY

ABSTRACT. For a given Coleman family of modular forms, we construct a formal model and prove the existence of a family of Galois representations associated to the Coleman family. As an application, we study the variations of Iwasawa λ - and μ -invariants of dual fine (strict) Selmer groups over the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} in Coleman families of modular forms. This generalizes an earlier work of Jha and Sujatha for Hida families.

RÉSUMÉ. Nous construisons un modèle formel pour toute famille de Coleman de formes modulaires, et nous montrons l'existence d'une famille de représentations galoisiennes qui lui est attachée. Grâce à cette construction nous étudions la variation des invariants d'Iwasawa λ et μ attachés aux deux des groupes de Selmer fins (strictes) le long de la \mathbb{Z}_p -extension cyclotomique de \mathbb{Q} dans des familles de Coleman de formes modulaires. Nos résultats généralisent ceux obtenus par Jha et Sujatha dans le cas de familles de Hida de formes modulaires ordinaires.

1. INTRODUCTION

Throughout the article, we denote by p a prime number greater than 3, and we let N be a positive integer that is prime to p .

In the celebrated papers [Col97a] and [Col97b], Coleman constructed p -adic families of elliptic cuspforms with fixed slope, often referred to as *Coleman families*. Coleman's theory is a non-ordinary generalization of Hida's theory, yet some of the properties of Coleman families are not completely parallel to those of Hida families. For example, Hida families are usually defined over the whole weight space, but Coleman families are usually defined only locally in the weight space. Also, a Hida family is defined over an Iwasawa algebra, or over a complete and semi-local ring which is finite over an Iwasawa algebra, whereas a Coleman family as constructed in [Col97b] is defined over an affinoid algebra. In particular, Wiles' method of pseudo-representations does not apply *verbatim* to attach a family of Galois representations to a Coleman family because affinoid algebras are not semi-local. Yet, this is the more natural setting to study Iwasawa-theoretic questions, that are the main focus of our work. Our aim is to analyse the behaviour of some Iwasawa invariants in Coleman families, and for this we need a convenient algebraic framework to simultaneously treat the “big” p -adic Galois representations over the whole Coleman family, as well as the “specialized” ones attached to each cuspform in the family. Moreover, the study of integrality

1991 *Mathematics Subject Classification*. Primary 11F33, 11R23; Secondary 11F80.

Key words and phrases. fine Selmer group, Iwasawa invariants, Coleman family, modular forms.

questions, or growth of denominators, is crucial for defining Iwasawa’s μ -invariant and this analysis makes sense only over integral domains where p is not invertible. Thus we devote the first half of the paper to detailing a construction of an integral formal model for Coleman families over a complete local $\mathcal{O}_{\mathcal{K}}$ -algebra $\mathbf{A}_{\mathcal{K}}^{\circ}$ where $\mathcal{O}_{\mathcal{K}}$ is the ring of integers of a finite extension \mathcal{K} of \mathbb{Q}_p . We do not recall a precise definition of $\mathbf{A}_{\mathcal{K}}^{\circ}$ here, rather pointing the reader to (4.1). The spectrum of $\mathbf{A}_{\mathcal{K}}^{\circ}$ contains a certain countable infinite set $Z_{(k_0,i)}[r_0]$ which is identified with a subset of $\mathbb{Z}_{\geq 2}$ (see (4.3) for the definition of $Z_{(k_0,i)}[r_0]$), so that for every $k \in Z_{(k_0,i)}[r_0]$ we can consider the unique classical cuspform f_k that belongs to the Coleman family—or that passes through it, as one can say from a more geometrical viewpoint. Denote by ρ_k the continuous, two-dimensional \mathbb{Q}_p -representation attached by Deligne to f_k : our first result goes in the direction of p -adically interpolating these Galois representations. Explicitly, we prove the following theorem that attaches a “big” Galois representation to the given Coleman family by the method of pseudo-representation: we refer to Theorem 4.3 for the precise statement and the proof, here it suffices to say that S is a finite set of rational primes containing p , and P_k is the prime ideal in $\mathbf{A}_{\mathcal{K}}^{\circ}$, corresponding to the point of $\text{Spec } \mathbf{A}_{\mathcal{K}}^{\circ}$ identified with the integer k , and such that $\mathbf{A}_{\mathcal{K}}^{\circ}/P_k$ is a finite \mathbb{Z}_p -module.

Theorem A. *There exists a free $\mathbf{A}_{\mathcal{K}}^{\circ}$ -module \mathbf{T} of rank two with a continuous action of unramified-outside- S Galois group $G_{\mathbb{Q},S}$ such that the representation*

$$\rho: G_{\mathbb{Q},S} \longrightarrow \text{Aut}_{\mathbf{A}_{\mathcal{K}}^{\circ}}(\mathbf{T})$$

satisfies $\pi_{\rho} = \pi$ (in a suitable basis). In particular, ρ modulo the ideal P_k corresponding to k is isomorphic to a lattice of ρ_k for all $k \in Z_{(k_0,i)}[r_0]$.

The construction of a big Galois representation attached to a Coleman family is not new. Already in the foundational articles [Col97b] and [CM98] the authors described it, and another approach (based on the theory of modular symbols) was later proposed by Bellaïche in [Bel12]. In these works, the Galois representation is supported on a rank-2 module over a Tate algebra, or possibly over the subalgebra of power-bounded elements in such a Tate algebra. This ring is not semi-local and therefore these constructions were not suited for our specialization techniques for fine Selmer groups in the setting of Iwasawa theory. The definition of a formal model for the Galois representation attached to a Coleman family that we present in this paper was first introduced in the preprint [NO16], that has received some citations but has finally not been published. We take the opportunity to present the construction here, together with its Iwasawa-theoretic applications. Independently from ours, another construction of a formal model is explained in [BL21, Remark 1.4 and Remark A.1] and relies on Loeffler–Zerbes’ work [LZ16], where the representation arises in the cohomology of “big” étale sheaves on modular curves as defined in [AIS15]. Our approach differs from theirs, as it mainly relies on the theory of pseudo-representations as in Wiles’ original work [Wil88] (see also Hida’s monograph [Hid93]). It has the advantage that it does not use any specific context of elliptic modular forms so that it can be generalised to the setting of Coleman families associated to more general Shimura varieties.

In the second half of this article, we analyse some Iwasawa-theoretic properties of the family of Galois representations constructed in the above theorem. This generalizes results

of Greenberg–Vatsal (see [GV00]) and of Emerton–Pollack–Weston (see [EPW06]), who studied the variations of Iwasawa λ - and μ -invariants of dual Selmer groups over the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} for families of p -ordinary modular forms. More precisely, let $\bar{r}: G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\kappa)$ be an absolutely irreducible modular Galois representation of $G_{\mathbb{Q}}$ over a finite field κ of characteristic p . Assume that \bar{r} is p -ordinary and p -distinguished in the sense that the restriction of \bar{r} to a decomposition subgroup at p is reducible and non-scalar (see [EPW06, p. 523]), and let $\mathcal{H}(\bar{r})$ be the Hida family of \bar{r} . Emerton–Pollack–Weston showed that if the μ -invariant for the dual Selmer group for some $f_0 \in \mathcal{H}(\bar{r})$ is trivial, then it is so for all $f \in \mathcal{H}(\bar{r})$. Further, they showed that, under the assumption that the μ -invariant is zero, the λ -invariant is constant across a branch in the Hida family $\mathcal{H}(\bar{r})$.

Our goal is to study an Emerton–Pollack–Weston type result for the *fine* Selmer group in a Coleman family. The fine Selmer group is a much studied object in Iwasawa theory and occurs in the formulation of the Iwasawa Main Conjecture for elliptic curves and modular forms. In [CS05], Coates and Sujatha initiated a systematic study of the fine Selmer group and formulated conjectures about its structure over a p -adic Lie extension. Under certain hypotheses, they have shown that Iwasawa’s μ -invariant Conjecture for the cyclotomic extension of a number field is equivalent to Conjecture A on the dual fine Selmer group of an elliptic curve (see [CS05, Theorem 3.4]). Conjecture A says that the μ -invariant of the dual fine Selmer group of an elliptic curve over the cyclotomic \mathbb{Z}_p -extension of a number field is trivial. Observe that the Pontryagin dual of the fine Selmer group over the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} is proved to be torsion by a deep result of Kato [Kat04, Theorem 12.4] regardless of whether f is ordinary at p or not. An interested reader might also want to view the Iwasawa Main Conjecture framed in terms of dual fine Selmer group; see [Kur02, Conjecture 6.1] or [Kob03, Proposition 7.1 (ii)] and [Kat04, Conjecture 12.10]. This is the version of the Iwasawa Main Conjecture without p -adic L -functions, framed in terms of fine Selmer groups, and this is what is usually considered when dealing with non-ordinary families (see [Nak20, Section 5]). This justifies the use of fine Selmer groups in this paper; the construction of plus and minus Selmer groups over a Coleman family is unattainable by known techniques.

Variations of Iwasawa invariants for the dual fine Selmer group in Hida families of ordinary modular forms have been studied by Jha and Sujatha in [JS11]. In this article, we go beyond their work and study these variations for the dual fine Selmer group when the modular forms vary in a Coleman family. We define a universal “big” fine Selmer group for the Galois representation \mathbf{T} produced in Theorem A, and we show that it is cotorsion over a two-variables Iwasawa algebra. Further, we specialize this big fine Selmer group at classical points and study its relation with the classical fine Selmer group (see Theorem 6.3). In this context, our main result is the following, where ρ is as in Theorem A:

Theorem B (see Theorems 7.1 and 7.3). *Assume that the residual representation of ρ when restricted to $G_{\mathbb{Q}_p}$ is irreducible. Given any $k \in Z_{(k_0, i)}[r_0]$, the μ -invariant of the dual fine Selmer group associated to ρ_k is zero if and only if the same is true for any other classical modular form in the Coleman family. Under the assumption that the μ -invariant of one, or of any, form is zero, the λ -invariants are constant for all but finitely many classical modular forms in the Coleman family.*

Other authors have studied the question of the variation of Iwasawa invariants in families of congruent modular forms. In [Kim09], Kim studied Iwasawa λ - and μ -invariants for the dual signed Selmer groups over the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} for elliptic curves at supersingular primes. He proved that they are constant for a family of elliptic curves with the same residual representation if the μ -invariant of any of them is zero. Kim's result was later generalized by the first-named author and Sujatha in [NS21] in a more general, but still supersingular, setting. Under Conjecture A and a few other additional assumptions they could give a criterion when the μ -invariant of the dual signed Selmer group vanishes purely in terms of a signed *residual* Selmer group (see [NS21, Theorem 4.13]). Variations of Iwasawa invariants for dual fine Selmer groups were also studied by Kim, Lee and Ponsinet in [KLP19] but there the authors considered a family of modular forms with fixed weight and varying tame levels, contrary to our set-up. Hatley and Lei considered the case of the variations of Iwasawa λ - and μ -invariants for dual signed Selmer groups for non-ordinary congruent modular forms and proved a result similar to Kim's (see [HL19, Theorem 4.6]). In [LS18], Lim and Sujatha studied fine Selmer groups of two congruent Galois representations over an admissible p -adic Lie extension and showed that, under appropriate congruence conditions, if the dual fine Selmer group of one form is pseudo-null, the same holds for the other form (see [LS18, Theorem 3.7]).

The structure of the paper is as follows. In Section 2, we prepare some generalities on affinoid algebras and on the weight space in relation to Coleman families. Based on this preparation, we construct a formal model of a given Coleman family in Section 3. In Section 4, we explain the construction of the Galois representation giving the proof of Theorem A (in the precise formulation of Theorem 4.3). In Section 5, we review basic facts on Galois deformations of Coleman families and construct the big fine Selmer group associated to a Coleman family. The specialization theorem at an arithmetic prime is given in Section 6. We use this specialization theorem to show that the big fine Selmer group is cotorsion over a two-variables Iwasawa algebra. Our results about variations of the Iwasawa invariants, summarized in Theorem B, are obtained in Section 7.

ACKNOWLEDGEMENT

The authors thank Adrian Iovita and Somnath Jha for answering several questions and for their comments on an earlier draft of this article. We warmly thank the two referees for a very carefully reading and for suggesting several corrections that helped improving the presentation and the mathematics of our manuscript. The third author is supported by the Inspire research grant.

2. PRELIMINARIES ON AFFINIDS AND WEIGHT SPACES

We refer the reader to [BGR84] for our conventions and basic results about rigid analytic spaces in the sense of Tate. Recall that p is a fixed odd prime and $N \geq 1$ satisfies $(p, N) = 1$.

Let K be a complete subfield of \mathbb{C}_p . The field K can be either a finite or an infinite extension of \mathbb{Q}_p . Let \mathcal{X} be an affinoid space defined over K . We write $\mathcal{A}_{\mathcal{X}}$ for the

ring of analytic functions on \mathcal{X} and $\mathcal{A}_{\mathcal{X}}^{\circ}$ for the subring of power-bounded elements (see [BGR84, Section 1.2.5]). They will always be endowed with their Gauß semi-norm (which is a norm and coincides with the sup-norm if \mathcal{X} is reduced). When K is a discrete valuation field, the ring of power-bounded elements $\mathcal{A}_{\mathcal{X}}^{\circ}$ is noetherian because it is a quotient of $\mathcal{O}_K\langle T_1, \dots, T_n \rangle$ and $\mathcal{O}_K\langle T_1, \dots, T_n \rangle$ is the p -adic completion of a polynomial algebra over \mathcal{O}_K . The ring $\mathcal{A}_{\mathcal{X}} = \mathcal{A}_{\mathcal{X}}^{\circ}[\frac{1}{p}]$ is noetherian whether K is a discrete valuation field or not. For every maximal ideal $\mathfrak{m} \subseteq \mathcal{A}_{\mathcal{X}}$, the quotient $\mathcal{A}_{\mathcal{X}}/\mathfrak{m}$ is a finite extension of K by [BGR84, Section 6.1.2, Corollary 3] and $\mathcal{A}_{\mathcal{X}}^{\circ}/\mathfrak{m}^{\circ}$ is a domain which is finite over \mathcal{O}_K by [BGR84, Section 6.1.3, Proposition 3].

We consider the following definition:

Definition 2.1. *Let K be a complete subfield of \mathbb{C}_p and let \mathcal{X} be an affinoid space over K . A subset \mathcal{Z} of the set of K -valued points $\mathcal{X}(K)$ is said to be Zariski-dense in \mathcal{X} if we have $U(K) \cap \mathcal{Z} \neq \emptyset$ for every non-empty Zariski-open subspace $U \subseteq \mathcal{X}$.*

Given $x_0 \in K$ and $r \in p^{\mathbb{Q}}$, we denote by $\mathcal{B}[x_0, r]_K$ and $\mathcal{B}(x_0, r)_K$, respectively, the closed and open ball of radius r and centre x_0 , seen as K -rigid analytic spaces (see [dJ95, Section 7] for a description of the second space). We note that we normalize the p -adic absolute value $|\cdot|$ so that $|p| = \frac{1}{p}$. For example, in the case $r = 1$ and $x_0 \in K$, $\mathcal{A}_{\mathcal{B}[x_0, 1]_K}^{\circ}$ is isomorphic to the following ring of restricted power series with coefficients in \mathcal{O}_K :

$$\mathcal{O}_K\langle T - x_0 \rangle = \left\{ \sum_{i=0}^{\infty} c_i (T - x_0)^i \in \mathcal{O}_K[[T - x_0]] \mid \lim_{i \rightarrow \infty} |c_i| = 0 \right\}.$$

Finally, given any complete subfield $L \subseteq \mathbb{C}_p$, we also need the notation $B[a, r]_L$ and $B(a, r)_L$ for the set of all $x \in L$ such that $|x - a| \leq r$ (respectively, such that $|x - a| < r$). When $K = \mathbb{Q}_p$, we denote $B[x_0, r]_{\mathbb{Q}_p}$ (resp. $B(x_0, r)_{\mathbb{Q}_p}$) by $B[x_0, r]$ (resp. $B(x_0, r)$) dropping the subscript.

Lemma 2.2. *Let K be a complete subfield of \mathbb{C}_p which is a discrete valuation field.*

- (1) *When \mathcal{X} is a reduced affinoid defined over K and $f \in \mathcal{A}_{\mathcal{X}}$ vanishes on every point of a Zariski-dense subset Z , we have $f = 0$.*
- (2) *Let $x_0 \in K$. Every infinite set inside $\mathcal{B}[x_0, 1]_K(K)$ is Zariski-dense.*

Proof. For the first assertion, suppose $f \neq 0$ and consider the Zariski-open subset $U_f = \{x \in \mathcal{X} \text{ such that } f(x) \neq 0\}$ of \mathcal{X} . Since \mathcal{X} is reduced and $f \neq 0$, we have $U_f \neq \emptyset$. By the assumption that $Z \subset \mathcal{X}$ is a Zariski-dense subset, we have $Z \cap U_f(K) \neq \emptyset$. For any point $z \in Z \cap U_f(K)$, we have $f(z) = 0$, contradicting the definition of U_f .

We pass to the second assertion. By the Weierstraß Preparation Theorem ([BGR84, Section 5.2.2, Theorem 1]), every function $f \in \mathcal{A}_{\mathcal{B}[x_0, 1]_K}$ can be factored as $f = P \cdot U$ where $P \in K[T - x_0]$ is a polynomial and $U \in \mathcal{A}_{\mathcal{B}[x_0, 1]_K}^{\times}$ is an invertible power series which does not vanish on $\mathcal{B}[x_0, 1]_K$. It follows that every such f has only finitely many zeroes and that $\mathcal{A}_{\mathcal{B}[x_0, 1]_K}$ is a PID, showing that non-trivial Zariski-closed sets in $\mathcal{B}[x_0, 1]_K$ consist of finitely many points. \square

One of the main rigid spaces of interest for us is the weight space \mathscr{W}_N , which is isomorphic to $\varphi(Np)$ copies of $\mathscr{B}(1, 1)_K$ indexed by

$$\mathfrak{D} = \text{Hom}((\mathbb{Z}/Np\mathbb{Z})^\times, \mathbb{C}_p^\times).$$

For generalities about \mathscr{W}_N , we refer to [Col97b, Section B1] and to [CM98, Section 1.4]. For more detailed accounts, we refer to [Gou88, Chapter I.3, §4 and Appendix] and [Buz07, p. 103]. By definition, the weight space satisfies

$$\mathscr{W}_N(\mathbb{C}_p) = \text{Hom}_{\text{cont}}(\varprojlim_n (\mathbb{Z}/Np^n\mathbb{Z})^\times, \mathbb{C}_p^\times).$$

Following Coleman and Mazur, we give the following definition:

Definition 2.3. Denote by $\omega: \mu_p(\mathbb{Z}_p) \rightarrow \mathbb{Z}_p^\times$ the Teichmüller character and denote the projection $x \mapsto x/\omega(x)$ by $\langle\langle \cdot \rangle\rangle: \mathbb{Z}_p^\times \rightarrow 1+p\mathbb{Z}_p$. For every integer $k \in \mathbb{Z}$ and every character $\chi \in \mathfrak{D}$ of finite order, the point $\chi \langle\langle \cdot \rangle\rangle^k \in \mathscr{W}_N(\mathbb{Q}_p)$ is called an accessible weight-character with coordinates (χ, k) .

As detailed in [CM98, Definition in Section 1.4] the accessible weight-characters are parametrized by the rigid analytic subspace $\mathscr{W}_N^* = \mathfrak{D} \times \mathscr{B}^* \subseteq \mathscr{W}_N$ where \mathscr{B}^* is the subdisk of $\mathscr{B}(1, 1)$ which is the isomorphic image of $\mathscr{B}(0, p^{\frac{p-2}{p-1}})$ via the map $s \mapsto (1+p)^s$, so

$$(2.1) \quad \mathscr{B}(0, p^{\frac{p-2}{p-1}}) \cong \mathscr{B}^* \subseteq \mathscr{B}(1, 1).$$

In the notation introduced in Definition 2.3, the character $\chi \langle\langle \cdot \rangle\rangle^k$ is represented by the point $(\chi, (1+p)^k) \in \mathscr{W}_N(\mathbb{Q}_p)$ which gets mapped to (χ, k) by the identification in (2.1); we see that the word “coordinates” comes from seeing \mathscr{W}_N^* as $\varphi(Np)$ -copies of $\mathscr{B}(0, p^{\frac{p-2}{p-1}})$. From now on we systematically write points in the weight space through their coordinates. In particular, for every fixed $\chi \in \mathfrak{D}$, the weights of characters with Nebentypus χ will be regarded as points in $\mathscr{B}(0, p^{\frac{p-2}{p-1}})$ rather than in $\mathscr{B}(1, 1)$.

The assumption $(N, p) = 1$ allows us to consider the group $\text{Hom}((\mathbb{Z}/p)^\times, \mathbb{C}_p^\times)$ as being a subgroup of \mathfrak{D} . It thus makes sense, for each $0 \leq j \leq p-2$, to interpret ω^j as an element of \mathfrak{D} . The characters $x \mapsto x^k$, which are accessible with coordinates $(1, k)$, are then the elements of $\mathscr{W}_N^*(\mathbb{Q}_p)$ which belong to the ω^k -th copy of \mathscr{B}^* .

3. p -ADIC FAMILIES OF MODULAR FORMS

We start with a classical eigencuspform $f \in S_{k_0}(\Gamma_1(Np), \varepsilon)$ of weight k_0 , level Np and Nebentypus ε , which we factor as a product $\varepsilon = \varepsilon_N \omega^{k_0-i}$ for some character ε_N of conductor divisible by N and some $0 \leq i \leq p-1$. Our main references concerning p -adic modular forms and p -adic families thereof are [Col97b, Part B] as well as [Gou88], in particular Section II.3 *ibid.* for the definition of the U_p -operator.

Definition 3.1. Let f be a p -adic modular form of level Np which is an eigenvector with respect to the U_p -operator. We define the slope of f to be the p -adic valuation of the U_p -eigenvalue of f . It is a non-negative rational number.

We assume that the slope of f is $0 \leq \alpha \leq k_0 - 1$. In [Col97b], [Col97a] and [CM98] Coleman and Coleman–Mazur have built a theory of families of p -adic modular forms of slope α interpolating f , a part of which is stated in Theorem 3.2 below. Let us introduce some notation. Given an element $k_0 \in \mathbb{Q}_p$, an integer i and an element $r \in p^{\mathbb{Q}}$ with $r < p^{\frac{p-2}{p-1}}$, we denote by $\mathcal{X}_{(k_0,i)}[r]$ the affinoid subspace of the weight space defined as

$$\mathcal{X}_{(k_0,i)}[r_0] := \{\varepsilon_N \omega^i\} \times \mathcal{B}[k_0, r] \subseteq \mathcal{W}_N^*.$$

Let us consider a finite extension \mathcal{K} of \mathbb{Q}_p , which will play the role of the field of coefficients of motives associated to cuspforms f in the given Coleman family. We note that \mathcal{K} has nothing to do with the field of definition of motives associated to cuspforms f in the given Coleman family, for which we can choose \mathbb{Q}_p . From now on, we denote by $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]/\mathcal{K}}^\circ$ (resp. $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]/\mathcal{K}}$) the extension of coefficients $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]}^\circ \otimes_{\mathbb{Z}_p} \mathcal{O}_{\mathcal{K}}$ (resp. $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]} \otimes_{\mathbb{Q}_p} \mathcal{K}$) where $\mathcal{O}_{\mathcal{K}}$ is the ring of integers of \mathcal{K} . We have the following result thanks to [Col97b]:

Theorem 3.2 ([Col97b]). *Suppose that f is a classical normalized cuspidal eigenform, of weight k_0 , level $\Gamma_1(Np)$, slope $\alpha < k_0 - 1$, Nebentypus $\varepsilon = \varepsilon_N \omega^{i-k_0}$ and which is new away from p . When $i = 0$, suppose moreover that $a^2 \neq \varepsilon_N(p)p^{k_0-1}$, where a is the U_p -eigenvalue of f .*

Then, there is a radius $r_0 < p^{\frac{p-2}{p-1}}$ lying in $p^{\mathbb{Q}}$ and analytic functions $a_n \in \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r_0]/\mathcal{K}}^\circ$, indexed by $n \in \mathbb{N}$, such that the following statements hold:

- (1) *For all integers $k \in \mathcal{X}_{(k_0,i)}[r_0](\mathbb{Q}_p)$ satisfying $k > \alpha + 1$, the series*

$$\sum_{n=1}^{\infty} a_n(k) q^n \in \mathcal{K}[[q]]$$

coincides with the q -expansion of a classical normalized cuspidal eigenform of level Np , weight k , slope α and character $\varepsilon_N \omega^{i-k}$.

- (2) *The series $\sum_{n=1}^{\infty} a_n(k_0) q^n \in \mathcal{K}[[q]]$ coincides with the q -expansion of f .*
- (3) *The space $\mathcal{X}_{(k_0,i)}[r_0]$ is a_p -small in the sense of [Kis03, (5.2)].*

Remark 3.3. At first glance, it might look better to write $a_n^{(r_0)}$ for the analytic functions appearing in the statement, since the radius r_0 is not uniquely associated to f and these functions might *a priori* depend on its choice. However, we will prove in Corollary 3.6 below that this is not necessary.

Remark 3.4. In this paper, the property that $\mathcal{X}_{(k_0,i)}[r_0]$ be a_p -small will not be used. Nevertheless, point (3) of Theorem 3.2 is essential when dealing with interpolation of crystalline periods along the Coleman family, adapting techniques from [Kis03].

Proof. In [Col97b, pp. 465–467] (and especially along the proof of Corollary B5.7.1 *ibidem*), Coleman attached to the f chosen above the space $\mathcal{X}_{(k_0,i)}[r_0]$ with a radius $r_0 \in p^{\mathbb{Q}}$ small enough. In [Col97b, pp. 465–467], this space is denoted simply by B and a crucial step is to

take a finite étale affinoid algebra $R_{(k_0,i)}[r_0]$ over $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r_0]}$ whose associated affinoid space parametrizes families of p' -new forms of slope α in the sense of [Col97b, p. 467, Definition]. As in [Col97b], we write $X(R_{(k_0,i)}[r_0]) \rightarrow \mathcal{X}_{(k_0,i)}[r_0]$ for the affinoid space associated to $R_{(k_0,i)}[r_0]$.

Then the statements (1)–(3) are in [Col97b, Corollary B5.7.1] *verbatim*, except for the condition that all a_n be power-bounded and that all forms above be normalized. In Lemma B5.3 *ibidem* it is shown that the Hecke eigenvalues of an overconvergent cuspidal eigenform are bounded by 1 if the form is normalized. We thus deduce the result by observing that $a_1 = 1$, which follows from $a_1 = T(1) = 1$ by the construction given in Theorem B5.7 *ibidem*. As discussed in [Kis03, §5.2] it is always possible to shrink a disk around a point in order to get a smaller one which is a_p -small, and we define this smaller radius as the constant r_0 of the statement. \square

Definition 3.5. *Let f be a form as in Theorem 3.2 and let $0 < r \leq r_0$ be smaller than or equal to the radius constructed *ibid*. The collection of the rigid functions $\{a_n\}_{n \in \mathbb{N}}$ on $\mathcal{X}_{(k_0,i)}[r]$ is called the Coleman family of slope α and radius r passing through the form f . The formal power series*

$$\sum_{n=1}^{\infty} a_n q^n \in \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]/\mathcal{K}}^{\circ} \llbracket q \rrbracket$$

is said to be the Fourier expansion of the Coleman family. For each $x \in \mathcal{X}_{(k_0,i)}[r]$, we denote by f_x the overconvergent modular form whose expansion is $\sum a_n(x)q^n$. Sometimes, we also refer to the collection of all these forms as the Coleman family of slope α through the form f .

Consider the subset $\mathcal{Z}_{(k_0,i)}[r]$ of accessible weight-characters in $\mathcal{X}_{(k_0,i)}[r]$ defined as

$$(3.1) \quad \mathcal{Z}_{(k_0,i)}[r] = \left(\{\varepsilon_N \omega^i\} \times \mathbb{Z}_{>\alpha+1} \right) \cap \mathcal{X}_{(k_0,i)}[r] \subseteq \mathcal{W}_N^*(\mathbb{Q}_p).$$

By an abuse of notation, we write k for elements $(\varepsilon_N \omega^i, k)$ of $\mathcal{Z}_{(k_0,i)}[r]$ and we denote the p -adic Deligne representation attached to f_k in [Del71] by ρ_k .

Consider now a form f of weight k_0 satisfying the assumption of Theorem 3.2 and a radius r_0 as constructed *ibid*.

Corollary 3.6. *Let $0 < r < r_0$ be an element of $p^{\mathbb{Q}}$. For $n \geq 1$ the functions*

$$\text{res}_{\mathcal{X}_{(k_0,i)}[r]}^{\mathcal{X}_{(k_0,i)}[r_0]}(a_n) \in \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]/\mathcal{K}}^{\circ}$$

are the Fourier expansion of a Coleman family of slope α and radius r passing through the form f .

Proof. This follows from the definitions, since for every $k \in \mathcal{Z}_{(k_0,i)}[r](\mathbb{Q}_p)$ the series

$$\sum_{n=1}^{\infty} \text{res}_{\mathcal{X}_{(k_0,i)}[r]}^{\mathcal{X}_{(k_0,i)}[r_0]}(a_n)(k) q^n$$

is the q -expansion of a form of the required type. \square

Thanks to the above corollary, we can unambiguously speak about the Fourier coefficients a_n of a Coleman family without referring to the radius; observe also that shrinking an a_p -small disk to one of smaller radius preserves a_p -smallness. We recall the following lemma:

Lemma 3.7. *Let $r \in p^{\mathbb{Q}}$ be a radius satisfying $r \leq r_0$. If the function $G \in \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]}/\mathcal{K}$ vanishes on $\mathcal{Z}_{(k_0,i)}[r_0]$, it is everywhere zero.*

Proof. The subset $\mathcal{Z}_{(k_0,i)}[r] \subseteq \mathcal{X}_{(k_0,i)}[r](\mathbb{Q}_p)$ is Zariski-dense in $\mathcal{X}_{(k_0,i)}[r]$ thanks to the second assertion of Lemma 2.2. Hence, the assertion is an immediate consequence of the first assertion of Lemma 2.2. \square

4. p -ADIC FAMILY OF GALOIS REPRESENTATIONS

The main result of this section is Theorem 4.3 which produces an *integral* Galois representation with values in $\mathcal{A}_{\mathcal{X}}^{\circ}/\mathcal{K}$ – for a given Coleman family and a suitable affinoid \mathcal{X} – that specializes to the Deligne representations attached to the classical eigenforms which belong to the given Coleman family. The most important ingredient is to construct pseudo-representations associated to a given Coleman family: once we have a pseudo-representation over our affinoid algebras, a standard argument allows us to recover a Galois representation from this pseudo-representation. Since we are not aware of a reference on constructing pseudo-representations over affinoid algebras, we construct them here by using a formal structure of Coleman families.

Let us fix a Coleman family as in Theorem 3.2 and let $r_0 \in p^{\mathbb{Q}}$ be the radius constructed *ibid.* Let K be a complete subfield of \mathbb{C}_p and take an element $e_0 \in K$ such that $r_0 = |e_0|$. Define \mathbf{A}_K° to be the complete local ring

$$(4.1) \quad \mathbf{A}_K^{\circ} = \mathcal{O}_K \left[\left[\frac{T - k_0}{e_0} \right] \right] = \left\{ \sum_{i=0}^{\infty} c_i \left(\frac{T - k_0}{e_0} \right)^i \mid c_i \in \mathcal{O}_K \right\}.$$

For any $r \in p^{\mathbb{Q}}$ with $r < r_0$, a power series F in \mathbf{A}_K° converges on $B[k_0, r]_K \subsetneq B(k_0, r_0)_K$ and its evaluations at each point $x \in B[k_0, r]_K$ satisfies $|F(x)| \leq 1$. We consider elements of \mathbf{A}_K° as functions on $\mathcal{X}_{(k_0,i)}[r]$, and this induces, by restriction, a ring homomorphism $\mathbf{A}_K^{\circ} \xrightarrow{\text{res}} \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_K}^{\circ}$. For each radius $r < r_0$, we choose $e_r \in \mathbb{C}_p$ such that $|e_r| = r$. By [BGR84, §6.1.5], there is an isomorphism

$$\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_{\mathbb{C}_p}}^{\circ} \cong \left\{ \sum_{i=0}^{\infty} c_i \left(\frac{T - k_0}{e_r} \right)^i \in \mathcal{O}_{\mathbb{C}_p} \left[\left[\frac{T - k_0}{e_r} \right] \right] \mid \lim_{i \rightarrow \infty} |c_i| = 0 \right\}.$$

Similarly as above, the inclusion $\{\varepsilon_N \omega^i\} \times \mathcal{B}(k_0, r_0)_K \subset \mathcal{X}_{(k_0,i)}[r_0]_K$ induces a homomorphism $\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r_0]_K}^{\circ} \rightarrow \mathbf{A}_K^{\circ}$ where we have a presentation

$$\mathcal{A}_{\mathcal{X}_{(k_0,i)}[r_0]_K}^{\circ} \cong \left\{ \sum_{i=0}^{\infty} c_i \left(\frac{T - k_0}{e_0} \right)^i \in \mathcal{O}_K \left[\left[\frac{T - k_0}{e_0} \right] \right] \mid \lim_{i \rightarrow \infty} |c_i| = 0 \right\}.$$




FIGURE 1.

The composition

$$(4.2) \quad \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r_0]_K}^\circ \longrightarrow \mathbf{A}_K^\circ \xrightarrow{\text{res}} \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_K}^\circ$$

corresponds to the inclusions

$$\mathcal{B}[k_0, r]_{\mathbb{C}_p} \subset \mathcal{B}(k_0, r_0)_K \subset \mathcal{B}[k_0, r_0]_K.$$

In Figure 1, there is a sketch of the radii that have occurred so far in our construction.

When the radius $r \in p^{\mathbb{Q}}$ tends to r_0 from below, we have

$$\mathbf{A}_K^\circ \subset \mathbf{A}_{\mathbb{C}_p}^\circ = \varprojlim_{\substack{r \rightarrow r_0 \\ r < r_0}} \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_{\mathbb{C}_p}}^\circ = \bigcap_{r < r_0} \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_{\mathbb{C}_p}}^\circ.$$

From now on, we denote $\mathbf{A}_{\mathbb{Q}_p}^\circ$ by \mathbf{A}° when $K = \mathbb{Q}_p$ dropping the subscript K . We summarize the above observations in the following proposition:

Proposition 4.1. *Maintain the same assumptions and notations of Theorem 3.2. There exists a finite extension \mathcal{K} of \mathbb{Q}_p such that, for every $n \in \mathbb{N}$, there is a unique function $A_n \in \mathbf{A}_{\mathcal{K}}^\circ$ such that the image of A_n via the restriction $\mathbf{A}_{\mathcal{K}}^\circ \hookrightarrow \mathcal{A}_{\mathcal{X}_{(k_0,i)}[r]_{\mathcal{K}}}^\circ$ coincides with the function a_n obtained in Theorem 3.2 on a smaller radius r .*

Proof. In Theorem 3.2, we obtained a Coleman family over $\mathcal{X}_{(k_0,i)}[r_0]$. By shrinking such a Coleman family to $\mathcal{X}_{(k_0,i)}[r]$ for any $r \in p^{\mathbb{Q}}$ with $r < r_0$ and by taking the limit $r \rightarrow r_0$, we showed in Corollary 3.6 that there exists $A_n \in \mathbf{A}_{\mathcal{K}}^\circ$ such that the image of A_n via the

inclusion $\mathbf{A}_{\mathcal{K}}^{\circ} \hookrightarrow \mathcal{A}_{\mathcal{X}(k_0, i)[r]}^{\circ}/\mathcal{K}$ coincides with a_n obtained in Theorem 3.2. This completes the proof of the proposition. \square

Based on the existence of the formal structure $\mathbf{A}_{\mathcal{K}}^{\circ}$, we introduce the notion of 2-dimensional pseudo-representation based on [Wil88]. We remark that other types of pseudo-representations were later introduced by Taylor in [Tay91] and, much more recently, by Chenevier in [Che11] but for our purposes Wiles' approach seems to be the best suited.

Given a topological group G , a topological ring R in which 2 is invertible, and three functions $A, D: G \rightarrow R$ and $\Xi: G \times G \rightarrow R$, we consider the following four properties given in [Wil88, Lemma 2.2.3]:

- (I) A, D and Ξ are continuous functions.
- (II) The relations

$$A(\sigma\tau) = A(\sigma)A(\tau) + \Xi(\sigma, \tau),$$

$$D(\sigma\tau) = D(\sigma)D(\tau) + \Xi(\tau, \sigma),$$

$$\Xi(\sigma\tau, \rho\gamma) = A(\sigma)A(\gamma)\Xi(\tau, \rho) + A(\gamma)D(\tau)\Xi(\sigma, \rho) + A(\sigma)D(\rho)\Xi(\tau, \gamma) + D(\tau)D(\rho)\Xi(\sigma, \gamma)$$

hold for all elements $\sigma, \tau, \gamma, \rho \in G$.

- (III) The relations $A(\mathbf{1}) = D(\mathbf{1}) = 1$ and $\Xi(\sigma, \mathbf{1}) = \Xi(\mathbf{1}, \tau) = 0$ hold for all elements $\sigma, \tau \in G$ where $\mathbf{1}$ is the unit element of G .

- (IV) The relation $\Xi(\sigma, \tau)\Xi(\rho, \eta) = \Xi(\sigma, \eta)\Xi(\rho, \tau)$ holds for all elements $\sigma, \tau, \rho, \eta \in G$.

We call a triple (A, D, Ξ) satisfying the above conditions (I)–(IV) a pseudo-representation.

Given a continuous representation $\rho: G \rightarrow \mathrm{GL}_2(R)$, by fixing a basis of R^2 we can write

$$\rho(\sigma) = \begin{pmatrix} a(\sigma) & b(\sigma) \\ c(\sigma) & d(\sigma) \end{pmatrix} \quad \text{for } \sigma \in G$$

and the triple $\pi_{\rho} = (A(\sigma) = a(\sigma), D(\sigma) = d(\sigma), \Xi(\sigma, \tau) = b(\sigma)c(\tau))$ is easily checked to be a pseudo-representation; observe that attaching a pseudo-representation to a representation depends on the choice of a basis. Extending the notation introduced in Definition 3.5, and in the set-up introduced *ibid*, we simply denote by π_k the pseudo-representation π_{ρ_k} attached to ρ_k in some chosen basis.

Let $Z_{(k_0, i)}[r_0]$ be the set of points inside $\mathcal{W}_N^*(\mathbb{Q}_p)$ defined as the intersection

$$(4.3) \quad Z_{(k_0, i)}[r_0] = \mathcal{B}(k_0, r_0)(\mathbb{Q}_p) \cap \mathcal{Z}_{(k_0, i)}[r_0]$$

where $\mathcal{Z}_{(k_0, i)}[r_0]$ is the set defined in (3.1). According to the convention established at the end of Section 2, weights in $Z_{(k_0, i)}[r_0]$ will be identified with integers and denoted by a single coordinate.

Proposition 4.2. *Let S be the finite set of primes of \mathbb{Q} consisting of the primes $\{\ell : \ell \mid N\}$, p and ∞ , and denote by $G_{\mathbb{Q}, S}$ the Galois group of the maximal extension of \mathbb{Q} unramified outside S . Then, there exists a continuous pseudo-representation*

$$\pi = (A, D, \Xi): G_{\mathbb{Q}, S} \rightarrow \mathbf{A}_{\mathcal{K}}^{\circ}$$

interpolating the pseudo-representations π_k attached to members of the Coleman family of slope α through f . In other words, for each $k \in Z_{(k_0, i)}[r_0]$, the evaluation $\text{ev}_k \circ \pi = (\text{ev}_k \circ A, \text{ev}_k \circ D, \text{ev}_k \circ \Xi)$ coincides with the pseudo-representation π_k .

Proof. The argument relying on pseudo-representations of rank two à la Wiles is more or less standard and the proof goes in a parallel manner as the proof given in the monograph [Hid93, §7.5], so we only give an outline.

Let us choose a complex conjugation $c \in G_{\mathbb{Q}, S}$. For each $k \in Z_{(k_0, i)}[r_0]$, we fix a basis for the representation ρ_k so that c is represented by the matrix $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$. Following [Wil88, Lemma 2.3.3], we define functions

$$A_{(k)}, D_{(k)}: G_{\mathbb{Q}, S} \longrightarrow \mathcal{K}$$

by the matrix representation

$$\rho_k(g) = \begin{pmatrix} A_{(k)}(g) & B_{(k)}(g) \\ C_{(k)}(g) & D_{(k)}(g) \end{pmatrix}.$$

We also set $\Xi_{(k)}: G_{\mathbb{Q}, S} \times G_{\mathbb{Q}, S} \longrightarrow \mathcal{K}$ by $\Xi_{(k)}(\gamma_1, \gamma_2) = A_{(k)}(\gamma_1 \gamma_2) - A_{(k)}(\gamma_1)A_{(k)}(\gamma_2)$. The functions $A_{(k)}, D_{(k)}, \Xi_{(k)}$ are continuous for every $k \in Z_{(k_0, i)}[r_0]$ thanks to continuity of the Deligne representation ρ_k .

For any $k \in Z_{(k_0, i)}[r_0]$, we denote by $P_k \subset \mathbf{A}_{\mathcal{K}}^{\circ}$ the kernel of the evaluation map $\mathbf{A}_{\mathcal{K}}^{\circ} \longrightarrow \mathcal{K}$ at k . Let us denote the function $A_{(k)} + D_{(k)}$ by Tr_{P_k} . Since $Z_{(k_0, i)}[r_0]$ is a countable set, we choose a numbering

$$Z_{(k_0, i)}[r_0] = \{k_1, k_2, \dots, k_s, \dots\}.$$

For $k_1, k_2 \in Z_{(k_0, i)}[r_0]$, consider the map

$$(4.4) \quad \mathbf{A}_{\mathcal{K}}^{\circ}/P_{k_1} \oplus \mathbf{A}_{\mathcal{K}}^{\circ}/P_{k_2} \longrightarrow \mathbf{A}_{\mathcal{K}}^{\circ}/(P_{k_1} + P_{k_2}), \quad (x, y) \mapsto (x \bmod P_{k_2}) - (y \bmod P_{k_1})$$

whose kernel is isomorphic to $\mathbf{A}_{\mathcal{K}}^{\circ}/(P_{k_1} \cap P_{k_2})$. Let ℓ be a prime number outside S . We have $\text{Tr}_{P_k}(\text{Frob}_{\ell}) = a_{\ell}(k)$ for every $k \in Z_{(k_0, i)}[r_0]$ and the Fourier coefficients $a_{\ell}(k_i)$ glue together when k_i varies. Hence the values $\text{Tr}_{P_k}(\text{Frob}_{\ell})$ glue together when k varies, which is true for any prime number ℓ outside S . By the sequence (4.4), we have a continuous function $\text{Tr}_{P_{k_1} \cap P_{k_2}}: G_{\mathbb{Q}, S} \longrightarrow \mathbf{A}_{\mathcal{K}}^{\circ}/(P_{k_1} \cap P_{k_2})$ whose value $\text{Tr}_{P_{k_1} \cap P_{k_2}}(\text{Frob}_{\ell})$ is congruent to $\text{Tr}_{P_{k_1}}(\text{Frob}_{\ell})$ (resp. $\text{Tr}_{P_{k_2}}(\text{Frob}_{\ell})$) mod P_{k_1} (resp. mod P_{k_2}) for every prime number ℓ outside S . Since the set of Frobenius elements $\{\text{Frob}_{\ell}\}_{\ell \notin S}$ is dense in $G_{\mathbb{Q}, S}$, it follows that the value $\text{Tr}_{P_{k_1} \cap P_{k_2}}(\sigma)$ is congruent to $\text{Tr}_{P_{k_1}}(\sigma)$ (resp. $\text{Tr}_{P_{k_2}}(\sigma)$) mod P_{k_1} (resp. mod P_{k_2}) for every $\sigma \in G_{\mathbb{Q}, S}$.

By an inductive argument, for each natural number s , we have a continuous function

$$\text{Tr}_{P_{k_1} \cap P_{k_2} \cap \dots \cap P_{k_s}}: G_{\mathbb{Q}, S} \longrightarrow \mathbf{A}_{\mathcal{K}}^{\circ}/(P_{k_1} \cap P_{k_2} \cap \dots \cap P_{k_s})$$

and reducing it (mod P_{k_i}) we recover the function $\text{Tr}_{P_{k_i}}$. Note that $\mathbf{A}_{\mathcal{K}}^{\circ} = \varprojlim_s \mathbf{A}_{\mathcal{K}}^{\circ}/(P_{k_1} \cap P_{k_2} \cap \dots \cap P_{k_s})$ since $\mathbf{A}_{\mathcal{K}}^{\circ}$ is complete and local. We can thus define the continuous function $\text{Tr}: G_{\mathbb{Q}, S} \longrightarrow \mathbf{A}_{\mathcal{K}}^{\circ}$ to be $\varprojlim_s \text{Tr}_{P_{k_1} \cap P_{k_2} \cap \dots \cap P_{k_s}}$.

We define the desired functions A and D as

$$A(\sigma) = \frac{\mathrm{Tr}(\sigma) - \mathrm{Tr}(c \cdot \sigma)}{2} \quad \text{and} \quad D(\sigma) = \frac{\mathrm{Tr}(\sigma) + \mathrm{Tr}(c \cdot \sigma)}{2}.$$

Since $\mathbf{A}_{\mathcal{K}}^{\circ}$ is complete and local, a similar argument yields the construction of Ξ , by first evaluating the value $b(g_1)c(g_2)$ at pairs $(g_1, g_2) = (\mathrm{Frob}_{\ell_1}, \mathrm{Frob}_{\ell_2})$ and then observing that the set of these pairs is dense in $G_{\mathbb{Q},S} \times G_{\mathbb{Q},S}$. We thus obtain a continuous function

$$\Xi: G_{\mathbb{Q},S} \times G_{\mathbb{Q},S} \longrightarrow \mathbf{A}_{\mathcal{K}}^{\circ}$$

that recovers the function $\Xi_{(k)}$ by taking the reduction modulo P_k of the function Ξ for every $k \in Z_{(k_0,i)}[r_0]$.

Setting $\pi := (A, D, \Xi)$, we need to check that the triple satisfies properties (II)–(IV) above, using the fact that $Z_{(k_0,i)}[r_0]$ is dense in the reduced \mathcal{X} . First, let us verify property (IV), namely that for each $g_1, g_2, h_1, h_2 \in G_{\mathbb{Q},S}$, the following holds:

$$(4.5) \quad \Xi(g_1, g_2)\Xi(h_1, h_2) - \Xi(g_1, h_2)\Xi(h_1, g_2) = 0.$$

We need to check that the function $\Xi(g_1, g_2)\Xi(h_1, h_2) - \Xi(g_1, h_2)\Xi(h_1, g_2)$ vanishes identically on \mathcal{X} . Since the aforementioned property (IV) for the pseudo-representation $\pi_k = (A_{(k)}, D_{(k)}, \Xi_k)$, we have

$$\begin{aligned} (\Xi(g_1, g_2)\Xi(h_1, h_2) - \Xi(g_1, h_2)\Xi(h_1, g_2))(k) \\ = \Xi_k(g_1, g_2)\Xi_k(h_1, h_2) - \Xi_k(g_1, h_2)\Xi_k(h_1, g_2) \end{aligned}$$

at each point k in the dense subset $Z_{(k_0,i)}[r_0]$. This proves the desired vanishing of (4.5) and the same argument shows the other properties (II) and (III). This completes the proof. \square

Theorem 4.3 below, which is the main result of this section, shows that the pseudo-representation that we have just constructed comes from a true representation.

Theorem 4.3. *Under the same assumptions and notations as in Theorem 3.2 and in Proposition 4.2, there exists a free $\mathbf{A}_{\mathcal{K}}^{\circ}$ -module \mathbf{T} of rank two with a continuous $G_{\mathbb{Q},S}$ -action such that the representation*

$$\rho: G_{\mathbb{Q},S} \longrightarrow \mathrm{Aut}_{\mathbf{A}_{\mathcal{K}}^{\circ}}(\mathbf{T})$$

satisfies $\pi_{\rho} = \pi$ (in a suitable basis). In particular, ρ modulo P_k is isomorphic to a lattice in the space V_k of ρ_k for all $k \in Z_{(k_0,i)}[r_0]$.

The following proof mainly relies on [Wil88, Lemma 2.2.3], see also [Hid93, Proposition 1, §7.5].

Proof. Start with a radius r_0 as in Theorem 3.2 and set $r = r_0|\varpi|$, where ϖ is a uniformizer of \mathcal{K} . Let $\pi = (A, D, \Xi)$ be the $\mathbf{A}_{\mathcal{K}}^{\circ}$ -valued pseudo-representation constructed in Proposition 4.2. There exists a pair of elements $\sigma, \tau \in G_{\mathbb{Q},S}$ such that $\Xi(\sigma, \tau)(k_0) \neq 0$. If not, the diagonal Galois representation $g \mapsto A(g)(k_0) \oplus D(g)(k_0)$ would have the same trace and determinant as the representation ρ_k , contradicting Ribet's result [Rib77, Theorem 2.3] stating that ρ_k be irreducible (see [Wil88, Lemma 2.2.3] or [Hid89, Proposition 1.1]). From now on, let us fix a pair σ, τ such that $\Xi(\sigma, \tau)(k_0) \in \mathcal{K}$ is of minimal valuation, say $\mu \in \mathbb{Q}$.

The element $\Xi(\sigma, \tau)$ can be decomposed as $\Xi(\sigma, \tau) = p^\mu V^{-1}$ with $V \in (\mathbf{A}_\mathcal{K}^\circ)^\times$. As in [Hid93, Proposition 1], we check that the map

$$\rho: g \mapsto \begin{pmatrix} A(g) & \Xi(g, \tau)Vp^{-\mu} \\ \Xi(\sigma, g) & D(g) \end{pmatrix}$$

is multiplicative, sends $\mathbf{1} \in G_{\mathbb{Q}, S}$ to $\mathbf{Id}_2 \in M_2(\mathbf{A}_\mathcal{K}^\circ)$ and takes values in $M_2(\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p)$. Hence we have a continuous group homomorphism $\rho: G_{\mathbb{Q}, S} \rightarrow \mathrm{GL}_2(\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p)$. In order to produce a finitely generated $\mathbf{A}_\mathcal{K}^\circ$ -submodule of $\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p$ which is ρ -stable, we follow the proof of continuity of [Hid89, Proposition 1.1]. Namely, define $\mathfrak{J} \subseteq \mathbf{A}_\mathcal{K}^\circ$ to be the ideal generated by all lower-left entries $\Xi(\sigma, g)$ for $g \in G_{\mathbb{Q}, S}$ and let \mathbf{T}' be the $\mathbf{A}_\mathcal{K}^\circ$ -submodule of $(\mathbf{A}_\mathcal{K}^\circ)^{\oplus 2} \subset (\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p)^{\oplus 2}$ generated by all vectors (x, y) with $x \in \mathbf{A}_\mathcal{K}^\circ, y \in \mathfrak{J}$. The $\mathbf{A}_\mathcal{K}^\circ$ -module \mathbf{T}' is finitely generated since $\mathbf{A}_\mathcal{K}^\circ$ is noetherian. Also, \mathbf{T}' is p -torsion free since \mathbf{T}' is contained in the module $(\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p)^{\oplus 2}$ which has no non-zero p -torsion element. Moreover, given ${}^t(x, y) \in \mathbf{T}'$ and $g \in G_{\mathbb{Q}, S}$, we have

$$\rho(g) \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} A(g)x + \Xi(g, \tau)Vp^{-\mu}y \\ \Xi(\sigma, g)x + D(g)y \end{pmatrix} \in \mathbf{T}'.$$

Indeed, the definition of \mathfrak{J} ensures $\Xi(\sigma, g)x \in \mathfrak{J}$, and the definition of μ guarantees that the element $\Xi(g, \tau)Vp^{-\mu}$ belongs to $\mathbf{A}_\mathcal{K}^\circ$. It follows that \mathbf{T}' is Galois stable. Finally, we have the equality $\mathbf{T}' \otimes \mathbb{Q}_p = (\mathbf{A}^\circ \otimes \mathbb{Q}_p)^{\oplus 2}$ because \mathfrak{J} contains the element $\Xi(\sigma, \tau) = p^{-\mu}V$ which becomes a unit after inverting p . Define \mathbf{T} to be the double dual $\mathrm{Hom}_{\mathbf{A}_\mathcal{K}^\circ}(\mathrm{Hom}_{\mathbf{A}_\mathcal{K}^\circ}(\mathbf{T}', \mathbf{A}_\mathcal{K}^\circ), \mathbf{A}_\mathcal{K}^\circ)$. We have a $G_{\mathbb{Q}, S}$ -equivariant $\mathbf{A}_\mathcal{K}^\circ$ -linear injection $\mathbf{T}' \hookrightarrow \mathbf{T}$ with finite cokernel. Since \mathbf{T} is a finitely generated reflexive module over a regular local ring $\mathbf{A}_\mathcal{K}^\circ$ of Krull dimension two, \mathbf{T} is a free $\mathbf{A}_\mathcal{K}^\circ$ -module of finite rank. Since $\mathbf{T} \otimes \mathbb{Q}_p$ is free of rank two over $\mathbf{A}_\mathcal{K}^\circ \otimes \mathbb{Q}_p$, the rank of \mathbf{T} over $\mathbf{A}_\mathcal{K}^\circ$ is two.

As for the interpolation property, recall that an irreducible representation of $G_{\mathbb{Q}, S}$ with values in a finite extension of \mathbb{Q}_p is uniquely determined by its trace and determinant. Thus $\mathbf{T}/P_k\mathbf{T}$ is isomorphic to a lattice inside V_k . \square

Note that $\iota: \mathbf{A}_\mathcal{K}^\circ \hookrightarrow \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ$ is continuous. In fact, $\mathbf{A}_\mathcal{K}^\circ$ is a local ring with maximal ideal $(\frac{T-k_0}{e_0}, \varpi)$ and $\mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ$ is endowed with the ϖ -adic topology, where ϖ denotes as before a uniformizer of \mathcal{K} . To show that ι is continuous, we need to show that $\iota^{-1}(\varpi \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ)$ contains the maximal ideal $(\frac{T-k_0}{e_0}, \varpi)$ of $\mathbf{A}_\mathcal{K}^\circ$. In fact, this implies that $\iota^{-1}(\varpi^n \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ)$ contains $(\frac{T-k_0}{e_0}, \varpi)^n$ for every n . The uniformizer ϖ is clearly contained in $\iota^{-1}(\varpi \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ)$ and $\frac{T-k_0}{e_0}$ is also contained in $\iota^{-1}(\varpi \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ)$ since we have $\frac{T-k_0}{e_0} = \varpi \frac{T-k_0}{e_0 \varpi} \in \varpi \mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ$. By extending the coefficients of the result obtained over $\mathbf{A}_\mathcal{K}^\circ$ in the Theorem 4.3 to $\mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ$, we obtain the following corollary:

Corollary 4.4. *Under the same assumptions and notations as in Theorem 4.3, there exists a free $\mathcal{A}_{\mathcal{X}(k_0, i)[r]/\mathcal{K}}^\circ$ -module \mathcal{T} of rank two with a continuous $G_{\mathbb{Q}, S}$ -action such that*

the representation

$$\rho: G_{\mathbb{Q},S} \longrightarrow \text{Aut}_{\mathcal{A}_{\mathcal{X}(k_0,i)}^{[r]}/\mathcal{K}}^{\circ}(\mathcal{T})$$

satisfies $\pi_{\rho} = \pi$ (in a suitable basis). In particular, ρ modulo \mathfrak{m}_k is isomorphic to ρ_k for all $k \in Z_{(k_0,i)}[r_0]$ where \mathfrak{m}_k is the unique maximal ideal of $\mathcal{A}_{\mathcal{X}(k_0,i)}^{[r]}/\mathcal{K}$ such that $\mathfrak{m}_k \cap \mathbf{A}_{\mathcal{K}}^{\circ} = P_k$.

5. THE FINE SELMER GROUP AND GALOIS DEFORMATIONS

We stick to the notation introduced in Theorem 3.2 and in Definition 3.5. Let F be a finite extension of \mathbb{Q} and fix $k \in Z_{(k_0,i)}[r_0]$ (see (4.3)). The fine Selmer group of f_k over F is defined by

$$\mathcal{R}(F, V_{f_k}/T_{f_k}) = \text{Ker} \left(H^1(\mathbb{Q}_S/F, V_{f_k}/T_{f_k}) \rightarrow \bigoplus_{v \in S} H^1(F_v, V_{f_k}/T_{f_k}) \right),$$

where T_{f_k} is the lattice of the representation V_{f_k} of ρ_k from Theorem 4.3. The fine Selmer over the extension $\mathbb{Q}(\mu_{p^\infty})$ is defined by

$$(5.1) \quad \mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), V_{f_k}/T_{f_k}) = \varinjlim \mathcal{R}(F, V_{f_k}/T_{f_k}),$$

where the limit is over all finite extensions of \mathbb{Q} contained in $\mathbb{Q}(\mu_{p^\infty})$. Set $(T_{f_k})^*(1) = \text{Hom}_{\text{cont}}(T_{f_k}, \mathcal{O}_{\mathcal{K}}) \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(1)$ and let $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), (T_{f_k})^*(1))$ be the Pontryagin dual of the group $\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), V_{f_k}/T_{f_k})$. Since p is odd,

$$\text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}) \cong \Delta \times (1 + p\mathbb{Z}_p)$$

where Δ is a finite group of order $p-1$. For a character η of Δ , by a result of Kato [Kat04, Theorem 12.4(1)], we know that the η -isotypic component $(\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), (T_{f_k})^*(1)))^\eta$ is torsion over the Iwasawa algebra $\mathbb{Z}_p[[1 + p\mathbb{Z}_p]]$. For simplicity we will restrict ourselves to the case $\mathbb{Q}(\mu_{p^\infty})$; one could have also worked with the dual fine Selmer group over the cyclotomic extension of any abelian number field where Kato's result is true.

Remark 5.1. For supersingular elliptic curves, the dual fine Selmer group is very close to the torsion subgroup of the whole classical dual Selmer group as suggested by the result [Win89, Corollary 2.5] of Wingberg, saying that given a supersingular elliptic curve E/F and assuming the finiteness of $\text{III}(E/F_n)_{p^\infty}$ for all n , the torsion part of the dual Selmer group of E over F_{cyc} is pseudo-isomorphic to the dual fine Selmer group of E over F_{cyc} . Therefore, philosophically, under appropriate hypotheses, the dual fine Selmer group is not far from the torsion part of the dual classical Selmer group. Under certain hypotheses, another evidence in this direction was also provided by Billot over the \mathbb{Z}_p^2 -extension $F_\infty = E[p^\infty]$ of F , when E/F is an elliptic curve with complex multiplication by an imaginary quadratic field K in F that has good supersingular reduction for primes above p (see [Bil86, Theorem 3.23]). In [Mat20, Theorem 1.1], Matar gave a Galois-theoretic proof of Wingberg's result with a slightly different hypothesis. Wingberg's result has been generalized by Lei and Lim to modular forms which are non-ordinary at p under appropriate hypotheses (see [LL22, Theorem 3.4]).

Let $\Lambda_{\text{cyc}} := \mathbb{Z}_p[[\text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q})]]$ be the Iwasawa algebra of the group $\text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q})$. Let $\Lambda_{\text{cyc}}(\tilde{\chi})$ be a free Λ_{cyc} -module of rank one on which the absolute Galois group $G_{\mathbb{Q}}$ of \mathbb{Q} acts via the cyclotomic character

$$\tilde{\chi}: G_{\mathbb{Q}} \rightarrow \text{Gal}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}) \hookrightarrow \Lambda_{\text{cyc}}^{\times}.$$

The following definition is adapted from Greenberg (see [Gre06] and [Gre94]).

Definition 5.2. Write $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ} := \mathbf{A}_{\mathcal{K}}^{\circ} \widehat{\otimes}_{\mathbb{Z}_p} \Lambda_{\text{cyc}}$, and let \mathbf{T} be as in Theorem 4.3. The cyclotomic deformation of \mathbf{T} is defined as

$$\tilde{\mathbf{T}} := \mathbf{T} \widehat{\otimes}_{\mathbb{Z}_p} \Lambda_{\text{cyc}}(\tilde{\chi})$$

regarded as a module over $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$. Set $\tilde{\mathbb{D}} := \tilde{\mathbf{T}} \otimes_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}} \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$ where $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ} = \text{Hom}_{\mathcal{O}_{\mathcal{K}}}(\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}, \mathcal{K}/\mathcal{O}_{\mathcal{K}})$ is the Pontryagin dual of $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$. We endow $\tilde{\mathbb{D}}$ with the discrete topology.

Finally, set $(\tilde{\mathbf{T}})^{\star} = \text{Hom}_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}}(\tilde{\mathbf{T}}, \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ})$ to be the $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$ -linear dual of $\tilde{\mathbf{T}}$, and let $\mathbf{T}^{\star} = \text{Hom}_{\mathbf{A}_{\mathcal{K}}^{\circ}}(\mathbf{T}, \mathbf{A}_{\mathcal{K}}^{\circ})$ be the $\mathbf{A}_{\mathcal{K}}^{\circ}$ -linear dual of \mathbf{T} .

Note that

$$\begin{aligned} \text{Hom}_{\mathcal{O}_{\mathcal{K}}}(\tilde{\mathbb{D}}, \mathcal{K}/\mathcal{O}_{\mathcal{K}}(1)) &\cong \text{Hom}_{\mathcal{O}_{\mathcal{K}}}(\tilde{\mathbf{T}} \otimes_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}} \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}, \mathcal{K}/\mathcal{O}_{\mathcal{K}}(1)) \\ &\cong \text{Hom}_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}}(\tilde{\mathbf{T}}, \text{Hom}_{\mathcal{O}_{\mathcal{K}}}(\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}, \mathcal{K}/\mathcal{O}_{\mathcal{K}})) \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(1) \\ &\cong \text{Hom}_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}}(\tilde{\mathbf{T}}, \widehat{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}})(1) \\ &\cong \text{Hom}_{\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}}(\tilde{\mathbf{T}}, \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ})(1) = (\tilde{\mathbf{T}})^{\star}(1). \end{aligned}$$

Similarly, one can define

$$\mathbb{D} = \mathbf{T} \otimes_{\mathbf{A}_{\mathcal{K}}^{\circ}} \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$$

where $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ}$ is the Pontryagin dual of $\mathbf{A}_{\mathcal{K}}^{\circ}$.

Suppose $\phi: \mathbf{A}_{\mathcal{K}}^{\circ} \rightarrow \mathcal{O}_{\mathcal{K}}$ is any continuous surjection, set $P_{\phi} = \text{Ker}(\phi)$ and $T_{\phi} = \mathbf{T}/P_{\phi}\mathbf{T}$. Define

$$\mathbb{D}_{\phi} := \mathbf{T}_{\phi} \otimes_{\mathcal{O}_{\mathcal{K}}} \widehat{\mathcal{O}}_{\mathcal{K}}.$$

By Greenberg [Gre06, p. 336], we have $\mathbb{D}[P_{\phi}] \cong \mathbb{D}_{\phi}$, where $\mathbb{D}[P_{\phi}]$ denotes the submodule of \mathbb{D} annihilated by P_{ϕ} .

Similarly, by *loc. cit.*, if $\tilde{\phi}: \widehat{\mathbf{A}}_{\mathcal{K}}^{\circ} \rightarrow \mathcal{O}_{\mathcal{K}}$ is a continuous surjection and $P_{\tilde{\phi}}$ denotes $\text{Ker}(\tilde{\phi})$, let $\mathbf{T}_{\tilde{\phi}} := \tilde{\mathbf{T}}/P_{\tilde{\phi}}\tilde{\mathbf{T}}$, which is a free $\mathcal{O}_{\mathcal{K}}$ -module of rank two. Then,

$$\tilde{\mathbb{D}}[P_{\tilde{\phi}}] \cong \tilde{\mathbb{D}}_{\tilde{\phi}} := \mathbf{T}_{\tilde{\phi}} \otimes_{\mathcal{O}_{\mathcal{K}}} \widehat{\mathcal{O}}_{\mathcal{K}}.$$

Following [Gre06, p. 336], we define the fine Selmer group of $\tilde{\mathbb{D}}$ as

$$\mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}}) = \text{Ker} \left(H^1(\mathbb{Q}_S/\mathbb{Q}, \tilde{\mathbb{D}}) \rightarrow \bigoplus_{v \in S} H^1(\mathbb{Q}_v, \tilde{\mathbb{D}}) \right).$$

Finally, write $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^{\star}(1))$ for the Pontryagin dual $\text{Hom}_{\mathcal{O}_{\mathcal{K}}}(\mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}}), \mathcal{K}/\mathcal{O}_{\mathcal{K}})$ of $\mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}})$.

6. SPECIALIZATION AT ARITHMETIC PRIMES

We continue to work in the setting introduced in §§3–4, and we fix a weight $k \in Z_{(k_0, i)}[r_0]$. Recall from §4 that P_k is a height-one prime ideal of \mathbf{A}_K° . Let $J_k = P_k \widehat{\mathbf{A}}_K^\circ$ be the corresponding height-one prime ideal of $\widehat{\mathbf{A}}_K^\circ$. We have a map

$$(6.1) \quad \frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))} \xrightarrow{s_k} \mathcal{Y}\left(\mathbb{Q}, \frac{(\tilde{\mathbf{T}})^*(1)}{J_k (\tilde{\mathbf{T}})^*(1)}\right)$$

which arises from taking Pontryagin dual of the map

$$(6.2) \quad \mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}}[J_k]) \xrightarrow{\widehat{s}_k} \mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}})[J_k].$$

Given any $v \in S$, let $G_{\infty, v}$ be the Galois group $\text{Gal}(\overline{\mathbb{Q}}_v / \mathbb{Q}_{\infty, v})$ where $\mathbb{Q}_{\infty, v} = \mathbb{Q}(\mu_{p^\infty})_v$.

Lemma 6.1. *For all $v \in S$, we have*

$$H_0(G_{\mathbb{Q}_v}, (\tilde{\mathbf{T}})^*) \cong H_0(G_{\infty, v}, \mathbf{T}^*)^\iota$$

as \mathbf{A}_K° -modules, where the exponent ι denotes that the Galois group acts by the involution $g \mapsto g^{-1}$.

Proof. The Pontryagin duals of $(\tilde{\mathbf{T}})^*$ and \mathbf{T}^* are $\tilde{\mathbb{D}}$ and \mathbb{D} , respectively. So the Pontryagin duals of $(\tilde{\mathbf{T}}^*)_{G_{\mathbb{Q}_v}}$ and $(\mathbf{T}^*)_{G_{\infty, v}}^\iota$ are $\tilde{\mathbb{D}}^{G_{\mathbb{Q}_v}}$ and $(\mathbb{D}^{G_{\infty, v}})^\iota$. Therefore, we have to show that

$$(6.3) \quad \tilde{\mathbb{D}}^{G_{\mathbb{Q}_v}} \cong (\mathbb{D}^{G_{\infty, v}})^\iota.$$

For this we will follow the line of arguments presented in [Gre94, p. 214-215] (see also [Gre06, Remark 5.9.2]). Note that

$$\text{Hom}_{\mathcal{O}_K}(\widehat{\mathbf{A}}_K^\circ, \mathcal{K}/\mathcal{O}_K) = \text{Hom}_{\mathcal{O}_K}(\mathbf{A}_K^\circ \widehat{\otimes}_{\mathbb{Z}_p} \Lambda_{\text{cyc}}, \mathcal{K}/\mathcal{O}_K).$$

Since we are considering continuous maps and the topology on $\mathcal{K}/\mathcal{O}_K$ is discrete, we have

$$\text{Hom}_{\mathcal{O}_K}(\mathbf{A}_K^\circ \widehat{\otimes}_{\mathbb{Z}_p} \Lambda_{\text{cyc}}, \mathcal{K}/\mathcal{O}_K) = \text{Hom}_{\mathcal{O}_K}(\mathbf{A}_K^\circ \otimes_{\mathbb{Z}_p} \Lambda_{\text{cyc}}, \mathcal{K}/\mathcal{O}_K).$$

Since the Hom functor and the tensor product form an adjoint pair, we deduce

$$\text{Hom}_{\mathcal{O}_K}(\mathbf{A}_K^\circ \otimes_{\mathbb{Z}_p} \Lambda_{\text{cyc}}, \mathcal{K}/\mathcal{O}_K) = \text{Hom}_{\mathbb{Z}_p}(\Lambda_{\text{cyc}}, \widehat{\mathbf{A}}_K^\circ).$$

Therefore

$$\tilde{\mathbb{D}} = \left(\mathbf{T} \otimes_{\mathbb{Z}_p} \Lambda_{\text{cyc}}(\tilde{\chi}) \right) \otimes_{\widehat{\mathbf{A}}_K^\circ} \text{Hom}_{\mathbb{Z}_p}(\Lambda_{\text{cyc}}, \widehat{\mathbf{A}}_K^\circ).$$

Now $\tilde{\mathbf{T}} = \mathbf{T} \otimes_{\mathbb{Z}_p} \Lambda_{\text{cyc}}(\tilde{\chi})$ is a free module of rank 2 over $\widehat{\mathbf{A}}_K^\circ$ and hence

$$\begin{aligned} \tilde{\mathbb{D}} &\cong \text{Hom}_{\mathbb{Z}_p}(\Lambda_{\text{cyc}}, \mathbf{T} \otimes_{\mathbf{A}_K^\circ} \widehat{\mathbf{A}}_K^\circ)(\tilde{\chi}) \\ &\cong \text{Hom}_{\mathbb{Z}_p}(\Lambda_{\text{cyc}}(\tilde{\chi}^{-1}), \mathbb{D}). \end{aligned}$$

We deduce that

$$\tilde{\mathbb{D}}^{G_{\infty, v}} \cong \text{Hom}_{G_{\infty, v}}(\Lambda_{\text{cyc}}(\tilde{\chi}^{-1}), \mathbb{D}) \cong \text{Hom}_{\mathbb{Z}_p}(\Lambda_{\text{cyc}}(\tilde{\chi}^{-1}), \mathbb{D}^{G_{\infty, v}}).$$

Writing Γ_v for the local cyclotomic Galois group $\text{Gal}(\mathbb{Q}_{\infty,v}/\mathbb{Q}_v) = G_{\mathbb{Q}_v}/G_{\infty,v}$, this implies that

$$\begin{aligned}\tilde{\mathbb{D}}^{G_{\mathbb{Q}_v}} &\cong \text{Hom}_{\Gamma_v}(\Lambda_{\text{cyc}}(\tilde{\chi}^{-1}), \mathbb{D}^{G_{\infty,v}}), \\ &\cong \text{Hom}_{\Gamma_v}(\Lambda_{\text{cyc}}(\tilde{\chi}), (\mathbb{D}^{G_{\infty,v}})^\iota).\end{aligned}$$

The argument that follows is identical to that in [Gre94, pp. 214–215]. One can identify $\text{Hom}_{\Gamma_v}(\Lambda_{\text{cyc}}(\tilde{\chi}), (\mathbb{D}^{G_{\infty,v}})^\iota)$ with $\text{Hom}_{\Lambda_{\text{cyc}}}(\Lambda_{\text{cyc}}, (\mathbb{D}^{G_{\infty,v}})^\iota)$ considering $\Lambda_{\text{cyc}}(\tilde{\chi})$ and $(\mathbb{D}^{G_{\infty,v}})^\iota$ as Λ_{cyc} -modules. But any element $f \in \text{Hom}_{\Lambda_{\text{cyc}}}(\Lambda_{\text{cyc}}, (\mathbb{D}^{G_{\infty,v}})^\iota)$ is determined completely by $f(1)$ and hence there is an isomorphism $\text{Hom}_{\Lambda_{\text{cyc}}}(\Lambda_{\text{cyc}}, (\mathbb{D}^{G_{\infty,v}})^\iota) \cong (\mathbb{D}^{G_{\infty,v}})^\iota$ as Λ_{cyc} -modules. This completes the proof of (6.3). \square

Let M be the maximal ideal of $\mathbf{A}_{\mathcal{K}}^\circ$, and set $\kappa = \mathbf{A}_{\mathcal{K}}^\circ/M$. For the rest of the article, we will need the following hypothesis.

Hypothesis 6.2. *The residual κ -representation ρ/M is irreducible when restricted to $G_{\mathbb{Q}_p}$, where ρ is the Galois representation constructed in Theorem 4.3.*

Theorem 6.3. *Under Hypothesis 6.2, the map s_k in (6.1) is surjective and $\text{Ker}(s_k)$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all $k \in Z_{(k_0,i)}[r_0]$. Furthermore, $\text{Ker}(s_k)$ is finite for all but finitely many k .*

Proof. We have the following commutative diagram with exact rows.

$$(6.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}}[J_k]) & \longrightarrow & H^1(\mathbb{Q}_S/\mathbb{Q}, \tilde{\mathbb{D}}[J_k]) & \longrightarrow & \bigoplus_{v \in S} H^1(\mathbb{Q}_v, \tilde{\mathbb{D}}[J_k]) \\ & & \downarrow \widehat{s_k} & & \downarrow g_k & & \downarrow h_k \\ 0 & \longrightarrow & \mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}})[J_k] & \longrightarrow & H^1(\mathbb{Q}_S/\mathbb{Q}, \tilde{\mathbb{D}})[J_k] & \longrightarrow & \bigoplus_{v \in S} H^1(\mathbb{Q}_v, \tilde{\mathbb{D}})[J_k] \end{array}$$

Observe that $\mathbb{D}[M]$ (the submodule of \mathbb{D} annihilated by M) is a finite dimensional κ -representation and, by Hypothesis 6.2,

$$H^0(\mathbb{Q}_S/\mathbb{Q}, \mathbb{D}[M]) = H^0(\mathbb{Q}_S/\mathbb{Q}, \mathbb{D})[M] = 0.$$

Note that $\Lambda_{\text{cyc}}(\tilde{\chi})$ is isomorphic to $p - 1$ copies of the Iwasawa algebra in one variable over \mathbb{Z}_p . Hence, the residual κ -representation $\tilde{\mathbf{T}}/M\tilde{\mathbf{T}}$ associated to $\tilde{\mathbf{T}}$ is isomorphic to a direct sum of $p - 1$ copies of the irreducible residual representation of $\mathbf{T}/M\mathbf{T}$. Therefore, any κ -irreducible subquotient of the residual representation of $\tilde{\mathbf{T}}/M\tilde{\mathbf{T}}$ is isomorphic to the irreducible residual representation $\mathbf{T}/M\mathbf{T}$. Applying [Gre06, Proposition 3.4], we deduce that the map g_k is an isomorphism and hence $\widehat{s_k}$ is injective (that is, s_k is surjective). In the following, we will analyse $\widehat{\text{Ker}(h_k)}$ and show that $\widehat{\text{Ker}(h_k)}$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all k , and is finite for all but finitely many k . By the snake lemma, the same conclusion will hold for $\widehat{\text{Coker}(\widehat{s_k})} \cong \widehat{\text{Ker}(s_k)}$. We remark that the Pontryagin dual of $\text{Ker}(h_k)$ is isomorphic to $\bigoplus_{v \in S} (\tilde{\mathbf{T}}^*)_{G_{\mathbb{Q}_v}}[J_k]$.

By Lemma 6.1, we obtain

$$\bigoplus_{v \in S} (\tilde{\mathbf{T}}^*)_{G_{\mathbb{Q}_v}}[J_k] = \bigoplus_{v \in S} (\mathbf{T}^*)_{G_{\infty, v}}^{\iota}[P_k].$$

If P_k does not divide the characteristic ideal of the torsion subgroup of $\bigoplus_{v \in S} (\mathbf{T}^*)_{G_{\infty, v}}^{\iota}$ as an $\mathbf{A}_{\mathcal{K}}^{\circ}$ -module, then we obtain that $\bigoplus_{v \in S} (\mathbf{T}^*)_{G_{\infty, v}}^{\iota}[P_k]$ is finite (see [Och05, Lemma 4.1] for a generalization of this to modules over Iwasawa algebras of several variables). Note also that, for all k , $\bigoplus_{v \in S} (\mathbf{T}^*)_{G_{\infty, v}}^{\iota}[P_k]$ is a finitely generated $\mathbf{A}_{\mathcal{K}}^{\circ}/P_k$ -module, and thus finitely generated over $\mathcal{O}_{\mathcal{K}}$ because $\mathbf{A}_{\mathcal{K}}^{\circ}/P_k$ is finite over $\mathcal{O}_{\mathcal{K}}$. This shows that $\widehat{\text{Coker}}(\widehat{s_k})$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all k , and it is finite for all but finitely many k .

Let us denote

$$\tilde{T}_{P_k} := (\mathbf{T}/P_k\mathbf{T}) \otimes_{\mathbb{Z}_p} \Lambda_{\text{cyc}}(\tilde{\chi})$$

which is the cyclotomic deformation of $\mathbf{T}/P_k\mathbf{T} \cong T_{f_k}$. By [Gre06, p. 357],

$$\tilde{\mathbb{D}}[J_k] \cong \tilde{T}_{P_k} \otimes_{\Lambda_{\text{cyc}}} \widehat{\Lambda}_{\text{cyc}}.$$

It follows from classical results of Greenberg (see, for example, [Gre06, (2), p. 342]) that

$$(6.5) \quad \mathcal{R}(\mathbb{Q}, \tilde{T}_{P_k} \otimes_{\Lambda_{\text{cyc}}} \widehat{\Lambda}_{\text{cyc}}) \cong \mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), V_{f_k}/T_{f_k})^{\iota},$$

where $\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), V_{f_k}/T_{f_k})$ is the fine Selmer group defined in (5.1). Note that Greenberg's result is over the cyclotomic extension. But our Λ_{cyc} is isomorphic to $\mathbb{Z}_p[[\mathbb{Z}_p^{\times}]]$ and hence we recover the fine Selmer group over $\mathbb{Q}(\mu_{p^\infty})$. See also [JS11, Proposition 3] where Jha and Sujatha prove (6.5) in the ordinary case. The proof remains essentially the same in our case. Therefore,

$$\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), V_{f_k}/T_{f_k})^{\iota} \xrightarrow{\widehat{s_k}} \mathcal{R}(\mathbb{Q}, \tilde{\mathbb{D}})[J_k]$$

is injective, $\widehat{\text{Coker}}(\widehat{s_k})$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all k , and is finite for all but finitely many k . Dually, the map

$$(6.6) \quad \frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))} \xrightarrow{s_k} \mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^{\iota}$$

is surjective and $\text{Ker}(s_k)$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all k . Furthermore, $\text{Ker}(s_k)$ is finite for all but finitely many k . This completes the proof of Theorem 6.3. \square

Recall $\mathbb{Z}_p^{\times} \cong \Delta \times (1 + p\mathbb{Z}_p)$ where Δ is a finite group of order $p - 1$. For a Dirichlet character $\eta: \Delta \rightarrow \mathbb{Z}_p^{\times}$, let e_{η} be the idempotent attached to η . Given a $\mathbb{Z}_p[[\mathbb{Z}_p^{\times}]]$ -module M define its η -isotypic component to be

$$M^{\eta} = e_{\eta} \cdot M,$$

considered as a $\mathbb{Z}_p[[1 + p\mathbb{Z}_p]]$ -module. As in [LP20, p. 3], we say that M has rank r as a $\mathbb{Z}_p[[\mathbb{Z}_p^{\times}]]$ -module if M^{η} has rank r as a $\mathbb{Z}_p[[1 + p\mathbb{Z}_p]]$ -module for all the characters η of Δ . Hence, M is torsion as a $\mathbb{Z}_p[[\mathbb{Z}_p^{\times}]]$ -module if M^{η} is torsion as a $\mathbb{Z}_p[[1 + p\mathbb{Z}_p]]$ -module for every character η .

Proposition 6.4. *For every character η of Δ , the component $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^{\eta}$ of the dual fine Selmer group is $\widehat{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -torsion.*

Proof. Fix an η as in the statement. By (6.6), choose k such that the map

$$\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))} \xrightarrow{s_k} \mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\iota$$

is surjective and the kernel of s_k is finite. Using the Poitou–Tate exact sequence (see for example [PR95, §A.3.1]) and a deep result of Kato [Kat04, Theorem 12.4], we know that $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))$ is torsion as a Λ_{cyc} -module. Now suppose $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ is not $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -torsion. Then by the structure theorem $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ can be decomposed into an $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -free part of rank at least 1 and an $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -torsion part. Recall that $J_k = P_k \widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ}$ is a height one prime ideal of $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ}$, where P_k is a height one prime ideal of $\mathbf{A}_{\mathcal{K}}^{\circ}$. Hence the $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -free part of $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ modulo J_k will contain a free $\Lambda_{\text{cyc}}^\eta$ module of rank at least 1. But this is impossible as the kernel of s_k is finite and $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is $\Lambda_{\text{cyc}}^\eta$ -torsion. \square

7. VARIATIONS OF IWASAWA μ AND λ -INVARIANTS

As before, in this section, we work under Hypothesis 6.2, and in the setting of §§3–4. From Proposition 6.4 and from the discussion before Theorem B, we know that for every character η of Δ , the component $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ of the fine Selmer group is $\widetilde{\mathbf{A}}_{\mathcal{K}}^{\circ \eta}$ -torsion and the specialization $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is also $\Lambda_{\text{cyc}}^\eta$ -torsion for all points $k \in Z_{(k_0, i)}[r_0]$.

Theorem 7.1. *For every character η of Δ , the following are equivalent:*

- (1) *The fine Selmer group $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ is a finitely generated $\mathbf{A}_{\mathcal{K}}^{\circ}$ -module.*
- (2) *The classical dual fine Selmer group $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for all points $k \in Z_{(k_0, i)}[r_0]$. That is, the μ -invariants of $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is zero for all points $k \in Z_{(k_0, i)}[r_0]$.*
- (3) *Part (2) above holds for some k . That is, there exists k such that the dual fine Selmer group $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is a finitely generated $\mathcal{O}_{\mathcal{K}}$ -module for that k (equivalently, its μ -invariant is zero).*

Proof. Fix a character η as in the statement.

We first show that (2) and (3) are equivalent. Fix a weight $k \in Z_{(k_0, i)}[r_0]$ and let $A_{f_k} = V_{f_k}/T_{f_k}$. As before, let ϖ be a uniformizer of $\mathcal{O}_{\mathcal{K}}$. We have the following Kummer sequence associated to multiplication by ϖ :

$$0 \rightarrow A_{f_k}[\varpi] \rightarrow A_{f_k} \xrightarrow{\varpi} A_{f_k} \rightarrow 0.$$

This gives us the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \left(\frac{H^0(\mathbb{Q}_S/\mathbb{Q}(\mu_{p^\infty}), A_{f_k})}{\varpi} \right)^\eta & \longrightarrow & H^1(\mathbb{Q}_S/\mathbb{Q}(\mu_{p^\infty}), A_{f_k}[\varpi])^\eta & \longrightarrow & \left(H^1(\mathbb{Q}_S/\mathbb{Q}(\mu_{p^\infty}), A_{f_k})[\varpi] \right)^\eta \longrightarrow 0 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ 0 & \longrightarrow & \left(\frac{\oplus_{v \in S} H^0(\mathbb{Q}(\mu_{p^\infty})_v, A_{f_k})}{\varpi} \right)^\eta & \longrightarrow & \oplus_{v \in S} H^1(\mathbb{Q}(\mu_{p^\infty})_v, A_{f_k}[\varpi])^\eta & \longrightarrow & \left(\oplus_{v \in S} H^1(\mathbb{Q}(\mu_{p^\infty})_v, A_{f_k})[\varpi] \right)^\eta \longrightarrow 0 \end{array}$$

Each of the first terms in the two horizontal exact sequences is finite. By definition, the kernel of β is $\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), A_{f_k}[\varpi])^\eta$ which is the η -component of the fine Selmer group corresponding to the module $A_{f_k}[\varpi]$ over the extension $\mathbb{Q}(\mu_{p^\infty})$. Clearly, the kernel of γ is $(\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), A_{f_k}[\varpi]))^\eta$. Therefore, by the snake lemma $(\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), A_{f_k}[\varpi]))^\eta$ is finite if and only if $\mathcal{R}(\mathbb{Q}(\mu_{p^\infty}), A_{f_k}[\varpi])^\eta$ is finite; but the latter only depends on the residual representation and all forms in the Coleman family have the same residual representation.

Now we show that (1) implies (3). By Theorem 6.3 and (6.6), there exists a point k such that

$$\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))} \xrightarrow{s_k} \mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^\star(1))^\iota$$

has finite kernel and is surjective. Taking η -isotypic components, (1) implies that the quotient $\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta}$ is finitely generated over $\mathbf{A}_K^\circ/P_k \cong \mathcal{O}_K$, and hence so is the component $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^\star(1))^\eta$. Next we show that (3) implies (1). By Nakayama's lemma, it suffices to show that $\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta}$ is a finitely generated \mathcal{O}_K -module. But $\text{Ker}(s_k)$ is a quotient of $\widehat{\text{Ker}(h_k)}$ (where h_k is as in diagram (6.4)) which is a finitely generated \mathcal{O}_K -module by the proof of Theorem 6.3. \square

Remark 7.2. Recall that Conjecture A of Coates and Sujatha deals with the case of elliptic curves defined over a number field F . It says that the μ -invariant of the dual fine Selmer group over the cyclotomic \mathbb{Z}_p -extension of F is trivial (see [CS05, p. 822]). This is equivalent to saying that this dual fine Selmer group is a finitely generated \mathbb{Z}_p -module. Hence the fact that $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^\star(1))^\eta$ is a finitely generated \mathcal{O}_K -module (namely, part (3) of Theorem 7.1) is essentially Conjecture A generalized for modular forms. Explicit examples of elliptic curves and modular forms with good *ordinary* reduction at the prime p satisfying Conjecture A are given in [Ari14]. For another approach to Coates and Sujatha's Conjecture A and variations of Iwasawa invariants beyond the ordinary case, see [NS21]; see [KNS22] for its relations with Greenberg's Generalized Conjecture.

Theorem 7.3. *Assume any of the equivalent conditions in Theorem 7.1. Then the λ -invariants of $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^\star(1))^\eta$ are equal for all but finitely many points $k \in Z_{(k_0, i)}[r_0]$.*

Proof. By Proposition 6.4, $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))$ is torsion as an $\widetilde{\mathbf{A}}_K^\circ$ -module but it may not be torsion over \mathbf{A}_K° . Write H for the characteristic ideal of the \mathbf{A}_K° -torsion submodule of $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta$, and factor it as $H = \prod_{1 \leq i \leq d} I_i^{m_i}$ where the I_i are height one prime ideal of $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^\star(1))^\eta$, and $m_i \geq 1$. Let

$$\Sigma = \{I_1, \dots, I_d\} \cup \{P_k \text{ such that } \text{Ker}(s_k) \text{ is infinite}\}.$$

Let k be such that P_k is not in Σ . The λ -invariant of $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is the \mathbb{Z}_p -rank of $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$. By (6.6) and by the choice of k , we have a surjective map

$$\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta} \xrightarrow{s_k} \mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^{\iota, \eta}$$

whose kernel is finite. Therefore, the λ -invariant of $\mathcal{Y}(\mathbb{Q}(\mu_{p^\infty}), T_{f_k}^*(1))^\eta$ is the \mathbb{Z}_p -rank of $\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}$. We will show that the \mathbb{Z}_p -rank of $\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}$ is equal to the $\mathbf{A}_{\mathcal{K}}^\circ$ -rank of the $(\mathbf{A}_{\mathcal{K}}^\circ)^\eta$ -module $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ and that the latter is independent of k . The theorem will then follow.

Consider $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ as a module over $\mathbf{A}_{\mathcal{K}}^\circ$. By the structure theorem of modules over Iwasawa algebras, we can decompose $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$ into an $\mathbf{A}_{\mathcal{K}}^\circ$ -free part of a certain rank r and an $\mathbf{A}_{\mathcal{K}}^\circ$ -torsion part of the form $\bigoplus_{i=1}^m \mathbf{A}_{\mathcal{K}}^\circ / I_i^{m_i}$. Recall that P_k is a height one prime ideal of $\mathbf{A}_{\mathcal{K}}^\circ$; it is the kernel of the evaluation map $\mathbf{A}_{\mathcal{K}}^\circ \rightarrow \mathcal{O}_{\mathcal{K}}$ at k . Also, note that we have chosen k such that P_k is not in Σ . Therefore, P_k is relatively prime to $I_i^{m_i}$ and hence $\mathbf{A}_{\mathcal{K}}^\circ / (I_i^{m_i}, P_k)$ is finite and hence has $\mathcal{O}_{\mathcal{K}}$ -rank equal to 0. Therefore, the $\mathcal{O}_{\mathcal{K}}$ -rank of $\frac{\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}{J_k \mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta}$ is r , which is the $\mathbf{A}_{\mathcal{K}}^\circ$ -rank of $\mathcal{Y}(\mathbb{Q}, (\tilde{\mathbf{T}})^*(1))^\eta$. \square

8. EXAMPLES

In this section, we sketch how to find examples satisfying the conditions of Theorems 7.1 and 7.3.

If there is any weight $k \in Z_{(k_0, i)}[r_0]$ such that the residual representation of ρ_{f_k} restricted to $G_{\mathbb{Q}_p}$ of the specialization f_k of the Coleman family at the weight k is irreducible, then the residual representation attached to the whole family ρ restricted to $G_{\mathbb{Q}_p}$ is irreducible and hence Hypothesis 6.2 is satisfied. Now, [Edi92, Theorem 2.6] says that the residual representation of ρ_{f_k} restricted to $G_{\mathbb{Q}_p}$ is irreducible if f_k is a cuspidal eigenform whose p -th Fourier coefficient a_p is 0 in $\overline{\mathbb{F}}_p$ and $2 \leq k < p + 1$. Hence there are plenty of examples satisfying Hypothesis 6.2. An explicit example is the elliptic curve 17A (labelled as in [Pol]); there are actually 4 curves in this isogeny class 17A, take any one of them). This curve has supersingular reduction at the prime $p = 11$, and the associated modular form is a weight 2 cuspidal modular form (see [LMF13]).

The next important hypothesis that we have made in Theorem 7.3 is a generalization of Conjecture A (see Remark 7.2). In the following, we will explain how to find examples when Conjecture A is true for elliptic modular forms at supersingular primes. Suppose p is a prime of good supersingular reduction for an elliptic curve E over \mathbb{Q} . Recall from [Kob03, Theorem 7.3 (i)] that the dual fine Selmer group of E over the cyclotomic extension of \mathbb{Q} is a quotient of the dual plus/minus signed Selmer groups which are torsion modules over the cyclotomic Iwasawa algebra. Recall also that the μ -invariant is additive along a short exact sequence of finitely generated and torsion Iwasawa modules. Hence if the dual signed Selmer groups over the cyclotomic extension of \mathbb{Q} have μ -invariant zero, then it is so for the dual

fine Selmer group as well; hence Conjecture A is valid in those examples. Pollack in [Pol] has already computed Iwasawa invariants of analytic plus/minus signed p -adic L -functions, and via the signed Iwasawa main conjectures, they are equal to the Iwasawa invariants of the dual signed Selmer groups. Recall that for CM elliptic curves over \mathbb{Q} and for primes $p > 3$ of supersingular reduction, the signed Iwasawa main conjectures have been proved by Pollack and Rubin in [PR04]. The signed Iwasawa main conjectures for supersingular elliptic curves over \mathbb{Q} have been completely settled now for odd primes (see [Wan21] for the case $a_p = 0$ and [Spr16] for the general case). So one can see from Pollack's table [Pol] that the dual signed Selmer groups for the curve 17A and prime $p = 11$ has trivial μ -invariants.

REFERENCES

- [AIS15] Fabrizio Andreatta, Adrian Iovita, and Glenn Stevens, *Overconvergent Eichler-Shimura isomorphisms*, J. Inst. Math. Jussieu **14** (2015), no. 2, 221–274.
- [Ari14] Chandrakant S. Aribam, *On the μ -invariant of fine Selmer groups*, J. Number Theory **135** (2014), 284–300.
- [Bel12] Joël Bellaïche, *Critical p -adic L -functions*, Invent. Math. **189** (2012), no. 1, 1–60. MR 2929082
- [BGR84] Sigfried Bosch, Ulrich Güntzer, and Reinhold Remmert, *Non-Archimedean analysis*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 261, Springer-Verlag, Berlin, 1984, A systematic approach to rigid analytic geometry.
- [Bil86] Patrick Billot, *Quelques aspects de la descente sur une courbe elliptique dans le cas de réduction supersingulière*, Compositio Math. **58** (1986), no. 3, 341–369.
- [BL21] Kâzım Büyükboduk and Antonio Lei, *Interpolation of generalized Heegner cycles in Coleman families*, Journal of the London Mathematical Society **104** (2021), no. 4, 1682–1716.
- [Buz07] Kevin Buzzard, *Eigenvarieties, L -functions and Galois representations*, London Math. Soc. Lecture Note Ser., vol. 320, Cambridge Univ. Press, Cambridge, 2007, pp. 59–120.
- [Che11] Gaëtan Chenevier, *The p -adic analytic space of pseudocharacters of a profinite group, and pseudorepresentations over arbitrary rings*, Automorphic Forms and Galois Representations (London Mathematical Society, ed.), EPSRC Durham Symposium, London Mathematical Society, 2011, pp. 1–56.
- [CM98] Robert F. Coleman and Barry Mazur, *The eigencurve*, Galois representations in arithmetic algebraic geometry (Durham, 1996), London Math. Soc. Lecture Note Ser., vol. 254, Cambridge Univ. Press, Cambridge, 1998, pp. 1–113.
- [Col97a] Robert F. Coleman, *On the coefficients of the characteristic series of the U -operator*, Proc. Nat. Acad. Sci. U.S.A. **94** (1997), no. 21, 11129–11132.
- [Col97b] ———, *p -adic Banach spaces and families of modular forms*, Invent. Math. **127** (1997), no. 3, 417–479.
- [CS05] John Coates and Ramdorai Sujatha, *Fine Selmer groups of elliptic curves over p -adic Lie extensions*, Math. Ann. **331** (2005), no. 4, 809–839.
- [Del71] Pierre Deligne, *Formes modulaires et représentations l -adiques*, Séminaire Bourbaki. Vol. 1968/69: Exposés 347–363, Lecture Notes in Math., vol. 175, Springer, Berlin, 1971, pp. Exp. No. 355, 139–172.
- [dJ95] Aise J. de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*, Publications mathématiques de l’IHES **82** (1995), 5–95.
- [Edi92] Bas Edixhoven, *The weight in Serre’s conjectures on modular forms*, Inventiones mathematicae **109** (1992), no. 1, 563–594.
- [EPW06] Matthew Emerton, Robert Pollack, and Tom Weston, *Variation of Iwasawa invariants in Hida families*, Invent. Math. **163** (2006), no. 3, 523–580.
- [Gou88] Fernando Q. Gouvêa, *Arithmetic of p -adic modular forms*, Lecture Notes in Mathematics, vol. 1304, Springer-Verlag, Berlin, 1988.

- [Gre94] Ralph Greenberg, *Iwasawa theory and p -adic deformations of motives*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 193–223.
- [Gre06] ———, *On the structure of certain Galois cohomology groups*, Doc. Math. (2006), no. Extra Vol., 335–391.
- [GV00] Ralph Greenberg and Vinayak Vatsal, *On the Iwasawa invariants of elliptic curves*, Invent. Math. **142** (2000), no. 1, 17–63.
- [Hid89] Haruzo Hida, *Nearly ordinary Hecke algebras and Galois representations of several variables*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 115–134.
- [Hid93] ———, *Elementary theory of L -functions and Eisenstein series*, London Mathematical Society Student Texts, vol. 26, Cambridge University Press, Cambridge, 1993.
- [HL19] Jeffrey Hatley and Antonio Lei, *Arithmetic properties of signed Selmer groups at non-ordinary primes*, Ann. Inst. Fourier (Grenoble) **69** (2019), no. 3, 1259–1294.
- [JS11] Somnath Jha and Ramdorai Sujatha, *On the Hida deformations of fine Selmer groups*, J. Algebra **338** (2011), 180–196.
- [Kat04] Kazuya Kato, *p -adic Hodge theory and values of zeta functions of modular forms*, Astérisque (2004), no. 295, ix, 117–290, Cohomologies p -adiques et applications arithmétiques. III.
- [Kim09] Byoung Du Kim, *The Iwasawa invariants of the plus/minus Selmer groups*, Asian J. Math. **13** (2009), no. 2, 181–190.
- [Kis03] Mark Kisin, *Overconvergent modular forms and the Fontaine–Mazur conjecture*, Invent. Math. **153** (2003), no. 2, 373–454.
- [KLP19] Chan-Ho Kim, Jaehoon Lee, and Gautier Ponsinet, *On the Iwasawa invariants of Kato’s zeta elements for modular forms*, <https://arxiv.org/abs/1909.01764>, 2019.
- [KNS22] Debanjana Kundu, Filippo A. E. Nuccio Mortarino Majno di Capriglio, and Ramdorai Sujatha, *Structure of fine Selmer groups in abelian p -adic Lie extensions*, <https://hal-cnrs.archives-ouvertes.fr/hal-03769801v3>, September 2022, preprint.
- [Kob03] Shin-ichi Kobayashi, *Iwasawa theory for elliptic curves at supersingular primes*, Invent. Math. **152** (2003), no. 1, 1–36.
- [Kur02] Masato Kurihara, *On the Tate Shafarevich groups over cyclotomic fields of an elliptic curve with supersingular reduction. I*, Invent. Math. **149** (2002), no. 1, 195–224.
- [LL22] Antonio Lei and Meng Fai Lim, *On fine Selmer groups and signed Selmer groups of elliptic modular forms*, Bull. Aust. Math. Soc. **105** (2022), no. 3, 419–430.
- [LMF13] LMFDB Collaboration, *The L -functions and Modular Forms Database*, <http://www.lmfdb.org>, 2013.
- [LP20] Antonio Lei and Gautier Ponsinet, *On the Mordell–Weil ranks of supersingular abelian varieties in cyclotomic extensions*, Proc. Amer. Math. Soc. Ser. B **7** (2020), 1–16.
- [LS18] Meng Fai Lim and Ramdorai Sujatha, *Fine Selmer groups of congruent Galois representations*, J. Number Theory **187** (2018), 66–91.
- [LZ16] David Loeffler and Sarah Livia Zerbes, *Rankin–Eisenstein classes in Coleman families*, Res. Math. Sci. **3** (2016), Paper No. 29, 53.
- [Mat20] Ahmed Matar, *On the Λ -cotorsion subgroup of the Selmer group*, Asian Journal of Mathematics **24** (2020), no. 3, 437–456.
- [Nak20] Kentaro Nakamura, *Zeta morphisms for rank 2 universal deformations*, <https://arxiv.org/abs/2006.13647>, 2020, preprint.
- [NO16] Filippo A. E. Nuccio Mortarino Majno di Capriglio and Tadashi Ochiai, *Coleman Map in Coleman Families*, <https://arxiv.org/abs/1606.04441v3>, 2016.
- [NS21] Filippo A. E. Nuccio Mortarino Majno di Capriglio and Ramdorai Sujatha, *Residual supersingular Iwasawa theory and signed Iwasawa invariants*, Rendiconti del Seminario Matematico di Padova (2021), accepted for publication, DOI 10.4171/RSMUP/111.
- [Och05] Tadashi Ochiai, *Euler system for Galois deformations*, Ann. Inst. Fourier (Grenoble) **55** (2005), no. 1, 113–146.
- [Pol] Robert Pollack, *Tables of Iwasawa invariants of elliptic curves*, <https://math.bu.edu/people/rpollack/Data/data.html>.

- [PR95] Bernadette Perrin-Riou, *Fonctions L p -adiques des représentations p -adiques*, Astérisque (1995), no. 229, 198.
- [PR04] Robert Pollack and Karl Rubin, *The main conjecture for CM elliptic curves at supersingular primes*, Ann. of Math. (2) **159** (2004), no. 1, 447–464. MR 2052361
- [Rib77] Kenneth A. Ribet, *Galois representations attached to eigenforms with Nebentypus*, Modular functions of one variable, V (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), Springer, Berlin, 1977, pp. 17–51. Lecture Notes in Math., Vol. 601.
- [Spr16] Florian Sprung, *The Iwasawa Main Conjecture for elliptic curves at odd supersingular primes*, arXiv preprint arXiv:1610.10017 (2016).
- [Tay91] Richard Taylor, *Galois representations associated to Siegel modular forms of low weight*, Duke Math. J. **63** (1991), no. 2, 281–332.
- [Wan21] Xin Wan, *Iwasawa main conjecture for supersingular elliptic curves and BSD conjecture*, arXiv preprint arXiv:1411.6352v8 (2021).
- [Wil88] Andrew Wiles, *On ordinary λ -adic representations associated to modular forms*, Invent. Math. **94** (1988), no. 3, 529–573.
- [Win89] Kay Wingberg, *Duality theorems for abelian varieties over \mathbf{Z}_p -extensions*, Algebraic number theory, Adv. Stud. Pure Math., vol. 17, Academic Press, Boston, MA, 1989, pp. 471–492.

(F. A. E. Nuccio) UNIVERSITÉ JEAN MONNET SAINT-ÉTIENNE, CNRS, INSTITUT CAMILLE JORDAN UMR 5208, F-42023, SAINT-ÉTIENNE, FRANCE

Email address: `filippo.nuccio@univ-st-etienne.fr`

(T. Ochiai) DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1 OOKAYAMA, MEGURO-KU, TOKYO 152-8551, JAPAN

Email address: `ochiai@math.titech.ac.jp`

(J. Ray) HARISH-CHANDRA RESEARCH INSTITUTE CHHATNAG ROAD, JHUNSI, PRAYAGRAJ (ALLAHABAD) 211 019 INDIA

Email address: `jishnuray@hri.res.in; jishnuray1992@gmail.com`