



**HAL**  
open science

## Measuring cortisol in fish scales to study stress in wild tropical tuna

Emmanuelle Roque d'Orbcastel, Yvan Bettarel, Marion Dellinger, Bastien Sadoul, Thierry Bouvier, Justin Monin, Laurent Dagorn, Benjamin Geffroy

► **To cite this version:**

Emmanuelle Roque d'Orbcastel, Yvan Bettarel, Marion Dellinger, Bastien Sadoul, Thierry Bouvier, et al.. Measuring cortisol in fish scales to study stress in wild tropical tuna. *Environmental Biology of Fishes*, 2021, 104, pp.725-732. 10.1007/s10641-021-01107-6 . hal-03357766

**HAL Id: hal-03357766**

**<https://hal.science/hal-03357766>**

Submitted on 29 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Measuring cortisol in fish scales to study stress in wild tropical tuna**

2 **Emmanuelle Roque d'Orbcastel<sup>1</sup>, Yvan Bettarel<sup>1</sup>, Marion Dellinger<sup>1,2</sup>, Bastien Sadoul<sup>1,3</sup>, Thierry**  
3 **Bouvier<sup>1</sup>, Justin Monin Amandé<sup>4,5</sup>, Laurent Dagorn<sup>1</sup>, Benjamin Geffroy<sup>1</sup>**

4 <sup>1</sup> MARBEC Univ Montpellier, CNRS, Ifremer, IRD, Sète, France

5 <sup>2</sup> Department of Aquaculture and Fish Biology, Hólar University, Hólar, Iceland

6 <sup>3</sup> ESE, Ecology and Ecosystem Health, Inrae, Institut Agro, Rennes, Brittany, France

7 <sup>4</sup> Centre de Recherches Océanologiques (CRO), DRAV, 29, Rue des pêcheurs, BP V 18, Abidjan, Côte  
8 d'Ivoire

9 <sup>5</sup> Cocody 2 plateaux Aghien, 01 BP 3012 Abidjan 01, Abidjan, Côte d'Ivoire

10 Author to whom correspondence should be addressed: Tel.: +33467130414 and email:  
11 [emmanuelle.roque@ifremer.fr](mailto:emmanuelle.roque@ifremer.fr)

12

## 13 **Abstract**

14 Cortisol is recognized as a physiological indicator of stress in fish. However, this hormone is typically  
15 measured in plasma samples. In this study, cortisol content was evaluated for the first time in the scales  
16 of tropical tuna. The sample included 20 skipjack tuna (*Katsuwonus pelamis*) and 25 yellowfin tuna  
17 (*Thunnus albacares*) collected in the Atlantic (the Gulf of Guinea off the Ivory Coast) and the Indian  
18 Ocean (off Reunion Island), and their scales were analyzed using an ELISA test. The results indicated  
19 that on average, cortisol concentration (1) did not show any geographical pattern, (2) was independent  
20 of fish size, and (3) was significantly higher in skipjack (mean:  $4.75 \pm 6.56 \text{ ng g}^{-1}$ ) than in yellowfin  
21 (mean:  $1.65 \pm 1.85 \text{ ng g}^{-1}$ ), although the difference was mainly due to four skipjack individuals. Larger  
22 datasets would be needed to confirm any species difference in cortisol concentration. The particularly  
23 high cortisol concentration observed in four individuals may be due to fish from different schools,  
24 suggesting the need for future sampling from free-swimming schools to investigate intra- and inter-  
25 school variability in cortisol concentration. This study opens the door for future research, including the  
26 collection of scales on tagged individuals, to investigate the links between chronic stress and behavior  
27 in these species.

28 **Key words** fish scales · cortisol · stress · welfare · skipjack tuna · yellowfin tuna

## 29 **Introduction**

30 The capacity of fish to cope with environmental challenges is of primary importance in a rapidly  
31 changing world. The stress axis plays a central role in the coping capacity of fish, as it diverts energy  
32 from growth and reproduction to cope with acute environmental changes (Sadoul and Vijayan 2016).  
33 The two stress-related response mechanisms in fish are brain-sympathetic chromaffin cells (BSC) and  
34 the hypothalamus-pituitary-interrenal (HPI) axis. The underlying mechanisms of the stress axis in fish  
35 have been well described (Barton 2002; Shreck and Tort 2016). At the physiological level, the primary  
36 response to a stressor in fish is neuroendocrinal and involves the activation by the central nervous system

37 of the sympathetic nervous system, releasing neurotransmitters and hormones such as catecholamine  
38 and corticosteroids (mainly cortisol in fish). Following the stimulus, plasma cortisol increases after a  
39 period of time ranging from a few minutes to an hour, depending on the species and the intensity and/or  
40 duration of the stimuli (Barton 2002). Cortisol is considered a good quantitative indicator of acute and  
41 chronic physiological stress (Ellis et al. 2012). Until recently, this hormone was mainly measured in the  
42 plasma of unstressed and stressed fish, providing a basal and high cortisol level. Yet plasma measures  
43 of cortisol can only provide insight into the stress level of the fish at a given time. In an ecological  
44 context, an integrative value of stress over a long period is much more valuable, allowing a more  
45 comprehensive idea of the life history of the animal. Recently, new matrices have emerged to measure  
46 cortisol; these are being used as relatively non-invasive markers of stress (Sadoul and Geffroy 2019).  
47 Among these markers, fish scales are the most promising matrix as they provide a record of the  
48 integrated stress response over time (i.e. from weeks to months, Aerts et al. 2015) and a grid to highlight  
49 behavioral ecology regarding exposure to stress factors such as temperature (Goikoetxea et al. 2020),  
50 since the metabolic clearance rate in scales is much slower than in plasma (Laberge et al. 2019). Previous  
51 studies have shown that intra-specific differences in basal cortisol are genetically driven (Øverli et al.  
52 2002a) and are generally aligned with differences in behavior or coping style, in which reactive  
53 individuals display higher basal cortisol levels compared to proactive individuals (Koolhaas et al. 1999).  
54 In trout, for instance, long-term cortisol administration was shown to decrease both locomotor activity  
55 and aggression (Øverli et al. 2002b). This could be transcribed at the scale level with cortisol  
56 accumulation.

57 Tuna are present in all oceans, and as top predators they play a major ecological role in offshore pelagic  
58 ecosystems. Worldwide, tuna species are exploited by both artisanal and industrial fisheries, using  
59 different types of fishing gear. Two tuna species, skipjack (SKJ, *Katsuwonus pelamis*) and yellowfin  
60 (YFT, *Thunnus albacares*), are among the most landed species in the world (FAO 2018). Although the  
61 biology and ecology of tuna have been extensively investigated, particularly to provide scientific advice  
62 for fishery management and the conservation of stocks, studies measuring cortisol in tuna are scarce and  
63 limited to only one species (Atlantic bluefin tuna, *Thunnus thynnus*) in one environment (Kirchhoff et  
64 al. 2011; Addis et al. 2013; Tracey et al. 2016). By measuring the basal levels of plasma cortisol, these  
65 studies showed a rapid accumulation of cortisol in response to an intense one-off event (capture) and  
66 focused on the short-term stress reaction to capture. In general, captured fish are exposed to stressful  
67 stimuli, e.g. physical trauma and physiological stress (Skomal 2007), which is worth investigating  
68 mainly in an ethical context. Our study chose another line of approach, using, for the first time, fish  
69 scales as a matrix to measure cortisol in two wild marine fish, skipjack and yellowfin tuna (male and  
70 female), in order to describe the causal effects of size and location on this indicator.

## 71 **Materials and methods**

72 The sample for the study was based on individuals from two tuna species, YFT and SKJ, captured by  
73 handline around anchored fish-aggregating devices (FADs) in the Atlantic (Gulf of Guinea off the Ivory  
74 Coast) and the Indian Ocean (off Reunion Island) (Fig. 1A and B) on 10–11 July and 26–29 September  
75 2018, respectively. The total time between the bite on the hook and the killing of a tuna was between 2  
76 and 10 minutes. To minimize the pain, we euthanized the animals as soon as they were boarded, with a  
77 blow to the head that immediately killed the tuna (following European directive 2010/63/UE). The  
78 number of fish sampled was  $n=22$  in the Gulf of Guinea (16 YFT, 6 SKJ) and  $n=23$  off Reunion Island  
79 (9 YFT, 14 SKJ).

80 Onboard, the fork length (FL in cm, from the tip of the upper jaw to the caudal fin fork) and body mass  
81 (W in kg) were measured by scientists. Scale samples (under the pectoral fin) were taken by scraping  
82 the flank with a blunt scalpel and these were collected in plastic Eppendorf tubes, which were  
83 immediately immersed in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until analysis. The handlers wore gloves  
84 during the entire process to avoid external contamination.

85 The cortisol extraction from the scale samples ( $n=45$ ) was adapted from two different protocols (Aerts  
86 and De Saeger 2017; Carbajal et al. 2018). Scales were washed by adding 1.5 mL of isopropanol in each  
87 tube, agitated with a ceramic ball mill at 10 Hz for 3 min in a Retsch MM 400 tissue lyzer, and  
88 centrifuged at 14,000 g and  $4^{\circ}\text{C}$  for 1 min. The supernatant was then discarded. This washing process  
89 was repeated three times to fully remove the covering mucus. The samples were then stored at  $-20^{\circ}\text{C}$   
90 for 1 h and lyophilized (2 h freezing phase down to  $-40^{\circ}\text{C}$ , followed by a rapid increase up to  $40^{\circ}\text{C}$   
91 lyophilization phase in a 1000 mbar vacuum overnight). Once dried, subsamples were aliquoted in a  
92 new 2 mL Eppendorf tube and precisely weighed (mean weight and SE:  $95.5 \pm 9$  mg). One ceramic mill  
93 ball was then added to each tube and samples were minced to powder in the tissue lyzer for 1 min at 30  
94 Hz. Cortisol extraction was performed by adding 1 mL of methanol to each tube, vortexed for 10 sec  
95 and shaken overnight (18 h) at room temperature. Then the samples were centrifuged at 12,000 g for 3  
96 min, and the supernatant transferred to a borosilicate glass evaporation tube. This last step was repeated  
97 after adding 1 mL of methanol to ensure maximum cortisol extraction. The contents of the glass tube  
98 were then evaporated to dryness by nitrogen flow at  $40^{\circ}\text{C}$  using a MULTIVAP® Nitrogen evaporator  
99 (Organomation®). All samples were then reconstituted in 150  $\mu\text{L}$  of 0.01% Bovine Serum Albumine  
100 (BSA) in Phosphate Buffered Saline solution (PBS – 2.7 M KCl; 0.137 M NaCl; 1.8 mM  $\text{KH}_2\text{PO}_4$ ;  
101  $\text{Na}_2\text{HPO}_4$ ), vortexed 3 times for 15 seconds and stored at  $4^{\circ}\text{C}$  until the ELISA assay. The cortisol was  
102 quantified using a competitive ELISA measurement described in Faught et al. (2016) and adapted from  
103 a protocol validated for zebrafish (*Danio rerio*) (Yeh et al. 2013). This involved coating 96-well plates  
104 with a cortisol monoclonal antibody ( $1.6 \mu\text{g}\cdot\text{mL}^{-1}$ ; East Coast Bio, ME, USA) in PBS and storing them  
105 at  $4^{\circ}\text{C}$  for 16 hours. The cross-reactivity of the antibody is 0.4% for cortisone, 7.7% for 11-deoxycortisol  
106 and 21.9% for corticosterone (Sadoul and Geffroy 2019). While this is a high level of cross-reactivity  
107 with corticosterone, it should be noted that levels of this hormone have been reported to be relatively

108 low in fish (at least 10 times lower than cortisol, as measured in plasma) (Kiilerich and Prunet 2011), so  
109 the specificity of the monoclonal antibody used here and in other fish studies (Yeh et al. 2013; Faught  
110 et al. 2016; Best et al. 2017) should only marginally change results. Standards were obtained by serially  
111 diluting hydrocortisone (Sigma) in PBS (0–25 ng.mL<sup>-1</sup>), and 50µL of either standards or samples were  
112 added to the wells in duplicate. Each well then received cortisol conjugated with horseradish peroxidase  
113 (HRP) and filled with a detection solution that turns blue when in contact with HRP. Sulfuric acid (1N)  
114 was used to stop the reaction. Absorption was read at 450 nm (EL800 Universal Microplate Reader,  
115 BIO-TEK INSTRUMENTS, INC.). Cortisol ELISA concentrations (in ng.mL<sup>-1</sup>) were normalized by the  
116 sample weight of each sample (expressed in ng of cortisol per g of scales). The intra-assay CV for the  
117 two plates was below 10%. The standard curve was performed with 0.5, 1, 2, 5, and 10 ng.mL<sup>-1</sup>, and the  
118 cortisol concentration measured in the scales was correlated with expected cortisol values for plate 1  
119 ( $r^2=0.95$ ) and plate 2 ( $r^2=0.98$ ).

120 We first compared cortisol concentration in scales between species caught at the same site and tested  
121 for a potential fish-size effect. Then we pooled all the data together to test whether the cortisol  
122 concentration in scales varied as a function of size, location or species in the same linear mixed model.  
123 We added the ELISA plate as a random factor, as samples were run on separate ELISA plates. The  
124 interaction was removed from the model when not significant. To ensure that all fixed effects were taken  
125 into account similarly, we used a type II Analysis of Variance with Satterthwaite’s method. All statistical  
126 analyses were performed using R software 3.2.2 (R Development Core Team 2013) and package “lme4”  
127 for linear mixed models (Bates et al. 2015). Maps were designed using the “ggmap” package (Kahle  
128 and Wickham 2013).

## 129 **Results and discussion**

130 We did not detect significant differences in cortisol concentration in scales as a function of fish length  
131 for either species (YFT:  $t=-0.24$ ;  $p=0.8$ ; SKJ:  $t=-0.14$ ;  $p=0.9$ ) (Table 1). Within each site, the cortisol  
132 concentration in SKJ scales was higher (although not significantly) than that in YFT scales ( $p=0.17$  in  
133 the Gulf of Guinea and  $p=0.09$  off Reunion Island), so that when pooling all individuals together in the  
134 model accounting for all fixed effects (Cortisol ~ Location + Length + Species + (1|PlateScales)),  
135 cortisol concentration was significantly higher ( $t=1.97$ ;  $p=0.0558$ ) in SKJ scales compared to YFT  
136 scales, independently of their size and location (Fig. 2). However, it is worth noting that this pattern was  
137 driven by 4 SKJ individuals between 54 and 59 cm in FL (3 individuals from Reunion Island and 1  
138 individual from the Gulf of Guinea). According to the sampling size of this study, neither the  
139 size:species interactions nor the size:location interactions were found to be significant ( $p=0.5$  &  $0.6$   
140 respectively).

141 Cortisol in scales has been proven to be an effective marker of chronic stress in various fish species  
142 (Aerts et al. 2015; Carbajal et al. 2019, Laberge et al. 2019; Hanke et al. 2020). Yet the mechanisms of  
143 cortisol clearance in scales are not well described, and to our knowledge only one recent study on

144 goldfish (*Carassius auratus*) has provided insights regarding its rate (Laberge et al. 2019). That study  
145 demonstrated that following a cortisol treatment, plasma levels went back to initial values after only 3  
146 days, while it took 27 days to return to the basal level in scales. This implies that cortisol is somehow  
147 catabolized at a much slower rate in scales than in plasma, likely because this steroid is embedded in  
148 calcified structures that protect it. The cortisol values found for both species in our study were well  
149 within the range of those reported for other fish species, e.g. European seabass (*Dicentrarchus labrax*)  
150 (from 1.7 to 3.4 ng g<sup>-1</sup>, Aerts and De Saeger 2017) and goldfish (*Carassius auratus*) (from 0.25 to 2.10  
151 ng.g<sup>-1</sup>, Carbajal et al. 2018). Although our study relied on a small dataset, the cortisol concentrations in  
152 scales were significantly higher in skipjack (mean: 4.75 ± 6.56 ng.g<sup>-1</sup>) than in yellowfin (mean: 1.65 ±  
153 1.85 ng.g<sup>-1</sup>) for individuals of the same size. Interestingly, the cortisol concentration did not increase  
154 with fish size (i.e. age), which argues for a slow – but active – renewal of cortisol in this matrix, as  
155 demonstrated by Laberge et al. (2019), rather than an accumulation over time.

156 Adult skipjack and juvenile yellowfin tuna (the size classes sampled in our study) are known to share  
157 many anatomical, physiological and behavioral similarities. They have similar swimming modes,  
158 thermal biology, increased rate functions (e.g. standard metabolic rate, aerobic capacity, heart rate and  
159 gut clearance) and cardiac physiology (Graham and Dickson 2004), which distinguish them from many  
160 other teleost species. They have the same vertical habitat utilization (Schaefer et al. 2009) and usually  
161 feed on the same prey species (Jaquemet et al. 2011), although some diet differences have been observed  
162 in specific oceanic regions (Olson et al. 2016). They both form large schools and associate with floating  
163 objects, but differences in their fine-scale FAD-associative behavior have been observed (Rodriguez-  
164 Tress et al. 2017). Considering the strong similarities between these two species, one might expect  
165 similar cortisol metabolism, resulting in similar concentrations of cortisol in scales. Larger datasets are  
166 needed to confirm the differences between these two species. In parallel, experiments on captive tuna  
167 should be conducted in order to investigate the time evolution of cortisol concentration over several  
168 days and weeks. If a species difference in cortisol concentration is confirmed by larger datasets, this  
169 may be explained by species-specific cortisol metabolism. This would imply intrinsic differences in the  
170 production and/or clearance of cortisol (dependent upon cortisol-binding proteins, target tissue  
171 receptors, tissue uptake and catabolism of cortisol) (Mommensen et al. 1999). The production/clearance  
172 of cortisol and the rate of scale formation (and therefore the associated integration of cortisol) in these  
173 two species have not yet been investigated and represent a perspective for future studies.

174 In our dataset, however, the difference in cortisol concentration observed between the two species was  
175 mainly due to four skipjack individuals. Interestingly, comparable high values were also detected in  
176 some extremely stressed European seabass (Aerts and de Saeger 2017) and goldfish (Laberge et al.  
177 2019). These high concentrations of cortisol in scales may be the consequence of specific stressful  
178 events experienced by these individuals, such as attacks by predators or fasting periods that might have  
179 occurred over the preceding days or weeks. Understanding why some individuals of the same species

180 and the same size, in the same fish aggregation around a FAD, have significantly higher concentrations  
181 of cortisol (about five times higher) would be valuable to study on a physiological and behavioral level.  
182 Considering that the clearance rate of cortisol in scales is very slow, it may be that individuals with high  
183 concentrations of cortisol belong to different schools than the fish with lower cortisol concentrations.  
184 Tuna aggregations at FADs usually consist of several schools (Josse et al. 2000; Dagorn et al. 2007;  
185 Moreno et al. 2007). As fish from the same school are likely to have a relatively long common history  
186 (although the stability of a tuna school appears to vary quite a lot, see Bayliff 1988 and Hilborn 1991),  
187 it can be assumed that fish from the same school have comparable concentrations of cortisol as they  
188 have usually experienced the same events. This hypothesis could be investigated by comparing intra-  
189 and inter-school variability of cortisol concentrations in individuals of the same species sampled from  
190 different free-swimming schools caught by purse seiners (as sampling tuna associated with FADs does  
191 not ensure that they are from the same school).

## 192 **Conclusion**

193 This preliminary study provides the first measurements of cortisol in the scales of wild tuna. While  
194 further studies are needed to better understand the mechanisms of scale cortisol clearance and  
195 metabolism in the selected species (e.g. a comparison of the cortisol level in scales and in other  
196 biological matrices: mucus, plasma, muscle, etc.), this data offers a good start for designing new  
197 experiments. In the same way Willis and Hobday (2008) suggested taking bioelectrical impedance  
198 analysis (BIA) measurements to measure the condition of tuna during tagging operations, we suggest  
199 sampling a few scales from individuals during conventional or electronic tagging. This would be a quick  
200 and harmless operation that should not disrupt tagging operations and would allow measuring the  
201 markers of chronic stress (as opposed to the stress due to capture, measured by cortisol in plasma) of  
202 tagged individuals. Comparing the large-scale movements (from conventional or archival tags) or the  
203 fine-scale movements within arrays of FADs (from acoustic tags) for fish with different concentrations  
204 of cortisol would provide key insights into the ecology of these species, particularly the possible links  
205 between stress response and behavior.

206

207 ***Supplementary Information.*** *Not applicable.*

## 208 ***Acknowledgments***

209 This study was conducted in the framework of the Tuna MicrobiOME (THE MOME) project, with  
210 funding from Montpellier University of Excellence (MUSE) and the Marine Biodiversity and  
211 Conservation (MARBEC) research group.

212 The sampling and slaughtering of animals was carried out by professional fishermen working for the  
213 French National Research Institute for Sustainable Development (IRD) Exploited Tropical Pelagic  
214 Ecosystems Observatory, which is ISO 9001:2015 certified.

215 The authors would also like to thank the professional fishermen from La Réunion Saint Gilles Fishing  
216 Club for their help in fish sampling. We also thank Aurélie Guillou (IRD), who assisted us in sampling  
217 some specimens. We are grateful to Antoinette Adingra, Nadège Kouadio and Constance Diaha,  
218 colleagues from the Oceanology Research Center of Abidjan, for their contribution in the fish sex  
219 assessment.

220 Finally, many thanks to the editor (Dr. Margaret Docker) and referees, who provided very constructive  
221 comments on the initial version of this manuscript, and to Elise Bradbury for the English proofreading.

#### 222 ***Authors' contributions***

223 E.R.O. and B.G. designed the study, E.R.O., Y.B. and T.B. collected the samples, and M.D., B.S, B.G.  
224 and E.R.O. performed the analyses. All authors prepared the manuscript and approved its submission.

#### 225 ***Funding***

226 The Tuna MicrobiOME project (THE MOME) receives funding from Montpellier University of  
227 Excellence (MUSE) and the Marine Biodiversity and Conservation (MARBEC) research group.

228 ***Code availability.*** *Not applicable.*

229 ***Data availability.*** *Not applicable.*

#### 230 ***Declarations***

#### 231 ***Ethics approval***

232 All sampling and animal handling respected good scientific practice and animal welfare rules. To  
233 minimize suffering, we euthanized the animals as soon as they were boarded. The number of fish  
234 sampled was limited to the strict number necessary.

235 ***Consent to participate.*** *Not applicable.*

236 ***Consent for publication.*** *Not applicable.*

237 ***Conflict of Interest/ competing interests.*** The authors declare that they have no conflict of interest  
238 concerning this manuscript and no competing financial interests that could influence the work reported  
239 in this paper.

240

241


242

243

244


245





246  
 247  
 248  
 249  
 250  
 251  
 252  
 253

**Fig. 1.** Tuna species, skipjack (*Katsuwonus pelamis*) and yellowfin (*Thunnus albacares*) were sampled in A) the Gulf of Guinea (Atlantic Ocean), 27 nautical miles off Abidjan (Ivory Coast) on July 2018 and B) La Réunion Island (Indian Ocean), 10 nautical miles from shore, on September 2018. Red dots indicate the sampling location of the Fish Aggregating Devices (FAD) where tunas have been captured.



254  
255  
256  
257  
258  
259  
260

**Fig. 2.** Cortisol concentrations ( $\text{ng}\cdot\text{g}^{-1}$ ) measured in the scales of the two tuna species *K. pelamis* (SKJ) and *T. albacares* (YFT), in the Atlantic and Indian oceans, correlated to the fish length (in cm).

261 **Table 1.** Biometrics data (number, length, weight) of the two tuna species (sp), skipjack (SKJ,  
 262 *Katsuwonus pelamis*) and yellowfin (YFT, *Thunnus albacares*)

	sp	nb	♂	♀	ND	length (cm) min-max	length (cm) average ± SD	weight (kg) min-max	weight (kg) average ± SD
Gulf of Guinea	YFT	16	7	6	3	46.5-66	58.08±5.22	1.6-4.6	3.13±0.87
Gulf of Guinea	SKJ	6	3	3	0	56-66	59.78±3.95	2.9-4.4	3.52±0.52
La Réunion Island	YFT	9	3	6	0	61.5-69.6	64.61±2.77	4.1-5.5	4.51±0.45
La Réunion Island	SKJ	14	4	10	0	41.4-60	51.57±5.62	1.2-4.0	2.45±0.79

263  
 264

265 **References**

- 266 Addis P, Corrias S, Garau C, Secci M (2013) Physiologic Responses to Stress and Changes in Atlantic  
267 Bluefin Tuna (*T. thynnus*) Meat Color During Trap Fisheries Capture and Processing in Sardinia  
268 (W. Mediterranean). J Aquat Food Prod Technol 22: 298–309.  
269 <https://doi.org/10.1080/10498850.2011.647596>
- 270 Aerts J, Metz J R, Ampe B, Decostere A, Flik G, Saeger SD (2015) Scales tell a story on the stress  
271 history of fish. PLoS One 10, e0123411. <https://doi.org/10.1371/journal.pone.0123411>
- 272 Aerts J, De Saeger S (2017) Quantification of glucocorticoids in fish scales as biomarkers for chronic  
273 stress. United States Patent 9, 638,703 B2
- 274 Barton BA (2002) Stress in fishes: a diversity of responses with particular reference to changes in  
275 circulating corticosteroids. Integr Comp Biol 42: 517–525. <https://doi.org/10.1093/icb/42.3.517>
- 276 Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. J  
277 Stat Softw 67: 1–48. <http://dx.doi.org/10.18637/jss.v067.i01>.
- 278 Bayliff WH (1988) Integrity of schools of skipjack tuna, *Katsuwonus pelamis*, in the eastern Pacific  
279 Ocean, as determined from tagging data. Fish Bull NOAA-NMFS 86(4): 631-643
- 280 Best C, Kurrasch DM, Vijayan MM (2017). Maternal cortisol stimulates neurogenesis and affects larval  
281 behaviour in zebrafish. Sci Rep 7(1), 1-10. <https://doi.org/10.1038/srep40905>
- 282 Carbajal A, Monclús L, Tallo-Parra O, Sabes-Alsina M, Vinyoles D, Lopez-Bejar M (2018) Cortisol  
283 detection in fish scales by enzyme immunoassay: Biochemical and methodological validation. J  
284 Appl Ichthyol 34: 967–970. <https://doi.org/10.1111/jai.13674>
- 285 Carbajal A, Reyes-López FE, Tallo-Parra O, Lopez-Bejar M, Tort L (2019). Comparative assessment  
286 of cortisol in plasma, skin mucus and scales as a measure of the hypothalamic-pituitary-interrenal  
287 axis activity in fish. Aquaculture, 506, 410-416. <https://doi.org/10.1016/j.aquaculture.2019.04.005>
- 288 Dagorn L, Holland KN, Itano DG (2007) Behaviour of yellowfin (*Thunnus albacares*) and bigeye (*T.*  
289 *obesus*) tuna in a network of fish aggregating devices (FADs). Mar Biol 151: 595–606.  
290 <https://doi.org/10.1007/s00227-006-0511-1>
- 291 Ellis T, Yildiz HY, López Olmeda J, Spedicato MT, Tort L, Øverli Ø, Martins CIM (2012) Cortisol and  
292 finfish welfare. Fish Physiol Biochem 38: 163–188. <https://doi.org/10.1007/s10695-011-9568-y>
- 293 Faught E, Best C, Vijayan MM (2016). Maternal stress-associated cortisol stimulation may protect  
294 embryos from cortisol excess in zebrafish. R Soc Open Sci 3(2), 160032.  
295 <https://doi.org/10.1098/rsos.160032>
- 296 Goikoetxea A, Sadoul B, Blondeau-Bidet E, Aerts J, Blanc MO, Parrinello H, Geffroy B (2020).  
297 Genetic pathways underpinning hormonal stress responses in fish exposed to short-and long-term  
298 warm ocean temperatures. Ecol Indic 120, 106937. <https://doi.org/10.1016/j.ecolind.2020.106937>
- 299 Graham JB, Dickson KA (2004) Tuna Comparative physiology. J Exp Biol 207 (Pt 23): 4015-4024.  
300 <https://doi.org/10.1242/jeb.01267>
- 301 Hanke I, Hassenrück C, Ampe B, Kunzmann A, Gärdes A, Aerts J (2020). Chronic stress under  
302 commercial aquaculture conditions: Scale cortisol to identify and quantify potential stressors in  
303 milkfish (*Chanos chanos*) mariculture. Aquaculture, 526, 735352.  
304 <https://doi.org/10.1016/j.aquaculture.2020.735352>
- 305 Hilborn R (1991) Modeling the stability of fish schools: exchange of individual fish between schools of  
306 skipjack tuna (*Katsuwonus pelamis*). Can J Fish Aquat Sci 48: 1081-1091.  
307 <https://doi.org/10.1139/f91-128>
- 308 Jaquemet S, Portier M, Ménard F (2011) Do drifting and anchored Fish Aggregating Devices (FADs)  
309 similarly influence tuna feeding habits? A case study from the western Indian Ocean. Fish Res 107:  
310 283-290. <https://doi.org/10.1016/j.fishres.2010.11.011>

311 Josse E, Dagorn L, Bertrand A (2000) Typology and behaviour of tuna aggregations around fish  
312 aggregating devices from acoustic surveys in French Polynesia. *Aquat Liv Resour* 13: 183–192

313 Kahle D, Wickham H (2013) ggmap: Spatial Visualization with ggplot2. *R J* 5(1): 144-161

314 Kiilerich P, Prunet P (2011). Corticosteroids, in: *Encyclopedia of Fish Physiology: From Genome to*  
315 *Environment*. Volume 2. Gas Exchange, Internal Homeostatis, and Food Uptake. Academic Press  
316 - Elsevier, p. np.

317 Kirchhoff NT, Leef MJ, Ellis D, Purser J, Nowak BF (2011) Effects of the first two months of ranching  
318 on the health of southern bluefin tuna *Thunnus maccoyii*. *Aquac* 315: 207–212.  
319 <https://doi.org/10.1016/j.aquaculture.2011.02.036>

320 Koolhaas JM, Korte SM, De Boer SF, Van Der Vegt BJ, Van Reenen CG, Hopster H, De Jong IC, Ruis  
321 MAW, Blokhuis HJ (1999). Coping styles in animals: Current status in behavior and stress-  
322 physiology. *Neurosci Biobehav Rev*, 23(7), 925–935. [https://doi.org/10.1016/S0149-](https://doi.org/10.1016/S0149-7634(99)00026-3)  
323 [7634\(99\)00026-3](https://doi.org/10.1016/S0149-7634(99)00026-3)

324 Laberge F, Yin-Liao I, Bernier NJ (2019) Temporal profiles of cortisol accumulation and clearance  
325 support scale cortisol content as an indicator of chronic stress in fish. *Conserv Physiol* 7(1) coz052.  
326 <https://doi.org/10.1093/conphys/coz052>

327 Mommsen T, Vijayan M, Moon T (1999) Cortisol in teleosts: dynamics, mechanisms of action, and  
328 metabolic regulation, *Rev Fish Biol Fish* 9 (3): 211–268.  
329 <https://doi.org/10.1023/A:1008924418720>

330 Moreno G, Dagorn L, Sancho G, Itano D (2007) Fish behaviour from fishers' knowledge: The case  
331 study of tropical tuna around drifting fish aggregating devices (DFADs). *Can J Fish Aquat Sci* 64:  
332 10.1139/F07-113. <https://doi.org/10.1139/F07-113>

333 Olson RJ, Young JW, Ménard F, Potier M, Allain V, Goñi N, Logan JM, Galván-Magaña F (2016)  
334 Chapter Four - Bioenergetics, Trophic Ecology, and Niche Separation of Tunas. *Adv Mar Biol* 74:  
335 199-344. <https://doi.org/10.1016/bs.amb.2016.06.002>

336 Øverli Ø, Pottinger TG, Carrick TR, Øverli E, Winberg S (2002a) Differences in behaviour between  
337 rainbow trout selected for high- and low-stress responsiveness. *J Exp Biol* 205, 391-395.  
338 <https://doi.org/10.1242/jeb.205.3.391>

339 Øverli Ø, Kotzian S, Winberg S (2002b) Effects of cortisol on aggression and locomotor activity in  
340 rainbow trout. *Horm Behav* 42, 53-61. <https://doi.org/10.1006/hbeh.2002.1796>

341 Rodriguez-Tress P, Capello M, Forget F, Soria M, Beeharry SP, Dussooa N, Dagorn L (2017)  
342 Associative behavior of yellowfin *Thunnus albacares*, skipjack *Katsuwonus pelamis* and bigeye  
343 tuna *Thunnus obesus* at fish aggregating devices (FADs) off the coast of Mauritius. *MEPS* 570.  
344 <https://doi.org/10.3354/meps12101>

345 Sadoul B, Geffroy B (2019) Measuring cortisol, the major stress hormone in fishes. *J Fish Biol* 94 (4):  
346 540-555. <https://doi.org/10.1111/jfb.13904>

347 Sadoul B, Vijayan MM (2016) 5—Stress and Growth. In C. B. Schreck, L. Tort, A. P. Farrell, & C. J.  
348 Brauner (Ed.), *Fish Physiol* 35: 167-205 Academic Press

349 Schaefer KM, Fuller DW, Block BA (2009) Vertical Movements and Habitat Utilization of Skipjack  
350 (*Katsuwonus pelamis*), Yellowfin (*Thunnus albacares*), and Bigeye (*Thunnus obesus*) Tunas in the  
351 Equatorial Eastern Pacific Ocean, Ascertained Through Archival Tag Data. In: Nielsen J.L.,  
352 Arrizabalaga H., Fragoso N., Hobday A., Lutcavage M., Sibert J. (eds) *Tagging and Tracking of*  
353 *Marine Animals with Electronic Devices. Reviews: Methods and Technologies in Fish Biol Fish*  
354 9. Springer, Dordrecht. <https://doi.org/10.1007/s00227-010-1524-3>

355 Schreck CB, Tort L (2016) Farrell AP, Brauner CJ (eds) *Biology of stress in fish*, *Fish Physiol* 35.  
356 Academic Press

- 357 Skomal GB (2007) Evaluating the physiological and physical consequences of capture on post-release  
358 survivorship in large pelagic fishes. *Fish Manag Ecol* 14: 81–89. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2400.2007.00528.x)  
359 [2400.2007.00528.x](https://doi.org/10.1111/j.1365-2400.2007.00528.x)
- 360 Tracey S, Hartmann K, Mc Allister J, Conron S, Leef M (2016) Capture-induced physiological stress  
361 and post-release survival of recreationally caught Southern Bluefin Tuna. *Can J Fish Aquat Sci* 73  
362 (10). <https://doi.org/10.1139/cjfas-2015-0516>
- 363 Willis J, Hobday AJ (2008) Application of bioelectrical impedance analysis as a method for estimating  
364 composition and metabolic condition of southern bluefin tuna (*Thunnus maccoyii*) during  
365 conventional tagging, *Fish Res* 93 (1-2): 64-71. <http://dx.doi.org/10.1016/j.fishres.2008.02.010>
- 366 Yeh CM, Glöck M, Ryu S (2013) An optimized whole-body cortisol quantification method for assessing  
367 stress levels in larval zebrafish. *PLoS One* 8, e79406. <https://doi.org/10.1371/journal.pone.0079406>