



**HAL**  
open science

# Hearing sensitivity of primates: recurrent and episodic positive selection in hair cells and stereocilia protein-coding genes

Andreia Moreira, Myriam Croze, Franklin Delehelle, Sylvain Cussat-Blanc, Hervé Luga, Catherine Mollereau, Patricia Balaesque

## ► To cite this version:

Andreia Moreira, Myriam Croze, Franklin Delehelle, Sylvain Cussat-Blanc, Hervé Luga, et al.. Hearing sensitivity of primates: recurrent and episodic positive selection in hair cells and stereocilia protein-coding genes. *Genome Biology and Evolution*, 2021, 13 (8), 10.1093/gbe/evab133 . hal-03357817

**HAL Id: hal-03357817**

**<https://hal.science/hal-03357817>**

Submitted on 29 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1  
2  
3  
4 **Hearing sensitivity of primates: recurrent and episodic positive selection in hair cells**  
5  
6 **and stereocilia protein-coding genes**  
7  
8  
9

10 Andreia Moreira<sup>1,2</sup>, Myriam Croze<sup>1,3\*</sup>, Franklin Delehelle<sup>1,2,4\*</sup>, Sylvain Cussat-Blanc<sup>2</sup>, Hervé  
11 Luga<sup>2</sup>, Catherine Mollereau<sup>1,5</sup> and Patricia Balaesque<sup>1,6+</sup>  
12  
13  
14  
15  
16

17 1. Anthropologie Moléculaire et Imagerie de Synthèse (AMIS), CNRS UMR5288, Université  
18 de Toulouse, Université Toulouse III Paul Sabatier, Faculté de Médecine Purpan, 37 allées  
19 Jules Guesde, 31073 Toulouse, France  
20  
21

22 2. Institut de Recherche en Informatique de Toulouse (IRIT), CNRS UMR5505, Université  
23 Toulouse III Paul Sabatier, Bâtiment 1R1, 118 route de Narbonne, 31400 Toulouse, France  
24  
25

26 3. Current address: Division of EcoScience and Department of Life Science, Ewha Womans  
27 University, Seoul, Korea  
28  
29

30 4. Current address: Institut de Biologie de l'Ecole Normale Supérieure (IBENS), CNRS  
31 UMR8197, École Normale Supérieure, Paris, France  
32  
33

34 5. Current address: Centre de Recherches sur la Cognition Animale (CRCA), CNRS  
35 UMR5169, Université de Toulouse, Université Toulouse III - Paul Sabatier, Bâtiment 4R4,  
36 118 route de Narbonne, 31062 Toulouse, France  
37  
38

39 6. Current address: Evolution et Diversité Biologique (EDB), CNRS UMR5174, Université de  
40 Toulouse, Université Toulouse III - Paul Sabatier, CNRS, IRD, Bâtiment 4R1, 118 route de  
41 Narbonne, 31062 Toulouse cedex 9, France  
42  
43  
44

45 \* These authors contributed equally to the work  
46  
47

48 + corresponding author: Patricia Balaesque, Université de Toulouse, Université III Paul  
49 Sabatier, 118 route de Narbonne, 31062 Toulouse, France; [patricia.balaesque@univ-  
50 tlse3.fr](mailto:patricia.balaesque@univ-tlse3.fr)  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1. Abstract

The large spectrum of hearing sensitivity observed in primates results from the impact of environmental and behavioral pressures to optimize sound perception and localization. Although evidence of positive selection in auditory genes has been detected in mammals including in Hominoids, selection has never been investigated in other primates. We analyzed 123 genes highly expressed in the inner ear of 27 primate species and tested to what extent positive selection may have shaped these genes in the order Primates tree. We combined both site and branch-site tests to obtain a comprehensive picture of the positively selected genes (PSGs) involved in hearing sensitivity, and drew a detailed description of the most affected branches in the tree. We chose a conservative approach, and thus focused on confounding factors potentially affecting PSG signals (alignment, GC-biased gene conversion, duplications, heterogeneous sequencing qualities). Using site tests, we showed that around 12% of these genes are PSGs, an  $\alpha$  selection value consistent with average human genome estimates (10-15%). Using branch-site tests, we showed that the primate tree is heterogeneously affected by positive selection, with the black snub-nosed monkey, the bushbaby, and the orangutan, being the most impacted branches. A large proportion of these genes is inclined to shape hair cells and stereocilia, which are involved in the mechanotransduction process, known to influence frequency perception. Adaptive selection, and more specifically recurrent adaptive evolution, could have acted in parallel on a set of genes (ADGRV1, USH2A, PCDH15, PTPRQ and ATP8A2) involved in stereocilia growth and the whole complex of bundle links connecting them, in species across different habitats, including high altitude and nocturnal environments.

## 2. Keywords

Inner ear expressed genes, Positive selection, Branch-site test, Stereocilia, Primates, Hearing

### 3. Significance statement

Positive selection signals, known to fix mutations beneficially and shaping certain physiological traits, are revealed by comparing genome sequences of related species. Here, we investigate how positive selection may have shaped auditory genes, and focused on inner ear expressed genes of primates. We found recurrent and episodic signals of positive selection in the primate tree, and observed a concentration of signals in species living at high altitude and in nocturnal species. Physically, these signals manifest themselves in the alteration of genes expressed in hair cells and stereocilia, which are involved in frequency perception.

### 4. Introduction

Audition constitutes a key gate between organisms and their acoustic environments. Hearing sensitivity varies between placental mammal species, influenced by a species' body size, middle and inner ear features, and continuous adaptation to their acoustic biotope (Vater et al., 2004; Bernardi et al., 2019; Köppl and Manley, 2018). Primates diverged from the rest of the mammals around 67.8 mya (Springer et al., 2012) and colonized very contrasting acoustic biotopes, from rainforest to savanna. Their adaptation to these new biotopes and associated sound propagation modalities (Brown and Waser, 2017) has likely had repercussions on their hearing sensitivity, optimizing sensitivity for their lifestyles (including circadian rhythms, i.e., nocturnal, diurnal or cathemeral) and for inter-individual communication (Ramsier and Rauschecker, 2017), in a process similar to what has been reported in subterranean mammal species (Heffner, 2004; Charlton et al., 2019). Therefore, the acoustic properties of a habitat may have acted as a direct source of selection (Brown and Waser, 2017), similarly to how primate calls and the corresponding hearing frequency sensitivity are driven by natural selection (Ramsier et al., 2012). Although the precise morphology of the ear varies across primate species (Coleman and Colbert, 2010), all therian mammal ears have the same three-

1 part structure: an external ear, a middle ear, and a coiled inner ear (cochlea). The external ear  
2 modulates sound perception (useful for localizing prey or congener) (Coleman and Ross, 2004;  
3 Dominy et al., 2004; Fleagle, 2013), while the middle and inner ear are known to influence  
4 perception on a spectral basis, weighing signals according to their frequential components. In  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
therian mammals, the middle and inner ear are thought to have co-evolved to improve high-  
frequency sensitivity (Coleman and Boyer, 2012; Köppl and Manley, 2018; Manley, 2017).  
The ancestors of extant primates are believed to have been nocturnal, small-sized animals with  
low inter-aural distances and short cochlea, features associated with high-frequency sensitivity  
and similar to those found in small mammals and rodents (Cao et al., 2015; Charlton et al.,  
2019; Heffner, 2004; Nummela, 2017). Nowadays, Strepsirrhini and tarsiers exhibit the same  
characteristics and are also more sensitive to high frequencies (Bernardi et al., 2019).  
Following the trend, New World monkeys, Platyrrhini, discriminate high frequencies better than  
their Catarrhini relatives (Old World monkeys and Apes) whose larger external ears and  
cochlea are more sensitive to lower frequencies (Coleman and Ross, 2004; Heffner, 2004;  
Nummela, 2017). Besides morphological features, the length of the outer hair cells, housed  
within the cochlea in the organ of Corti, is also linked to hearing sensitivity (Fettiplace, 2017).  
The length varies not only along the cochlea, but also between species. Species more sensitive  
to higher frequencies typically exhibit relatively shorter cells and stereocilia, whereas species  
displaying a lower frequency sensitivity have longer cells (Fettiplace, 2017). These differences  
in cellular morphology may be evolutionarily selected to adapt hearing to specific frequency  
ranges and improve communication. Moreover, the somatic electromotility of the outer hair  
cells plays a fundamental role in cochlear amplification (Köppl and Manley, 2018), and genes  
implicated in this function (e.g., *SLC26A5*, a gene encoding for prestin) are found to be under  
positive selection in some mammalian lineages (Franchini and Elgoyhen, 2006; Pisciotto et  
al., 2019). Due to the diversity in frequency sensitivity observed in the order Primates, we  
hypothesized that positive selection may have shaped genes expressed in the inner ear. Across

1 the primate tree, and depending on the approach used and the number of species considered,  
2  
3 3% (Van Der Lee et al., 2017) to 13% (Yin et al., 2020) of protein-coding genes were found to  
4  
5 present sites under positive selection. They were mainly enriched in proteins of immunity,  
6  
7 olfaction, and gustatory and auditory perception (Van Der Lee et al., 2017; Yin et al., 2020).  
8  
9 Focusing on the hearing system, evidence of molecular adaptation impacting three genes of the  
10  
11 auditory genome (OTOG,TECTA, MYO15) was found in a cross-species comparison focusing  
12  
13 on "deafness" genes in 69 mammals, including echolocating species such as bats and cetaceans  
14  
15 (Kirwan et al., 2013). However, although natural selection and convergent evolution have been  
16  
17 examined in echolocating species (Li et al., 2008; Parker et al., 2013), no specific research has  
18  
19 been conducted on selective adaptation in the evolution of primate auditory genes. In this paper,  
20  
21 we focused on 123 genes highly expressed in the inner ear and 27 primate species to obtain a  
22  
23 deep evolutionary insight into hearing adaptation. Using site test analyzes, we assessed whether  
24  
25 positive selection has influenced the auditory genes, and more specifically, the fraction  
26  
27 expressed in the inner ear where signal amplification occurs. Using branch-site tests, we  
28  
29 investigated which branches of the primate tree were the most susceptible to selection  
30  
31 regarding this set of genes. We chose a conservative approach, and focused on the potential  
32  
33 confounding factors affecting PSG signals (namely alignment errors, multiple comparisons  
34  
35 tests, genomic duplications, and gBGC). We also investigated and discussed the impact of  
36  
37 heterogeneous sequencing quality, and demography when data availability allowed. Our results  
38  
39 show that adaptive selection acts on inner ear expressed genes, and more specifically on genes  
40  
41 coding for stereocilia structures inside hair bundles.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

## 52 **5. Results**

53  
54 We analyzed 123 genes expressed in the inner ear of 27 primate species using both site tests,  
55  
56 determining selection signals at the base level, and branch-site tests, determining selection  
57  
58 signals at the phylogenetic branch level. The five steps of analyzes and results, including  
59  
60

1 evolutionary analyzes and filtering for confounding factors are summarized in Figure 1.  
2  
3  
4  
5

### 6 **5.1 Phylogenetic analyzes: the 123 auditory genes super-tree and $\omega$ values**


7

8 The phylogenetic super-tree obtained from the concatenated 319,074 bp of the 123 selected  
9 genes is similar to the well-established primate phylogeny (Supplementary Figure 34)  
10 (Perelman et al., 2011; Springer et al., 2012). The number of nucleotide substitutions per codon  
11 ranges from 0.01 to 0.18 across all branches analyzed, in accordance with the whole-genome  
12 estimate based on nine primate species (Van Der Lee et al., 2017).  $\omega$  values were calculated  
13 for each gene and the median values fluctuate from 0.05 to 0.1 (Supplementary Table 14) in  
14 accordance with Gibbs et al. (2007). The overall  $\omega$  for the whole tree is 0.157 and coherent  
15 with purifying selection being the main driving force acting on both primate and mammal  
16 genomes as a whole (Gibbs et al., 2007; Kosiol et al., 2008; Mikkelsen et al., 2005; Scally et  
17 al., 2012; Van Der Lee et al., 2017).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **5.2 Site test analyzes and global estimate of inner ear expressed PSGs in the primate tree**

34

35 Signals of positive selection were detected on 52/123 genes using site test analyzes, including  
36 both the genes rejecting the M1a and M7 null models, as well as those rejecting only the M7  
37 null model (Figure 1). After correcting for alignments and multiple tests, 41 PSGs remained  
38 (33% of the initial set), including 15 genes rejecting both the M1a and M7 models, and 26  
39 genes rejecting only the M7 model (Supplementary Table 6). Among the 41 PSGs detected in  
40 the site test analysis, 30 were positive to Bayes Empirical Bayes (Supplementary Table 6). In  
41 total, 68 out of 319,074 sites tested were found to be under positive selection (0.021%).  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**Figure 1.** Overview of our comparative evolutionary analysis procedure for conservative inference of positively selected genes and branches. 1. Selection of 123 genes from the EARxpressed repository; 2. Multiple sequence alignments and maximum likelihood-based phylogenetic analyzes; 3. Site tests (PAML) and confounding factor analyzes (alignment and multiple test corrections); 4. Branch-site tests (PAML) and confounding factor analyzes (alignment and multiple test corrections); 5. Additional analyzes of complexifying factors (duplications and gBGC).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 5.3 Branch-site test analyzes: are specific branches affected by episodic positive selection?

The branch-site test determined which branches of the primate tree were affected by episodic or continuous positive selection. A total of 72 PSGs were detected (Figure 1, Supplementary Table 15). After correction for alignments and multiple comparison issues, 42 PSGs remained (Supplementary Table 16). After a final visual inspection, performed using the Bioedit tool (Hall, 1999), 31 PSGs were considered as being true signals (*i.e.*, 25% of the initial gene set, protein alignment in Supplementary Figure 1-31) and were placed on the cladogram of primate species (Figure 2). These 31 PSGs represent 45 selective signatures distributed on 12 distinct internal branches and 33 distinct terminal branches. From the 6,383 branches analyzed, 45 showed evidence of positive selection, resulting in a total of 0.70% impacted branches. These 31 validated PSGs are heterogeneously distributed in the whole tree, and are detected preferentially in the black snub-nosed monkey (6 PSGs), the bushbaby (5 PSGs) and the orangutan (5 PSGs). Some sub-families and families such as Hominoidea, Hylobatidae,



**Figure 2:** Cladogram of primate species displaying the 31 validated PSGs of branch-site test analyzes after multiple tests plus alignment corrections and a final visual inspection. PSGs potentially impacted by duplications and gBGC are mentioned in italics. Branch lengths are unscaled.

and Cercopithecoidea are almost unaffected (Figure 2). We analyzed the detailed functional

1 enrichment for these 31 genes and found they were mainly annotated with GO terms related  
2 to stereocilium organization and neuron projection, suggesting structural changes in hair cells  
3 and more specifically in stereocilium structure. Branch-site test analyzes identified the  
4 branches displaying the strongest signals, and also the occurrence and temporal scheme of  
5 these selective signals. We defined three types of PSGs, the "One-time PSG", the "Recurrent-  
6 PSG" and the "Continuous-PSG". Out of the 31 validated PSGs, most were detected only once  
7 in the whole primate tree and defined as "One-time PSG" (20/31 PSGs: 65% of all PSG  
8 detected) suggesting either the punctual nature of this signal or the lack of genomic  
9 information in adjacent branches to make them detectable on deeper branches. Five  
10 "Recurrent-PSGs" were detected on the whole tree: ADGRV1 (Baboon, bushbaby, black snub-  
11 nosed monkey), USH2A (Bushbaby, orangutan), PCDH15 (Black lemur and tarsier), PTPRQ  
12 (Black lemur and orangutan) and ATP8A2 (Black snub-nosed monkey and proboscis  
13 monkey). These five genes code for membrane proteins constitutive of the hair bundles, and  
14 more specifically of ankle links (ADGRV1, USH2A), the tip and kinocilial links (PCDH15)  
15 and shaft connectors (PTPRQ). Nucleotide changes observed affecting the same gene in  
16 several lineages might suggest recurrent pressures on this specific cellular component. In the  
17 case of ADGRV1 and PTPRQ, the amino acid changes are predicted to have a functional  
18 impact (Polyphen2, Supplementary Table 7). They are located in the extracellular domain of  
19 the proteins involved in the links. For PCDH15, the site under selection is located in a highly  
20 variable intracellular domain, although it is predicted to be potentially damaging. Finally, two  
21 genes STOX1 and KCNQ4 were identified as "Continuous PSGs" in the Papionini/baboon  
22 respectively, snub-nosed monkey (genus *Rhinopithecus*)/black snub-nosed monkey. The gene  
23 STOX1 is involved in regulating the proliferation of inner ear epithelial cells (Chen et al.,  
24 2019) and KCNQ4 is a voltage-gated potassium channel that plays an essential role in  
25 maintaining ion homeostasis and regulating hair cell membrane potential. Interestingly, two  
26 of the amino acid changes in KCNQ4 correspond to human polymorphisms, and could suggest  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 potential balancing selection acting in humans (Supplementary Table 7). To complete the  
2  
3 branch-site analyzes, we explored the potential impact of the duplications and the gBGC on  
4  
5 the remaining 31 PSGs. Out of these 31 PSGs, 27 PSGs correspond to 33 external branches  
6  
7 that could be investigated. Using RPHAST, we showed that biased gene conversion (gBGC)  
8  
9 could have affected four genes (ADGRV1, COL10A1, USH2A and KCNQ4) out of the 27  
10  
11 PSGs, specifically in the bushbaby and the black snub-nosed monkey (Supplementary Table  
12  
13 S10). However, the probability of detecting gBGC is systematically associated with the same  
14  
15 probability of detecting either positive selection (e.g., KCNQ4 for the black snub-nosed  
16  
17 monkey and COL10A1 for the bushbaby) or relaxation of purifying selection (e.g., ADGRV1  
18  
19 and USH2A for the bushbaby). None of the signals detected display a unique probability of  
20  
21 detecting gBGC. In total, 25 PSGs could still be considered as true PSGs after filtering for  
22  
23 gBGC (Supplementary Table 10). We explored the potential impact of duplication by  
24  
25 analyzing only the PSGs associated with identified BEB sites: 23 PSGs, corresponding to 65  
26  
27 identified BEB sites in external branches were tested. Due to partially available genomic  
28  
29 information for some species, the analyzes were restricted to 12 PSGs out of these 23, further  
30  
31 limiting our conclusions (Figure 2, Supplementary Table 8). BEB and SD intersections were  
32  
33 tested for 12 PSGs, representing 12 external branches, 17 exons, and an associated 37 BEB  
34  
35 sites. Most of the PSGs were considered as true PSGs, and only one duplicated exon  
36  
37 intersected with a BEB in ADGRV1 in the black snub-nosed monkey and was considered as  
38  
39 a potentially false PSG (Supplementary Table 8; Supplementary Table 9). Although the  
40  
41 differences in genome quality, annotations, and SD detection between species could limit our  
42  
43 current investigation, at this stage, neither duplication nor gBGC could be considered as a  
44  
45 strong confounding factor. More genomic information would be required for these tests to be  
46  
47 conclusive.  
48  
49  
50  
51  
52  
53  
54  
55

56 **5.4 Site and branch-site tests: shared-PSG signatures between both approaches** We  
57  
58 used site and branch-site tests in parallel to detect: 1) evidence of adaptive selection on the  
59  
60

1 whole primate phylogeny using site tests; and 2) the most impacted PSG/branch couples using  
2 branch-site tests. A total of 35 shared PSGs were revealed by both approaches before filtering  
3 for confounding factors which resulted in a remaining 13 PSGs (Venn diagram in  
4 Supplementary Figure 35). Five of these 13 PSGs have been already reported in the literature  
5 as PSGs or accelerated genes involved in auditory function in Homininae (Humans,  
6 Chimpanzees, Gorillas) and others mammals (ADGRV1) (Pisciottano et al., 2019; Scally et  
7 al., 2012), CLIC5 (Cao et al., 2015), MYO15A (Kirwan et al., 2013), RPGRIP1L (Van Der  
8 Lee et al., 2017), and SRRM4 (Lindblad-Toh et al., 2011). Among the filtered PSGs, CDH23,  
9 TMC1, and OTOF have also been reported in the literature (Scally et al., 2012) and could have  
10 been over-filtered using our procedure. Interestingly, the genes detected in the Homininae  
11 branch were not detected before or after filtering for confounding factors; this might be  
12 explained by our use of more recent versions of the concerned genomes compared to Scally et  
13 al. (2012), removing potentially ambiguous sites or due to the higher number of primate  
14 species analysed here allowing a more accurate detection of adaptive selection (McBee et al.,  
15 2015). New PSGs have been revealed by our study including AADACL2, CLCA2, WFS1,  
16 COL10A1, LMO7, SLC52A3, SRRM4, and STOX1. Two of these genes are involved in cell  
17 adhesion: CLCA2 a calcium activated chloride channel regulator, and LMO7 described to  
18 organize the actin network in the cuticular plate and cell junctions (Du et al., 2019). The  
19 splicing (SRRM4) and transcription (STOX1) factors are involved in gene regulation. WFS1  
20 is a reticulum endoplasmic cation channel playing a role in  $Ca^{2+}$  homeostasis, and is  
21 responsible for the hearing disorder Wolfram Syndrome Type I. In the case of the deacetylase  
22 AADACL2 and the collagen COL10A1, several sites are found under selection and these  
23 nucleotide sites are polymorphic in humans and might suggest that balancing selection could  
24 be active at the intra-specific level in humans (Supplementary Table 7).

## 5.5 Impact of sequencing quality on PSG detection

We tested the impact of heterogeneous genome quality on PSG detection by comparing a set of metrics with the number of PSGs detected. To be objective and in the absence of NG50 and Q-score metrics for all primates, we chose to run the QUASt-LG tool on raw data to generate the quality metrics (Supplementary Table 12) (Kuderna et al., 2020). The relative position of the analyzed genome was assessed using these metrics and a PCA; no significant association between PSG number and genome quality was observed: PCA coordinates were not detected for axis 1 (Kruskal-Wallis test, chi-squared = 5.15, p-value < 0.08) nor axis 2 (Kruskal-Wallis test, chi-squared = 1.5 p-value < 0.47). We concluded that the number of PSGs detected is not associated to the relative quality of the genome analyzed.

## 5.6 Impact of demographic history on detected PSGs

The retention of the adaptive allele and the efficacy of selection depend on population genetics and the demographic history of each allele; thus, the large differences in effective population sizes of the primate species analyzed (from very large in owl monkeys (Schrago and Seuánez, 2019) to relatively low for the *Rhinopithecus* genus (Zhou et al., 2014) might have impacted the heterogeneous fixation of the adaptive alleles. We used the IUCN status as a proxy for demographic history, and found no relation between IUCN status and the number of PSGs detected (Kruskal-Wallis chi-squared = 0.88068, df = 4, p-value = 0.9273). Although we cannot rule out that IUCN status is a poor proxy for current census size and effective population size, IUCN status does provide an approximate index of recent population demography. Further investigations on species-specific demography should be planned, especially on long-term effective population size which should be a more valid indicator compared to current census size or effective population size (Cagan et al., 2016).

## 6. Discussion

In this paper, we comprehensively and conservatively analyzed a chosen set of 123 inner ear expressed genes in 27 primate species. We performed both site and branch-site test analyzes, aiming to establish selection patterns both at the gene and at the clade/lineage levels. We found signals of selective pressure, and, building upon the expression characteristics and the environment of the concerned species, we hypothesize some macroscopical factors from which this selective pressure may stem.

### 6.1 Questioning auditory gene selection and the impact of confounding factors

**Auditory gene selection:** We chose to analyze a subset of genes highly expressed in the inner ear, gathered in the EARxpress repository to include the maximum number of primate species and to give us the opportunity to explore alignment by visual inspection due to the highly heterogeneous sequencing qualities. By focusing on genes displaying a strong expression pattern in the inner ear, but also known to be transcribed and associated with "hearing" Gene Ontology term (GO), we aimed to target key auditory genes, without any *a priori* in regard to the literature. We confirmed the impact of adaptive selection on a few auditory genes previously revealed in Hominae and other mammals. We highlighted new genes and new species to consider in relation with their functions and ecology, respectively. Further analyzes on a much larger set of genes expressed in hearing would help to progress further in identifying the key gene network involved in auditory function for different species and their associated habitat and social structure.

**Impact of alignment and multiple test corrections:** We estimated the relative impact of the main sources of ambiguities stemming from the large number of species analyzed. For both site and branch-site tests, correcting for multiple tests was the most severe filtering step,

1 removing 8% and 22% of the initially captured PSGs, respectively (Figure 1). In contrast,  
2  
3 correcting for alignment ambiguities using automatic filtering methods had a marginal impact  
4  
5 on the final list of the selected PSGs, impacting no more than 1% of the initial candidates.  
6  
7 However, the final visual inspection had a stronger impact, filtering out more than 10% of the  
8  
9 remaining PSGs in branch-site test analyzes. Similar filters have been used in previous studies  
10  
11 (Daub et al., 2017; Van Der Lee et al., 2017), in which alignment ambiguities due to incorrect  
12  
13 sequences, frameshift or alternative exons are often mentioned as the most problematic sources  
14  
15 of false PSGs (Kosiol et al., 2008; Mallick et al., 2009; Markova-Raina and Petrov, 2011;  
16  
17 Wong et al., 2008). If we had considered only automatic filtering methods (HMMCleaner and  
18  
19 GUIDANCE), without considering visual inspection, this would have had much less of an  
20  
21 impact on our results, as has been noted in the literature (Fletcher and Yang, 2010). The drastic  
22  
23 effect of this step attests the need for better automated methods to highlight alignment failures  
24  
25 or additional control procedures.  
26  
27  
28  
29  
30  
31  
32

33 **Exploring the impact of gBGC and segmental duplications on PSG signals:** We tested the  
34  
35 impact of gBGC and SD on PSG signals. gBGC is known to affect around 10% of the primate  
36  
37 genomes (Galtier et al., 2009); however, it was not detected as acting independently in our  
38  
39 study, but rather as always coinciding with positive selection, as also noticed by Van Der Lee  
40  
41 et al. (2017). Similarly, SDs as a proxy for duplication processes modestly impact our  
42  
43 estimates, as also shown in Daub et al. (2017). However, the large fraction of unplaced  
44  
45 scaffolds in many genomes, limits the output of realistic SDs maps and the conclusions that we  
46  
47 can draw from it (from 0.37% to 100%, in base pairs, of non-chromosomal top-level scaffolds,  
48  
49 Supplementary Table 9). We found evidence of duplication for only one exon containing  
50  
51 positively selected sites ADGRV1 in the black snub-nosed monkey (Supplementary Table 8).  
52  
53 This result indicates either the low incidence of SDs in the set of analyzed genes, or the large  
54  
55 underestimation of duplications in absence of reliable structural variation maps for the genomes  
56  
57  
58  
59  
60




1 of the primates analyzed in our study. More investigations are needed to determine whether  
2 evidence of duplication for ADGRV1 in the black snub-nosed monkey could have potentially  
3 biased our results by considering a very close paralogous gene, or whether this gene could be  
4 Copy Number Variant (CNV) and should be considered as a driving force for this gene. The  
5 significant relationship between duplications, CNV and adaptive traits (Prunier et al., 2017)  
6 and especially of the key role played by CNV in climatic and environmental adaptation should  
7 be kept in mind for future studies focusing on auditory genes, especially for species living in  
8 extreme environments, such as the black snub-nosed monkey. GC-biased gene conversion, also  
9 known to mimic positive selection through the biased fixation of GC sites has been investigated  
10 previously (Lartillot, 2013; Ratnakumar et al., 2010; Romiguier and Roux, 2017). In our study,  
11 none of the PSGs detected display an unambiguous gBGC signal, but are rather always, when  
12 present, equiprobable with a positive selection signal and could not help us to conclude on any  
13 potential impact of this force here. Collecting both high-quality assembled genomes and  
14 associated genome annotation, for the low-profile primate genomes studied here, and more  
15 specifically in Strepsirrhini, would contribute to better discussions on the impact of duplication  
16 and gene conversion on PSG estimates.

## 6.2 The impact of adaptive selection on auditory genes

45 We found that 12% of the 123 genes analyzed show evidence of positive selection when  
46 considering the M1a-M7 models, and 21.1% when considering the M7 model alone. This  
47 estimate is four times higher than genome-wide PSGs estimates based on nine primate species  
48 (Van Der Lee et al., 2017), and nearly identical to the genome-wide estimate based on 12  
49 primate species (Yin et al., 2020). This result stresses the importance of the number of species  
50 considered, and the procedure used to treat confounding factors (Arbiza et al., 2006). We chose  
51 a conservative approach, favoring to err in the direction of under-detection. However, the  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 proportion of PSGs found in this study appears quite high despite neurological and auditory  
2 genes being one of the most enriched sensory families with 8.5% of genome-wide PSG  
3 estimates in mammals, compared to chemical perception (6%), smell (3.75%), or taste (2%)  
4 (Kosiol et al., 2008). The impact of positive selection on the auditory genes in primates could  
5 reflect the adaptation of the inner ear to the spectra of perceived frequencies encountered in the  
6 ecology of each species (Brown and Waser, 2017; Ramsier et al., 2012; Ramsier and  
7 Rauschecker, 2017) that ranges from from 28 Hz to 65 kHz. Hearing sensitivity in these clades  
8 has evolved in parallel with key changes in morphology and cellular structure of the inner ear  
9 (Köppl and Manley, 2018; Vater et al., 2004), detectable by selective signatures on the genomic  
10 fraction involved in inner ear functioning (Kosiol et al., 2008; Pisciotano et al., 2019). This  
11 evolution could be linked to the sound propagation in their respective biotope and species-  
12 specific needs in term of acoustic communication (Waser and Brown, 1986). Five genes  
13 (ADGRV1, USH2A, PCDH15, PTPRQ and ATP8A2), predicted to be functionally associated  
14 (Supplementary Figure 36), showed evidence of adaptive selection in several lineages and  
15 could reflect convergent adaptive evolution acting on the whole complex of links connecting  
16 the stereocilia (Figure 3 and 4). Most of the PSGs show species-specific patterns (20/31 One-  
17 time PSG) and might reflect the need to adapt differently at the genetic level, whatever the  
18 environment, a result that has already been detected and discussed in primates (Cagan et al.,  
19 2016). Alternatively, these regions may represent cases of convergent evolution, where the  
20 same phenotype is positively selected across species (Cagan et al., 2016). Specific targets in  
21 the inner ear are discussed later in this paper.



**Figure 3:** Examples of the position of selected PSG amino acid (AA) sites (in black square) in the protein sequences of the four genes: ADGRV1, KCNQ4, PCDH15 and PTPRQ.


### 6.3 PSG realms: hair cells, stereocilia and mechanotransduction

The mechanotransduction process occurs in hair cells and stereocilia structures; atmospheric waves are transformed into a nervous message that is then sent to the brain. These cells and structures are also involved in signal amplification and modification, and alterations of these cellular components could explain the different modalities of hearing perception (frequencies and amplitude) between primate families (Coleman and Ross, 2004; Fettiplace, 2017). Each hair cell carries hundreds of stereocilia, organized either in three rows at their apical pole (OHC scheme), or in a staircase pattern with multiple rows (IHC scheme), connected to their neighbors by extracellular filaments, hair bundle links (Fettiplace, 2017; Richardson and Petit, 2019). The development of hair bundles is characterized by transitory links such as ankle links, shaft connectors, horizontal top connectors, tip links, and kinocilial links (Richardson and Petit, 2019). All these types of links are required to maintain the structure, function, and normal

1 development of hair bundles (Richardson and Petit, 2019). The latter consist of actin-rich  
2 stereocilia and a microtubule-based kinocilium. The PSGs protocadherin 15 (PCDH15) and  
3  
4 cadherin 23 (CDH23) form the kinociliary link complex between the kinocilium and the  
5  
6 longest stereocilia, as well as the tip links that connect stereocilia (Fettiplace, 2017; Richardson  
7  
8 and Petit, 2019) (Figure 4). The ADGRV1 (also called VLGR1 or GPR98) and USH2A  
9  
10 compounds are found at the base of the stereocilia, where they are thought to form ankle links  
11  
12 (Richardson and Petit, 2019). Myosins apply tension on the actin filaments and are also  
13  
14 implicated in the stereocilia length (Michalski and Petit, 2015). Myosin 6 (MYO6) is highly  
15  
16 concentrated in the cuticular plate at the apical hair cell surface, but is also localized in  
17  
18 stereocilia. MYO7A is expressed throughout stereocilia and is notably present in ankle links in  
19  
20 some auditory epithelial. Most of the PSGs detected correspond to key proteins involved in  
21  
22 mechanotransduction (Fettiplace, 2017) and hair bundle structure, and the recurrent PSGs  
23  
24 suggest recurrent changes affecting these genes in multiple lineages. Among the stereocilia  
25  
26 links, the component of the shaft connectors, PTPRQ, is also positively selected and plays an  
27  
28 important role in the maturation of hair bundles (Richardson and Petit, 2019). In our study,  
29  
30 most of the components of these links are positively selected, showing that the development of  
31  
32 the stereocilia and their cohesion could have undergone a specific adaptation in some lineages.  
33  
34 The selection signatures of this type of link component is not specific to a given primate hearing  
35  
36 range, but is found multiple times in their phylogenetic tree, which points to the recurrent  
37  
38 adaptation of this set of genes to external constraints. Further investigations on the relation  
39  
40 between transient links, actin-binding proteins, and stereocilia length will be of importance to  
41  
42 link observed hearing sensitivity differences to molecular and cellular characteristics.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

#### 54 **6.4 Adaptation of hair cells and hair bundles to extrinsic or intrinsic parameters**

55  
56  
57  
58  
59  
60



**Figure 4:** (A) Cochlear duct and its cellular structure including hair cell stereociliary bundles; (B) details of the stereocilium and hair bundles, and the structure and location of link types including kinocilial links, tip links, horizontal top connectors, shaft connectors, and ankle links (b); (C) Zoom on the tip link connecting two stereocilium - adapted from Fettiplace (2017) and Richardson and Petit (2019).

1 primate phylogenetic tree. Going further in interpreting such heterogeneous distribution in  
2  
3 regard to the ecological parameters would require accessing the whole genome auditory-linked  
4  
5 PSGs for populations and species living in distinctively identified ecological niches.  
6  
7 Nevertheless, it worth noticing that the most affected species is the black snub-nosed monkey  
8  
9 which lives at the highest altitude of any known non-human primate. High altitude is  
10  
11 responsible for various physiological adaptations in many organisms (Ai et al., 2014; Beall,  
12  
13 2006; Qiu et al., 2012); however, few studies have supported the strong effects of high altitude  
14  
15 on the inner ear (Cingi et al., 2010; Fan et al., 2016). The exception is the domestic Yak: this  
16  
17 species lives at a similar altitude range as the black snub-nosed monkey and presents an  
18  
19 accelerated rate of evolution for its genes involved in sound perception (Qiu et al., 2012). If  
20  
21 altitude is a key factor modeling auditory genes, we should notice that the black snub-nosed  
22  
23 monkey and its closest relative, the golden snub-nosed monkey, displaying a totally different  
24  
25 PSG pattern, have divergent phylogenetic histories and are now living in different habitats. The  
26  
27 black snub-nosed monkey lives at higher altitudes (3,400 to 4,600 meters) and faces  
28  
29 environmental challenges such as low temperature and oxygen levels (Zhou et al., 2016), while  
30  
31 the golden snub-nosed monkey lives at more moderate elevations (1,500 to 3,400 m). To  
32  
33 determine whether PSGs detected in the black snub-nosed monkey reflect a process of  
34  
35 physiological adaptation and modulation in frequency or amplitude perception abilities, and  
36  
37 the potential existence of an altitude threshold, auditory brainstem responses or other methods  
38  
39 to estimate the hearing sensitivity spectrum would have to be tested and compared on several  
40  
41 individuals for these two species. The second most affected species is the bushbaby, a nocturnal  
42  
43 species that exhibits a higher sensitivity to low frequencies (around 250 Hz) compared with  
44  
45 diurnal species (Bernardi et al., 2019; Masters, 1991). Specific constraints associated with a  
46  
47 nocturnal life might have modulated the molecular network responsible for  
48  
49 mechanotransduction explaining the enrichment in PSGs for this species. In addition,  
50  
51 specificity between species could explain the different sets of PSGs detected. Different  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 biochemical or regulatory networks could be involved; the connectivity and centrality of the  
2 genes in these networks might differ between networks, thus modulating the effect size of the  
3 genes (Olson-Manning et al., 2012). This could be especially true for genes included in more  
4 than one pathway and those influencing multiple phenotypes, such as those co-involved in  
5 vision detection for these species, that may experience pleiotropic effects. Some PSGs that  
6 form hair bundle links are also implicated in vision; for example, CLIC5, PCDH15, USH2A,  
7 and ADGRV1 are all involved in Usher syndrome, congenital deafness due to stereocilia  
8 disorganization with associated progressive retinitis pigmentosa (Fettiplace, 2017). This might  
9 suggest specific hearing sensitivity characteristics associated to nocturnal life, together with a  
10 co-adaptation of vision-linked expressed genes to optimize light perception. Together with  
11 ADGRV1, USH2A, RPGRIP1L, CLIC5 and WFS1 are implicated in both hearing and visual  
12 perception, suggesting that these genes had a continuous and complex evolution pattern during  
13 primate evolution, with various adaptation phases when species converted from nocturnal to  
14 diurnal activities. Supporting this hypothesis would require more investigations on both  
15 physiological and genomic components (Nummela, 2017).

## 37 **7. Material and Methods**

### 38 **7.1 Genes expressed in the inner ear: EARxpress, a dedicated database**

39 In the absence of a comprehensive repository of all genes involved in auditory perception and,  
40 more specifically, inner ear expression, we gathered all publicly available information on these  
41 genes, in primates and the mouse. Information on auditory genes were retrieved from four  
42 papers focusing on the genes expressed in the inner ear of *Macaca fascicularis* (Mutai et al.,  
43 2018), and on the transcriptomic (Liu et al., 2014), developmental (Scheffer et al., 2015), and  
44 proteomic expression of inner ear genes in the mouse (Hickox et al., 2017). We gathered 2367  
45 genes and consolidated them in our EARxpress database (Supplementary Table 1). We  
46 retrieved orthologous loci in the human genome using Biomart when possible, and Blast when  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 no orthologous sequences could be detected. Genes were considered orthologous when their  
2  
3 sequence identity exceeded 80% and the e-value of the Blast matches remained below 0.05  
4  
5 (Kosiol et al., 2008). To explore the potential role played by adaptive selection on genes  
6  
7 expressed in the inner ear in the order Primates, we chose to focus our attention on a restricted  
8  
9 number of genes that would have a greater probability of playing a role in the functioning of  
10  
11 the inner ear and to include the maximum number of primate species. We prioritized genes  
12  
13 over-expressed in the inner ear with regard to other tissues and selected first the 32 genes  
14  
15 expressed in *M. fascicularis* (Mutai et al., 2018). Then, we selected 95 additional genes  
16  
17 common to at least three or two of the remaining studies (Liu et al., 2014; Hickox et al., 2017;  
18  
19 Scheffer et al., 2015), or associated with a GO term related to "hearing". This selection was  
20  
21 made without any *a priori* regarding the PSGs detected in the literature or their specific  
22  
23 functions in the inner ear. Out of these 127 pre-selected genes, three genes (COCH, MAF,  
24  
25 SLC17A8) were common between the two data sources and were considered only once. The  
26  
27 DEFB122 was excluded from the following analyzes because it is a pseudogene in humans and  
28  
29 it could not be used as a functional gene reference to retrieve functional orthologous genes in  
30  
31 other non-human primate species. In this study, we considered 123 genes in the phylogenetic  
32  
33 tree construction and selective pressures analyzes (Supplementary Table 2), following the  
34  
35 pipeline detailed in Figure 1.  
36  
37  
38  
39  
40  
41  
42  
43  
44

## 45 **7.2 Orthologous identification, alignment quality and phylogenetic tree** 46 47 **reconstruction**

48  
49 The orthologous coding DNA sequences (CDS) of all available non-human primate  
50  
51 species were retrieved from the UCSC Genome Browser (Karolchik et al., 2004) using the  
52  
53 Multiz tool. In total, 27 available species were retrieved: *Homo sapiens* (Human), *Colobus*  
54  
55 *angolensis palliates* (Angolan colobus), *Papio anubis* (Olive baboon), *Eulemur macaco* (Black  
56  
57 lemur), *Rhinopithecus bieti* (Black snub-nosed monkey), *Pan paniscus* (Bonobo), *Otolemur*  
58  
59  
60



1 *garnetti* (Northern greater bushbaby), *Pan troglodytes* (Chimpanzee), *Propithecus coquereli*  
2 (Coquerel's sifaka), *Macaca fascicularis* (Crab-eating macaque), *Mandrillus leucophaeus*  
3 (Drill), *Nomascus leucogenys* (Northern white-cheeked gibbon), *Rhinopithecus roxellana*  
4 (Golden snub-nosed monkey), *Gorilla gorilla gorilla* (Western lowland gorilla), *Chlorocebus*  
5 *sabaeus* (Green monkey), *Aotus nancymaae* (Ma's night monkey), *Callithrix jacchus* (Common  
6 marmoset), *Microcebus murinus* (Mouse lemur), *Pongo pygmaeus abelii* (Sumatran  
7 orangutan), *Macaca nemestrina* (Pig-tailed macaque), *Nasalis larvatus* (Proboscis monkey),  
8 *Macaca mulatta* (Rhesus macaque), *Eulemur flavivrons* (Sclater's lemur), *Cercocebus atys*  
9 (Sooty mangabey), *Saimiri boliviensis* (Black-capped squirrel monkey), *Tarsius syrichta*  
10 (Philippine tarsier), and *Cebus capucinus imitator* (White-faced sapajou) (Supplementary  
11 Table 3). We downloaded multiple sequence alignments using Multiz Align from the Table  
12 Browser according to the GENCODE v24 track. When multiple transcripts variants were  
13 available, we chose the variant featuring the longest open reading frame compared to the human  
14 sequence (hg38/GRCh38, Dec. 2013). Details on the human transcripts are provided in  
15 Supplementary Table 2; for seven genes and a few species, orthologous sequences were  
16 missing and the corresponding positions were filled with unspecified nucleotides (N)  
17 (Supplementary Table 4). The GUIDANCE2 server was used to generate a new realignment  
18 for each gene using MAFFT (Kato and Standley, 2013). To assess the robustness of the  
19 alignments, we ran 100 bootstrap tree iterations (Penn et al., 2010; Sela et al., 2015).  
20 GUIDANCE2 ranged from 0.98 to 1 for 121 genes, with two genes, LOR and NEFH, falling  
21 below at 0.92 and 0.94 respectively, which might suggest further misalignments regarding  
22 these two genes. Additionally, regardless of their GUIDANCE score, all alignments were  
23 filtered and cleaned with HMMCleaner, using the default parameters (Di Franco et al., 2019).  
24 Ambiguous nucleotide positions were masked and replaced automatically by HMMCleaner  
25 using gaps and were considered in the following analyzes. We controlled that aligned cDNA  
26 sequences corresponded to an ORF by the systematic control of a methionine and a stop codon  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(see Supplementary data in Figure 1-31); these sequences were also compared with those from the NCBI website to be certain that the ORF was present. The last filtering step consisted in a simple visual inspection of the protein sequences alignment by using BioEdit (URL: <http://www.mbio.ncsu.edu/BioEdit/>) in order to eliminate artefacts due to alternative transcription or unreliable alignment surrounding the positive selected residues (Van der lee 2017; Wong 2008). We built the phylogenetic super-tree from the concatenated-alignment obtained with the 123 genes using RAxML-NG v.0.6.0 (Kozlov et al., 2019), using the mouse as the outgroup (Supplementary Figure 32). To preserve the maximum number of positions in the concatenated tree, no specific filtering procedure was applied. We used the best-fit model of nucleotide substitution given by Akaike's information criteria corrected (AICc) in the JModelTest2 (Darriba et al., 2012) to build the concatenated alignment tree with the most adapted model of nucleotide substitution, and used the APE R package to unroot the tree (Paradis et al., 2004; R Core Team, 2013).

### 7.3 Evolutionary analyzes: site and branch-site tests

We used codeml from the PAML v4.9h software suite (Yang, 2007) to detect the signatures of positive selection. We used the codon frequency model F3X4 and the combination of the following codeml parameters: NSsites = 0 1 2 7 8, CodonFreq = 2, clock = 0, fix kappa = 0, kappa = 0.3, fix omega = 0, omega = 0.4, method = 0, cleandata = 0. We estimated the ratio of nonsynonymous/synonymous substitutions per site (dN/dS or  $\omega$ ) using maximum likelihood (ML). The dN/dS ratio was computed for each gene, using the M0 model with the gene alignment and the phylogenetic tree built from the concatenated alignment. The branch-site test used the codon frequency model F3X4, and for the alternative model (model A, positive selection), we used the following codeml parameters: model = 2, NSsites = 2, fix kappa = 0, kappa = 2, fix-omega = 0, omega = 1, method = 0, cleandata = 0. The null uses the same parameters as model A, except fix-omega was fixed to 1.

### 7.3.1 Site test analyzes

First, we searched for genes exhibiting sites under positive selection. Since all amino acid sites in a protein are not necessarily expected to be under the same selective pressure, we assumed heterogeneous classes of sites characterized by different  $\omega$  (Yang and Swanson, 2002). The random variation of  $\omega$  was accounted for by using different statistical distributions implemented by Nielsen and Yang (1998) and Yang et al. (2000). We used two tests known to be particularly effective: the first compares the null model M1a (neutral;  $\omega < 1$ ) and the alternative model M2a (selection;  $\omega > 1$ ), and the second compares the null model M7 (beta distribution for  $\omega < 1$ ) and the alternative model M8 (beta &  $\omega > 1$ ). In both cases, each of the two models was separately fitted to data and the log-likelihood was computed. Then the LRT (log-likelihood ratio statistic) was approximated by a chi-square ( $\chi^2$ ) distribution of the two models with their respective degrees of freedom. The p-values obtained for the two site tests were corrected using the Benjamini-Hochberg method with a criterion of  $FDR \leq 0.05$  (Benjamini and Hochberg, 1995). All parameters of the evolutionary models are detailed in Supplementary Table 5. After the maximum likelihood estimates (MLEs) of the model parameters were obtained, we used Bayes Empirical Bayes (BEB), selected for its ability to accommodate uncertainties in the MLEs of the parameters (Yang et al., 2005) to infer the most likely site classes for every site with  $\omega > 1$ . We considered genes that rejected null models M1a or M7 to be positively selected, and sites with probability values  $\geq 0.95$  in BEB analyzes as evolving under positive selection. GO terms enrichment and interaction of proteins were retrieved from the STRING database (Szklarczyk et al., 2019). PSGs revealed by site test analyzes are listed in Supplementary Table 6.

### 7.3.2 Branch-site test analyzes

Second, we aimed to detect which branches of the primate tree (Zhang et al., 2005) of

1 the selected auditory genes are or were subject to positive selection. We define the "foreground  
2 branch" as the branch being tested for selection and all other branches as being the "background  
3 branches". The null model allows  $\omega$  to vary among sites, but fixes a set of sites where  $\omega = 1$   
4 for the background and foreground branches. The alternative model lets  $\omega$  be greater than 1 in  
5 the foreground branch, but forces  $\omega = 1$  in the background branches for the same sites. We  
6 computed the log-likelihood for each branch three times, then took the highest log-likelihood  
7 for the null and alternative models, as recommended by Yang and Dos Reis (2010). We then  
8 computed the LRT and compared it with a chi-squared distribution with one degree of freedom.  
9 Since we tested a total of 52 branches for 123 genes, including 27 terminal branches and 25  
10 internal branches, we corrected the p-values obtained from the chi-squared test as  
11 recommended by Anisimova and Yang (2007) using the p.adjust function and the FDR (False  
12 Discovery Rate) model (Benjamini and Hochberg, 1995; R Core Team, 2013). The last filtering  
13 step consisted in a visual inspection of the protein sequence alignments to eliminate artifacts  
14 due to alternative transcription or unreliable alignments surrounding the positively selected  
15 residues (Van Der Lee et al., 2017); a site was filtered out if a gap was present within three  
16 amino acids around it (Zhou et al., 2014) or if the sequence alignment surrounding the position  
17 displayed a high sequence divergence compared to the other species. For these validated PSGs,  
18 we assessed the functional consequences of the amino acid changes using Polyphen-2, a  
19 software that uses machine-learning classification to estimate the probability of a SNP being  
20 damaging based on: 1) protein sequence annotation; 2) structure; and 3) an evolutionary  
21 conserved multi-species alignment (Adzhubei et al., 2013). We checked whether these sites  
22 were present in the functional domain of the corresponding protein. The PSGs revealed by  
23 branch-site test analyzes are listed in Supplementary Table 7. All validated PSGs were assigned  
24 to one of three categories: One-time PSG, a gene being observed positively selected in only  
25 one terminal branch; Recurrent PSG, a gene being determined to be positively selected in at  
26 least two independent branches; and Continuous PSG, a gene whose selective signal is detected  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
uninterruptedly in at least two external and internal connected branches, suggesting a selective  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
signal detectable over a longer time period.

### 7.3.3 Mitigating potential confounding factors on false detection

In addition to the previously detailed measures taken to reduce the impact of sequencing and assembly errors (MAFFT, GUIDANCE and HMMCleaner), we investigated the potential impact of large amplicon duplications (Segmental duplications), gBGC, heterogeneous sequencing quality and, demographic histories on PSG detection. Another potential source of false PSGs is Multi-Nucleotide Mutations (MNM), multiple and closely spaced substitutions in a sequence (Schridder, 2011). If MNM can be due to low sequencing quality in the detection of positive selection in our study, these adjacent substitutions could have either a mutational or a selective origin (Stoletzki and Eyre-Walker, 2011). New models incorporating MNMs have been developed and these may help to determine the cause of adjacent substitutions (Venkat et al, 2018). However, it is still a challenge to distinguish between nucleotides affected by authentic positive selection from those caused by neutral fixation of MNMs (especially in the particular case of the amino acid serine; Rogozin et al, 2016); in addition, this bias has mainly been discussed in branch-site, and not in site analyses still poorly explored, and used in our paper (Dong et al, 2021). In consequence, we decided to rely on the conservative pipeline we had designed to attenuate the impact of all false positives, to avoid an over-filtration of potential PSG, especially when MNVs are shared, or partially shared, between neighbor species (Averof et al, 2000).

**Impact of large amplicon duplications on PSGs:** We investigated the impact of large duplications (segmental duplications) on PSGs since they can 1) lead to misidentify orthologous exons across species (Conrad and Antonarakis, 2007; Blake and Barolo, 2014); and 2) be a fertile recombination substrate (Bailey and Eichler, 2006; Marques-Bonet et al.,

2009) they commonly trigger gene conversion events, which, in turn, may alter estimated evolutionary profiles. We used ASGART, an efficient tool we have developed, to quickly establish a SD map for any genome (Delehelle et al, 2018). SDs were defined as sets of segments longer than 1000bp exhibiting more than a 90% identity rate between each other. To assess which genes could be potentially affected by SDs, we used ASGART to establish *de novo* exhaustive SD maps for all the concerned genomes. ASGART was run with its default settings (probe-size = 20, gap-size = 100, min-length = 1000), which are tuned toward relatively recent and well-conserved SDs, both direct and palindromic. All genes displaying evidence of positive selection, with at least one exon intersecting with a SD, were mentioned as duplicated in the final PSG list (Supplementary Table 8). Detecting duplication events was only possible for 881 exons from 24 PSGs, when the appropriate exon annotation was available (Supplementary Table 8 and 9).

**Impact of gBGC:** A region evolving at an accelerated rate on a particular branch of a phylogenetic tree is interpreted as evidence of an underlying positive selection pressure. However, selective pressure is not the only process leading to this result; gBGC events, through the biased fixation of G and C nucleotides, can also lead to the same observation. To mitigate gBGC-originated perturbation in our branch-site tests, we tested whether: 1) selection pressure-originated patterns of substitutions; 2) biased gene conversion; or 3) a mix of these two pressures were most likely to be responsible for the observed positive branch-site test results. We used the RPHAST package (Hubisz et al., 2011; Kostka et al., 2012) to compare a neutral model of evolution with three alternative tree models reflecting these possibilities. We implemented the three models and computed the likelihoods values corresponding to each of them; Likelihood Ratio Test (LRT) and the p-values were computed and corrected by the False Discovery Rate method (Benjamini and Hochberg, 1995). The lowest p-value associated with the highest LRT for these three tests was chosen as the best fitting scenario, and PSGs were

1 filtered accordingly. Results from the best-fit model and estimated parameters are provided in  
2  
3  
4 Supplementary Table 10.  
5  
6  
7

### 8 **Genome quality assessment: impact of heterogeneous genome quality on PSG detection?**

9  
10 Our study involves 27 genomes sequenced by multiple teams using various sequencing and  
11  
12 assembly methods (Supplementary Table 11); this variation may influence our results. In order  
13  
14 to estimate the potential impacts of these technicalities, we computed a set of metrics using the  
15  
16 QUASt-LG toolkit (Mikheenko et al., 2018). QUASt-LG computes two categories of metrics:  
17  
18 first, a set of length measures based on the physical characteristics of the contigs and scaffolds,  
19  
20 namely the N\* and L\* numbers, as well as the distribution of the lengths of the contigs and  
21  
22 scaffolds. Second, leveraging the BUSCO database and when genome annotations are  
23  
24 available, the toolkit assesses the presence or absence of a curated set of orthologous and  
25  
26 paralogous gene families that are hypothesized to be present and well conserved in the  
27  
28 concerned species. Using the default settings and, when available, the same genome  
29  
30 annotations as used for the rest of the study, the computation amounted to 123 linear CPU-  
31  
32 hours (Xeon Gold 6140@2.30GHz), most of them being consumed during the BUSCO genes  
33  
34 search. All metrics computed with QUASt-LG are summarized in Supplementary Table 11.  
35  
36 To avoid using specific metrics to characterize genome quality, we performed a PCA to plot  
37  
38 the different genomes used according to their characteristics (Supplementary Figure 33). The  
39  
40 genomes are discriminated along axis 1 by length metrics (N\*, L\*) and along axis 2 by  
41  
42 BUSCO, GC content, *i.e.*, more biologically oriented metrics. To avoid any *a priori* on the  
43  
44 informativeness of the metrics and due to the ongoing debate of which metrics should be used  
45  
46 to assess genome quality (Jauhal and Newcomb, 2021), each genome was considered using its  
47  
48 spatial coordinates on the first two axes of the PCA analyzes: no correlation was detected  
49  
50 between the number of PSGs and the coordinates along axes 1 and 2, suggesting the absence  
51  
52 of an association between the PSG detected and the quality metrics considered here  
53  
54  
55  
56  
57  
58  
59  
60

1 (Supplementary Table 12).  
2  
3  
4  
5

6 **Impact of demography on PSG detection:** To assess whether demographic history could  
7 explain the large differences in PSG allele fixation, we tested the impact of demographic  
8 history on PSG detection. Precise estimates of census population size and effective population  
9 size ( $N_e$ ) were not available for all the 27 species; thus, we used the IUCN status, a proxy  
10 metric for recent demographic history, combining current census size and information on  
11 habitat fragmentation (Supplementary Table 13). We tested whether the number of inferred  
12 PSGs was correlated to IUCN status and found no relation between status rank and the number  
13 of PSGs detected for each species (Kruskal-Wallis chi-squared = 0.6056, df = 5, p-value =  
14 0.9877).  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

## 29 **8. Supplementary material**

  
30  
31  
32

33 Supplementary data are available at Genome Biology and Evolution online.  
34  
35  
36  
37

## 38 **9. Data Availability Statement**

  
39  
40  
41  
42

43 The genomes used, together with the structural and functional annotation of genes, are  
44 available at <https://genome-euro.ucsc.edu>. GO terms were extracted from <https://string-db.org>.  
45 All figures and tables generated during the study are included in the paper and/or in the  
46 supplementary material online. Sequence alignments for PSG are available at Zenedo, DOI:  
47 10.5281/zenodo.4744361. Additional data related to this paper may be requested from the  
48 authors.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## 10. Acknowledgments

This project was supported by two CNRS-MITI interdisciplinary research programs: the white PEPS AUDEVOL (Projets Exploratoires Premier Soutien) and the DEFI X-LiFE (Branches méconnues du vivant) granted to PB. AM and FD were supported by a PhD studentship from the Ministry of Research (France) and MC by a post-doc fellowship (AESOP plus, an Erasmus program promoting the France and South-Africa exchanges). This work used HPC resources from CALMIP (grant P1434) and we thank them for their support.

## References

- Adzhubei I, Jordan DM, Sunyaev SR. 2013. Predicting functional effect of human missense mutations using polyphen-2. *Current protocols in human genetics*. 76(1):7-20.
- Ai H, et al. 2014. Population history and genomic signatures for high-altitude adaptation in Tibetan pigs. *BMC genomics*. 15(1):1-14.
- Anisimova M, Yang Z. 2007. Multiple hypothesis testing to detect lineages under positive selection that affects only a few sites. *Mol Biol Evol*. 24(5):1219-1228.
- Arbiza L, Dopazo J, Dopazo H. 2006. Positive selection, relaxation, and acceleration in the evolution of the human and chimp genome. *PLoS Comput Biol*. 2(4):e38.
- Averof M, Rokas A, Wolfe KH, Sharp PM. 2000. Evidence for a high frequency of simultaneous double-nucleotide substitutions. *Science*. 287(5456):1283-1286.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Bailey JA, Eichler EE. 2006. Primate segmental duplications: crucibles of evolution, diversity and disease. *Nat Rev Genet.* 7(7):552-564.
- Beall CM. 2006. Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude hypoxia. *Integr Comp Biol.* 46(1):18-24.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological).* 57(1):289-300.
- Bernardi M, Couette S, Chateau Smith C, Montuire S. 2019. Middle ear pneumatization in nonhuman primates: A comparative analysis. *Am J Phys Anthropol.* 169(3):540-556.
- Blake VM, Barolo S. 2014. Genome evolution: How sister genes grow apart. *Curr Biol.* 24(15):R695-R697.
- Brown, CH, Waser PM. 2017. Primate habitat acoustics. In: Quam RM, Ramsier MA, Fay RR, Popper AN, editors. *Primate hearing and communication.* Springer Cham. p. 79-107.
- Cagan A, et al. 2016. Natural selection in the great apes. *Mol Biol Evol.* 33(12):3268-3283.
- Cao X, Sun YB, Irwin DM, Wang GD, Zhang YP. 2015. Nocturnal to diurnal transition in the common ancestor of haplorrhines: evidence from genomic-scan for positively selected genes. *J Genet Genomics.* 42(1):33-37.
- Charlton BD, Owen MA, Swaisgood RR. 2019. Coevolution of vocal signal characteristics and

1 hearing sensitivity in forest mammals. *Nat Commun.* 10(1):1-7.

2  
3  
4  
5  
6 Chen Y, Zhang S, Chai R, Li H. 2019. Hair cell regeneration. In: Li H, Chai R, editors. *Hearing*  
7  
8 *Loss: Mechanisms, Prevention and Cure*. Springer Singapore p. 1-16.

9  
10  
11  
12  
13 Cingi C, Erkan AN, Rettinger G. 2010. Ear, nose, and throat effects of high altitude. *Eur Arch*  
14  
15 *Oto-rhino-l.* 267(3):467-471.

16  
17  
18  
19  
20 Coleman MN, Boyer DM. 2012. Inner ear evolution in primates through the Cenozoic:  
21  
22 implications for the evolution of hearing. *Anat Rec.* 295(4):615-631.

23  
24  
25  
26  
27 Coleman MN, Colbert MW. 2010. Correlations between auditory structures and hearing  
28  
29 sensitivity in non-human primates. *J Morphol.* 271(5):511-532.

30  
31  
32  
33  
34  
35 Coleman MN, Ross CF. 2004. Primate auditory diversity and its influence on hearing  
36  
37 performance. *Anat Rec Part A.* 281(1):1123-1137.

38  
39  
40  
41  
42 Conrad B, Antonarakis SE. 2007. Gene duplication: a drive for phenotypic diversity and cause  
43  
44 of human disease. *Annu Rev Genom Hum Genet.* 8:17-35.

45  
46  
47  
48  
49 Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics  
50  
51 and parallel computing. *Nat Methods.* 9(8):772-772.

52  
53  
54  
55  
56 Daub JT, Moretti S, Davydov II, Excoffier L, Robinson-Rechavi M. 2017. Detection of  
57  
58 pathways affected by positive selection in primate lineages ancestral to humans. *Mol Biol Evol.*  
59  
60 34(6):1391-1402.

1  
2  
3  
4 Delehelle F, Cussat-Blanc S, Alliot JM, Luga H, Balaesque P. 2018. ASGART: fast and  
5 parallel genome scale segmental duplications mapping. *Bioinformatics*. 34(16):2708-2714.  
6  
7

8  
9  
10 Di Franco A, Poujol R, Baurain D, Philippe H. 2019. Evaluating the usefulness of alignment  
11 filtering methods to reduce the impact of errors on evolutionary inferences. *BMC Evol Biol*.  
12 19(1):1-17.  
13  
14  
15

16  
17  
18  
19  
20 Dominy NJ, Ross CF, Smith TD. 2004. Evolution of the special senses in primates: past,  
21 present, and future. *Anat Rec Part A*. 281(1):1078-1082.  
22  
23  
24

25  
26  
27 Dong X, Liang Q, Li J, Feng, P. 2021. Positive selection drives the evolution of a primate  
28 bitter taste receptor gene. *Ecol Evol*. 00:1-9.  
29  
30  
31

32  
33  
34 Du TT, et al. 2019. LMO7 deficiency reveals the significance of the cuticular plate for hearing  
35 function. *Nature Commun*. 10(1):1-15.  
36  
37  
38

39  
40  
41 Fan D, Ren H, Danzeng D, Li H, Wang P. 2016. Influence of high-altitude hypoxic  
42 environments on the survival of cochlear hair cells and spiral ganglion neurons in rats. *Biomed*  
43 *Rep*. 5(6):681-685.  
44  
45  
46  
47

48  
49  
50 Fettiplace R. 2017. Hair cell transduction, tuning, and synaptic transmission in the mammalian  
51 cochlea. *Compr Physiol*. 7(4):1197-1227.  
52  
53  
54

55  
56  
57  
58 Fleagle JG. 2013. Chapter 2 - the primate body. In: Fleagle JG, editor. *Primate Adaptation and*  
59 *Evolution*. Academic Press San Diego. p. 9-33.  
60

1  
2  
3  
4 Fletcher W, Yang Z. 2010. The effect of insertions, deletions, and alignment errors on the  
5  
6 branch-site test of positive selection. *Mol Biol Evol.* 27(10):2257-2267.  
7  
8

9  
10 Franchini LF, Elgoyhen AB. 2006. Adaptive evolution in mammalian proteins involved in  
11  
12 cochlear outer hair cell electromotility. *Mol Phylogenet Evol.* 41(3):622-635.  
13  
14

15  
16 Galtier N, Duret L, Glémin S, Ranwez V. 2009. GC-biased gene conversion promotes the  
17  
18 fixation of deleterious amino acid changes in primates. *Trends Genet.* 25(1):1-5.  
19  
20  
21

22  
23 Gibbs RA, et al. 2007. Evolutionary and biomedical insights from the rhesus macaque genome.  
24  
25  
26 *Science.* 316(5822):222-234.  
27  
28

29  
30 Hall TA. 1999. BioEdit: A User-Friendly Biological Sequence Alignment Editor and Analysis  
31  
32 Program for Windows 95/98/NT. *Nucl Acid S.* 41:95-98.  
33  
34

35  
36 Heffner RS. 2004. Primate hearing from a mammalian perspective. *Anat Rec Part A.*  
37  
38  
39 281(1):1111-1122.  
40  
41

42  
43 Hickox AE, et al. 2017. Global analysis of protein expression of inner ear hair cells. *J Neurosci.*  
44  
45  
46 37(5):1320-1339.  
47  
48

49  
50 Hubisz MJ, Pollard KS, Siepel A. 2011. PHAST and RPHAST: phylogenetic analysis with  
51  
52 space/time models. *Brief Bioinform.* 12(1):41-51.  
53  
54

55  
56 Jauhal AA, Newcomb RD. 2021. Assessing genome assembly quality prior to downstream  
57  
58  
59  
60

1 analysis: N50 versus BUSCO. *Mol Ecol Resour.* 00:1–6.

2  
3  
4  
5  
6 Karolchik D, et al. 2004. The UCSC Table Browser data retrieval tool. *Nucleic Acids Res.*  
7  
8 32(suppl\_1):D493-D496.  
9

10  
11  
12  
13 Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:  
14  
15 improvements in performance and usability. *Mol Biol Evol.* 30(4):772-780.  
16  
17

18  
19  
20 Kirwan JD, et al. 2013. A phylomedicine approach to understanding the evolution of auditory  
21  
22 sensory perception and disease in mammals. *Evol Appl.* 6(3):412-422.  
23  
24

25  
26  
27 Köppl C, Manley GA. 2018. A functional perspective on the evolution of the cochlea. *CSH*  
28  
29 *Perspect Med.* a033241.  
30

31  
32  
33 Kosiol C, et al. 2008. Patterns of positive selection in six mammalian genomes. *PLoS Genet.*  
34  
35 4(8):e1000144.  
36  
37

38  
39  
40 Kostka D, Hubisz MJ, Siepel A, Pollard KS. 2012. The role of GC-biased gene conversion in  
41  
42 shaping the fastest evolving regions of the human genome. *Mol Biol Evol.* 29(3):1047-1057.  
43  
44

45  
46  
47 Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a fast, scalable  
48  
49 and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics.* 35(21):  
50  
51 4453-4455.  
52  
53

54  
55  
56 Kuderna LF, Esteller-Cucala P, Marques-Bonet T. 2020. Branching out: what omics can tell  
57  
58 us about primate evolution. *Curr Opin Genet Dev.* 62:65-71.  
59  
60

1  
2  
3  
4 Lartillot N. 2013. Phylogenetic patterns of GC-biased gene conversion in placental mammals  
5 and the evolutionary dynamics of recombination landscapes. *Mol Biol Evol.* 30(3):489-502.  
6  
7

8  
9  
10 Li G, et al. 2008. The hearing gene Prestin reunites echolocating bats. *Proc Natl Acad Sci USA.*  
11  
12 105(37):13959-13964.  
13  
14

15  
16  
17 Lindblad-Toh K., et al. 2011. A high-resolution map of human evolutionary constraint using  
18  
19 29 mammals. *Nature.* 478(7370):476-482.  
20  
21

22  
23  
24 Liu H, et al. 2014. Characterization of transcriptomes of cochlear inner and outer hair cells. *J*  
25  
26 *Neurosci.* 34(33):11085-11095.  
27  
28

29  
30  
31 Mallick S, Gnerre S, Muller P, Reich D. 2009. The difficulty of avoiding false positives in  
32  
33 genome scans for natural selection. *Genome Res.* 19(5):922-933.  
34  
35

36  
37  
38 Manley GA. 2017. Comparative auditory neuroscience: understanding the evolution and  
39  
40 function of ears. *J Assoc Res Oto.* 18(1):1-24.  
41  
42

43  
44  
45 Markova-Raina P, Petrov D. 2011. High sensitivity to aligner and high rate of false positives  
46  
47 in the estimates of positive selection in the 12 Drosophila genomes. *Genome Res.* 21(6): 863-  
48  
49 874.  
50  
51

52  
53  
54 Marques-Bonet T, Girirajan S, Eichler EE. 2009. The origins and impact of primate segmental  
55  
56 duplications. *Trends Genet.* 25(10):443-454.  
57  
58  
59  
60

1 Masters JC. 1991. Loud calls of *Galago crassicaudatus* and *G. garnettii* and their relation to  
2 habitat structure. *Primates*. 32(2):153-167.  
3

4  
5  
6  
7  
8 McBee RM, Rozmiarek SA, Meyerson NR, Rowley PA, Sawyer SL. 2015. The effect of  
9 species representation on the detection of positive selection in primate gene data sets. *Mol Biol*  
10 *Evol*. 32(4):1091-1096.  
11  
12

13  
14  
15  
16  
17 Michalski N, Petit C. 2015. Genetics of auditory mechano-electrical transduction. *Pflüg Arch*  
18 *Eur J Phy*. 467(1):49-72.  
19  
20

21  
22  
23  
24 Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome  
25 assembly evaluation with QUASt-LG. *Bioinformatics*. 34(13):i142–i150.  
26  
27

28  
29  
30  
31 Mikkelsen T, et al. 2005. Initial sequence of the chimpanzee genome and comparison with the  
32 human genome. *Nature*. 437(7055):69-87.  
33  
34

35  
36  
37  
38 Mutai H, et al. 2018. Gene expression dataset for whole cochlea of *Macaca fascicularis*. *Sci*  
39 *Rep*. 8(1):1-9.  
40  
41

42  
43  
44  
45 Nielsen R, Yang Z. 1998. Likelihood models for detecting positively selected amino acid sites  
46 and applications to the HIV-1 envelope gene. *Genetics*. 148(3):929-936.  
47  
48

49  
50  
51  
52 Nummela S. 2017. The Primate Peripheral Auditory System and the Evolution of Primate  
53 Hearing. In: Quam RM, Ramsier MA, Fay RR, Popper AN, editors. *Primate hearing and*  
54 *communication*. Springer Cham. p. 13-45.  
55  
56  
57  
58  
59  
60



1 Olson-Manning CF, Wagner MR, Mitchell-Olds T. 2012. Adaptive evolution: evaluating  
2 empirical support for theoretical predictions. *Nat Rev Genet.* 13(12):867-877.  
3  
4  
5

6  
7  
8 Paradis E, Claude J, Strimmer K. 2004. APE: analyses of phylogenetics and evolution in R  
9 language. *Bioinformatics.* 20(2):289-290.  
10  
11  
12

13  
14  
15 Parker J, et al. 2013. Genome-wide signatures of convergent evolution in echolocating  
16 mammals. *Nature.* 502(7470):228-231.  
17  
18  
19

20  
21  
22 Penn O, et al. 2010. GUIDANCE: a web server for assessing alignment confidence scores.  
23  
24  
25  
26  
27  
28

29 Perelman P., et al. 2011. A molecular phylogeny of living primates. *Plos Genet.* 7(3):  
30 e1001342.  
31  
32  
33  
34  
35

36 Pisciotto F, et al. 2019. Inner ear genes underwent positive selection and adaptation in the  
37 mammalian lineage. *Mol Biol Evol.* 36(8):1653-1670.  
38  
39  
40  
41  
42

43 Prunier J, et al. 2017. Gene copy number variations in adaptive evolution: the genomic  
44 distribution of gene copy number variations revealed by genetic mapping and their adaptive  
45 role in an undomesticated species, white spruce (*Picea glauca*). *Mol Ecol.* 26(21):5989-6001.  
46  
47  
48  
49  
50

51 Qiu Q, et al. 2012. The yak genome and adaptation to life at high altitude. *Nat genet.* 44(8):946-  
52 949.  
53  
54  
55  
56  
57

58 R Core Team. 2013. R: A language and environment for statistical computing.  
59  
60

1  
2  
3  
4 Ramsier MA, Cunningham AJ, Finneran JJ, Dominy NJ. 2012. Social drive and the evolution  
5 of primate hearing. *Philos T R Soc B*. 367(1597):1860-1868.  
6  
7  
8

9  
10 Ramsier MA, Rauschecker JP. 2017. Primate audition: Reception, perception, and ecology. In:  
11 Quam RM, Ramsier MA, Fay RR, Popper AN, editors. *Primate hearing and communication*.  
12 Springer Cham. p. 47-77  
13  
14  
15  
16  
17

18  
19 Ratnakumar A, et al. 2010. Detecting positive selection within genomes: the problem of biased  
20 gene conversion. *Philos T R Soc B*. 365(1552):2571-2580.  
21  
22  
23  
24  
25

26 Richardson GP, Petit C. 2019. Hair-bundle links: genetics as the gateway to function. *CSH*  
27 *Perspect Med*. 9(12):a033142.  
28  
29  
30  
31

32 Rogozin IB, et al. 2016. Evolutionary switches between two serine codon sets are driven by  
33 selection. *Proc Natl Acad Sci USA*. 113(46):13109-13113.  
34  
35  
36  
37  
38

39 Romiguier J, Roux C. 2017. Analytical biases associated with GC-content in molecular  
40 evolution. *Front Genet*. 8:16.  
41  
42  
43  
44  
45

46 Scally A, et al. 2012. Insights into hominid evolution from the gorilla genome sequence.  
47 *Nature*. 483(7388):169-175.  
48  
49  
50  
51

52 Scheffer DI, Shen J, Corey DP, Chen ZY. 2015. Gene expression by mouse inner ear hair cells  
53 during development. *J Neurosci*. 35(16):6366-6380.  
54  
55  
56  
57  
58  
59  
60

1 Schrago CG, Seuánez HN. 2019. Large ancestral effective population size explains the difficult  
2 phylogenetic placement of owl monkeys. *Am J Primatol.* 81(3):e22955.

3  
4  
5  
6  
7  
8 Sela I, Ashkenazy H, Katoh K, Pupko T. 2015. GUIDANCE2: accurate detection of unreliable  
9 alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids Res.*  
10 43(W1):W7-W14.

11  
12  
13  
14  
15  
16  
17 Springer MS, et al. 2012. Macroevolutionary dynamics and historical biogeography of primate  
18 diversification inferred from a species supermatrix. *Plos one.* 7(11):e49521.

19  
20  
21  
22  
23  
24 Stoletzki N, Eyre-Walker, A. 2011. The Positive Correlation between dN/dS and dS in  
25 Mammals Is Due to Runs of Adjacent Substitutions. *Mol Biol Evol.* 28(4): 1371-1380.

26  
27  
28  
29  
30  
31 Szklarczyk D, et al. 2019. STRING v11: protein–protein association networks with increased  
32 coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*  
33 *Acids Res.* 47(D1):D607-D613.

34  
35  
36  
37  
38  
39  
40 Van Der Lee R, Wiel L, Van Dam TJ, Huynen MA. 2017. Genome-scale detection of positive  
41 selection in nine primates predicts human-virus evolutionary conflicts. *Nucleic Acids Res.*  
42 45(18):10634-10648.

43  
44  
45  
46  
47  
48  
49 Vater M, Meng J, Fox RC. 2004. Hearing organ evolution and specialization: early and later  
50 mammals. In: Manley GA, Fay RR, editors. *Evolution of the vertebrate auditory system.*  
51 Springer, New York. p. 256-288.

52  
53  
54  
55  
56  
57  
58  
59 Waser PM, Brown CH. 1986. Habitat acoustics and primate communication. *Am J Primatol.*  
60

1 10(2):135-154.  
2  
3  
4  
5

6 Wong KM, Suchard MA, Huelsenbeck JP. 2008. Alignment uncertainty and genomic analysis.  
7  
8 *Science*. 319(5862):473-476.  
9

10  
11  
12 Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*.  
13 24(8):1586-1591.  
14  
15  
16  
17  
18

19 Yang, Z, Dos Reis M. 2010. Statistical properties of the branch-site test of positive selection.  
20  
21 *Mol Biol Evol*. 28(3):1217-1228.  
22  
23  
24  
25

26 Yang, Z, Swanson WJ. 2002. Codon-substitution models to detect adaptive evolution that  
27 account for heterogeneous selective pressures among site classes. *Mol Biol Evol*. 19(1):49-57.  
28  
29  
30  
31  
32

33 Yang Z, Nielsen R, Goldman N, Pedersen AMK. 2000. Codon-substitution models for  
34 heterogeneous selection pressure at amino acid sites. *Genetics*. 155(1):431-449.  
35  
36  
37  
38  
39

40 Yang Z, Wong WS, Nielsen R. 2005. Bayes empirical Bayes inference of amino acid sites  
41 under positive selection. *Mol Biol Evol*. 22(4):1107-1118.  
42  
43  
44  
45  
46

47 Yin Y, et al. 2020. The draft genome of mandrill (*Mandrillus sphinx*): An old World monkey.  
48  
49 *Sci Rep*. 10(1):1-8.  
50  
51  
52  
53

54 Zhang J, Nielsen R, Yang Z. 2005. Evaluation of an improved branch-site likelihood method  
55 for detecting positive selection at the molecular level. *Mol Biol Evol*. 22(12): 2472-2479.  
56  
57  
58  
59  
60

1 Zhou X, et al. 2014. Whole-genome sequencing of the snub-nosed monkey provides insights  
2 into folivory and evolutionary history. *Nat genet.* 46(12):1303-1310.  
3  
4  
5  
6  
7

8 Zhou X, et al. 2016. Population genomics reveals low genetic diversity and adaptation to  
9 hypoxia in snub-nosed monkeys. *Mol Biol Evol.* 33(10):2670-2681.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1. Overview of our comparative evolutionary analysis procedure for conservative inference of positively selected genes and branches. 1. Selection of 123 genes from the EARxpressed repository; 2. Multiple sequence alignments and maximum likelihood-based phylogenetic analyses; 3. Site tests (PAML) and confounding factor analyses (alignment and multiple test corrections); 4. Branch-site tests (PAML) and confounding factor analyses (alignment and multiple test corrections); 5. Additional analyzes of complexifying factors (duplications and gBGC).

412x633mm (300 x 300 DPI)




Figure 2. Cladogram of primate species displaying the 31 validated PSGs of branch-site test analyzes after multiple tests plus alignment corrections and a final visual inspection. PSGs potentially impacted by duplications and gBGC are mentioned in italics. Branch lengths are unscaled.

592x385mm (300 x 300 DPI)




Figure 3. Examples of the position of selected PSG amino acid (AA) sites (in black square) in the protein sequences of the four genes: ADGRV1, KCNQ4, PCDH15 and PTPRQ.

555x375mm (300 x 300 DPI)






Figure 4. (A) Cochlear duct and its cellular structure including hair cell stereociliary bundles; (B) details of the stereocilium and hair bundles, and the structure and location of link types including kinocilial links, tip links, horizontal top connectors, shaft connectors, and ankle links (b); (C) Zoom on the tip link connecting two stereocilium - adapted from Fettiplace (2017) and Richardson and Petit (2019).

197x292mm (300 x 300 DPI)