

Examples of finite time blow up in mass dissipative reaction-diffusion systems with superquadratic growth

Michel Pierre, Didier Schmitt

▶ To cite this version:

Michel Pierre, Didier Schmitt. Examples of finite time blow up in mass dissipative reaction-diffusion systems with superquadratic growth. Discrete and Continuous Dynamical Systems - Series A, 2023, 43 (3&4), pp.1686-1701. 10.3934/dcds.2022039 . hal-03360258v2

HAL Id: hal-03360258 https://hal.science/hal-03360258v2

Submitted on 20 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Manuscript submitted to AIMS' Journals Volume **X**, Number **0X**, XX **200X** doi:10.3934/xx.xxxxxx

pp. **X–XX**

EXAMPLES OF FINITE TIME BLOW UP IN MASS DISSIPATIVE REACTION-DIFFUSION SYSTEMS WITH SUPERQUADRATIC GROWTH

MICHEL PIERRE

Ecole Normale Supérieure de Rennes and Institut de Recherche Mathématique de Rennes, Campus de Ker Lann, 35170-Bruz, France

DIDIER SCHMITT

Université de Lorraine, CNRS, Institut Élie Cartan de Lorraine, BP 70239, 54506 Vandœuvre-lès-Nancy Cedex, France

ABSTRACT. We provide explicit examples of finite time L^{∞} -blow up for the solutions of 2 × 2 reaction-diffusion systems for which three main properties hold: positivity is preserved for all time, the total mass is uniformly controlled and the growth of the nonlinear reaction terms is superquadratic. They are obtained by choosing the space dimension large enough. This is to be compared with recent global existence results of uniformly bounded solutions for the same kind of systems with quadratic or even slightly superquadratic growth depending on the dimension. Such blow up may occur even with homogeneous Neumann boundary conditions. All these L^{∞} -blowing up solutions may be extended as weak global solutions. Blow up examples are also provided in space dimensions one, two and three with various growths.

4 1. Introduction. The main goal of this paper is to provide examples of blow up
5 in finite time for 2 × 2 reaction-diffusion systems of the form

$$\begin{cases} \partial_t u_1 - d_1 \Delta u_1 = f_1(u_1, u_2) \text{ on } (0, T) \times B_N, \\ \partial_t u_2 - d_2 \Delta u_2 = f_2(u_1, u_2) \text{ on } (0, T) \times B_N, \\ \text{"good " boundary conditions on } (0, T) \times \partial B_N, \\ u_1(0, \cdot) = u_1^0 \ge 0, \ u_2(0, \cdot) = u_2^0 \ge 0, \end{cases}$$
(1)

⁶ where B_N is the open unit ball in \mathbb{R}^N , $d_1, d_2 \in (0, +\infty)$, $f_1, f_2 : [0, +\infty)^2 \to \mathbb{R}$ are ⁷ regular nonlinearities such that the nonnegativity of the solutions to (1) is preserved ⁸ for all time which, as it is well-known, is equivalent to

(P) quasipositivity:
$$f_1(0, s_2) \ge 0, f_2(s_1, 0) \ge 0, \forall s_1, s_2 \in [0, +\infty),$$
 (2)

9 and such that the following 'mass dissipativity' condition holds

(M) dissipativity: $f_1(s_1, s_2) + f_2(s_1, s_2) \le 0, \forall s_1, s_2 \in [0, +\infty).$ (3)

¹⁰ This last property implies that the $L^1(B_N)$ -norm of the solutions $u_1(t), u_2(t)$ of (1)

¹¹ does not blow up in finite time (= control of the total mass for all time).

²⁰²⁰ Mathematics Subject Classification. Primary: 35B44,35K57,35K40;Secondary: 35K91.

Key words and phrases. Reaction-diffusion, blow up, global existence, dissipative evolution, quasipositivity.

It is well-known that, for $u_1^0, u_2^0 \in L^{\infty}(B_N)^+$, there exist $T \in (0, +\infty]$ and a classical solution to (1) on [0, T). If all the d_i are equal, then $T = +\infty$ so that global classical existence holds. However, if the d_i are different from each other, it is known that the two conditions (**P**), (**M**) are in general not sufficient to provide global classical solutions on $[0, +\infty)$ as proved in [11], [12] where examples of solutions blowing up in $L^{\infty}(B_N)$ in finite time are described for this class of systems.

⁸ A main new result proved here (see Theorem 2.1) is that actually, for all d =⁹ $2 + \eta, \eta > 0$, there exists a choice of $d_1, d_2 \in (0, +\infty)$, of the space dimension N and ¹⁰ of the nonlinearities f_1, f_2 satisfying (**P**), (**M**) and the growth condition (**G**) below, ¹¹ for which the solution to (1) blows up in $L^{\infty}(B_N)$ in finite time $T < +\infty$:

(G) d-growth: $|f_i(s_1, s_2)| \le C_0 + C_1(s_1 + s_2)^d, \ \forall s_1, s_2 \in [0, +\infty), i = 1, 2,$ (4)

where $C_0, C_1 \in [0, +\infty)$. The point is that finite time blow up may occur for any superquadratic growth.

This result has to be analyzed in parallel with the recent results in [4], [5] saying 14 that, for any dimension N, there exists $\eta > 0$, depending on the dimension N, such 15 that, if the growth of f_1, f_2 at infinity is at most $d = 2 + \eta$, then all solutions of (1) 16 are global in time: in particular, the $L^{\infty}(B_N)$ -norm of $u_1(t), u_2(t)$ is bounded on 17 all intervals [0, T]. Note that these global existence results are proved in [4], [5] even 18 19 for all $m \times m$ systems, $m \ge 2$, with the properties (**P**), (**M**), (**G**). Theorem 2.1 shows that these results are in some sense optimal. Besides [4], [5], several papers had 20 provided global existence results of classical solutions for close to quadratic systems 21 like [7], [6], [2], [14]. More results may also be found in [3], [15], [8], [1], [13]. We re-22 fer to the introduction in [4] where a nice state of art on these questions is described. 23 24

²⁵ Obviously, given $\eta > 0$ and f_1, f_2 with a $2+\eta$ growth, to obtain blow up examples, ²⁶ it is necessary to work in space dimensions N which are higher and higher when η ²⁷ is smaller and smaller.

Four more facts are also interesting about these blow up examples.

1) First it turns out that, not only the L^{∞} -norm blows up at time t = T, but also the L^m -norm for all $m \ge N(1+\eta)/2$, N large.

2) Next these *T*-blowing up solutions can nevertheless be extended as global weak solutions on $[0, +\infty)$ of the corresponding system (1) (see Section 7).

33 3) We also prove (see Theorem 2.2)) that the same kind of solutions provide 34 blow up examples for the following kind of systems:

$$\begin{pmatrix}
\partial_t u_1 - d_1 \Delta u_1 = c_1(t, x) u_1^{\alpha} u_2^{\beta} \text{ on } (0, T) \times B_N, \\
\partial_t u_2 - d_2 \Delta u_2 = c_2(t, x) u_1^{\alpha} u_2^{\beta} \text{ on } (0, T) \times B_N, \\
\text{"good " boundary conditions on } (0, T) \times \partial B_N, \\
u_1(0, \cdot) = u_1^0 \ge 0, \ u_2(0, \cdot) = u_2^0 \ge 0,
\end{cases}$$
(5)

where $c_1 + c_2 \leq 0, c_1, c_2 \in L^{\infty}(Q_T)$ and $\alpha + \beta = 2 + \eta$ with $\eta > 0$ as small as we want.

4) The boundary conditions of the examples provided in Theorem 2.1 are nonhomogeneous boundary conditions. We also describe blowing up examples for systems with homogeneous Neumann boundary conditions, up to replacing f_i , i = 1, 2 by more general (still regular) nonlinearities $g_i = g_i(t, x, s_1, s_2)$. This is indicated in Theorem 2.3.

An optimal dependence of N in terms of η or d is not explicitly known in the main result of Theorem 2.1. What comes out in the proof of this theorem [see (28)], is that one should choose N large enough so that

$$\frac{2N-\theta}{N-\theta} \le d = 2+\eta \quad \Leftrightarrow \quad N \ge \theta(1+\eta^{-1}),$$

1 where $\theta \in (4/3, +\infty)$. This means that we should have $N > 4(1 + \eta^{-1})/3$.

Another natural question is, given the dimension N, to find the optimal growth which provides global existence (or blow up in finite time) for nonlinearities satisfying the main properties (**P**), (**M**). As proved in [11], [12], examples of blow up may occur even in dimension N = 1 by choosing a large enough growth for the nonlinearity. It is known that in dimension N = 1, global existence is proved for at most cubic growth [see [6], [15]], but the optimal growth is not known.

To progress in this understanding, we give in Section 8 explicit blow up examples in the following cases :

$$N = 1, d = 6; N = 2, d = 7/2; N = 3, d = 3.$$

8 2. Statement of the main results. Some notation. Here B_N denotes the open 9 euclidian unit ball in \mathbb{R}^N , $Q_T = (0, T) \times B_N$, $\Sigma_T = (0, T) \times \partial B_N$. We also denote by 10 $C^{\infty}(A_1, A_2)$ the family of C^{∞} -mappings from A_1 to A_2 where $A_i \subset \mathbb{R}^{N_i}, i = 1, 2,$ 11 and N_1, N_2 are positive integers. If $A_2 = \mathbb{R}$, we simply denote $C^{\infty}(A_1)$.

Theorem 2.1. Let $d := 2 + \eta, \eta \in (0, +\infty)$ and $T \in (0, +\infty)$. Then there exist $f_1, f_2 \in C^{\infty}([0, +\infty)^2)$ satisfying $(\mathbf{P}) + (\mathbf{M}) + (\mathbf{G}), d_1, d_2 \in (0, +\infty)$, a dimension N large enough, $u_1^0, u_2^0 \in C^{\infty}(\overline{B_N})^+$, $\alpha_1, \alpha_2 \in C^{\infty}([0, T])^+$ and u_1, u_2 nonnegative C^{∞} -solutions of (1) with

$$u_1(t,x) = \alpha_1(t), \ u_2(t,x) = \alpha_2(t) \ on \ \Sigma_T,$$

and such that

$$\lim_{t \to T^{-}} \|u_1(t)\|_{L^m(B_N)} = \lim_{t \to T^{-}} \|u_2(t)\|_{L^m(B_N)} = +\infty,$$

12 for $m \ge N(d-1)/2$. Moreover, for λ close to 1

$$(\mathbf{M}_{\lambda}) \ f_1(s_1, s_2) + \lambda f_2(s_1, s_2) \le 0, \ \forall s_1, s_2 \in [0, +\infty).$$
(6)

Remark 2.1. Despite the blow up of $u_1(t)$, $u_2(t)$ at time t = T, it turns out that the 13 solutions u_i provided by Theorem 2.1 can be extended to the whole interval $[0, +\infty)$ 14 as weak global solutions of the system (1) on the whole interval $[0, +\infty)$. By weak 15 solution, we mean that the nonlinear terms $f_i(u_1, u_2), i = 1, 2$ are in $L^1((0, \tau) \times B_N)$ 16 for all $\tau \in (0, +\infty)$ and u_1, u_2 are solutions in the sense of distributions or in the 17 sense of the 'variation of constants' formula. This is a consequence of the fact that, 18 in the blow up examples, not only $f_1 + f_2 \leq 0$ holds, but also $f_1 + \lambda f_2 \leq 0$ for some 19 $\lambda \neq 1$. This provides the L¹-bound on the nonlinear terms and this is sufficient for 20 the existence of weak global solutions. All this is made more precise in Section 7. 21

Theorem 2.2. Let $d := 2 + \eta, \eta \in (0, +\infty), \alpha, \beta \in (1, +\infty)$ with $\alpha + \beta = d$ and $T \in (0, +\infty)$. Then there exist a dimension N large enough, $d_1, d_2 \in (0, +\infty)$ and

$$\begin{cases} c_1, c_2 \in C^{\infty}\left([0,T] \times \overline{B_N}\right) \cap L^{\infty}(Q_T), \\ u_1^0, u_2^0 \in C^{\infty}(\overline{B_N})^+, \ \alpha_1, \alpha_2 \in C^{\infty}([0,T])^+, \end{cases}$$

and u_1, u_2 nonnegative C^{∞} -solutions of the system (5) with $u_i = \alpha_i, i = 1, 2$ on Σ_T and such that

$$c_1(t,x) + c_2(t,x) \le 0, \quad \forall (t,x) \in Q_T,$$
$$\lim_{t \to T^-} \|u_1(t)\|_{L^m(B_N)} = \lim_{t \to T^-} \|u_2(t)\|_{L^m(B_N)} = +\infty$$

 $\quad \text{ for } m \geq N(d-1)/2.$

2 Remark 2.2. The solutions u_1, u_2 appearing in this result are the same as those in 3 Theorem 2.1. Only the interpretation of the nonlinear part changes. In particular, 4 they blow up in $L^m(B_N)$ as $t \to T^-$ for the same values of m.

It turns out that we can also use the same kind of solutions to construct new blowing up solutions for similar systems, but with *homogeneous Neumann boundary conditions*. Here the nonlinearities depend also on (t, x). This is the purpose of the next theorem.

9 **Theorem 2.3.** Let $d := 2 + \eta, \eta \in (0, +\infty)$ and $T \in (0, +\infty)$. Then there exist $\tau_0 \in [0, T), g_1, g_2 \in C^{\infty}([\tau_0, T) \times \overline{B_N} \times [0, +\infty)^2)$ satisfying $(\mathbf{P}') + (\mathbf{M}'_{\lambda}) + (\mathbf{G}')$ below, 11 $d_1, d_2 \in (0, +\infty)$, a dimension N large, $v_1^0, v_2^0 \in C^{\infty}(\overline{B_N})^+$ and v_1, v_2 nonnegative 12 C^{∞} -solutions of

$$\begin{aligned}
\partial_t v_1 &- d_1 \Delta v_1 = g_1(t, x, v_1, v_2) in (\tau_0, T) \times B_N, \\
\partial_t v_2 &- d_2 \Delta v_2 = g_2(t, x, v_1, v_2) in (\tau_0, T) \times B_N, \\
\partial_\nu v_1 &= 0 = \partial_\nu v_2 on (\tau_0, T) \times B_N, \\
v_i(\tau_0, \cdot) &= v_i^0 \ge 0,
\end{aligned}$$
(7)

and such that

$$\lim_{t \to T^{-}} \|v_1(t)\|_{L^m(B_N)} = \lim_{t \to T^{-}} \|v_2(t)\|_{L^m(B_N)} = +\infty,$$

13 for $m \ge N(d-1)/2$.

Here the conditions $(\mathbf{P}') + (\mathbf{M}'_{\lambda}) + (\mathbf{G}')$ are the same as $(\mathbf{P}) + (\mathbf{M}_{\lambda}) + (\mathbf{G})$, but twith a (t, x)-dependence, namely,

$$\begin{aligned} \forall (t,x) \in [\tau_0, T] \times \overline{B_N}, \ \forall (s_1, s_2) \in [0, +\infty)^2, \\ (\mathbf{P}') \ g_1(t,x,0,s_2) \ge 0, \ g_2(t,x,s_1,0) \ge 0, \\ (\mathbf{M}'_{\lambda}) \ g_1(t,x,s_1,s_2) + \lambda g_2(t,x,s_1,s_2) \le 0, \ \forall \lambda \text{ close to } 1, \\ (\mathbf{G}') \ |g_1(t,x,s_1,s_2)| + |g_2(t,x,s_1,s_2)| \le C_0 + C_1(s_1+s_2)^d. \end{aligned}$$

$$(8)$$

16

17 3. Steps of the proof of Theorem 2.1. The main idea in the proof of Theorem 2.1

18 is similar to the approach in [11], [12]. It consists in working with functions u_1, u_2 of the 19 form

$$u_i(t,x) = \frac{a_i(T-t) + b_i r^2}{[(T-t) + r^2]^{\gamma}}, \ r = |x|, i = 1, 2,$$
(9)

where a_1, a_2, b_1, b_2 will be well-chosen in $(0, +\infty)$, as well as $\gamma \in (1, 2)$ and the dimension N (large enough), in such a way that there exist $d_1, d_2 \in (0, +\infty)$ and two functions f_1, f_2 as described in Theorem 2.1 for which

$$\partial_t u_i - d_i \Delta u_i = f_i(u_1, u_2), \quad \text{in } (0, T) \times B_N, \ i = 1, 2.$$
 (10)

The values of u_1, u_2 on ∂B_N are obviously C^{∞} -functions. Moreover the $L^{\infty}(B_N)$ -norm of $u_1(t), u_2(t)$ blows up as $t \to T^-$ since $\gamma > 1$. More precisely, we directly check (see Lemma 3.1) that

$$\lim_{t \to T^{-}} \|u_i(t)\|_{L^m(B_N)} = +\infty, \quad \forall m \ge \frac{N}{2(\gamma - 1)}, \quad i = 1, 2.$$
(11)

Thus Theorem 2.1 will be proved if we can achieve (10) with regular functions f_1, f_2 satisfying (**P**), (**M**), (**G**) with $d = 2 + \eta \ge \gamma/(\gamma - 1)$. It will be a consequence of the three

1 following lemmas.

2

8

17

First note that $u_i, i = 1, 2$, is of the following form:

$$u_i(t,x) = (T-t)^{1-\gamma} \rho_i\left(\frac{r^2}{T-t}\right), \ \rho_i(\sigma) = \frac{a_i + b_i\sigma}{(1+\sigma)^{\gamma}}, i = 1, 2.$$
(12)

- 4 We have the following technical lemma.
- 5 Lemma 3.1. Let u_1, u_2 be given by (9). For i = 1, 2, we have in $(0, T) \times B_N$

$$\partial_t u_i - d_i \Delta u_i = (T - t)^{-\gamma} (1 + \sigma)^{-(\gamma + 2)} [A_i + B_i \sigma + C_i \sigma^2], \ \sigma = r^2 / (T - t),$$
(13)

6 where A_i, B_i, C_i are real numbers given by

$$\begin{pmatrix}
A_i = (\gamma - 1)a_i - 2d_iN(b_i - \gamma a_i), \\
B_i = (\gamma - 1)[2a_i + b_i(1 + 2d_iN + 4d_i)] + (b_i - \gamma a_i)[4d_i(\gamma + 1) - 2d_iN + 1], \\
C_i = (\gamma - 1)[a_i + b_i(1 + 2d_iN - 4\gamma d_i)] + b_i - \gamma a_i.
\end{cases}$$
(14)

7 As a consequence

$$\partial_t u_1 - d_1 \Delta u_1 + \partial_t u_2 - d_2 \Delta u_2 = (T - t)^{-\gamma} (1 + \sigma)^{-(\gamma + 2)} [A + B\sigma + C\sigma^2], \quad (15)$$

$$A = (\gamma - 1)a - 2NE + 2N\gamma G, B = (\gamma - 2)a + \gamma b + [2N(\gamma - 2) + 8\gamma]E - 2\gamma[2(\gamma + 1) - N]G, C = -a + \gamma b + 2(\gamma - 1)(N - 2\gamma)E,$$
(16)

9 where only the four following quantities are involved

$$a := a_1 + a_2, \ b := b_1 + b_2, E := b_1 d_1 + b_2 d_2, \ G := d_1 a_1 + d_2 a_2.$$
 (17)

10 Moreover, $\lim_{t\to T^-} \|u_i(t)\|_{L^m(B_N)} = +\infty$ if $m \ge N/2(\gamma - 1)$.

The proof of this lemma is elementary (see below). Next we check that $\partial_t u_i - d_i \Delta u_i$ may be written as a function of u_1, u_2 . This is the purpose of the following lemma. Here \mathbb{Q} denotes the set of rational numbers.

14 Lemma 3.2. Let u_1, u_2 be given by (9). Assume

$$\gamma \in (1,2) \cap \mathbb{Q}, \quad a_1 b_2 - a_2 b_1 \neq 0. \tag{18}$$

15 Then for i = 1, 2, there exists a C^{∞} -function $P_i : (0, +\infty)^2 \to \mathbb{R}^2$ which is homogeneous 16 of degree $\gamma' = \gamma/(\gamma - 1)$ and such that

$$\partial_t u_i - d_i \Delta u_i = P_i(u_1, u_2) \quad in \ (0, T) \times B_N, \tag{19}$$

$$|P_i(s_1, s_2)| \le K(s_1 + s_2)^{\gamma'}, \ \forall (s_1, s_2) \in (0, +\infty)^2, \ for \ some \ K \in (0, +\infty),$$
(20)

$$P_i(a_1 + b_1\sigma, a_2 + b_2\sigma) = (1 + \sigma)^{\frac{\gamma}{\gamma - 1}} [A_i + B_i\sigma + C_i\sigma^2], \ \forall \sigma \in (0, +\infty).$$
(21)

19 **Remark 3.1.** By homogeneous of degree γ' , we mean that

$$P_i(\lambda s_1, \lambda s_2) = \lambda^{\gamma'} P_i(s_1, s_2), \quad \forall \lambda \in (0, +\infty), (s_1, s_2) \in (0, +\infty)^2.$$
(22)

20 This lemma will be proved below. But let us continue describing the scheme of the proof

of Theorem 2.1. A main step is now to prove that we can choose the various parameters $a_i, b_i, d_i, i = 1, 2, d, \gamma$ and the dimension N such that the functions P_1, P_2 appearing in Lemma 3.2 satisfy

$$P_1(s_1, s_2) + P_2(s_1, s_2) \le 0, \ \forall (s_1, s_2) \in (0, +\infty)^2.$$
(23)

By (21), this inequality will be satisfied for $(s_1, s_2) = (a_1 + b_1\sigma, a_2 + b_2\sigma), \sigma \in (0, +\infty)$ if and only if

$$A + B\sigma + C\sigma^2 \le 0, \ \forall \sigma \in (0, +\infty),$$

where A, B, C are defined in (15) and (16) of Lemma 3.1. It will be the case if

$$A < 0, \ C < 0, \ B^2 < 4AC.$$
 (24)

And we then construct the P_i so that this inequality extends to all $(s_1, s_2) \in (0, +\infty)^2$.

The main point in the proof of Theorem 2.1 is that we can choose the various parameters in the definition (9) so that (24) holds for A, B, C as defined in Lemma 3.1. This is the purpose of the following main lemma where we use the following notation.

Notation. Writing a function $D : \mathbb{R} \to \mathbb{R}$ as $D(N) = O(N^{-\alpha}), \alpha \in \mathbb{R}$, means that

$$\limsup_{N \to +\infty} |D(N)| N^{\alpha} < +\infty$$

5 Lemma 3.3. In the definition (9) of u_1, u_2 , let us choose

$$\begin{cases} a_1 = 1/N^2, \ a_2 = 2N - N^{-2}, \ b_1 = b_2 = 1/\sqrt{N}, \\ d_1 = \sqrt{N}, \ d_2 = 1/N^3, \ \gamma = 2 - \theta/N, \ \theta \in (4/3, +\infty) \cap \mathbb{Q}. \end{cases}$$
(25)

6 Then as $N \to +\infty$, the values A, B, C, as defined in Lemma 3.1, satisfy

$$\begin{cases} A = -2\theta + O(N^{-\frac{1}{2}}) < 0, \quad C = -(8+2\theta) + O(N^{-\frac{1}{2}}) < 0, \\ B^2 - 4AC = 64(4-3\theta) + O(N^{-\frac{1}{2}}) < 0, \\ A + B\sigma + C\sigma^2 \le A - B^2/4C = \frac{8(4-3\theta)}{4+\theta} + O(N^{-\frac{1}{2}}) < 0, \forall \sigma \in [0, +\infty). \end{cases}$$
(26)

- 7 Moreover (18) is satisfied and the functions P_1, P_2 can be chosen in Lemma 3.2 so that

$$P_1(s_1, s_2) + \lambda P_2(s_1, s_2) \le 0, \ \forall (s_1, s_2) \in (0, +\infty)^2,$$
(27)

9 for λ close to 1.

4. Proof of Theorem 2.1. Theorem 2.1 is a consequence of Lemmas 3.1, 3.2, 3.3. We
will prove these lemmas below. But let us first show how these lemmas imply Theorem 2.1.

- 13 We consider the functions u_1, u_2 as defined in (9) with the choice of parameters
- 14 $a_1, a_2, b_1, b_2, d_1, d_2, \gamma$ given in Lemma 3.3 with $\theta \in (4/3, +\infty) \cap \mathbb{Q}$ and N large enough so 15 that

$$\gamma' = \frac{2N - \theta}{N - \theta} \le d = 2 + \eta.$$
⁽²⁸⁾

16 Since by assumption $\theta \in \mathbb{Q}$, we have $\gamma \in \mathbb{Q}$. Moreover $\gamma \in (1, 2)$ for $N > \theta$. For this choice 17 of the parameters, $a_2b_1 - a_1b_2 = 2N^{1/2} + O(N^{-5/2}) \neq 0$ for N large.

Thus by Lemma 3.2, there exist two C^{∞} -functions $P_1, P_2: (0, +\infty)^2 \to \mathbb{R}$ such that

$$\partial_t u_i - d_i \Delta u_i = P_i(u_1, u_2), \quad in \ (0, T) \times B_N,$$

$$|P_i(s_1, s_2)| \le K(s_1 + s_2)^{\gamma'}, \forall (s_1, s_2) \in (0, +\infty)^2, i = 1, 2, \text{ for some } K \in (0 + \infty).$$

- By Lemma 3.3, we can even choose P_1, P_2 so that (27) holds for λ close to 1.
- 19

Note that there exists $m_N \in (0, +\infty)$ such that

$$u_i(t,x) \ge m_N, \ \forall (t,x) \in [0,T] \times \overline{B}_N, \ i=1,2.$$

Indeed, we may choose $m_N = (T+1)^{1-\gamma} \min\{a_1, a_2, b_1, b_2\}$ as it can be seen by writing

$$u_{i}(t,x) = (T - t + r^{2})^{1 - \gamma} \frac{a_{i} + b_{i}\sigma}{1 + \sigma}$$

Let us introduce $\varphi \in C^{\infty}([0, +\infty), [0, +\infty))$ such that

$$0 \leq \varphi \leq 1, \ \varphi \equiv 0 \ on \ [0, m_N/2], \ \varphi \equiv 1 \ on \ [m_N, +\infty).$$

Let us denote $f_i(s_1, s_2) := \varphi(s_1)\varphi(s_2)P_i(s_1, s_2), i = 1, 2$ on $[0, +\infty)^2$. Then

$$\partial_t u_i - d_i \Delta u_i = f_i(u_1, u_2), \ in \ (0, T) \times B_N,$$

and f_1, f_2 are exactly as described in Theorem 2.1. More precisely, and for future reference, we even have

$$f_i(s_1, s_2) = 0, \ \forall (s_1, s_2) \in [0, +\infty)^2 \setminus [m_N/2, +\infty)^2.$$
 (29)

Next we have $u_i^0 \in C^{\infty}(\overline{B}_N)$ and $\alpha_i \in C^{\infty}([0,T])$ for i = 1, 2 since

$$\begin{cases} u_i^0(x) = (T+r^2)^{-\gamma}(a_iT+b_ir^2), \\ \alpha_i(t) = (T-t+1)^{-\gamma}(a_i(T-t)+b_i), \ i=1,2. \end{cases}$$

Now, we know by Lemma 3.1, that the $L^m(B_N)$ -norm of $u_i(t), i = 1, 2$ blows up if $m \ge N[2(\gamma - 1)]^{-1}$. Here γ is chosen [see (28)] so that $\gamma' \le d$, that is $(\gamma - 1)^{-1} \le d - 1$. If $m \ge N(d-1)/2$, then $m \ge N[2(\gamma - 1)]^{-1}$ and the $L^m(B_N)$ -norm of $u_i(t)$ blows up. This ends the proof of the main Theorem 2.1, assuming Lemmas 3.1, 3.2, 3.3.

5 6

Let us now prove these three lemmas.

7

Proof of Lemma 3.1. This result is derived by technical but easy computations. We may use (see (12)) $u_i = (T-t)^{1-\gamma} \rho_i(\sigma), \sigma = r^2/(T-t)$ so that

$$\partial_t u_i = (\gamma - 1)(T - t)^{-\gamma} \rho_i(\sigma) + \frac{r^2}{(T - t)^2} (T - t)^{1 - \gamma} \rho_i'(\sigma),$$

$$\partial_r u_i = (T - t)^{1 - \gamma} \left(\frac{2r}{T - t}\right) \rho_i'\left(\frac{r^2}{T - t}\right),$$

$$\partial_{rr} u_i = (T - t)^{1 - \gamma} \left[\frac{4r^2}{(T - t)^2} \rho_i''\left(\frac{r^2}{T - t}\right) + \frac{2}{T - t} \rho_i'\left(\frac{r^2}{T - t}\right)\right],$$

$$\frac{N - 1}{r} \partial_r u_i + \partial_{rr} u_i = (T - t)^{-\gamma} \left[2N\rho_i'(\sigma) + 4\sigma\rho_i''(\sigma)\right].$$

 $\partial_t u_i - d_i \Delta u_i = (T - t)^{-\gamma} \left[(\gamma - 1)\rho_i(\sigma) + (\sigma - 2d_i N)\rho_i'(\sigma) - 4d_i \sigma \rho_i''(\sigma) \right].$ With the choice of $\rho_i(\sigma) = (a_i + b_i \sigma)/(1 + \sigma)^{\gamma}$, we have

$$\rho_i'(\sigma) = \frac{b_i - \gamma a_i + b_i(1-\gamma)\sigma}{(1+\sigma)^{\gamma+1}},$$
$$\rho_i''(\sigma) = \frac{b_i(1-\gamma)}{(1+\sigma)^{\gamma+1}} - \frac{(\gamma+1)[b_i - \gamma a_i + b_i(1-\gamma)\sigma]}{(1+\sigma)^{\gamma+2}}.$$

8 This leads to

$$\partial_t u_i - d_i \Delta u_i = (T - t)^{-\gamma} (1 + \sigma)^{-(\gamma + 2)} \left[A_i + B_i \sigma + C_i \sigma^2 \right], \tag{30}$$

9 where A_i, B_i, C_i are given as stated in Lemma 3.1.

By summing for i = 1, 2 these two expressions, we obtain the claims (15)-(16)-(17) of Lemma 3.1. It is important to notice that this sum can be written only in terms of the

12 four parameters a, b, E, G.

Finally we compute the $L^m(B_N)$ -norm of $u_i(t)$. For some $C_N \in (0, +\infty)$

$$\int_{B_N} u_i^m(t,x) dx = C_N (T-t)^{m(1-\gamma)} \int_0^1 r^{N-1} \rho_i^m \left(r^2 / (T-t) \right) dr.$$

By setting $\sigma = r^2/(T-t) \Leftrightarrow r = \sqrt{T-t}\sqrt{\sigma}$, we have

$$\int_{B_N} u_i^m(t,x) dx = \frac{C_N}{2} (T-t)^{\frac{N}{2} + m(1-\gamma)} \int_0^{(T-t)^{-1}} \sigma^{\frac{N}{2}-1} \rho_i^m(\sigma) d\sigma.$$

As $\sigma^{\frac{N}{2}-1}\rho_i^m(\sigma)$ is equivalent to $\sigma^{\frac{N}{2}-1+m(1-\gamma)}$, up to a positive constant, as $\sigma \to +\infty$, we have

$$\frac{N}{2} < m(\gamma - 1) \Rightarrow \int_0^{+\infty} \sigma^{\frac{N}{2} - 1} \rho_i^m(\sigma) d\sigma < +\infty.$$

13 As a consequence, in this case $\int_{B_N} u_i^m(t,x) dx$ behaves like $(T-t)^{\frac{N}{2}+(1-\gamma)m}$, up to a con-

14 stant, as $t \to T^-$ and therefore tends to $+\infty$. The conclusion is the same in the equality 15 case $N = 2m(\gamma - 1)$ since then $\int_{B_N} u^m(t, x) dx$ behaves like $|\log(T - t)|$ as $t \to T^-$. 1 **Proof of Lemma 3.2.** We are looking for two functions $P_i(\cdot, \cdot)$, i = 1, 2 such that

$$\partial_t u_i - d_i \Delta u_i = P_i(u_1, u_2) = P_i\left((T-t)^{1-\gamma} \rho_1(\sigma), (T-t)^{1-\gamma} \rho_2(\sigma) \right), \ \sigma = r^2/(T-t).$$
(31)

According to (13) in Lemma 3.1, and by the expected γ' -homogeneity of P_i , this is equivalent to

$$(1+\sigma)^{-(\gamma+2)}[A_i + B_i\sigma + C_i\sigma^2] = (1+\sigma)^{\gamma^2/(1-\gamma)}P_i(a_1 + b_1\sigma, a_2 + b_2\sigma),$$

$$\Leftrightarrow (1+\sigma)^{\frac{2-\gamma}{\gamma-1}} [A_i + B_i \sigma + C_i \sigma^2] = P_i(a_1 + b_1 \sigma, a_2 + b_2 \sigma), \,\forall \sigma \in [0, +\infty).$$
(32)

Let us write $\gamma' = p/q \in (2, +\infty) \cap \mathbb{Q}$ with p, q two coprime positive integers. Then

$$\gamma = p/(p-q), \ (2-\gamma)/(\gamma-1) = (p-2q)/q > 0.$$

3 Thus (32) may be rewritten

$$(1+\sigma)^{\frac{p-2q}{q}}[A_i + B_i\sigma + C_i\sigma^2] = P_i(a_1 + b_1\sigma, a_2 + b_2\sigma),$$
(33)

4 which also implies

$$(1+\sigma)^{p-2q}[A_i + B_i\sigma + C_i\sigma^2]^q = \{P_i(a_1 + b_1\sigma, a_2 + b_2\sigma)\}^q.$$
(34)

Here p > 2q and the left-hand side is a polynomial of degree at most p in σ . It is easy to see that, since $a_1b_2 - a_2b_1 \neq 0$, the family of p + 1 polynomials

$$\{(a_1 + b_1\sigma)^k (a_2 + b_2\sigma)^{p-k}, k = 0, ..., p\}$$

is a linear basis of the linear space of polynomials of degree at most p. Indeed, let us analyze the relation

$$\sum_{k=0}^{p} \mu_k (a_1 + b_1 \sigma)^k (a_2 + b_2 \sigma)^{p-k} \equiv 0, \ \mu_k \in \mathbb{R}, \ \forall k = 0, ..., p.$$

Assume by contradiction that there exists $r \in \{0, ..., p\}$ such that

$$\mu_k = 0, \ \forall k = 0, ..., r - 1, \ \mu_r \neq 0.$$

Then after dividing by $(a_1 + b_1 \sigma)^r$, the above relation may be rewritten

$$\sum_{k=r}^{p} \mu_k (a_1 + b_1 \sigma)^{k-r} (a_2 + b_2 \sigma)^{p-k} = 0.$$

5 If r = p, this obviously implies $\mu_r = 0$ whence a contradiction. If r < p, choosing $\sigma =$

6 $-a_1/b_1$ leads to $\mu_r(a_2 - b_2a_1/b_1)^{p-r} = 0$, whence again $\mu_r = 0$ thanks to the assumption 7 (18) on $a_i, b_i, i = 1, 2$, and this is a contradiction.

8 We are now going to define the convenient function P_i . Going back to the polynomial 9 of degree at most p involved in (34), we can claim that there exist $\lambda_k^i \in \mathbb{R}, k = 0, ..., p$ such 10 that for all $\sigma \in \mathbb{R}$

$$(1+\sigma)^{p-2q}[A_i+B_i\sigma+C_i\sigma^2]^q = \sum_{k=0}^p \lambda_k^i (a_1+b_1\sigma)^k (a_2+b_2\sigma)^{p-k}.$$
 (35)

11 Assume first that q is odd. Let us define

$$\widetilde{P}_i(s_1, s_2) := \left\{ \sum_{k=0}^p \lambda_k s_1^k s_2^{p-k} \right\}^{1/q}, \quad (s_1, s_2) \in [0, +\infty)^2.$$
(36)

12 Note that \widetilde{P}_i is homogeneous of degree p/q and

$$(1+\sigma)^{\frac{p-2q}{q}}[A_i+B_i\sigma+C_i\sigma^2] = \widetilde{P}_i(a_1+b_1\sigma,a_2+b_2\sigma).$$
(37)

13 According to (30), this implies [see also (31), (32)]

$$\partial_t u_i - d_i \Delta u_i = \tilde{P}_i(u_1, u_2) \text{ in } Q_T.$$
(38)

Moreover, we directly see on the definition (36) of \widetilde{P}_i that for $(s_1, s_2) \in [0, +\infty)$

$$|\widetilde{P}_i(s_1, s_2)| \le \left\{ \sum_{k=0}^p |\lambda_k| \right\}^{1/q} (s_1 + s_2)^{p/q} = C(s_1 + s_2)^{\gamma'}$$

Now, for the regularity of \tilde{P}_i , it is obvious from its definition (36) that \tilde{P}_i is C^{∞} around each point (s_1, s_2) for which $\tilde{P}_i(s_1, s_2) \neq 0$. We are going to modify it into a new function P_i which will be in $C^{\infty}((0, +\infty)^2)$ with the same main properties. For this, let us assume without loss of generality in (18) that $a_1b_2 - a_2b_1 > 0$ or $b_2/b_1 > a_2/a_1$ and let us consider the function

$$s \in (0, +\infty) \to X := \frac{a_2 + b_2 s}{a_1 + b_1 s} \in I := \left(\frac{a_2}{a_1}, \frac{b_2}{b_1}\right).$$

1 It is C^{∞} with derivative $(a_1b_2 - a_2b_1)/(a_1 + b_1s)^2 > 0$ on $(0, +\infty)$. Thus its inverse 2 $\psi: I \mapsto (0, +\infty)$ is also C^{∞} .

From (36), we may write

$$\widetilde{P}_i(s_1, s_2) = s_1^{p/q} \widetilde{P}_i(1, s_2/s_1),$$

3 and thanks to the relation (37) and the p/q homogeneity of \widetilde{P}_i , we have

$$\widetilde{P}_{i}(1,X) = [a_{1} + b_{1}\psi(X)]^{-p/q} [1 + \psi(X)]^{(p-2q)/q} [A_{i} + B_{i}\psi(X) + C_{i}\psi(X)^{2}].$$
(39)

Whence the C^{∞} -property of $\left[X \in I \mapsto \widetilde{P}_i(1, X)\right]$ and therefore of $\widetilde{P}_i(s_1, s_2)$ on \mathcal{I} where we set

$$\mathcal{I} := \{ (s_1, s_2) \in (0, +\infty)^2; s_2/s_1 \in I \}.$$

Actually, $X \in \mathbb{R} \mapsto \widetilde{P}_i(1, X)$ is continuous and does not vanish in a neighborhood of the extremities of I, namely

$$X = a_2/a_1 \Leftrightarrow \psi(X) = 0, \ \widetilde{P}_i(1, X) = A_i a_1^{-p/q},$$
$$X = b_2/b_1 \Leftrightarrow \psi(X) = +\infty, \ \widetilde{P}_i(1, X) = C_i b_1^{-p/q}.$$

- 4 Thus, as noticed before thanks to the definition (36), $X \mapsto \widetilde{P}_i(1, X)$ is C^{∞} even on some
- 5 open interval J containing \overline{I} . We now introduce a function $\chi \in C^{\infty}([0, +\infty))$ such that:

$$0 \le \chi \le 1, \ \chi \equiv 1 \ on \ I, \ \chi \equiv 0 \ on \ [0, +\infty) \setminus J.$$
 (40)

And we set

9

$$P_i(s_1, s_2) := s_1^{p/q} \chi(s_2/s_1) \widetilde{P}_i(1, s_2/s_1) = \chi(s_2/s_1) \widetilde{P}_i(s_1, s_2).$$

⁶ Thus P_i is C^{∞} on all of $(0, +\infty)^2$ and coincides with \tilde{P}_i on \mathcal{I} . It follows that we may ⁷ replace \tilde{P}_i by P_i in the relations (37) and (38). This ends the proof of Lemma 3.2 when q⁸ is odd.

10 Assume now that q is even. We start again from the relation (35) from which we deduce

$$(1+\sigma)^{(p-2q)/q} [A_i + B_i \sigma + C_i \sigma^2] = \widetilde{P}_i (a_1 + b_1 \sigma, a_2 + b_2 \sigma) sign\{A_i + B_i \sigma + C_i \sigma^2\}, \quad (41)$$

11 where we again set

$$\widetilde{P}_{i}(s_{1},s_{2}) := \left\{ \sum_{k=0}^{p} \lambda_{k}^{i} s_{1}^{k} s_{2}^{p-k} \right\}^{1/q} = s_{1}^{p/q} \widetilde{P}_{i}(1,s_{2}/s_{1}), \ \forall (s_{1},s_{2}) \in (0,+\infty)^{2},$$
(42)

and where

$$sign(r) = 1, \forall r \in (0, +\infty), \ sign(0) = 0, \ sign(r) = -1, \forall r \in (-\infty, 0).$$

12 We deduce from (41) and the p/q homogeneity of P_i that, for all $X \in I$,

$$\begin{cases} \widetilde{P}_{i}(1,X)sign\left(A_{i}+B_{i}\psi(X)+C_{i}\psi(X)^{2}\right)\\ = (a_{1}+b_{1}\psi(X))^{-p/q}(1+\psi(X))^{(p-2q)/q}[A_{i}+B_{i}\psi(X)+C_{i}\psi(X)^{2}] \end{cases}$$
(43)

and this function of X is C^{∞} on I. Since $\tilde{P}_i(1, X)$ does not vanish near the extremities of the interval I, it is also locally C^{∞} there. Moreover $sign(A_i + B_i\psi(X) + C_i\psi(X)^2)$ is constant as X tends to these extremities from inside I. It follows that the function of X appearing in (43) may be extended in a C^{∞} - way to an open interval J containing \overline{I} . Using the same function χ as above (see (40)), by setting

$$P_i(s_1, s_2) := s_1^{p/q} \chi(s_2/s_1) \widetilde{P}_i(1, s_2/s_1) sign\left(A_i + B_i \psi(s_2/s_1) + C_i \psi^2(s_2/s_1)\right)$$

we define a $C^\infty\text{-}\mathrm{function}$ on $(0,+\infty)^2$ which is $p/q\text{-}\mathrm{homogeneous}$ and satisfies

$$P_i(a_1 + b_1\sigma, a_2 + b_2\sigma) = (1 + \sigma)^{(p-2q)/q} [A_i + B_i\sigma + C_i\sigma^2], \ \forall \sigma \in [0, +\infty)$$

and therefore (see (31), (32), (33))

$$\partial_t u_i - d_i \Delta u_i = P_i(u_1, u_2) \text{ on } Q_T$$

This ends the proof of Lemma 3.2.

Proof of Lemma 3.3. As a consequence of the choice of the parameters in Lemma 3.3, we obtain

$$a = a_1 + a_2 = 2N, \ b = b_1 + b_2 = 2/\sqrt{N},$$

$$E = d_1b_1 + d_2b_2 = \sqrt{N}(1/\sqrt{N}) + N^{-3}(1/\sqrt{N}) = 1 + O(N^{-7/2}),$$

$$G = d_1a_1 + d_2a_2 = \sqrt{N}N^{-2} + N^{-3}[2N - (1/N^2)] = O(N^{-3/2}).$$

It follows from the formulas (16) in Lemma 3.1 that for N large

$$\begin{split} A &= [1 - \theta/N] 2N - 2N [1 + O(N^{-7/2})] + 2N [2 - \theta/N] O(N^{-3/2}) = -2\theta + O(N^{-1/2}), \\ C &= -2N + [2 - \theta/N] 2/\sqrt{N} + 2[1 - \theta/N] [N - 4 + 2\theta/N] [1 + O(N^{-7/2})] = -(8 + 2\theta) + O(N^{-1/2}), \\ B &= \begin{cases} [-\theta/N] 2N + [2 - \theta/N] [2/\sqrt{N}] + [2N(-\theta/N) + 8(2 - \theta/N)] [1 + O(N^{-7/2})] \\ -2[2 - \theta/N] [2(3 - \theta/N) - N] [O(N^{-3/2})] = 16 - 4\theta + O(N^{-1/2}). \end{cases} \\ B^2 - 4AC = 16[4 - \theta]^2 - 8\theta(8 + 2\theta) + O(N^{-1/2}) = 64[4 - 3\theta] + O(N^{-1/2}). \end{split}$$

3 With the choice of $\theta > 4/3$, we obtain $A < 0, C < 0, B^2 - 4AC < 0$ for N large as claimed 4 in Lemma 3.3.

Since the maximum of $\sigma \in \mathbb{R} \mapsto A + B\sigma + C\sigma^2$ is reached for $\sigma = -B/2C$, we deduce $A + B = + C\sigma^2 \in A - B^2/4C$ $K = O(N^{-\frac{1}{2}}) \forall \sigma \in [0, +\infty)$ (44)

$$A + B\sigma + C\sigma^2 \le A - B^2/4C = -K_\theta + O(N^{-\frac{1}{2}}), \forall \sigma \in [0, +\infty)$$

$$\tag{44}$$

6 with $K_{\theta} := 8(3\theta - 4)/(4 + \theta) > 0.$

On the other hand, $a_2b_1 - a_1b_2 = 2N^{1/2} + O(N^{-5/2}) \neq 0$ and $\gamma \in (1,2) \cap \mathbb{Q}$ so that (18) is satisfied and we may apply Lemma 3.2. Using now the relation (21) in this Lemma 3.2 together with (44), we obtain that

$$(P_1 + P_2)(a_1 + b_1\sigma, a_2 + b_2\sigma) \le -K_{\theta} + O(N^{-\frac{1}{2}}), \ \forall \sigma \in [0, +\infty).$$

By the p/q-homogeneity of $P_1 + P_2$, we deduce that

$$(P_1 + P_2)(1, X) \le -K_{\theta} a_1^{-p/q} + O(N^{-\frac{1}{2}}), \ \forall X \in I,$$

where I is defined as in the previous proof of Lemma 3.2. Since I is bounded, so is $P_2(1, X)$ for $X \in I$. It follows that for λ close to 1 and N large enough, we may claim that

$$(P_1 + \lambda P_2)(1, X) \le -\frac{K_{\theta} a_1^{-p/q}}{2}, \ \forall X \in I.$$

Up to reducing J and the support of the function χ in (40) of the proof of Lemma 3.2, we may claim that

$$(P_1 + \lambda P_2)(1, X) \le -\frac{K_\theta a_1^{-p/q}}{4}, \ \forall X \in J$$

By homogeneity, this implies that

$$(P_1 + \lambda P_2)(s_1, s_2) \le 0, \ \forall (s_1, s_2) \in (0, +\infty)^2.$$

7 This ends the proof of Lemma 3.3.

1 5. Proof of Theorem 2.2. We use the same functions u_1, u_2 as those introduced in the 2 proof of Theorem 2.1 and defined in (9), and we choose γ as in (28), namely such that

$$\gamma' \le d \quad \Leftrightarrow \quad \gamma \ge d/(d-1). \tag{45}$$

We deduce from (30) that

$$\partial_t u_i - d_i \Delta u_i = \frac{A_i (T-t)^2 + B_i (T-t) r^2 + C_i r^4}{(T-t+r^2)^{\gamma+2}}, \ i = 1, 2.$$

Given α , β as indicated in the statement of Theorem 2.2, we may understand this expression as being of the form

$$\partial_t u_i - d_i \Delta u_i = c_i(t, x) u_1^{\alpha} u_2^{\beta},$$

where we define

$$\begin{split} c_i(t,x) &:= \frac{A_i(T-t)^2 + B_i(T-t)r^2 + C_i r^4}{(T-t+r^2)^{\gamma+2}} u_1^{-\alpha} u_2^{-\beta}, \\ \Rightarrow \ c_i(t,x) &= \frac{(T-t+r^2)^{\gamma(d-1)-2} [A_i(T-t)^2 + B_i(T-t)r^2 + C_i r^4]}{[a_1(T-t) + b_1 r^2]^{\alpha} [a_2(T-t) + b_2 r^2]^{\beta}}. \end{split}$$

Obviously c_i is C^{∞} on $[0,T) \times \overline{B_N}$. Let us check that it is in $L^{\infty}(Q_T)$ as follows. Let $H := T - t + r^2$. We use

$$\Gamma - t \le H, r^2 \le H, a_i(T - t) + b_i r^2 \ge \min\{a_i, b_i\}H.$$

Thus for some $K_1, K_2 \in (0, +\infty)$, and using that $\gamma(d-1) - 2 \ge \gamma(d-1) - d \ge 0$, we have

$$|c_i(t,x)| \le \frac{H^{\gamma(d-1)-2}[[A_i] + |B_i| + |C_i|]H^2}{K_1 H^{\alpha+\beta}} \le K_2 H^{\gamma(d-1)-d} \le K_2 (T+1)^{\gamma(d-1)-d}.$$

3 Finally, we see that the sign of $c_1 + c_2$ is the same as the sign of $A(T-t)^2 + B(T-t)r^2 + Cr^4$, that is $a_1 + a_2 \leq 0$ since $A \leq 0$, $C \leq 0$, $B^2 = AAC \leq 0$ by Lemma 2.2. This and the proof

4 that is $c_1 + c_2 \leq 0$ since $A < 0, C < 0, B^2 - 4AC < 0$ by Lemma 3.3. This ends the proof 5 of Theorem 2.2.

6 6. Proof of Theorem 2.3. Let u_1, u_2 be the solutions obtained in Theorem 2.1. Their 7 normal derivatives on ∂B_N are given as follows

$$\partial_{r} u_{1}(t,1) = \frac{2[(b_{1}-\gamma a_{1})(T-t)-(\gamma-1)b_{1}]}{(T-t+1)^{\gamma+1}}, \quad \partial_{r} u_{2}(t,1) = \frac{2[(b_{2}-\gamma a_{2})(T-t)-(\gamma-1)b_{2}]}{(T-t+1)^{\gamma+1}}, \\ b_{1} = N^{-1/2} = b_{2}, \quad b_{1}-\gamma a_{1} = N^{-1/2} - 2N^{-2} + O(N^{-3}), \\ b_{2}-\gamma a_{2} = -4N + O(1).$$
(46)

8 Obviously for N large

$$\begin{cases} \partial_r u_1(t,1) \le 0, \ \forall t \in [\tau_0,T], \ \tau_0 := (T - \gamma + 1)^+ (< T), \\ \partial_r u_2(t,1) \le 0, \ \forall t \in [0,T]. \end{cases}$$
(47)

9 We now introduce the solutions $\beta_i, i = 1, 2$ of

$$\begin{cases} \partial_t \beta_i - d_i \Delta \beta_i = 0 \ in \ (\tau_0, T) \times B_N, \\ \partial_\nu \beta_i = -\partial_r u_i(t, 1) \ on \ (\tau_0, T) \times \partial B_N, \\ \beta_i(\tau_0, \cdot) \equiv 0. \end{cases}$$
(48)

By the C^{∞} -property of $\partial_r u_i(t, 1)$, these solutions are also in $C^{\infty}(\overline{Q_T})$. Moreover, thanks to (47), and by maximum principle, we have $\beta_i(t, x) \ge 0$ on $[\tau_0, T] \times B_N$. We now introduce for i = 1, 2

$$v_i(t,x) := u_i(t,x) + \beta_i(t,x) \ge 0$$
 so that $\partial_{\nu} v_i = \partial_r v_i(t,1) = 0$ on Σ_T .

And according to Theorem 2.1 and the definition of β_i , we have

$$\partial_t v_i - d_i \Delta v_i = \partial_t u_i - d_i \Delta u_i = f_i(u_1, u_2) = g_i(t, x, v_1, v_2) \text{ in } Q_T, \ i = 1, 2,$$

where we set

$$g_i(t, x, s_1, s_2) := f_i(s_1 - \beta_1(t, x), s_2 - \beta_2(t, x)).$$

Here the functions f_i are extended by 0 outside $[0, +\infty)^2$ and with the help of (29), we see that they are C^{∞} on \mathbb{R}^2 . The functions g_i defined in this way satisfy $(\mathbf{P}'), (\mathbf{M}'_{\lambda}), (\mathbf{G}')$ since, by nonnegativity of β_i :

$$\begin{aligned} g_1(t, x, 0, s_2) &= f_1(-\beta_1(t, x), s_2 - \beta_2(t, x)) = 0, \\ g_2(t, x, s_1, 0) &= f_2(s_1 - \beta_1(t, x), -\beta_2(t, x)) = 0, \\ (g_1 + \lambda g_2)(t, x, s_1, s_2) &= (f_1 + \lambda f_2)(s_1 - \beta_1(t, x), s_2 - \beta_2(t, x)) \le 0 \ [using \ (6)], \\ |g_i(t, x, s_1, s_2)| &= |f_i(s_1 - \beta_1(t, x), s_2 - \beta_2(t, x))| \\ &\leq C_0 + C_1[(s_1 - \beta_1(t, x))^+ + (s_2 - \beta_2(t, x)^+]^d \\ &\Rightarrow |g_i(t, x, s_1, s_2)| \le C'_0 + C'_1|s_1 + s_2|^d, \end{aligned}$$

1 where C'_0, C'_1 depend on $C_0, C_1, d, \|\beta_i\|_{L^{\infty}(Q_T)}$. Finally, the $L^m(B_N)$ -norm of $v_i(t)$ blows 2 up as $t \to T^-$ together with the $L^m(B_N)$ -norm of $u_i(t)$. This ends the proof of Theorem 3 2.3.

4 7. About global weak solutions. Let us prove here that the solutions u_1, u_2 obtained 5 in Theorem 2.1 may be extended to global weak solutions of the same system. For this we 6 will use the approach of the previous section and, as a first step, we extend the functions 7 v_1, v_2 to weak solutions on $[0, +\infty)$. This will provide extensions of u_1, u_2 as well. 8 Let us start by extending the functions β_i as the solutions on $[0, +\infty)$ of

$$\begin{cases} \partial_t \beta_i - d_i \Delta \beta_i = 0 \text{ in } (\tau_0, +\infty) \times B_N, \\ \partial_\nu \beta_i = \begin{cases} -\partial_r u_i(t, 1) \text{ on } (\tau_0, T) \times \partial B_N \\ -\partial_r u_i(T, 1) \text{ on } (T, +\infty) \times \partial B_N, \end{cases} \\ \beta_i(\tau_0, \cdot) \equiv 0. \end{cases}$$

$$(49)$$

These extensions are C^{∞} on $[\tau_0, +\infty) \times \overline{B_N}$. We also extend the nonlinear function g_i as follows for $(t, x) \in (\tau_0, +\infty) \times \overline{B_N}$

$$g_i(t, x, s_1, s_2) := f_i(s_1 - \beta_1(t, x), s_2 - \beta_2(t, x)), \ \forall (s_1, s_2) \in [0, +\infty)^2.$$

9 They satisfy the same properties $(\mathbf{P}'), (\mathbf{M}'_{\lambda}), (\mathbf{G}')$ on $[\tau_0, +\infty)$. We now apply Theorem 10 5.9 in [10] to the system

$$\begin{cases}
\partial_t v_1 - d_1 \Delta v_1 = g_1(t, x, v_1, v_2) in(\tau_0, +\infty) \times B_N, \\
\partial_t v_2 - d_2 \Delta v_2 = g_2(t, x, v_1, v_2) in(\tau_0, +\infty) \times B_N, \\
\partial_\nu v_1 = 0 = \partial_\nu v_2 on(\tau_0, +\infty) \times B_N, \\
v_i(\tau_0, \cdot) = v_i^0 = u_i(\tau_0, \cdot) + \beta(\tau_0, \cdot).
\end{cases}$$
(50)

Indeed, thanks to the property (\mathbf{M}'_{λ}) , any good approximation of this system with bounded nonlinearities g_i^n in place of g_i and the same regular data at $t = \tau_0$, will provide approximate regular solutions v_i^n such that, for all $S \in [\tau_0, +\infty)$

$$\sup_{n \ge 1} \int_{(\tau_0, S) \times B_N} |g_i^n(t, x, v_1^n(t, x), v_2^n(t, x))| < +\infty, \ i = 1, 2,$$

We may for instance choose

$$g_i^n(t, x, s_1, s_2) := \frac{g_i(t, x, s_1, s_2)}{1 + n^{-1}[(|g_1| + |g_2|)(t, x, s_1, s_2)]},$$

so that $|g_i^n| \leq n$. These approximate functions g_i^n satisfy also $(\mathbf{P}'), (\mathbf{M}'_{\lambda})$ and it is easy to prove that these properties imply the above $L^1((\tau_0, S) \times B_N)$ -bounds (see e.g. Proposition 5.1 in [10]). As a consequence, and as proved in [10], these approximate solutions v_i^n converge to a *weak solution* of (50) which means that

$$v_i(t) = S_i(t - \tau_0)v_i^0 + \int_{\tau_0}^t S_i(t - s)g_i\left(s, \cdot, v_1(s, \cdot), v_2(s, \cdot)\right) \, ds, \ i = 1, 2,$$

where $S_i(t)$ denotes the semigroup generated by the operator $-d_i\Delta$ on B_N with homogeneous Neumann boundary conditions.

These weak solution coincides on the interval $[\tau_0, T)$ with the classical solution of (7) as found in Theorem 2.3. Indeed, since $v_i^0 \in L^{\infty}(B_N)$, it is classical to prove that the approximate solution (v_1^n, v_2^n) stays uniformly bounded, independently of n, on some interval $[\tau_0, \tau_1] \subset [\tau_0, T]$. To see it, we might for instance use the inequalities

$$\partial_t v_i^n - d_i \Delta v_i^n \le |g_i(t, x, v_1^n, v_2^n)| \le C_0' + C_1' |v_1^n + v_2^n|^d.$$

1 Thus the limit (v_1, v_2) is therefore a classical solution on $[\tau_0, \tau_1]$. By uniqueness of classical

2 solutions for the reaction-diffusion system (7), both solutions coincide at least on $[\tau_0, \tau_1]$,

3 and subsequently on the whole interval $[\tau_0, T]$ by a classical continuity argument.

Finally, if we now set $u_i(t,x) := v_i(t,x) - \beta_i(t,x), i = 1, 2$ on the whole domain $[\tau_0, +\infty) \times \overline{B_N}$, we find a global weak solution of the initial system (1) on $[0, +\infty)$ which extends the (classical) solution obtained on [0,T) in Theorem 2.1. It remains to get convinced that $u_i \ge 0$ for all t. This can be seen by noticing that $u_i^n := v_i^n - \beta_i$ is a classical solution on $[\tau_0, +\infty)$ of

$$\partial_t u_i^n - d_i \Delta u_i^n = g_i^n(t, x, v_1^n, v_2^n) = f_i^n(u_1^n, u_2^n),$$

4 with boundary conditions $\partial_{\nu} u_i^n = -\partial_r \beta_i \ge 0$ with a quasi-positive nonlinearity (f_1^n, f_2^n) .

5 Thus $u_i^n \ge 0$ and this is preserved at the limit.

8. Some more explicit examples in small dimensions. We use here Lemma 3.1 and Lemma 3.2. All blow up examples we are going to describe here are defined as in (9) so that they blow up as $t \to T^-$. Moreover by Lemma 3.2, we know that we can then find P_1, P_2 so that (19) holds, namely

$$\partial_t u_i - d_i \Delta u_i = P_i(u_1, u_2), \text{ in } Q_T, \ i = 1, 2.$$

6 The goal is then to choose all parameters involved in the definition (9) well enough so that 7 (24) holds, namely

$$A < 0, \ C < 0, \ B^2 - 4AC < 0, \tag{51}$$

8 where A, B, C are defined in Lemma 3.1. Then, exactly as in the proof of Theorem 2.1, 9 we prove that the previous inequalities implies that $P_1 + P_2 \leq 0$ and that P_1, P_2 can then 10 slightly be modified into functions f_1, f_2 satisfying (**P**), (**M**) and (1). Moreover the growth 11 at infinity of f_1, f_2 is at most γ' .

The following technical proposition is very useful to help finding examples of blow up. Notations are as in Lemma 3.1.

14 **Proposition 8.1.** Let $a, b, E, G \in (0, +\infty)$. Then we can find $a_i, b_i, d_i, i = 1, 2 \in (0, +\infty)$ 15 such that

$$+a_2 = a, b_1 + b_2 = b, b_1d_1 + b_2d_2 = E, a_1d_1 + a_2d_2 = G,$$
 (52)

$$a_1b_2 - a_2b_1 = \frac{aE - bG}{d_2 - d_1}.$$
(53)

Proof of Proposition 8.1. Let us choose d_1, d_2 such that

 $0 < d_1 < \min\{Eb^{-1}, Ga^{-1}\} \le \max\{Eb^{-1}, Ga^{-1}\} < d_2.$

Then, given the four parameters a, b, E, G, a solution to (52) is given by

$$a_1 = \frac{ad_2 - G}{d_2 - d_1}, a_2 = \frac{G - ad_1}{d_2 - d_1}, b_1 = \frac{bd_2 - E}{d_2 - d_1}, b_2 = \frac{E - bd_1}{d_2 - d_1}.$$

- 17 We easily check that, with this choice, the relations (52), (53) are satisfied and all param-
- eters are positive. The relation (53) will be used to check (18).

19 Remark 8.1. In what follows, we give several explicit choices of parameters which lead

- to blow up examples according to the analysis of the previous sections. Thanks to the
- 21 Proposition 8.1, it is sufficient to provide the values of a, b, E, G such that (51) holds for
- 22 the corresponding values of A, B, C.

 a_1

8.1. Blow up with a cubic growth in dimension N = 3. Here we choose with $\epsilon \in (0, 1)$ small enough,

$$\gamma = 3/2, \ a = 10, \ b = 1, \ E = 1, \ G = \epsilon$$

Then, $aE - bG = 10 - \epsilon \neq 0$ and

$$A = (\gamma - 1)a - 6E + 6\gamma G = 5 - 6 + 9\epsilon = -1 + 9\epsilon < 0,$$

$$C = -a + \gamma b + 2(\gamma - 1)(3 - 2\gamma)E = -10 + 3/2 + 0 = -17/2 < 0,$$

$$B = (\gamma - 2)a + \gamma b + [6(\gamma - 2) + 8\gamma]E - 2\gamma[2(\gamma + 1) - 3]G = 11/2 - 6\epsilon.$$

$$B^{2} - 4AC = \frac{11^{2}}{4} - 4\frac{17}{2} + 0(\epsilon) = -15/4 + O(\epsilon) < 0.$$

1 Thus (51) holds so that this provides a blowing up example with growth $\gamma' = 3$. By 2 Lemma 3.1, for $m \geq 3$, the $L^m(B_N)$ -norm of the solution blows up as $t \to T^-$. Note 3 that since all inequalities are strict in this example, we could slightly modify it to obtain 4 a blowing up example with growth $3 - \sigma$ for some small $\sigma > 0$.

8.2. Blow up with a 7/2-growth in dimension N = 2. An example with such a growth was already mentioned in [12]. We give here another one, easier to compute. Let us choose the following parameters with $\epsilon > 0$ small enough.

$$\gamma = 7/5, \ a = 4, \ b = \epsilon, E = 1, G = \epsilon.$$

then $aE - bG = 4 - \epsilon^2 \neq 0$ and

$$\begin{split} A &= \frac{2}{5}a - 4E + \frac{28}{5}G = \frac{8}{5} - 4 + O(\epsilon) = -\frac{12}{5} + O(\epsilon) < 0, \\ C &= -a + \frac{7}{5}b - \frac{16}{25}E = -4 - \frac{16}{25} + O(\epsilon) = -\frac{116}{25} + O(\epsilon) < 0, \\ B &= -\frac{3}{5}a + \frac{7}{5}b + \frac{44}{5}E - \frac{14^2}{25}G = -\frac{12}{5} + \frac{44}{5} + O(\epsilon) = \frac{32}{5} + O(\epsilon). \\ B^2 - 4AC &= \frac{32^2}{25} - 4\frac{12}{5}\frac{116}{25} + O(\epsilon) = \frac{1}{125}[5120 - 5568] + O(\epsilon) = -\frac{448}{125} + O(\epsilon) < 0. \end{split}$$

Thus this provides a blow up example with growth $\gamma' = 7/2$ and an $L^m(B_N)$ -norm blowing up for $m \ge 5/2$. Again, since all inqualities are strict, we could find another example with growth $7/2 - \sigma$ for some $\sigma > 0$.

8.3. Blow up with a 6-growth in dimension N = 1. Here we choose

$$\gamma = 6/5, \ a = 5, \ b = 1/2, \ E = 1, G = 1/10.$$

A = 1 - 2 + 6/25 = -19/25 < 0,

then $aE - bG = 5 - 1/20 \neq 0$ and

$$C = -5 + 3/5 + 2(1/5)(1 - 12/5) = -22/5 - 14/25 = -124/25 < 0,$$

$$B = -4 + 3/5 + 8 - 12.17/250 = 473/125.$$

$$125^{2}[B^{2} - 4AC] = 473^{2} - 4.95.620 = 223729 - 235600 = -11871 < 0.$$

8 This provides a blow up example with growth $\gamma' = 6$ and with an $L^m(B_N)$ -norm 9 blowing up for $m \ge 5/2$. As above again, we could improve it to a lower growth $6 - \sigma$ for 10 some $\sigma > 0$.

REFERENCES

- [1] J. Cañizo, L. Desvillettes and K. Fellner, Improved duality estimates and applications to reaction-diffusion equations, Comm. in P.D.E. 39 (6) (2014), 1185-1204. 3 4 [2] M.C. Caputo, T. Goudon and A.F. Vasseur, Solutions of the 4-species quadratic reaction-5 diffusion systems are bounded and C^{∞} -smooth, in any space dimension, Analysis & PDE, **12** (7) (2019), 1773–1804. [3] B.P. Cupps, J. Morgan and B.Q. Tang, Uniform boundedness for reaction-diffusion systems with mass dissipation, SIAM J. Math. Anal., 53 (1) (2021), 323-350. 8 [4] K. Fellner, J. Morgan and B.Q. Tang, Uniform-in-time bounds for quadratic reaction-diffusion 10 systems with mass dissipation in higher dimensions, Discrete & Continuous Dynamical Systems - S, **14** (2) (2021), 635–651. [5] K. Fellner, J. Morgan and B.Q. Tang, Global classical solutions to quadratic systems with mass control in arbitrary dimensions, Annales IHP C, Ana. Non Linéaire, 37 (2020), 281-307. T. Goudon and A. Vasseur, Regularity analysis for systems of reaction-diffusion equations, [6]Ann. Sci. Éc. Norm. Sup., 43 (4) (2010), 117–142. Ya.I. Kanel', Solvability in the large of a system of reaction-diffusion equations with the [7]balance condition, Differentsialýe Uravneniya, 26 (1990), 448-458. E.-H. Laamri, Global existence of classical solutions for a class of reaction-diffusion systems, [8] Acta Appl. Math. 115 (2) (2011), 153-165. [9] M. Pierre, Weak solutions and supersolutions in L^1 for reaction-diffusion systems, J. Evol. 20 Equ. 3 (2003), 153-168. [10] M. Pierre, Global Existence in Reaction-Diffusion Systems with Dissipation of Mass : a Survey, Milan J. Math. 78 (2) (2010), 417-455. [11] M. Pierre and D. Schmitt, Blow up in reaction-diffusion systems with dissipation of mass, SIAM J. Math. Anal., 28 (1997), 259-269. [12] M. Pierre and D. Schmitt, Blow up in reaction-diffusion systems with dissipation of mass, SIAM Review 42 (1) (2000), 93-106. 28 [13] M. Pierre, T. Suzuki and Y. Yamada, Dissipative reaction-diffusion systems with quadratic growth, Indiana Univ. Math. J., 68 (2019), 291-322. Ph. Souplet, Global existence for reaction-diffusion systems with dissipation of mass and 30 [14]quadratic growth, J. Evol. Equ. , 18 (2018), 1713–1720. 32 B.Q. Tang, Global classical solutions to reaction-diffusion systems in one and two dimensions, [15]Comm. in Math. Sc., 16 (2) (2018), 411-423. Received xxxx 20xx; revised xxxx 20xx. E-mail address: michel.pierre@ens-rennes.fr
- 36 E-mail address: didier.schmitt@univ-lorraine.fr

1

2

6

7

9

11

12 13

14

15

16

17

18

19

21 22

23

24 25

26 27

29

31

33

34