



**HAL**  
open science

## High temperature patterns at the onset of seed maturation determine seed yield and quality in oilseed rape (*Brassica napus* L.) in relation to sulphur nutrition

Lethicia Magno Massuia de Almeida, Jean-Christophe Avice, Annette Morvan Bertrand, Marie H el ene Wagner, Mar ia Reyes Gonz alez-Centeno, Pierre-Louis Teissedre, Jean Jacques Bessoule, Marina Le Gu edard, Tae Hwan Kim, Alain Mollier, et al.

### ► To cite this version:

Lethicia Magno Massuia de Almeida, Jean-Christophe Avice, Annette Morvan Bertrand, Marie H el ene Wagner, Mar ia Reyes Gonz alez-Centeno, et al.. High temperature patterns at the onset of seed maturation determine seed yield and quality in oilseed rape (*Brassica napus* L.) in relation to sulphur nutrition. *Environmental and Experimental Botany*, 2021, 185, pp.1-14. 10.1016/j.envexpbot.2021.104400 . hal-03363680

**HAL Id: hal-03363680**

**<https://hal.science/hal-03363680>**

Submitted on 4 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 Original article

2 **High temperature patterns at the onset of seed maturation determine seed yield and quality in**  
3 **oilseed rape (*Brassica napus* L.) in relation to sulphur nutrition**

4

5 Lethicia Magno Massuia de Almeida<sup>1</sup>, Jean-Christophe Avice<sup>1</sup>, Annette Morvan Bertrand<sup>1</sup>, Marie  
6 H el ene Wagner<sup>2</sup>, Mar ıa Reyes Gonz alez-Centeno<sup>3,4</sup>, Pierre-Louis Teissedre<sup>3,4</sup>, Jean Jacques Bessoule<sup>5</sup>,  
7 Marina Le Gu edard<sup>5,6</sup>, Tae Hwan Kim<sup>1,7</sup>, Alain Mollier<sup>8</sup>, Sophie Brunel-Muguet<sup>1\*</sup>

8

9 1. Normandie Universit e, UNICAEN, INRAE, UMR 950 Ecophysiologie V eg etale, Agronomie et  
10 nutriments N, C, S, Esplanade de la Paix, CS14032, 14032 Caen Cedex 5, France

11 2. Station Nationale d'Essais de Semences, GEVES, 49071 Beaucauz e France

12 3. Universit e de Bordeaux, Unit e de Recherche C enologie, EA 4577, ISVV, 33882 Villenave d'Ornon,  
13 France

14 4. INRAE, USC 1366 C enologie, ISVV, 33882 Villenave d'Ornon, France

15 5. Univ. Bordeaux, CNRS, Laboratoire de Biog en ese Membranaire (LBM), UMR 5200, 71, avenue  
16 Edouard Bourlaux, 33883 Villenave d'Ornon Cedex, France

17 6. LEB Aquitaine Transfert-ADERA, 71, avenue Edouard Bourlaux, 33883 Villenave d'Ornon Cedex

18 7. Environment-Friendly Agriculture Research Center (EFARC), Department of Animal Science,  
19 Institute of Agricultural Science and Technology, College of Agriculture & Life Science, Chonnam  
20 National University, Buk-Gwangju, P.O. Box 205, Gwangju 500-600, South Korea

21 8. ISPA, Bordeaux Sciences Agro, INRAE, F-33140, Villenave d'Ornon, France

22

23 \* Corresponding author: lethicia.magno-massuia-dealmeida@unicaen.fr

24

25

26

1 **Abstract**

2 High temperatures during the crop reproductive stage impact seed yield and quality. The changing  
3 climate will require consideration of the effects of high temperature events that differ from their  
4 intensity, their duration and their frequency over the seed quality-building stages. The impact of these  
5 features deserve to be investigated at the light of induced thermo-sensitization which can lead to  
6 alleviate expected negative impacts. In our work, maturing seeds of the sulphur-demanding crop,  
7 oilseed rape, were exposed to several temperature sequences that varied in intensity, duration and  
8 frequency at the onset of seed maturation. Results-measured in seeds that were at the onset of  
9 maturation when the temperature stress occurred-indicated that (i) the longer the cumulated duration of  
10 the temperature stress, the more negatively impacted the quality criteria with decreased fatty acids  
11 (FAs) concentration, increased  $\omega 6$ :  $\omega 3$  ratio, lower seed membrane integrity and increased seed  
12 dormancy and (ii) a mild stress event prior to heat peaks had an alleviating effect on the negative  
13 impact of the later heat peaks (priming effect) on seed nitrogen, desiccation tolerance and the  
14 phytohormones involved in thermoinhibition. sulphur restriction was positive on FAs, proteins  
15 concentrations and negative on breaking dormancy. In addition, sulphur supply interfered with  
16 temperature modality, features such that positive impact of sulphur limitation on boosting oxidative  
17 response were cancelled with intense late heat peaks. This work provides insights to define  
18 thermopriming protocols in relation to the timing of quality building processes, their respective  
19 optimal temperature and adequate sulphur supply.

20

21 **Key words:** oilseed rape, seed quality, high temperature, sulphur, repeated stresses, priming, stress  
22 memory, thermotolerance.

23

24 **Running head:** Thermotolerance for seed quality in oilseed rape

25

26 **Main abbreviations:** S: sulphur; C: carbon; N: nitrogen; FA: fatty acid; UFA: unsaturated FA; SFA:  
27 saturated FA; ABA: abscisic acid; GA3: gibberellic acid; IAA: indole-3-acetic acid; SA: salicylic acid;

1 SSP: seed storage protein; ROS: reactive oxygen species; TSW: thousand seed weight; DW: dry  
2 weight.

### 3 **Introduction**

4 Evidence for increased frequency of stress events such as heat waves has been observed over recent  
5 decades. Based on the last Intergovernmental Panel on Climate Change (IPCC) report (Hoegh-  
6 Guldberg et al., 2018), heat waves are expected to become more frequent, to last longer and to increase  
7 in intensity during the reproductive phase of economically important crops (Christidis et al., 2015;  
8 Trnka et al., 2014). These new climatic patterns have led to attempts to decipher crop behavior and  
9 final performance in the light of recurring stresses. While the effects of extreme/mild environmental  
10 stresses have been widely investigated from molecular to whole plant levels (Kotak et al., 2007;  
11 Ohama et al., 2016; Wahid et al., 2007), far fewer studies have tackled the issue of understanding the  
12 effects of their recurrence throughout the crop season. Indeed, the overall magnitude of the plant  
13 response to successive stresses might not match the effects induced by individual stressing events  
14 because (i) the first stress triggers physiological and metabolic adjustments that bring the plant to a  
15 modified status (including phenology) when the stress recurs, which leads to different responses to the  
16 second stress and (ii) a mild stress prior to further similar stresses can induce stress memory that lasts  
17 the duration of the crop season (i.e. intra-generational memory, Ding, Fromm & Avramova 2012) and  
18 is sometimes transmitted to offspring (i.e. transgenerational memory, Molinier, Ries, Zipfel & Hohn  
19 2006; Wang *et al.* 2016; Hatzig, Nuppenau, Snowdon & Schiebl 2018; Kinoshita & Seki 2014; Crisp,  
20 Ganguly, Eichten, Borevitz & Pogson 2016; Kumar 2018). Stress memory is defined as the process of  
21 storage and retrieval of information acquired during an initial exposure to stress (Crisp et al., 2016;  
22 “Stress priming, memory, and signalling in plants,” 2019). This information acts as a priming process  
23 with beneficial effects when the stress recurs and it can lead to earlier, more rapid, intense, and  
24 sensitive responses that help plants to acclimate in changing environments (Kinoshita and Seki, 2014).  
25 Underlying mechanisms include epigenetic regulation, transcriptional priming, primed conformation  
26 of proteins, and/or specific hormonal or metabolic signatures. (Groot et al., 2016; Hatzig et al., 2018;  
27 Molinier et al., 2006; Wang et al., 2016) This means that as the climate changes, the effects of repeated

1 stresses should be harnessed as a crop improvement strategy because this approach has promise for  
2 inducing acclimation to heat stress (Wang and Liiang, 2017).

3 In winter oleaginous crops, the reproductive phase occurs during spring, which might expose  
4 flowering, grain filling and grain maturation to high temperature events. Due to its indeterminate  
5 growth, winter oilseed rape plants display flowers and growing pods in different proportions  
6 throughout the reproductive phase. Consequently, heat stress can impact reproductive organs by  
7 limiting number and size, which in turn leads modification to carbon (C) partitioning in favor of  
8 already developed organs that have passed the sensitive stage, and this makes analysis of the direct  
9 effects of heat stress on the organs more complex (Guilioni et al., 1997). While heat stress at flowering  
10 limits pollination (Sage et al., 2015) and/or induces early pod abortion resulting in yield losses in  
11 oilseed rape (Morrison and Stewart, 2002; Young et al., 2004), heat stress that occurs during seed  
12 filling and maturation affects seed storage compounds quantitatively and qualitatively, leading to seed  
13 quality alteration. Few studies have reported the effects on seed quality in oilseed rape of repeated heat  
14 stress events that might be erratic and fluctuate as predicted in climate change models in field  
15 conditions (Deng & Scarth 1998; Baux *et al.* 2013).

16 Seed quality encompasses a range of criteria related to nutritional and physiological characteristics  
17 (i.e. germination behaviors and storage capacity). In winter oilseed rape, oil content, fatty acid (FA)  
18 profiles, and protein content are major nutritional criteria for edible oil and cakes used in human and  
19 animal consumption, respectively. Its oil contains higher unsaturated FA (UFA) content than other oil  
20 crops (sunflower, soybean), which makes it a healthy edible oil for human consumption (Aguirrezábal  
21 et al., 2015). Contrasting variations have been observed in oil content according to the temperature  
22 intensity, to the timing of stress exposure and to the pools of seeds analyzed (main stem vs. bulk) (e.g.  
23 increased in Brunel-Muguet *et al.* (2015); decreased in Canvin (1965); Aksouh *et al.* (2001); Aksouh-  
24 Harradj *et al.* (2006)). High temperatures are known to induce decreases in poly-UFAs in favor of  
25 saturated FAs (SFA, mainly C16:0 and C18:0) and mono-UFAs, and increases in the  $\omega 6:\omega 3$  ratio (i.e.  
26 C18:2/C18:3 ratio), as a result of temperature-triggered impairment of desaturase enzyme activity i.e.  
27 oleic and linoleic desaturases (Aksouh-Harradj et al., 2006; Baux et al., 2013, 2008; Brunel-Muguet et

1 al., 2015; Gauthier et al., 2017; Schulte et al., 2013). By contrast, seed nitrogen (N) and protein  
2 concentrations in the oil-free meal is usually negatively correlated with total oil content (Aksouh-  
3 Harradj et al., 2006; Aksouh et al., 2001) as observed in other oil crops (in soybean, N concentration  
4 Chebrolu *et al.* (2016); protein concentration, Dornbos & Mullen (1992)). Additionally, other seed  
5 characteristics related to physiological quality, i.e. seed storage capacity and germination behavior,  
6 were investigated but not to any great extent (Brunel-Muguet et al., 2015). A drastic degradation of  
7 seed storage capacity has been observed using seed conductivity and the ratio of soluble sugars  
8 ([stachyose and raffinose]:sucrose), abscisic acid (ABA) and gibberellic acid (GA3) as proxies (Bailly  
9 et al., 2001; Brunel-Muguet et al., 2015) in seeds from long-term heat-stressed mother plants. Other  
10 phytohormones were shown to be involved in the control of secondary dormancy, defined as failure in  
11 the germination process of mature and non-dormant seeds under adverse conditions (Pekrun et al.,  
12 1997). Recent studies have highlighted the role of indole-3-acetic acid (IAA), whose concentrations  
13 increase in dormancy-induced seeds (Liu et al., 2019; Shu et al., 2016; Tuan et al., 2019). Although  
14 several studies have reported that salicylic acid (SA) enhanced germination in *Arabidopsis* seeds by  
15 reducing oxidative damage (Chitnis et al., 2014; Lee and Park, 2010) and inhibited germination  
16 because of higher oxidative stress (Xie et al., 2007).

17 In Brassica species, sulphur (S) nutrition determines yield components and seed quality because of  
18 their high S requirements throughout the crop cycle (Brunel-Muguet et al., 2015; D'Hooghe et al.,  
19 2014). In addition to its well-known implication in the synthesis and signaling of stress tolerance-  
20 controlling phytohormones (Hasanuzzaman et al., 2018), S might be involved in the acquisition of  
21 thermotolerance mediated by epigenetic regulation (Bokszczanin et al., 2013). This is based on  
22 evidence for S involvement in DNA methylation (through the role of S-adenosylmethionine as a donor  
23 of methyl groups, Meng *et al.* 2018), one of the key epigenetic markers that supports stress memory,  
24 thus making the analysis of S supply relevant in the context of epigenetic memory.

25 In our study we focused on the effects of high spring temperatures on seeds at the onset of maturation  
26 to deepen our knowledge of this seed quality-determining stage in relation to S nutrition. Our  
27 assumptions were that the effects of high temperature at advanced seed filling can greatly vary

1 depending on whether plants are exposed to a mild heat stress event that primes them to withstand  
2 later heat peaks and that S nutrition might impact the ability of the plants to endure heat stress.  
3 Overall, our experimental design addresses the following questions: (i) what are the quantitative  
4 effects of different high temperature sequences applied at advanced seed filling/onset of seed  
5 maturation on seed yield, quality criteria and stress response indicators? (ii) is there any beneficial  
6 effect from a mild stress event that occurs prior to later repeated intense heat peaks? (iii) to what  
7 extent do the effects of successive high temperature events applied to maturing pods differ from the  
8 effect of individual events? (iv) what are the underlying defense pathways triggered by temperature  
9 stress? and (v) how does sulphur nutrition impact heat stress responses through acquisition of  
10 thermotolerance?

11

## 12 **Materials and Methods**

### 13 *Experimental treatments and growth conditions*

14 Seeds of *Brassica napus* L. (cv. Aviso) were germinated in vermiculite in October 2016 under  
15 greenhouse conditions. After five weeks the seedlings were transplanted into pots containing perlite  
16 and vermiculite (2:1, v/v) for seven weeks and seedlings were grown as described in Poisson *et al.*  
17 (2019). Afterwards, seedlings were subjected to a 12-week period of vernalization in a cold chamber  
18 (standard model, Froid & Mesures, Beaucouzé, France) maintained at 4°C (night) and 8°C (day) with  
19 artificial light during the day (10h day/14h night) and supplied with a 25% Hoagland solution without  
20 sulphur, to prevent the plants from building substantial S reserves which would later impact their  
21 nutritional status when applying the contrasting sulphur supplies.

22 Then, the plants were transferred into the greenhouse (Caen, France, 49°11'09 N, 0°21'32 W) and  
23 subjected to a thermoperiod of 20°C (day) and 15°C (night) without additional light. The plants were  
24 manually provided with two N applications with an NH<sub>4</sub>NO<sub>3</sub> solution: 100 kg N/ha at the end of  
25 vernalization (Growing Stage 30 (GS30), stem elongation, Lancashire *et al.* 1991) and 50 kg N/ha at  
26 early flowering (GS60, bud formation) assuming a plant density of 40 plants.m<sup>-2</sup>. The two contrasting  
27 S supplies i.e. High Sulphur and Low Sulphur were manually applied at the end of vernalization

1 (GS30) as usually provided in the field. Plants were supplied with a solution of  $\text{MgSO}_4$  containing 75  
2 kg  $\text{SO}_3$ /ha (High Sulphur) and 25 kg  $\text{SO}_3$ /ha (Low Sulphur), which represent, respectively, 100% and  
3 33% of the conventional supply.

4 Four temperature modalities (Temp-modalities) were applied to plants at stage GS72 (i.e. 20% of the  
5 pods having reached their maximum size) for 17 days. The Temp-modalities were designed to assess  
6 the effects of a mild temperature event prior to a more intense temperature event including daily heat  
7 peaks (Figure 1). In our complete design, we tested five Temp-modalities which included a Temp-  
8 control modality that is natural thermoperiod conditions in the greenhouse (Figure 1a). The other four  
9 Temp-modalities were the following: (i) the early mild stress modality (Figure 1b) composed of 5  
10 early full days under mild warming [i.e.  $25.3^\circ\text{C} \pm 1.8$  (day, 16h) /  $21.7^\circ\text{C} \pm 0.7$  (night, 8h)] and  
11 followed by 12 days under natural thermoperiod (mean, maximum and minimum temperatures over  
12 the natural thermoperiod being  $15.7^\circ\text{C} \pm 2.7$ ,  $28.4^\circ\text{C}$  and  $11.5^\circ\text{C}$ , respectively), (ii) the 3 late heat  
13 peaks modality composed of 14 days under natural thermoperiod followed by 3 days under mild  
14 warming with a daily heat peak applied for 5 hours (i.e.  $31.4^\circ\text{C} \pm 1.7$  between 11 am till 4 pm, Figure  
15 1c), (iii) the 4 late heat peaks modality composed of 10 days under natural thermoperiod followed by 7  
16 days under mild warming with daily heat peaks on days 12, 15, 16 and 17 (Figure 1d), and (iv) the  
17 priming modality composed of 5 days under mild warming followed by 5 days under natural  
18 thermoperiod and eventually 7 days under mild warming with daily heat peaks on days 12, 15, 16 and  
19 17 (Figure 1e).

20





1 and monitored, according to the experimental design (Figure 1). The temperature intensities were  
2 chosen according to previous studies that used similar ranges to mimic mild and intense temperature  
3 treatments under controlled conditions (Aksouh-Harradj et al., 2006). Temperatures were recorded  
4 hourly with temperature probes (105T Campbell, Campbell Scientific Ltd., Leicestershire, UK).  
5 Throughout the 17 days, the incident Photosynthetically Active Radiation (PAR<sub>i</sub>) values was recorded  
6 (every 15 minutes) in the greenhouse and processed to calculate the daily PAR<sub>i</sub> values (Supplemental  
7 Data, Table S1). No substantial difference in daily PAR<sub>i</sub> were observed between the greenhouse units  
8 with the natural thermoperiod conditions and the higher temperature conditions (mild warming and  
9 heat peaks).

10 Because mixed-age pods were present throughout the temperature sequences, on the day before the  
11 beginning of the temperature modalities exposure we have labeled each branch of the plants to identify  
12 the two categories of pods: (i) pods whose length was above 5 cm (pods<sub>L≥5cm</sub>) and (ii) pods whose  
13 length was shorter than 5 cm (pods<sub>L<5cm</sub>). Indeed, preliminary experiments (Supplemental Data, Table  
14 S2) indicated that pods<sub>L≥5cm</sub> (or longer) contain seeds that have reached at least half their final fresh  
15 weight (about 55%), which coincides with advanced seed-filling development and the onset of seed  
16 maturation (Borisjuk et al., 2013). When the pods started desiccating, they were carefully and  
17 individually wrapped with plastic pouches to avoid seed dispersal and the mixing of seeds between the  
18 pod categories.

19

#### 20 ***Seed yield and components***

21 At maturity, the seeds from the two categories of pods were weighed after freeze-drying for dry weight  
22 (DW) measurements. To determine the individual seed weight (Thousand Seed Weight (TSW)), we  
23 weighed and photographed seeds from both pod categories so as to score their number using image  
24 analysis algorithms (ImageJ Software, Schindelin et al. 2012).

25

#### 26 ***Biochemical characteristics of seeds from pods<sub>L≥5cm</sub>***

1 In the following sections, biochemical characteristics of each individual plant ( $n=4$ , and  $n=3$  only for  
2 hormones) were measured solely on seeds from pods<sub>L $\geq$ 5cm</sub>.

3

#### 4 *Seed carbon, nitrogen and sulphur concentrations*

5 Seeds of each individual plant were pooled and dried for 48h at 50°C. The dried seeds were ground  
6 manually and the resulting powder (around 3 mg per sample) was placed into tin capsules for analysis.

7 The percentage of total carbon, nitrogen, and sulphur per mg DW of the seeds were determined with a  
8 C/N/S analyzer (EA3000, Euro Vector, Milan, Italy) linked to a continuous flow isotope mass  
9 spectrometer (IRMS, Isoprime, GV Instrument, Manchester, UK).

10

#### 11 *Seed fatty acid concentration*

12 Oil and fatty acid profile contents (C16:0, C18:0, C18:1, C18:2, C18:3, C20:0, C20:1 and C22:1) were  
13 determined as described in Marchand *et al.* (2016). Approximately 200 mg of seeds from each plant  
14 were suspended in 1 mL of methanol/toluene/H<sub>2</sub>SO<sub>4</sub> solution (100:20:2.5; v/v) containing C17:0 as  
15 internal standard (5  $\mu\text{g mL}^{-1}$ ), overnight at 85°C for transmethylation. After cooling, 500  $\mu\text{L}$  of hexane  
16 was added and the hexane phase containing the resulting fatty acid methyl esters (FAMES) was  
17 recovered for gas chromatography analysis combined with flame ionization detection (GC-FID). The  
18 FAMES (1  $\mu\text{L}$ ) were injected into an Agilent 7890 gas chromatograph equipped with a Carbowax  
19 column (15m by 0.53mm, 1.2 m) (Alltech Associates, Deerfield, IL) and FID system. FAMES were  
20 identified by comparing their retention times with those of commercial standards (Sigma, St. Louis,  
21 MO) and quantified using ChemStation (Agilent) to calculate the peak areas.

22

#### 23 *Seed storage protein concentration*

24 Protein analysis was performed in two steps: (i) 20 g of ground seeds from plants of each treatment  
25 was previously stored at -20°C before protein extraction and (ii) the Bradford assay. The Bradford  
26 assay is based on the use of a standard range with several dilutions of the reference protein (Bovine  
27 Serum Albumin, BSA). Thus, after dilution (by 10) the samples were placed in a microplate and the

1 measure is made at 570 nm after addition of the Bradford reagent. The calibration straight line allowed  
2 the total protein concentrations to be calculated by considering the mass and volume of the sample.  
3 The detailed steps for extraction and the Bradford assays are described for leaves in Akmouche *et al.*  
4 (2019).

5

#### 6 *Seed soluble sugar concentrations*

7 Soluble sugars were extracted from 50 mg lyophilized and ground seeds, with 1 mL methanol/water  
8 (80:20, v/v) and 40 µL melicitose used as the internal sugar standard. Glucose, fructose, sucrose,  
9 raffinose, and stachyose contents were quantified using High Performance Liquid Chromatography  
10 (HPLC) on a cation exchange column (Sugar-PAK, 300 X 6.5 mm, Millipore Waters, Milford, MA,  
11 USA) eluted at 0.5 mL min<sup>-1</sup> and 85 °C with 0.1 mM Ca-EDTA in water and quantified using a  
12 refractive index detector (2410 Differential Refractometer, Millipore Waters, Waters Corporation,  
13 MA, USA), according to Brunel-Muguet *et al.* (2015). The [raffinose+stachyose]:sucrose ratio was  
14 used as an indicator of seed drying tolerance i.e. the higher the value, the more tolerant the seed is to  
15 desiccation (Bailly *et al.*, 2001).

16

#### 17 *Analysis of stress signaling and seed dormancy-related phytohormones*

18 For each treatment, 50 mg of ground seeds were used to quantify 2-*cis*, 4-*trans*-abscisic acid,  
19 gibberellic acid, indole-3-acetic acid and salicylic acid. Previously freeze-dried samples were mixed  
20 with 500 µL of extraction solvent [2-propanol/H<sub>2</sub>O/ concentrated HCl (2:1:0.002, v/v/v)] and then  
21 analyzed by HPLC-MS as described in (Pan *et al.*, 2010). Because the dynamics of ABA and GA3  
22 controls the balance between dormancy and germination, the ABA:GA3 ratio was used as a proxy for  
23 seed dormancy under stress condition (Debeaujon and Koornneef, 2000; Finkelstein, 2013).

24

#### 25 *Seed conductivity measurements*

26 Conductivity measurements were performed according to Brunel-Muguet *et al.* (2015). For each  
27 treatment, measurements with the electrolytes were made with 20 pre-weighed seeds. At room

1 temperature (20°C), the seeds were placed in 5 mL of ultrapure water for 16 hours. Conductivity was  
2 measured with a portable electrochemical analyzer (Consort C931, UK).

3

#### 4 *Determination of seed total phenolic content and antioxidant capacity*

5 Prior to the extraction, seeds from plants of each treatment were oven-dried at 60°C for at least 48  
6 hours and then ground. The extraction procedure was adapted from Szydłowska-Czerniak, Amarowicz  
7 & Szłyk (2010) in terms of solvent, extraction time and sample:solvent ratio. Seeds were extracted  
8 with methanol:H<sub>2</sub>O (50:50, v/v) at a ratio of 20:1 (mg/mL) by using mechanical stirring for 1 h at 25  
9 °C. Total phenolic content was spectrophotometrically determined with a modified Folin-Ciocalteu  
10 method (González-Centeno et al., 2015) and was expressed as the mean of six determinations of  
11 sinapic acid in mg equivalents per DW g of seeds. To measure the antioxidant capacity, seed extracts  
12 were diluted at a ratio 1/4 with methanol:H<sub>2</sub>O (50:50, v/v) according to the methodology previously  
13 described by González-Centeno *et al.* (2012). ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic  
14 acid)] assay values were expressed as mg of Trolox equivalent antioxidant capacity per g of seeds  
15 DW.

16

#### 17 *Statistical analyses*

18 Two-way ANOVAs to test temperature (Temp) stress and S nutrition effects, both considered as  
19 independent factors, were performed on the measured variables. The Temp-modalities, sulphur  
20 condition and Temp x Sulphur interaction effects were analyzed using R software (version 4.0.2).  
21 Prior to the ANOVAs, residues independency and normality, and homogeneity of variances were  
22 previously tested (Dubin-Watson, Shapiro-Wilk, Bartlett and Levene tests respectively). ANOVAs  
23 tables presented the means for each Temp-modality (gathering the values of both sulphur supplies) and  
24 for each sulphur supplies (gathering the values of the five Temp-conditions including the Temp-  
25 control) (Tables 1, 2 and 3). Mean comparison tests were performed differently according to whether  
26 interaction effects were detected or not. When no interaction effects were observed, Tukey tests were  
27 performed amongst the five temperature modalities (Tables 1, 2 and 3). Otherwise, when interaction

1 effects were observed, comparison amongst the 10 treatments (crossing the five temperature  
2 modalities and the two sulphur conditions) were performed as one-way ANOVA and considering 10  
3 independent treatments for the Tukey tests (Figures 2, 3 and 4). The last table in Supplemental data  
4 (Table S3) summarizes the mean  $\pm$  se of the all measured variables for each of the 10 treatments in  
5 order to provide reference values in a given combination “sulphur supply/Temperature modality”.

6

## 7 **Results**

### 8 *Plant growth and yield components*

9 Table 1 displays yield and yield components i.e. seed number and TSW for the overall seeds (Total)  
10 and the two categories of seeds collected separately (from pods<sub>L $\geq$ 5cm</sub> and pods<sub>L<5cm</sub>). No significant  
11 Temp x Sulphur interaction effect was observed for any of the yield related-measured variables. The  
12 total seed yield (including seeds from both categories of pods) ranged from 8.2 to 9.8 g plant<sup>-1</sup>, with no  
13 significant sulphur effect. Decreases in the total seed yield were observed for all the four Temp-  
14 modalities, being significant only in the priming modality (-16.3% relative to Temp-control) . These  
15 decreases were due mostly to the seed yield from pods<sub>L<5cm</sub>, (though not significant) since seed yield  
16 from pods<sub>L $\geq$ 5cm</sub> remained almost unchanged compared to the Temp-control plants. The total seed  
17 numbers were not impacted by the Temp-modality but a slight sulphur effect was observed (p<0.05),  
18 with Low Sulphur values being 5% lower than High Sulphur values on average. The TSWs from  
19 pods<sub>L<5cm</sub> ranged from 3.0 g (3 late heat peaks modality) to 3.4 g (Temp-control), whereas the  
20 minimum and maximum values from pods<sub>L $\geq$ 5cm</sub> were 3.2 g (priming modality) and 3.7 g (3 late heat  
21 peaks modality). Under both sulphur supplies, no Temp-modality effects on the TSW were observed  
22 in either pod category. These results confirmed that (i) seeds from pods<sub>L $\geq$ 5cm</sub> had reached the onset of  
23 seed maturation and (ii) seeds developing in pods<sub>L<5cm</sub> or seeds from fertilized flowers that developed  
24 after the temperature sequences all benefited from increased sink strength, due to young pod abortion  
25 or reductions in late flowering.

26

### 27 *Nutritional seed quality criteria*

1 *Seed C, N and S concentrations*

2 Table 2 displays C, N and S seed concentrations. Similar to yield components, no Temp x Sulphur  
3 interaction effect was observed. The seed C concentration was impacted by Temp-modalities  
4 ( $p < 0.001$ ) but no sulphur effect was observed. The priming modality displayed the lowest value (-  
5 5.4% from Temp-control) whereas the Temp-control displayed the highest value. Similar to C, the  
6 seed N concentration was only affected by Temp-modalities ( $p < 0.001$ ). Under both sulphur  
7 conditions, extreme N values were observed for the priming modality, which had the highest ranking  
8 (+21.9 % compared to Temp-control). These observations showed that the early mild stress had a  
9 positive effect over the late heat peaks, which suggested an alleviating effect. As expected, the seed S  
10 concentration was significantly impacted by the sulphur supply ( $p < 0.001$ ), with Low Sulphur values  
11 being 32% lower than High Sulphur values on average. While no significant differences were  
12 observed among the Temp-modalities, the S in the priming modality was slightly higher than in Temp-  
13 control (+19.9%).

14

15 *Fatty Acids concentrations and profiles*


16 Total fatty acids, saturated FAs and unsaturated FAs concentrations and the  $\omega 6:\omega 3$  ratios are displayed  
17 in Table 2 and in Figure 2. For total FAs concentration, no Temp x Sulphur interaction effect was  
18 observed on FAs concentration. Effects of Temp-modalities were observed ( $p < 0.01$ ) with values  
19 ranging from 33% to 41% of DW for priming and 3 late heat peaks modalities respectively (Table 2).  
20 Values under Low Sulphur were significantly higher ( $p < 0.001$ ) on average (41 %DW) than values  
21 under High Sulphur (35 %DW). Temp-modality effect was observed in both SFAs and UFAs  
22 concentrations, with a significant effect of Temp x Sulphur interaction only observed on SFAs  
23 ( $p < 0.01$ ) (Table 2, Figure 2). The SFAs (including C16:0, C18:0, C20:0, C22:0) concentrations ranged  
24 from 3.0% to 3.6%DW and the UFAs (including C16:1, C18:1, C18:2, C18:3, C20:1) concentrations  
25 ranged from 30% to 38%DW. For both SFAs and UFAs concentrations, the priming modality and the  
26 3 late heat peaks displayed the lowest and highest values respectively.

1 A highly significant sulphur effect was observed in both SFAs and UFAs concentrations, with values  
2 under Low Sulphur (3.6 %DW of SFAs and 38% DW of UFAs) being significantly higher on average  
3 than values under High Sulphur (3.1 %DW of SFAs and 32% DW of UFAs) ( $p < 0.001$ ) (Table 2,  
4 Figure 2). Overall, the priming modality had the greatest impact on decreasing FAs concentrations, as  
5 a result of the lowest decreases in SFAs and UFAs concentrations, and the early mild stress event did  
6 not alleviate the negative effects of later heat peaks on total FAs concentration. In addition, the 4 late  
7 heat peaks sequence had a greater impact than the 3 late heat peaks sequence on total FAs, SFAs and  
8 UFAs meaning the more intense events, the more impacting.

9 The  $\omega 6:\omega 3$  ratio (i.e. C18:2/C18:3 ratio) is commonly used as an indicator of the edible quality of  
10 vegetable oils. This ratio was significantly impacted by the Temp-modalities, but no significant  
11 Sulphur nor Temp x Sulphur interaction effects were observed (Table 2). The highest and lowest ratios  
12 were observed on the priming and early mild stress modalities respectively (+9.3 and -8.8% compared  
13 to the Temp-control). In addition, the temperature effect on 3 late heat peaks modality was less  
14 negative than the early mild stress modality (Table 2). These results indicated that the number of  
15 desaturations decreased with greater duration of stress exposure (priming modality) and intensity and  
16 earliness of the heat stress event (4 late heat peaks modality) under both sulphur conditions.

17





1

2 **Figure 2:** Boxplot and ANOVA table for saturated FAs (SFAs) measured on seeds from pods  $L_{\geq 5cm}$   
 3 under the 10 treatments (crossing temperature modalities and sulphur conditions). SFAs displayed  
 4 highly significant interaction (Temp x Sulphur) effects. Letters indicate the ranking amongst the 10  
 5 treatments. P-values and levels of significance are given for Temp, Sulphur and Temp x Sulphur  
 6 interaction effects. Levels of significance: ns non-significant,  $p < 0.05^*$ ,  $p < 0.01^{**}$ ,  $p < 0.001^{***}$ . SFAs:  
 7 saturated fatty acids; 3\_LHP: 3 late heat peaks modality; 4\_LHP: 4 late heat peaks modality.

8

9 *Seed storage protein concentrations*

10 The seed storage protein (SSP) concentration ranged from 123 mg/g DW to 168 mg/g DW, with  
 11 values under High Sulphur condition being significantly lower on average (106 mg/g DW) than values  
 12 under Low Sulphur condition (185 mg/g DW) (Table 2,  $p < 0.001$ ). In contrast, no significant effects of  
 13 Temp-modalities on SSP were observed although the lowest values were observed under the late heat  
 14 peaks modalities (Table 2). The increase under the early mild stress modality might result from  
 15 reductions in the number of growing sinks due to pod abortion, reduced seed filling and/or impaired  
 16 pollination.

17

1 ***Physiological seed quality-related criteria***

2 *The soluble sugar composition as an indicator of desiccation tolerance acquisition*

3 Table 3 presents the concentration of the main soluble sugar in seeds i.e. sucrose, raffinose and  
4 stachyose. For the three main soluble sugars, only temperature effects were observed ( $p < 0.001$ ). All  
5 the Temp-modalities led to decreased soluble sugar concentrations under both sulphur conditions.  
6 Sucrose, raffinose and stachyose concentrations were the highest in the Temp-control and the lowest  
7 in the priming and early mild stress modalities. Under the priming modality, sucrose, raffinose and  
8 stachyose concentrations were respectively 54%, 55% and 52% lower than the Temp-control.  
9 Similarly, under the early mild stress modality these concentrations were respectively 59%, 55% and  
10 56% lower than under the Temp-control. These results indicated that the early mild stress had a  
11 negative impact on the sugar soluble concentrations. The [raffinose+stachyose]:sucrose ratio, used as a  
12 proxy of seed desiccation tolerance, ranged from 0.35 (4 late heat peaks modality) to 0.46 (early mild  
13 stress modality) without significant Sulphur effect (Table 3). These results indicate that whatever the  
14 sulphur supply, the 4 late heat peaks modality was the most detrimental to acquisition of seed  
15 desiccation tolerance. However, these data also highlighted that the prior event of mild stress  
16 alleviated the negative effects of late heat peaks since the value under the priming modality was higher  
17 than under the early mild stress modality.

18

19 *Seed conductivity as indicator of membrane permeability*

20 Seed conductivity values ranged from 0.8  $\mu\text{S}/\text{mg DW}$  (Temp-control) to 2.9  $\mu\text{S}/\text{mg DW}$  (priming  
21 modality), with a highly significant temperature effect ( $p < 0.001$ , Table 3). No effect of Sulphur nor  
22 Temp x Sulphur interaction effects were observed. Extreme values were observed in Temp-control  
23 (the lowest) and in priming modality (the highest, +262% compared to Temp-control). The other three  
24 Temp-modalities (early mild stress, 3 and 4 late heat peaks modalities) led to similar values than under  
25 the Temp-control. These results indicate that seed conductivity is highly responsive to the duration of  
26 the stressing period, because seeds from temperature modalities displaying the shortest numbers of  
27 days where temperature was higher than the control temperature, were the least negatively impacted.

1


2 *Phytohormone changes as indicators of temperature-induced seed dormancy*

3 Figure 3 displays the results of the main hormones involved with dormancy in seeds (i.e. IAA, SA and  
4 ABA:GA3 ratio). Temp-modality effects, as well as Sulphur and Temp x Sulphur interaction effects  
5 were observed for the three variables with highly significance ( $p<0.001$ ). The ABA:GA3 ratio ranged  
6 from 0.51 (early mild stress modality) to 0.70 (4 late heat peaks modality), with significant lower  
7 values in High Sulphur (0.47 on average) than in Low Sulphur condition (0.78 on average, that is 66%  
8 higher). The interaction effects pointed out that according to the sulphur condition, the Temp-modality  
9 effect was different i.e. while the early mild stress modality and the 3 late heat peaks modality under  
10 high sulphur were ranked the same, these Temp-modalities under Low Sulphur were highly  
11 contrasting (with values being much higher under the 3 late heat peaks modality) (Figure 3).

12 A high ABA:GA3 ratio indicates increased secondary seed dormancy, which is induced by  
13 thermoinhibition. As expected, under high temperature this ratio increased, but under High Sulphur  
14 condition, the modality that displayed the more days with temperature above the control temperature  
15 (priming modality) was the most negatively impacted (highest value), whereas under Low Sulphur  
16 condition, the late heat peaks events (4 or 3) induced the highest ratios. In addition, under Low  
17 Sulphur condition, the early mild stress tended to alleviate the negative effects of late heat peaks by  
18 lowering the ratio (lower values in priming and early mild stress modalities than in late heat peaks  
19 modalities).

20 Seed concentrations of IAA and SA were measured to investigate their variation under the different  
21 temperature sequences and sulphur supplies. Consistent with the ABA:GA3 ratio, high concentrations  
22 of IAA and SA were observed under high temperatures. IAA and SA ranged from 147 ng/g DW to  
23 330 ng/g DW and 50 ng/g DW to 109 ng/g DW respectively, with Temp-control displaying the lowest  
24 values and 4 late heat peaks modality displaying the highest values for both measured variables  
25 (Figure 3). Both hormones concentrations were significantly lower in Low Sulphur than in High  
26 Sulphur condition, with High Sulphur values in IAA being 17% higher than Low Sulphur and values  
27 in SA being 9% higher than Low Sulphur. By contrast to the ABA:GA3 ratio, Sulphur limitation

1 decreased IAA concentrations except under Temp-control and under the early mild stress modalities  
2 with values being respectively 16% and 73% higher in Low Sulphur than in High Sulphur (Figure 3).  
3 Under High Sulphur condition, the early mild stress event prior to the 4 late heat peaks (priming  
4 modality) led to alleviate the strong increase in IAA concentration observed in the late heat peaks  
5 solely (4 late heat peaks modality), thus suggesting an effective priming effect. The interaction effects  
6 pointed out that according to the sulphur condition, the Temp-modality effect was different i.e. while  
7 the priming modality led to higher value than under the early mild stress modality under High Sulphur,  
8 it led to lower value than the early mild stress modality under Low Sulphur (Figure 3).  
9 As observed for IAA concentrations, SA concentrations decreased under Low Sulphur except under  
10 the Temp-control and under the early mild stress modalities. Similar to IAA concentrations, the early  
11 mild stress event prior to the 4 late heat peaks (Priming modality) led to alleviate the strong increase in  
12 SA concentration observed under the 4 late heat peaks solely (4 late heat peaks modality), although  
13 this was only observed under the High Sulphur condition.



1

Variable	Temp effect	Sulphur effect	Temp x Sulphur
SA	0.000 ***	0.000 ***	0.000 ***
IAA	0.000 ***	0.000 ***	0.000 ***
ABA:GA3	0.000 ***	0.000 ***	0.000 ***

2

3 **Figure 3:** Boxplots and ANOVA table for seed physiological quality variables measured on seeds  
 4 from pods  $\geq 5\text{cm}$  under the 10 treatments (crossing temperature modalities and sulphur conditions).  
 5 Hormones (SA, IAA) concentration and ABA:GA3 ratio displayed highly significant interaction

1 (Temp x Sulphur) effects. For each variable, letters indicate the ranking amongst the 10 treatments. P-  
2 values and levels of significance are given for Temp, Sulphur and Temp x Sulphur interaction effects.  
3 Levels of significance: ns non-significant,  $p < 0.05^*$ ,  $p < 0.01^{**}$ ,  $p < 0.001^{***}$ . 3\_LHP: 3 late heat peaks  
4 modality; 4\_LHP: 4 late heat peaks modality; SA: salicylic acid; IAA: indole-3-acetic acid; ABA:  
5 abscisic acid; GA3: gibberellic acid.


6

### 7 ***Total phenolic content and antioxidant capacity in seeds as stress response indicators***

8 Figure 4 displays phenolic concentrations and the antioxidant activity (ABST values) measured in  
9 mature seeds from pods<sub>L≥5cm</sub>. Phenolic concentrations ranged from 3.6 to 4.2 mg SA/g seed DW with  
10 no significant effect of S supply. By contrast, significant Temp-modality ( $p < 0.01$ ) and Temp x  
11 Sulphur interaction ( $p < 0.05$ ) effects were observed (Figure 4). Under High Sulphur condition,  
12 extreme seed phenolic concentrations were observed under the early mild stress modality (+2%  
13 compared to Temp-control in High Sulphur) and under the 4 late heat peaks modality (-12% compared  
14 to Temp-control in High Sulphur) whereas under Low Sulphur, extreme values were observed under  
15 the early mild stress modality (+26% compared to Temp-control in Low Sulphur) and 3 late heat peaks  
16 modality (similar value to the Temp-control in Low Sulphur). These results illustrate the Temp x  
17 Sulphur interactions and the benefits of Sulphur restriction in specific temperature modalities to boost  
18 phenolic concentrations.

19 The antioxidant capacity measured with the ABTS assays revealed Temp-modality effects ( $p < 0.01$ ),  
20 Sulphur effects ( $p < 0.05$ ) and Temp x Sulphur interaction effects ( $p < 0.05$ ). Temp-modality rankings  
21 were similar to those observed for seed phenolic concentrations with extreme values being observed in  
22 the early mild stress modality (highest values) and late heat peaks modalities (3 or 4, lowest values)  
23 (Figure 4). Overall, under both Sulphur conditions, increased antioxidant capacities were observed in  
24 the early mild stress modality while decreased antioxidant capacities (or levels similar to the Temp-  
25 control) were observed when the stress occurred later, and despite the number of heat peaks. In  
26 addition, while lower oxidative response than the Temp-control were observed in the priming modality  
27 under High Sulphur conditions, a higher response was observed under Low Sulphur (Figure 4). The

1 results also indicated that the early mild stress event prior to the 4 late heat peaks (priming modality)  
2 were beneficial to the sharp decrease in ABTS values observed under the 4 late heat peaks alone, both  
3 Sulphur conditions. Regarding the effects of Sulphur conditions, limitation led to decreased  
4 antioxidant capacity in the seeds with mean values being 17.9 mg Trolox/g under High Sulphur and  
5 16.8 mg Trolox/g under Low Sulphur (i.e. -6%) which suggest a crucial role for the Sulphur supply in  
6 favoring oxidative responses, not only under the Temp-control condition but also in the 3 late heat  
7 peaks modality (Figure 4). Eventually, the Temp x Sulphur interaction effects were observed because  
8 Sulphur conditions modified the ranking of the temperature modalities compared the Temp-control,  
9 i.e. under High Sulphur ABTS values were lower in the priming and 4 late heat peaks modalities than  
10 the Temp-control but higher under Low Sulphur. Taking into account the values of all the treatments,  
11 antioxidant capacities of the seeds were positively correlated to both phenolic concentration ( $r=0.93$ ,  
12  $p<0.001$ ) and the [raffinose+stachyose]:sucrose ratio ( $r=0.64$ ,  $p<0.05$ ) (Supplemental Data, Figure  
13 S1).



Variable	Temp effect	Sulphur effect	Temp x Sulphur
Polyphenols	0.002 **	0.603	0,021 *
ABTS	0.002 **	0.016 *	0.010 *

1

2 **Figure 4:** Boxplots and ANOVA table for phenolic concentrations and ABTS values measured on  
3 seeds from pods<sub>L≥ 5cm</sub> under the 10 treatments (crossing temperature modalities and sulphur  
4 conditions). Both variables displayed highly significant interaction (Temp x Sulphur) effects. For each  
5 variable, letters indicate the ranking amongst the 10 treatments. P-values and levels of significance are  
6 given for Temp, Sulphur and Temp x Sulphur interaction effects. Levels of significance: ns non-  
7 significant,  $p < 0.05^*$ ,  $p < 0.01^{**}$ ,  $p < 0.001^{***}$ . SA\*: synaptic acid, ABTS: 2,2'-azino-bis (3-  
8 ethylbenzothiazoline-6-sulfonic acid).



1

## 2 **Discussion**

### 3 *Yield components were only impacted in reproductive plant parts developed after the stress*

4 Yield components were not drastically impacted by the high temperature sequences applied at GS72  
5 (Table 1). No effect of temperature was observed on yield of pods<sub>L $\geq$ 5cm</sub>, meaning that these seeds  
6 experienced a normal filling period. Therefore, the effect of the heat sequences on total seed yield  
7 ( $p < 0.05$ ) was the consequence of a yield reduction in seeds from pods<sub>L $<$ 5cm</sub>, which resulted from  
8 reduced seed number and individual seed weight, even though neither component reduction was  
9 significant. Indeed, abortions of young pods during the Temp-modality exposure and/or failure of  
10 pollination in the flowers present during this period might have occurred (Young et al., 2004). The  
11 reproductive adaptation might also have reduced the incidence of late flowering (specific to  
12 indeterminate species) to the benefit of filling seeds in already developed pods. These observations are  
13 consistent with prior studies in pea (another indeterminate growth species) that demonstrated heat-  
14 stress modification of seed distribution along the main stem, which led to larger quantities of seeds on  
15 the basal parts than on the upper parts where reproductive organs are younger. These results explained  
16 the maintenance of seed yield from basal reproductive parts due to a higher allocation rate of  
17 carbohydrates and excluded the direct effects of heat stress on developing seeds (Guilioni et al., 2003).  
18 The non-significant effect of sulphur on seed yield components, other than in the total seed number,  
19 indicates that the Low Sulphur supply in our study satisfied growth rate and yield requirements.

20

### 21 *Priming effects on quality criteria are dependent on the timing of the stress events*

22 Our initial objective was to pinpoint any beneficial effect from a mild stress event that occurs prior to  
23 repeated later intense heat peaks. This could be accounted for by a beneficial heat stress memory  
24 generated by induced thermo-sensitization. Our work highlights that the alleviating effects of the early  
25 mild stress event prior to later heat peaks are not observed for all the measured quality variables. Our  
26 results pointed out that these alleviating effects are likely determined by (i) whether the expected

1 priming event occurs concomitantly with biosynthesis of seed storage compounds, and (ii) whether the  
2 temperature is adequate to enhance the targeted process.

3 Figure 5 presented the Temp-modality effects on yield components and nutritional and physiological  
4 quality criteria measured in maturing seeds, for the Temp-modalities (early mild stress, 3 or 4 late heat  
5 peaks and priming modality) and the sulphur limitation. The priming modality significantly decreased  
6 seed C, total FAs, SFAs and UFAs concentrations under both Sulphur conditions. These observations  
7 suggest that the more the duration of the heat stress sequence (i.e. sequence that records the more days  
8 with temperature higher than the control temperature), the most detrimental for maintaining C  
9 concentration and hence FAs synthesis, since the priming modality (that cumulated the highest number  
10 of days with temperature above the Temp-control modality) induced a greater decrease than the early  
11 mild stress or the late heat peaks alone. These results correlate with earlier studies (Aksouh-Harradj et  
12 al., 2006; Aksouh et al., 2001) of intense temperature exposure during seed maturation, but contrast  
13 with others that highlighted increased total FAs concentrations under a long mild stressing event  
14 (Brunel-Muguet et al., 2015).

15 Disruption of FAs accumulation in oilseed rape was pointed out under intense heat stress across a  
16 range of temperatures similar to those imposed under our heat-peak conditions. This was shown to be  
17 the consequence of photosynthesis inhibition and downregulation of BnWRI1 gene expression, which  
18 is a key regulator of FA biosynthesis (Huang et al., 2019; Ruuska et al., 2002). Our results also  
19 revealed higher total FAs, saturated FAs and unsaturated FAs concentrations under Low Sulphur  
20 condition (Table 2, Figure 2), which contrasted with prior results (Brunel-Muguet et al., 2015). These  
21 differences can be explained in the current work by the lower Sulphur supply and the different  
22 approaches to seed sampling and analysis (on maturing seeds exposed to the heat stress modalities).

23 The negative effects of high temperature – mainly of duration of high temperature sequence (Figure 5)  
24 - on total FAs and unsaturated FAs also indicated that (i) FA biosynthesis was concomitant with the  
25 17-days high-temperature treatment starting approximately 3 weeks after the onset of flowering  
26 (Figure 1), with FAs levels rising during the late storage stage (i.e. 20 after pollination (Borisjuk et al.,  
27 2013; Niu et al., 2009)) and (ii) the later the heat peaks, the lower the effects. The increased  $\omega 6:\omega 3$

1 ratio under the priming and 4 late heat peaks modalities is also detrimental to oil quality, and is known  
2 to result from impaired functioning of the oleic and linoleic desaturases (Aksouh-Harradj et al., 2006).  
3 As usually observed, the accumulation of lipids and proteins (herein linked to seed N concentration)  
4 are negatively correlated because they are competitive processes that have spatial and temporal  
5 overlaps (Borişjuk et al., 2013; Grami and Stefansson, 1977). Therefore, consistently, while the  
6 priming modality decreased the total FA concentration, the seed N concentration remained similar to  
7 Temp-control under both Sulphur conditions (Table 2). A decrease in seed storage protein  
8 concentrations under the late heat peaks modalities were observed, although not significant, suggesting  
9 that the high concentrations observed in the priming modality could be interpreted as a priming effect  
10 that helped to overcome further negative effects of the late heat peaks. This observation supports the  
11 hypothesis of compensatory effects between the pods that were maturing and the ones that were still  
12 developing during the 17-days temperature sequence. Indeed, seed N accumulation was likely to be  
13 impaired in filling seeds, and this resulted in N reallocation towards filled and maturing seeds. In  
14 contrast, while the seed storage protein concentrations were not significantly increased under high  
15 temperature, they increased under Low Sulphur condition as previously observed (Brunel-Muguet et  
16 al., 2015).

17 Ultimately, seed physiological characteristics were also highly dependent on the features of the high  
18 temperature sequences (duration, intensity, frequency), which shapes the dynamics of biosynthesis of  
19 storage compounds involved in seed dormancy and stress tolerance. Seed storability and desiccation  
20 tolerance can be estimated by two proxies i.e. membrane conductivity and the  
21 [raffinose+stachyose]:sucrose ratio. Our results highlighted strong negative effects of high temperature  
22 on seed conductivity, indicating degradation of membrane permeability (Table 3). However, the  
23 [raffinose+stachyose]:sucrose ratio remained unchanged in the priming and in the early mild stress  
24 modalities, which suggests acquisition of desiccation tolerance irrespective of Sulphur supply. The  
25 beneficial effects of the early mild stress on desiccation tolerance were maintained over the late heat  
26 peaks which points out its alleviating effect.

1 Several phytohormones control seed dormancy, as indicated by the ABA:GA3 ratio, which has been  
2 shown to vary according to stresses imposed on the parent plant during seed development (Brunel-  
3 Muguet et al., 2015). A high ABA:GA3 ratio indicates higher seed dormancy, which is expected for  
4 efficient seed storage before favorable environmental conditions permit the seeds to germinate. The  
5 priming modality and the late heat peaks modalities had the greatest impact on the ABA:GA3 ratio  
6 (Figures 3 and 5). Along with the non-significant effect of the early mild stress modality, these  
7 observations indicate that the intensity and timing of the stress exposure are determining on the  
8 ABA:GA3 ratio. In this example, the temperature of the mild stress was not deleterious on the  
9 ABA:GA3 ratio, or this early sequence occurs before the hormone syntheses, thus leading to no  
10 observed effects. These observations also pointed out that the temperature of the late heat peaks was  
11 deleterious on the ABA:GA3 ratio and that these heat peaks occurred when these hormones were  
12 synthesized. Therefore, our results raised questions not only about the compound-specific  
13 biosynthesis/maturation temperatures but also the synchronization between these compound specific-  
14 processes and the temperature event applied in the aim to induce an effective thermo-sensitization  
15 effects.

Criteria		Temp-modality				S condition
		Early Mild Stress	3 Heat Peaks	4 Heat Peaks	Priming	Low Sulphur
YIELD COMPONENTS	Total Seed Yield	-10%	-6%	-5%	-16%	
	Total Seed number					-5%
	Seeds Yield from pods $L < 5cm$	-40%	-16%	-22%	-40%	
NUTRITIONAL QUALITY	Seed Carbon	-2%	-2%	-3%	-5%	
	Seed Nitrogen				+22%	
	Seed Sulphur	+3%	+4%	+2%	+20%	-32%
	Total Fatty Acids	+3%	+5%	-5%	-15%	+17%
	Saturated FAs		+9%		-9%	+16%
	Unsaturated FAs			-5%	-17%	+19%
	$\omega 6:\omega 3$ ratio	-9%	+1%	+7%	+9%	
	Protein concentration					+75%
PHYSIOLOGICAL QUALITY	[Raff+Stach]:Suc ratio	+12%		-15%	+2%	
	Seed conductivity				+262%	
	ABA:GA3 ratio	-5%	+28%	+30%	+28%	+66%



1

2 **Figure 5:** Summary of the effects of the four temperature sequences (early mild stress, 3 and 4 late  
3 heat peaks and priming modality) and the Sulphur conditions on yield components, nutritional and  
4 physiological quality criteria. Indicators of quality were measured on seeds from pods $L \geq 5cm$  at the  
5 beginning of stress exposure. The effects are given in reference to the ANOVAs in Tables 1, 2, 3 and  
6 Figures 2 and 3. As illustrated in the legend, colors indicate the trends by level of increase or decrease  
7 to the Temp-control modality. Numbers in boxes display the relative difference between the Temp-  
8 modality and the Temp-control (for temperature modalities) and between Low Sulphur and High  
9 Sulphur (for Sulphur conditions). FAs: fatty acids, Raff: raffinose, Stach: stachyose, Suc: sucrose,  
10 ABA: abscisic acid, GA3: gibberellic acid.

11

### 12 *Modulation of antioxidant capacities and thermotolerance acquisition as stress defense strategy*

13 Heat stress is known to trigger oxidative bursts that lead to a wide spectrum of responses including  
14 enzymatic and non-enzymatic components such as polyphenols, hormones and sugars, which have  
15 been shown to scavenge reactive oxygen species (ROS) (Nishizawa et al., 2008; Serrano et al., 2019;

1 Soares et al., 2019; Soengas et al., 2018). In our study, measurements of the antioxidant capacity  
2 indicated complex results that differed according to the S supply (Figure 4). In the High Sulphur  
3 condition, the greatest increase were observed under the early mild stress modality and not under the  
4 priming nor the late heat peaks modalities, likely to phenolic concentrations. These observations  
5 contrasted with prior studies which indicated enhancement of subcellular antioxidant activities using  
6 other antioxidant systems as proxies (such as enzymes superoxide dismutase, glutathione reductase,  
7 and peroxidase in mitochondria and chloroplasts of wheat leaves) under multiple heat priming  
8 sequences prior to later high temperature events (Wang et al., 2014).

9 Our findings also pointed out that(Wang et al., 2014) the measured concentrations of raffinose and  
10 stachyose, which have been reported to protect against oxidative damage (Nishizawa et al., 2008),  
11 were not increased under early mild stress modality (irrespective of the Sulphur conditions), and so it  
12 was not possible to ascertain their role in the antioxidant defense pathways in our conditions (Table 3).  
13 But as discussed above, this soluble sugar-related results should be interpreted carefully regarding  
14 their timing and optimal biosynthesis within the maturing seeds (Baud et al., 2002; Leprince et al.,  
15 2017). Indeed, our results might suggest that the temperature events (whether mild or intense) were  
16 above the temperature threshold of the stepwise transfer reactions involving raffinose synthase and  
17 stachyose synthase, leading to impairment of sugar biosynthesis and thus the sugar-mediated oxidative  
18 response (Gangl et al., 2015).

19 Phytohormones such as IAA and SA have been demonstrated to interact with ROS during stress  
20 tolerance as stress-signaling cues (Balfagón et al., 2019; Bielach et al., 2017; Clarke et al., 2004;  
21 Prerostova et al., 2020; Sharma and Laxmi, 2016) and have also been linked to stress thermotolerance  
22 (Bokszczanin et al., 2013; Clarke et al., 2009; Shu et al., 2016; Tuan et al., 2019) because increases in  
23 the levels of these compounds under high temperature are associated with greater stress-induced  
24 dormancy and thermoinhibition (Toh et al., 2012). (Bokszczanin et al., 2013; Clarke et al., 2009)

25 In our study, we observed drastic increases in IAA and SA in mature seeds collected from pods  
26 exposed to the longest cumulated high temperature (priming modality) or late heat peaks at the onset  
27 of maturation under High Sulphur conditions and to a lesser extent under Low Sulphur conditions

1 (Figure 3). The 4 late heat peak modality induced the highest increase in IAA and SA, while the early  
2 mild stress modality remained similar to the control under both Sulphur conditions. These  
3 observations pointed out that the early mild stress event prior to 4 late heat peaks allowed their  
4 negative effects on IAA and SA concentrations to be alleviated thus leading to lower thermoinhibition.  
5 Our results also highlighted that the late 3 heat peaks modality had less effect than the late 4 heat  
6 peaks on IAA and SA, which suggests that the shorter the period of intense stressing, the lower the  
7 impact on the seed hormones concentration. (Bokszczanin et al., 2013; Clarke et al., 2009)

8

### 9 *Impact of S availability on thermotolerance acquisition*

10 Firstly, the level of sulphur limitation used in our experiment was mild enough not to impact seed  
11 yield and components as only a slight negative effect on the total seed number was observed (Figure  
12 5). The effects of sulphur restriction were observed on nutritional quality criteria i.e. FAs-related  
13 variables, seed storage proteins and physiological quality criteria i.e. ABA:GA3 ratio, which were all  
14 increased under sulphur limitation. Interactions effects were only observed for saturated FAs and the  
15 ABA:GA3 ratio. They highlighted that late heat peaks or early mild stress events led to increased  
16 unsaturated FAs concentrations under Low Sulphur but not under High Sulphur (Figure 2). The  
17 ABA:GA3 ratio was only increased under Low Sulphur with the late heat peaks (Figure 3).

18 The variables related to thermoinhibition (i.e. phytohormones IAA and SA) and oxidative responses  
19 were globally decreased by sulphur limitation (Figures 3 and 4). Nevertheless, the sense of variation or  
20 the extent of the decreases varied according to the temperature modality. Indeed, Temp x Sulphur  
21 interaction effects were observed for IAA, SA concentrations and the ABA:GA3 ratio, phenolic  
22 concentrations and ABTS values. Our results pinpointed increased antioxidant capacities with early  
23 mild stress event (observed for the priming modality and the mild stress modality) under Low Sulphur  
24 compared to the natural thermoperiod whereas similar or slightly lower values to the natural  
25 thermoperiod were observed with the early mild stress event under High Sulphur (Figure 4). For  
26 phytohormones IAA and SA, the temperature modalities globally increased the concentrations under  
27 the two Sulphur conditions although this trend was more pronounced under High Sulphur than under

1 Low Sulphur. Nevertheless, only under Low Sulphur, the early mild stress event alone trigger  
2 increased concentrations compared to Temp-control. Sulphur nutrition has been reported to improve  
3 antioxidant defenses due to S-containing antioxidant compounds such as the redox-active cysteine  
4 residues of glutathione and thioredoxin (Mukwevho et al., 2014) and also specific phytohormones  
5 (Bashir et al., 2015; Hasanuzzaman et al., 2018; Xia et al., 2015). IAA biosynthesis relies on the  
6 effectiveness of S metabolism via adequate levels of glutathione (Kopriva et al., 2016) and is  
7 improved when S is limiting (Nikiforova et al., 2003). SA is also known to interact with S during SA  
8 homeostasis regulation (Baek et al., 2010). When comparing both Sulphur conditions under the natural  
9 thermoperiod, our results are in line with these findings as a slight increase was observed under Low  
10 Sulphur. Our study also highlighted how the intensity, the timing and the duration of heat stress can  
11 restrict the boosting effects of sulphur restriction.

12

### 13 **Conclusion**

14 Our study analyzed the effects of different high temperature protocols applied at the onset of seed  
15 maturation on yield components, seed nutritional and physiological quality under two Sulphur  
16 supplies. The initial working hypothesis was to highlight inducing thermotolerance protocols based on  
17 stress memory acquisition. Our results pinpointed that the effects of duration, timing and intensity of  
18 the stressing events that designed the different temperature protocols were determining as they led to  
19 different impacts on the measured quality criteria. Our results showed that thermo-sensitization  
20 protocols must require: (i) the optimal temperature to promote a targeted process and (ii) the  
21 synchronization between the expected thermoprimering event and the underlying biosynthesis and  
22 maturation processes specific to the targeted storage compound. Our experiments were also designed  
23 to observe whether sulphur nutrition (which is essential to Brassica species) interfered with expected  
24 thermo-sensitization effects. The level of sulphur limitation in our assay was mild enough to impact a  
25 few nutritional (fatty acids, seed storage protein concentration) and physiological (IAA, SA,  
26 ABA:GA3 ratio) quality criteria, as well as the antioxidant capacity. This raises the question of the  
27 direct role of sulphur in the cascade of events leading to the hormones biosynthesis and the indirect



1 role of sulphur in stress memory mediated by epigenetic regulation. In contrast, sulphur restriction  
2 optimized seed oil concentrations in stress-exposed maturing seeds, especially when the stress was  
3 delayed, thus highlighting the need to satisfy sulphur requirements in certain climatic contexts.  
4 Overall, we foresee the need for trade-offs to optimize quality criteria that are dependent on features of  
5 temperature such as intensity, frequency and timing of application. High temperature acclimation  
6 strategies should also include sulphur supply which restricted levels can either improve seed quality  
7 criteria (e.g. total FAs, saturated and unsaturated FAs) in temperature-stressed maturing seed, lessen  
8 the positive effects of a thermoprimer profile (e.g. indicators of desiccation tolerance) or amplify the  
9 oxidative responses to high temperature stress when applied at sensitive stages during the maturation  
10 process.

11

## 12 **Acknowledgments**

13 This study was funded by the Environment and Agronomy division of INRAE (funding “Pari  
14 Scientifique INRAE”). The authors would like to thank Magali Bodereau, Josiane Pichon, Josette  
15 Bonnefoy, Marine Lechevrel, Théo Lemercier, Julien Mignot and Christophe Muguet for their  
16 technical assistance and help with data analysis. They also thank Dr. Bae for the hormone analyses,  
17 Nicolas Elie from the CEMABIO platform for seed image analysis and Laurence Cantrill for editing  
18 the manuscript and providing relevant suggestions.

## 1 Bibliography

- 2 Aguirrezábal L., Martre P., Pereyra-Irujo G., Echarte M.M. & Izquierdo N. (2015) Improving grain  
3 quality: ecophysiological and modeling tools to develop management and breeding strategies.  
4 *Crop Physiology*, 423–465.
- 5 Akmouche Y., Cheneby J., Lamboeuf M., Elie N., Laperche A., Bertheloot J., Brunel-Muguet S.  
6 (2019) Do nitrogen- and sulphur-remobilization-related parameters measured at the onset of the  
7 reproductive stage provide early indicators to adjust N and S fertilization in oilseed rape  
8 (*Brassica napus* L.) grown under N- and/or S-limiting supplies? *Planta* **250**, 2047–2062.
- 9 Aksouh-Harradj N.M., Campbell L.C. & Mailer R.J. (2006) Canola response to high and moderately  
10 high temperature stresses during seed maturation. *Canadian Journal of Plant Science* **86**, 967–  
11 980.
- 12 Aksouh N.M., Jacobs B.C., Stoddard F.L. & Mailer R.J. (2001) Response of canola to different heat  
13 stresses. *Australian Journal of Agricultural Research* **52**, 817–824.
- 14 Baek D., Pathange P., Chung J. et al. (2010) A stress-inducible sulphotransferase sulphonates salicylic  
15 acid and confers pathogen resistance in *Arabidopsis*. *Plant, Cell & Environment* **33**, 1383–1392.
- 16 Bailly C., Audigier C., Ladonne F., Wagner M.H., Coste F., Corbineau F. & Côme D. (2001) Changes  
17 in oligosaccharide content and antioxidant enzyme activities in developing bean seeds as related  
18 to acquisition of drying tolerance and seed quality. *Journal of Experimental Botany* **52**, 701–708.
- 19 Balfagón D., Sengupta S., Gómez-Cadenas A., Fritschi F.B., Azad R.K., Mittler R. & Zandalinas S.I.  
20 (2019) Jasmonic acid is required for plant acclimation to a combination of high light and heat  
21 stress. *Plant physiology* **181**, 1668–1682.
- 22 Bashir H., Ibrahim M.M., Bagheri R., Ahmad J., Arif I.A., Baig M.A. & Qureshi M.I. (2015)  
23 Influence of sulfur and cadmium on antioxidants, phytochelatins and growth in Indian mustard.  
24 *AoB PLANTS* **7**.
- 25 Baud S., Boutin J.P., Miquel M., Lepiniec L. & Rochat C. (2002) An integrated overview of seed  
26 development in *Arabidopsis thaliana* ecotype WS. *Plant Physiology and Biochemistry* **40**, 151–  
27 160.
- 28 Baux A., Colbach N., Allirand J.M., Jullien A., Ney B. & Pellet D. (2013) Insights into temperature  
29 effects on the fatty acid composition of oilseed rape varieties. *European Journal of Agronomy*  
30 **49**, 12–19.
- 31 Baux A., Hebeisen T. & Pellet D. (2008) Effects of minimal temperatures on low-linolenic rapeseed  
32 oil fatty-acid composition. *European Journal of Agronomy* **29**, 102–107.
- 33 Bielach A., Hrtyan M. & Tognetti V.B. (2017) Plants under stress: Involvement of auxin and  
34 cytokinin. *International Journal of Molecular Sciences* **18**.
- 35 Bokszczanin K., Fragkostefanakis S., Bostan H., Bovy A., Chaturvedi P., Chiusano M.L., Winter P.  
36 (2013) Perspectives on deciphering mechanisms underlying plant heat stress response and  
37 thermotolerance. *Frontiers in Plant Science* **4**, 315.
- 38 Borisjuk L., Neuberger T., Schwender J., Heinzl N., Sunderhaus S., Fuchs J., Rolletschek H. (2013)

- 1 Seed architecture shapes embryo metabolism in oilseed rape. *The Plant cell* **25**, 1625–40.
- 2 Bruce T.J.A., Matthes M.C., Napier J.A. & Pickett J.A. (2007) Stressful “memories” of plants:  
3 Evidence and possible mechanisms. *Plant Science* **173**, 603–608.
- 4 Brunel-Muguet S., D’Hooghe P., Bataillé M.-P., Larré C., Kim T.-H., Trouverie J., Dürr C. (2015)  
5 Heat stress during seed filling interferes with sulfur restriction on grain composition and seed  
6 germination in oilseed rape (*Brassica napus* L.). *Frontiers in Plant Science* **6**.
- 7 Canvin D.T. (1965) The effect of temperature on the oil content and fatty acid composition of the oils  
8 from several oil seed crops. *Canadian Journal of Botany* **43**, 63–69.
- 9 Chebrolu K.K., Fritschi F.B., Ye S., Krishnan H.B., Smith J.R. & Gillman J.D. (2016) Impact of heat  
10 stress during seed development on soybean seed metabolome. *Metabolomics* **12**, 1–14.
- 11 Chitnis V.R., Gao F., Yao Z., Jordan M.C., Park S. & Ayele B.T. (2014) After-ripening induced  
12 transcriptional changes of hormonal genes in wheat seeds: The cases of brassinosteroids,  
13 ethylene, cytokinin and salicylic acid. *PLoS ONE* **9**, 1–14.
- 14 Christidis N., Jones G.S. & Stott P.A. (2015) Dramatically increasing chance of extremely hot  
15 summers since the 2003 European heatwave. **5**, 46–49.
- 16 Clarke S.M., Cristescu S.M., Miersch O., Harren F.J.M., Wasternack C. & Mur L.A.J. (2009)  
17 Jasmonates act with salicylic acid to confer basal thermotolerance in *Arabidopsis thaliana*. *New*  
18 *Phytologist* **182**, 175–187.
- 19 Clarke S.M., Mur L.A.J., Wood J.E. & Scott I.M. (2004) Salicylic acid dependent signaling promotes  
20 basal thermotolerance but is not essential for acquired thermotolerance in *Arabidopsis thaliana*.  
21 *Plant Journal* **38**, 432–447.
- 22 Crisp P.A., Ganguly D., Eichten S.R., Borevitz J.O. & Pogson B.J. (2016) Reconsidering plant  
23 memory: Intersections between stress recovery, RNA turnover, and epigenetics. *Science*  
24 *Advances* **2**, e1501340–e1501340.
- 25 D’Hooghe P., Dubousset L., Gallardo K., Kopriva S., Avice J.-C. & Trouverie J. (2014) Evidence for  
26 proteomic and metabolic adaptations associated with alterations of seed yield and quality in  
27 sulfur-limited *Brassica napus* L. *Molecular & Cellular Proteomics* **13**, 1165–1183.
- 28 D’Hooghe P., Escamez S., Trouverie J. & Avice J.-C. (2013) Sulphur limitation provokes  
29 physiological and leaf proteome changes in oilseed rape that lead to perturbation of sulphur,  
30 carbon and oxidative metabolisms. *BMC plant biology* **13**, 23.
- 31 Debeaujon I. & Koornneef M. (2000) *Gibberellin requirement for arabidopsis seed germination is*  
32 *determined both by testa characteristics and embryonic abscisic acid 1*.
- 33 Deng X. & Scarth R. (1998) Temperature effects on fatty acid composition during development of  
34 low-linolenic oilseed rape (*Brassica napus* L.). *Journal of the American Oil Chemists’ Society* **75**,  
35 759–766.
- 36 Ding Y., Fromm M. & Avramova Z. (2012) Multiple exposures to drought “train” transcriptional  
37 responses in *Arabidopsis*. *Nature communications* **3**, 740.
- 38 Dornbos D.L. & Mullen R.E. (1992) Soybean seed protein and oil contents and fatty acid composition

- 1 adjustments by drought and temperature. *Journal of the American Oil Chemists Society* **69**, 228–  
2 231.
- 3 Finkelstein R. (2013) Abscisic acid synthesis and response. *The Arabidopsis Book* **11**, e0166.
- 4 Gangl R., Behmüller R. & Tenhaken R. (2015) Molecular cloning of AtRS4, a seed specific  
5 multifunctional RFO synthase/galactosylhydrolase in *Arabidopsis thaliana*. *Frontiers in Plant*  
6 *Science* **6**, 789.
- 7 Gauthier M., Pellet D., Monney C., Herrera J.M., Rougier M. & Baux A. (2017) Fatty acids  
8 composition of oilseed rape genotypes as affected by solar radiation and temperature. *Field*  
9 *Crops Research* **212**, 165–174.
- 10 González-Centeno M.R., Comas-Serra F., Femenia A., Rosselló C. & Simal S. (2015) Effect of power  
11 ultrasound application on aqueous extraction of phenolic compounds and antioxidant capacity  
12 from grape pomace (*Vitis vinifera* L.): Experimental kinetics and modeling. *Ultrasonics*  
13 *Sonochemistry* **22**, 506–514.
- 14 González-Centeno M.R., Jourdes M., Femenia A., Simal S., Rosselló C. & Teissedre P.-L. (2012)  
15 Proanthocyanidin composition and antioxidant potential of the stem winemaking byproducts  
16 from 10 different grape varieties (*Vitis vinifera* L.). *Journal of Agricultural and Food Chemistry*  
17 **60**, 11850–11858.
- 18 Grami B. & Stefansson B. (1977) Gene action for protein and oil content in summer rape. *Canadian*  
19 *Journal of Plant Science* **57**.
- 20 Groot M.P., Kooke R., Knob N., Vergeer P., Keurentjes J.J.B., Ouborg N.J. & Verhoeven K.J.F.  
21 (2016) Effects of multi-generational stress exposure and offspring environment on the expression  
22 and persistence of transgenerational effects in *Arabidopsis thaliana*. *PLoS ONE* **11**, 1–16.
- 23 Guilioni L., Wéry J. & Lecoeur J. (2003) High temperature and water deficit may reduce seed number  
24 in field pea purely by decreasing plant growth rate. *Functional Plant Biology* **30**, 1151–1164.
- 25 Guilioni L., Wery J. & Tardieu F. (1997) Heat stress-induced abortion of buds and flowers in pea: Is  
26 sensitivity linked to organ age or to relations between reproductive organs? *Annals of Botany* **80**,  
27 159–168.
- 28 Hasanuzzaman M., Bhuyan M.H.M.B., Mahmud J.A., Nahar K., Mohsin S.M., Parvin K. & Fujita M.  
29 (2018) Interaction of sulfur with phytohormones and signaling molecules in conferring abiotic  
30 stress tolerance to plants. *Plant Signaling and Behavior* **13**, 1–5.
- 31 Hatzig S. V., Nuppenau J.N., Snowdon R.J. & Schiebl S. V. (2018) Drought stress has  
32 transgenerational effects on seeds and seedlings in winter oilseed rape (*Brassica napus* L.). *BMC*  
33 *Plant Biology* **18**, 1–13.
- 34 Hilker M. & Schmölling T. (2019) Stress priming, memory, and signalling in plants. *Plant Cell and*  
35 *Environment* **42**, 753–761.
- 36 Hoegh-Guldberg O., Jacob D., Taylor M., Bindi M., Brown S., Camilloni I., Zhou G. (2018) Impacts  
37 of 1.5°C Global Warming on Natural and Human Systems. In In: Global Warming of 1.5°C. An  
38 IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and  
39 related global greenhouse gas emission pathways, in the context of strengthening the global

- 1 response to the threat of climate change. (eds V. Masson-Delmotte, P. Zhai, H. Pörtner, D.  
2 Roberts, J. Skea, P. Shukla, ... T. Waterfield), p. 32 pp.
- 3 Huang R., Liu Z., Xing M., Yang Y., Wu X., Liu H. & Liang W. (2019) Heat stress suppresses  
4 brassica napus seed oil accumulation by inhibition of photosynthesis and BnWRI1 pathway.  
5 *Plant and Cell Physiology* **60**, 1457–1470.
- 6 Kinoshita T. & Seki M. (2014) Epigenetic memory for stress response and adaptation in plants. *Plant*  
7 *and Cell Physiology* **55**, 1859–1863.
- 8 Kopriva S., Talukdar D., Takahashi H., Hell R., Sirko A., D'Souza S.F. & Talukdar T. (2016)  
9 Editorial: Frontiers of sulfur metabolism in plant growth, development, and stress response.  
10 *Frontiers in Plant Science* **6**, 1220.
- 11 Kotak S., Larkindale J., Lee U., von Koskull-Döring P., Vierling E. & Scharf K.D. (2007) Complexity  
12 of the heat stress response in plants. *Current Opinion in Plant Biology* **10**, 310–316.
- 13 Kumar S. (2018) Epigenetic memory of stress responses in plants. *J. Phytochem. Biochem* **2**, e102.
- 14 Lancashire P.D., Bleiholder H., Van Den Boom T., Langelüddeke P., Stauss R., Weber E.,  
15 Witzemberger A. (1991). A uniform decimal code for growth stages of crops and weeds. *Annals*  
16 *of applied Biology* **119**, 561-601.
- 17 Lee S. & Park C.-M. (2010) Modulation of reactive oxygen species by salicylic acid in Arabidopsis  
18 seed germination under high salinity. *Plant Signaling & Behavior* **5**, 1534.
- 19 Leprince O., Pellizzaro A., Berriri S. & Buitink J. (2017) Late seed maturation: Drying without dying.  
20 *Journal of Experimental Botany* **68**, 827–841.
- 21 Liu L., Liu F., Chu J. et al. (2019) A transcriptome analysis reveals a role for the indole GLS-linked  
22 auxin biosynthesis in secondary dormancy in rapeseed (*Brassica napus* L.). *BMC Plant Biology*  
23 **19**, 1–18.
- 24 Marchand L., Pelosi C., González-Centeno M.R. et al. (2016) Trace element bioavailability, yield and  
25 seed quality of rapeseed (*Brassica napus* L.) modulated by biochar incorporation into a  
26 contaminated technosol. *Chemosphere* **156**.
- 27 Meng J., Wang L., Wang J. et al. (2018) Methionine adenosyltransferase 4 mediates DNA and histone  
28 methylation. *Plant Physiology* **177**, pp.00183.2018.
- 29 Molinier J., Ries G., Zipfel C. & Hohn B. (2006) Transgeneration memory of stress in plants. *Nature*  
30 **442**, 1046–1049.
- 31 Morrison M.J. & Stewart D.W. (2002) Heat stress during flowering in summer Brassica. *Crop Science*  
32 **42**, 797–803.
- 33 Mukwevho E., Ferreira Z. & Ayeleso A. (2014) Potential role of sulfur-containing antioxidant systems  
34 in highly oxidative environments. *Molecules* 2014, Vol. 19, Pages 19376-19389 **19**, 19376–  
35 19389.
- 36 Nikiforova V., Freitag J., Kempa S., Adamik M., Hesse H. & Hoefgen R. (2003) Transcriptome  
37 analysis of sulfur depletion in *Arabidopsis thaliana*: Interlacing of biosynthetic pathways  
38 provides response specificity. *Plant Journal* **33**, 633–650.

- 1 Nishizawa A., Yabuta Y. & Shigeoka S. (2008) Galactinol and raffinose constitute a novel function to  
2 protect plants from oxidative damage. *Plant Physiology* **147**, 1251–1263.
- 3 Niu Y., Wu G.-Z., Ye R., Lin W.-H., Shi Q.-M., Xue L.-J., ... Xue H.-W. (2009) Global analysis of  
4 gene expression profiles in *Brassica napus* developing seeds reveals a conserved lipid  
5 metabolism regulation with *Arabidopsis thaliana*. *Molecular Plant* **2**, 1107–1122.
- 6 Ohama N., Sato H., Shinozaki K., Yamaguchi-Shinozaki K., Lesk C., et al. (2016) Transcriptional  
7 regulatory network of plant heat stress response. *Trends in Plant Science* **0**, 84–87.
- 8 Pan X., Welti R. & Wang X. (2010) Quantitative analysis of major plant hormones in crude plant  
9 extracts by high-performance liquid chromatography–mass spectrometry. *Nature Protocols* **5**,  
10 986–992.
- 11 Pekrun C., Lutman P.J.W. & Baeumer K. (1997) Germination behaviour of dormant oilseed rape seeds  
12 in relation to temperature. *Weed Research* **37**, 419–431.
- 13 Poisson E., Trouverie J., Brunel-Muguet S., Akmouche Y., Pontet C., Pinochet X. & Avice J.C. (2019)  
14 Seed yield components and seed quality of oilseed rape are impacted by sulfur fertilization and  
15 its interactions with nitrogen fertilization. *Frontiers in Plant Science* **10**.
- 16 Prerostova S., Dobrev P.I., Kramna B., Gaudinova A., Knirsch V., Spichal L., Zatloukal M., Vankova  
17 R. (2020) Heat acclimation and inhibition of Cytokinin degradation positively affect heat stress  
18 tolerance of *Arabidopsis*. *Frontiers in Plant Science* **11**, 1–14.
- 19 Ruuska S.A., Girke T., Benning C. & Ohlrogge J.B. (2002) Contrapuntal networks of gene expression  
20 during *Arabidopsis* seed filling. *The Plant Cell* **14**, 1191–1206.
- 21 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for  
22 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 23 Sage T.L., Bagha S., Lundsgaard-Nielsen V., Branch H.A., Sultmanis S. & Sage R.F. (2015) The  
24 effect of high temperature stress on male and female reproduction in plants. *Field Crops*  
25 *Research* **182**, 30–42.
- 26 Schindelin, J.; Arganda-Carreras, I. & Frise, E. et al. (2012), "Fiji: an open-source platform for  
27 biological-image analysis", *Nature methods* 9(7): 676-682.
- 28 Schulte L.R., Ballard T., Samarakoon T., Yao L., Vadlani P., Staggenborg S. & Rezac M. (2013)  
29 Increased growing temperature reduces content of polyunsaturated fatty acids in four oilseed  
30 crops. *Industrial Crops and Products* **51**, 212–219.
- 31 Serrano N., Ling Y., Bahieldin A. & Mahfouz M.M. (2019) Thermopriming reprograms metabolic  
32 homeostasis to confer heat tolerance. *Scientific Reports* **9**, 1–14.
- 33 Sharma M. & Laxmi A. (2016) Jasmonates: Emerging players in controlling temperature stress  
34 tolerance. *Frontiers in Plant Science* **6**, 1129.
- 35 Shu K., Liu X., Xie Q. & He Z. (2016) Two faces of one seed: hormonal regulation of dormancy and  
36 germination. *Molecular Plant* **9**, 34–45.
- 37 Soares C., Carvalho M.E.A., Azevedo R.A. & Fidalgo F. (2019) Plants facing oxidative challenges—  
38 A little help from the antioxidant networks. *Environmental and Experimental Botany* **161**, 4–25.

- 1 Soengas P., Rodríguez V.M., Velasco P. & Cartea M.E. (2018) Effect of Temperature Stress on  
2 Antioxidant Defenses in Brassica oleracea. *ACS Omega* **3**, 5237–5243.
- 3 Szydłowska-Czerniak A., Amarowicz R. & Szlyk E. (2010) Antioxidant capacity of rapeseed meal  
4 and rapeseed oils enriched with meal extract. *European Journal of Lipid Science and Technology*  
5 **112**, 750–760.
- 6 Toh S., Kamiya Y., Kawakami N., Nambara E., McCourt P. & Tsuchiya Y. (2012) Thermoinhibition  
7 uncovers a role for strigolactones in arabidopsis seed germination. *Plant and Cell Physiology* **53**,  
8 107–117.
- 9 Trnka M., Rötter R.P., Ruiz-Ramos M., Kersebaum K.C., Olesen J.E., Žalud Z. & Semenov M.A.  
10 (2014) Adverse weather conditions for European wheat production will become more frequent  
11 with climate change. *Nature Climate Change* **4**, 637–643.
- 12 Tuan P.A., Yamasaki Y., Kanno Y., Seo M. & Ayele B.T. (2019) Transcriptomics of cytokinin and  
13 auxin metabolism and signaling genes during seed maturation in dormant and non-dormant  
14 wheat genotypes. *Scientific Reports* **9**, 1–7.
- 15 Wahid A., Gelani S., Ashraf M. & Foolad M.R. (2007) Heat tolerance in plants: An overview.  
16 *Environmental and Experimental Botany* **61**, 199–223.
- 17 Wang X., Cai J., Liu F., Dai T., Cao W., Wollenweber B. & Jiang D. (2014) Multiple heat priming  
18 enhances thermo-tolerance to a later high temperature stress via improving subcellular  
19 antioxidant activities in wheat seedlings. *Plant Physiology and Biochemistry* **74**, 185–192.
- 20 Wang X. & Liang D. (2017) Priming: A promising strategy for crop production in response to future  
21 climate. *Journal of Integrative Agriculture* **16**, 2709–2716.
- 22 Wang X., Xin C., Cai J., Zhou Q., Dai T., Cao W. & Jiang D. (2016) Heat priming induces trans-  
23 generational tolerance to high temperature stress in wheat. *Frontiers in Plant Science* **7**, 501.
- 24 Xia X.J., Zhou Y.H., Shi K., Zhou J., Foyer C.H. & Yu J.Q. (2015) Interplay between reactive oxygen  
25 species and hormones in the control of plant development and stress tolerance. *Journal of*  
26 *Experimental Botany* **66**, 2839–2856.
- 27 Xie Z., Zhang Z.-L., Hanzlik S., Cook E. & Shen Q.J. (2007) Salicylic acid inhibits gibberellin-  
28 induced alpha-amylase expression and seed germination via a pathway involving an abscisic-  
29 acid-inducible WRKY gene. *Plant Molecular Biology* **64**, 293–303.
- 30 Young L.W., Wilen R.W. & Bonham-Smith P.C. (2004) High temperature stress of Brassica napus  
31 during flowering reduces micro- and megagametophyte fertility, induces fruit abortion, and  
32 disrupts seed production. *Journal of Experimental Botany* **55**, 485–495.

33

34

## 1 Tables

Factor\Variables	Total			Pods L < 5cm			Pods L ≥ 5cm		
	Yield	Seed number	TSW	Yield	Seed number	TSW	Yield	Seed number	TSW
Temp-modality									
Control	9.8 a	2784 ns	3.3 ns	3.2 ns	956 ns	3.4 ns	6.6 ns	1829 ns	3.6 ns
Early mild stress	8.8 ab	2565 ns	3.1 ns	1.9 ns	615 ns	3.1 ns	6.9 ns	1950 ns	3.6 ns
3 late heat peaks	9.2 ab	2723 ns	2.9 ns	2.7 ns	927 ns	3.0 ns	6.5 ns	1796 ns	3.7 ns
4 late heat peaks	9.3 ab	2724 ns	3.1 ns	2.5 ns	785 ns	3.1 ns	6.8 ns	1939 ns	3.5 ns
Priming	8.2 b	2497 ns	3.2 ns	1.9 ns	545 ns	3.2 ns	6.2 ns	1952 ns	3.2 ns
<i>se</i>	0.4	80	0.3	0.5	0.5	0.3	0.6	158	0.2
Sulphur									
High Sulphur	9.1 ns	2730 a	3.1 ns	2.6 ns	809 ns	3.1 ns	6.5 ns	1921 ns	3.4 ns
Low Sulphur	9.0 ns	2587 b	3.1 ns	2.3 ns	722 ns	3.2 ns	6.7 ns	1865 ns	3.6 ns
<i>se</i>	0.3	52	0.2	0.3	0.3	0.2	0.4	97.0	0.1
Temp effect	<b>0.047</b> *	0.068	0.874	0.309	0.082	0.886	0.944	0.912	0.393
Sulphur effect	0.769	<b>0.048</b> *	0.981	0.514	0.424	0.745	0.732	0.684	0.372
Temp x Sulphur	0.342	0.573	0.090	0.195	0.061	0.244	0.526	0.130	0.822

2

3 **Table 1:** Yield components distinguishing the two pools of pods (i.e. pods<sub>L < 5cm</sub> and pods<sub>L ≥ 5cm</sub> at the beginning of the Temp-modality application). Results are  
4 presented by factor (Temp-modality, Sulphur condition). For a given variable, different letters (Tukey multiple comparisons test) indicate the ranking among  
5 Temp-modalities (including Temp-control) or between the two Sulphur conditions. P-values and levels of significance are given for Temp, Sulphur and Temp  
6 x Sulphur interaction effects. Levels of significance: ns non-significant,  $p < 0.05$ \*,  $p < 0.01$ \*\*,  $p < 0.001$ \*\*\*. SE: standard error, TSW: Thousand Seed Weight.

7



	Seed C, N and S concentrations (%DW)			Oil contents and fatty acids (%DW)			Protein (mg/g DW)	
	Carbon	Nitrogen	Sulphur	Fatty acids	SFA	UFA	ω6:ω3 ratio	SSP
Temp-modality								
Control	59.2 a	3.2 b	0.311 ns	39 a	3.3 -	36 a	2.05 ab	150 ns
Early mild stress	58.2 ab	3.5 b	0.321 ns	40 a	3.3 -	37 a	1.87 b	168 ns
3 late heat peaks	58.2 ab	3.2 b	0.323 ns	41 a	3.6 -	38 a	2.08 ab	136 ns
4 late heat peaks	57.2 bc	3.4 b	0.316 ns	37 ab	3.3 -	34 ab	2.20 a	123 ns
Priming	56.0 c	3.9 a	0.373 ns	33 b	3.0 -	30 b	2.24 a	150 ns
<i>se</i>	0,5	0,1	0,027	2	0.1	2	0,06	21
Sulphur								
High Sulphur	57,7 ns	3,5 ns	0,392 a	35 b	3.1 -	32 b	2,05 ns	106 b
Low Sulphur	57,8 ns	3,4 ns	0,266 b	41 a	4.6 -	38 a	2,12 ns	185 a
<i>se</i>	0,4	0,1	0,009	1	0.1	1	0,05	10
Temp effect	<b>0,001</b> **	<b>0,000</b> ***	<b>0,013</b> *	<b>0,002</b> **	<b>0,001</b> **	<b>0,003</b> **	<b>0,001</b> **	0,278
Sulphur effect	0,829	0,550	<b>0,000</b> ***	<b>0,000</b> ***	<b>0,000</b> ***	<b>0,000</b> ***	0,225	<b>0,000</b> ***
Temp x Sulphur	0,613	0,574	0,296	0,112	<b>0,009</b> **	0,129	0,387	0,233

1

2 **Table 2:** Nutritional seed quality criteria measured on seeds from pods<sub>L≥ 5cm</sub>. Results are presented by factor (Temp-modality, Sulphur condition). For a given  
3 measured variable, different letters (Tukey multiple comparisons test) indicate the ranking among Temp-modalities (including Temp-control) or between the  
4 two Sulphur conditions. As SFAs are variables with highly significant interaction (Temp x Sulphur) effects, results for each individual treatment are displayed  
5 in Figure 2. P-values and levels of significance are given for Temp, Sulphur and Temp x Sulphur interaction effects. Levels of significance: ns non-significant,  
6  $p < 0.05$ \*,  $p < 0.01$ \*\*\*,  $p < 0.001$ \*\*\*. SE: standard error, DW: dry weight, SSP: Seed Storage Protein.

7

Factor\Variables	Soluble sugar				Conductivity
	Sucrose (%DW)	Raffinose (%DW)	Stachyose (%DW)	[Raffinose+stachyose] : sucrose	Seed conductivity (μS/mg)
Temp-modality					
Control	7.4 a	0.47 a	2.5 a	0.41 b	0.8 b
Early mild stress	3.0 c	0.21 c	1.1 b	0.46 ab	1.0 b
3 late heat peaks	6.1 ab	0.38 ab	2.1 a	0.41 b	1.0 b
4 late heat peaks	4.6 bc	0.26 bc	1.4 b	0.35 c	1.0 b
Priming	3.4 c	0.21 c	1.2 b	0.42 ab	2.9 a
<i>se</i>	0.5	0.03	0.1	0.01	0.2
Sulphur					
High Sulphur	4.9 ns	0.32 ns	1.7 ns	0.42 ns	1.4 ns
Low Sulphur	4.8 ns	0.29 ns	1.6 ns	0.40 ns	1.3 ns
<i>se</i>	0.5	0.03	0.2	0.01	0.2
Temp effect	<b>0.000 ***</b>	<b>0.000 ***</b>	<b>0.000 ***</b>	<b>0.000 ***</b>	<b>0.000 ***</b>
Sulphur effect	0.850	0.850	0.620	0.054	0.336
Temp x Sulphur	0.247	0.383	0.431	0.058	0.658

1

2 **Table 3:** Seed physiological quality values measured on seeds from pods<sub>L≥ 5cm</sub>. Results are presented by factor (Temp-modality, Sulphur condition). For a  
3 given measured variable, different letters (Tukey multiple comparisons test) indicate the ranking among Temp-modalities (including Temp-control) or  
4 between the two Sulphur conditions. P-values and levels of significance are given for Temp, Sulphur and Temp x Sulphur interaction effects. Levels of  
5 significance: ns non-significant,  $p < 0.05^*$ ,  $p < 0.01^{**}$ ,  $p < 0.001^{***}$ . SE: standard error, DW: dry weight.

1 **Supplemental Data**

Period covering the 17 day of heat treatment	Daily PARi in the heated unit (MJ/day)	Daily PARi in the control unit (MJ/day)	Ratio
April 27 <sup>th</sup>	5.1	7.4	1.4
April 28 <sup>th</sup>	5.2	7.2	1.4
April 29 <sup>th</sup>	3.3	3.7	1.1
April 30 <sup>th</sup>	5.0	5.9	1.2
May 1 <sup>st</sup>	5.5	7.1	1.3
May 2 <sup>nd</sup>	6.7	7.1	1.0
May 3 <sup>rd</sup>	5.3	5.3	1.0
May 4 <sup>th</sup>	5.8	5.8	1.0
May 5 <sup>th</sup>	3.5	3.4	1.0
May 6 <sup>th</sup>	6.5	7.5	1.1
May 7 <sup>th</sup>	4.4	5.0	1.1
May 8 <sup>th</sup>	4.4	10.1	2.3
May 9 <sup>th</sup>	4.8	9.8	2.0
May 10 <sup>th</sup>	4.8	4.5	0.9
May 11 <sup>th</sup>	5.7	5.5	1.0
May 12 <sup>th</sup>	6.7	5.4	0.8
May 13 <sup>th</sup>	6.1	9.4	1.5
May 14 <sup>th</sup>	5.1	7.4	1.4
May 15 <sup>th</sup>	5.2	7.2	1.4

2

3

4 **Table S1:** Daily incident Photosynthetically Active Radiation values (PARi, MJ/day) in the heated  
5 unit and in the control unit throughout the 17 days of Temp- modality application (from the 27<sup>th</sup> of  
6 April until the 15<sup>th</sup> of May).

7

8

9

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19


Harvest date	Harvest 1	Harvest 2
Mean Seed FW	3.31 (0.76)	5.98 (0.99)
p-value	<b>0.02*</b>	

**Table S2:** Mean individual seed (*n*) fresh weight (FW, mg/seed) at the beginning (Harvest 1, *n*=212) and at the end of Temp-modality exposure (Harvest 2, *n*=250) over 17 days. FWs are given for Temp-control plants. Standard deviation into brackets. Level of significance of the ANOVA test for the date effect:  $p < 0.05^*$ .


Sulphur condition	High Sulphur					Low Sulphur					Temp effect	Sulphur effect	Temp x Sulphur
	Control	Early Mild Stress	3 Late Heat Peaks	4 Late Heat Peaks	Priming	Control	Early Mild Stress	3 Late Heat Peaks	4 Late Heat Peaks	Priming			
<b>Yield components (per plant)</b>													
Total Seed Yield (g)	9.92 (0.57)	8.82 (0.82)	9.55 (0.74)	9.61 (0.55)	7.52 (0.91)	9.70 (1.35)	8.71 (0.64)	8.77 (1.61)	8.95 (0.46)	8.78 (1.87)	<b>0.04*</b>	0.77	0.34
Total Seed Number	2774 (203)	2672 (184)	2802 (249)	2882 (192)	2519 (168)	2794 (328)	2457 (253)	2643 (183)	2565 (121)	2474 (124)	0.07	<b>0.04*</b>	0.57
Thousand Seed Weight (g)	3.53 (0.29)	3.21 (0.34)	3.42 (0.34)	3.35 (0.28)	2.85 (0.65)	3.36 (0.46)	3.40 (0.34)	3.17 (0.62)	3.31 (0.15)	3.54 (1.51)	0.87	0.98	0.09
<b>Seed C, N and S concentrations (%DW)</b>													
Carbon	59.8 (0.7)	58.1 (0.9)	57.9 (1.7)	57.3 (1.6)	55.4 (1.3)	58.7 (1.8)	58.3 (1.2)	58.4 (1.9)	57.0 (1.0)	56.6 (1.8)	<b>0.00***</b>	0.83	0.61
Nitrogen	3.36 (0.23)	3.44 (0.22)	3.22 (0.34)	3.32 (0.24)	3.97 (0.19)	3.11 (0.50)	3.51 (0.09)	3.18 (0.20)	3.46 (0.25)	3.79 (0.33)	<b>0.00***</b>	0.55	0.57
Sulfur	0.39 (0.05)	0.38 (0.05)	0.39 (0.05)	0.40 (0.04)	0.42 (0.02)	0.24 (0.04)	0.27 (0.03)	0.26 (0.01)	0.24 (0.01)	0.33 (0.04)	<b>0.01*</b>	<b>0.00***</b>	0.30
<b>Oil contents and fatty acids (%DW)</b>													
Total Fatty Acid	39.3 (4.5)	36.4 (1.1)	37.3 (1.1)	31.2 (4.8)	30.1 (3.2)	39.2 (8.1)	44.5 (1.0)	45.1 (4.1)	42.6 (5.0)	35.0 (1.9)	<b>0.00**</b>	<b>0.00***</b>	0.11
$\omega 6:\omega 3$ ratio	2.03 (0.12)	1.83 (0.10)	2.05 (0.11)	2.08 (0.20)	2.29 (0.21)	2.07 (0.29)	1.91 (0.08)	2.11 (0.14)	2.33 (0.21)	2.19 (0.09)	<b>0.00**</b>	0.22	0.38
<b>Protein concentration (mg/g DW)</b>													
Seed Storage Protein	91.0 (15.2)	149.9 (55.6)	79.9 (48.7)	93.2 (13.0)	118.3 (16.6)	208.3 (48.4)	186.3 (66.7)	192.5 (36.2)	153.7 (38.2)	182.6 (33.7)	0.28	<b>0.00***</b>	0.23
<b>Soluble sugars concentrations (%DW)</b>													
Sucrose	6.52 (0.80)	2.95 (0.25)	6.42 (2.17)	5.04 (1.76)	3.69 (0.82)	8.28 (0.84)	2.99 (0.88)	5.80 (2.10)	4.14 (1.06)	3.02 (0.38)	<b>0.00***</b>	0.85	0.25
Raffinose	0.44 (0.08)	0.21 (0.02)	0.41 (0.13)	0.30 (0.13)	0.25 (0.07)	0.50 (0.05)	0.21 (0.05)	0.35 (0.15)	0.21 (0.05)	0.17 (0.02)	<b>0.00***</b>	0.85	0.38
Stachyose	2.35 (0.38)	1.15 (0.15)	2.09 (0.52)	1.51 (0.58)	1.40 (0.41)	2.71 (0.25)	1.14 (0.32)	2.11 (0.76)	1.21 (0.31)	1.00 (0.10)	<b>0.00***</b>	0.62	0.43
[raff+stach]:sucr	0.43 (0.02)	0.46 (0.02)	0.40 (0.03)	0.35 (0.03)	0.44 (0.04)	0.39 (0.02)	0.45 (0.03)	0.42 (0.02)	0.34 (0.01)	0.39 (0.02)	<b>0.00***</b>	0.05	0.06
<b>Seed conductivity (<math>\mu</math>S/mg DW)</b>													
Seed Conductivity	0.87 (0.45)	1.28 (0.73)	1.19 (0.55)	0.97 (0.50)	2.86 (1.57)	0.78 (0.16)	0.66 (0.13)	0.81 (0.27)	1.08 (0.78)	3.01 (0.93)	<b>0.00***</b>	0.34	0.66
<b>Hormone concentrations (ng/g DW)</b>													
Indole-3-Acetic Acid	134.3 (1.2)	121.7 (1.1)	278.2 (6.6)	380.3 (1.8)	268.9 (4.3)	160.1 (6.9)	211.5 (1.9)	164.6 (1.2)	279.5 (3.6)	192.7 (1.9)	<b>0.00***</b>	<b>0.00***</b>	<b>0.00***</b>
Salicylic Acid	40.8 (0.6)	42.6 (0.3)	96.6 (1.6)	135.6 (0.5)	103.1 (2.5)	59.3 (2.6)	76.1 (0.7)	73.5 (0.9)	82.5 (0.9)	96.1 (1.1)	<b>0.00***</b>	<b>0.00***</b>	<b>0.00***</b>
ABA:GA3 ratio	0.28 (0.01)	0.41 (0.00)	0.42 (0.01)	0.49 (0.01)	0.74 (0.01)	0.79 (0.03)	0.60 (0.01)	0.95 (0.00)	0.91 (0.01)	0.63 (0.02)	<b>0.00***</b>	<b>0.00***</b>	<b>0.00***</b>
<b>Phenolic content and antioxidant capacity</b>													
Polyphenols	3.96 (0.15)	4.12 (0.18)	3.79 (0.23)	3.44 (0.49)	3.76 (0.30)	3.34 (0.48)	4.26 (0.20)	3.35 (0.26)	3.71 (0.36)	4.11 (0.15)	<b>0.00**</b>	0.60	<b>0.01*</b>
ABST	18.70 (1.04)	19.14 (0.87)	18.18 (1.93)	15.66 (1.97)	17.72 (0.76)	15.29 (1.31)	18.68 (1.91)	15.47 (0.96)	16.40 (1.75)	18.17 (0.84)	<b>0.00**</b>	<b>0.01*</b>	<b>0.00**</b>

1

2 **Table S3:** Summary of the mean value  $\pm$  se of the all measured variables for each of the 10 treatments in order to provide reference values in a given  
3 combination “Sulphur supply/Temperature modality”. Standard deviation into brackets. P-values and levels of significance are given for Temp, Sulphur and  
4 Temp x Sulphur interaction effects. Levels of significance: ns non-significant,  $p < 0.05^*$ ,  $p < 0.01^{**}$ ,  $p < 0.001^{***}$ . DW: dry weight.



Pearson correlation:  
 $R=0,9309$   
 $P<0.0001$



Pearson correlation:  
 $R=0,6387$   
 $P=0,0468$

1

2

3 **Figure S1:** Illustration of the correlation between ABTS value related to (a) polyphenols and (b)

4 soluble sugars ratio. SA: synaptic acid, ABTS: 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid).