



HAL
open science

Stabilization of the Wave Equation through Nonlinear Dirichlet Actuation

Nicolas Vanspranghe, Francesco Ferrante, Christophe Prieur

► **To cite this version:**

Nicolas Vanspranghe, Francesco Ferrante, Christophe Prieur. Stabilization of the Wave Equation through Nonlinear Dirichlet Actuation. 2021. hal-03363940v1

HAL Id: hal-03363940

<https://hal.science/hal-03363940v1>

Preprint submitted on 4 Oct 2021 (v1), last revised 26 Aug 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

STABILIZATION OF THE WAVE EQUATION THROUGH NONLINEAR DIRICHLET ACTUATION *

NICOLAS VANSPRANGHE¹, FRANCESCO FERRANTE² AND CHRISTOPHE PRIEUR¹

Abstract. In this paper, we consider the problem of nonlinear (in particular, saturated) stabilization of the high-dimensional wave equation in the Dirichlet boundary condition. The wave dynamics are subject to a dissipative nonlinear velocity feedback and generate a strongly continuous semigroup of contractions on the optimal energy space $L^2(\Omega) \times H^{-1}(\Omega)$. It is first proved that any solution to the closed-loop equations converges to zero in the aforementioned topology. Secondly, under the condition that the feedback nonlinearity has linear growth around zero, polynomial energy decay rates are established for solutions with smooth initial data. This constitutes new Dirichlet counterparts to well-known results pertaining to nonlinear stabilization in $H^1(\Omega) \times L^2(\Omega)$ of the wave equation in the Neumann boundary conditions.

1991 Mathematics Subject Classification. 35L05, 93C20, 93D15, 93D20.

The dates will be set by the publisher.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^d$ ($d \geq 2$) be a bounded domain with smooth boundary Γ . Given a relatively open nonempty subset Γ_0 of Γ , we consider the wave equation subject to non-homogeneous Dirichlet boundary condition:

$$\partial_{tt}u(x, t) - \Delta u(x, t) = 0 \quad \text{in } \Omega \times (0, +\infty), \quad (1.1a)$$

$$u|_{\Gamma}(\sigma, t) = -g(U(\sigma, t)) \quad \text{on } \Gamma_0 \times (0, +\infty), \quad (1.1b)$$

$$u|_{\Gamma}(\sigma, t) = 0 \quad \text{on } \Gamma_1 \times (0, +\infty), \quad (1.1c)$$

where $\Gamma_1 \triangleq \Gamma \setminus \Gamma_0$, U represents a control input, and g is a real nondecreasing function fulfilling the following assumption.

Assumption 1. The scalar mapping g satisfies the following properties:

- (i) g is globally Lipschitz continuous and nondecreasing;
- (ii) $g(s) = 0$ if and only if $s = 0$.

Keywords and phrases: Wave equation, boundary stabilization, saturating feedback, asymptotic stability, polynomial decay rates.

* This work has been partially supported by MIAI@Grenoble Alpes (ANR-19-P3IA-0003). A preliminary version of this work, containing the well-posedness and asymptotic stability results, has appeared in the Proceedings of the Third IFAC Conference on Modelling, Identification and Control of Nonlinear Systems, Tokyo, Japan, September 2021.

¹ Univ. Grenoble Alpes, CNRS, Grenoble INP, GIPSA-lab, 38000 Grenoble, France; e-mail: name.surname@gipsa-lab.fr.

² Department of Engineering, University of Perugia, 06125 Perugia, Italy; e-mail: francesco.ferrante@unipg.it.

Background. The general problem of this paper is the feedback stabilization of the control system (1.1) in presence of a static pointwise nonlinearity g . Consider the velocity feedback

$$U(\sigma, t) = -\partial_\nu[A^{-1}u'](\sigma, t), \quad (1.2)$$

where ∂_ν denotes the outward normal derivative and A^{-1} is the inverse of the positive “minus Laplacian with homogeneous Dirichlet boundary condition” operator. The corresponding linear feedback system (i.e., when g is the identity) was first introduced by Lasiecka and Triggiani in [9], with initial data lying in the energy space

$$\mathcal{H} \triangleq L^2(\Omega) \times H^{-1}(\Omega). \quad (1.3)$$

The choice of state space is motivated by optimal regularity results for second-order hyperbolic equations with Dirichlet boundary data in $L^2(0, +\infty; L^2(\Gamma))$ – see [11]. It was proved in [9] that the linear version of (1.1)-(1.2) gives rise to an exponentially stable semigroup of operators on \mathcal{H} under the assumption that the whole boundary is actuated (i.e., $\Gamma = \Gamma_0$) and that Ω satisfies suitable geometrical conditions. The proof relies on the analysis of a new variable p defined as

$$p \triangleq A^{-1}u' \quad (1.4)$$

which is smoother and solves a wave-type equation as well. The result was later refined by the same authors in [10] where feedback acting only on a subset of the boundary is allowed and, most importantly, specific geometrical conditions related to the analysis of the p -variable by multipliers are relaxed. This was achieved by the mean of another change of variable operating at the level of pseudodifferential calculus. In short, after transposing problem (1.1)-(1.2) to the half-space via partition of unity and truncating the solution with respect to the time variable, one defines a new variable w by

$$\mathcal{F}[w](\xi, \omega; x) = \lambda(\xi, \omega)\mathcal{F}[u](\xi, \omega; x), \quad \xi \in \mathbb{R}^{d-1}, \quad \omega \in \mathbb{R}, \quad x \geq 0, \quad (1.5)$$

where \mathcal{F} denotes the Fourier transform in both tangential and time variables and λ is a carefully constructed symbol. While transformations (1.4) and (1.5) are quite different in nature, both enable computations on variables with $H^1(\Omega) \times L^2(\Omega)$ -regularity.

As far as we know, there has been no attempt to extend the stability analysis of the closed-loop system (1.1)-(1.2) to the *nonlinear* case. Yet, one can see problem (1.1)-(1.2) as a natural Dirichlet counterpart to the wave equation with nonlinear Neumann boundary dissipation

$$\partial_{tt}u(x, t) - \Delta u(x, t) = 0 \quad \text{in } \Omega \times (0, +\infty) \quad (1.6a)$$

$$\partial_\nu u(\sigma, t) = -g(\partial_t u(\sigma, t)) \quad \text{on } \Gamma_0 \times (0, +\infty), \quad (1.6b)$$

$$u|_\Gamma(\sigma, t) = 0 \quad \text{on } \Gamma_1 \times (0, +\infty), \quad (1.6c)$$

which, in contrast, have been extensively studied in the literature. To cite only a few, when the nonlinearity g has linear growth at infinity, *uniform* decay of the $H^1(\Omega) \times L^2(\Omega)$ -energy of solutions to (1.6) can be achieved, as in [20] or [12] – see also [8] and the references therein, or more recently [5]. In the one-dimensional settings, arguments based on Riemann invariants are available, and the decay of the energy can be analyzed via appropriate iterated sequences. See for instance [1], where g is allowed to be a multivalued monotone mapping, or [18], where it is proved, in particular, that exponential or polynomial uniform decay cannot be achieved when g represents a pointwise saturation mapping – see also [19] or [14] for a stability analysis in the saturated case.

Outline of the paper and contributions. This paper aims at bridging the gap between Neumann and Dirichlet boundary conditions as far as nonlinear boundary stabilization is concerned. First, we prove that the nonlinear dynamics (1.1)-(1.2) generate a strongly continuous semigroup of contractions on the energy space \mathcal{H} (Theorem 2.1) that is globally asymptotically stable around the zero equilibrium (Theorem 2.2). The proof relies on LaSalle’s invariance principle and unique continuation for the wave equation.

Next, having in mind the more specific problem of *saturated* boundary stabilization, in Section 3, we work under the assumption that g has linear growth around zero (see Assumption 2 below). Then, by analogy with

the Neumann case, we focus on *non-uniform* decay rates for solutions with “smooth” initial data. We establish a polynomial decay rate for smooth solutions (Theorem 3.1) that holds under standard geometrical conditions – see Assumption 3 below, which is however always satisfied when $\Gamma_0 = \Gamma$, i.e., the whole boundary is actuated. To do so, we consider the change of variable (1.4) and we derive appropriate integral inequalities using suitable multipliers. Note that the question of uniform stability when g has linear growth at infinity is out of the scope of the paper – this is discussed in Section 4 below. Throughout the paper, one can think of the “hard” saturation mapping sat_S with threshold $S > 0$, defined by

$$\text{sat}_S(s) \triangleq \begin{cases} s & \text{if } |s| \leq S, \\ S \frac{s}{|s|} & \text{otherwise,} \end{cases} \quad (1.7)$$

as a prototype nonlinearity satisfying Assumptions 1 and 2, highlighting the fact that no differentiability condition on g is prescribed.

Notation and elements from elliptic theory. We end this section by introducing some notation and recalling useful results from elliptic theory.

First, if H is a given Hilbert space, we denote by $\|\cdot\|_H$ its norm, and its scalar product is written $(\cdot, \cdot)_H$. For $T > 0$, we denote by $W^{1,p}(0, T; H)$ the subspace of $L^p(0, T; H)$ composed of (classes of) H -valued functions ϕ such that, for some ξ in H and ψ in $L^p(0, T; H)$, $\phi(t) = \xi + \int_0^t \psi(s) ds$ almost everywhere in $(0, T)$. Such class ϕ is identified with its continuous representative and we say that $\phi' = \psi$ in the sense of H -valued distributions. Note that vector-valued integrals are intended in the sense of Bochner. Also, the space of bounded linear operators between two normed spaces E and F is denoted by $\mathcal{L}(E, F)$.

In this paper, all scalar functions are real-valued. The notation dx indicates the standard Lebesgue measure on \mathbb{R}^d while $d\sigma$ denotes the induced surface measure on Γ . By $H^s(\Omega)$ (resp. $H^s(\Gamma)$) we denote the $L^2(\Omega)$ -based (resp. $L^2(\Gamma)$ -based) Sobolev space of order s . The space of compactly supported and infinitely differentiable functions on Ω is written $C_c^\infty(\Omega)$. We also recall that $H_0^1(\Omega)$ is defined as the closure of $C_c^\infty(\Omega)$ in $H^1(\Omega)$. Furthermore, $H^{-1}(\Omega)$ is the topological dual of $H_0^1(\Omega)$.

The unbounded operator A can be defined¹ as follows: having set

$$\mathcal{D}(A) \triangleq H^2(\Omega) \cap H_0^1(\Omega), \quad (1.8)$$

we let $Au \triangleq -\Delta u \in L^2(\Omega)$ for all $u \in \mathcal{D}(A)$. Then, A is a strictly positive self-adjoint densely defined operator on $L^2(\Omega)$. Its domain $\mathcal{D}(A)$ is equipped with the norm $\|A \cdot\|_{L^2(\Omega)}$, which is equivalent to the norm induced by $H^2(\Omega)$. As a positive operator, A possesses fractional powers A^s , $s \in \mathbb{R}$ – see for instance [17, Chapter II, Section 2.1]. Those are also strictly positive self-adjoint operators. For $s \geq 0$, $\mathcal{D}(A^s)$ are dense subsets of $L^2(\Omega)$, which we equip with the norm $\|A^s \cdot\|_{L^2(\Omega)}$. In particular, we have $\mathcal{D}(A^{1/2}) = H_0^1(\Omega)$, with

$$\|\nabla w\|_{L^2(\Omega)}^2 = \|A^{1/2} w\|_{L^2(\Omega)}^2 = \|w\|_{H_0^1(\Omega)}^2 \quad \text{for all } w \in H_0^1(\Omega). \quad (1.9)$$

Then, we let $\mathcal{D}(A^{-s}) \triangleq \mathcal{D}(A^s)'$ and we can extend A^s as an isomorphism between $L^2(\Omega)$ and $\mathcal{D}(A^{-s})$. Here, $H^{-1}(\Omega)$ is equipped with the scalar product

$$(v_1, v_2)_{H^{-1}(\Omega)} \triangleq (A^{-1/2} v_1, A^{-1/2} v_2)_{L^2(\Omega)}, \quad (1.10)$$

which induces a norm equivalent to the dual one; we also recover $\mathcal{D}(A^{-1/2}) = H^{-1}(\Omega)$. Throughout the paper, we use the following chain of continuous embeddings:

$$\mathcal{D}(A) \hookrightarrow H_0^1(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow H^{-1}(\Omega) \hookrightarrow \mathcal{D}(A^{-1}). \quad (1.11)$$

¹Alternatively, A can be defined as a duality mapping between $H_0^1(\Omega)$ and $H^{-1}(\Omega)$, in which case (1.8) is recovered *a posteriori* by applying elliptic regularity theory.

Finally, we define the Dirichlet map D , which is a right inverse for the trace operator. For any f in $H^{1/2}(\Gamma)$, there exists a unique u in $H^1(\Omega)$ solving $-\Delta u = 0$ and $u|_\Gamma = f$; and we let $Df \triangleq u$. The mapping D defined in this way belongs to $\mathcal{L}(H^{1/2}(\Omega), H^1(\Omega))$; also, we define its adjoint D^* by $(D^*u, f)_{L^2(\Gamma)} = (u, Df)_{L^2(\Omega)}$ for all u in $L^2(\Omega)$ and f in $L^2(\Gamma)$. Extensions on fractional Sobolev spaces are denoted with the same symbols:

$$D \in \mathcal{L}(H^s(\Gamma), H^{s+1/2}(\Omega)), \quad D^* \in \mathcal{L}(H^s(\Omega), H^{s+1/2}(\Gamma)) \quad \text{for all } s \in \mathbb{R}. \quad (1.12)$$

2. WELL-POSEDNESS AND ASYMPTOTIC STABILITY

In this section, we give the operator-theoretic formulation of the evolution problem (1.1a) with initial data in $L^2(\Omega) \times H^{-1}(\Omega)$ and feedback control (1.2). After that, we state and prove the well-posedness and asymptotic stability properties of the feedback system (1.1)-(1.2).

2.1. Operator model and well-posedness

We shall recast the closed-loop evolution equations (1.1)-(1.2) into a first-order abstract Cauchy problem on the energy space \mathcal{H} and state well-posedness results by relying on nonlinear semigroup theory.

With a little abuse of notation, we denote by g the (nonlinear) Lipschitz mapping on $L^2(\Gamma)$ defined by $g(f)(\sigma) \triangleq g(f(\sigma))$ for any f in $L^2(\Gamma)$. We also define a projection operator P on $L^2(\Gamma)$ by $[Pf](\sigma) = \mathbb{1}_{\Gamma_0}(\sigma)f(\sigma)$ for any $f \in L^2(\Gamma)$. From the Green formula, it follows that

$$-D^*Ap = \partial_\nu p \quad \text{for all } p \in \mathcal{D}(A). \quad (2.1)$$

Therefore, the boundary condition associated with the feedback law (1.2) can be rewritten as follows:

$$u|_\Gamma = -Pg(D^*u'). \quad (2.2)$$

Next, we introduce the unbounded nonlinear operator \mathcal{A} associated with the closed-loop system (1.1)-(1.2). Recalling the chain of embeddings (1.11) and that A maps $L^2(\Omega)$ onto $\mathcal{D}(A^{-1})$, we define \mathcal{A} by

$$\mathcal{D}(\mathcal{A}) \triangleq \{[u, v] \in \mathcal{H} : v \in L^2(\Omega), A[u + DPg(D^*v)] \in H^{-1}(\Omega)\} \quad (2.3a)$$

$$\mathcal{A}[u, v] \triangleq [-v, Au + ADPg(D^*v)]. \quad (2.3b)$$

Equivalently, $\mathcal{D}(\mathcal{A})$ is the set of all $[u, v]$ in $L^2(\Omega) \times L^2(\Omega)$ such that $u + DPg(D^*v)$ belongs to $H_0^1(\Omega)$.

Note that the feedback law (1.2) appears as a natural choice when (formally) differentiating the energy functional

$$\mathcal{E}(u, v) \triangleq \frac{1}{2} \{ \|u\|_{L^2(\Omega)}^2 + \|v\|_{H^{-1}(\Omega)}^2 \}, \quad [u, v] \in \mathcal{H}, \quad (2.4)$$

along ‘‘trajectories’’ of the open-loop system (1.1). Indeed, this leads to the energy identity

$$\frac{d}{dt} \mathcal{E}(u, u') = \int_{\Gamma_0} g(U(t)) \partial_\nu [A^{-1}u'] \, d\sigma \quad (2.5)$$

and since g satisfies Assumption 1, we see that (1.2) renders the energy \mathcal{E} nonincreasing along the trajectories.

In the sequel, we employ the standard nonlinear semigroup terminology: by a *strong* solution to (1.1)-(1.2), we mean an absolutely continuous \mathcal{H} -valued function $[u, v]$ that satisfies $[u(t), v(t)] \in \mathcal{D}(\mathcal{A})$ for all $t \geq 0$ and

$$\frac{d}{dt} [u, v] + \mathcal{A}[u, v] = 0 \quad \text{a.e.} \quad (2.6)$$

in the sense of strong differentiation in \mathcal{H} ; by a *generalized* solution to (1.1)-(1.2), we mean a continuous \mathcal{H} -valued function $[u, v]$ that is, on each interval $[0, T]$, the uniform limit of some sequence of strong solutions.

Theorem 2.1 (Hadamard well-posedness). *The nonlinear operator \mathcal{A} is densely defined and maximal monotone. Thus, $-\mathcal{A}$ is the infinitesimal generator of a strongly continuous semigroup $\{\mathcal{S}_t\}$ of (nonlinear) contractions on the energy space \mathcal{H} . For all initial data $[u_0, v_0]$ in \mathcal{H} , there exists a unique generalized solution $[u, u'] \in \mathcal{C}(\mathbb{R}^+, \mathcal{H})$ to (1.1)-(1.2). If $[u_0, v_0]$ belongs to $\mathcal{D}(\mathcal{A})$, then $[u, u']$ is a strong solution to (1.1)-(1.2). Furthermore,*

(i) *Strong solutions satisfy the inequality*

$$\|\mathcal{A}[u(t), u'(t)]\|_{\mathcal{H}} \leq \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}} \quad \text{for all } t \geq 0; \quad (2.7)$$

(ii) *Strong solutions satisfy the energy identity*

$$\frac{d}{dt} \mathcal{E}(u, u') = - \int_{\Gamma_0} g(D^* u') D^* u' \, d\sigma = \int_{\Gamma_0} g(-\partial_\nu[A^{-1} u']) \partial_\nu[A^{-1} u'] \, d\sigma \quad (2.8)$$

in the scalar distribution sense on $(0, +\infty)$.

Remark 2.1. If we also assume that, say, $|g(s)| \geq \alpha|s|$ for all $s \in \mathbb{R}$ and some $\alpha > 0$, the energy identity (2.8) provides a uniform estimate of the $L^2(0, +\infty; L^2(\Gamma_0))$ -norm of $\partial_\nu[A^{-1} u']$ for strong solutions. From there, one can prove that (2.8) holds for generalized solution as well by passing to the limit and recovering the traces $u|_\Gamma$ and $\partial_\nu[A^{-1} u']$ in $L^2(0, +\infty; L^2(\Gamma_0))$ – see for instance [2] for similar arguments in the Neumann boundary conditions.

Proof of Theorem 2.1. Once proven that \mathcal{A} is maximal monotone, existence and uniqueness of strong and generalized solutions to (1.1)-(1.2), together with the appropriate semigroup properties, follow from Kato's theorem and standard nonlinear semigroup theory – see, e.g., [16, Chapter IV].

Step 1: Monotonicity. Let $[u_1, v_1]$ and $[u_2, v_2]$ in $\mathcal{D}(\mathcal{A})$. Then,

$$\begin{aligned} (\mathcal{A}[u_1, v_1] - \mathcal{A}[u_2, v_2], [u_1, v_1] - [u_2, v_2])_{\mathcal{H}} &= -(v_1 - v_2, u_1 - u_2)_{L^2(\Omega)} \\ &+ (A^{1/2}[u_1 - u_2 + DPg(D^* v_1) - DPg(D^* v_2)], A^{-1/2}[v_1 - v_2])_{L^2(\Omega)}. \end{aligned} \quad (2.9)$$

Now we use that $A^{-1/2}[v_1 - v_2]$ belongs to $\mathcal{D}(A^{1/2})$ and that $A^{1/2}$ is self-adjoint to obtain

$$\begin{aligned} (\mathcal{A}[u_1, v_1] - \mathcal{A}[u_2, v_2], [u_1, v_1] - [u_2, v_2])_{\mathcal{H}} &= (DPg(D^* v_1) - DPg(D^* v_2), v_1 - v_2)_{L^2(\Omega)} \\ &= (Pg(D^* v_1) - Pg(D^* v_2), D^* v_1 - D^* v_2)_{L^2(\Gamma)} \\ &= (g(D^* v_1) - g(D^* v_2), D^* v_1 - D^* v_2)_{L^2(\Gamma_0)} \geq 0, \end{aligned} \quad (2.10)$$

the right-hand side being nonnegative by nondecreasingness of g , which proves that \mathcal{A} is monotone.

Step 2: Range condition. Let $\lambda > 0$ and $[f_1, f_2] \in \mathcal{H}$. To solve the equation $\mathcal{A}[u, v] + \lambda[u, v] = [f_1, f_2]$, it suffices to find $v \in L^2(\Omega)$ such that

$$\lambda^{-1} v + DPg(D^* v) + \lambda A^{-1} v = A^{-1} f_2 - \lambda^{-1} f_1. \quad (2.11)$$

This is seen by substituting $-v + \lambda u = f_1$ into the second coordinate of the equation and applying A^{-1} to the result. If such $v \in L^2(\Omega)$ is found, then $[u, v]$ belongs to $\mathcal{D}(\mathcal{A})$ since $A^{-1} f_2 \in H_0^1(\Omega)$ and $DPg(D^* v) \in H^1(\Omega)$.

We define a nonlinear operator Θ on $L^2(\Omega)$ by $\Theta(v) \triangleq \lambda^{-1} v + DPg(D^* v) + \lambda A^{-1} v$ for all $v \in L^2(\Omega)$. Then, Θ enjoys the following properties:

- (i) Θ maps bounded sets into bounded sets;
- (ii) $(\Theta(v_1) - \Theta(v_2), v_1 - v_2)_{L^2(\Omega)} \geq 0$ for all v_1 and v_2 in $L^2(\Omega)$;
- (iii) The scalar function $t \mapsto (\Theta(v_1 + tv_2), v_2)_{L^2(\Omega)}$ is continuous for all v_1 and v_2 in $L^2(\Omega)$.

Also, we have

$$(\Theta(v), v)_{L^2(\Omega)} \geq \lambda^{-1} \|v\|_{L^2(\Omega)}^2 \quad \text{for all } v \in L^2(\Omega). \quad (2.12)$$

Thus, it follows from [16, Lemma 2.1 and Theorem 2.1] that Θ is onto. Consequently, the equation $\mathcal{A}[u, v] + \lambda[u, v] = [f_1, f_2]$ has a solution in $\mathcal{D}(\mathcal{A})$.

Step 3: Denseness of the domain. Let $[u, v] \in \mathcal{H}$ and $\epsilon > 0$. Since $A^{-1}v \in H_0^1(\Omega)$ and $C_c^\infty(\Omega)$ is dense in $H_0^1(\Omega)$, we can pick $\phi \in \mathcal{D}(\Omega)$ such that

$$\|A^{-1}v - \phi\|_{H_0^1(\Omega)}^2 \leq \epsilon, \quad \text{and thus } \|v - A\phi\|_{H^{-1}(\Omega)}^2 \leq C\epsilon \quad (2.13)$$

where $C > 0$ comes from $A \in \mathcal{L}(H_0^1(\Omega), H^{-1}(\Omega))$. Besides, there exists $\psi \in C_c^\infty(\Omega)$ such that $\|u - \psi\|_{L^2(\Omega)}^2 \leq \epsilon$. Since $\phi \in C_c^\infty(\Omega) \subset \mathcal{D}(A)$, we have $A\phi \in L^2(\Omega)$ and also, using (2.1), $g(D^*A\phi) = g(-\partial_\nu \phi) = 0$. Thus, $[\psi, A\phi] \in \mathcal{D}(\mathcal{A})$; also, we have $\|[u, v] - [\psi, A\phi]\|_{\mathcal{H}}^2 \leq (1 + C)\epsilon$. It is now proved that $\mathcal{D}(\mathcal{A})$ is dense in \mathcal{H} .

Step 4: Energy identity. Let $[u, v]$ be a strong solution to (1.1)-(1.2). We recall that $\mathcal{E}(u, u') = \frac{1}{2}\|[u, u']\|_{\mathcal{H}}^2$. Consequently, by the chain rule, $\mathcal{E}(u, u')$ is an absolutely continuous scalar function and

$$\frac{d}{dt}\mathcal{E}(u, u') = (\mathcal{A}[u, u'], [u, u'])_{\mathcal{H}} \quad \text{a.e.} \quad (2.14)$$

Thus, the desired identity (2.8) follows from (2.10). \square

2.2. Additional properties of the semigroup

Now, we establish some compactness and regularity properties that are useful in the proof of the stability results presented in Subsections 2.3 and 3.1. We start by introducing the following proposition, which enables us to prove asymptotic stability of the feedback system (1.1)-(1.2) using LaSalle's invariance principle.

Proposition 2.1 (Compactness). *For any $\lambda > 0$, the (nonlinear) resolvent operator $(\mathcal{A} + \lambda \text{id})^{-1}$ is well-defined on \mathcal{H} and compact. In particular, for all initial data $[u_0, v_0] \in \mathcal{H}$, the (semi)trajectory $\{\mathcal{S}_t[u_0, v_0]\}_{t \geq 0}$ is relatively compact in \mathcal{H} .*

Proof. Assume for a moment that $(\mathcal{A} + \lambda \text{id})^{-1}$ is well-defined and compact for some $\lambda > 0$. Then, since $\mathcal{A}(0) = 0$, relative compactness of the trajectories follows from [4, Theorem 3].

Let $\lambda > 0$. We already know from the proof of Theorem 2.1 that the equation

$$\mathcal{A}[u, v] = [f_1, f_2] \quad (2.15)$$

has a solution in $\mathcal{D}(\mathcal{A})$ for all $[f_1, f_2] \in \mathcal{H}$.

Step 1: Uniqueness. Consider two solutions $[u_1, v_1]$ and $[u_2, v_2]$ to (2.15). Then, we recall from (2.11) in the proof of Theorem 2.1 that

$$\lambda^{-1}[v_1 - v_2] + DP[g(D^*v_1) - g(D^*v_2)] + \lambda A^{-1}[v_1 - v_2] = 0. \quad (2.16)$$

Taking the scalar product in $L^2(\Omega)$ of (2.16) with $v_1 - v_2$ yields

$$\lambda^{-1}\|v_1 - v_2\|_{L^2(\Omega)}^2 + (g(D^*v_1) - g(D^*v_2), v_1 - v_2)_{L^2(\Gamma_0)} + \lambda\|v_1 - v_2\|_{H^{-1}(\Omega)}^2 = 0. \quad (2.17)$$

In particular, since g is nondecreasing, we infer from (2.17) that $v_1 = v_2$; thus, $[u_1, v_1] = [u_2, v_2]$ and $(\mathcal{A} + \lambda \text{id})^{-1}$ is well-defined.

Step 2: Compactness of the resolvent operator. In what follows, we let $[u, v] \triangleq (\mathcal{A} + \lambda \text{id})^{-1}[f_1, f_2]$ and we look for estimates of $[u, v] \in \mathcal{D}(\mathcal{A})$ in stronger norms. First, as in the previous step, we obtain

$$\lambda^{-1}\|v\|_{L^2(\Omega)}^2 + (g(D^*v), v)_{L^2(\Gamma_0)} + \lambda\|v\|_{H^{-1}(\Omega)}^2 = (A^{-1/2}f_2, A^{-1/2}v)_{L^2(\Omega)} - \lambda^{-1}(f_1, v)_{L^2(\Omega)}, \quad (2.18)$$

where it is used that $A^{-1/2}$ is self-adjoint. From (2.18), using Cauchy-Schwarz and Young inequalities with appropriate constants, we obtain the estimate

$$\|v\|_{L^2(\Omega)}^2 \leq \frac{1}{2}\|f_2\|_{H^{-1}(\Omega)}^2 + \|f_1\|_{L^2(\Omega)}^2. \quad (2.19)$$

The remainder of the proof relies on elliptic regularity theory and in particular [13, Théorème 10.1]. Since $[u, v] \in \mathcal{D}(\mathcal{A})$, we know that $u + DPg(D^*v)$ belongs to $H_0^1(\Omega)$ and

$$u + DPg(D^*v) = A^{-1}f_2 - \lambda A^{-1}v. \quad (2.20)$$

Picking an arbitrary test-function ϕ in $\mathcal{C}_c^\infty(\Omega) \subset \mathcal{D}(A)$, taking the scalar product in $L^2(\Omega)$ of (2.20) with $A\phi$ and using again that $D^*A\phi = -\partial_\nu\phi = 0$ leads to $-\Delta u = f_2 - \lambda v$ in the sense of distributions on Ω . Besides, since $-\Delta u \in H^{-1}(\Omega)$, $u|_\Gamma$ is well-defined in $H^{-1/2}(\Gamma)$; then, we infer from $u + DPg(D^*v) \in H_0^1(\Omega)$ that $u|_\Gamma = -Pg(D^*v) \in L^2(\Gamma)$. Applying the aforementioned theorem, we obtain $u \in H^{1/2}(\Omega)$ along with the estimate

$$\|u\|_{H^{1/2}(\Omega)}^2 \leq C_1 \left\{ \|Pg(D^*v)\|_{L^2(\Gamma)}^2 + \|f_2 - \lambda v\|_{H^{-1}(\Omega)}^2 \right\} \quad (2.21)$$

where $C_1 > 0$ is solution independent. Since P and g are Lipschitz continuous on $L^2(\Gamma)$, $g(0) = 0$ and D^* is linear continuous from $L^2(\Omega)$ into $L^2(\Gamma)$, plugging (2.19) into (2.21) yields

$$\|u\|_{H^{1/2}(\Omega)}^2 \leq C_1 C_2 \|v\|_{L^2(\Omega)}^2 + 2C_1 \lambda^2 \|v\|_{H^{-1}(\Omega)}^2 + 2C_1 \|f_2\|_{H^{-1}(\Omega)}^2, \quad (2.22)$$

where $C_2 > 0$ is some other constant.

Combining (2.19) and (2.22), we see that $(\mathcal{A} + \text{id})^{-1}$ maps bounded sets of $\mathcal{H} = L^2(\Omega) \times H^{-1}(\Omega)$ into bounded sets of $H^{1/2}(\Omega) \times L^2(\Omega)$, the latter being compactly embedded into \mathcal{H} . Thus, the result is proved. \square

The next proposition is meant for use in Section 3, where we work under additional assumptions on Ω ; however, since it is a direct continuation of the proof of Proposition 2.1, we introduce it here.

Proposition 2.2 (Regularity). *Suppose that $\overline{\Gamma_0} \cap \overline{\Gamma_1} = \emptyset$. Then, the following explicit characterization of $\mathcal{D}(\mathcal{A})$ holds:*

$$\mathcal{D}(\mathcal{A}) = \{[u, v] \in \mathcal{H} : v \in L^2(\Omega), u \in H^1(\Omega), u|_\Gamma = -\mathbb{1}_{\Gamma_0} g(D^*v)\}. \quad (2.23)$$

Thus, strong solutions $[u, u']$ take values in $H^1(\Omega) \times L^2(\Omega)$. Furthermore, there exists a constant $K > 0$ such that any strong solution to (1.1)-(1.2) satisfies

$$\|[u(t), u'(t)]\|_{H^1(\Omega) \times L^2(\Omega)} \leq K \|\mathcal{A}[u(0), u'(0)]\|_{\mathcal{H}} \quad \text{for all } t \geq 0. \quad (2.24)$$

Remark 2.2. In particular, (2.24) implies that for some constant $K' > 0$,

$$\|[u_0, v_0]\|_{\mathcal{H}} \leq K' \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}} \quad \text{for all } [u_0, v_0] \in \mathcal{D}(\mathcal{A}). \quad (2.25)$$

Proof of Proposition 2.2. Let us prove the left inclusion in (2.23). Recalling calculations made in Proposition 2.1, we already know that elements $[u, v]$ in $\mathcal{D}(\mathcal{A})$ satisfy $-\Delta u \in H^{-1}(\Omega)$ and $u|_\Gamma = -Pg(D^*v)$. Therefore, in comparison with the proof of Proposition 2.1, it suffices to show that $u|_\Gamma$ belongs to $H^{1/2}(\Gamma)$ instead of $L^2(\Gamma)$ and apply the elliptic regularity theorem to gain the desired extra half-unit of regularity. By virtue of (1.12), we have $D^*v \in H^{1/2}(\Gamma)$.

First, recall that pointwise Lipschitz nonlinearities such as g map bounded sets of $H^{1/2}(\Gamma)$ into bounded sets of $H^{1/2}(\Gamma)$. Indeed, using the definition of Sobolev spaces on manifold by local charts and the Sobolev-Slobodeckij characterization of the fractional spaces $H^s(\mathbb{R}^{d-1})$ (see [6]), we know that for a given f in $H^{1/2}(\Gamma)$,

$$g(f) \in H^{1/2}(\Gamma) \text{ if and only if } \iint_{\mathbb{R}^{d-1} \times \mathbb{R}^{d-1}} \frac{|\phi_i(x_1)g([f \circ \psi_i](x_1)) - \phi_i(x_2)g([f \circ \psi_i](x_2))|^2}{\|x_1 - x_2\|^d} dx_1 dx_2 < +\infty, \quad (2.26)$$

for all suitable (ϕ_i, ψ_i) , where the functions $\phi_i \in C_c^\infty(\mathbb{R}^{d-1})$ are chosen from a partition of unity subordinate to some (finite) covering of Γ and the functions ψ_i are corresponding local representations of the surface.

The integral term in (2.26) is finite because g and ϕ_i are globally Lipschitz continuous; hence, $g(f) \in H^{1/2}(\Omega)$. Furthermore, taking the square root of $\|\cdot\|_{L^2(\mathbb{R}^{d-1})}^2$ plus the integral term in (2.26) defines a norm on $H^{1/2}(\mathbb{R}^{d-1})$ equivalent to the one given by interpolation. Thus, after coming back to functions on Γ , it follows from (2.26) that

$$\|g(f)\|_{H^{1/2}(\Gamma)} \leq K \|f\|_{H^{1/2}(\Gamma)} \quad \text{for all } f \in H^{1/2}(\Gamma), \quad (2.27)$$

where K is some positive constant coming from the Lipschitz continuity of g and norm equivalence.

Next we have to check that $P \in \mathcal{L}(H^{1/2}(\Gamma))$. Again, this is a consequence of (2.26): we observe that since $\overline{\Gamma_0} \cap \overline{\Gamma_1} = \emptyset$, there exists $m > 0$ such that $\|x_1 - x_2\| > m$ whenever $(\psi_i(x_1), \psi_i(x_2)) \in [\Gamma_0 \times \Gamma_1] \cup [\Gamma_1 \times \Gamma_0]$.

Finally, combining (1.12) for $s = 1/2$, the estimate (2.27), the fact that $P \in \mathcal{L}(H^{1/2}(\Gamma))$ together with the elliptic regularity theorem, we obtain the stronger estimate

$$\begin{aligned} \|u\|_{H^1(\Omega)} &\leq C \{ \|\Delta u\|_{H^{-1}(\Omega)} + \|v\|_{L^2(\Omega)} \} \\ &= C \{ \|A[u + DPg(D^*v)]\|_{H^{-1}(\Omega)} + \|v\|_{L^2(\Omega)} \} \leq C' \|A[u, v]\|_{\mathcal{H}} \end{aligned} \quad (2.28)$$

where C and C' are some positive constants that do not depend on $[u, v]$. The property (2.24) readily follows from (2.28) and (2.7). \square

2.3. Asymptotic stability

Next, we state the second main result of the section, which asserts that the zero equilibrium of the closed-loop system (1.1)-(1.2) is globally asymptotically stable.

Theorem 2.2 (Asymptotic stability of the closed-loop system). *Let $[u_0, v_0] \in \mathcal{H}$. Then,*

$$\|\mathcal{S}_t[u_0, v_0]\|_{\mathcal{H}} \rightarrow 0 \quad \text{as } t \rightarrow +\infty. \quad (2.29)$$

Together with the contraction property of $\{\mathcal{S}_t\}$, (2.29) implies that 0 is a globally asymptotically stable equilibrium point for the feedback system (1.1)-(1.2).

Proof. By the contraction property of the semigroup $\{\mathcal{S}_t\}$ and denseness of $\mathcal{D}(\mathcal{A})$ in \mathcal{H} , it suffices to prove (2.29) for initial data $[u_0, v_0]$ in $\mathcal{D}(\mathcal{A})$.

To do so, we use a Lasalle-type invariance approach. Let us recall the classical line of arguments. We consider the ω -limit set $\omega([u_0, v_0])$ of $[u_0, v_0]$, which can be characterized as follows: $[w_0, z_0] \in \mathcal{H}$ belongs to $\omega([u_0, v_0])$ if there exists an increasing sequence $\{t_n\} \in \mathbb{R}^{\mathbb{N}}$ such that $t_n \rightarrow +\infty$ and

$$\mathcal{S}_{t_n}[u_0, v_0] \rightarrow [w_0, z_0] \quad \text{in } \mathcal{H} \text{ as } n \rightarrow +\infty. \quad (2.30)$$

Recall that $\{\mathcal{S}_t[u_0, v_0]\}_{t \geq 0}$ is relatively compact in \mathcal{H} . Therefore, $\omega([u_0, v_0])$ is a nonempty (positively) invariant compact set, and $\text{dist}(\mathcal{S}_t[u_0, v_0], \omega([u_0, v_0])) \rightarrow 0$ as $t \rightarrow +\infty$ – see [7, Théorème 1.1.8]. Moreover, since $t \mapsto \|A[\mathcal{S}_t[u_0, v_0]]\|_{\mathcal{H}}$ is bounded, it follows from [3, Lemma 2.3] and (2.30) that $\omega([u_0, v_0]) \subset \mathcal{D}(\mathcal{A})$. Besides, since $\mathcal{E}(\mathcal{S}_t[u_0, v_0])$ is bounded and nonincreasing with respect to t , it must converge to some $\mathcal{E}_\infty \geq 0$ as t goes to $+\infty$. By (2.30) and continuity of \mathcal{E} , we have $\mathcal{E}(w_0, z_0) = \mathcal{E}_\infty$ for any $[w_0, z_0] \in \omega([u_0, v_0])$.

The remainder consists in proving that $\omega([u_0, v_0])$ is reduced to $\{0\}$. Let $[w_0, z_0] \in \omega([u_0, v_0])$; we write $[w(t), w'(t)] = \mathcal{S}_t[w_0, z_0]$ and we notice that $\mathcal{E}(w(t), w'(t)) = \mathcal{E}_\infty$ for all $t \geq 0$. Furthermore, $[w, w']$ is a strong solution to (1.1)-(1.2) and we infer from the energy identity (2.8) that

$$\int_0^\tau \int_{\Gamma_0} g(-\partial_\nu[A^{-1}w']) \partial_\nu[A^{-1}w'] \, d\sigma \, dt = 0 \quad \text{for all } \tau \geq 0. \quad (2.31)$$

Letting $p \triangleq A^{-1}w' \in \mathcal{C}(\mathbb{R}^+, H_0^1(\Omega)) \cap L^\infty(0, +\infty; \mathcal{D}(A))$, (2.31) together with the properties of g leads to

$$\partial_\nu p(\sigma, t) = 0 \quad \text{for a.e. } (\sigma, t) \in \Gamma_0 \times (0, +\infty). \quad (2.32)$$

Next, we recall that $w' \in W^{1,\infty}(0, +\infty; H^{-1}(\Omega))$ and, using (2.32) together with the operator-theoretic formulation of (1.1)-(1.2), we obtain $w'' + Aw = 0$. Hence, $p \in W^{1,\infty}(0, +\infty; H_0^1(\Omega))$ satisfies $p' + Ap = 0$, which in turn implies that $p \in W^{2,\infty}(0, +\infty; H^{-1}(\Omega))$ and solves $p'' + Ap = 0$ in $H^{-1}(\Omega)$, i.e., the standard variational formulation of the wave equation with homogeneous Dirichlet boundary condition. In particular, $p \in \mathcal{C}^1(\mathbb{R}^+, L^2(\Omega))$ and solves the following boundary value problem:

$$\partial_{tt}p - \Delta p = 0 \quad \text{in } \Omega \times (0, +\infty), \quad (2.33a)$$

$$p|_\Gamma = 0 \quad \text{on } \Gamma \times (0, +\infty), \quad (2.33b)$$

$$\partial_\nu p = 0 \quad \text{on } \Gamma_0 \times (0, +\infty). \quad (2.33c)$$

The subset Γ_0 being relatively open in Γ , a unique continuation argument for waves yields $p = 0$ – for instance, one can directly apply [15, Théorème 2]. Therefore, $w' = 0$, $Aw = 0$ and finally $w = 0$, which concludes the proof. \square

3. POLYNOMIAL DECAY RATES FOR STRONG SOLUTIONS

This section is dedicated to the analysis of the decay rate of strong solutions under additional assumptions on the feedback nonlinearity and the geometry of the problem.

3.1. Statement of the result and outline of the proof

In what follows, we work under stronger assumptions that are given next.

Assumption 2. There exist positive constants S , α_1 and α_2 such that

$$\alpha_1|s| \leq |g(s)| \leq \alpha_2|s| \quad \text{for all } |s| \leq S. \quad (3.1)$$

Assumption 3. The domain $\Omega \subset \mathbb{R}^d$ with smooth boundary $\Gamma = \Gamma_0 \cup \Gamma_1$ satisfies the following conditions:

(1) The boundary is such that

$$\overline{\Gamma_0} \cap \overline{\Gamma_1} = \emptyset; \quad (3.2)$$

(2) There exists a point $x_0 \in \mathbb{R}^d$ such that, setting $h(x) \triangleq x - x_0$,

$$h \cdot \nu \leq 0 \quad \text{on } \Gamma_1. \quad (3.3)$$

Then, we can estimate the decay rate of each strong solution.

Theorem 3.1 (Non-uniform polynomial decay rate). *Let $r \geq \max\{d-1, 2\}$. Under Assumptions 2 and 3, strong solutions $[u, u']$ to (1.1)-(1.2) satisfy*

$$\mathcal{E}(u(t), u'(t)) \leq C_u t^{-\frac{2}{r-1}} \quad \text{for all } t \geq 0. \quad (3.4)$$

where C_u is a positive constant depending only on $\|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}$.

Theorem 3.1 is a Dirichlet counterpart to non-uniform polynomial decay results that are well-known in the Neumann boundary conditions – see, e.g., [8, Theorem 9.10]. Let us introduce the following notation: if $[u, u']$ is a given solution to (1.1)-(1.2), we define a (continuous) function \mathcal{E}_u over \mathbb{R}^+ by

$$\mathcal{E}_u(t) \triangleq \mathcal{E}(u(t), u'(t)). \quad (3.5)$$

Here, polynomial decay rate is obtained by applying the following classical lemma to the (nonincreasing) energy \mathcal{E}_u of each solution – see [8, Theorem 9.1] for a proof.

Lemma 3.1. *Let $E : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a nonincreasing function. Assume that there exist two positive constants γ and T such that*

$$\int_{\tau}^{+\infty} E^{\gamma+1}(t) dt \leq TE(0)^\gamma E(\tau) \quad \text{for all } \tau \geq 0. \quad (3.6)$$

Then,

$$E(t) \leq E(0) \left(\frac{T + \gamma t}{T + \gamma T} \right)^{-1/\gamma} \quad \text{for all } t \geq T. \quad (3.7)$$

We already know from Section 2 that $\mathcal{E}_u(t)$ converges to 0 as t goes to $+\infty$. Our subsequent efforts focus on estimating

$$\int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2}(t) dt \quad \text{for arbitrary } 0 \leq \tau_1 \leq \tau_2, \quad (3.8)$$

where we recall that

$$\mathcal{E}_u(t) = \frac{1}{2} \{ \|u\|_{L^2(\Omega)}^2 + \|u'\|_{H^{-1}(\Omega)}^2 \}. \quad (3.9)$$

As mentioned in the introduction, the proof is based on an analysis of the variable p defined by

$$p = A^{-1}u' \quad (3.10)$$

which solves, at least formally, the following boundary-value problem:

$$\partial_{tt}p - \Delta p = -\partial_t[DPg(-\partial_\nu p)] \quad \text{in } \Omega \times (0, +\infty), \quad (3.11a)$$

$$p|_\Gamma = 0 \quad \text{on } \Gamma \times (0, +\infty). \quad (3.11b)$$

If u' takes values in $L^2(\Omega)$, recalling the formula (2.1), we have

$$-\partial_\nu p = -\partial_\nu[A^{-1}u'] = D^*Ap = D^*u'. \quad (3.12)$$

To alleviate notation, in the sequel we denote by Φ the term

$$\Phi(t) \triangleq DPg(D^*u'(t)). \quad (3.13)$$

In regards to (2.2), we see that $-\Phi$ is the harmonic extension of the trace $u|_\Gamma$. As mentioned earlier, the p -variable is smoother, which permits, in regards to the wave-type equation (3.11a) satisfied by p , the use of a differential multiplier technique to obtain estimates of the integral over time of

$$\frac{1}{2} \int_{\Omega} \|\nabla p\|^2 + |p'|^2 dx \quad (3.14)$$

premultiplied by an appropriate power of \mathcal{E}_u . The quantity (3.14) is the natural energy of $[p, p']$ at the $H^1(\Omega) \times L^2(\Omega)$ -level (i.e., the standard variational framework); from there, we will be able to deduce a suitable integral estimate of the energy \mathcal{E}_u associated with the less regular u -variable.

Remark 3.1. In fact, since we want to avoid differentiating terms involving g so that our results remain valid when the nonlinearity is only continuous, we will rather multiply an integrated version of (3.11a), namely the formula $u = -[p + \Phi]$, by the time derivative of the multiplier – see Lemma 3.2 below. In particular, p' need not be continuous.

3.2. The multiplier identity

In this subsection, we give more precise properties of the p -variable and derive an expression of (3.8) in the form of a identity obtained by applying an appropriate multiplier to (3.11a). We recall that $\mathcal{D}(A)$ is $H^2(\Omega) \cap H_0^1(\Omega)$ equipped with the norm $\|A \cdot\|_{L^2(\Omega)}$, which is equivalent to the one induced by $H^2(\Omega)$.

Lemma 3.2. *Let $[u, u']$ be a strong solution. The corresponding functions p and Φ enjoy the regularity*

$$p \in L^\infty(0, +\infty; \mathcal{D}(A)) \cap W^{1,\infty}(0, +\infty; H_0^1(\Omega)), \quad \Phi \in L^\infty(0, +\infty; H^1(\Omega)), \quad (3.15)$$

Also, the following identity holds:

$$u = -[p' + \Phi] \in \mathcal{C}(\mathbb{R}, L^2(\Omega)). \quad (3.16)$$

Proof. We infer from Proposition 2.2 that $[u, u'] \in L^\infty(0, +\infty; H^1(\Omega) \times L^2(\Omega))$. As a consequence, since A^{-1} is continuous from $L^2(\Omega)$ into $\mathcal{D}(A)$, we get

$$p = A^{-1}u' \in L^\infty(0, +\infty; \mathcal{D}(A)). \quad (3.17)$$

Besides, $u' : \mathbb{R}^+ \rightarrow H^{-1}(\Omega)$ is absolutely continuous and

$$u'' + Au = -ADPg(D^*u') \quad \text{in } H^{-1}(\Omega) \text{ a.e.} \quad (3.18)$$

Thus, applying A^{-1} to(3.18) yields

$$p' = A^{-1}u'' \in L^\infty(0, +\infty; H_0^1(\Omega)), \quad (3.19)$$

and also

$$u = -[p' + \Phi], \quad \text{and } \Phi \in L^\infty(0, +\infty; H^1(\Omega)), \quad (3.20)$$

which concludes the proof. \square

Define the usual wave multiplier as follows:

$$\mathcal{M}p \triangleq 2h \cdot \nabla p + (d-1)p, \quad (3.21)$$

where $h(x) = x - x_0$ as defined in Assumption 3 and d is the space dimension. Since p satisfies a wave equation, we know that $\int_{\tau_1}^{\tau_2} \int_{\Omega} \|\nabla p\|^2 + |p'| dx dt$ can be estimated by multiplying (3.11a) by $\mathcal{M}p$ and integrating over $\Omega \times (\tau_1, \tau_2)$. Since we are looking for estimates of \mathcal{E}_u at the power $(r+1)/2$, we premultiply $\mathcal{M}p$ by \mathcal{E}_u at the power $(r-1)/2$. Thus, we shall multiply (3.11a) by

$$\mathcal{E}_u^{(r-1)/2}(t)\mathcal{M}p(x, t). \quad (3.22)$$

The resulting identity is given in the next lemma.

Lemma 3.3 (Multiplier identity). *The following equality holds for any $0 \leq \tau_1 \leq \tau_2$:*

$$\begin{aligned} 2 \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} dt &= \mathcal{E}_u^{(r-1)/2} \int_{\Omega} u \mathcal{M}p dx \Big|_{\tau_1}^{\tau_2} + \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Gamma} (h \cdot \nu) |\partial_\nu p|^2 d\sigma - \int_{\Gamma_0} (h \cdot \nu) |g(D^*u')|^2 d\sigma dt \\ &\quad - \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} (d+1)\Phi u + \Phi[2h \cdot \nabla u] dx dt - \frac{(r-1)}{2} \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u \mathcal{M}p dx dt. \end{aligned} \quad (3.23)$$

Proof. The proof is split into four steps.

Step 1: Integration by parts with respect to time. First, by linearity and continuity of \mathcal{M} , $\mathcal{M}p$ belongs to $W^{1,\infty}(0, +\infty; L^2(\Omega))$. On the other hand, recall that \mathcal{E}_u is bounded and absolutely continuous with

$$\mathcal{E}'_u = - \int_{\Gamma_0} g(D^*u')D^*u' d\sigma \quad \text{a.e.}, \quad \mathcal{E}'_u \in L^\infty(0, +\infty), \quad (3.24)$$

because strong solutions are Lipschitz continuous with respect to time. Thus, $\mathcal{E}_u^{(r-1)/2}$ belongs to $W^{1,\infty}(0, +\infty)$ and $\mathcal{E}_u^{(r-1)/2}\mathcal{M}p$ belongs to $W^{1,\infty}(0, +\infty; L^2(\Omega))$ with

$$[\mathcal{E}_u^{(r-1)/2}\mathcal{M}p]' = \mathcal{E}_u^{(r-1)/2}\mathcal{M}p' + \frac{(r-1)}{2}\mathcal{E}'_u\mathcal{E}_u^{(r-3)/2}\mathcal{M}p \quad \text{a.e. in } L^2(\Omega). \quad (3.25)$$

Now, it follows from (3.16) that $p' + \Phi$ belongs to $W^{1,\infty}(0, +\infty; L^2(\Omega))$ and

$$- [p' + \Phi]' = u' = Ap. \quad (3.26)$$

Let $0 \leq \tau_1 \leq \tau_2$. Recall that, since $p \in \mathcal{D}(A)$, $Ap = -\Delta p \in L^2(\Omega)$. Thus, taking the scalar product of (3.26) with $\mathcal{E}_u^{(r-1)/2}\mathcal{M}p$ in $L^2(\tau_1, \tau_2; L^2(\Omega))$ and using the integration by parts formula in $W^{1,2}(\tau_1, \tau_2; L^2(\Omega))$ leads to

$$\begin{aligned} - \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \Delta p \mathcal{M}p \, dx \, dt &= \mathcal{E}_u^{(r-1)/2} \int_{\Omega} u \mathcal{M}p \, dx \Big|_{\tau_1}^{\tau_2} - \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} u \mathcal{M}p' \, dx \, dt \\ &\quad - \frac{(r-1)}{2} \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u \mathcal{M}p \, dx \, dt. \end{aligned} \quad (3.27)$$

Step 2: Multiplier technique for the wave equation. In what follows, we apply classical vector calculus identities to recover the $H_0^1(\Omega) \times L^2(\Omega)$ -energy of $[p, p']$. Since p takes values in $H^2(\Omega)$, Rellich's identity yields

$$\int_{\Omega} \Delta p [2h \cdot \nabla p] \, dx = (d-2) \int_{\Omega} \|\nabla p\|^2 \, dx + \int_{\Gamma} \partial_\nu p [2h \cdot \nabla p] \, d\sigma - \int_{\Gamma} (h \cdot \nu) \|\nabla p\|^2 \, d\sigma. \quad (3.28)$$

Furthermore, $p|_{\Gamma} = 0$; thus

$$\nabla p = (\partial_\nu p)\nu \quad \text{on } \Gamma. \quad (3.29)$$

Combining (3.28) and (3.29), we obtain

$$\int_{\Omega} \Delta p [2h \cdot \nabla p] = (d-2) \int_{\Omega} \|\nabla p\|^2 \, dx + \int_{\Gamma} (h \cdot \nu) |\partial_\nu p|^2 \, d\sigma. \quad (3.30)$$

On the other hand,

$$\int_{\Omega} \Delta p (d-1)p \, dx = -(d-1) \int_{\Omega} \|\nabla p\|^2 \, dx, \quad (3.31)$$

where we use again that p vanishes on the boundary. Summing (3.30) and (3.31) yields

$$\int_{\Omega} \Delta p \mathcal{M}p \, dx = - \int_{\Omega} \|\nabla p\|^2 \, dx + \int_{\Gamma} (h \cdot \nu) |\partial_\nu p|^2 \, d\sigma. \quad (3.32)$$

Coming back to (3.27) and recalling (3.16), let us write

$$\begin{aligned}
\int_{\Omega} u\mathcal{M}p' &= - \int_{\Omega} p' \mathcal{M}p' \, dx - \int_{\Omega} \Phi \mathcal{M}p' \, dx \\
&= - \int_{\Omega} p' [2h \cdot \nabla p'] + (d-1)|p'|^2 \, dx - \int_{\Omega} \Phi \mathcal{M}p' \, dx \\
&= \int_{\Omega} |p'|^2 \, dx - \int_{\Omega} \Phi \mathcal{M}p' \, dx,
\end{aligned} \tag{3.33}$$

where we use the identity

$$\int_{\Omega} \phi [2h \cdot \nabla \phi] \, dx = \int_{\Gamma} (h \cdot \nu) (\phi|_{\Gamma})^2 \, d\sigma - \int_{\Omega} (\phi)^2 \operatorname{div} h \, dx \quad \text{for all } \phi \in H^1(\Omega) \tag{3.34}$$

together with $\operatorname{div} h = d$ and $p|_{\Gamma} = 0$. Therefore, combining (3.27) with (3.32) and (3.33) leads to

$$\begin{aligned}
\int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \|\nabla p\|^2 + |p'|^2 \, dx \, dt &= \mathcal{E}_u^{(r-1)/2} \int_{\Omega} u\mathcal{M}p \, dx \Big|_{\tau_1}^{\tau_2} + \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \Phi \mathcal{M}p' \, dx \, dt \\
&+ \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Gamma} (h \cdot \nu) |\partial_{\nu} p|^2 \, d\sigma \, dt - \frac{(r-1)}{2} \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u\mathcal{M}p \, dx \, dt.
\end{aligned} \tag{3.35}$$

Step 3: Additional terms. Here, we put the terms involving Φ into a form suitable for further estimation. It follows from (3.16) that

$$\int_{\Omega} \Phi \mathcal{M}p' \, dx = - \int_{\Omega} \Phi \mathcal{M}u \, dx - \int_{\Omega} \Phi \mathcal{M}\Phi \, dx. \tag{3.36}$$

Applying (3.34) to Φ , similarly to (3.33), we obtain

$$\begin{aligned}
- \int_{\Omega} \Phi \mathcal{M}\Phi \, dx &= \int_{\Omega} |\Phi|^2 \, dx - \int_{\Gamma} (h \cdot \nu) (\Phi|_{\Gamma})^2 \\
&= \int_{\Omega} |\Phi|^2 \, dx - \int_{\Gamma_0} (h \cdot \nu) |g(D^* u')|^2 \, d\sigma,
\end{aligned} \tag{3.37}$$

where we use that, by definition,

$$\Phi|_{\Gamma} = g(D^* A p) = g(D^* u') \text{ on } \Gamma_0, \quad \Phi|_{\Gamma} = 0 \text{ on } \Gamma_1. \tag{3.38}$$

On the other hand, we recall that

$$- \int_{\Omega} \Phi \mathcal{M}u \, dx = - \int_{\Omega} \Phi [2h \cdot \nabla u] + (d-1)\Phi u \, dx. \tag{3.39}$$

Plugging (3.37) and (3.39) into (3.35) leads to

$$\begin{aligned}
\int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \|\nabla p\|^2 + |p'|^2 \, dx \, dt &= \mathcal{E}_u^{(r-1)/2} \int_{\Omega} u\mathcal{M}p \, dx \Big|_{\tau_1}^{\tau_2} - \frac{(r-1)}{2} \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u\mathcal{M}p \, dx \, dt \\
&+ \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Gamma} (h \cdot \nu) |\partial_{\nu} p|^2 \, d\sigma - \int_{\Gamma_0} (h \cdot \nu) |g(D^* A p)|^2 \, d\sigma - \int_{\Omega} \Phi [2h \cdot \nabla u] + (d-1)\Phi u - |\Phi|^2 \, dx \, dt.
\end{aligned} \tag{3.40}$$

Step 4: Conclusion. We finish the proof by rewriting (3.40) as an estimate of $\int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} dt$. This is done as follows. First, by definition of the p -variable,

$$\int_{\Omega} \|\nabla p\|^2 dx = \|A^{1/2}p\|_{L^2(\Omega)}^2 = \|A^{-1/2}u'\|_{L^2(\Omega)}^2 = \|u'\|_{H^{-1}(\Omega)}^2. \quad (3.41)$$

On the other hand, it immediately follows from $u = -[p' + \Phi]$ that

$$\int_{\Omega} |p'|^2 dx = \int_{\Omega} |u|^2 dx + \int_{\Omega} 2\Phi u + |\Phi|^2 dx \quad \text{a.e.} \quad (3.42)$$

Summing (3.41) and (3.42) yields

$$\int_{\Omega} \|\nabla p\|^2 + |p'|^2 dx = 2\mathcal{E}_u + \int_{\Omega} 2\Phi u + |\Phi|^2 dx \quad \text{a.e.} \quad (3.43)$$

Plugging (3.43) into (3.40), we get the desired identity. \square

3.3. Estimates of the right-hand side and conclusion

Our goal in this subsection is to establish an integral inequality in the form of

$$(1 - \mu) \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} dt \leq K_u \{\mathcal{E}_u(\tau_1) + \mathcal{E}_u(\tau_2)\} \quad \text{for all } 0 \leq \tau_1 \leq \tau_2, \quad (3.44)$$

where K_u is a constant that may depend on the initial data and μ is a sufficiently small constant that may depend on u as well. Assuming that (3.44) holds, we let τ_2 go to $+\infty$ to obtain

$$\int_{\tau}^{+\infty} \mathcal{E}_u^{(r+1)/2} dt \leq \frac{K_u}{1 - \mu} \mathcal{E}_u(\tau) \quad \text{for all } \tau \geq 0. \quad (3.45)$$

Then, Theorem 3.1 follows readily from Lemma 3.1 if we choose

$$\gamma = \frac{(r-1)}{2} \quad \text{and} \quad T = \frac{K_u}{1 - \mu} \mathcal{E}_u(0)^{-(r-1)/2}. \quad (3.46)$$

To prove (3.44), we shall examine each term in the multiplier identity (3.23) and derive estimates in terms of

- Either directly $\mathcal{E}_u(\tau_1)$ or $\mathcal{E}_u(\tau_2)$;
- Or the boundary dissipation term $\int_{\tau_1}^{\tau_2} \int_{\Gamma_0} g(D^*u') D^*u' d\sigma$ which is nonnegative and can be integrated, since

$$-\mathcal{E}'_u = \int_{\Gamma_0} g(D^*u') D^*u' d\sigma = - \int_{\Gamma_0} g(-\partial_\nu p) \partial_\nu p d\sigma; \quad (3.47)$$

- And also $\int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} dt$ premultiplied by small μ so that it can be absorbed in the left-hand side.

Remark 3.2. In what follows, we shall denote by K , K' , etc. generic constants that *do not* depend on the initial data.

In practice, we can write estimates in terms of $\int_{\tau_1}^{\tau_2} \int_{\Gamma_0} |g(D^*u')|^2 d\sigma dt$ instead of $\int_{\tau_1}^{\tau_2} \int_{\Gamma_0} g(D^*u') D^*u' d\sigma dt$ since

$$0 \leq \int_{\Gamma_0} |g(D^*u')|^2 d\sigma \leq K \int_{\Gamma_0} |g(D^*u') D^*u'| d\sigma = K \int_{\Gamma_0} g(D^*u') D^*u' d\sigma \quad (3.48)$$

by Lipschitz continuity and nonincreasingness of g , together with $g(0) = 0$.

Moreover, recalling (3.3) in Assumption 3 and looking at the sign of each term, we observe that $|\partial_\nu p|$ in (3.23) need not be estimated on the uncontrolled boundary Γ_1 .

That being said, let us start by estimating the term involving $2h \cdot \nabla u$ in (3.23). This is done in the following lemma.

Lemma 3.4. *Suppose that $r \geq 2$. Then, there exists a positive constant K such that*

$$\left| \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \Phi[2h \cdot \nabla u] \, dx \, dt \right| \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{1/2} \left\{ \int_{\tau_1}^{\tau_2} \frac{1}{\mu} \int_{\Gamma_0} |g(D^* u')|^2 \, d\sigma + \mu \mathcal{E}_u^{(r-2)/2}(0) \mathcal{E}_u^{(r+1)/2} \, dt \right\} \quad (3.49)$$

for all $\tau_2 \geq \tau_1 \geq 0$ and $\mu > 0$.

Proof. We start by writing

$$(\Phi, 2h \cdot \nabla u)_{L^2(\Omega)} = (DPg(D^* u'), 2h \cdot \nabla u)_{L^2(\Omega)} = (g(D^* u'), D^*[2h \cdot \nabla u])_{L^2(\Gamma_0)}. \quad (3.50)$$

Thus, applying the Cauchy-Schwarz inequality yields

$$\left| \int_{\Omega} \Phi[2h \cdot \nabla u] \, dx \right| \leq \|g(D^* u')\|_{L^2(\Gamma_0)} \|D^*[2h \cdot \nabla u]\|_{L^2(\Gamma_0)} \leq \|g(D^* u')\|_{L^2(\Gamma_0)} \|D^*[2h \cdot \nabla u]\|_{L^2(\Gamma)}. \quad (3.51)$$

Next, recall from (1.12) that

$$D^* \in \mathcal{L}(H^{-1/2}(\Omega), L^2(\Gamma)). \quad (3.52)$$

Therefore, it follows from (3.51) that

$$\left| \int_{\Omega} \Phi[2h \cdot \nabla u] \, dx \right| \leq K \|g(D^* Ap)\|_{L^2(\Gamma_0)} \|2h \cdot \nabla u\|_{H^{-1/2}(\Omega)}. \quad (3.53)$$

Linear interpolation between the Sobolev spaces $L^2(\Omega)$ and $H^{-1}(\Omega)$ leads to

$$\|2h \cdot \nabla u\|_{H^{-1/2}(\Omega)} \leq K \|2h \cdot \nabla u\|_{L^2(\Omega)}^{1/2} \|2h \cdot \nabla u\|_{H^{-1}(\Omega)}^{1/2} \quad (3.54)$$

First, by Proposition 2.2,

$$\|2h \cdot \nabla u\|_{L^2(\Omega)}^{1/2} \leq K \|h\|_{L^\infty(\Omega)^d}^{1/2} \|\nabla u\|_{L^2(\Omega)^d}^{1/2} \leq K' \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{1/2}. \quad (3.55)$$

Besides, since $2h \cdot \nabla u$ belongs to $L^2(\Omega)$, we have

$$\langle 2h \cdot \nabla u, w \rangle_{H^{-1}(\Omega), H_0^1(\Omega)} = (2h \cdot \nabla u, w)_{L^2(\Omega)} \quad \text{for all } w \in H_0^1(\Omega). \quad (3.56)$$

Let us write

$$\begin{aligned} \int_{\Omega} [2h \cdot \nabla u] w \, dx &= 2 \sum_{i=1}^d \int_{\Omega} h_i \partial_{x_i} u w \, dx \\ &= -2 \sum_{i=1}^d \int_{\Omega} u [w \partial_i h_i + h_i \partial_i w] \, dx + \sum_{i=1}^d \int_{\Gamma} \nu_i h_i u w \, d\sigma \\ &= -2 \int_{\Omega} u [w \operatorname{div} h + h \cdot \nabla w] \, dx, \end{aligned} \quad (3.57)$$

where h_i (resp. ν_i) denotes the i -th coordinate of the vector field h (resp. the outward normal vector ν). Recall that $h \in \mathcal{C}^1(\overline{\Omega})$. Then, using the Poincaré inequality on w , we obtain that for some $K > 0$,

$$|(2h \cdot \nabla u, w)_{L^2(\Omega)}| \leq K \|u\|_{L^2(\Omega)} \|w\|_{H_0^1(\Omega)}. \quad (3.58)$$

By norm equivalence between $H^{-1}(\Omega)$ and $H_0^1(\Omega)'$, we infer from (3.58) that

$$\|2h \cdot \nabla u\|_{H^{-1}(\Omega)} \leq K \|u\|_{L^2(\Omega)}. \quad (3.59)$$

Coming back to (3.53), combining (3.55) and (3.59) yields

$$\begin{aligned} \left| \int_{\Omega} Dg(D^*u') [2h \cdot \nabla u] \, dx \right| &\leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{1/2} \|g(D^*u')\|_{L^2(\Gamma_0)} \|u\|_{L^2(\Omega)}^{1/2} \\ &\leq K' \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{1/2} \|g(D^*u')\|_{L^2(\Gamma_0)} \mathcal{E}_u^{1/4}. \end{aligned} \quad (3.60)$$

Therefore, since $\mathcal{E}_u \geq 0$, we have

$$\left| \mathcal{E}_u^{(r-1)/2} \int_{\Omega} Dg(D^*u') 2h \cdot \nabla u \, dx \right| \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{1/2} \|g(D^*u')\|_{L^2(\Gamma_0)} \mathcal{E}_u^{(r-1)/2+1/4}. \quad (3.61)$$

Applying Young's inequality with a parameter $\mu > 0$, we obtain

$$\|g(D^*u')\|_{L^2(\Gamma)} \mathcal{E}_u^{(r-1)/2+1/4} \leq \frac{1}{2\mu} \|g(D^*u')\|_{L^2(\Gamma)}^2 + \frac{\mu}{2} \mathcal{E}_u^{r-1/2}. \quad (3.62)$$

It is assumed that $r \geq 2$. Thus, letting $\eta \triangleq (r-1/2) - (r+1)/2 \geq 0$, by nonincreasingness of \mathcal{E}_u , we have

$$\mathcal{E}_u^{r-1/2} = \mathcal{E}_u^\eta \mathcal{E}_u^{(r+1)/2} \leq \mathcal{E}_u^\eta(0) \mathcal{E}_u^{(r+1)/2}. \quad (3.63)$$

Plugging (3.63) into (3.62) and integrating over (τ_1, τ_2) yields the desired result. \square

Next, we deal with the term involving $|\partial_\nu p| = |D^*u'|$ on the controlled boundary Γ_0 . Here, the arguments are very similar to those employed in the case of saturated Neumann feedback – see [8, Theorem 9.10] or [19, Theorem 3].

Lemma 3.5. *Suppose that $r \geq d-1$. Then, there exists $K > 0$ and $\eta \in (0, 1)$ such that*

$$\begin{aligned} \left| \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Gamma_0} |\partial_\nu p|^2 \, d\sigma \, dt \right| &\leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{2-\eta} \left\{ \mu^{1/\eta} \int_{\tau_1}^{\tau_2} \int_{\Gamma_0} g(D^*u') D^*u' \, d\sigma \, dt + \mu^{-1/(1-\eta)} \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} \, dt \right\} \\ &\quad + K \int_{\tau_1}^{\tau_2} \int_{\Gamma_0} |g(D^*u')|^2 \, d\sigma \, dt \end{aligned} \quad (3.64)$$

for all $0 \leq \tau_1 \leq \tau_2$ and $\mu > 0$.

Proof. For each $t \geq 0$, we set

$$\Gamma_t^1 \triangleq \{\sigma \in \Gamma_0 : |\partial_\nu p(\sigma, t)| \leq S\} \quad \text{and} \quad \Gamma_t^2 \triangleq \Gamma_0 \setminus \Gamma_t^1. \quad (3.65)$$

Then,

$$\left| \int_{\Gamma_0} (h \cdot \nu) |\partial_\nu p|^2 \, d\sigma \right| \leq K \int_{\Gamma_t^0} |\partial_\nu p|^2 \, d\sigma + K \int_{\Gamma_t^1} |\partial_\nu p|^2 \, d\sigma. \quad (3.66)$$

Using (3.1) in Assumption 2, we estimate the first term in (3.66) as follows:

$$\int_{\Gamma_t^0} |\partial_\nu p|^2 \, d\sigma \leq \alpha_1^{-2} \int_{\Gamma_t^0} |g(-\partial_\nu p)|^2 \, d\sigma \leq \alpha_1^{-2} \int_{\Gamma_0} |g(-\partial_\nu p)|^2 \, d\sigma = \alpha_1^{-2} \int_{\Gamma_0} |g(D^*u')|^2 \, d\sigma. \quad (3.67)$$

Let us examine the second term in (3.66). Setting a parameter $\eta \in (0, 1)$ to be tuned later on, we have

$$\int_{\Gamma_t^1} |\partial_\nu p|^2 d\sigma = \int_{\Gamma_t^1} |D^* u'|^2 d\sigma = \int_{\Gamma_t^1} |D^* u'|^{2-\eta} \frac{|g(D^* u') D^* u'|^\eta}{|g(D^* u')|^\eta} d\sigma. \quad (3.68)$$

Equation (3.68) makes sense since $|g(D^* u')| \geq \min\{g(S), -g(-S)\} > 0$ on Γ_t^1 . In fact, we have

$$\begin{aligned} \int_{\Gamma_t^1} |D^* u'|^2 d\sigma &\leq \min\{g(S), -g(-S)\}^{-\eta} \int_{\Gamma_t^1} |D^* u'|^{2-\eta} |g(D^* u') D^* u'|^\eta d\sigma \\ &\leq K \int_{\Gamma_0} |D^* u'|^{2-\eta} |g(D^* u') D^* u'|^\eta d\sigma. \end{aligned} \quad (3.69)$$

Using Hölder's inequality with conjugates $1/\eta$ and $1/(1-\eta)$, we infer from (3.69) that

$$\int_{\Gamma_t^1} |D^* u'|^2 d\sigma \leq K \left(\int_{\Gamma_0} |D^* u'|^{\frac{2-\eta}{1-\eta}} \right)^{1-\eta} \left(\int_{\Gamma_0} g(D^* u') D^* u' d\sigma \right)^\eta. \quad (3.70)$$

Now, $[u, u']$ being a strong solution to (1.1)-(1.2), we recall from Proposition 2.2 that u' takes values in $L^2(\Omega)$ and $\|u'(t)\|_{L^2(\Omega)} \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}$ for all $t \geq 0$. The continuity property (1.12) yields

$$\|D^* u'(t)\|_{H^{1/2}(\Gamma)} \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}} \quad \text{for all } t \geq 0. \quad (3.71)$$

In what follows, we rely on (fractional) Sobolev inequalities – see [6, Theorem 6.5 and Theorem 6.9]. First, we consider the case $d \geq 3$, where we recall that d denotes the space dimension. We have the continuous embedding

$$H^{1/2}(\Gamma_0) \hookrightarrow L^q(\Gamma_0) \quad \text{for all } q \in \left[2, \frac{2(d-1)}{d-2} \right] \triangleq I_d. \quad (3.72)$$

Furthermore, since $r+1 \geq d$, if we choose $\eta = 2/(r+1)$, some computations yield $(2-\eta)/(1-\eta) \in I_d$; hence

$$H^{1/2}(\Gamma_0) \hookrightarrow L^{\frac{2-\eta}{1-\eta}}(\Gamma_0). \quad (3.73)$$

If $d = 2$, then the embedding (3.72) holds in fact for any $q \in [2, +\infty)$; therefore, (3.73) is valid as well. Coming back to (3.70), combining (3.73) with (3.71) yields

$$\mathcal{E}_u^{(r-1)/2} \int_{\Gamma_t^1} |D^* u'|^2 d\sigma \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{2-\eta} \mathcal{E}_u^{(r-1)/2} \left(\int_{\Gamma_0} g(D^* u') D^* u' d\sigma \right)^\eta. \quad (3.74)$$

Applying the Young inequality with conjugates $1/\eta$ and $1/(1-\eta)$, we get

$$\mathcal{E}_u^{(r-1)/2} \int_{\Gamma_t^1} |D^* u'|^2 d\sigma \leq K \|\mathcal{A}[u_0, v_0]\|_{\mathcal{H}}^{2-\eta} \left\{ \mu^{-\frac{1}{1-\eta}} \mathcal{E}_u^{\frac{r-1}{2(1-\eta)}} + \mu^{\frac{1}{\eta}} \int_{\Gamma_0} g(D^* u') D^* u' d\sigma \right\} \quad \text{for all } \mu > 0. \quad (3.75)$$

Since $(r-1)/2(1-\eta) = (r+1)/2$, we conclude the proof by combining (3.75) and (3.67) together with (3.66). \square

At this point, the proof of Theorem 3.1 is almost complete. Estimates of the remaining terms in (3.23) are given in the next lemmas. Following our remarks at the beginning of the subsection, we claim that Theorem 3.1 is proved once those are established.

Lemma 3.6. *There exists a positive constant K such that*

$$\left| \left[\mathcal{E}_u^{(r-1)/2} \int_{\Omega} u \mathcal{M} p \, dx \right]_{\tau_1}^{\tau_2} \right| \leq K \mathcal{E}_u^{(r-1)/2}(0) \{ \mathcal{E}_u(\tau_1) + \mathcal{E}_u(\tau_2) \} \quad \text{for all } 0 \leq \tau_1 \leq \tau_2. \quad (3.76)$$

Proof. Let $\tau \geq 0$. Then,

$$\begin{aligned} \left| \mathcal{E}_u^{(r-1)/2}(\tau) \int_{\Omega} u(\tau) \mathcal{M} p(\tau) \, dx \right| &\leq \mathcal{E}_u^{(r-1)/2}(\tau) \|u(\tau)\|_{L^2(\Omega)} \|\mathcal{M} p(\tau)\|_{L^2(\Omega)} \\ &\leq K \mathcal{E}_u^{(r-1)/2}(\tau) \|u(\tau)\|_{L^2(\Omega)} \|p(\tau)\|_{H_0^1(\Omega)} \\ &\leq K \mathcal{E}_u^{(r-1)/2}(\tau) \|u(\tau)\|_{L^2(\Omega)} \|u'(\tau)\|_{H^{-1}(\Omega)} \\ &\leq K \mathcal{E}_u^{(r-1)/2}(0) \mathcal{E}_u(\tau), \end{aligned} \quad (3.77)$$

where it used that \mathcal{E}_u is nonincreasing. Equation (3.76) readily follows from the triangular inequality. \square

Lemma 3.7. *There exists a positive constant K such that*

$$\left| \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u \mathcal{M} p \, dx \, dt \right| \leq K \mathcal{E}_u^{(r-1)/2}(0) \{ \mathcal{E}_u(\tau_1) + \mathcal{E}_u(\tau_2) \} \quad \text{for all } 0 \leq \tau_1 \leq \tau_2. \quad (3.78)$$

Proof. Again, we write

$$\left| \int_{\Omega} u \mathcal{M} p \, dx \right| \leq K \mathcal{E}_u. \quad (3.79)$$

Therefore,

$$\begin{aligned} \left| \int_{\tau_1}^{\tau_2} \mathcal{E}'_u \mathcal{E}_u^{(r-3)/2} \int_{\Omega} u \mathcal{M} p \, dx \, dt \right| &\leq K \int_{\tau_1}^{\tau_2} (-\mathcal{E}'_u) \mathcal{E}_u^{(r-1)/2} \, dt = -K \int_{\tau_1}^{\tau_2} \left[\frac{2}{r+1} \mathcal{E}_u^{(r+1)/2} \right]' \, dt \\ &= \frac{2K}{r+1} \{ \mathcal{E}_u^{(r+1)/2}(\tau_1) - \mathcal{E}_u^{(r+1)/2}(\tau_2) \}. \end{aligned} \quad (3.80)$$

The desired inequality follows from the nonincreasingness of \mathcal{E}_u and (3.80). \square

Lemma 3.8. *There exists a positive constant K such that*

$$\left| \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \Phi u \, dx \, dt \right| \leq K \left\{ \frac{1}{\mu} \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} \, dt + \mu \mathcal{E}_u^{(r-1)/2}(0) \int_{\tau_1}^{\tau_2} \int_{\Gamma_0} |g(D^* u')|^2 \, d\sigma \, dt \right\} \quad (3.81)$$

for all $0 \leq \tau_1 \leq \tau_2$ and $\mu > 0$.

Proof. First, using Cauchy-Schwarz and Young inequalities, we obtain

$$\begin{aligned} \left| \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \int_{\Omega} \Phi u \, dx \, dt \right| &\leq \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \|\Phi\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)} \, dt \\ &\leq \frac{1}{2\mu} \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r-1)/2} \|\Phi\|_{L^2(\Omega)}^2 \, dt + \mu \int_{\tau_1}^{\tau_2} \mathcal{E}_u^{(r+1)/2} \, dt. \end{aligned} \quad (3.82)$$

Next, recall that $\Phi = DPg(D^* u')$ and that D continuously maps $L^2(\Gamma)$ into $L^2(\Omega)$. Therefore,

$$\|\Phi\|_{L^2(\Omega)}^2 \leq K \|Pg(D^* u')\|_{L^2(\Gamma)}^2 = K \|g(D^* u')\|_{L^2(\Gamma_0)}^2 \quad (3.83)$$

and we conclude the proof by plugging (3.83) into (3.82) and using that \mathcal{E}_u is nonincreasing. \square

4. CONCLUDING REMARKS

In this section, we discuss the results of our paper and give some comments and perspectives.

- Theorem 3.1 deals with the decay rate of strong solutions to (1.1)-(1.2), which remain bounded in a stronger norm (here, in $H^1(\Omega) \times L^2(\Omega)$). In particular, this enables the use of Sobolev embeddings to obtain estimates of the boundary term $\partial_\nu[A^{-1}u']$ in $L^\infty(0, +\infty; L^q(\Gamma))$ for some appropriate q . In view of the energy identity (2.8), and as done in Lemma 3.5, we can then derive an estimate involving only the “dissipation term” $g(-\partial_\nu[A^{-1}u'])$ and lower-order energy terms, even though no lower bound on nonlinearity g is prescribed at infinity. Here, using the terminology of [18], the feedback is allowed to be *weak*, i.e., $g(s)/s$ can go to 0 as $|s|$ goes to infinity, as it is the case when g represents a saturation mapping; then, loss of uniformity is to be expected. More precisely, coming back to the Neumann problem, the one-dimensional version of (1.6) with g given by (1.7) is known to possess weak solutions that decay to zero (in the natural energy space $H^1(\Omega) \times L^2(\Omega)$) slower than any exponential or polynomial, whereas strong solutions decay exponentially to zero but in a non-uniform way – see [18, Theorem 4.1] or also [1, Theorem 59]. Proving a similar result in our Dirichlet case would be interesting.
- Putting aside the matter of saturated feedback and assuming if needed that g has linear growth at infinity, we see that, unfortunately, the strategy followed here is not sufficient to prove *uniform* decay of solutions to (1.1)-(1.2). Indeed, while estimating the term $(\Phi, 2h \cdot \nabla u)_{L^2(\Omega)}$ as in Lemma 3.4 is good enough for the purpose of proving Theorem 3.1, it requires, again, that solutions remain bounded in a norm stronger than that of the energy space \mathcal{H} . If, instead of (3.49), one manages to prove something in the likes of

$$\int_0^\tau \int_\Omega \Phi[2h \cdot \nabla u] dx dt \leq K(\tau) \int_0^\tau |g(D^*u')|^2 d\sigma dt + K'\{\mathcal{E}_u(0) + \mathcal{E}_u(\tau)\} + \epsilon \int_0^\tau \mathcal{E}_u dt \quad (4.1)$$

for some $\tau > 0$, where $K(\tau)$ is allowed to depend on τ and ϵ can be chosen sufficiently small, then, by remarking that the multiplier identity (3.23) is still valid with the time-varying weight $\mathcal{E}(u, u')^{(r-1)/2}$ replaced by the constant 1, one could easily adapt the rest of our proof to obtain exponential uniform stability. By following the proof of [9, Lemma 3.3], we can prove such estimate when g is the identity, at least under some specific geometrical conditions; however, the argument breaks down in the nonlinear case. Therefore, as mentioned in the introduction, the problem of uniform stability is still open.

REFERENCES

- [1] Yacine Chitour, Swann Marx, and Guilherme Mazanti. One-dimensional wave equation with set-valued boundary damping: well-posedness, asymptotic stability, and decay rates. 2020.
- [2] Igor Chueshov, Matthias Eller, and Irena Lasiecka. On the attractor for a semilinear wave equation with critical exponent and nonlinear boundary dissipation. *Communications in Partial Differential Equations*, 27(9-10):1901–1951, January 2002.
- [3] Michael G. Crandall and Amnon Pazy. Semi-groups of nonlinear contractions and dissipative sets. *Journal of functional analysis*, 3(3):376–418, 1969.
- [4] C.M. Dafermos and M. Slemrod. Asymptotic behavior of nonlinear contraction semigroups. *Journal of Functional Analysis*, 13(1):97–106, May 1973.
- [5] Moez Daoulati, Irena Lasiecka, and Daniel Toundykov. Uniform energy decay for a wave equation with partially supported nonlinear boundary dissipation without growth restrictions. *Discrete & Continuous Dynamical Systems-S*, 2(1):67, 2009.
- [6] Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bulletin des sciences mathématiques*, 136(5):521–573, 2012.
- [7] Alain Haraux. *Systemes dynamiques dissipatifs et applications*, volume 17. Masson, 1991.
- [8] Vilmos Komornik. *Exact controllability and stabilization: the multiplier method*, volume 36. Elsevier Masson, 1994.
- [9] I. Lasiecka and R. Triggiani. Uniform Exponential Energy Decay of Wave Equations in a Bounded Region with $L_2(0, +\infty; L_2(\Gamma))$ -Feedback Control in the Dirichlet Boundary Conditions. *Journal of Differential Equations*, 66:34c–390, 1987.
- [10] I. Lasiecka and R. Triggiani. Uniform stabilization of the wave equation with Dirichlet or Neumann feedback control without geometrical conditions. *Applied Mathematics and Optimization*, 25(2):189–224, March 1992.

- [11] Irena Lasiecka, J.-L. Lions, and Roberto Triggiani. Non homogeneous boundary value problems for second order hyperbolic operators. *Journal de Mathématiques pures et Appliquées*, 65(2):149–192, 1986.
- [12] Irena Lasiecka and Daniel Tataru. Uniform boundary stabilization of semilinear wave equations with nonlinear boundary damping. *Differential and integral Equations*, 6(3):507–533, 1993.
- [13] Jacques-Louis Lions and E. Magenes. Problemes aux limites non homogenes. II. In *Annales de l’institut Fourier*, volume 11, pages 137–178, 1961.
- [14] Christophe Prieur, Sophie Tarbouriech, and Joao M. Gomes da Silva Jr. Wave Equation With Cone-Bounded Control Laws. *IEEE Transactions on Automatic Control*, 61(11):3452–3463, November 2016.
- [15] Luc Robbiano. Théorème d’unicité adapté au contrôle des solutions des problèmes hyperboliques. *Communications in partial differential equations*, 16(4-5):789–800, 1991.
- [16] Ralph Edwin Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49. American Mathematical Soc., 2013.
- [17] Roger Temam. *Infinite-dimensional dynamical systems in mechanics and physics*, volume 68. Springer Science & Business Media, 2012.
- [18] Judith Vancostenoble and Patrick Martinez. Optimality of energy estimates for the wave equation with nonlinear boundary velocity feedbacks. *SIAM Journal on Control and Optimization*, 39(3):776–797, 2000.
- [19] Cheng-Zhong Xu and Gen Qi Xu. Saturated Boundary Feedback Stabilization of a Linear Wave Equation. *SIAM Journal on Control and Optimization*, 57(1):290–309, January 2019.
- [20] Enrike Zuazua. Uniform Stabilization of the Wave Equation by Nonlinear Boundary Feedback. *SIAM Journal on Control and Optimization*, 28(2):466–477, March 1990.