



The socio-ecological legacies of centuries-old charcoal making practices in a mountain forest of the Northern Pyrenees

Léonel Fouédjeu, Sandrine Paradis-Grenouillet, Laurent Larrieu, Mélanie Saulnier, Sylvain Burri, Vanessa Py-Saragaglia

► To cite this version:

Léonel Fouédjeu, Sandrine Paradis-Grenouillet, Laurent Larrieu, Mélanie Saulnier, Sylvain Burri, et al.. The socio-ecological legacies of centuries-old charcoal making practices in a mountain forest of the Northern Pyrenees. *Forest Ecology and Management*, 2021, 502, 10.1016/j.foreco.2021.119717 . hal-03365433

HAL Id: hal-03365433

<https://hal.science/hal-03365433>

Submitted on 5 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title: The socio-ecological legacies of centuries-old charcoal making practices in a mountain forest of the Northern Pyrenees

Léonel Fouédjeu^{1*}; Sandrine Paradis-Grenouillet^{2,3}; Laurent Larrieu^{4,5}, Mélanie Saulnier^{1,6}; Sylvain Burri⁷ and Vanessa Py-Saragaglia¹

¹ GEODE, LABEX DRIHM, UMR 5602 CNRS. Université Toulouse Jean Jaurès, 5, Allées A. Machado, 31058 TOULOUSE Cedex 1, France

² Éveha, bureau d'étude en archéologie préventive, 31 rue Soyouz 87000 Limoges

³ Université de Limoges, CNRS, GEOLAB, 39 E rue Camille Guérin, F-87036 LIMOGES

⁴ Université de Toulouse, INRAE, UMR DYNAFOR, Castanet-Tolosan, France

⁵ CNPF-CRPF Occitanie, Tarbes, France

⁶ Česká zemědělská univerzita v Praze, c, Kamýcká 129, 165 00, Praha 6, Suchbát, Czech Republic

⁷ TRACES, UMR 5608 CNRS Université Toulouse Jean Jaurès, 5, Allées A. Machado, 31058 TOULOUSE Cedex 1, France

16 Highlights

- 17 • Past charcoal making practices were reconstructed through analysis of charcoal remains
- 18 • These practices were transmitted over centuries without significant change
- 19 • Management shifted from coppice-with-standard to coppice-selection to high forest
- 20 • The resulting forest homogenization induced a decrease in biodiversity hosting capacity
- 21 • The socio-cultural legacy of this ancient forest is currently threatened

22

Abstract

Centuries of charcoal making has profoundly shaped European mountain forest ecosystems. However, it remains difficult to assess this impact due to a lack of knowledge about the full operational sequence and related silvicultural systems. To accurately reconstruct such practices and shed light on the resulting legacies, we carried out an interdisciplinary study in the Bernadouze forest, a 46 ha mountain forest of European beech (*Fagus sylvatica* L.) in the French Pyrenees. We performed a multi-proxy analysis of 7990 charcoal fragments from 28 charcoal kilns to assess the long-term changes in forest composition and structure, but also in harvesting practices and related silvicultural systems. In addition, we assessed the legacy of such practices on the forest's main features, by evaluating the biodiversity on 13 one-hectare plots using the Index of Potential Biodiversity (IBP). The 18 radiocarbon dates showed that charcoal making took place at least from the 9th-10th c. to the 19th-20th c. Beech and silver fir (*Abies alba* Mill.) were the main species used. We discerned sustainable silvicultural practices performed over the centuries without significant change. Charcoal burners harvested well calibrated wood pieces mostly from the end of the growing season until the early spring, and charred after a seasoning period. The forest was managed as a beech coppice with fir standard before being progressively transformed, between the mid 15th-mid 17th c. period, into a monospecific beech coppice, probably treated in a coppice selection system. While this management allowed forest cover to be continuously sustained, it also resulted in the homogeneity of the forest today, both in composition and structure, and led to low hosting capacity for biodiversity. Nevertheless, this ancient forest, which constitutes a high biocultural legacy due to immemorial use rights, needs dedicated management.

Key words: charcoal making practices, technical operational sequence, forest management, anthracology, IBP, centuries-old legacies

45 Introduction

46 Current forest research emphasizes the importance of developing more holistic approaches to account for the
47 long-term legacies of human activity (Bergès and Dupouey, 2017; Paradis-Grenouillet et al., 2018a; Rostain and
48 Saulieu, 2018; Bergès and Dupouey, 2020). An evolving framework of interdisciplinary research, called
49 historical ecology, has been developing over the past 40 years with significant advances over the last two
50 decades (Crumley, 1994; Szabó, 2015; Watkins, 2015; Bergès and Dupouey, 2020). In this conceptual field, the
51 forest is considered a constructed heritage, resulting from interactions between societies and the environment
52 over time. This diachronic perspective allows (i) the study of current forest stands to be carried out in historical
53 depth, (ii) a better comprehension of their evolution in the medium and long term, and finally (iii) an
54 understanding of the current structure and functioning of ecosystems in order to better guide conservation
55 objectives and to predict future dynamics (Peterken, 1993; Kirby and Watkins, 1998; Rackham, 2003; Cevalasco
56 and Moreno, 2015; Szabó, 2015; Bürgi et al., 2017).

57 Among key analytical approaches developed in the field of historical forest ecology, the study of soil and
58 archaeological charcoal, called anthracology, provided tools to renew our knowledge of woodland landscapes
59 from a historical and socio-environmental perspective in the *longue durée* of forest ecosystems (Nelle et al.,
60 2013; Feiss et al., 2017; Dufraisie and Coubray, 2018; Moreno et al., 2019).

61 In the early 1990s, researchers became increasingly interested in charcoal remains from artisanal and industrial
62 activities, particularly those recorded in former charcoal making terraces, so-called charcoal kilns (CKs). Such
63 charcoal remains testified to pre-industrial logging-induced spatio-temporal changes in the structure and
64 composition of forest ecosystems (Hillebrecht, 1982; Bielenin, 1992; Pott et al., 1992; Davasce, 2000;
65 Montanari et al., 2000; Bonhôte et al., 2002; Ludemann and Nelle, 2002). Contrary to the long-accepted idea
66 that pre-industrial woodfuel supply activities heavily contributed to land clearance, past societies were shown to
67 have introduced management systems that, while degrading forest ecosystems, nonetheless maintained tree
68 cover and woodland resources (Deforce et al., 2013; Py et al., 2013; Samojlik et al., 2013; Schmidt et al., 2016;
69 Benatti et al., 2018).

70 A better knowledge of forest management history and woodland changes induced by charcoal making
71 necessarily requires an in-depth understanding of the resulting practices (logging, species and wood caliber
72 selection, etc.) and their temporalities (seasonality, cutting rotation, etc.). Available data mainly rely on
73 ethnographic observation (Acovitsioti-Hameau, 2001; Hanus, 2007; Burri, 2008; Burri et al., 2010), modern

74 scholarly treatises (e.g. Biringuccio, 1556; Della Fratta Montalbano, 1678; Duhamel Du Monceau, 1761;
75 Bellenghi, 1816; Marié de L'Isle, 1835; Malepeyre, 1836; Valerius, 1851; Percy, 1864; Svedelius et al., 1875;
76 Dromart, 1880), and/or experimental approaches (Poggi and Métaillé, 1998; Fregni and Maccaferri, 2010;
77 Paradis-Grenouillet, 2012). However, the only way to develop an understanding of local practices and, more
78 broadly, to learn how people worked before modern times, is to study CK remains. In this perspective, wood
79 scientists are developing dendrometric and dendrochronological tools to accurately reconstruct the entire
80 technical operational sequence of charcoal production, from wood selection and removal to silviculture
81 management and the charring process (Ludemann and Nelle, 2002; Strachan et al., 2013; Raab et al., 2015;
82 Dupin, 2018; Paradis-Grenouillet et al., 2018a; 2018b; Fouédjeu et al., 2021).

83 This study proposes an innovative combination of archaeological, anthracological, and ecological approaches to
84 characterise past charcoal making practices and better understand how related silvicultural systems have driven
85 shifts in the forest trajectories using the well-known example of the Northern central Pyrenees. In this area, the
86 rarefaction of silver fir (*Abies alba* Mill.) and the expansion of beech (*Fagus sylvatica* L.) are commonly
87 considered a consequence of harvesting selection for the production of charcoal needed in the historical iron
88 metallurgy industry (Davassee, 2000; Bonhôte et al., 2002; Py-Saragaglia et al., 2018; 2019). Beech, able to
89 regrow from stump or stool, would have been favoured by coppicing (Davassee and Galop, 1990; Davassee,
90 2000). However, it remains difficult to precisely date the shift from a mixed beech-fir forest to a monospecific
91 beech forest. In addition, while the development of beech coppice silviculture is a common working hypothesis,
92 there is no solid evidence for this from before the 17th c. (Davassee, 2000). Therefore, the origins of coppicing in
93 a northern Pyrenean beech-fir forest, its rise and fall and its relation to charcoal production, need further
94 investigation. Furthermore, the extent to which the legacies of forest conversion has constrained current
95 biodiversity is still unexamined.

96 To address all of these issues, we applied an interdisciplinary approach in the forest of Bernadouze (Suc-et-
97 Sentenac, Ariège), which is representative of most Northern Pyrenean mountain forests deemed to have a long-
98 lasting charcoal making history. We suggest that the current forest structure and composition result from many
99 centuries of charcoal burning practices. We test this assumption by suggesting (i) that charcoal-making history,
100 related forest management practices and their evolution can be reconstructed using charcoal analyses, and (ii)
101 that current biodiversity and habitat functions are still impacted by the legacies of this history of charcoal
102 production. Beyond these research hypotheses, we discuss how the insights of historical ecology could

contribute to improving future forest management and, more broadly, to taking forests as biocultural heritage in future policies.




Fig. 1: Sampling design of the Bernadouze forest green points represent the totality of recorded and sampled CKs (n=80); red points, CKs (n=17) dated by C14; red squares, CKs with dendro-anthracological analysis

(n=25); red circles, IBP plots (n=13); “fires” represent pedoanthracological pits (n=3); yellow points represent pastoral sites (n=5) .

1. Materials and methods

1.1. The survey area

The study area (1300 to 1450 m.a.s.l) detailed in Saulnier et al. (2019), is located on the northern slope of the Pyrenees in the upper Vicdessos valley, hosting a permanent Human-Environment Observatory (called in French “OHM Haut-Vicdessos”) aiming at studying long-term human-environment relationships (Fig. 1). Soils (rendisols, calcisols and brunisols), which developed on recrystallized limestone, are characterised by a silty-clayey texture in various proportions and by basic pH values. Current vegetation consists of three main beech stand types: (i) high forest resulting from the conversion of former selection coppice with individuals aged between 130-160 years, with rare and scattered fir trees (Py-Saragaglia et al., 2018; Fouédjeu et al., 2021), (ii) tall-grown coppices of old trees from a coppice selection method, and (iii) tall-grown coppices around the peat bog boundary that have been browsed in the past. A set-aside area (90 ha), which includes the study area and an adjacent peat bog, is managed by the French National Forest Office as a Managed Biological Reserve (RBD), which allows for occasional active management interventions for pre-defined conservation objectives. To this end, group-selection harvesting has been prescribed in the buffer zone surrounding the bog.

1.2. Archaeological survey, CK selection and sampling strategy

To reconstruct charcoal making chronology and its intensity through time, we carried out a systematic archaeological field survey that afforded the detection of 80 CKs on about 46 ha (Fig. 1). In addition, five undated pastoral sites were georeferenced.

The study area was divided into four Analysis Spatial Units (ASU) according to their exposure and topography (Fig.1). In each ASU, for charcoal analysis, we randomly selected 4 to 9 CKs (n=28) in order to be fully representative of the ASU studied, i.e. proportionally of the total numbers of CKs per ASU.

Because most CKs exhibited a thick layer of charcoal (ranging from 20 to 120 cm thick) mixed with a very black and homogeneous soil, generally without any stratigraphy, stepwise sampling was carried out using a soil auger, every 20 cm in depth from the first horizon (A) to the ochre sterile and continuous mineral layer as is

detailed in Py-Saragaglia et al. (2017). Samples for each 20 cm layer from all sampling points carried out on the same terrace (at least six per CK) work together and form the so-called “charcoal layer” (CL). A total of 57 CLs were analyzed (Fig. 1).

Among all studied CKs, we selected 18 charcoal samples from 17 CKs for radiocarbon analysis. These samples are constituted by a mix of several charcoal fragments - preferably young twigs with bark, charcoal fragments with bark, or bark alone -, dated as a single sample. This approach aims to reduce errors that may be associated with dating a single fragment that may be intrusive (Py et al., 2013). Radiocarbon dating by Accelerator Mass Spectrometry (AMS) was calibrated using the OxCal program, version 4.4 Intcal 20 database (Reimer et al., 2020).

1.3. Analysis of the charcoal dataset

Tree species used for charcoal production were identified at the species, genus, family or sub-family level using a reflected light microscope (Leica DM4 with magnifications of 100, 200, 500 and 1000), xylological atlases (Schweingruber, 1990; Vernet et al., 2001) and the reference collection of the GEODE laboratory.

To determine the annual timing of logging operations, we observed the state of progress of the last ring’s formation when charcoal has retained bark cells (Fig. 2A). We then grouped samples into two categories: (i) vegetative period (last ring ended between early wood and the end of early wood) and (ii) end of vegetative period (last ring ended between late wood and end of late wood). Hereafter, the timing of harvesting was calculated as:

$$T_h = ((N_{vp} - N_{evp}) / N_{cb}) * 100$$

where T_h is the timing of harvesting, N_{vp} and N_{evp} are, respectively, the number of charcoal samples for which the terminal ring ended in early wood/late early wood or in late wood, and N_{cb} is the number of charcoal samples with bark remains.




Fig. 2 Photos illustrating: terminal growth rings of charcoal fragments that retained their bark (A), hyphae fungi contamination (B), cellular collapse (C), and radial cracks (D)

In order to reconstruct harvested wood diameters, we applied the three-step method described in Paradis-Grenouillet et al. (2015) (see Appendix 1 for more details) (Fig.3) : (step 1) we measure the radius of curvature of charcoals with at least 3 rings, and a cross-section $\geq 3\text{mm}$ using the AnthracoloJ program; (step 2) we reconstitute the different diameters of logs composing the wood pile by comparing archaeological data with 405 mathematical simulations and the Total Deviation (TD) shows the reliability of comparison; (step 3) we compare the reconstituted wood pile to wood pile references established in current stands in order to characterize the shape/morphology of the harvested stands (i.e. different types of coppice, high forest).




Fig. 3. Illustration of the three steps required to find wood diameter (Paradis-Grenouillet et al., 2015, modified).

To assess the logging cyclicity and to reconstruct coppicing rotation, we systematically counted the number of rings observed on charcoal pieces, except for fragments where the number of rings was not readable, giving the minimum age of wood pieces used by charcoal burners.

In order to characterize the state of wood harvested (dead or alive) and test for an eventual storage period in forest, we characterized the degree of deterioration of the charred wood using (i) the mean number of radial cracks (RC) per cm² on the transverse plane of the charcoal (Fig. 2D) (Théry-Parisot and Henry, 2012), (ii) the degree of charred fungal hyphae infestation in cells (Moskal-del Hoyo et al., 2010; Saulnier et al., 2019) and (iii) the cellular collapse that informs about the decay stage of the wood (healthy or decayed) (Fig. 2B, 2C) (Henry and Théry-Parisot, 2014). Four contamination levels (Cont. 0 to Cont. 3) were defined (see appendix 4, Table 1). A Decay Index (DI) was calculated based on a weighted mean average (Théry-Parisot et al., 2016):

$$DI = (nb \text{ "Cont.1" } \times 1 + nb \text{ "Cont.2" } \times 2 + nb \text{ "Cont.3" } \times 3) / nb \text{ total.}$$

1.4. Assessment of current stand capacity to host biodiversity

To assess the effect of former charcoal practices on current stand biodiversity, a diagnosis was carried out using the Index of Biodiversity Potential (IBP; Larrieu and Gonin, 2008) on 13 one-hectare plots, centered around a subset of the assessed CKs. The IBP is a biodiversity evaluation tool according to Larsson (2001) that combines ten key historical, structural, and compositional factors for forest-dwelling species, which are easily and directly measurable in the field (for more details, see Gosselin and Larrieu (2020) and Appendix 2). It can be used as a rapid habitat assessment method to evaluate the hosting capacity of stands for forest biodiversity (Larrieu et al., 2012; Bouget et al., 2014; Larrieu et al., 2019a). The raw number of items (e.g. number of tree species, habitat-trees, etc.) counted when getting through the stand is compared with thresholds values to set a score for each factor which ranges from 0 to 5. Then, the sum of the scores given for each of the 10 factors (potential range 0-50) provides an index of the stand's hosting capacity (Larrieu and Gonin, 2008).

2. Results

2.1. Chronology of charcoal making activity

The 18 radiocarbon dates are distributed from the 9th-10th c. to the 19th-20th c., with a maximum number concentrated from the mid-15th to 19th c. We classified them into five historical periods based on radiocarbon ranges (Table 1; Fig. 4).

Table 1: Historical periods based on radiocarbon range of charcoal production in the Bernadouze forest

Historical periods based on radiocarbon ranges (1 or 2 σ = 70 to 95.4% of probability)	Number of dates	N° charcoal kiln
19th c. - 20th c.	2	4, 74
Mid-17th c. -19th c.	4	14, 17, 29, 63
Mid-15th c. - mid 17th c	7	1, 6, 7, 17, 19, 34, 79
End 13th c. - mid 15th c.	3	11, 27, 49
9th c. - 10th c.	2	8, 62




Fig. 4. Plot of radiocarbon dates for charcoal kilns in the Bernadouze forest distributed into five historical periods. Data were calibrated with OxCal v4.4.4 Bronk Ramsey (2021); r:5 Atmospheric data from Reimer et al. 2020.

2.2. Wood selection for charcoal production

2.2.1. Selected tree species and seasonality of logging

We identified two main species (Fig. 5 and 6): beech (n=5605) and fir (n=1438). We also identified 19 other tree taxa with relative frequencies below 1% (Fig. 5 and 7): (i) pioneer taxa, i.e. legume family (Fabaceae), birch (*Betula*), willow (*Salix*), grey alder (*Alnus incana* L.), alder (*Alnus*), juniper (*Juniperus*), (ii) post-pioneer taxa, i.e. wild cherry (*Prunus*), scots/mountain pine (*Pinus*), evergreen and deciduous oak (*Quercus*), apple sub-family (Rosaceae, Maloideae), mountain ash (*Rosaceae, Maloideae cf. Sorbus*), hazel (*Corylus*), ash (*Fraxinus*), rose (*Rosa*), elm (*Ulmus*), and, (iii) shade-tolerant tree taxa (so-called “dryads”), i.e. holly (*Ilex aquifolium* L.), common yew (*Taxus baccata* L.) and fir/juniper (*Abies alba/Juniperus*). When similar anthracological (tree taxa

2.2.2. Wood diameter reconstruction, number of rings and the minimum age of the charred wood

We restituted the radius curvature of 3003 charcoal pieces, i.e. those with a minimum transverse section of 3 mm, from 45 CLs (Appendix 1), including 2602 beech fragments distributed across 25 CKs (Appendix 4, Fig. 2) and 401 of fir, distributed across 4 CKs (Appendix 4, Fig. 3).

Data from 41 CLs out of the total of 43 CLs for beech, and from all CLs for fir highlight good correlation (TD between 20 and 40 %), sometimes very good (TD < 20%), with simulated data (Appendix 4, Fig. 2 and 3).

Results for beech show the dominance of small diameters (less than 5 cm), and the low presence of medium (5 to 10 cm) and large (10-20 cm) diameters (Fig. 6; Appendix 4, Fig. 2). Regarding fir, we observed a mix of all diameter classes with the dominance of the 2-15 cm diameters and a small proportion of very large wood (>20 cm) (Fig. 6; Appendix 4, Fig. 3).

The minimum number of rings was counted on 5534 beech charcoal samples and 1436 fir samples (Fig. 6; Appendix 3). We observed the predominance of charcoal with 1 to 5 tree rings preserved: 55.9% for beech and 66.1% for fir. Charcoal with 6-15 rings represented 37.1% for beech and 25.6% for fir. In addition, a few samples kept 16-30 rings: 6.2% for beech and 6.7% for fir. Charcoal with more than 30 rings is rare: 0.6% of beech and 1.3% of fir.

2.2% of charcoal (n=155) retained pith and bark: 140 beech and 15 fir (Appendix 3). For these samples, the total number of rings is known and, therefore, so is the age of the branch, twig, or stem. Beech charcoal had from 1 to 35 rings and fir charcoal from 3 to 21 rings.

2.2.3. State of wood before carbonization

The decay index (DI), with a range from 0 to 1.5, was calculated for 7043 charcoal pieces, of which 5605 beech and 1438 fir.

Globally, for both taxa (Fig. 6), we observed 25 CLs of beech and 13 CLs of fir with low decay ($0.1 < DI \leq 0.5$).

We also observed moderate decay ($0.5 < DI < 1$) in 30 beech CLs and 16 fir CLs and very high decay ($DI \geq 1$) in 6 CLs for beech and 12 for fir. Finally, very low decay ($DI \leq 0.1$) was observed in 1 CL for beech and 5 CLs for fir (Appendix 4, Fig. 4A,B).

247 In the same corpus, we identified RCs on 1421 beech charcoal pieces and 156 on fir charcoal pieces. A total of
248 5466 charcoal have no RC. The mean RC/cm² per CLs ranges from 0 to 13.6 for beech charcoal and from 0 to
249 53.1 for fir. Results show the use of seasoned wood in 53 CLs and green wood in 1 CL for beech (Appendix 4,
250 Fig.5A) and the use of seasoned wood in 36 CLs and green wood in 6 CLs for fir (Fig. 6; Appendix 4, Fig.5B).





Fig. 6. Chronological representation of radiocarbon dated charcoal kilns (CKs). The oldest CKs are at the bottom and the most recent at the top. All data concerning the taxa found through anthracological analysis have been represented, including data on harvest seasonality, wood diameter reconstruction, the decay and moisture stage of the wood, and the number of growth rings.

2.3. Current stand capacity to host biodiversity

The IBP total scores ranged from 22 to 30, i.e. from 40% to 60% of the potential maximum. Values were rather average for the management score (12-20/35) (Fig. 7A), though the context scores ranged rather high (7-15/15) (Fig. 7B). The number of tree-species ranged from 1 to 7 per plot; beech always dominated; silver fir occurred on three plots only. Vegetation generally presents little vertical stratification, with only two or three strata present (23%). Very few deadwood items occurred, particularly those with large diameters. Very large trees (diameter at breast height > 67.5 cm) are also rare, occurring on only four plots. However, habitat-trees are numerous (from 8 to 131/ha) and provide a high diversity of tree-related microhabitat types. Stand openness is high on average (34%), ranging from 1.5 to 90%. Aquatic and rocky habitats occur on 57 % and 100 % of the plots, respectively (Fig. 7A).

A/ IBP score (absolute value) for stand and current management



B/ IBP score (absolute value) for context




Fig. 7. IBP results (in %): A) related to the seven factors describing current stand; B) Related to the three factors describing context

269 3. Discussion

270 3.1. General trends in the historical charcoal manufacturing practices at Bernadouze

271 3.1.1. Mono- or multi-species CKs depend on wood availability and economic strategies

272 Generally the choice to use one or more species is driven by environmental availability in the supply area as
273 well as technical purposes, i.e. the quality of charcoal wanted and its final use (Davassee, 2000; Durand et al.,
274 2010; Adeniji et al., 2015). For example, beech wood was appreciated by some metallurgists because of its very
275 low ash content and fairly high density, which allows slower burning (Maillat, 1942; Bonhôte and Fruhauf,
276 1990a). Furthermore, as with other hardwoods, beech is known to produce a greater quantity of charcoal and to
277 increase charcoal productivity (Valerius, 1851, 222; Bonhôte and Fruhauf, 1990).

278 In the Bernadouze forest, half of studied CLs (n=32) are nearly monospecific (up to 90 % beech), and 29 CLs
279 are multi-specific, with the co-dominance of beech and fir in varying proportions (Fig. 5 and 6). In a
280 homogeneous environmental context (type of soil, climate, exposure), such results may reflect a change in forest
281 cover, and/or in silvicultural management, and/or in charcoal making practices (e.g. Davassee, 2000; Tolksdorf et
282 al., 2015). To clarify which hypothesis is correct, we need to know if the anthracological records reflect the
283 successive charring of monospecific wood piles with either fir or beech, or a mixture of two dominant species in
284 the same wood pile. An argument in favor of successive monospecific charring is that CL records generally
285 contain residues of several, rather than a single carbonization (Gebhardt, 2007; Dupin et al., 2019; Fouédjeu et
286 al., 2021). The vertical and lateral displacement of remaining charcoal pieces may mix charcoal from the
287 alternating charring of monospecific wood piles. In addition to this taphonomic consideration, mixing was
288 generally discouraged by scholars and forest engineers (Duhamel Du Monceau, 1761, 8; Valerius, 1851, 202).

289 However, the mixing of hardwood and softwood could occur when the amount of either wood species was
290 insufficient to make up a whole CK (Valerius, 1851, 224). For example, in Central Pyrenees (Ariège, Aude),
291 charcoal burners used a mix of beech with oak, hazelnut, or a weak proportion of fir ($\frac{1}{5}$ or less) during the 18th
292 and 19th c. (Lapassat, 1983; Fruhauf, 1990; Bonhôte, 1998, 220-223). Because of this practice, the possibility
293 that fir and beech were charred together in the same CK at Bernadouze cannot be excluded, but in highly
294 different proportions probably due to local wood availability and/or economic strategies.

In addition to the species, several other factors may affect the quality of charcoal produced: wood pile construction, caliber, age of wood pieces, tree growth pattern, moisture content, decay stage, etc. (Biringuccio, 1556, 84; Duhamel Du Monceau, 1761, 9-10; Percy, 1864, 201; Svedelius et al., 1875, 24).

3.1.2. The use of well-calibrated pieces of woods harvested between the end of the growing season and vegetative recovery

Our results highlight the preferential use of small-caliber wood (2 to 5 cm in diameter) for beech, and medium (5 to 10 cm) and large-caliber wood (15 cm) for fir (Fig. 6), as already evidenced in other areas (Nelle, 2002; Deforce et al., 2013; Paradis-Grenouillet et al., 2015; Dupin, 2018; Máliš et al., 2021). This use is in line with the recommendation given in some historical treatises, i.e. the best charcoal was made with small and medium caliber wood (+/- 5 to 10 cm in diameter) (Duhamel Du Monceau, 1761, 9; Marié de L'Isle, 1835, 110; Malepeyre, 1836, 128). Moreover, the mixing of small (less than 5 cm) and large woods allows a more homogeneous carbonization and avoids wood pile collapse, and small branches or twigs are also used to fill gaps left between large pieces of wood (e.g. Dromart, 1880; Lugli and Pracchia, 1995; Bond, 2007; Burri, 2008). Another common practice consisted in splitting some of the largest wood items before carbonization to facilitate their charring (Malepeyre, 1836, 128; Valerius, 1851, 222). To conclude, we cannot absolutely exclude the possibility that this was practiced at Bernadouze, insofar as the presence of large wood in the CKs was recorded.

The relative frequency of charcoal pieces that kept bark was low, probably resulting from both carbonization and taphonomic (fragmentation) processes. But 64% of these charcoal fragments revealed that, from the end of 13th c. to the 20th c., fuelwood for charcoal production was harvested between the end of the growing season (i.e. autumn yellowing of deciduous tree leaves) and vegetative recovery (leaf appearance), as already demonstrated by many anthracological and ethnoarchaeological studies (Montanari et al., 2002; Bond, 2007; Glais, 2007, 6; Py-Saragaglia et al., 2017). This globally corresponds to the recommendations in 18th and 19th c. technical treatises i.e from the fall until the first sap rises at the end of winter (Duhamel Du Monceau, 1761, 9; Bellenghi, 1816, 43; Marié de L'Isle, 1835, 269; Valerius, 1851, 204; Percy, 1864, 208; Svedelius et al., 1875, 35-36).

3.1.3. Wood moderately seasoned before the burning season to obtain quality charcoal

In Bernadouze, charcoal burners mainly used seasoned wood, implying a seasoning phase just after cutting. In the same way, the Decay Index (DI) shows that 81.9% of the wood was weakly to moderately decayed, suggesting that most charred wood was felled healthy and then decayed during a seasoning period. An experimental study has highlighted that contamination by wood-rotting fungi usually lasts two years after tree death and occurs only during the warmer months, i.e. the short summertime, in mountain areas (Théry-Parisot, 2001, 51; Py et al., 2015). Considering the low contamination levels, the seasoning period is estimated to be less than one year. Our interpretation is in line with almost all contemporary scholarly treatises reporting a seasoning period lasting from 2 to 12 months, even if a few treatises suggested, if necessary, the possibility of charring green wood (Della Fratta Montalbano, 1678). The length of seasoning depends on the cutting season, weather, and wood diameter (e.g. Biringuccio, 1556, 85; Duhamel Du Monceau, 1761, 10; Bellenghi, 1816, 43; Malepeyre, 1836, 113), but must not be too long in order to obtain moderately seasoned wood to ensure good quality charcoal (Duhamel Du Monceau, 1761, 10; Malepeyre, 1836, 113; Percy, 1864, 208-209; Svedelius et al., 1875, 35-36).

Around a fifth of the charcoal (18.1%) was heavily altered, implying that some rotten wood was charred. This practice is not in line with scholarly treatises, which advise against the use of rotten wood which degrade both charcoal productivity and quality (Valerius, 1851, 222; Percy, 1864, 209). Nevertheless, carbonization of dead wood was sometimes recommended to recycle forestry waste (Bellenghi, 1816, 43). In this sense, a 1302 law regulating charcoal production in Vicdessos required charcoal burners to use dry and dead trees before cutting living ones (Verna, 2001, 72, 82).

After seasoning, wood was very often charred as early as the following spring and continued until the first snows, as has been demonstrated for another Pyrenean valley (Bonhôte, 1998, 220-223; Py-Saragaglia et al., 2017) and prescribed in the historical treatises (Duhamel Du Monceau, 1761, 10; Malepeyre, 1836, 128; Valerius, 1851, 222). Such a claim has been made for other regions in Western Europe (e.g. Burri, 2012, 214-224; Hanus, 2018).

3.2. Some insights about silvicultural practices over time

Charcoal results show that charcoal production started in the 9th-10th c. in a mixed beech-fir forest and ended in the first half of the 20th c. in a mono-specific beech woodland with very few occurrences of fir. They suggest that the Bernadouze forest was managed, from the 9th-10th to the mid-17th c., as a beech coppice with fir

standards leading to a multi-level forest. In Europe, at least from the late Middle Ages onwards, coppicing was the most common silviculture treatment for broadleaved species to produce fuel (Peterken, 1993; Unrau et al., 2018). But, among the wide range of possible operating systems (e.g. simple coppice, coppice selection system, pollarding, etc.), it is still difficult to accurately characterize past coppice treatment from charcoal alone because of the lack of charcoal diameter references from these different kinds of beech coppices.

Reports from the Forest Administration mention beech coppice in 1669, 1807, and 1860, without specifying the type of coppice forest. At Bernadouze, the first explicit mention of a selection coppice system dates from 1892 (Chassinat, 1892), but we do not know when this system was implemented exactly. We hypothesize that the implementation of this system was consecutive to a strong harvesting of fir, occurring during the mid-15th-mid-17th c., resulting in poor fir regeneration and its progressive rarefaction in the overstory. With fir trees providing the shade which favors the growth of beech shoots, the coppice selection system and selective cuttings were technical solutions to respond to the increasing demand for charcoal from the mid-17th c. Moreover, the exposure to multiple cutting cycles could have led to the increasing shoot production per resprouting stump, but it also reduces shoot diameter. This hypothesis is supported by evolving diameters of beech charcoal, from a mix including large diameters from the 9th-10th c. to the mid 15th c., to the almost exclusive use of small diameters mixed with some medium diameters from the mid 15th-mid 17th c. (Fig. 6).

From the end of the 19th c., thinning, which consisted in removing undesirable individuals from the main stool as well as standing deadwood and decaying stems, was prescribed by forest managers to improve productivity in the Bernadouze forest (Chassinat, 1892; Rochebrunel, 1920; Noiriel, 1965). Thinning consisted in removing undesirable individuals from the main stool - such as defective works, poorly or over abundantly stemmed growth - as well as standing deadwood and decaying stems. Therefore, the partly altered charcoal found in the CKs could reflect from the recycling of this management technique's byproducts. While iron production-related demand gradually ceased in the 19th c. (Cantelaube, 2005, 693-734), harvesting for local needs (e.g. blacksmith shops) continued. Due to socio-economic changes in the 20th c., degraded beech coppice trees were gradually converted to high forest using cleaning-cuts and selective thinning (Rochebrunel, 1920; Noiriel, 1965).

In Bernadouze, the minimum harvesting age for beech ranges from 6 to 15 years (37.1%) and 16 to 30 years (6.1%) (Fig. 6). Possible coppice rotation would range from 6 to 30 years, with a dominance of wood harvested between 6 and 15 years. The over-representation of 1-5 years-old charcoal may result from fragmentation accentuated by (i) the scraping and cleaning of the terrace by charcoal burners before each carbonization, (ii)

taphonomic processes in soils, and (iii) the archaeological sampling and sieving (Chrzavzez, 2013; Blondel et al., 2018; Fouédjeu et al., 2021).

There is little documentation of former beech coppicing rotation in Ariège. During the 19th c., cutting cycles ranged from 15 to 18 years, according to Forest managers and Mines engineers (Cantelaube, 2005, 215). Currently, for coppice selection systems, undergrowth cutting rotation is commonly estimated at 6 to 12 years (total cycle of 36 years) in the western Italian Alps and the Apennines, and 10 to 15 years in the Pyrenees (total cycle c. 30 years) (Salvador, 1930; Coppini and Hermanin, 2007; Nocentini, 2009; Nicolescu et al., 2018). Such variability depends on (i) the target diameter and product (fuel wood, charcoal, poles), and (ii) the capacity for coppice regeneration, which varies according to species and site conditions (Buckley and Mills, 2015). Regarding the seasonality of harvesting, our results are in line with previous research which demonstrated that shoot growth and coppice production are optimized when shoots are cut between late winter and early spring, which corresponds to the best season for its growth (Nicolescu et al., 2018).

Regarding fir, 25.6% of charcoal is between 6-15 years and 6.8% is between 16-30 years old (Fig. 6). This does not match the expected age for standards because their rotation length is generally at least twice as long as that of coppices (Burley et al., 2004). According to Huffel (1926, 68), fir trees could be harvested from 40 years of age onwards. In the Eastern Pyrenees, the mid-17th c. Forest Law set the age of exploitability of the fir trees scattered in beech woodlands from 80 to 100 years (Fruhauf, 1980, 31; Bartoli, 2011, 141-142). Because of the difficulty of hauling fir timber due to topography and the lack of buoyant watercourses, 20th c. forest engineers recommended exploiting trees of about 45-50 cm in diameter at breast height (Salvador, 1930). When fir trees had reached the target diameter, they were harvested and only the best logs were transported to sawmills, while the crowns, tops, branches and slash could be recycled for charcoal production (Bonhôte and Fruhauf, 1990b). Despite fragmentation bias (which limits the number of tree rings retained), charcoal age results may reflect the use of such forestry waste.

3.3. The long- term legacy of historical human practices

3.3.1. An ancient but immature forest, with weak tree-species diversity and hosting capacity

Forest ancientness refers to the duration of uninterrupted wood cover, independent of the past and current management practices that may have changed forest features (Dupouey et al., 2002). Forest ancientness impacts communities for several taxa, in particular those species with weak recolonization capacity such as vascular

plants (Hermy et al., 1999), lichens (Rose, 1993), ground beetles (Assmann, 1999) and ectomycorrhizal fungi (Diedhiou et al., 2009).

Ancientness is usually evaluated by comparing current forest maps to older maps (Cateau et al., 2015). In France, the *Etat-Major* maps, established when forest cover was at its lowest in the mid-19th c., are considered to be the most accurate of available maps. On this map, our study area is represented with forest cover. The assessment of ancientness for longer periods is exceedingly difficult, due to the absence of continuous historical records. However, our results combined with a previous soil charcoal study provide evidence of continuous forest cover from the Middle-Late Bronze Age to the modern period, through the absence of identified charcoal from heathland species as well as the scarcity of pioneer species (Saulnier et al., 2019). Until the Carolingian period (610-722 AD), the forest hosted the three main European mountain dryads: beech, fir and yew (Saulnier et al. 2019). Then, the combined or successive effects of charcoal production and agro-sylvo-pastoral activities led to the progressive shift of forest composition: the disappearance of yew, the rarefaction of fir, and the favourisation of the more competitive beech eliminating post-pioneer light-demanding species in modern times (Py-Saragaglia et al., 2018; Saulnier et al., 2019).

Forest maturity refers to the state of biological development of the trees within a stand (age diameter and development stage of the oldest and largest trees), and also to the quantity and diversity of dead wood and tree-related microhabitats (TreMs) (Cateau et al., 2015). Deadwood and TreMs are pivotal resources for many forest-dwelling taxa (Stokland et al., 2012; Larrieu et al., 2018). Very large trees provide the entire range of TreMs (Larrieu et al., 2014b) including e.g. perched deadwood, a particularly important resource for saproxylic beetles (Bouget et al., 2011) that increases their functional diversity (Gossner et al., 2013). Maturity is heavily driven by past and present forest management. To display maturity attributes similar to a near-natural forest, the forest must not have been subject to harvest for several decades (Bouget et al., 2014; Paillet et al., 2015). When management is based on coppicing, more than 70 years of being set-aside are needed to recover both the amount and diversity of deadwood and TreMs significantly larger than those observed after harvest (Larrieu et al., 2017; 2019b).

At Bernadouze, regular and short-rotation logging, intended to remove all the shoots that have reached the target size, has not favoured the presence of dead trees (standing and on the ground), very large timber, and old trees (the oldest trees are 130-160 years old, as shown in Fouédjeu et al., 2021). Today, a large part of the forest is managed according to group-selection from former stumps, except for five plots where there is still old and

stunted coppice. However, given the geographical position (northern central Pyrenees) and the stationary conditions (north-facing mountain level, soil types), the potential habitat is a mixed forest of deciduous and coniferous trees dominated by beech, silver fir and yew (Bardat et al., 2004; <http://www.euforgen.org/species/>). This potential mixed forest would be host to a variety of secondary species (pioneer and post-pioneer species such as alder, European mountain ash, ash tree, willow etc.), more or less localized at particular stations (rock outcrops, wetlands with waterlogged soil) or openings associated with natural and/or anthropogenic disturbances (clearings, edges, and other areas with a well-developed herb layer composed of flowering plants, etc.). The loss of tree-species diversity provides a much lower hosting capacity because biodiversity associated with deciduous and coniferous trees do not fully overlap (e.g. Bouget et al., 2019). It can be assumed that, although secondary species are always scattered or spatially localized in natural beech-dominated forests (e.g. Commarmot et al., 2005), their extreme rarity at Bernadouze confers a decrease of hosting capacity, since most pioneer and post-pioneer species have partly gender-specific corteges (e.g. Ryvardeen and Melo, 2014; Bouget et al., 2019). Moreover, these secondary species are crucial for the silvigenetic cycle and for biodiversity, because they provide deadwood and dendromicrohabitats during the phases of the cycle when these attributes, provided by dryads, are scarce (Larrieu et al., 2014a). The absence of very large trees and the scarcity of deadwood also restrict the forest's hosting capacity. However, fairly high density and diversity of TreMs were observed, which is favourable to biodiversity as these structures host a wide range of taxa (Larrieu et al., 2018). This finding could be partly tied to the increased density of trunk-based cavities induced by the conversion of former stump coppices into high forest (Gouix, 2011).

3.3.2. Ensuring collective memory and cultural heritage

Ancientness contributes significantly to the patrimonialization of a forest. According to oral testimonies recorded in the CNRS documentary film “*Haut-Videssos, une vallée et des hommes*”, this heritage value has a significant socio-cultural and collective dimension (<https://www.youtube.com/watch?v=LjXtsRTaACk>). It might have contributed to reinforcing the feeling of belonging to a social group that shares an attachment to a forest which they inherited from medieval common rights granted to the population by their lords. Despite the nationalization of the forest and the erosion of common rights during the 19th c., this attachment is still present today. Indeed the last, large group-selection harvest, carried out in 2016, sparked considerable public outcry among the local population. From the perspective of their historical depth and their impact on forest cover, charcoal-making platforms are a key component - beside pastoral remains - of the heritage value of the

Bernadouze forest. Such vernacular heritage reflects the close relationship between society and forests. Today, it is immediately necessary to compile the living memory of the native inhabitants of the valley, in order (i) to better understand the importance of the place in the attachment of the local populations to the Bernadouze forest (Sebastien, 2020) and (ii) to pass it down to future generations through educational and cultural activities that capture the millennial history of this forest (e.g. discovery trails etc.). To be successful, these activities must include forest managers, scientists and inhabitants. In Europe and France, ancient forests are estimated to represent between 30-50% of the total forest area (Cateau et al., 2015; Bergès and Dupouey, 2017). It is urgent to conserve them (Jacquemin et al., 2014; Bergès and Dupouey, 2020). In order to offset the regular harvesting that has taken place over the centuries, and which has profoundly modified the natural cycles of the Bernadouze forest, managers have to consider these legacies as they implement management strategy. This management should also promote biocultural issues.

4. Conclusion and recommendations for the Bernadouze forest

Our interdisciplinary approach has allowed an accurate characterization of centuries-old charcoal-making practices, integrating the methods of collecting wood and managing the forest, which are essential for a better understanding of the legacy effects of these practices on current landscapes.

We highlighted how sustainable charcoal making practices emerged and took root from the end of the 13th to the 15th centuries without significant change. Starting in the Modern era (mid-15th-mid-17th c), socio-economic strategies heavily contributed to the forest homogenization processes through the conversion of coppice-with-standards (leading to the rarefaction of silver fir) into a beech coppice selection system. Finally, the shift from historical coppicing systems to beech high forest during the 20th c. led to the rejuvenation of stands, inducing a reduction in the biodiversity hosting capacity of the forest. The continuous removal of dead wood, in these successive management strategies, has likely contributed in great part to biodiversity loss, specifically in saproxylic species.

In order to reverse the effects of past practices and increase the forest's biodiversity hosting capacity, we recommend : (i) conserving all firs and secondary tree-species already present and increasing the density of the latter by creating gaps large enough to be suitable for saplings of pioneers, (ii) setting aside areas to reach high degrees of maturity at a local level, (iii) re-introducing yew in areas at a distance from pasture (potential conflict with pastoral uses must be considered), making it possible to recover a degree of naturalness at the site in the

long term, (iv) increasing deadwood components (e.g. by leaving the crowns of felled trees on the ground and avoiding regular harvesting of scattered fallen trees) and conserving a high density of habitat-trees to increase saproxylic resources significantly (most tree-related microhabitats contain decaying wood).

Aged coppices, which result from the extension of rotation cycles after the suspension of charcoal production, provide interesting ecological niches due to their heterogeneous structure and to the presence of dead wood. These coppices require adapted silviculture practices to maintain these characteristics and their associated ecological niches (Nicolescu et al., 2018).

From a scientific perspective, it is now necessary to renew research by considering the complexity of the multiple uses of the forest over time. To do so, other drivers must be investigated, such as pastoralism, timber production, and hunting, to understand their combined effect on forest ecosystems and on the shaping of the landscape. Finally, we suggest using this case study for scientific mediation related to the multi-secular relations between bloomeries and the dynamics of Pyrenean mountain forests.

Credit authorship statement

Léonel Fouédjeu: Field investigation, Laboratory data acquisition, Data analysis methodology, Data analysis, Writing, editing. **Sandrine Paradis-Grenouillet:** Supervision, Data analysis methodology, Data analysis, Writing, editing. **Laurent Larrieu:** Supervision, Field investigation, Data analysis, Writing, editing. **Mélanie Saulnier:** Supervision, Field investigation, Writing, editing. **Sylvain Burri:** Supervision, Field investigation, Writing, editing. **Vanessa Py-Saragaglia:** Supervision, Field investigation, Data analysis methodology, Writing, editing.

Acknowledgements

This paper forms part of L. Fouédjeu's doctoral thesis co-supervised by V. Py-Saragaglia and D. Galop and funded by the University Toulouse 2 Jean Jaurès. Financial support for field investigations was provided by the FODYNA and TRANSYLVE projects directed by V. Py-Saragaglia and S. Burri. These projects are part of the OHM-Haut Vicdessos interdisciplinary program led by D. Galop and (co)funded by the LabEx DRIIHM, the French program "Investissements d'Avenir" (ANR-11-LABX-0010) which is managed by the ANR. The authors

519 express their gratitude to S. Buscaino, V. Pescini, R. Cunill, E. Faure, N. Badache and V. Labbas for help during
520 fieldwork. Finally, we thank Evan Fisher for reviewing and correcting english.

521 References

- 522 Acovitsioti-Hameau, A., 2001. Pratiques et constructions de l'espace sylvopastoral dans le centre-Var : état
523 présent et rétrospective (thesis). Aix-Marseille 1.
- 524 Adeniji, O.A., Zaccheaus, O.S., Ojo, B.S., Adedeji, A.S., 2015. Charcoal Production and Producers' Tree
525 Species Preference in Borgu Local Government Area of Niger State, Nigeria. *Journal of Energy*
526 *Technologies and Policy* 5, 1.
- 527 Assmann, T., 1999. The ground beetle fauna of ancient and recent woodlands in the lowlands of north-west
528 Germany (Coleoptera, Carabidae). *Biodiversity & Conservation* 8, 1499–1517.
529 doi:10.1023/A:1008974413376
- 530 Bardat, J., Bioret, F., Botineau, M., Bouillet, V., Delpech, R., Géhu, J.-M., Haury, J., Lacoste, A., Rameau, J.-C.,
531 Royer, J.-M., Roux, G., Touffet, J., 2004. *Prodrome des végétations de France*. Museum National
532 d'Histoire Naturelle, 61, pp.171, 2004, Collection Patrimoines Naturels, 2-85653-563-1.
- 533 Bartoli, M., 2011. Louis de Froidour, 1626?-1685: notre héritage forestier, Les Dossiers Forestiers n°23. Office
534 national des forêts, Paris.
- 535 Bellenghi, A., 1816. Articoli sulla coltivazione dei boschi nel Piceno e nell'Umbria e sulla utilità degli alberi
536 indigeni di d. Albertino Bellenghi abate camaldolese ... Con note interessanti la medicina indigena, le
537 arti meccaniche, e l'economia domestica. Francesco Bourliè, Roma.
- 538 Benatti, A., Bal, M., Allée, P., Bosi, G., Dallai, D., Mercuri, A.M., 2018. Charcoal Kilns in the Northern
539 Apennines (Italy): Forest Exploitation by Past Societies in Mountain Areas. *Interdisciplinaria*
540 *Archaeologica - Natural Sciences in Archaeology* IX, 169–178. doi:10.24916/iansa.2018.2.4
- 541 Bergès, L., Dupouey, J., 2020. Historical ecology and ancient forests: Progress, conservation issues and
542 scientific prospects, with some examples from the French case. *Journal of Vegetation Science*
543 jvs.12846. doi:10.1111/jvs.12846
- 544 Bergès, L., Dupouey, J.-L., 2017. Écologie historique et ancienneté de l'état boisé : concepts, avancées et
545 perspectives de la recherche. *Revue Forestière Française* 297. doi:10.4267/2042/67863
- 546 Bielenin, K., 1992. Le combustible de la sidérurgie antique des Monts de Saint-Croix en Pologne. In: In J.-P
547 Métailié (Éd.), *Proto Industries et Histoire Des Forêts*, Les Cahiers de l'Isard, p.317-329.
- 548 Biringuccio, V., 1556. *La pyrotechnie, ou Art du feu : contenant dix livres, auxquels est amplement traité de*
549 *toutes sortes & diversité de minières, fusions & séparations des métaux, des formes & moules pour*
550 *getter artilleries, cloches & toutes autres figures.* chez Claude Fremy, à l'enseigne saint Martin, rue
551 saint Jacques.
- 552 Blondel, F., Cabanis, M., Girardclos, O., Paradis-Grenouillet, S., 2018. Impact of carbonization on growth rings:
553 Dating by dendrochronology experiments on oak charcoals collected from archaeological sites.
554 *Quaternary International* 463, 268–281. doi:10.1016/j.quaint.2017.03.030
- 555 Bond, J., 2007. Medieval Charcoal-burning in England. In: Klápšte, J., Sommer, P. (Eds.), *Arts and Crafts in*
556 *Medieval Rural Environment*. Brepols Publishers, Turnhout, pp. 277–294. doi:10.1484/M.RURALIA-

- 558 Bonhôte, J., 1998. Forges et forêts dans les Pyrénées ariégeoises : pour une histoire de l'environnement.
559 Pyrégraph. Aspet. 337p.
- 560 Bonhôte, J., Fruhauf, C., 1990a. La métallurgie au bois et les espaces forestiers dans les Pyrénées de l'Aude et
561 de l'Ariège. In: Forges et Forêts. Recherches Sur La Consommation Proto-Industrielle de Bois (Sous
562 La Direction de D. Woronoff), Recherches d'histoire et de Sciences Sociales. Ecole des Hautes Etudes
563 Sciences Sociales, Paris, France, pp. 151–212.
- 564 Bonhôte, J., Fruhauf, C., 1990b. Métallurgie au bois et espaces forestiers pyrénéens dans l'Aude et l'Ariège.
565 Revue géographique des Pyrénées et du Sud-Ouest. Sud-Ouest Européen 61, 459–474.
566 doi:10.3406/rgpso.1990.3225
- 567 Bonhôte, J., Davasse, B., Dubois, C., Izard, V., Métailié, J.-P., 2002. Charcoal kilns and environmental history
568 in the eastern Pyrenees (France). A methodological approach. Second international meeting of
569 anthracology, Oct 2000, Paris, France 219–228.
- 570 Bouget, C., Brin, A., Brustel, H., 2011. Exploring the “last biotic frontier”: Are temperate forest canopies
571 special for saproxylic beetles? Forest Ecology and Management 261, 211–220.
572 doi:10.1016/j.foreco.2010.10.007
- 573 Bouget, C., Larrieu, L., Brin, A., 2014. Key features for saproxylic beetle diversity derived from rapid habitat
574 assessment in temperate forests. Ecological Indicators 36, 656–664. doi:10.1016/j.ecolind.2013.09.031
- 575 Bouget, C., Brustel, H., Noblecourt, T., Zagatti, P., 2019. Les coléoptères saproxyliques de France: catalogue
576 écologique illustré. Muséum national d'Histoire naturelle, Paris, 744p. (Patrimoines naturels ; 79).
- 577 Buckley, P., Mills, J.E., 2015. Coppice silviculture: from the mesolithic to the 21st century. In: Kirby, K. J.;
578 Watkins, C. (Eds), “Europe’s Changing Woods and Forests. From Wildwood to Managed Landscapes”,
579 CABI, Oxfordshire, Boston, 77-92. pp. 77–92.
- 580 Bürgi, M., Bieling, C., Hackwitz, K. von, Kizos, T., Lieskovský, J., Martín, M.G., McCarthy, S., Müller, M.,
581 Palang, H., Plieninger, T., Printsmann, A., 2017. Processes and driving forces in changing cultural
582 landscapes across Europe. Landscape Ecology 32, 2097–2112. doi:10.1007/s10980-017-0513-z
- 583 Burley, J., Youngquist, J., Evans, J., 2004. Encyclopedia of forest sciences. Elsevier, Oxford.
- 584 Burri, S., 2008. Enquête ethnoarchéologique sur le mode de vie et le savoir-faire des derniers charbonniers de
585 Calabre (Italie) : entre tradition et modernité. Presented at the Cultures, Economies, Sociétés et
586 Environnement du début de la Préhistoire au Moyen-Age : travaux en cours, Actes de la troisième table
587 ronde des jeunes chercheurs en archéologie de la MMSH, Aix-en-Provence, 6 juin 2008, A. Boutet, C.
588 Defrasne, T. Lachenal (dir.), <http://www.mmsh.univ-aix.fr/ecoledoctorale/trjca/burri.htm>.
- 589 Burri, S., 2012. Vivre de l'inculte, vivre dans l'inculte en Basse Provence centrale à la fin du Moyen Âge :
590 Histoire, archéologie et ethnoarchéologie d'un mode de vie itinérant (These de doctorat). Aix-
591 Marseille.
- 592 Burri, S., Durand, A., Alilou, M., 2010. Way of Life and Technical Knowledge of Moroccan Charcoal Burners:
593 an Ethnoarchaeological Approach (Gharb and Middle Atlas Morocco), Lugli F., Stoppiello A. et
594 Biagetti S., Ethnoarchaeology Current Research and Field Methods Rome 13–14 May 2010 BAR
595 British Archaeological Reports 2472, 2013, 123-128.
- 596 Cantelaube, J., 2005. La forge à la catalane dans les Pyrénées ariégeoises: une industrie à la montagne, XVIIe-

597 XIXe siècle. CNRS, Université de Toulouse-Le Mirail, Toulouse. 805p.

598 Cateau, E., Larrieu, L., Vallauri, D., Savoie, J.-M., Touroult, J., Brustel, H., 2015. Ancienneté et maturité : deux
599 qualités complémentaires d'un écosystème forestier. *Comptes Rendus Biologies* 338, 58–73.
600 doi:10.1016/j.crv.2014.10.004

601 Cevasco, R., Moreno, D., 2015. Historical Ecology in Modern Conservation in Italy. In: Kirby, K.J. and
602 Watkins, Ch. (Eds). *Europe's Changing Woods and Forests. From Wildwood to Managed Landscapes*,
603 Cabi, Boston, Oxfordshire, 227-242. pp. 227–242.

604 Chassinat, E., 1892. Forêt domaniale de Suc-et-Sentenac, aménagement. (Inventaire - chap IV - n°33. Archives
605 départementales de l'Ariège. Cote archive: 7P1081. 46p).

606 Chrzavzez, J., 2013. Approche expérimentale de la conservation des charbons de bois dans les gisements
607 paléolithiques : processus post-dépositionnels, fragmentation et représentativité des assemblages
608 anthracologiques (These de doctorat). Nice.

609 Commarmot, B., Bachofen, H., Bundziak, Y., Bürgi, A., Ramp, B., Shparyk, Y., Sukhariuk, D., Viter, R., Zingg,
610 A., 2005. Structure of virgin and managed beech forests in Uholka (Ukraine) and Sihlwald
611 (Switzerland): a comparative study. *Forest Snow and Landscape Research* 45–56.

612 Coppini, M., Hermanin, L., 2007. Restoration of selective beech coppices: A case study in the Apennines
613 (Italy). *Forest Ecology and Management, Traditional knowledge, cultural heritage and sustainable*
614 *forest management* 249, 18–27. doi:10.1016/j.foreco.2007.04.035

615 Crumley, C.L. (Ed.), 1994. *Historical ecology: cultural knowledge and changing landscapes*, 1st ed. ed, School
616 of American Research advanced seminar series. School of American Research Press ; Distributed by
617 the University of Washington Press, Santa Fe, N.M. : [Seattle].

618 Davasé, B., 2000. Forêts, charbonniers et paysans dans les Pyrénées de l'est du moyen âge à nos jours: une
619 approche géographique de l'histoire de l'environnement. Geode. Toulouse. 287p.

620 Davasé, B., Galop, D., 1990. Le paysage forestier du Haut Vicdessos (Ariège) : l'évolution d'un milieu
621 anthropisé.

622 Deforce, K., Boeren, I., Adriaenssens, S., Bastiaens, J., De Keersmaecker, L., Haneca, K., Tys, D.,
623 Vandekerckhove, K., 2013. Selective woodland exploitation for charcoal production. A detailed analysis
624 of charcoal kiln remains (ca. 1300–1900 AD) from Zoersel (northern Belgium). *Journal of*
625 *Archaeological Science* 40, 681–689. doi:10.1016/j.jas.2012.07.009

626 Della Fratta Montalbano, M.A., 1678. *Pratica minerale. Trattato del March*. Bologna.

627 Diedhiou, A.G., Dupouey, J.-L., Buee, M., Dambrine, E., Laure, L., Garbaye, J., 2009. Response of
628 ectomycorrhizal communities to past Roman occupation in an oak forest. *Soil Biology and*
629 *Biochemistry* 41, 2206–2213.

630 Dromart, E., 1880. *Traité de la carbonisation des bois en forêts*, E. Lacroix (Paris). ed, Collection : Bibliothèque
631 des professions industrielles et agricoles ; n° 17.

632 Dufraisie, A., Coubray, S., 2018. De l'arbre à la forêt domestiquée : apport de l'anthraco-typologie. *Les*
633 *Nouvelles de l'archéologie* 54–61. doi:10.4000/nda.4533

634 Duhamel Du Monceau, H.-L., 1761. *Art du charbonnier, ou Manière de faire le charbon de bois. Description des*
635 *arts et métiers, faits ou approuvés par Messieurs de l'Académie royale des sciences*. Chez Desaint et
636 Saillant (ed.), Librairie, rue Saint Jean de Beauvais. p69.

637 Dupin, A., 2018. Caractérisation du charbonnage moderne et contemporain bisontin (Franche-Comté, France) et
 638 de son impact sur les peuplements forestiers : le cas de la forêt de Chailluz (Thèse de doctorat).
 639 Bourgogne Franche-Comté.

640 Dupin, A., Sordoillet, D., Fréville, K., Girardclos, O., Gauthier, E., 2019. The taphonomic characterization of a
 641 charcoal production platform. Contribution of an innovative pair of methods: Raman analysis and
 642 micromorphology. *Journal of Archaeological Science* 107, 87–99. doi:10.1016/j.jas.2019.05.003

643 Dupouey, J.-L., Sciama, D., Dambrine, É., Rameau, J.-C., Koerner, W., 2002. La Végétation des forêts
 644 anciennes. *Revue Forestière Française* 521. doi:10.4267/2042/4940

645 Durand, A., Duval, S., Vaschalde, C., 2010. Le charbonnage des Ericacées méditerranéennes : approches
 646 croisées archéologiques, anthracologiques et historiques. Editions ADPCA.

647 Feiss, T., Horen, H., Brasseur, B., Buridant, J., Gallet-Moron, E., Decocq, G., 2017. Historical ecology of
 648 lowland forests: Does pedoanthracology support historical and archaeological data? *Quaternary*
 649 *International* 457, 99–112. doi:10.1016/j.quaint.2016.10.029

650 Fouédjeu, L., Saulnier, M., Lejay, M., Dušátko, M., Labbas, V., Jump, A.S., Burri, S., Buscaino, S., Py-
 651 Saragaglia, V., 2021. High resolution reconstruction of modern charcoal production kilns: An
 652 integrated approach combining dendrochronology, micromorphology and anthracology in the French
 653 Pyrenees. *Quaternary International, Anthracology: Charcoal Science in Archaeology and*
 654 *Palaeoecology* 593–594, 306–319. doi:10.1016/j.quaint.2020.11.033

655 Fregni, C., Maccaferri, D., 2010. Come eravamo, “I Carbonai”, ricostruzione storica di una carbonaia [CD-
 656 ROM]. Appennino Cinema festival, CAI Gruppo Regionale Emilia-Romagna.

657 Fruhauf, C., 1980. Forêt et société : de la forêt paysanne à la forêt capitaliste en pays de Sault sous l’Ancien
 658 régime (vers 1670-1791).

659 Fruhauf, C., 1990. Les conséquences des aménagements des XVIIe et XVIIIe siècles sur l’aspect actuel de la
 660 forêt pyrénéenne. *Annales du Midi* 102, 587–598. doi:10.3406/anami.1990.2268

661 Gebhardt, A., 2007. Impact of charcoal production activities on soil profiles: the micromorphological point of
 662 view. *ArcheoSciences. Revue d’archéométrie* 127–136. doi:10.4000/archeosciences.833

663 Glais, P., 2007. Charbonniers de Brocéliande: l’art de la fouée. Les Amis de la Bibliothèque de Paimpont,
 664 Paimpont (1 Esplanade de Brocéliande, 35380).

665 Gosselin, F., Larrieu, L., 2020. Developing and using statistical tools to estimate observer effect for ordered
 666 class data: The case of the IBP (Index of Biodiversity Potential). *Ecological Indicators* 110, 105884.
 667 doi:10.1016/j.ecolind.2019.105884

668 Gossner, M.M., Lachat, T., Brunet, J., Isacsson, G., Bouget, C., Brustel, H., Brandl, R., Weisser, W.W., Müller,
 669 J., 2013. Current Near-to-Nature Forest Management Effects on Functional Trait Composition of
 670 Saproxyllic Beetles in Beech Forests: Functional Diversity of Beetles. *Conservation Biology* 27, 605–
 671 614. doi:10.1111/cobi.12023

672 Goux, N., 2011. Gestion forestière et biodiversité, les enjeux de conservation d’une espèce parapluie :
 673 *Limoniscus violaceus* (Coleoptera) (phdthesis). Université Pierre et Marie Curie - Paris VI.

674 Hanus, P., 2007. Urbain des bois, l’apprenti charbonnier. Les Cahiers de Framespa. doi:10.4000/framespa.440

675 Hanus, P., 2018. Charbonnier, fais fumer! Chronique ethnographique d’une fête du Vercors. In: Charbonnage,
 676 Charbonniers, Charbonnières. Confluence de Regards Autour d’un Artisanat Méconnu (Paradis-

- Grenouillet, S., Burri, S., Rouaud, R., Eds.). pp. 49–60.
- Henry, A., Théry-Parisot, I., 2014. From Evenk campfires to prehistoric hearths: charcoal analysis as a tool for identifying the use of rotten wood as fuel. *Journal of Archaeological Science* 52, 321–336. doi:10.1016/j.jas.2014.09.005
- Hermý, M., Honnay, O., Firbank, L., Grashof-Bokdam, C., Lawesson, J.E., 1999. An ecological comparison between ancient and other forest plant species of Europe, and the implications for forest conservation. *Biological Conservation* 91, 9–22. doi:10.1016/S0006-3207(99)00045-2
- Hillebrecht, M.-L., 1982. Die Relikte der Holzkohlewirtschaft als Indikatoren für Waldnutzung und Waldentwicklung: Untersuchungen an Beispielen aus Südniedersachsen, Göttinger geographische Abhandlungen. Goltze, Göttingen.
- Huffel, G., 1926. Les Méthodes de l'aménagement forestier en France. Étude historique.
- Jacquemin, F., Kervyn, T., Branquart, E., Delahaye, L., Dufrêne, M., Claessens, H., 2014. Les forêts anciennes en Wallonie. 1ère partie : Concepts généraux. *Forêt.Nature*.
- Kirby, K.J., Watkins, C. (Eds.), 1998. The ecological history of European forests. Presented at the International Conference on Advances in Forest and Woodland History, CAB International, Oxon, UK ; New York.
- Lapassat, R., 1983. L'industrie du fer dans les Pyrénées Orientales et Ariégeoises au XIXème siècle, Editions du Castillet. ed, Groupe de recherches historiques et archéologiques du Conflent, Prades. Perpignan.
- Larrieu, L., Gonin, P., 2008. L'indice de biodiversité potentielle (ibp) : une méthode simple et rapide pour évaluer la biodiversité potentielle des peuplements forestiers. *Revue Forestière Française*. doi:10.4267/2042/28373
- Larrieu, L., Gonin, P., Deconchat, M., 2012. Le domaine d'application de l'indice de biodiversité potentielle (IBP). *Revue Forestière Française*. doi:10.4267/2042/50657
- Larrieu, L., Cabanettes, A., Gonin, P., Lachat, T., Paillet, Y., Winter, S., Bouget, C., Deconchat, M., 2014a. Deadwood and tree microhabitat dynamics in unharvested temperate mountain mixed forests: A life-cycle approach to biodiversity monitoring. *Forest Ecology and Management* 334, 163–173. doi:10.1016/j.foreco.2014.09.007
- Larrieu, L., Cabanettes, A., Brin, A., Bouget, C., Deconchat, M., 2014b. Tree microhabitats at the stand scale in montane beech–fir forests: practical information for taxa conservation in forestry. *European Journal of Forest Research* 133, 355–367. doi:10.1007/s10342-013-0767-1
- Larrieu, L., Cabanettes, A., Goux, N., Burnel, L., Bouget, C., Deconchat, M., 2017. Development over time of the tree-related microhabitat profile: the case of lowland beech–oak coppice-with-standards set-aside stands in France. *European Journal of Forest Research* 136, 37–49. doi:10.1007/s10342-016-1006-3
- Larrieu, L., Bouget, C., Courbaud, B., Goux, N., Goulard, M., Lachat, T., Stillhard, J., 2018. Spatial patterns of tree-related microhabitats: key factors and ecological significance for the conservation of the associated biodiversity. In: *Proceedings of the 5th European Congress of Conservation Biology*. Presented at the 5th European Congress of Conservation Biology, Jyväskylä University Open Science Centre, Jyväskylä, Finland. doi:10.17011/conference/eccb2018/107340
- Larrieu, L., Gosselin, F., Archaux, F., Chevalier, R., Corriol, G., Dauffy-Richard, E., Deconchat, M., Gosselin, M., Ladet, S., Savoie, J.-M., Tillon, L., Bouget, C., 2019a. Assessing the potential of routine stand variables from multi-taxon data as habitat surrogates in European temperate forests. *Ecological*

717 Indicators 104, 116–126. doi:10.1016/j.ecolind.2019.04.085

718 Larrieu, L., Cabanettes, A., Gouix, N., Burnel, L., Bouget, C., Deconchat, M., 2019b. Post-harvesting dynamics
719 of the deadwood profile: the case of lowland beech-oak coppice-with-standards set-aside stands in
720 France. *European Journal of Forest Research* 138, 239–251. doi:10.1007/s10342-019-01164-8

721 Larsson, T.-B. (Ed.), 2001. Biodiversity evaluation tools for European forests, *Ecological bulletins*. Enckell
722 [u.a.], Lund.

723 Ludemann, T., Nelle, O., 2002. Die Wälder am Schauinsland und ihre Nutzung durch Bergbau und Köhlerei,
724 Schriftenreihe Freiburger forstliche Forschung. Forstliche Versuchs- und Forschungsanst. Baden-
725 Württemberg, Freiburg (Breisgau).

726 Lugli, F., Pracchia, S., 1995. Modelli e finalità nello studio della produzione di carbone di legna in archeologia,
727 Origini, XVIII, Roma, 1995, pp. 425-479.

728 Maillat, J., 1942. La carbonisation dans la Haute-Garonne. *Revue géographique des Pyrénées et du Sud-Ouest*
729 13, 265–276. doi:10.3406/rgps.1942.1177

730 Malepeyre, A.F., 1836. *Maison rustique du XIXe siècle: Arts agricoles*. 1836. Agriculture forestière, législation
731 et administration rurale. Bureau [du Journal d’agriculture pratique].

732 Máliš, F., Bobek, P., Hédli, R., Chudomelová, M., Petřík, P., Ujházy, K., Ujházyová, M., Kopecký, M., 2021.
733 Historical charcoal burning and coppicing suppressed beech and increased forest vegetation
734 heterogeneity. *Journal of Vegetation Science* 32, e12923. doi:https://doi.org/10.1111/jvs.12923

735 Marié de L’Isle, E.-B., 1835. *Manuel complet des marchands de bois et de charbons, ou Traité de ce commerce*
736 en général. La librairie encyclopédique de Roret, Paris.

737 Montanari, C., Prono, P., Scipioni, S., 2000. The study of charcoal-burning sites in the Apennine Mountains of
738 Liguria (NW Italy) as a tool for forest history. In: Agnoletti, M., Anderson, S. (Eds.), *Methods and*
739 *Approaches in Forest History. Papers Selected from a Conference Held in Florence, Italy in 1998*.
740 CABI, Wallingford, pp. 79–91. doi:10.1079/9780851994208.0079

741 Montanari, C.A., Scipioni, S., Calderoni, G., Leonardi, G., Moreno, D.T., 2002. Linking anthracology and
742 historical ecology: suggestion from a post-medieval site in the Ligurian Apennines (north-west Italy).
743 In: Thiébault, S. (Ed.), *Charcoal Analysis. Methodological Approaches, Palaeoecological Results and*
744 *Wood Uses*. *British Archaeological Reports*, Vol. 1063. Archaeopress, Oxford. pp. 209–217.

745 Moreno, D., Cevasco, R., Pescini, V., Gabellieri, N., 2019. The Archeology of Woodland Ecology:
746 Reconstructing Past Woodmanship Practices of Wooded Pasture Systems in Italy. In: Allende Álvarez,
747 F., Gomez-Mediavilla, G., López-Estébanez, N. (Eds.), *Silvicultures - Management and Conservation*.
748 IntechOpen. doi:10.5772/intechopen.86101

749 Moskal-del Hoyo, M., Wachowiak, M., Blanchette, R.A., 2010. Preservation of fungi in archaeological
750 charcoal. *Journal of Archaeological Science* 37, 2106–2116. doi:10.1016/j.jas.2010.02.007

751 Nelle, O., 2002. Charcoal burning remains and forest stand structure - Examples from the Black Forest (south-
752 west Germany) and the Bavarian Forest (south-east Germany). In: Thiébault, S. (Ed.), *Charcoal*
753 *Analysis, Methodological Approaches, Palaeoecological Studies and Wood Uses, Proceeding of the*
754 *Second International Meeting of Anthracology (Paris, September 2000)*, Oxford, Archaeopress: 201-
755 207 (BAR International Series 1063). pp. 201–207.

756 Nelle, O., Robin, V., Talon, B., 2013. *Pedoanthracology: Analysing soil charcoal to study Holocene*

757 palaeoenvironments. *Quaternary International, Pedoanthracology and Environmental Studies* 289, 1–4.
758 doi:10.1016/j.quaint.2012.11.024

759 Nicolescu, V.-N., Carvalho, J., Hochbichler, E., Bruckman, V.J., Piqué, M., Hernea, C., Viana, H., Štochlová,
760 P., Ertekin, M., Đodan, M., Dubravac, T., Vandekerckhove, K., Kofman, P.D., Rossney, D., Unrau, A.,
761 2018. Silvicultural Guidelines for European Coppice Forests. In: A. Unrau, G. Becker, R. Spinelli, D.
762 Lazdina, N. Magagnotti, V.N. Nicolescu, P. Buckley, D. Bartlett, P.D. Kofman (Eds.), *Coppice Forests*
763 *in Europe* (Pp. 46-63). Freiburg i. Br., Germany: Albert Ludwig University of Freiburg. COST Action
764 PF1301 EuroCoppice. Albert-Ludwigs-Universität Freiburg, Freiburg, pp. 46–63.

765 Nocentini, S., 2009. Structure and management of beech (*Fagus sylvatica* L.) forests in Italy. *iForest -*
766 *Biogeosciences and Forestry* 2, 105–113. Doi:10.3832/ifer0499-002

767 Noiriel, 1965. Projet de révision d'aménagement (1966-1980) de la forêt de Suc-et-Sentenac (Rapport de
768 présentation. *Eaux et forêts - serie 12, n°28 - (double)*. Archive départementale de l'Ariège. Cote:
769 J108.227. 56p).

770 Paillet, Y., Pernot, C., Boulanger, V., Debaive, N., Fuhr, M., Gilg, O., Gosselin, F., 2015. Quantifying the
771 recovery of old-growth attributes in forest reserves: A first reference for France. *Forest Ecology and*
772 *Management* 346, 51–64. doi:10.1016/j.foreco.2015.02.037

773 Paradis-Grenouillet, S., 2012. Étudier les forêts métallurgiques : analyses dendro-anthracologiques et approches
774 géohistoriques : exemple des forêts du mont Lozère et du Périgord-Limousin (Thèse de doctorat).
775 Limoges. 452p.

776 Paradis-Grenouillet, S., Allée, P., Vives, G.S., Ploquin, A., 2015. Sustainable management of metallurgical
777 forest on Mont Lozère (France) during the Early Middle Ages. *Environmental Archaeology* 20, 168–
778 183. doi:10.1179/1749631414Y.00000000050

779 Paradis-Grenouillet, S., Burri, S., Rouaud, R. (Eds.), 2018a. Charbonnage, charbonniers, charbonnières:
780 confluence de regards autour d'un artisanat méconnu. Presses universitaires de Provence, Aix-en-
781 Provence. p236.

782 Paradis-Grenouillet, S., Rouaud, R., Allée, P., 2018b. Retrouver le diamètre des bois charbonnés, validation de
783 la chaîne opératoire dendro-anthracologique par l'étude de trois charbonnières expérimentales. In:
784 Paradis-Grenouillet, S., Burri, S., Rouaud, R. (Eds.), *Charbonnage, Charbonniers, Charbonnières.*
785 *Confluence de Regards Autour d'un Artisanat Méconnu*. pp. 105–120.

786 Percy, J., 1864. *Traité complet de métallurgie*.

787 Peterken, G.F., 1993. *Woodland conservation and management*, 2nd ed. ed. Chapman & Hall, London ; New
788 York.

789 Poggi, G., Métailié, J.-P., 1998. Carbonai dell'Antola, video-cassetta 40', co-produzione CAV-UTM/Ente Parco
790 Antola, realizzazione Bruno Bastard. [https://www.canal-](https://www.canal-u.tv/video/universite_toulouse_ii_le_mirail/les_charbonniers_de_l_antola_jean_paul_metailie_guiseppi_na_poggi.4327)
791 [u.tv/video/universite_toulouse_ii_le_mirail/les_charbonniers_de_l_antola_jean_paul_metailie_guiseppi](https://www.canal-u.tv/video/universite_toulouse_ii_le_mirail/les_charbonniers_de_l_antola_jean_paul_metailie_guiseppi_na_poggi.4327)
792 [na_poggi.4327](https://www.canal-u.tv/video/universite_toulouse_ii_le_mirail/les_charbonniers_de_l_antola_jean_paul_metailie_guiseppi_na_poggi.4327).

793 Pott, R., Freund, H., Speier, M., 1992. Anthropogenic changes of landscape by extensive woodland management
794 and charcoal production in Siegerland (Northrhine-Westphalia, Germany. In: Métailié, J.-P. (Ed.),
795 *Protoindustries et Histoire Des Forêts, Les Cahiers de l'Isard*, p.163-183.

796 Py, V., Durand, A., Ancel, B., 2013. Anthracological analysis of fuel wood used for firesetting in medieval

797 metallic mines of the Faravel district (southern French Alps). *Journal of Archaeological Science* 40,
798 3878–3889. doi:10.1016/j.jas.2013.05.006

799 Py, V., Durand, A., Mocci, F., Walsh, K., 2015. Temps de la feuille, temps du boisillage ? Lire les données
800 anthracologiques au prisme de l'économie alpestre médiévale. In: *Traces Du Végétal*. Presses
801 Universitaires de Rennes, pp. 43–62.

802 Py-Saragaglia, V., Cunill Artigas, R., Métailié, J.-P., Ancel, B., Baron, S., Paradis-Grenouillet, S., Lerigoleur,
803 E., Badache, N., Barcet, H., Galop, D., 2017. Late Holocene history of woodland dynamics and wood
804 use in an ancient mining area of the Pyrenees (Ariège, France). *Quaternary International* 458, 141–157.
805 doi:10.1016/j.quaint.2017.01.012

806 Py-Saragaglia, V., Saulnier, M., Cunill Artigas, R., Fouédjeu, L., Paradis-Grenouillet, S., Buscaino, S., Galop,
807 D., 2018. Chapter 10: Long-term forest evolution and woodland uses in an ancient charcoal-production
808 forest of the eastern French Pyrenees: an interdisciplinary approach at high spatio-temporal resolution.
809 In: Aspe, C., Burri, S., Paradis-Grenouillet, S. (Eds.), *Into the Woods. Overlapping Perspectives on the*
810 *History of Ancient Forests*. Quae.

811 Py-Saragaglia, V., Burri, S., Fouédjeu, L., 2019. Les forêts montagnardes du versant nord des Pyrénées. In:
812 Bépoix, S., Richard, H. (Eds.), *La Forêt Au Moyen Âge*. Paris, France, pp. 276–299.

813 Raab, A., Takla, M., Raab, T., Nicolay, A., Schneider, A., Rösler, H., Heußner, K.-U., Bönisch, E., 2015. Pre-
814 industrial charcoal production in Lower Lusatia (Brandenburg, Germany): Detection and evaluation of
815 a large charcoal-burning field by combining archaeological studies, GIS-based analyses of shaded-
816 relief maps and dendrochronological age determination. *Quaternary International* 367, 111–122.
817 doi:10.1016/j.quaint.2014.09.041

818 Rackham, O., 2003. *Ancient woodland: its history, vegetation and uses in England*. Castlepoint, Dalbeattie,
819 Kirkcubrightshire.

820 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Ramsey, C.B., Butzin, M., Cheng, H.,
821 Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G.,
822 Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., Plicht, J. van der,
823 Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F.,
824 Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S.,
825 Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern
826 Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62, 725–757.
827 doi:10.1017/RDC.2020.41

828 Rochebrunel, 1920. Forêt domaniale de Suc-et-Sentenac, révision d'aménagement, service des aménagements et
829 du reboisement (Inventaire chpa IV n°77. Archive départementale de l'Ariège. 26p).

830 Rose, F., 1993. British Wildlife | Ancient British Woodlands and their Epiphytes. *Br. Wildl.* 5 83–93.

831 Rostain, S., Saulieu, G. de, 2018. *Ecologie historique*, Première édition. ed. Éditions de la Maison des sciences
832 de l'homme, Paris.

833 Ryvarden, L., Melo, I., 2014. Poroid fungi of Europe, Oslo : Fungiflora. ed, *Synopsis fungorum*, 31. 455p.

834 Salvador, J., 1930. Simples notes sur l'aménagement et l'exploitation des forêts pyrénéennes françaises. *Revue*
835 *géographique des Pyrénées et du Sud-Ouest* 1, 58–74. doi:10.3406/rgps.1930.3944

836 Samojlik, T., Rotherham, I.D., Jedrzejewska, B., 2013. Quantifying Historic Human Impacts on Forest

837 Environments: A Case Study in Bialowieza Forest, Poland. *Environmental History* 18, 576–602.
838 doi:10.1093/envhis/emt039

839 Saulnier, M., Cunill Artigas, R., Fouédjeu, L., Buscaino, S., Métailié, J.-P., Galop, D., Py-Saragaglia, V., 2019.
840 A study of late Holocene local vegetation dynamics and responses to land use changes in an ancient
841 charcoal making woodland in the central Pyrenees (Ariège, France), using pedoanthracology.
842 *Vegetation History and Archaeobotany*. doi:10.1007/s00334-019-00740-7

843 Schmidt, M., Mölder, A., Schönfelder, E., Engel, F., Fortmann-Valtink, W., 2016. Charcoal kiln sites,
844 associated landscape attributes and historic forest conditions: DTM-based investigations in Hesse
845 (Germany). *Forest Ecosystems* 3, 8. doi:10.1186/s40663-016-0067-6

846 Schweingruber, F.H., 1990. *Anatomy of european woods : An atlas for the identification of european trees,*
847 *shrubs and dwarf shrubs*, Haupt, Stuttgart. Birmensdorf:Eidgenössische Anstalt für das forstliche
848 Versuchswesen.

849 Sebastien, L., 2020. The power of place in understanding place attachments and meanings. *Geoforum* 108, 204–
850 216. doi:10.1016/j.geoforum.2019.11.001

851 Stokland, J.N., Siitonen, J., Jonsson, B.G., 2012. *Biodiversity in dead wood*, Ecology, biodiversity, and
852 conservation. Cambridge Univ. Press, Cambridge.

853 Strachan, S., Biondi, F., Lindström, S.G., McQueen, R., Wigand, P.E., 2013. Application of Dendrochronology
854 to Historical Charcoal-Production Sites in the Great Basin, United States. *Historical Archaeology* 47,
855 103–119. doi:10.1007/BF03377126

856 Svedelius, G., Anderson, R.B., Nicodemus, W.J.L., 1875. *Handbook for Charcoal Burners*. John Wiley & Son.
857 246p.

858 Szabó, P., 2015. Historical ecology: past, present and future. *Biological Reviews of the Cambridge*
859 *Philosophical Society* 90, 997–1014. doi:10.1111/brv.12141

860 Théry-Parisot, I., Henry, A., 2012. Seasoned or green? Radial cracks analysis as a method for identifying the use
861 of green wood as fuel in archaeological charcoal. *Journal of Archaeological Science* 39, 381–388.
862 doi:10.1016/j.jas.2011.09.024

863 Théry-Parisot, I., Henry, A., Chravzez, J., 2016. Apport de l'expérimentation à la compréhension des pratiques
864 sociétales en anthracologie : gestion et utilisation du bois de feu dans les sociétés préhistoriques.
865 *Cadernos do Lapaarq* 13, 484–509. doi:10.15210/LEPAARQ.V13I25.7410

866 Théry-Parisot, I.A., 2001. *Économie des combustibles au paléolithique : expérimentation, taphonomie,*
867 *anthracologie*. CNRS. Paris.

868 Tolksdorf, J.F., Elburg, R., Schröder, F., Knapp, H., Herbig, C., Westphal, T., Schneider, B., Fülling, A.,
869 Hemker, C., 2015. Forest exploitation for charcoal production and timber since the 12th century in an
870 intact medieval mining site in the Niederpöbel Valley (Erzgebirge, Eastern Germany). *Journal of*
871 *Archaeological Science: Reports* 4, 487–500. doi:10.1016/j.jasrep.2015.10.018

872 Unrau, A., Becker, G., Spinelli, R., Lazdina, D., Magagnotti, N., Nicolescu, V.-N., Buckley, P., Bartlett, D.,
873 Kofman, P.D. (Eds.), 2018. *Coppice Forests in Europe*, 1. Auflage. ed, COST Action PF1301
874 *EuroCoppice*. Albert-Ludwigs-Universität Freiburg, Freiburg.

875 Valerius, J.-B., 1851. *Traité théorique et pratique de la fabrication de la fonte*. Kissling et Company.

876 Verna, C., 2001. *Le temps des moulins: fer, technique et société dans les Pyrénées centrales (XIIIe-XVIe*

- 877 siècles). Publications de la Sorbonne, 2001, Paris, France.
- 878 Vernet, J.L., Ogereau, P., Figueiral, I., 2001. Guide d'identification des charbons de bois préhistoriques et
879 récents: sud-ouest de l'Europe : France, péninsule ibérique, et îles Canaries. CNRS, Paris.
- 880 Watkins, C., 2015. Methods and approaches in the study of woodland history. In: Kirby, K.J., Watkins, C.
881 (Eds.), Europe's Changing Woods and Forests: From Wildwood to Managed Landscapes. CABI,
882 Wallingford, pp. 18–32. doi:10.1079/9781780643373.0018