



HAL
open science

Antigenic waves of virus–immune coevolution

Jacopo Marchi, Michael Lässig, Aleksandra M Walczak, Thierry Mora

► **To cite this version:**

Jacopo Marchi, Michael Lässig, Aleksandra M Walczak, Thierry Mora. Antigenic waves of virus–immune coevolution. *Proceedings of the National Academy of Sciences of the United States of America*, 2021, 118 (27), pp.e2103398118. 10.1073/pnas.2103398118 . hal-03367761

HAL Id: hal-03367761

<https://hal.science/hal-03367761v1>

Submitted on 8 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Antigenic waves of virus-immune co-evolution

Jacopo Marchi^a, Michael Lässig^b, Aleksandra M. Walczak^{a,1}, and Thierry Mora^{a,1}

^aLaboratoire de physique de l'École normale supérieure, CNRS, PSL University, Sorbonne Université, and Université de Paris, 75005 Paris, France; ^bInstitute of Theoretical Physics, University of Cologne, 50937 Cologne, Germany

The evolution of many microbes and pathogens, including circulating viruses such as seasonal influenza, is driven by immune pressure from the host population. In turn, the immune systems of infected populations get updated, chasing viruses even further away. Quantitatively understanding how these dynamics result in observed patterns of rapid pathogen and immune adaptation is instrumental to epidemiological and evolutionary forecasting. Here we present a mathematical theory of co-evolution between immune systems and viruses in a finite-dimensional antigenic space, which describes the cross-reactivity of viral strains and immune systems primed by previous infections. We show the emergence of an antigenic wave that is pushed forward and canalized by cross-reactivity. We obtain analytical results for shape, speed, and angular diffusion of the wave. In particular, we show that viral-immune co-evolution generates a new emergent timescale, the persistence time of the wave's direction in antigenic space, which can be much longer than the coalescence time of the viral population. We compare these dynamics to the observed antigenic turnover of influenza strains, and we discuss how the dimensionality of antigenic space impacts on the predictability of the evolutionary dynamics. Our results provide a concrete and tractable framework to describe pathogen-host co-evolution.

1. Introduction

The evolution of viral pathogens under the selective pressure of its hosts' immunity is an example of rapid co-evolution. Viruses adapt in the usual Darwinian sense by evading immunity through antigenic mutations, while immune repertoires adapt by creating memory against previously encountered strains. Some mechanisms of in-host immune evolution, such as the affinity maturation process, are important for the rational design of vaccines. Examples are the seasonal human influenza virus, where vaccine strain selection can be informed by predicting viral evolution in response to collective immunity(1), as well as chronic infections such as HIV (2–5), where co-evolution occurs within each host. Because of the relatively short time scales of selection and strain turnover, these dynamics also provide a laboratory for studying evolution and its link to ecology (6).

It is useful to think of both viral strains and immune protections as living in a common antigenic space (6), corresponding to an idealized “shape space” of binding motifs between antibodies and their cognate epitopes (7). While the space of molecular recognition is high-dimensional, projections onto a low-dimensional effective shape space have provided useful descriptions of the antigenic evolution. In the example of influenza, neutralization data from hemagglutination-inhibition assays can be projected onto a two-dimensional antigenic space (8–10). Mapping historical antigenic evolution in this space suggests a co-evolutionary dynamics pushing the virus away from its past positions, where collective immunity has developed. Importantly, the evolution of influenza involves competitive interactions of antigenically distinct clades in the viral

population, generating a “Red Queen” dynamics of pathogen evolution (11, 12). Genomic analysis of influenza data has revealed evolution by clonal interference (13); this mode of evolution is well-known from laboratory microbial populations (14). In addition, the viral population may split into subtypes. Such splitting or “speciation” events, which are marked by a decoupling of the corresponding immune interactions, happened in the evolution of influenza B (15) and of noroviruses (16).

The joint dynamics of viral strains and the immune systems of the host population can be modeled using agent-based simulations (17, 18) that track individual hosts and strains. Such approaches have been used to study the effect of competition on viral genetic diversity (19), to study geographical effects (20), and the effect of vaccination (21). Alternatively, systems of coupled differential equations known as Susceptible-Infected-Recovered (SIR) models may be adapted to incorporate evolutionary mechanisms of antigenic adaptation (6, 22, 23). Agent-based simulations in 2 dimensions were used to recapitulate the ballistic evolution characteristic of influenza A (18), and to predict the occurrence of splitting and extinction events (24). In parallel, theory was developed to study the Red Queen effect (12, 25), based on the well established theory of the traveling fitness wave (26–28). While effectively set in one dimension, this class of models can nonetheless predict extinction and splitting events assuming an infinite antigenic genome (12).

In this work, we propose a co-evolutionary theory in an antigenic interaction space of arbitrary dimension d , which is described by joint non-linear stochastic differential equations coupling the population densities of viruses and of protected hosts. We show that these equations admit a d -dimensional antigenic wave solution, and we study its motion, shape, and

Significance Statement

Viruses, such as influenza, evolve under the selection of host immune systems. Previously infected individuals become immune, forcing the virus to find susceptible hosts or mutate, chasing it away in antigenic space. We formulate this viral escape process in terms of a low-dimensional wave moving in antigenic space. The dimensionality of the antigenic space impacts both the persistence, as well as stability of viral evolution. We uncover a new characteristic timescale for the persistence of the viral strain, which is an order of magnitude longer than individual host immunity, and emerges collectively from the pressure of the chasing immune systems. These results offer intuition about the antigenic turnover of viruses, and highlight the importance of the effective dimensionality of co-evolution.

The authors declare no competing interest.

¹To whom correspondence should be addressed: aleksandra.walczak@phys.ens.fr, thierry.mora@phys.ens.fr

63 stability, using simulations and analytical approximations.
 64 Based on these results, we discuss how canalization and pred-
 65 ctability of antigenic evolution depend on the dimensionality
 66 d .

67 2. Results

68 **A. Coarse-grained model of viral-immune co-evolution.** Our
 69 model describes the joint temporal evolution of populations
 70 of viruses and immune protections in some effective antigenic
 71 space of dimension d . Both viral strains and immune pro-
 72 tections are labeled by their position $\mathbf{x} = (x_1, \dots, x_d)$ (or
 73 “phenotype”) in that common antigenic space (Fig. 1A). In
 74 that space, viruses randomly move as a result of antigenic
 75 mutations and proliferate through infections of new hosts. Im-
 76 mune memories are added at the past positions of viruses.
 77 Immune memories distributed across the host population pro-
 78 vide protection that reduces the effective fitness of the virus.
 79 We coarse-grain that description by summarizing the viral pop-
 80 ulation by a density $n(\mathbf{x}, t)$ of hosts infected by a particular
 81 viral strain \mathbf{x} , and immunity by a density $h(\mathbf{x}, t)$ of immune
 82 memories specific to strain \mathbf{x} in the host population.

83 At each infection cycle, each host may infect R_0 unpro-
 84 tected hosts, where R_0 is called the basic reproduction number.
 85 However, a randomly picked host is susceptible to strain \mathbf{x}
 86 with probability $(1 - c(\mathbf{x}, t))^M$, where $c(\mathbf{x}, t)$ is the coverage
 87 of strain \mathbf{x} by immune memories of the population, and the
 88 number M of immune memories carried by each host. Because
 89 of cross-reactivity, which allows immune memories to confer
 90 protection against closely strains, immune coverage is given
 91 as a function of the density of immune memories:

$$92 \quad c(\mathbf{x}, t) = \frac{1}{M} \int d\mathbf{x}' h(\mathbf{x}', t) H(\mathbf{x} - \mathbf{x}'), \quad [1]$$

93 where $H(\mathbf{x} - \mathbf{x}') = \exp(-|\mathbf{x} - \mathbf{x}'|/r)$ is a cross-reactivity kernel
 94 describing how well memory \mathbf{x}' protects against strain \mathbf{x} , and
 95 r is the range of the coverage provided by cross-reactivity. In
 96 summary, the effective growth rate, or “fitness”, of the virus
 97 is given by $f(\mathbf{x}, t) \equiv \ln[R_0(1 - c(\mathbf{x}, t))^M]$.

The coupled dynamics of viruses and immune memories is
 then described by the stochastic differential equations (with
 time in units of infection cycles throughout):

$$98 \quad \partial_t n(\mathbf{x}, t) = f(\mathbf{x}, t)n(\mathbf{x}, t) + D\partial_{\mathbf{x}}^2 n + \sqrt{n(\mathbf{x}, t)}\eta(\mathbf{x}, t) \quad [2]$$

$$99 \quad \partial_t h(\mathbf{x}, t) = \frac{1}{N_h} \left[n(\mathbf{x}, t) - N(t) \frac{h(\mathbf{x}, t)}{M} \right]. \quad [3]$$

100 Here η is a Gaussian white noise in time and space,
 101 $\langle \eta(\mathbf{x}, t)\eta(\mathbf{x}', t') \rangle = \delta(\mathbf{x} - \mathbf{x}')\delta(t - t')$, accounting for demo-
 102 graphic noise (29). This stochastic term is crucial, as it will
 103 drive the evolution of the wave. The diffusion constant D
 104 describes the effect of infinitesimal mutations on the phenotype,
 105 $D = \mu\langle \delta x_1^2 \rangle / 2$, where μ is the mean number of mutations per
 106 cycle, and $\langle \delta x_1^2 \rangle$ the mean squared effect of each mutation along
 107 each antigenic dimension (assuming that mutations do not
 108 have a systematic bias, $\langle \delta x_1 \rangle = 0$). The continuous-diffusion
 109 assumption implied by Eq. 2 is only valid when there are
 110 many small mutation effects, $\mu \gg 1$ and $\delta x \ll r$, in contrast
 111 with regimes where mutations are rare but have a substantial
 fitness effect drawn from a distribution (25, 30). Our choice
 is simpler in that it describes the mutation process through




Fig. 1. **A simple model of viral-host co-evolution predicts the emergence of an antigenic wave.** **A.** Schematic of the co-evolution model. Viruses proliferate while effectively diffusing in antigenic space (here in 2 dimensions) through mutations, with coefficient D . Past virus positions are replaced by immune protections (light blue). Immune protections create a fitness gradient for the viruses (green gradient) favoring strains at the front. Both populations of viruses and immune populations are coarse-grained into densities in antigenic space. **B.** Snapshot of a numerical simulation of Eq. 2-3 showing the existence of a wave solution. The blue colormap represents the density of immune protections $h(\mathbf{x}, t)$ left behind by past viral strains. The current virus density $n(x)$ is shown in red. **C.** Close-up onto the viral population, showing fitness isolines. The wave moves in the direction of the fitness gradient (arrow) through the enhanced growth of stains at the edge of the wave (black dots). **D.** Distribution of fitness across the viral population (corresponding to the projection of B. along the fitness gradient). Parameters for B-D: $D/r^2 = 3 \cdot 10^{-9}$, $N_h = 10^8$, $\ln R_0 = 3$, $M = 1$.

a single parameter D . Along with the choice of the cross-reactivity kernel H , it also naturally preserves the isotropy of the antigenic space.

The total viral population size, or number of infected hosts, $N(t) = \int d\mathbf{x} n(\mathbf{x}, t)$ is subject to fluctuations. At the same time, the host population size N_h , remains constant because newly added memories (first term of right-hand side of Eq. 3) overwrite existing ones picked uniformly at random (second term of r.h.s. of Eq. 3). Since each host carries M immune receptors, we have $\int d\mathbf{x} h(\mathbf{x}, t) = M$.

If we assume that the system reaches an evolutionary steady state, with stable viral population size $N(t) = N$, then Eq. 3 can be integrated explicitly:

$$125 \quad h(\mathbf{x}, t) = \frac{M}{N} \int_{-\infty}^t \frac{dt'}{\tau} e^{-\frac{t-t'}{\tau}} n(\mathbf{x}, t'), \quad [4]$$

with $\tau = MN_h/N$. This equation shows how the density of protections reflects the past evolution of the viral population.

B. Antigenic waves. We simulated Eq. (2)-Eq. (3) on a square lattice (Methods) and found a stable wave solution (Fig. 1B-D). The wave has a stable population size N , and moves approximately ballistically through antigenic space, pushed from behind by the immune memories left in the trail of past viral strains (Fig. 1B). These memories exert an immune pressure on the viruses, forming a fitness gradient across the width of the wave (Fig. 1C), favoring the few strains that are furthest from immune memories, at the edge of the wave.

137 We assume that the solution of the coupled evolution equa- 176
 138 tions Eq. (2)-Eq. (3) takes the form of a moving quasispecies 177
 139 in a d -dimensional antigenic space, 178

$$140 \quad n(\mathbf{x}, t) = \frac{N}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(x_1 - vt)^2}{\sigma^2} \right] \rho(x_2, \dots, x_d). \quad [5]$$

141 Here, we have written the solution in a co-moving frame, in 179
 142 which a motion with constant speed v takes place in the 180
 143 direction of the coordinate x_1 , and fluctuations in the other 181
 144 dimensions, $\rho(x_2, \dots, x_d, t)$, centered around $x_i = 0$ for $i > 1$,
 145 are assumed to be independent. In the next sections, we will
 146 analyse solutions of this form. First, we will project the d -
 147 dimensional antigenic wave onto the one-dimensional fitness
 148 space; this projection produces a travelling fitness wave (26–
 149 28, 31, 32) that determines the antigenic speed v and the mean
 150 pair coalescence time $\langle T_2 \rangle$ of the viral genealogy. Second, we
 151 will study the shape of the d -dimensional quasispecies and
 152 determine the fluctuations in the transverse directions. These
 153 fluctuations produce a key result of this paper: immune in-
 154 teractions canalize the evolution of the antigenic wave; this
 155 constraint can be quantified by characteristic time scales gov-
 156 erning the transverse antigenic fluctuations. Canalization is
 157 most pronounced in spaces of low dimensionality d and, as we
 158 discuss below, affects the predictability of antigenic evolution.

159 **C. Speed of antigenic evolution.** Projected onto the fitness
 160 axis $f = f(\mathbf{x}, t)$, the solution is approximately Gaussian
 161 (Fig. 1D). This representation suggests a strong similarity to
 162 the fitness wave solution found in models of rapidly adapting
 163 populations with an infinite reservoir of beneficial mutations
 164 (26–28, 31, 32). To make the analogy rigorous, we must assume
 165 that the fitness gradient in antigenic space is approximately
 166 constant, meaning that fitness isolines are straight and equidis-
 167 tant. Mutations along the gradient direction have a fitness
 168 effect that is linear in the displacement, while mutations along
 169 perpendicular directions are neutral and can be treated inde-
 170 pendently. Note that while we will use this projection onto
 171 fitness to compute the speed of the antigenic wave, the under-
 172 lying antigenic wave remains in d dimensions; we will come
 173 back to transverse fluctuations in the next sections.

There are several models of fitness waves that differ in
 the assumptions on the statistics of mutational effects. Our
 assumption of diffusive motion makes our projected dynamics
 equivalent to that studied in ref. (32), which itself builds on
 earlier work (27). This equivalence results from the two key
 assumptions of the mutation model in antigenic space: muta-
 tions have a small effect, and their distribution is isotropic,
 meaning that there are as many deleterious as beneficial muta-
 tions. In the limit where the wave is small compared to the
 adaptation time scale, $v\tau \gg \sigma$, the wave may be replaced by
 a Dirac delta function at $\mathbf{x} = (vt, 0, \dots, 0)$ in Eq. 4. One can
 then calculate explicitly the immune density (upstream of the
 wave) and coverage (downstream of the wave, using Eq. 1):

$$174 \quad h(\mathbf{x}, t) \approx \frac{M}{v\tau} e^{-\frac{vt-x_1}{v\tau}} \Theta(vt - x_1) \delta(x_2) \cdots \delta(x_d), \quad [6]$$

$$175 \quad c(\mathbf{x}, t) \approx \frac{e^{-(x_1-vt)/r}}{1+v\tau/r}, \quad x_1 \geq vt, \quad x_{i>1} \ll r \quad [7]$$

where $\Theta(x) = 1$ for $x \geq 0$ and 0 otherwise. This idealized
 exponential trail of immune protections $h(\mathbf{x}, t)$ corresponds to

the blue trace of Fig. 1B, and the coverage or fitness gradient
 to the isolines of Fig. 1C.

In the moving frame of the wave, (u, x_2, \dots, x_d) , with $u =$
 $x_1 - vt$, the local immune protection and viral fitness can
 be expanded locally for $u, x_i \ll v\tau$ (see (25) for a similar
 treatment in a one-dimensional antigenic space):

$$182 \quad f((u, x_{i>1}); t) \approx \ln \left[R_0 \left(1 - \frac{e^{-u/r}}{1+v\tau/r} \right)^M \right] \approx f_0 + su, \quad [8]$$

where $f_0 = \ln R_0 - M \ln[1 + r/(v\tau)]$ is the average population
 fitness, and

$$184 \quad s = |\partial_{x_1} f| = \frac{M}{r} \left(R_0^{1/M} - 1 \right) \quad [9]$$

185 is the fitness gradient. Rescaling the antigenic variable x_1 as
 186 sx_1 , this process is equivalent to the evolution of a population
 187 where mutation effects are described by diffusion in fitness
 188 space with coefficient Ds^2 . This is precisely the model from
 189 which the fitness wave solution of Ref. (27, 32) was described
 190 (see SI Appendix). In the following we will use results from
 191 these works to describe the antigenic wave. However, we
 192 note that in the usual fitness wave theory, population is kept
 193 constant by construction, which implies that fitness is only
 194 relevant when compared to the mean of the population. By
 195 contrast, in our model population size is itself a dynamical
 196 variable, and fitness is defined as an absolute growth rate.
 197 In this version of the model, the fitness of the whole viral
 198 population undergoes continuous negative drift due to the
 199 constant adaptation of immune systems, encoded in the $-svt$
 200 term in Eq. 8. This negative fitness drift has an analogous
 201 effect to subtracting the mean fitness in models with constant
 202 population size, making the equivalence possible.
 203

The fitness wave theory allows us to make analytical pre-
 diction about the properties of the antigenic wave. Let us
 start with its population size N , which is regulated by how
 fast the immune system catches up with the wave. The im-
 mune turnover time τ in Eq. 4 is inversely proportional to N :
 the larger the population size, the faster immune memories
 are updated, increasing the immune pressure on current viral
 strains (lower f_0), and thus decreasing N . As the moving wave
 reaches a stable moving state, its size N becomes stable over
 time, giving the condition $(1/N)dN/dt = f_0 = 0$, which in
 turn constraints the ratio between the wave's size and speed:
 214

$$215 \quad \frac{N}{v} = \frac{MN_h}{r} \left(R_0^{1/M} - 1 \right) = N_h s. \quad [10]$$

216 But the fitness wave theory predicts that the speed of the
 217 wave itself depends on the population size. The larger N ,
 218 the more outliers at the nose of the fitness wave, and the
 219 further out they may jump in antigenic space, establishing
 220 fitter ancestors of the future population. This results in a
 221 fitness wave whose speed depends only weakly on population
 222 size and mutation rate (see (32) and SI Appendix),

$$223 \quad v_F \approx D_F^{2/3} \left[24 \ln(ND_F^{1/3}) \right]^{1/3}, \quad [11]$$

224 where $D_F = s^2 D$ and $v_F = sv$ are the diffusivity and wave
 225 speed in fitness space, which are related to their counterparts
 226 in antigenic space through the scaling factor s . Replacing this
 227 scaling into Eq. 11 yields a relation between antigenic speed
 228 and population size,

$$229 \quad v \approx D^{2/3} s^{1/3} \left[24 \ln(N(Ds^2)^{1/3}) \right]^{1/3}, \quad [12]$$




Fig. 2. Analytical prediction of wave properties. Shown are the numerical versus analytical predictions for the wave’s population size N (A), speed v (B), width σ along the wave’s direction of motion (C), and width σ_{\perp} in the direction perpendicular to motion (D), with $d = 2$ dimensions. Length are in units of the cross-reactivity range (so that $r = 1$, with no loss of generality). Parameters: $N_h = 10^8$ (squares), 10^{10} (circles), or 10^{12} (triangles); $\ln R_0 = 1$ (filled symbols) or 3 (empty symbols); $M = 1$ (small symbols) or 5 (large symbols).

which closes the system of equations: using the definition of s (Eq. 9), Eqs. 10 and 12 completely determine N and v as a function of the model’s parameters (through a transcendental equation, see SI Appendix). We validated these theoretical predictions for N and v by comparing them to numerical simulations, which show good agreement over a wide range of parameters (Fig. 2A-B). We note that the alternative fitness wave model of Desai and Fisher (28) predicts different scaling relations between speed and population size, including for an arbitrary distribution of fitness effects (30). The major difference with our description is that we assume infinitesimal and reversible fitness effects. Relaxing that assumption to account for rare but strong mutational effects would affect Eq. 12, but the dependence on N would still be logarithmic at most.

D. Shape of the antigenic wave. The width σ of the wave in the direction of motion is given by Fisher’s theorem, which relates the rate of change of the average fitness to its variance in the population: $\partial_t f = \text{Var}(f)$. In our description fitness and the antigenic dimension x_1 are linearly related with coefficient s , implying $s^2 \sigma^2 = sv$. The result of that prediction for σ is validated against numerical simulations in Fig. 2C.

The wave is led by an antigenic ‘nose’ formed by few outlying strains of reduced cross-reactivity with the concurrent immune population, generating high fitness. These strains have phenotype $u_c = s\sigma^4/4D = v^2/(4Ds)$ and fitness su_c . They serve as founder strains from which the bulk of the future population will derive some time $\sim u_c/v = \sigma^2/4D$ later (see SI Appendix). As a result, two strains taken at random

can trace back their most recent common ancestor to some average time $\langle T_2 \rangle = \alpha\sigma^2/2D$ in the past, where $\alpha \approx 1.66$ is a numerical factor estimated from simulations (32).

To explain the width σ_{\perp} of the wave in the other phenotypic dimensions than that of motion ($x_{i>1}$), we note that in these directions evolution is neutral. Two strains taken at random in the bulk are expected to have drifted, or ‘diffused’ in physical language, by an average squared displacement $\langle \Delta x_i^2 \rangle = 2DT_2$ from their common ancestor, so that their mean squared distance is $4D\langle T_2 \rangle = 2\alpha\sigma^2$ along x_i . If one assumes an approximately Gaussian wave of width σ_{\perp} , the mean square distance between two random strains along x_i should be equal to $2\sigma_{\perp}^2$. Equating the two estimates yields $\sigma_{\perp}^2 = \alpha\sigma^2$. Fig. 2D checks the validity of this prediction against simulations.

Both longitudinal and transversal fluctuations in antigenic space are instances of quantitative traits under interference selection generated by multiple small-effect mutations. The width of these traits is governed by the common relation $\langle \Delta x_i^2 \rangle = 2D\langle T_2 \rangle \sim \sigma^2$, which expresses the effective neutrality of the underlying genetic mutations (33). This relation says that antigenic variations in all dimensions scale in the same way with the model parameters, and the wave should have an approximately spherical shape. Consistently, here we find a wave with a fixed ratio $\alpha \approx 1.66$ between transverse and longitudinal variations. This implies a slightly asymmetric shape (which may be non-universal and depend on the microscopic assumptions of our mutation model).

In what parameter regime is our theory valid? The fitness wave theory we built upon is meant to be valid in the large population size, $N \gg 1$. In addition, we assumed that the fitness landscape was locally linear across the wave. This approximation should be valid all the way up to the tip of wave, given by u_c , since this is where the selection of future founder strains happen. This condition translates into $u_c \ll r$, implying $D \ll r^2/\ln(N)^2$ (using $u_c = v^2/(4Ds)$ and Eqs. 9,12), where D is in antigenic unit squared per infection cycle. This result means that one infection cycle will not produce enough mutations for the virus to leave the cross-reactivity range. In that limit, another assumption is automatically fulfilled, namely that the width of the wave be small compared to the span of immune memory: $\sigma \ll v\tau$. Our simulations, which run in the regime of very slow effective diffusion ($D/r^2 \lesssim 10^{-6}$) and have relatively large population sizes ($N \gtrsim 10^4$), satisfy these conditions. This explains the good agreement between analytics and numerics.

E. Equations of motion of the wave’s position. The wave solution allows for a simplified picture. The wave travels in the direction of the fitness gradient (or equivalent the gradient of immune coverage) with speed v (Fig. 3A). Occasionally the population splits into two separate waves that then travel away from each other and from their common ancestor (Fig. 3B). The tip of the wave’s nose, which contains the high-fitness individual that will seed the future population, determines its future position in antigenic space. In the directions perpendicular to the fitness gradient, this position diffuses neutrally with coefficient D . This motivates us to write effective equations




Fig. 3. Stochastic behaviour of the wave: diffusive motion, splits, and extinctions. **A.** The wave moves forward in antigenic space but is driven by its nose tip, which undergoes antigenic drift (diffusion) in directions perpendicular to its direction of motion. These fluctuations deviate that direction, resulting in effective angular diffusion. **B.** When antigenic drift is large, the wave may randomly split into subpopulations, creating independent waves going in different directions. Each wave can also go extinct as size fluctuations bring it to 0. **C.** Cartoon illustrating the wave's angular diffusion. Selection and drift combine to create an inertial random walk of persistence time t_{persist} . **D.** Analytical prediction (Eq. 17) for the persistence time, versus estimates from simulations. Symbols and colors are the same as in Fig. 2.

of motion for the mean position of the wave:

$$\frac{d\mathbf{x}}{dt} = -\left(v + \sqrt{2D_{\parallel}}\xi_{\parallel}(t)\right) \frac{\partial_{\mathbf{x}}c}{|\partial_{\mathbf{x}}c|} + \sqrt{2D}\xi_{\perp}(t), \quad [13]$$

$$c(\mathbf{x}, t) = \int_{-\infty}^t \frac{dt'}{\tau} e^{-\frac{t-t'}{\tau} - \frac{|\mathbf{x}-\mathbf{x}(t')|}{r}}, \quad [14]$$

where ξ_{\parallel} and ξ_{\perp} are Gaussian white noises in the directions along, and perpendicular to, the fitness gradient $\partial_{\mathbf{x}}f/|\partial_{\mathbf{x}}f| = -\partial_{\mathbf{x}}c/|\partial_{\mathbf{x}}c|$. D_{\parallel} is an effective diffusivity in the direction of motion resulting from the fluctuations at the nose tip. These fluctuations are different than suggested by D , as they involve feedback mechanisms between the wave's speed v , size N , and advancement of the fitness nose u_c . In the following, we do not consider these fluctuations, and focus on perpendicular fluctuations instead.

F. Angular diffusion and persistence of the antigenic wave.

In the description of Eqs. 13-14, the viral wave is pushed by immune protections left in its trail. The fitness gradient, and thus the direction of motion, points in the direction that is set by the wave's own path. This creates an inertial effect that stabilizes forward motion. On the other hand, fluctuations in perpendicular directions are expected to deviate the course of that motion, contributing to effective angular diffusion. To study this behaviour, we assume that motion is approximately straight in direction $x_1 = vt$, and study small fluctuations in the perpendicular directions, $\mathbf{x}_{\perp} = (x_2, \dots, x_d)$, with $|\mathbf{x}_{\perp}| \ll r$ (as illustrated in Fig. 3C). Eqs. Eq. (13)-Eq. (14) simplify to (see SI Appendix):

$$\partial_t \mathbf{x}_{\perp}(t) = \int_0^{+\infty} \frac{dt'}{T} \frac{d\mathbf{x}_{\perp}(t) - \mathbf{x}_{\perp}(t-t')}{t'} e^{-t'/T} + \sqrt{2D}\xi_{\perp}(t), \quad [15]$$

where $T = (v/r + 1/\tau)^{-1} = (r/v)R_0^{-1/M}$ is an effective memory timescale combining the host's actual immune memory, and the cross-reactivity with strains encountered in the past.

Eq. 15 may be solved in Fourier space. Defining $\tilde{\mathbf{x}}_{\perp}(\omega) = \int_{-\infty}^{+\infty} dt e^{i\omega t} \mathbf{x}_{\perp}(t)$, it becomes:

$$-i\omega \tilde{\mathbf{x}}_{\perp}(\omega) \left(1 + \frac{\ln(1 - i\omega T)}{i\omega T}\right) = \sqrt{2D}\tilde{\xi}_{\perp}(\omega). \quad [16]$$

To understand the behaviour at long times $\gg T$, we expand at small ω : $-\omega^2 \mathbf{x}_{\perp}(\omega) \approx \sqrt{8D}\tilde{\xi}_{\perp}(\omega)/T$ or equivalently in the temporal domain $\partial_t^2 \mathbf{x}_{\perp} \approx \sqrt{8D}\xi_{\perp}(t)/T$. This implies that the direction of motion, $\hat{\mathbf{e}} \sim \partial_{\mathbf{x}}f/|\partial_{\mathbf{x}}f| \sim \partial_t \mathbf{x}/|\partial_t \mathbf{x}|$, undergoes effective angular diffusion in the long run: $\partial_t \hat{\mathbf{e}} = \sqrt{8D}\xi_{\perp}(t)/(vT)$. The persistence time of that inertial motion,

$$t_{\text{persist}} = \frac{v^2 T^2}{4D} = \frac{r^2}{4D} R_0^{-2/M}, \quad [17]$$

does not depend explicitly on speed, population size, or the dimension of antigenic space. However, a larger diffusivity implies larger N and v while reducing the persistence time. Likewise, a larger reproduction number R_0 or smaller memory capacity M speeds up the wave and increases its size, but also reduces its persistence time. This implies that, for a fixed number of hosts N_h , larger epidemic waves not only move faster across antigenic space, but also change course faster.

This persistence time scales as the time it would take a single virus drifting neutrally to escape the cross-reactivity range, r^2/D . For comparison, the much shorter timescale for a population of viruses to escape from the cross-reactivity range r ,

$$t_{\text{escape}} = \frac{r}{v} = TR_0^{1/M} = \frac{N_h M}{N} (R_0^{1/M} - 1), \quad [18]$$

scales with the inverse incidence rate N_h/N . This is consistent with the whole population having been infected at least one every $\sim N_h/N$ infection cycles. This separation of time scales is consistent with the observation that evolution in the transverse directions is driven by neutral drift, which is much slower than adaptive evolution in the longitudinal direction. Both t_{persist} and t_{escape} are longer than the coalescence time of the viral population, $u_c/v \sim \sigma^2/4D$, since they reflect long-term memory from the immune system. However, while $t_{\text{escape}} \sim N_h/N$ is related to the re-infection period and is thus bounded by the hosts' immune memory (itself bounded by their lifetime, which we do not consider), t_{persist} can be longer than that. This is possible thanks to inertial effects, which are allowed by the high-order dynamics of Eq. 15 generated by the immune system. This very much like when, in mechanics, a massive object set in motion in a given direction will keep that direction without the need for an external force to maintain it.

The high-frequency behaviour of Eq. (16) has a logarithmic divergence, meaning that the total power of $\hat{\mathbf{e}}$ is infinite unless we impose a (ultraviolet) cutoff. Such a regularization emerges from the fine structure of the wave. While the motion of the wave is driven by its nose tip, the immune pressure only extends back to the recent past of the bulk of the distribution, which stands at a distance u_c away from the nose. In other words, there is a lag (and thus an gap u_c in antigenic space) between the most innovative variants that drive viral evolution, and the majority of currently circulating variants which drive host immunity. Mathematically, this implies that the domain

384 of integration of the first term in the right-hand side of Eq. (15)
 385 should start at $t_c = u_c/v$, which regularizes the divergence.
 386 A more careful analysis provided in the SI Appendix shows
 387 that this regularization does not affect the long-term diffusive
 388 behaviour of the wave.

389 **G. Canalization, speciations, and predictability of antigenic**
 390 **evolution.** We now examine how deflections of the wave in
 391 the transverse direction determines the predictability and
 392 stability of the viral quasi-species. Assuming $t \gg T$, angular
 393 diffusion causes motion to be deflected as (see SI Appendix)
 394 $\langle x_{\perp}^2 \rangle = \frac{8(d-1)D}{3T^2} t^3$. Crucially, this deflection depends on the
 395 dimension of the antigenic space, because the displacement
 396 acts additively in each of the transversal coordinates. Higher
 397 dimension means more deviation from the predictable course
 398 of the wave, and thus less predictability. We can define a
 399 predictability time scale

$$400 \quad t_{\text{predict}} \sim [8(d-1)/3]^{-1/3} T^{2/3} (r^2/D)^{1/3}, \quad [19]$$


401 which is the time it takes for prediction errors to become of
 402 the order of the cross-reactivity range. In low dimensions, this
 403 time scales as a weighted geometric mean between $t_{\text{escape}} \sim T$
 404 and $t_{\text{persist}} \sim r^2/D$. However, at high dimensions t_{predict} may
 405 be significantly reduced, causing loss of predictability even
 406 below t_{escape} . The prediction timescale is distinct from the
 407 previously discussed persistence time: t_{predict} involves the inte-
 408 grated displacement in the transversal direction, while t_{persist}
 409 quantifies the diffusion of the tangent velocity vector. Thus,
 410 t_{predict} may be interpreted as quantifying the predictability of
 411 the actual location of the next viral population in antigenic
 412 space, while t_{persist} gives the predictability of the general di-
 413 rection of evolution, which changes more slowly. Therefore,
 414 the persistence time is both harder to extract from data and
 415 less relevant for actionable predictions.

416 To get a sense of numbers, we can compare our results with
 417 epidemiological data, taking the evolution of influenza as an
 418 example, with an infection cycle time of 3 days. It is assumed
 419 that individuals lose immunity to the circulating strain of the
 420 flu within ~ 5 years ~ 500 cycles, meaning that the wave
 421 would travel a distance r in $t = 500$, i.e. $v/r \sim 2 \cdot 10^{-3}$.
 422 For instance, with $N_h = 10^9$ - 10^{10} , $R_0 = 2$, and $M = 1$,
 423 we may choose $D/r^2 = 3 \cdot 10^{-6}$ to get a speed of the same
 424 order, $v/r \sim 1.3 \cdot 10^{-3}$, and $t_{\text{persist}} \sim 2 \cdot 10^4 \sim 200$ years. By
 425 contrast, the predictability timescale t_{predict} is much shorter
 426 and depends on dimension, albeit slowly, ranging from ~ 20
 427 years for $d = 2$ to about 2 years for $d = 1000$. We stress that
 428 these numbers are obtained by scaling laws, and should not
 429 be taken as precise quantitative predictions.

430 Large deflections may also cause speciations, or splits, which
 431 occur when two substrains co-exist long enough to become
 432 independent from the immune standpoint. This happens when
 433 two sub-lineages see the difference of their transverse positions
 434 $\Delta \mathbf{x}_{\perp}$ become larger than $\Delta x_0 \sim r$, within some limited period
 435 given by the coalescence time. We estimated the rate of such
 436 splitting events using a saddle-point approximation (see SI
 437 Appendix):

$$438 \quad k_{\text{split}} \approx \sqrt{\frac{3}{8}} \frac{v^2}{4D} e^{-\mathcal{L}}, \quad \mathcal{L} = \alpha \left(\frac{s^3 R_0^{-2/M} D^2 r^4}{(d-1)v^5} \right)^{1/4} \quad [20]$$

439 with α some numerical factor. Simulations confirmed the
 440 validity of this scaling (Fig. 4a).



441 **Fig. 4. Rate of speciation.** **A.** Rescaled rate of splitting events, defined as the
 442 emergence of two substrains at distance $\Delta x_0 = 0.1r$ from each other in antigenic
 443 space, meaning that they are becoming antigenically independent. The predicted
 444 scaling, $k_{\text{split}} \sim (v^2/D)e^{-\mathcal{L}}$, as well as the definition of the collective variable
 445 \mathcal{L} as a function of the model parameters, are given by Eq. 20. The line shows a
 446 linear fit of the logarithm of the ordinate. **B.** Predicted rate of splitting as a
 447 function of the dimension d , for $R_0 = 2$, $M = 1$, $N_h = 10^9$, and $D/r^2 = 3 \cdot 10^{-6}$, with
 448 $\Delta x_0 = r$.

441 The splitting rate grows with the dimension (Fig. 4b),
 442 consistent with the intuition that departure from canalized
 443 evolution is easier when more directions of escape are available.
 444 Splitting events are expected to strongly affect our ability to
 445 predict the future course of the wave. However, the rarity
 446 of such events (exponential scaling of k_{split}) means that they
 447 will have a lower impact on predictability than deflections.
 448 These results provide a theoretical and quantitative basis from
 449 which to assess the effect of dimension on predictability, and
 450 possibly estimate d from antigenic time course data of real
 451 viral populations.

452 3. Discussion

453 In this work, we have developed an analytical theory for
 454 studying antigenic waves of viral evolution in response to
 455 immune pressure. We showed that predictability is limited
 456 by two features of antigenic evolution, transversal diffusion
 457 and lineage speciations of the antigenic wave, both of which
 458 explicitly depend on the dimensionality of antigenic space.

459 To derive these results, we explicitly embedded the anti-
 460 genic phenotype in a d -dimensional Euclidean space. This
 461 description is different from previous work that considered one-
 462 (25) or infinite-dimensional antigenic spaces (12). It allows
 463 for the possibility of compensatory mutations, and makes it
 464 easier to compare results with empirical studies of viral evo-
 465 lution projected onto low-dimensional spaces (8, 9). Unlike
 466 these studies, however, our work does not address the question
 467 how an effective dimension of antigenic space arises from the
 468 molecular architecture of immune interactions. Rather, we
 469 focused on the implications of the dimensionality of antigenic
 470 space for phenotypic evolution and its predictability.

471 Our results suggest a hierarchy of time scales for viral
 472 evolution. The shortest is the coalescence time $\langle T_2 \rangle$, which
 473 determines population turnover. Then comes t_{escape} , which
 474 is the time it takes the viral population to escape immunity
 475 elicited at a previous time point. The longest timescale is the
 476 persistence time t_{persist} , which governs the angular diffusion
 477 of the wave's direction, but has no bearing on the prediction
 478 of the actual position of the dominant strain in antigenic

space. That time scale is due to inertial effects. It does not rely directly on the hosts' immune memories, and may thus exceed their individual lifetimes. Finally, the prediction timescale t_{predict} , beyond which prediction accuracy falls below the resolution of cross-reactivity, scales between t_{escape} and t_{persist} at low dimensions. This time scales measures the predictability of transversal fluctuations, and is thus the most relevant for actual predictions of future dominant strains in antigenic space. Importantly, it decreases with the dimension of the antigenic space, and may become arbitrarily low at very high dimensions. The fact that the evolution of influenza strains are hard to predict beyond a year suggests that the effective dimension may indeed be large.

Our solution builds on the fitness wave solution for a diffusion model of mutation effects (27, 32). It implies a particular dependence of the wave's speed on the population size, Eq. 12. General distribution of non-infinitesimal mutational effects, such as considered in (30), would yield different expressions for the speed. However, we expect most of our other results to hold—in particular, all expression that do not carry an explicit logarithmic dependence on N , as well the effective equations of motion for the wave. Our results strongly rely on the assumption of a homogeneous, isotropic antigenic space. We expect our results to be affected by anisotropies (e.g. in the mutational or the cross-reactivity Kernels), or by structure in the *intrinsic* fitness landscape (i.e. not linked to immunity). Such structure may funnel the wave in preferred directions, hinder it, or favor its splitting. Generally, the local geometry and metric of the space is expected to determine the evolutionary behaviour. For instance, Yan et al. (12) assumed a Hamming distance metric in an effectively infinite antigenic space, meaning that any mutation is both an escape mutation and a candidate for a lineage split. By contrast, in our geometry, escape happens only in the direction of the wave, while splits originate from mutations perpendicular to that direction, due to the choice of a Euclidean metric. While our results emphasize the role of the effective dimension d , studying other geometrical effects is an interesting topic for future work.

Despite these caveats, it is interesting to ask whether the effective antigenic dimension d can be extracted from data. A possible scheme for doing so starts by inferring the effective model parameters. R_0 may be estimated from exponential epidemic growth in a susceptible population. Dependence of key quantities on M such as s is weak. M may be assumed to be of the order of the number of antigenically distinct infections encountered during a host's lifetime, $\sim 4-6$ (every 15 years). D/r^2 may be inferred from v/r , which can be estimated from cross-immunity assays or from the incidence rate N/N_h . Alternatively, since D/r^2 is the inverse time it takes for mutations to neutrally evade immunity, it could be estimated directly from genomic data by computing the time for unselected mutations (whose rate is inferred from synonymous mutations) to affect antigenic sites. Interestingly, if v/r and D/r^2 can be inferred independently, predictions about the wave's shape, width, angular diffusion and splitting do not depend on the particular choice of fitness wave theory. Assuming that all these parameters are known, the splitting rate, which depends sensitively on d (Fig. 4b), could be used to infer an effective dimension. Since splitting is rare and may not be observed in practice, one could define instead partial splits, where a sublineage diverges an antigenic distance $\Delta x_0 < r$

from the main lineage, for which the same scaling as Eq. 20 holds (see SI Appendix). Alternatively, our results could be used to check the consistency of dimensionality-reduction schemes based on serological assays (8–10), by testing our predicted relations between the speed of the wave, its width and length, and angular diffusion properties, and ask what choice of dimension best agrees with our theory.

Our framework should be applicable to general host-pathogens systems. For instance, co-evolution between viral phages and bacteria protected by the CRISPR-Cas system (34) is governed by the same principles of escape and adaptation as vertebrate immunity. Even more generally, our theory (Eqs. 2,3) may be relevant to the coupled dynamics of predators and preys interacting in space (geographical or phenotypic), opening potential avenues for experimental tests of these theories in synthetic microbial systems. Given the current context of the global SARS-CoV-2 pandemic, it is natural to ask whether our results could be applicable to predict its evolution. While our theory describes the long-term co-evolution of viral strains with the hosts' immune systems, in which most hosts have been exposed to at least on strain of the virus, SARS-CoV-2 is still in a phase of growth, and has not exhausted the reservoir of susceptible hosts. As the situation develops, it will be interesting to see whether its future evolution follows a Red Queen type of evolution like influenza, goes extinct, or splits into many antigenically independent sub-lineages. While our model may shed light on these questions, fine microscopic details such as geographical and population structure impose additional challenges for predictions.

4. Methods

We simulated discrete population dynamics of infected hosts $n(\mathbf{x}, t)$ and immune protections $n_h(\mathbf{x}, t) \equiv N_h h(\mathbf{x}, t)$ (all integers) on a $2D$ square lattice with lattice size Δx ranging from $10^{-5}r$ to $0.1r$. Each time step corresponds to a single infection cycle, $\Delta t = 1$. At each time step: (1) viral fitness f is computed at each occupied lattice site from the immune coverage Eq. 1; (2) viruses at each occupied lattice site are grown according to their fitness, $n(\mathbf{x}, t+1) \sim \text{Poisson}[(1+f\Delta t)n(\mathbf{x}, t)]$; (3) viruses are mutated by jumping to nearby sites on the lattice; (4) the immune system is updated according to a discrete version of Eq. 3, by implementing $n_h(\mathbf{x}, t+1) = n_h(\mathbf{x}, t) + n(\mathbf{x}, t)$ and then removing $N(t)$ protections at random (so that N_h remains constant).

To implement (1), we used a combination of exact computation of Eq. 1 and approximate methods, including one based on non-homogeneous fast Fourier transforms (35, 36). Details are given in the SI Appendix.

To implement (3), we drew the number of mutants at each occupied site from a binomial distribution $\text{Binomial}(n(\mathbf{x}, t), 1 - e^{-\mu\Delta t})$. The number of new mutations m affecting each of these mutants is drawn from a Poisson distribution of mean $\mu\Delta t$ conditioned on having at least 1 mutation. The new location of each mutant is drawn as $\mathbf{x} + \delta\mathbf{x}$, with $\delta\mathbf{x} = \text{round}(\sum_{i=1}^m \boldsymbol{\epsilon}_i)$ (rounding is applied to each dimension), where $\boldsymbol{\epsilon}_i$ is a vector of random orientation and modulus drawn from a Gamma distribution of mean $\delta \sim 2\Delta x$ and shape parameter 20. This distribution was chosen so as to maximize the number of non-zero jumps while maintaining isotropy. We then define $D = \mu \langle \delta x_1^2 \rangle / 2$.

To find the wave solution more rapidly, the viral population

was initialized as a Gaussian distribution centered at $(0, 0)$ with size N and width σ in all dimensions, to which 0.1% additional viruses are randomly added within the interval $(0; u_c)$ along x_1 (N , σ , and u_c being all given by the theory prediction). Immune protections are placed according to Eq. 6. The first 20,000 time steps serve to reach steady state and are discarded from the analysis. When a population extinction ($N = 0$) or explosion ($N = N_h/2$) occurs, the simulation is resumed at an earlier checkpoint to avoid re-equilibrating. Simulations are ended after $5 \cdot 10^6$ steps or after 20 consecutive extinctions or explosions from the same checkpoint.

In order to analyze the organization of viruses in phenotypic space, we save snapshots of the simulation at regular time intervals. For each saved snapshot we take all the coordinates with $n > 0$ and then cluster them into separate lineages through the python scikit-learn DBSCAN algorithm (37) (38) with the minimal number of samples $\text{min_samples} = 10$. The ϵ parameter defines the maximum distance between two samples that are considered to be in the neighborhood of each other. We perform the clustering for different values of ϵ and select the value that minimizes the variance of the 10th nearest neighbor distance. Clustering results are not sensitive to this choice. This preliminary clustering step is refined by merging clusters if their centroids are closer than the sum of the maximum distances of all the points in each cluster from the corresponding centroid.

From the clustered lineages we can easily obtain a series of related observables, such as its speed v obtained as the derivative of the center's position. The width of the lineage profile in the direction of motion σ as well as in the perpendicular direction σ_\perp are obtained by taking the standard deviation of the desired component of the distances of all the lineage viruses from the lineage centroid. Reported numbers are time averages of these observables. We can track their separate trajectories in antigenic space. A split of a lineage into two new lineages is defined when two clusters are detected where previously there was one, and their distance is larger than Δx_0 , the chosen threshold for calling a split.

To estimate the persistence time, we first subsample the trajectory so that the distance between consecutive points is bigger than $6(\langle\sigma\rangle + \text{std}(\sigma))$ so that fast fluctuations in the population size do not affect the inference. We take the resulting trajectory angles and smooth them with a sliding window of 5. Then we divide the trajectory into subsegments, and compute the angles mean squared displacement (MSD) over all lineages and all subsegments. We consider time lags only bigger than twice the typical smoothing time, and if the MSD trace is long enough we also require the time lag to be bigger than $2T$. Finally we only keep time lag bins with at least 10 datapoints. We fit the resulting time series to a linear function $ax + b$, and get the persistence time as $\frac{2}{a}$. We compute the reduced χ^2 as a goodness-of-fit score. Results are shown for simulations that had enough statistics to perform the fit, lasted at least 10^5 cycles, and had a reduced χ^2 below 3.

Acknowledgements. The study was supported by the European Research Council COG 724208 and ANR-19-CE45-0018 "RESP-REP" from the Agence Nationale de la Recherche and DFG grant CRC 1310 "Predictability in Evolution".

1. DH Morris, et al., Predictive Modeling of Influenza Shows the Promise of Applied Evolutionary Biology. *Trends Microbiol.* **26**, 102–118 (2018).

2. S Wang, et al., Manipulating the Selection Forces during Affinity Maturation to Generate Cross-Reactive HIV Antibodies. *Cell* **160**, 785–797 (2015).
3. JP Barton, et al., Relative rate and location of intra-host HIV evolution to evade cellular immunity are predictable. *Nat. Commun.* **7**, 1–10 (2016).
4. A Nourmohammad, J Otwinowski, JB Plotkin, Host-Pathogen Coevolution and the Emergence of Broadly Neutralizing Antibodies in Chronic Infections. *PLoS genetics* **12**, e1006171 (2016).
5. A Nourmohammad, C Eksin, Optimal evolutionary control for artificial selection on molecular phenotypes. *bioRxiv* (2019).
6. S Gandon, T Day, CJE Metcalf, BT Grenfell, Forecasting Epidemiological and Evolutionary Dynamics of Infectious Diseases. *Trends Ecol. Evol.* **31**, 776–788 (2016).
7. LA Segel, AS Perelson, Shape space: an approach to the evaluation of cross-reactivity effects, stability and controllability in the immune system. *Immunol. Lett.* **22**, 91–99 (1989).
8. DJ Smith, AS Lapedes, JCD Jong, Mapping the Antigenic and Genetic. *Science* **305**, 371–377 (2004).
9. T Bedford, et al., Integrating influenza antigenic dynamics with molecular evolution. *eLife* **2014**, 1–26 (2014).
10. JM Fonville, et al., Antibody landscapes after influenza virus infection or vaccination. *Sci. (New York, N.Y.)* **346**, 996–1000 (2014).
11. L Van Valen, A new evolutionary law. *Evol Theory* **1**, 1–30 (1973).
12. L Yan, RA Neher, BI Shraiman, Phylogenetic theory of persistence, extinction and speciation of rapidly adapting pathogens. *eLife* **8**, e44205 (2019).
13. N Strelkowa, M Lässig, Clonal interference in the evolution of influenza. *Genetics* **192**, 671–682 (2012).
14. PJ Gerrish, RE Lenski, The fate of competing beneficial mutations in an asexual population. *Genetica* **102/103**, 127–144 (1998).
15. PA Rota, et al., Lineages of Influenza Type B Virus since 1983. *Virology* **68**, 59–68 (1990).
16. PA White, Evolution of norovirus. *Clin. Microbiol. Infect.* **20**, 741–745 (2014).
17. NM Ferguson, AP Galvani, RM Bush, Ecological and immunological determinants of influenza evolution. *Nature* **422**, 428–433 (2003).
18. T Bedford, A Rambaut, M Pascual, Canalization of the evolutionary trajectory of the human influenza virus. *BMC Biol.* **10**, 38 (2012).
19. D Zinder, T Bedford, S Gupta, M Pascual, The Roles of Competition and Mutation in Shaping Antigenic and Genetic Diversity in Influenza. *PLoS Pathog.* **9** (2013).
20. F Wen, T Bedford, S Cobey, Explaining the geographical origins of seasonal influenza A (H3N2). *Proc. Royal Soc. B: Biol. Sci.* **283** (2016).
21. FT Wen, A Malani, S Cobey, The beneficial effects of vaccination on the evolution of seasonal influenza. *bioRxiv* (2017).
22. JR Gog, BT Grenfell, Dynamics and selection of many-strain pathogens. *PNAS* **2002** (2002).
23. K Koelle, M Kamradt, M Pascual, Understanding the dynamics of rapidly evolving pathogens through modeling the tempo of antigenic change : In fl uenza as a case study. *Epidemics* **1**, 129–137 (2009).
24. J Marchi, M Lässig, T Mora, AM Walczak, Multi-lineage evolution in viral populations driven by host immune systems. *Pathogens* **8**, 1–16 (2019).
25. IM Rouzine, G Rozhnova, Antigenic evolution of viruses in host populations. *PLoS Pathog.*, 1–16 (2018).
26. IM Rouzine, J Wakeley, JM Coffin, The solitary wave of asexual evolution. *Proc. Natl. Acad. Sci. United States Am.* **100**, 587–592 (2003).
27. E Cohen, DA Kessler, H Levine, Front propagation up a reaction rate gradient. *Phys. Rev. E - Stat. Nonlinear, Soft Matter Phys.* **72**, 1–11 (2005).
28. MM Desai, DS Fisher, Beneficial mutation-selection balance and the effect of linkage on positive selection. *Genetics* **176**, 1759–1798 (2007).
29. O Hallatschek, The noisy edge of traveling waves. *Proc. Natl. Acad. Sci. United States Am.* **108**, 1783–1787 (2011).
30. BH Good, IM Rouzine, DJ Balick, O Hallatschek, MM Desai, Distribution of fixed beneficial mutations and the rate of adaptation in asexual populations. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 4950–4955 (2012).
31. LS Tsimring, H Levine, DA Kessler, RNA virus evolution via a fitness-space model. *Phys. Rev. Lett.* **76**, 4440–4443 (1996).
32. Ra Neher, O Hallatschek, Genealogies of rapidly adapting populations. *Proc. Natl. Acad. Sci. United States Am.* **110**, 437–442 (2013).
33. T Held, D Klemmer, M Lässig, Survival of the simplest in microbial evolution. *Nat. Commun.* **10** (2019).
34. ER Westra, S Van Houte, S Gandon, R Whitaker, The ecology and evolution of microbial CRISPR-Cas adaptive immune systems. *Philos. Transactions Royal Soc. B: Biol. Sci.* **374** (2019).
35. J Keiner, S Kunis, D Potts, Using NFFT 3—a software library for various nonequispaced fast Fourier transforms. *ACM Transactions on Math. Softw. (TOMS)* **36**, 1–30 (2009).
36. D Potts, G Steidl, A Nieslony, Fast convolution with radial kernels at nonequispaced knots. *Numer. Math.* **98**, 329–351 (2004).
37. F Pedregosa, et al., Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).
38. M Ester, HP Kriegel, J Sander, X Xu, A density-based algorithm for discovering clusters in large spatial databases with noise. (AAAI Press), pp. 226–231 (1996).